EslB is required for cell wall integrity and cell division in *Listeria monocytogenes*

Jeanine Rismondo[a,*,#], Maria Yacoub[a], Ashima Wadhawan[b], Marc S. Dionne[b], Angelika

Gründling[a,#]

[a] Section of Molecular Microbiology and Medical Research Council Centre for Molecular

Bacteriology and Infection, Imperial College London, London SW7 2AZ.

[b] Department of Life Sciences, Medical Research Council Centre for Molecular Bacteriology

and Infection, Imperial College London, London SW7 2AZ.

[*] Present Address: Department of General Microbiology, GZMB, Georg-August-Universität

Göttingen, 37077 Göttingen, Germany

[#]To whom correspondence should be addressed:

Jeanine Rismondo – jrismon@gwdg.de, Angelika Gründling – a.grundling@imperial.ac.uk

Running title: Characterization of an *eslB* deletion strain

ABSTRACT

Lysozyme is an enzyme found in secretions including tears, saliva and mucus and is an

important component of the innate immune system. It functions by hydrolysing the

peptidoglycan layer of bacteria. The human pathogen *Listeria monocytogenes* is intrinsically

lysozyme resistant. The peptidoglycan N-deacetylase PgdA and O-acetyltransferase OatA are

two known factors contributing to its lysozyme resistance. Furthermore, it was shown that the

absence of components of an ABC transporter, here referred to as EslABC, leads to a reduction

in lysozyme resistance. However, the function of the transporter and how its activity is linked

to lysozyme resistance are still unknown. To investigate this further, a strain with a deletion in

eslB, coding for the membrane component of the ABC transporter, was constructed in the *L.*

monocytogenes strain 10403S. The *eslB* mutant showed as expected a 40-fold reduction in the

minimal inhibitory concentration to lysozyme. Using growth and autolysis assays, we further

show that the absence of EslB manifests in a growth defect in media containing high

concentrations of sugars and increased endogenous cell lysis. However, no growth defect could

be detected within host cells. On the other hand, the *eslB* mutant showed slight changes in its

peptidoglycan structure as well as a noticeable cell division defect and formed elongated cells.

Based on our results, we hypothesize that EslB is important for the maintenance of cell wall

integrity in *L. monocytogenes* under specific growth and stress conditions.

IMPORTANCE

The ABC transporter EslABC is associated with the intrinsic lysozyme resistance of *Listeria*

monocytogenes. However, the exact role of the transporter in this process and in the physiology

of *L. monocytogenes* is unknown. Using different assays to characterize an *eslB* deletion strain,

we found that the absence of EslB not only affects lysozyme resistance, but also endogenous

autolysis, cell division and the ability of the bacterium to grow in media containing high

concentrations of sugars. Our results indicate that EslB is by a yet unknown mechanism an

important determinant for cell wall integrity in *L. monocytogenes*.

INTRODUCTION

Gram-positive bacteria are surrounded by a complex cell wall, which is composed of a thick layer of peptidoglycan (PG) and cell wall polymers. The bacterial cell wall is important for the maintenance of cell shape, the ability of bacteria to withstand harsh environmental conditions and to prevent cell lysis (Höltje, 1998, van Heijenoort, 2001). Due to its importance, cell wall-targeting antibiotics such as β-lactam, glycopeptide, fosfomycin and moenomycin antibiotics are commonly used to treat bacterial infections (Koch, 2003, Sarkar et al., 2017). These cell-wall targeting antibiotics inhibit enzymes involved in different stages of the PG biosynthesis process (Sarkar et al., 2017). Moenomycin and β-lactam antibiotics, for instance, block the glycosyltransferase and transpeptidase activity of penicillin binding proteins, respectively, which are required for the polymerization and crosslinking of the glycan strands (Huber and Nesemann, 1968, Sauvage and Terrak, 2016, Tipper and Strominger, 1965). Peptidoglycan is also the target of the cell wall hydrolase lysozyme, which is a component of animal and human secretions such as tears and mucus. Lysozyme cleaves the glycan strands of PG by hydrolysing the 1,4-β-linkage between N-acetylmuramic acid (MurNAc) and N-acetylglucosamine (GlcNAc). This reaction leads to a loss of cell integrity and results in cell lysis (Callewaert and Michiels, 2010). The intracellular human pathogen *Listeria monocytogenes* is intrinsically resistant to lysozyme due to modifications of its PG. The N-deacetylase PgdA deacetylates GlcNAc residues, whereas MurNAc residues are acetylated by the O-acetyltransferase OatA (Boneca et al., 2007, Aubry et al., 2011). Consequently, deletion of either of these enzymes results in reduced lysozyme resistance (Boneca et al., 2007, Aubry et al., 2011). One or both of these enzymes are also present in other bacterial pathogens and important for lysozyme resistance, such as for instance PgdA in *Streptococcus pneumoniae,* OatA in *Staphylococcus aureus* and PgdA and OatA in *Enterococcus faecalis* (Benachour et al., 2012, Hébert et al., 2007, Vollmer and Tomasz, 2000, Bera et al., 2005). Besides enzymes that directly alter the peptidoglycan structure, a number of other factors have been shown to contribute to lysozyme resistance in diverse bacteria. For instance, the cell wall polymer wall teichoic acid and the two-component system GraRS contribute to lysozyme resistance in *S. aureus* (Bera et al., 2007, Herbert et al., 2007). In *E. faecalis*, the extracytoplasmic function sigma factor SigV is required for the upregulation of *pgdA* expression in the presence of lysozyme (Le Jeune et al., 2010, Benachour et al., 2012). Recently, other factors have also been identified, which contribute to the intrinsic lysozyme resistance of *L. monocytogenes* such as the predicted carboxypeptidase PbpX, the transcription factor DegU and the noncoding RNA Rli31 (Burke et al., 2014). DegU and Rli31 are involved in the regulation of *pgdA* and *pbpX* expression in *L. monocytogenes*

(Burke et al., 2014). Furthermore, components of a predicted ABC transporter encoded by the

lmo2769-6 operon in *L. monocytogenes* and here referred to as *eslABCR* for elongation, sugar-

and lysozyme sensitive phenotype (Fig. 1), have been associated with lysozyme resistance

(Burke et al., 2014, Durack et al., 2015, Toledo-Arana et al., 2009). An *eslB* mutant strain was

also shown to be more sensitive to cefuroxime and cationic antimicrobial peptides (Burke et

al., 2014).

ABC transporters can either act as importers or exporters. Importers are involved in the

uptake of sugars, peptides or other metabolites, which are recognized by substrate binding

proteins. On the other hand, toxic compounds such as antibiotics can be exported by ABC

exporters (Higgins, 1992, Berntsson et al., 2010, Tanaka et al., 2018). ABC transporters are

usually composed of homo- or heterodimeric cytoplasmic nucleotide-binding domain (NBD)

proteins, also referred to as ATP-binding cassette proteins, and homo- or heterodimeric

transmembrane domain (TMD) proteins (Locher, 2016). In ABC exporters belonging to the B-

family, such as the *S. aureus* multidrug transporter Sav1866, NBDs and TMDs are fused

(Dawson and Locher, 2006, Locher, 2016). Other ABC transporters, e.g. type I and type II

ABC importers, consist of a complex of heterodimers of NBD- and TMD-containing proteins

(Locher, 2016). In addition to NBDs and TMDs, type I and type II ABC importers have an

extracellular substrate binding protein (SBP), which is important for the delivery of specific

substrate molecules to the transporter (Davidson et al., 2008). Type III or energy-coupling

factor (ECF) transporters use a membrane-integrated S component for substrate-binding

(Mächtel et al., 2019, Slotboom, 2014). The *esl* operon encodes EslA, the NBD protein, EslB,

the TMD protein forming part of the ABC transporter, EslC, a membrane protein of unknown

function and EslR, the RpiR-type transcriptional regulator (Fig. 1). So far, it is not known

whether the predicted ABC transporter EslABC acts as an importer or exporter and its exact

cellular function has not been identified. Furthermore, it has not been investigated whether

EslC is a component of the ABC transporter encoded in the *esl* operon, e.g. a protein similar to

the S component of the ECF transporters, or whether it functions independent from EslAB.

Here, we show that the absence of EslB, the transmembrane part of the ABC transporter, not

only leads to an increased lysozyme sensitivity, but also to an increased endogenous cell lysis

and a cell division defect. We hypothesize that EslB may be required for the maintenance of

the cell wall integrity of *L. monocytogenes* during stress conditions.

MATERIALS AND METHODS

Bacterial strains and growth conditions. All strains and plasmids used in this study are listed in Table S1. *Escherichia coli* strains were grown in Luria-Bertani (LB) medium and *Listeria monocytogenes* strains in brain heart infusion (BHI) medium at 37°C unless otherwise stated. If necessary, antibiotics and supplements were added to the medium at the following concentrations: for *E. coli* cultures, ampicillin (Amp) at 100 µg/ml, chloramphenicol (Cam) at 20 µg/ml and kanamycin (Kan) at 30 µg/ml, and for *L. monocytogenes* cultures, Cam at 10 µg/ml, erythromycin (Erm) at 5 µg/ml, Kan at 30 µg/ml and IPTG at 1 mM.

Strain and plasmid construction. All primers used in this study are listed in Table S2. For the markerless inframe deletion of *lmo2768* (*lmrg_01927*, *eslB*), approximately 1kb-DNA fragments up- and downstream of the *eslB* gene were amplified by PCR using primer pairs ANG2532/2533 and ANG2534/2535. The resulting PCR products were fused in a second PCR using primers ANG2532/2535, the product was cut with BamHI and XbaI and ligated with plasmid pKSV7 that had been cut with the same enzymes. The resulting plasmid pKSV7-Δ*eslB* was recovered in *E. coli* XL1-Blue yielding strain ANG4236. The plasmid was subsequently transformed into *L. monocytogenes* strain 10403S and *eslB* deleted by allelic exchange using a previously described procedure (Camilli et al., 1993). The deletion of *eslB* was verified by PCR. The deletion procedure was performed with two independent transformants and resulted in the construction of two independent *eslB* mutant strains 10403SΔ*eslB*$_{(1)}$ (ANG4275) and 10403SΔ*eslB*$_{(2)}$ (ANG5662). For complementation analysis, pIMK3-*eslB* was constructed, in which the expression of *eslB* can be induced by IPTG. The gene *eslB* was amplified using primer pair ANG2812/ANG2813, cut with NcoI and SalI and fused with pIMK3 that had been cut with the same enzymes. The resulting plasmid pIMK3-*eslB* was recovered in *E. coli* XL1-Blue yielding strain ANG4647. Due to difficulties in preparing electrocompetent cells of *L. monocytogenes eslB* mutant strains, plasmid pIMK3-*eslB* was first electroporated into the wildtype *L. monocytogenes* strain 10403S yielding strain 10403S pIMK3-*eslB* (ANG4678). In the second step, *eslB* was deleted from the genome of strain ANG4678 resulting in the construction of the first *eslB* complementation strain 10403SΔ*eslB*$_{(1)}$ pIMK3-*eslB* (ANG4688, short 10403SΔ*eslB*$_{(1)}$ compl.). In addition, complementation plasmid pPL3e-P$_{eslA}$-*eslABC* was constructed. To this end, the *eslABC* genes including the upstream promoter region were amplified by PCR using primers ANG3349/ANG3350. The resulting PCR product was cut with SalI and BamHI and fused with plasmid pPL3e that had been cut with the same enzymes.

Plasmid pPL3e-P$_{eslA}$-*eslABC* was recovered in *E. coli* XL1-Blue yielding strain ANG5660.

Next, plasmid pPL3e-P$_{eslA}$-*eslABC* was transformed into *E. coli* SM10 yielding strain

ANG5661. Lastly, SM10 pPL3e-P$_{eslA}$-*eslABC* was used as a donor strain to transfer plasmid

pPL3e-P$_{eslA}$-*eslABC* by conjugated into *L. monocytogenes* strain 10403SΔ*eslB*$_{(2)}$ (ANG5662)

using a previously described method (Lauer et al., 2002). This resulted in the construction of

the second *eslB* complementation strain 10403SΔ*eslB*$_{(2)}$ pPL3e-P$_{eslA}$-*eslABC* (ANG5663, short

10403SΔ*eslB*$_{(2)}$ compl.).

Whole genome sequencing. Genomic DNA of *L. monocytogenes* was extracted using the

FastDNA^{TM} Kit (MP Biomedicals) and libraries for sequencing prepared using the Illumina

Nextera DNA kit. The samples were sequenced at the London Institute of Medical Sciences

using an Illumina MiSeq machine and a 150 paired end Illumina kit. The reads were trimmed,

mapped to the *L. monocytogenes* 10403S reference genome (NC_017544) and single

nucleotide polymorphisms (SNPs) with a frequency of at least 80% and small deletions (zero

coverage) identified using CLC workbench genomics (Qiagen).

Growth analysis. *L. monocytogenes* strains were grown overnight in 5 ml BHI medium at

37°C with shaking. The next day, these cultures were used to inoculate 15 ml fresh BHI

medium or BHI medium containing 0.5 M sucrose, fructose, glucose, maltose, galactose or

sodium chloride to an OD$_{600}$ of 0.05. The cultures were incubated at 37°C with shaking at 180

rpm, OD$_{600}$ readings were taken every hour for 8 h. Averages values and standard deviations

from three independent experiments were plotted.

Fluorescence and phase contrast microscopy. Overnight cultures of the indicated *L.*

monocytogenes strains were diluted 1:100 in BHI medium and grown for 3 h at 37°C. For

staining of the bacterial membrane, 100 µl of these cultures were mixed with 5 µl of 100 µg/ml

nile red solution and incubated for 20 min at 37°C. The cells were washed twice with PBS and

subsequently suspended in 50 µl of PBS. 1-1.5 µl of the different samples were subsequently

spotted on microscope slides covered with a thin agarose film (1.5 % agarose in distilled water),

air-dried and covered with a cover slip. Phase contrast and fluorescence images were taken at

1000x magnification using the Zeiss Axio Imager.A1 microscope coupled to the AxioCam

MRm and processed using the Zen 2012 software (blue edition). The nile red fluorescence

signal was detected using the Zeiss filter set 00. The length of 300 cells was measured for each

experiment and the median cell length was calculated. Averages values and standard deviations

of the median cell length of three independent experiments were plotted.

Determination of minimal inhibitory concentration (MIC). The minimal inhibitory

concentration for the cell wall-acting antibiotics penicillin and moenomycin and the cell wall

hydrolase lysozyme was determined in 96-well plates using a microbroth dilution assay.

Approximately 10^4 *L. monocytogenes* cells were used to inoculate 200 µl BHI containing two-

fold dilutions of the different antimicrobials. The starting antibiotic concentrations were: 0.025

µg/ml for penicillin G, 0.2 µg/ml moenomycin and 10 mg/ml or 0.25 mg/ml for lysozyme. The

96-well plates were incubated at 37°C with shaking at 500 rpm in a plate incubator

(Thermostar, BMG Labtech) and OD_{600} determined after 24 hours of incubation. The MIC

value refers to the antibiotic concentration at which bacterial growth was inhibited by >90%.

Peptidoglycan isolation and analysis. Overnight cultures of 10403S$\Delta eslB_{(1)}$ and

10403S$\Delta eslB_{(1)}$ compl. were diluted in 1 L BHI broth (supplemented with 1 mM IPTG for

strain 10403S$\Delta eslB_{(1)}$ compl.) to an OD_{600} of 0.06 and incubated at 37°C. At an OD_{600} of 1,

bacterial cultures were cooled on ice for 1h and the bacteria subsequently collected by

centrifugation. The peptidoglycan was purified, digested with mutanolysin and the

muropeptides analyzed by HPLC using an Agilent 1260 infinity system, as previously

described (de Jonge et al., 1992, Corrigan et al., 2011). Peptidoglycan of the wildtype *L.*

monocytogenes strain 10403S was purified and analyzed in parallel. The chromatogram of the

same wild-type control strain was recently published (Rismondo et al., 2019) and also used as

part of this study, since all strains were analyzed at the same time. The major peaks 1-6 were

assigned according to previously published HPLC spectra (Burke et al., 2014, Rismondo et al.,

2015), with peaks 2, 4, 5 and 6 corresponding to N-deacetylated GlcNAc residues. Peaks 1-2

correspond to monomeric and peaks 4-6 to dimeric (crosslinked) muropeptide fragments. The

Agilent Technology ChemStation software was used to integrate the areas of the main

muropeptide. For quantification, the sum of the peak areas was set to 100% and the area of

individual peaks was determined. The sum of values for peaks 3-6 corresponds to the %

crosslinking, whereas the deacetylation state was calculated by adding up the values for peaks

4, 5 and 6. Averages values and standard deviations were calculated from three independent

extractions.

Autolysis assays. *L. monocytogenes* strains were diluted in BHI or BHI medium supplemented with 0.5 M sucrose to an OD_{600} of 0.05 and grown for 4 h at 37°C. Cells were collected by centrifugation and resuspended in 50 mM Tris-HCl, pH 8 to an OD_{600} of 0.7-0.9 and incubated at 37°C. For penicillin- and lysozyme-induced lysis, 25 µg/ml penicillin G or 2.5 µg/ml lysozyme was added to the cultures. Autolysis was followed by determining OD_{600} readings every 15 min.

Cell culture. Bone marrow-derived macrophages (BMMs) were extracted from female C57BL/6 mice as described previously (Stapels et al., 2018). BMMs were a gift from Charlotte S. C. Michaux and Sophie Helaine. 5×10^5 BMMs were seeded into each well of a 24-well plate and grown overnight in 500 µl high glucose Dulbecco's Modified Eagle Medium (DMEM) at 37°C and 5% CO_2. *L. monocytogenes* strains were grown overnight without shaking in 2 ml BHI medium at 30°C. The next morning, bacteria were opsonized with 8% mouse serum (Sigma) at room temperature for 20 min and BMMs were infected for one hour at a multiplicity of infection (MOI) of 2. BMMs were washed with PBS and 1 ml DMEM containing 40 µg/ml gentamycin was added to kill extracellular bacteria. After 1 h, cells were washed with PBS and covered with 1 ml DMEM containing 10 µg/ml gentamycin. The number of recovered bacteria was determined 2, 4, 6 and 8 h post infection. To this end, BMMs were lysed using 1 ml PBS containing 0.1% (v/v) triton X-100 and serial dilutions were plated on BHI agar plates. The number of colony forming units (CFUs) was determined after incubating the plates overnight at 37°C. Three technical repeats were performed for each experiment and average values calculated. Average values and standard deviations from three independent experiments were plotted.

***Drosophila melanogaster* infections.** Fly injections were carried out with microinjection needles produced from borosilicate glass capillaries (World Precision Instruments TW100-4) and a needle puller (Model PC-10, Narishige). Injections were performed using a Picospritzer III system (Parker Hannifin), and the injection volume was calibrated by expelling a drop of liquid from the needle into a pot of mineral oil and halocarbon oil (both Sigma). The expelled drop was measured using the microscope graticule to obtain a final injection volume of 50 nanolitres (nl). Flies were then anesthetized with CO_2 and injected with either 50 nl of bacterial suspension in PBS or sterile PBS. 5-7-day old age matched male flies were used for all experiments. Flies were grouped into uninjected control, wounding control (injection with sterile PBS), and flies infected with *L. monocytogenes*. Each group consisted of 58-60 flies.

All survival experiments were conducted at 29°C. Dead flies were counted daily. Food vials

were placed horizontally to reduce the possibility of fly death from flies getting stuck to the

food, and flies were transferred to fresh food every 3-4 days. For the quantification of the

bacterial load, 16 flies per condition and per bacterial strain were collected at the indicated time

points. The flies were homogenised in 100 µl of TE-buffer pH 8 containing 1% Triton X-100

and 1% Proteinase K (NEB, P8107S). Homogenates were incubated for 3 h at 55°C followed

by a 10 min incubation step at 95°C. Following incubation, qPCR was carried out using the

actA gene specific primers EGD-E_ActA_L1 and EGD-E_ActA_R1 to determine the number

of bacterial colony forming units. PCR was performed with Sensimix SYBR Green no-ROX

(Bioline) on a Corbett Rotor-Gene 6000. The cycling conditions were as follows: Hold 95°C

for 10 min, then 45 cycles of 95°C for 15 s, 57°C for 30 s, 72°C for 30 s, followed by a melting

curve. Gene abundances were calculated as previously described (Sharrock et al., 2019).

RESULTS

Deletion of *eslB* in *L. monocytogenes* leads to lysozyme resistance and a slight increase in

peptidoglycan crosslinking.

Previously it has been shown that *L. monocytogenes* strains with mutations in the *eslABCR*

operon (Fig. 1A) display decreased resistance towards the cell wall hydrolase lysozyme (Burke

et al., 2014, Durack et al., 2015). The *eslABC* operon codes for a predicted ABC transporter,

however, the function of this transporter and how its activity is linked to lysozyme resistance

are still unknown. To investigate this further, a strain with a markerless in-frame deletion in

eslB, coding for the membrane component of the ABC transporter, was constructed in the *L.*

monocytogenes strain 10403S. Confirming previous results, the *eslB* deletion strain had a 40-

fold lower MIC for lysozyme as compared to the wildtype strain (Fig. 2A) (Durack et al., 2015,

Burke et al., 2014). This phenotype could be complemented and strain 10403SΔ*eslB₍₁₎* compl.,

in which *eslB* is expressed from an IPTG-inducible promoter, is even slightly more resistant to

lysozyme as compared to the wildtype strain (Fig. 2A). Next, we tested whether the resistance

towards two other cell wall-targeting antibiotics, namely penicillin and moenomycin, is

changed upon deletion of *eslB*. The MIC values obtained for the wildtype, *eslB* deletion and

eslB complementation strains were comparable (Fig. 2B-C), suggesting that the deletion of

eslB does not lead to a general sensitivity to all cell wall-acting antimicrobials but is specific

to lysozyme. In *L. monocytogenes*, lysozyme resistance is achieved by the modification of the

peptidoglycan (PG) by N-deacetylation via PgdA and O-acetylation via OatA (Boneca et al.,

2007, Aubry et al., 2011). To assess whether deletion of *eslB* impacts the N-deacetylation and

crosslinking of PG, PG was isolated from wildtype 10403S, the *eslB* deletion and

complementation strains, digested with mutanolysin and the muropeptides analysed by high

performance liquid chromatography (HPLC). This analysis revealed a slight increase in the PG

crosslinking in the *eslB* mutant strain (68±0.53%) compared to the wildtype (65.47±0.31%)

and the complementation strain grown in the presence of IPTG (64.57±2.3%) (Fig. 3). The PG

isolated from the *eslB* deletion strain was also slightly more deacetylated (71.54±0.21%) as

compared to the wildtype (67.17±0.31%) and the complementation strain (67±2.27%) (Fig. 3),

which should theoretically result in an increase and not decrease in lysozyme resistance. Taken

together, while slight changes in the PG structure were observed in the *eslB* mutant strain, our

results suggest that the lysozyme sensitivity of the *eslB* deletion strain is not caused by a defect

in PgdA enzyme activity and increased N-acetylation of GlcNAc residues.

Deletion of *eslB* results in increased endogenous and lysozyme-induced lysis.

The observed lysozyme sensitivity of the *eslB* deletion strain could also be a result of an

increased autolysis due to an impaired cell integrity. To test this, autolysis assays were

performed. To this end, the *L. monocytogenes* wildtype strain 10403S, the *eslB* deletion and

complementation strains were grown for 4 h in BHI medium, bacteria collected by

centrifugation and suspended in 50 mM Tris-HCl buffer (pH 8). After 2 h incubation at 37°C,

the OD_{600} of the suspensions of the wildtype and *eslB* complementation strain had dropped to

89.9±1.6% and 86.5±2.9% of the initial OD_{600}, respectively. (Fig. 4A). Enhanced endogenous

cell lysis was observed for the *eslB* mutant strain and the OD_{600} of the suspensions had dropped

to 68.8±1.7% (Fig. 4A). Interestingly, the addition of penicillin had no impact on the autolysis

rate of any of the strains tested (Fig. 4B). On the other hand, the addition of 2.5 μg/ml lysozyme

increased the autolysis rate of all strains, but had a particularly drastic effect on the *eslB* mutant

strain. After 30 min, the OD_{600} reading of a suspension of the *eslB* deletion strain had dropped

to 50.3±10.2%. For the wildtype and *eslB* complementation strains, it took 90 min to see a 50%

reduction in OD_{600} readings of the suspensions (Fig. 4B). These results show that the absence

of EslB leads to an increased endogenous autolysis of *L. monocytogenes* cells that can be

further enhanced by the addition of lysozyme.

Deletion of *eslB* results in a growth defect in high sugar media.

The bacterial cell wall is an important structure to maintain the cell integrity and to prevent lysis due to the high internal turgor pressure or when bacteria experience changes in the external osmolality. Cell wall defects leading to an impaired cell integrity could affect the growth of bacteria in environments with high osmolalities, e.g. in the presence of high salt or sugar concentrations. Next, we compared the growth of the wildtype, the *eslB* mutant and complementation strains at 37°C in different media. No growth difference could be observed between the strains tested, when grown in BHI medium (Fig. 5A). However, the *eslB* deletion strain grew slower in BHI medium containing 0.5 M sucrose as compared to the wildtype and the *eslB* complementation strain (Fig. 5B). A similar growth phenotype could be observed when the strains were grown in BHI medium containing either 0.5 M fructose, glucose, maltose or galactose (Fig. S1). In contrast, the presence of 0.5 M NaCl did not affect the growth of the *eslB* deletion strain (Fig. 5C). These results suggest that the observed growth defect seen for the *eslB* mutant is not solely caused by the increase in external osmolality, but rather seems to be specific to the presence of high concentrations of sugars.

Next, we wanted to determine which impact the growth in the presence of high levels of sucrose has on endogenous bacterial autolysis rates. To this end, the wildtype 10403S, *eslB* mutant and complementation strains were grown in BHI medium supplemented with 0.5 M sucrose, cell suspensions prepared in Tris-buffer and used in autolysis assays. While the wildtype and complementation strain showed similar autolysis rates following growth in BHI sucrose medium (Fig. 6A) as after growth in BHI medium (Fig 4A), the *eslB* mutant that had been transferred into Tris-buffer after growth in BHI 0.5 M sucrose medium lysed rapidly (Fig. 6A). The lysis of the *eslB* mutant strain could be further enhanced by the addition of 25 µg/ml penicillin, a concentration which only acts bacteriostatic on the *L. monocytogenes* wildtype strain 10403S (Fig. 6B). These findings strengthen the hypothesis that the absence of EslB results in an impaired cell wall integrity.

The *eslB* deletion strain is impaired in cell division, but not in virulence.

The increased endogenous autolysis together with the observed changes in the PG structure of the *eslB* deletion strain could result in an increased sensitivity to autolysins. The major autolysins of *L. monocytogenes* are p60 and NamA (MurA), which hydrolyze PG and are required for daughter cell separation during cell division (Pilgrim et al., 2003, Carroll et al., 2003). Absence of either p60 or NamA results in the formation of cell chains (Carroll et al., 2003, Pilgrim et al., 2003). We thus wondered whether deletion of *eslB* causes changes in the

cell morphology in *L. monocytogenes*. Microscopic analysis revealed that cells lacking EslB

are significantly longer with a median cell length of 3.36 ± 0.49 μm as compared to the *L.*

monocytogenes wildtype strain, which produced cells with a length of 1.91 ± 0.02 μm (Fig. 7),

highlighting that the absence of EslB therefore results in a cell division defect.

Next, we wanted to assess whether the impaired cell integrity and the observed cell division

defect would also affect the virulence of the *L. monocytogenes eslB* mutant. Of note, in a

previous study, it was shown that deletion of *eslA*, coding for the ATP-binding protein

component of the ABC transporter, has no effect on the cell-to-cell spread of *L. monocytogenes*

(Durack et al., 2015). Before assessing the contribution of EslB to the virulence potential of *L.*

monocytogenes, we wanted to ensure that the *eslB* mutant we constructed did not contain any

other genomic alteration that could impact the pathogenicity of this strain. Our whole genome

sequencing (WGS) analysis revealed that strain 10403S∆*eslB(1)* contained an additional small

deletion in gene *lmo2396* coding for an internalin protein with a leucine-rich repeat (LRR) and

a mucin-binding domain (Table S3). While to the best of our knowledge, the contribution of

Lmo2396 to the pathogenicity of *L. monocytogenes* has not yet been investigated, other

internalins are important and well-established virulence factors (Gaillard et al., 1991, Bierne et

al., 2007) and hence our original *eslB* mutant strain 10403S∆*eslB(1)* might not be ideal to

investigate the contribution of EslB to the virulence potential of *L. monocytogenes*. Our WGS

analysis also revealed a single point mutation in gene *lmo2342*, coding for a pseudouridylate

synthase in the complementation strain 10403S∆*eslB(1)* compl. (Table S3). Since we identified

an additional mutation in a gene coding for a potential virulence factor in our first *eslB* mutant

strain, we constructed a second independent *eslB* mutant, 10403S∆*eslB(2)*. We also constructed

a second complementation strain, strain 10403S∆*eslB(2)* p*PeslA-eslABC* (or short

10403S∆*eslB(2)* compl.), in which the *eslABC* genes are expressed from the native *eslA*

promoter from a chromosomally integrated plasmid. Our WGS analysis revealed that strain

10403S∆*eslB(2)* did not contain any secondary mutations (Table S3). A 1-bp deletion in gene

lmo2022 encoding a predicted NifS-like protein required for NAD biosynthesis, was identified

in strain 10403S∆*eslB(2)* compl. (Table S3), which if non-complementable phenotypes are

observed needs to be kept in mind.

Next, we confirmed that our second *eslB* mutant strain 10403S∆*eslB(2)*, which does not contain

any additional chromosomal mutations, showed the same lysozyme sensitivity, growth defect

in BHI sucrose medium and formed elongated cells as our original *eslB* mutant strain

10403S∆*eslB(1)* (Fig. S2). Furthermore, all phenotypes could be reverted back to wildtype

levels in our newly constructed complementation strain 10403SΔ*eslB(2)* compl., in which *eslB*

is expressed along with *eslA and eslC* from their proceeding native promoter (Fig. S2).

To determine whether EslB is involved in the virulence of *L. monocytogenes*, primary mouse

macrophages were infected with wildtype 10403S, the *eslB* mutant 10403SΔ*eslB(2)* and

complementation strain 10403SΔ*eslB(2)* compl.. All three strains showed a comparable

intracellular growth pattern (Fig. 8A), suggesting that EslB does not impact the ability of *L.*

monocytogenes to grow in primary mouse macrophages. Next, we assessed the ability of the

eslB deletion strains to kill *Drosophila melanogaster* as lysozyme is one important component

of its innate immune response (Daffre et al., 1994). All uninfected flies (U/C) and 96.6% of

the flies that were injected with PBS survived the duration of the experiment (Fig. 8B). Flies

infected with strains 10403SΔ*eslB(1)* and 10403SΔ*eslB(2)* showed an increase in survival in

comparison to the wildtype strain. However, these differences were not statistically significant

(Fig. 8B). In addition, the difference in killing by the 10403SΔ*eslB(2)* could not be

complemented by the expression of *eslABC* in strain 10403SΔ*eslB(2)* compl. (Fig. 8B). Finally,

no statistically significant differences could be observed for the bacterial load of flies infected

with the different *L. monocytogenes* strains (Fig. 8C). These results indicate that, while EslB

does not impact the ability of *L. monocytogenes* to infect and kill mammalian macrophages or

Drosophila melanogaster, it nonetheless impacts the cell division and cell wall integrity of *L.*

monocytogenes by a yet unknown mechanism.

DISCUSSION

Over the past years, several determinants contributing to the intrinsic lysozyme resistance of

L. monocytogenes have been described (Boneca et al., 2007, Aubry et al., 2011, Burke et al.,

2014, Durack et al., 2015). One of these is a predicted ABC transporter encoded as part of the

eslABCR operon (Burke et al., 2014, Durack et al., 2015). In this study, we aimed to further

investigate the role of the ABC transporter EslABC in *L. monocytogenes* by constructing and

characterizing an *eslB* deletion strain, which also in our hands showed a significant increase in

lysozyme sensitivity. We further found that the PG isolated from the *eslB* mutant was slightly

more crosslinked and also the fraction of deacetylated GlcNAc residues was slightly increased

as compared to the PG isolated from the wildtype strain 10403S. Deacetylation of GlcNAc

residues in PG is achieved by the N-deacetylase PgdA and has been shown to lead to increased

lysozyme resistance (Boneca et al., 2007). Since we saw a slight increase in the deacetylation

of GlcNAc residues in the *eslB* mutant strain, our results indicate that the lysozyme sensitivity

phenotype of the *eslB* deletion strain is independent of PgdA and that this enzyme functions properly in the mutant strain. Lysozyme also acts as a cationic antimicrobial peptide (CAMP), which permeabilizes the bacterial membrane (Herbert et al., 2007, Ibrahim et al., 2002). Cells lacking EslB might therefore be more susceptible to this non-enzymatic activity of lysozyme. Indeed, a previous study showed that disruption of *eslB* affects the resistance to CAMPs (Burke et al., 2014). Another possible explanation for the increased lysozyme sensitivity of the *eslB* deletion strain could be an impaired cell wall integrity, perhaps specifically at the cell division site. Absence of EslB leads to the formation of elongated cells, however, it is currently not clear how the function of EslABC is linked to cell division of *L. monocytogenes*. EslABC could potentially affect the localization of cell division-specific proteins and deletion of *eslB* could lead to a delayed assembly of the divisome, which could lead to an altered PG biosynthesis at the division site and an impaired cell integrity. Indeed, cells of the *eslB* mutant lysed more rapidly as compared to the *L. monocytogenes* wildtype strain 10403S when shifted from BHI broth to Tris-buffer. The autolysis of cells lacking EslB was strongly induced following growth in BHI supplemented with 0.5 M sucrose prior to the incubation in Tris-buffer. These results indicate that the *eslB* mutant is sensitive to an osmotic downshift and we hypothesize that this is due to an impaired cell integrity.

Growth comparisons in different media also revealed that the absence of EslB results in a reduced growth in BHI broth containing high concentrations of mono- or disaccharides. One could speculate that the EslABC transporter might be a sugar transporter with a broad sugar spectrum. The genomic locus of *eslB* supports this hypothesis as the *eslABC* genes are co-transcribed with *eslR* coding for a RpiR family transcriptional regulator (Toledo-Arana et al., 2009). Members of the RpiR transcriptional regulators family are usually involved in the control of sugar phosphate metabolic pathways (Sorensen and Hove-Jensen, 1996, Jaeger and Mayer, 2008, Poulsen et al., 1999, Kim et al., 1997, Yamamoto et al., 2001). The RpiR regulator MurR regulates the MurNAc etherase MurQ in *Escherichia coli* that is essential for the catabolism of MurNAc, a component of the bacterial cell wall (Jaeger and Mayer, 2008). *murQ* is transcribed in an operon with *murP*, coding for a MurNAc-specific EIIBC of the phosphotransferase system (PTS), which is responsible for the import of MurNAc from the growth medium (Dahl et al., 2004). This links the import of sugars by a PTS transporter, whose activity is regulated by the RpiR family transcription factor MurR, with the cell wall biosynthesis in *E. coli*. Accordingly, the transport of sugars by EslABC could therefore impact processes involved in cell wall biosynthesis and the maintenance of cell integrity. Future studies will aim to investigate this further.

A reduction in lysozyme resistance is often associated with reduced virulence. An *E. faecalis*
strain with a deletion in the gene coding for the peptidoglycan deacetylase PgdA, showed a
reduced ability to kill *Galleria mellonella* (Benachour et al., 2012). Similarly, a *S. pneumoniae*
pgdA mutant showed a decreased virulence in a mouse model of infection (Vollmer and
Tomasz, 2000). In our study, we found that inactivation of EslB does not affect the intracellular
growth of *L. monocytogenes* in primary mouse macrophages or the ability to kill *Drosophila*
melanogaster. These observations are consistent with a previous report that another component
of the EslABC transporter, EslA, is dispensable for the ability of *L. monocytogenes* to spread
from cell to cell (Durack et al., 2015). Previously, it was also shown that combined inactivation
of PgdA and OatA reduced the ability of *L. monocytogenes* to grow in bone-marrow derived
macrophages, whereas inactivation of PgdA alone had no impact on the virulence of *L.*
monocytogenes (Rae et al., 2011). We therefore reason that the reduction in lysozyme
resistance upon deletion of *eslB* is not strong enough to affect the ability of *L. monocytogenes*
to grow and survive in primary macrophages and flies.

Taken together, we could show that EslB is not only important for the resistance towards
lysozyme, its absence also affects the autolysis, cell division and ability of *L. monocytogenes*
to grow in media containing high concentrations of sugars. We hypothesize that the function
of the ABC transporter EslABC is directly or indirectly linked to processes involved in the cell
wall biosynthesis and maintenance of cell integrity in *L. monocytogenes*.

468
469

ACKNOWLEDGEMENTS
We thank Ivan Andrew and Jaspreet Haywood from the CSC Genomics Laboratory,
Hammersmith Hospital, for their help with the whole genome sequencing and Annika Gillis
for help with the genome sequence analysis. We would also like to thank Charlotte S. C.
Michaux and Sophie Helaine for the bone marrow-derived macrophages. This work was
funded by the Wellcome Trust grant 210671/Z/18/Z and MRC grant MR/P011071/1 to AG,
the German research foundation (DFG) grant RI 2920/1-1 to JR, and the Wellcome Trust grant
207467/Z/17/Z and MRC grant MR/R00997X/1 to MSD.

FIGURES AND FIGURE LEGENDS

Figure 1: Schematic representation of the *L. monocytogenes eslABCR* operon and cellular location of the encoded proteins. (A) Genomic arrangement of the *eslABCR* operon in *L. monocytogenes*. Arrowheads indicate the orientation of the genes. Small black arrow indicates the promoter identified in a previous study (Toledo-Arana et al., 2009). (B) Schematic of the ABC transporter composed of the NBD protein EslA, which hydrolyses ATP, and the TMD protein EslB, and the cytoplasmic RpiR family transcription regulator EslR. The membrane topology of EslC, a protein of unknown function, was predicted using the TMHMM server 2.0 (Sonnhammer et al., 1998). The *eslB* gene and EslB protein, which were investigated as part of the study, are highlighted in pink.

Figure 2: Impact of *eslB* deletion on resistance towards cell wall-targeting antimicrobials.

(A-C) Minimal inhibitory concentration (MIC) of *L. monocytogenes* strains 10403S (wt),

10403SΔ*eslB(1)* and 10403SΔ*eslB(1)* compl. towards (A) lysozyme, (B) penicillin G and (C)

moenomycin. Strain 10403SΔ*eslB(1)* compl. was grown in the presence of 1 mM IPTG. The

results of four independent experiments are shown. For statistical analysis, a one-way ANOVA

followed by a Dunnett's multiple comparisons test was used (** $p \leq 0.01$).

Figure 3: Deletion of *eslB* leads to slight changes in the peptidoglycan structure. (A) HPLC analysis of muropeptides derived from mutanolysin digested peptidoglycan isolated from strains 10403S (wt), 10403SΔ*eslB(1)* and 10403SΔ*eslB(1)* compl. (the latter grown in the presence of 1 mM IPTG). The muropeptide spectrum of the wildtype strain 10403S has been previously published (Rismondo et al., 2019) and was used again as part of this study as all strains were analyzed at the same time. Major muropeptide peaks are numbered 1-6 according to previously published HPLC spectra (Burke et al., 2014, Rismondo et al., 2015), with numbers shown in red likely corresponding to muropeptides with N-deacetylated GlcNAc residues and peaks 1-2 corresponding to monomeric and 4-6 to dimeric (crosslinked) muropeptide fragments. (B) Quantification of the relative abundance of muropeptide peaks 1-6 for peptidoglycan isolated of strains 10403S (wt), 10403SΔ*eslB(1)* and 10403SΔ*eslB(1)* compl.. For quantification, the sum of the peak areas was set to 100% and the area of individual peaks was determined. Averages values and standard deviations were calculated from three independent peptidoglycan extractions and plotted. For statistical analysis, a two-way ANOVA followed by a Dunnett's multiple comparisons test was used (** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$).

Figure 4: An *L. monocytogenes eslB* deletion strain shows increased endogenous and lysozyme-induced autolysis. Autolysis assays were performed with *L. monocytogenes* strains 10403S (wt), 10403SΔ*eslB(1)* and 10403SΔ*eslB(1)* compl.. Bacteria were grown for 4 h in BHI medium (containing 1 mM IPTG for 10403SΔ*eslB(1)* compl.) and subsequently bacterial suspensions prepared in (A) 50 mM Tris HCl pH 8, (B) 50 mM Tris HCl pH 8 containing 25 µg/ml penicillin, or (C) 50 mM Tris HCl pH 8 with 2.5 µg/ml lysozyme. Cell lysis was followed by taking OD_{600} readings every 15 min. The initial OD_{600} reading for each bacterial suspension was set to 100% and subsequent readings are shown as % of the initial OD_{600} reading. The average % OD_{600} values and standard deviations were calculated from three independent experiments and plotted.

Figure 5: Addition of sucrose but not NaCl negatively impacts the growth of the *L.*

***monocytogenes eslB* mutant strain**. (A-C). Bacterial growth curves. *L. monocytogenes* strains

10403S (wt), 10403S$\Delta eslB_{(1)}$ and 10403S$\Delta eslB_{(1)}$ compl. were grown in (A) BHI broth, (B)

BHI broth containing 0.5 M sucrose or (C) BHI broth containing 0.5 M NaCl. Strain

10403S$\Delta eslB_{(1)}$ compl. was grown in the presence of 1 mM IPTG. OD_{600} readings were

determined at hourly intervals and the average values and standard deviations from three

independent experiments calculated and plotted.

Figure 6: Pre-growth in sucrose containing medium stimulates the endogenous and
penicillin-induced autolysis rate of an *L. monocytogenes eslB* mutant strain. Autolysis
assays were performed with *L. monocytogenes* strains 10403S (wt), 10403S$\Delta eslB_{(1)}$ and
10403S$\Delta eslB_{(1)}$ compl.. Bacteria were grown for 4 h in BHI medium containing 0.5 M sucrose
(containing 1 mM IPTG for 10403S$\Delta eslB_{(1)}$ compl.) and subsequently bacterial suspensions
prepared in (A) 50 mM Tris HCl pH 8, or (B) 50 mM Tris HCl pH 8 containing 25 μg/ml
penicillin. Cell lysis was followed by taking OD_{600} readings every 15 min. The initial OD_{600}
reading for each bacterial suspension was set to 100% and subsequent readings are shown as
% of the initial OD_{600} reading. The average % OD_{600} values and standard deviations were
calculated from three independent experiments and plotted.

Figure 7: The *L. monocytogenes eslB* mutant has a cell division defect and bacteria have

an increased cell length. (A) Microscopy images of *L. monocytogenes* strains 10403S (wt),

10403SΔ*eslB*$_{(1)}$ and 10403SΔ*eslB*$_{(1)}$ compl.. Bacterial membranes were stained with nile red

and cells analyzed by phase contrast and fluorescence microscopy. Scale bar is 2 μm.

Representative images from three independent experiments are shown. (B) Cell length of *L.*

monocytogenes strains 10403S (wt), 10403SΔ*eslB*$_{(1)}$ and 10403SΔ*eslB*$_{(1)}$ compl.. The cell

length of 300 cells per strain was measured and the median cell length calculated. Three

independent experiments were performed, and the average values and standard deviation of the

median cell length plotted. For statistical analysis, a one-way ANOVA analysis followed by a

Dunnett's multiple comparisons test was used (** $p \leq 0.01$).

Figure 8: Impact of the deletion of *eslB* on the virulence of *L. monocytogenes*. (A) Intracellular growth of *L. monocytogenes* strains 10403S (wt), 10403SΔ*eslB*(2) and 10403SΔ*eslB*(2) compl. in mouse bone marrow-derived macrophages (BMMs). For the infection, bacteria were grown overnight at 30°C statically and used on the next day to infect BMMs at an MOI of 2. The number of intracellular bacteria was determined 2, 4, 6 and 8 h post infection as described in the methods section. Three technical repeats were performed per experiment and strain and average values calculated. The average CFU count/ml and standard deviations from three independent experiments were calculated and plotted. (B) Survival curve

581 of flies infected with *L. monocytogenes*. Flies were infected with *L. monocytogenes* strains

582 10403S (wt), 10403SΔ*eslB(1)*, 10403SΔ*eslB(2)* and 10403SΔ*eslB(2)* compl.. Uninjected control

583 flies (U/C) and flies injected with PBS were used as controls. Fly death was monitored daily.

584 For statistical analysis, a one-way ANOVA followed by a Dunnett's multiple comparisons test

585 was used (*** $p \leq 0.001$, **** $p \leq 0.0001$). (C) Bacterial quantification. 16 flies infected with the

586 indicated *L. monocytogenes* strain were collected 24 and 48 h post infection and bacterial load

587 (CFU) determined as described in the methods section. For statistical analysis, a nested one-

588 way ANOVA followed by a Dunnett's multiple comparisons test was used. The observed

589 differences were not statistically significant.

REFERENCES

AUBRY, C., GOULARD, C., NAHORI, M. A., CAYET, N., DECALF, J., SACHSE, M., BONECA, I. G., COSSART, P. & DUSSURGET, O. 2011. OatA, a peptidoglycan O-acetyltransferase involved in *Listeria monocytogenes* immune escape, is critical for virulence. *J Infect Dis,* 204**,** 731-40.

BENACHOUR, A., LADJOUZI, R., LE JEUNE, A., HÉBERT, L., THORPE, S., COURTIN, P., CHAPOT-CHARTIER, M. P., PRAJSNAR, T. K., FOSTER, S. J. & MESNAGE, S. 2012. The lysozyme-induced peptidoglycan N-acetylglucosamine deacetylase PgdA (EF1843) is required for *Enterococcus faecalis* virulence. *J Bacteriol,* 194**,** 6066-73.

BERA, A., BISWAS, R., HERBERT, S., KULAUZOVIC, E., WEIDENMAIER, C., PESCHEL, A. & GÖTZ, F. 2007. Influence of wall teichoic acid on lysozyme resistance in *Staphylococcus aureus. J Bacteriol,* 189**,** 280-3.

BERA, A., HERBERT, S., JAKOB, A., VOLLMER, W. & GÖTZ, F. 2005. Why are pathogenic staphylococci so lysozyme resistant? The peptidoglycan O-acetyltransferase OatA is the major determinant for lysozyme resistance of *Staphylococcus aureus. Mol Microbiol,* 55**,** 778-87.

BERNTSSON, R. P., SMITS, S. H., SCHMITT, L., SLOTBOOM, D. J. & POOLMAN, B. 2010. A structural classification of substrate-binding proteins. *FEBS Lett,* 584**,** 2606-17.

BIERNE, H., SABET, C., PERSONNIC, N. & COSSART, P. 2007. Internalins: a complex family of leucine-rich repeat-containing proteins in *Listeria monocytogenes. Microbes Infect,* 9**,** 1156-66.

BONECA, I. G., DUSSURGET, O., CABANES, D., NAHORI, M. A., SOUSA, S., LECUIT, M., PSYLINAKIS, E., BOURIOTIS, V., HUGOT, J. P., GIOVANNINI, M., COYLE, A., BERTIN, J., NAMANE, A., ROUSSELLE, J. C., CAYET, N., PREVOST, M. C., BALLOY, V., CHIGNARD, M., PHILPOTT, D. J., COSSART, P. & GIRARDIN, S. E. 2007. A critical role for peptidoglycan N-deacetylation in *Listeria* evasion from the host innate immune system. *Proc Natl Acad Sci U S A,* 104**,** 997-1002.

BURKE, T. P., LOUKITCHEVA, A., ZEMANSKY, J., WHEELER, R., BONECA, I. G. & PORTNOY, D. A. 2014. *Listeria monocytogenes* is resistant to lysozyme through the regulation, not the acquisition, of cell wall-modifying enzymes. *J Bacteriol,* 196**,** 3756-67.

CALLEWAERT, L. & MICHIELS, C. W. 2010. Lysozymes in the animal kingdom. *J Biosci,* 35**,** 127-60.

CAMILLI, A., TILNEY, L. G. & PORTNOY, D. A. 1993. Dual roles of *plcA* in *Listeria monocytogenes* pathogenesis. *Mol Microbiol,* 8**,** 143-57.

CARROLL, S. A., HAIN, T., TECHNOW, U., DARJI, A., PASHALIDIS, P., JOSEPH, S. W. & CHAKRABORTY, T. 2003. Identification and characterization of a peptidoglycan hydrolase, MurA, of *Listeria monocytogenes*, a muramidase needed for cell separation. *J Bacteriol,* 185**,** 6801-8.

CORRIGAN, R. M., ABBOTT, J. C., BURHENNE, H., KAEVER, V. & GRÜNDLING, A. 2011. c-di-AMP is a new second messenger in *Staphylococcus aureus* with a role in controlling cell size and envelope stress. *PLoS Pathog,* 7**,** e1002217.

DAFFRE, S., KYLSTEN, P., SAMAKOVLIS, C. & HULTMARK, D. 1994. The lysozyme locus in Drosophila melanogaster: an expanded gene family adapted for expression in the digestive tract. *Mol Gen Genet,* 242**,** 152-62.

DAHL, U., JAEGER, T., NGUYEN, B. T., SATTLER, J. M. & MAYER, C. 2004. Identification of a phosphotransferase system of *Escherichia coli* required for growth on N-acetylmuramic acid. *J Bacteriol,* 186**,** 2385-92.

DAVIDSON, A. L., DASSA, E., ORELLE, C. & CHEN, J. 2008. Structure, function, and evolution of
bacterial ATP-binding cassette systems. *Microbiol Mol Biol Rev,* 72**,** 317-64, table of
contents.
DAWSON, R. J. & LOCHER, K. P. 2006. Structure of a bacterial multidrug ABC transporter.
Nature, 443**,** 180-5.
DE JONGE, B. L., CHANG, Y. S., GAGE, D. & TOMASZ, A. 1992. Peptidoglycan composition of a
highly methicillin-resistant *Staphylococcus aureus* strain. The role of penicillin binding
protein 2A. *J Biol Chem,* 267**,** 11248-54.
DURACK, J., BURKE, T. P. & PORTNOY, D. A. 2015. A prl mutation in SecY suppresses secretion
and virulence defects of *Listeria monocytogenes secA2* mutants. *J Bacteriol,* 197**,** 932-
42.
GAILLARD, J. L., BERCHE, P., FREHEL, C., GOUIN, E. & COSSART, P. 1991. Entry of *L.
monocytogenes* into cells is mediated by internalin, a repeat protein reminiscent of
surface antigens from gram-positive cocci. *Cell,* 65**,** 1127-41.
HÉBERT, L., COURTIN, P., TORELLI, R., SANGUINETTI, M., CHAPOT-CHARTIER, M. P., AUFFRAY,
Y. & BENACHOUR, A. 2007. *Enterococcus faecalis* constitutes an unusual bacterial
model in lysozyme resistance. *Infect Immun,* 75**,** 5390-8.
HERBERT, S., BERA, A., NERZ, C., KRAUS, D., PESCHEL, A., GOERKE, C., MEEHL, M., CHEUNG, A.
& GOTZ, F. 2007. Molecular basis of resistance to muramidase and cationic
antimicrobial peptide activity of lysozyme in staphylococci. *PLoS Pathog,* 3**,** e102.
HIGGINS, C. F. 1992. ABC transporters: from microorganisms to man. *Annu Rev Cell Biol,* 8**,**
67-113.
HÖLTJE, J. V. 1998. Growth of the stress-bearing and shape-maintaining murein sacculus of
Escherichia coli. *Microbiol Mol Biol Rev,* 62**,** 181-203.
HUBER, G. & NESEMANN, G. 1968. Moenomycin, an inhibitor of cell wall synthesis. *Biochem
Biophys Res Commun,* 30**,** 7-13.
IBRAHIM, H. R., AOKI, T. & PELLEGRINI, A. 2002. Strategies for new antimicrobial proteins and
peptides: lysozyme and aprotinin as model molecules. *Curr Pharm Des,* 8**,** 671-93.
JAEGER, T. & MAYER, C. 2008. The transcriptional factors MurR and catabolite activator
protein regulate N-acetylmuramic acid catabolism in *Escherichia coli*. *J Bacteriol,* 190**,**
6598-608.
KIM, C., SONG, S. & PARK, C. 1997. The D-allose operon of *Escherichia coli* K-12. *J Bacteriol,*
179**,** 7631-7.
KOCH, A. L. 2003. Bacterial wall as target for attack: past, present, and future research. *Clin
Microbiol Rev,* 16**,** 673-87.
LAUER, P., CHOW, M. Y., LOESSNER, M. J., PORTNOY, D. A. & CALENDAR, R. 2002.
Construction, characterization, and use of two *Listeria monocytogenes* site-specific
phage integration vectors. *J Bacteriol,* 184**,** 4177-86.
LE JEUNE, A., TORELLI, R., SANGUINETTI, M., GIARD, J. C., HARTKE, A., AUFFRAY, Y. &
BENACHOUR, A. 2010. The extracytoplasmic function sigma factor SigV plays a key role
in the original model of lysozyme resistance and virulence of *Enterococcus faecalis*.
PLoS One, 5**,** e9658.
LOCHER, K. P. 2016. Mechanistic diversity in ATP-binding cassette (ABC) transporters. *Nat
Struct Mol Biol,* 23**,** 487-93.
MÄCHTEL, R., NARDUCCI, A., GRIFFITH, D. A., CORDES, T. & ORELLE, C. 2019. An integrated
transport mechanism of the maltose ABC importer. *Res Microbiol,* 170**,** 321-337.

PILGRIM, S., KOLB-MAURER, A., GENTSCHEV, I., GOEBEL, W. & KUHN, M. 2003. Deletion of
the gene encoding p60 in *Listeria monocytogenes* leads to abnormal cell division and
loss of actin-based motility. *Infect Immun,* 71**,** 3473-84.
POULSEN, T. S., CHANG, Y. Y. & HOVE-JENSEN, B. 1999. D-Allose catabolism of *Escherichia coli*:
involvement of alsI and regulation of als regulon expression by allose and ribose. *J*
Bacteriol, 181**,** 7126-30.
RAE, C. S., GEISSLER, A., ADAMSON, P. C. & PORTNOY, D. A. 2011. Mutations of the *Listeria*
monocytogenes peptidoglycan N-deacetylase and O-acetylase result in enhanced
lysozyme sensitivity, bacteriolysis, and hyperinduction of innate immune pathways.
Infect Immun, 79**,** 3596-606.
RISMONDO, J., HALBEDEL, S. & GRÜNDLING, A. 2019. Cell Shape and Antibiotic Resistance Are
Maintained by the Activity of Multiple FtsW and RodA Enzymes in *Listeria*
monocytogenes. *MBio,* 10.
RISMONDO, J., MOLLER, L., ALDRIDGE, C., GRAY, J., VOLLMER, W. & HALBEDEL, S. 2015.
Discrete and overlapping functions of peptidoglycan synthases in growth, cell division
and virulence of *Listeria monocytogenes*. *Mol Microbiol,* 95**,** 332-51.
SARKAR, P., YARLAGADDA, V., GHOSH, C. & HALDAR, J. 2017. A review on cell wall synthesis
inhibitors with an emphasis on glycopeptide antibiotics. *Medchemcomm,* 8**,** 516-533.
SAUVAGE, E. & TERRAK, M. 2016. Glycosyltransferases and Transpeptidases/Penicillin-
Binding Proteins: Valuable Targets for New Antibacterials. *Antibiotics (Basel),* 5.
SHARROCK, J., ESTACIO-GOMEZ, A., JACOBSON, J., KIERDORF, K., SOUTHALL, T. D. & DIONNE,
M. S. 2019. *fs(1)h* controls metabolic and immune function and enhances survival via
AKT and FOXO in *Drosophila*. *Dis Model Mech,* 12.
SLOTBOOM, D. J. 2014. Structural and mechanistic insights into prokaryotic energy-coupling
factor transporters. *Nat Rev Microbiol,* 12**,** 79-87.
SONNHAMMER, E. L., VON HEIJNE, G. & KROGH, A. 1998. A hidden Markov model for
predicting transmembrane helices in protein sequences. *Proc Int Conf Intell Syst Mol*
Biol, 6**,** 175-82.
SORENSEN, K. I. & HOVE-JENSEN, B. 1996. Ribose catabolism of *Escherichia coli*:
characterization of the *rpiB* gene encoding ribose phosphate isomerase B and of the
rpiR gene, which is involved in regulation of *rpiB* expression. *J Bacteriol,* 178**,** 1003-11.
STAPELS, D. A. C., HILL, P. W. S., WESTERMANN, A. J., FISHER, R. A., THURSTON, T. L., SALIBA,
A. E., BLOMMESTEIN, I., VOGEL, J. & HELAINE, S. 2018. *Salmonella* persisters
undermine host immune defenses during antibiotic treatment. *Science,* 362**,** 1156-
1160.
TANAKA, K. J., SONG, S., MASON, K. & PINKETT, H. W. 2018. Selective substrate uptake: The
role of ATP-binding cassette (ABC) importers in pathogenesis. *Biochim Biophys Acta*
Biomembr, 1860**,** 868-877.
TIPPER, D. J. & STROMINGER, J. L. 1965. Mechanism of action of penicillins: a proposal based
on their structural similarity to acyl-D-alanyl-D-alanine. *Proc Natl Acad Sci U S A,* 54**,**
1133-41.
TOLEDO-ARANA, A., DUSSURGET, O., NIKITAS, G., SESTO, N., GUET-REVILLET, H., BALESTRINO,
D., LOH, E., GRIPENLAND, J., TIENSUU, T., VAITKEVICIUS, K., BARTHELEMY, M.,
VERGASSOLA, M., NAHORI, M. A., SOUBIGOU, G., REGNAULT, B., COPPEE, J. Y., LECUIT,
M., JOHANSSON, J. & COSSART, P. 2009. The *Listeria* transcriptional landscape from
saprophytism to virulence. *Nature,* 459**,** 950-6.

VAN HEIJENOORT, J. 2001. Formation of the glycan chains in the synthesis of bacterial
peptidoglycan. *Glycobiology,* 11**,** 25R-36R.
VOLLMER, W. & TOMASZ, A. 2000. The *pgdA* gene encodes for a peptidoglycan N-
acetylglucosamine deacetylase in *Streptococcus pneumoniae. J Biol Chem,* 275**,** 20496-
501.
YAMAMOTO, H., SERIZAWA, M., THOMPSON, J. & SEKIGUCHI, J. 2001. Regulation of the *glv*
operon in *Bacillus subtilis*: YfiA (GlvR) is a positive regulator of the operon that is
repressed through CcpA and cre. *J Bacteriol,* 183**,** 5110-21.
740