The parasite *Schistocephalus solidus* secretes proteins with putative host manipulation functions

Chloé Suzanne Berger[1,2,3], Jérôme Laroche[2], Halim Maaroufi[2], Hélène Martin[1,2,4], Kyung-Mee Moon[5], Christian R. Landry[1,2,4,6,7], Leonard J. Foster[5] and Nadia Aubin-Horth[1,2,3]

[1]Département de Biologie, Université Laval, Québec, QC, Canada

[2]Institut de Biologie Intégrative et des Systèmes (IBIS), Université Laval, Québec, QC, Canada

[3]Ressources Aquatiques Québec (RAQ), Institut des sciences de la mer de Rimouski, Québec, Canada

[4]Département de Biochimie, Microbiologie et Bioinformatique, Université Laval, Québec, QC, Canada

[5]Department of Biochemistry & Molecular Biology, Michael Smith Laboratories, University of British Columbia, Vancouver, Canada V6T 1Z4

[6]PROTEO, Le réseau québécois de recherche sur la fonction, la structure et l'ingénierie des protéines, Université Laval, Québec, Canada

[7]Centre de Recherche en Données Massives (CRDM), Université Laval, Québec, Canada

Corresponding author: Nadia Aubin-Horth (Nadia.Aubin-Horth@bio.ulaval.ca)

ABSTRACT

Background: Manipulative parasites are thought to liberate molecules in their external environment acting as manipulation factors with biological functions implicated in their host's physiological and behavioural alterations. These manipulation factors are part of a complex mixture called the secretome. While the secretomes of various parasites have been described, there is very little data for a putative manipulative parasite. It is necessary to study the molecular interaction between a manipulative parasite and its host to better understand how such alterations evolve.

Methods: Here, we used proteomics to characterize the secretome of a model cestode with a complex life cycle based on trophic transmission. We studied *Schistocephalus solidus* during the life stage in which behavioural changes take place in its obligatory intermediate fish host, the threespine stickleback (*Gasterosteus aculeatus*). We produced a novel genome sequence and assembly of *S. solidus* to improve protein coding gene prediction and annotation for this parasite. We then described the whole worm's proteome and its secretome during fish host infection using LC-MS/MS.

Results: A total of 2 290 proteins were detected in the proteome of *S. solidus*, and 30 additional proteins were detected specifically in the secretome. We found that the secretome contains proteases, proteins with neural and immune functions, as well as proteins involved in cell communication. We detected Receptor-type tyrosine-protein phosphatases, which were reported in other parasitic systems to be manipulation

factors. We also detected 12 *S. solidus*-specific proteins in the secretome that may play

important roles in host-parasite interactions.

Conclusions: Our results suggest that *S. solidus* liberates molecules with putative

host manipulation functions in the host and that many of them are species specific.

Keywords: *Schistocephalus solidus*, secretome, proteomics, manipulation factor,

parasite, behaviour

BACKGROUND

Parasites have major impacts on their hosts, including on their morphology (1), physiology (2) and behaviour (3, 4). To induce these complex changes in their hosts, it has been proposed that parasites produce, store and release manipulation factors that interfere with the host physiological and central nervous systems (5–7). These manipulation factors are thought to be part of a complex mixture of molecules called the secretome, which is a key element of parasite-host interactions (6). The secretome of a parasite includes lipids (8), nucleic acids (9) and proteins (10), which are sometimes protected inside extracellular vesicles (11). Using molecular and bioinformatics approaches, the proteomic fraction of secretomes of parasites infecting humans (12) and livestock (13) have been described, both in terms of protein composition and function (14,15) (see Table 1 for a review).

The secretomes that have been examined so far are enriched in peptidases and proteases (12,15), which are known to weaken the host immunity barriers. Other secreted proteins, such as paramyosin in the blood fluke *Schistosoma mansoni*, have been shown to help the parasite to escape the host immune response, while secreted proteins involved in calcium functions have impacts on the host neural activity (12). In the context of behavioural manipulation, the secretome is a logical potential source of manipulation factors. However, the secretome content of a behaviour-manipulating parasite has rarely been investigated, to the point that secretomes are referred to as "the missing link in parasite manipulation" (7). The literature contains several reports

from which it is possible to infer a list of putative manipulation factors, which would

target the neural and the immune systems of the hosts and induce behavioural changes

(Table 1). Our knowledge regarding if and how many proteins with neural and immune

functions can be found in the secretomes of manipulative parasites is very limited, and

is based in many cases on inferred proteins rather than actual detection.

One particularly powerful model to study behavioural manipulation is the cestode

Schistocephalus solidus (16). This parasite exhibits a complex life cycle based on

trophic transmission that includes three hosts: a copepod, the threespine stickleback

(*Gasterosteus aculeatus*, obligatory intermediate fish host) and a fish-eating bird, in

which *S. solidus* reproduces (16,17). *S. solidus* infects the threespine stickleback's

abdominal cavity through the ingestion of a parasitized copepod (18). The

consequences of the infection by *S. solidus* on the threespine stickleback's morphology

(19), physiology (20), immune system (21), and behaviour (22) are well-documented.

For example, sticklebacks infected by *S. solidus* show drastic behavioural changes that

result in a loss of the anti-predator response (16): infected fish are more exploratory

(23), less anxious (24) and bolder in the presence of a predator (25) than non-infected

fish.

Most of these behavioural alterations seen in infected fish appear after several weeks,

when the worm has grown to reach the infective stage within its intermediate host (*i.e.*

larger than 50 mg) (26). The infective stage coincides with the time at which *S. solidus*

is ready to reproduce in the bird (16,27), which also generally corresponds to the

activation of the immune response in the host. In the first phase of infection, the

adaptive immune response is generally not activated in the fish. It is only when the

worm reaches the infective stage that an ineffective up-regulation of the respiratory

burst activity occurs (21). Nevertheless, activation of the immune response through the

production of reactive oxygen species (ROS) by granulocytes has been shown to occur

early during infection, depending on the genotype of the stickleback population (126).

Several studies have suggested that the manipulation of the stickleback's behaviour

increases the worm's transmission rate to its final avian host (16). Yet, the adaptive

value for *S. solidus* of such behavioural changes has never been demonstrated (28),

and it is possible that these behavioural modifications in the fish may solely result from

a side effect of infection (7), such as the effect of the parasite mass burden or of the

activation of the host immune response (24, 109). To demonstrate that behavioural

changes in the host are the result of direct parasitic manipulation, the first step requires

to determine if the parasite can liberate molecules in its external environment, and if

yes, to study their functions in relation with the host's phenotype perturbations.

This host-parasite system was used for the first experimental demonstration of a

complex life cycle of a trophically transmitted parasite in 1790 (as reviewed in (16)). A

rich body of work on this model system during the past 50 years has shown, using

enzymatic assays, that the activity of proteases (29) and transferases (30) are required

for *S. solidus* survival and growth. Furthermore, a partial genome of the worm is

available (31) and extensive transcriptome data has been produced (32,33).

Quantification of the transcriptome dynamics across the life stages has uncovered that

when the worm reaches the infective stage in its fish host, genes involved in neural

pathways and sensory perception have higher expression levels, compared to the

earlier stages in the same host, which are characterized by upregulation of growth-

related pathways (33). Furthermore, vesicles are present inside the *S. solidus*

tegument, as shown through scanning and transmission electron microscopy (34).

Moreover, *S. solidus* excretes (through passive mechanisms) or secretes (through

active mechanisms) molecules, including (uncharacterized) proteins, and these

secretions are sufficient to affect its fish host behaviour (35). Finally, a well-annotated

genome of the threespine stickleback host is also available (36), which is important to

adequately differentiate proteins coming from the host and the parasite. The

Schistocephalus-stickleback system is thus appropriate to test the presence of

manipulation factors. However, the nature of the protein content of the *S. solidus*

secretome has never been explored (7, 35). Analyzing the proteomic fraction of the

secretome of *S. solidus,* and its potential enrichment in manipulation factors involved in

neuronal and immune functions, will help us to understand if the behavioural changes of

the host could be induced by parasitic manipulation through the secretome.

Here, we characterized the worm's whole-body proteome and the protein fraction of the

secretome of *S. solidus* using mass spectrometry (LC-MS/MS) (37). We focused on

individuals in the infective stage of development so that the secretome may include

manipulation factors that could be associated with the fish host's behavioural changes.

In helminth parasites, a portion of proteins from the proteome are passively released in

the external environment as metabolic waste products, contributing to an important

fraction of their secretome content (38). Therefore, the secretome is generally a subset

of the proteome in terms of protein content (39). We thus expected that the secretome

of *S. solidus* would also mainly be a subset of its proteome. However, in the context of

behavioural manipulation, we hypothesized that proteins could also be actively liberated

by the parasite in its external environment (38), so that they would be enriched in the

secretome compared to the proteome. Based on what has been described in previous

parasitic systems (references reviewed in Table 1), if *S. solidus* manipulates stickleback

behaviour with its secretions as we hypothesize, then its secretome would include

proteases, as well as proteins with neural and immune functions. Because the worm is

not in direct contact with the vertebrate host brain (16), we also expected to detect

proteins involved in cell communication, cell-cell signaling or transport functions. These

proteins would mediate the communication of the worm with its host brain to induce

potential neural and immune changes, and ultimately behavioural alterations.

As the genome of *S. solidus* is used during LC-MS/MS as a reference database, a more

thorough annotation than the one publicly available (31) of the genome could allow us to

detect more proteins, including potential manipulation factors. We therefore used a

multi-pronged approach combining genomics with proteomics (described in Figure 1).

We first sequenced the genome of *S. solidus* using a combination of long and short

reads to combine longer contigs and improved annotation. Then, we investigated the

global proteomic composition of the proteome and secretome of *S. solidus* using LC-

MS/MS.

Figure 1. The multi-pronged approach designed to detect potential manipulation factors in the secretome of *Schistocephalus solidus*. On the left, the proteomic approach based on LC-MS/MS aims at describing the global proteomic composition of the proteome and the secretome of *S. solidus*. On the right, the genomic approach aims at producing a novel genome sequence and assembly of *S. solidus* to improve protein coding gene prediction and annotation for this parasite. The genome, whose quality is improved, can be used as a new reference database to infer proteins in the proteome and the secretome of *S. solidus*. Combined together, these approaches allow to characterize the secretome of *S. solidus* at the infective stage, including the uncovering of proteins detected only in that fraction and not in its proteome, thus representing potential manipulation factors.

METHODS

Genome sequencing of *Schistocephalus solidus*

<u>Worm collection</u>

We caught threespine sticklebacks from Lac Témiscouata (Québec, 47°40′33″N 68°50′15″O) where fish are known to be infected by *S. solidus* (24), using minnow traps in July 2015. Fish were brought to the "Laboratoire Aquatique de Recherche en Sciences Environnementales et Médicales" at Université Laval (Québec, Canada) and were maintained under a 12h:12h Light:Dark cycle and a water temperature of 15°C. Fish were fed daily with a mixture of blood worms and Artemia. After 10 months, one fish (out of 40) exhibited the morphological changes typically induced by *S. solidus* (19) and was consequently sacrificed to collect the worm. The fish was euthanized with an overdose of MS-222 (75 mg/L mg/kg) and dissected to confirm the infection by *S. solidus*. The worm was immediately put in ethanol 90% and stored at 4°C until genome sequencing. All the other fish were later used during a behavioural experiment (24).

<u>DNA extraction and sequencing</u>

Genomic DNA was extracted using DNeasy Blood and Tissue kit (Qiagen Inc., Valencia, CA, USA) from ~20 mg of tissues. After lysis, we added 4 µL of RNase A (10 mg/mL). Elution was done twice in 100 µL of elution buffer. To reach the desired

concentration for the Nanopore library preparation and Illumina sequencing, we

concentrated DNA with a SpeedVac Concentrator (ThermoFisher, Waltham, MA, USA)

for one hour.

Illumina HiseqX sequencing

The DNA libraries for Illumina sequencing were prepared using a Shotgun PCR-free

library preparation (Lucigen) Illumina Library at the McGill University and Genome

Quebec Innovation Center (Montréal, Canada). Sequencing was performed at the

same center on an Illumina HiSeqX instrument, using paired-ends reads (PE150). A

total of 157 475 128 reads were obtained.The estimated genome size is 540 MB (31).

Oxford Nanopore Technnonologies (MinION) sequencing

Library preparation for *Oxford Nanopore* sequencing was done with a PCR-free ligation

sequencing kit SQK-LSK108 (ONT, Oxford, UK). Briefly, approximately 2.5 µg of high

molecular weight DNA was repaired using the NEBNext FFPE Repair Mix (NEB,

Ipswich, Ma, USA) for 15 min at 20°C before purification with Ampure XP beads. The

repaired DNA was end-prepped with the NEBNext Ultra II End Repair/dA-Tailing

Module (NEB, Ipswich, Ma, USA), for 30 min at 20°C and 30 min at 65 °C, according to

the manufacturer's instructions and purified with Ampure XP beads. Adapter mix (ONT,

Oxford, UK) and Blunt/TA Ligation Master Mix (NEB, Ipswich, Ma, USA) were added to

the purified end-prepped DNA and incubated for 15 min at room temperature. Library

was then purified with Ampure XP beads and ABB wash buffer and recovered in Elution

Buffer (ONT, Oxford, UK). Approximately 333 ng of the purified library was loaded onto

a primed R9.4 SpotOn Flow cell (FLO-MIN106), along with RBF (ONT, Oxford, UK) and

Library Loading Beads (ONT, Oxford, UK). Sequencing was performed with a MinION

Mk1B sequencer running for 48 hours and the MinKNOW software (provided by ONT,

Oxford, UK) was used to control the sequencing process. Four libraries were

sequenced using this protocol. Base calling was performed with albacore

(read_fast5_basecaller.py, ONT Albacore Sequencing Pipeline Software v2.3.1) on all

fast5 files. A total of 4 636 932 read sequences resulted from the base calling step for a

total of 14 535 522 365 nucleotides.

<u>Genome assembly and annotation</u>

All fastq ONT files were pooled in one file prior to assembly. ONT sequences were

assembled using Flye v2.4 <u>(40,41)</u>. The final assembly produced 17 882 scaffolds, the

largest scaffold being 919 337 nt, with a N50 of 121 189 nt. The mean coverage across

scaffolds was 20X. A first phase of correction (polishing) was carried out with

nanopolish (v0.10.2) (https://github.com/jts/nanopolish). A total of 3 473 881 changes

were applied. A second correction phase with one Illumina HiSeqX paired-end

sequence library (20X coverage) was also done with Pilon (v1.22) <u>(42)</u>. A total of 4 889

938 changes helped to improve the quality of the assembly sequences. Scaffolds that

had an average coverage of less than 10X coverage and those that were shorter than

500 bp were removed because they were of limited use prior to genome annotation.

These represented less than 1% of the data and all contigs that contained hits with the

transcripts from S. solidus (33) remained after this selection. This left a total of 15 357

scaffolds and 625 207 408 nucleotides. This Whole Genome Shotgun project has been

deposited at DDBJ/ENA/GenBank under the accession WEDG00000000.1 and the

assembly ID ASM1759139v1.

Completeness of the genome assembly

We used a dataset of 24 765 transcripts from *S. solidus* that we previously published

(33) and mapped them on the *de novo* assembly using GMAP (v2019-03-15) (71) as

implemented in the pipeline GAWN v0.3.2 (https://github.com/enormandeau/gawn). An

assembly quality analysis was performed using the Benchmarking Universal Single-

Copy Orthologs (BUSCO) and BUSCO groups were searched in the metazoa database

(43).

Protein-coding gene prediction

To find out the proportion of repeated regions in the genome and to get a masked

assembly prior to running BRAKER2 to predict protein coding genes, we built a

RepeatModeler (1.0.8) database (44) based on the new genome sequence and ran

RepeatMasker (4.0.6) based on that database (45).

We used BRAKER2 (46–51) for protein-coding gene prediction using two approaches.

The first approach was *ab initio* as it was not based on external data to find Open

Reading Frames (ORFs) but only on genes predicted by GeneMark-ES that are

selected for training Augustus (46, 47). The second approach used the alignment (bam

files) of the transcripts from *S. solidus* (33) on the genome. The two sets of ORFs

obtained with the two approaches were merged and duplicate sequences were

removed, which allowed to obtain a final number of predicted ORFs. We used a BLAST

analysis to quantify the percentage of the *S. solidus* transcriptome we previously

published (33) that was found in the genes predicted using Augustus and vice-versa.

Identification of sequences specific to the new assembly

The predicted ORFs were locally aligned using BLAST+ (52,53) against a database of

43 058 protein sequences from *Schistocephalus solidus* obtained from release 230 of

NCBI (March 2019, https://www.ncbi.nlm.nih.gov). From the predicted ORFs, we

selected those that had no blast match or that had no significant match based on the

fact that the length of the alignment was less that 80% of the length of the query or that

had less than 90% of identical nucleotides over the length of the query (54). These

unmatched sequences that were specific to the assembly were used as one of the

reference databases during LC-MS/MS analysis (see below "Protein identification").

Genome annotation

A functional annotation of the predicted ORFs that were obtained with BRAKER2 (46–

51) was performed using Hmmer (version 3.3) (55) against PFAM domain database

(release 32) (56) and using orthology assignment with eggNOG-mapper (Evolutionary

Genealogy of Genes: Non-supervised Orthologous Groups) (version 2.0.1) (57).

Mass spectrometry characterization of the worm proteome and secretome

<u>Worm and secretome collection for mass spectrometry analysis</u>

Sampling of experimental individuals

Whole worms were collected from wild-caught fish acclimatized to laboratory conditions. Juvenile sticklebacks came from Lac Témiscouata (Québec, 47°40'33"N 68°50'15"O), the same lake that was used to collect a worm for genome sequencing. Fish were caught using a seine in August 2016. They were brought to the "Laboratoire Aquatique de Recherche en Sciences Environnementales et Médicales" at Université Laval where they were raised for one year in 80 L tanks under a Light:Dark photoperiod of 12 h:12 h and a temperature of 15°C reflecting their natural environment conditions (Québec, Canada). Fish were fed daily with brine shrimps.

Collection of proteome and secretome samples

In summer of 2017, 51 fish were individually isolated in 2 L tanks. The day following isolation, fish were injected with 100 µL of PBS (Phosphate Buffered Saline, pH 7.4 Life Technologies) in their abdominal cavity in order to sample their fluids to detect infection by *S. solidus* following the method described in (58). This protocol was repeated the next day. If fish were detected as infected, they were euthanized with an overdose of MS-222 (75 mg/L mg/kg) and dissected to confirm the infection by *S. solidus* and to

collect the worm and its secretome. Fish sex, size and mass, and *S. solidus* mass and

number in each fish were noted. We found that 5 fish were infected, each harbouring a

worm whose weight was above 50 mg (worm 1 = 485.3 mg; worm 2 = 504.1 mg; worm

3 = 286.5 mg; worm 4 = 544.5 mg; worm 5 = 220.9 mg). All the worms used to collect

proteome and secretome samples were far above the mass threshold of 50 mg, which

is commonly used to define the parasitic infective stage that coincides with the

appearance of the behavioural changes in the fish host (16, 26, 27). Furthermore,

behavioural changes in fish infected by worms of similar masses were previously

reported in the Temiscouata population (35). Therefore, we considered these worms

and secretome samples to be representative of the manipulative stage of *S. solidus*.

We only selected fish that were infected by a single worm to prevent potential effects of

multiple infections on the proteomic content of the secretome.

The worm secretome was collected according to a protocol adapted from (59). Each

worm was rinsed with 1 mL of PBS (Phosphate Buffered Saline, pH 7.4 Life

Technologies) to remove fish fluids and then immediately put in a 2 mL tube of PBS.

The tube was covered with aluminium foil to protect the worm from light and placed in a

water recipient at the same temperature as the fish tanks (15°C) for 2 hours. The worm

was removed from the tube and a tablet of Complete, Mini Protease Inhibitor (Sigma)

was added to the tube to protect the proteins from the protease activity. The worm

tissues were then snap frozen in liquid nitrogen and stored dried at -80°C for proteomics

analysis. The liquid in which the worm was incubated (PBS + potential secretome

collected) was stored at -20°C. Further experiments were therefore conducted with 5

worms and their respective secretome.

<u>Preparation of worm tissues and secretome for in-gel digestion</u>

Worms were individually washed with 2 mL of PBS (Phosphate Buffered Saline, pH 7.4

Life Technologies) to remove potential remaining contaminants from the fish. They were

then cut into three equal pieces that were individually put in a tube of 700 µL of lysis

buffer (4% (w/v) SDS in 100 mM Tris pH 8 - 10 mM Dithiothreitol DTT). Six sterile

ceramic beads (size 2.8 mm) were added to each lysis tube and the samples were

homogenized for 20 seconds at 6 000 rpm using a Precellys Homogenizer (Bertin

Technologies). Homogenization was repeated three times. Samples were put on ice for

1 min between each run. Samples were then spun at 10 000 x g at 4°C for 10 min. For

each individual worm, the three homogenates obtained were pooled together and

redistributed into equal volumes into 2 tubes. Samples were heated at 95°C for 10 min,

then spun for 10 min at room temperature. The supernatant of each sample was

collected into a new tube and protein concentration was calculated using the extinction

coefficient measured with a Nanodrop 1000 spectrophotometer (A280 nm; Thermo

Scientific). The concentration obtained for each worm lysate was respectively: 24.14

mg/mL; 22.03 mg/mL; 15.66 mg/mL; 28.53 mg/mL; and 11.37 mg/mL. Worm lysates

were kept at -20°C before performing in-gel digestion.

For the secretome, the protein concentration of each of the liquids in which the worms

were incubated ("secretome" samples) was measured using a Pierce Coomassie

(Bradford) Protein Assay Kit (Fisher Scientific). Bovine Serum Albumin (BSA) at 2

mg/mL was used as standard (7-point standard curve ranging from 0 to 25 µg/mL). The

concentrations obtained were: 13.11 µg/mL; 15.09 µg/mL; 9.83 µg/mL; 16.59 µg/mL;

and 7.46 µg/mL. For each secretome sample, 10 µg of proteins were precipitated with

Trichloroacetic Acid (TCA) (60). Precipitated samples were directly used for in gel-

digestion.

<u>In-gel digestion</u>

For each sample, 50 µg of proteins from worm lysate and 10 µg of proteins from

secretome were resolved on a 12% SDS-PAGE with a BenchMark protein ladder

(Invitrogen) and a negative control (SDS-PAGE loading buffer and water). Migration

was performed during 60 min at 175 V for the worm lysates, and during 30 min at 175 V

for secretomes. Coomassie blue G250 (PMID: 15174055) was used for staining

overnight (Additional file 1). Each migration lane was cut into 5 fractions for worm

lysates, and 3 fractions for secretomes. In-gel digestion was performed on these

fractions according to a previously developed protocol (61). Briefly, after cutting the gel

into slices, proteins were reduced using 10 mM Dithiothreitol (DTT) for 45 min at 56°C,

then alkylated with 55 mM iodoacetamide (IAA) for 30 min at room temperature in the

dark. Digestion was performed overnight at 37°C with trypsin (Promega V5113; 0.1-1 µg

of trypsin depending on the gel staining intensity). The next day, peptides were

extracted using an organic solvent (100 μL acetonitrile) and a step-wise protocol (40%

acetonitrile then 100% acetonitrile), and dried (< 50 μL). Following in-gel digestion, a

STAGE-TIP protocol using C18 extraction disks (3M Empore) was performed to desalt

the samples (62,63). Samples were acidified with trifluoroacetic acid (TFA) (pH < 2.5)

before being passed through the tip. At the end of the STAGE-TIP protocol, samples

were completely dried using a Vacufuge Plus concentrator (Eppendorf) and stored at -

20°C until LC-MS/MS analysis.

<u>LC-MS/MS analysis</u>

<u>The peptides were run on analyse using a quadrupole time of flight mass spectrometer</u>

<u>(Impact II; Bruker Daltonics) coupled to an Easy nano LC 1000 HPLC (Thermo Fisher).</u>

<u>More information about the LC-MS/MS analysis can be found in additional information</u>

<u>(Additional file 2).</u>

<u>Protein identification</u>

We searched the detected mass spectra against the *S. solidus* genome using

MaxQuant (version 1.6.1.0) (65). Two searches were independently performed: the first

search used the larger proteome of *S. solidus* (43 058 entries, downloaded June 21,

2018 and updated May 6, 2019) from the Universal Protein Resource release 2018-05

and 2019-03 (UniProt https://www.uniprot.org/) as a reference database, which includes

proteins predicted from the partial genome of *S. solidus* that was then currently

available (31), as well as proteins predicted from the *de novo* transcriptome (32,33). For

the second search, we used the unmatched sequences that we reported to be specific

to our genome assembly (36 140 sequences, see Results) as a reference database.

The search included common contaminants and variable modifications of methionine

oxidation, and N-acetylation of the proteins. The data was filtered for matches passing

1% false discovery rate set by MaxQuant (65). We included the larger proteome of the

threespine stickleback host (29 032 entries, downloaded June 21, 2018) from the

Universal Protein Resource release 2018-05 (UniProt https://www.uniprot.org/) in the

search in order to include proteins originating from the fish host during the MaxQuant

data search. Subsequent analyses were performed with all the proteins inferred using

the larger proteome of *S. solidus* and the unmatched sequences of the new genome as

reference databases.

<u>Data analysis of the proteome</u>

Data analysis was performed with Python custom scripts (version 3.6.4) using Jupyter

notebooks (version 5.4.0). A template of the custom script is available in additional

information (additional file 3). Proteins detected in each worm sample were retrieved

using MaxQuant. We filtered out of the dataset protein IDs that were solely attributed to

the threespine stickleback (Additional file 4), protein IDs with REV coding (reverse hits

for False Discovery Rate filtering) and protein IDs with CON coding (contaminant hits

that were added into the search). In some cases, several protein IDs were found by

MaxQuant for a specific protein because of high sequence similarities between them.

Research on NCBI (https://www.ncbi.nlm.nih.gov/) and Uniprot

(https://www.uniprot.org/) databases demonstrated that the proteins IDs detected for

one protein were probably isoforms with identical functions. These multiple protein IDs

were kept during annotation to obtain exhaustive functional information, but only the first

ID of each protein was kept for enrichment analyses to avoid an over-representation of

specific processes or functions (see below). This final dataset was used to describe the

global composition of the proteome of *S. solidus.*

We performed two distinct enrichment analyses for the proteins detected in at least one

worm sample, and for the proteins detected in all worm samples. For proteins with

several protein IDs, we kept only the first ID in order to prevent enrichment bias during

analysis. Enrichment analysis was performed using the tool Funrich (version 3.1.3) (66).

We constructed a custom reference database using the protein IDs and the Gene

Ontology (GO) annotation (biological process, cellular component, molecular function)

of all the proteins described in the larger proteome of *S. solidus* available on Uniprot

(https://www.uniprot.org/ 43 058 entries). P-values for enrichment were obtained with a

hypergeometric test corrected with the Bonferroni method.

<u>Data analysis of the secretome</u>

Data analysis was performed with Python using the same approach as for the

proteome. We separated proteins into two categories: proteins that were shared

between the proteome and the secretome samples, and those that were found only in a

secretome sample. For the proteins that were shared between the proteome and the

secretome samples, we performed two distinct enrichment analyses: one for the

proteins detected in at least one secretome sample and one for the proteins detected in

all secretome samples. The enrichment analysis was performed using the tool Funrich

(version 3.1.3) (66) using the same approach as for the proteome.

We investigated the annotation of proteins found only in the secretome and not the

proteome using a three-step approach. During the first step, the protein ID obtained with

the MaxQuant analysis was searched in Uniprot to retrieve its information if available

(protein name and sequence, corresponding coding gene, protein function, localization

and/or structural information). During the second step, we inferred the function of the

protein based on sequence homologies using the Basic Local Alignment Search Tool

(BLAST) on NCBI (https://blast.ncbi.nlm.nih.gov/Blast.cgi). We used Blastp (protein-

protein) using the Non-redundant protein sequences (nr) database (August 2018) and

Tblastn (protein-translated nucleotide) using the Nucleotide collection (nr/nt) database

(August 2018). During the last step, when no information was found with the previous

steps or to confirm the information previously found, we used the Pfam database

(version 32.0) (56) to infer the function of the protein based on domain organization

homologies. In some cases, several protein IDs were found by MaxQuant for a specific

protein because of high sequence similarities between these IDs. We found by applying

the previous approach that in all of these cases, the putatively redundant protein IDs

detected for one protein were indeed isoforms with identical functions. These multiple

protein IDs were nevertheless kept during annotation (see below) to obtain exhaustive

functional information.

Annotation of proteins

In order to obtain an exhaustive annotation of proteins, we conducted complementary

approaches based on sequence and structure analyses (as described below) (67-78)

for the proteins detected in the proteome and/or enriched in the secretome for which

few or no annotation was available. These analyses were also performed for the

proteins that were detected using the unmatched sequences specific to our genome

assembly as a reference database, for which limited or no annotation was available.

Protein sequence analysis

We predicted secreted proteins using the SignalP-5.0 (classical secretory proteins with

option "eukaryota", http://www.cbs.dtu.dk/services/SignalP/) and SecretomeP 1.0f (non-

classical pathway secreted proteins with option "mammalian",

http://www.cbs.dtu.dk/services/SignalP-1.1/#submission) servers (67, 68). In addition,

we predicted transmembrane helices in proteins using TMHMM Server v. 2.0

(http://www.cbs.dtu.dk/services/TMHMM-2.0/) (69). The determination of protein

domains was performed using HmmScan (http://hmmer.org/) against profile-HMM

databases such as Pfam, Gene3D and Superfamily (70). We predicted protein

disordered regions using the PrDOS software (http://prdos.hgc.jp/about.html) (71).

Protein structure analysis

To construct 3D models of proteins, we searched their homologs in PDB database

using BlastP, Delta-Blast and HHpred (70). Then, 3D structure models were built by

homology modelling based on their homologous structures (PDB ID: 4xp4_A) using

MODELLER (72). The quality of the models was assessed by Ramachandran plot

analysis through PROCHECK (73). The images were generated with PyMOL software

(http://pymol.org/) (74).

RESULTS

Genome sequencing of *Schistocephalus solidus*

<u>Genome assembly and completeness</u>

By combining short and long reads produced with Oxford Nanopore technology, we

produced a genome assembly corresponding to 4 944 scaffolds. These scaffolds

summed to over 500 Mb (80% of the assembled genome). To test the completeness of

our assembly, we mapped all the available transcripts of *S. solidus* (33) on our *de novo*

assembly. We found that more than 99.6% (24 676/24 765) of transcripts mapped onto

the new assembly. The remaining 89 transcripts that were not found were blasted

(blastn) against the assembly dataset. A total of 46 transcripts had a hit on 28 scaffolds

but these hits were weak regarding the alignment length. This left 43 orphan transcripts

that had no match in the assembly.

We also searched BUSCO groups to investigate the completeness of the genome

assembly using a complementary approach. A total of 649 complete BUSCOS (66%)

were found out of the 978 BUSCO groups in the metazoa database, which is close to

the proportions obtained from the assembly of other cestode genomes (*e.g.* 72.6% in

the genome of *Schistosoma japonicum* (79); 73.2% in the genome of *Schistosoma*

haematobium (80)). The busco duplication rate was 0.8%. Overall, these results

suggest that the coding genome was well-represented in the genome assembly. Finally,

we identified a total of 56% of elements that appeared to be repeated in the new

genome of *S. solidus* (Additional file 5), which is in accordance with previous studies

that demonstrated that the genomes of most flatworms include large numbers of

repetitive elements (81).

Number of putative genes

We first performed gene prediction using an *ab initio* approach, which gave a total of 21

780 ORFs. A second approach based on bam files from the transcript alignments of *S.*

solidus (33) led to the identification of 30 103 ORFs. The two sets were merged and we

found that 9 103 ORFs overlapped between the two sets, with a highly variable percent

identity (ranging from 1% to 100%). Because of this variation in percent identity and in

order to obtain exhaustive information, we decided to keep duplicate sequences for the

rest of the analysis, which left a total of 51 883 ORF sequences identified. When

comparing these predicted genes to the previously published transcriptome of *S.*

solidus, we found that 77% of the transcripts were found in these predicted genes, and

that 66% of the predicted genes were found in the transcriptome.

 Identification of sequences specific to the new assembly

The 51 883 ORF sequences were aligned against a database of 43 058 protein

sequences of *Schistocephalus solidus* obtained from NCBI. We identified 19 853

sequences that had no blast match and 16 287 sequences that had no significant match

(E value < 1e-5). The total unmatched sequences were therefore 36 140. These

unmatched sequences are potential protein coding genes not represented in public

databases of *S. solidus* and were therefore used as one of the reference databases in

the LC-MS/MS analysis.

Annotation of the new expanded *S. solidus* genome

The 51 883 ORF sequences were annotated using domain information (56) and

orthology assignment (57). First, an important fraction of the predicted sequences (35

576 ORFs; 68.6%) did not have a match in any database and therefore functions could

not be assigned to them. We refer to them as putative *S. solidus*-specific genes

(Additional file 6). We found that 16 307 ORFs were successfully assigned a putative

biological function (Additional file 6: 7 796 ORFs had a match on both databases, 4 591

ORFs had a match on the domain database only and 3 920 ORFs had a match on the

orthology database only). Enrichment analyses on the new expanded *S. solidus*

genome were conducted to identify potential class of genes that could be over-

represented, and that could be either associated with *S. solidus* parasitic life style or

with the phenotypic changes in the fish host. Overall, enrichment analyses showed that

the functions were related to environment sensing: 2.5% of ORFs are *rd3* genes known

to be expressed in photoreceptor cells (82); cell division, growth and development: 1.7%

of ORFs encode for proteins involved in the movement of microtubules and 1.2% of

ORFs are atrophin coding genes involved in development (83); as well as cell

physiology: 1% of ORFs encode for PARP proteins involved in various cell physiological

processes (84).

543   Characterization of the proteome

Using mass spectrometry (LC-MS/MS), we detected 2 290 proteins in samples of whole

S. solidus tissues, among which 1 467 proteins were detected in all worm samples

(Additional file 7). Among these 2 290 proteins, 246 proteins were only detected using

the new genome of *S. solidus* as a reference database during the LC-MS/MS analysis,

with 113 proteins detected in all worms. The new genome sequence and annotation

therefore provide a significant additional resource for *S. solidus* functional genomics.

Most of the 2 290 proteins were functionally annotated, with the exception of the 246

proteins previously mentioned, for which functions were further investigated using a 3-

step approach based on sequence, structure and phylogenetic analyses (see below).

Using this approach, we found that the 2 290 proteins of *S. solidus* included 40 proteins

encoded by 27 genes that did not have any sequence or domain similarities with other

known species, and for which functions could not be described (*i.e.* 1.7 % of putative *S.*

solidus-specific proteins in the proteome, compared to 68.6% of putative *S. solidus*-

specific genes in the new genome) (Additional file 8).

Enrichment analyses in terms of biological process, cellular component and molecular

function (GO terms) were performed for all the proteins detected in at least one of the

worm samples, and also for the proteins detected in all worm samples. For proteins with

several protein IDs, we kept only the first ID in order to prevent enrichment bias. We

found that 14.9% of proteins detected in all worm samples were involved in protein

metabolism processes, namely "translation" [GO:0006412] and "protein folding"

[GO:0006457] ($p < 0.001$). These are typically highly expressed proteins, which explains

their over-representation here (81). Also, we found that 10.2% of proteins detected in all

worm samples referred to mechanisms important in the functioning and the regulation of

the cell cycle: "microtubule based process" [GO:0007017], with the microtubules having

major roles in cell division and growth, and "tricarboxylic acid cycle" [GO:0006099], the

latter referring to abundant metabolic enzymes that are involved in respiration by

producing ATP ($p < 0.001$) (Figure 2A). We also found that 14.4% of proteins detected in

all worm samples were proteins with GTP binding [GO:0005525] and GTPase activity

[GO:0003924] ($p < 0.001$) (Figure 2B). Consistent with all the functions described above,

we found that 23.2% of proteins detected in all worm samples were localized in the

cytoplasm ($p < 0.001$).

Figure 2. The proteome and secretome of *S. solidus* include proteins involved in

protein metabolism/cell growth/energy intake. Results that were significant

(p<0.001) for the enrichment analyses performed in terms of biological process (**A.**) and

molecular function (**B.**). In each case, analysis was performed for the proteins detected

in all worm (outer chart) and secretome (inner chart) samples.

The prediction of the functions of the 246 proteins that were detected using the new

genome as a reference database was further investigated with sequence and structure

analyses (Additional file 7). Their functions were in accordance with the previous

enrichment results, as we found for example proteins involved in microtubules (protein

ID: g20896.t1) or with ATPase activity (protein ID: g10622.t1). Additionally, we found

that six proteins were peptidases or proteases with conserved active sites, suggesting

that they are functional (Table 2), and that a protein contained an amidase domain. This

protein "g7530.t1" is a Fatty acid amide hydrolase like (SsFAAH-like). FAAH enzymes

degrade signalling lipids of the endocannabinoid class (94).

Additionally, 141 proteins were assigned both to the worm and to the fish host during

LC-MS/MS analysis (Additional file 9). As it was not possible to directly determine if the

141 proteins were produced by the worm or the fish, we searched if these proteins

could be core proteins, *i.e.* proteins conserved in all eukaryotes and whose functions

are well characterized (85). Using Blastp and Pfam tools, we found that the 141 proteins

had sequence similarities with 248 previously reported eukaryotic core proteins (85)

(mean Expect-value 2.45e-11), and that they were composed of conserved domains.

We therefore concluded that these proteins were expressed both in the worm and in the

fish because of their fundamental functions in the cell but that their peptide similarities

did not allow to determine their origin. Enrichment analysis performed on these 141

proteins confirmed that they were involved in cell physiology and in energy production

as found for the whole proteome.

Table 2. Peptidases detected in the proteome of *S. solidus* using its new expanded genome.

Peptidase name	ID	Signal peptide (amino acid)	Active site
Cathepsin B	g20295.t1	1 to 26	Q131, C137, H306, N326
Aminopeptidase M17	g13261.t1	1 to 30	K280, R355
Aminopeptidase M17	g19803.t1	1 to 17	K284, R359
Aminopeptidase M17	g13265.t1	1 to 19	K317, R392
Peptidase family M49	g12315.t1	No	E438, E439, H443, E495
Peptidase M16B	g29053.t1	No	E71, E141, L267, R368

The name, the identification number from the new genome assembly (ID), the length of the signal peptide (in numbers of amino acids) and the name of the conserved active sites are indicated for each peptidase.

Characterization of the secretome

A total of 1 568 proteins were detected in the secretome samples (Additional file 7). The

numbers ranged between 781 and 1 183 proteins depending on the sample, with 459

proteins detected in all secretome samples. As expected, the secretome of *S. solidus*

was mostly a subset of the worm proteome, both in terms of protein number and

functions, with a few exceptions described below. The whole protein content of the

secretome of a given worm represented up to 59% of the whole protein content of the

proteome of that same worm. In total, 1 538 unique proteins were shared between at

least one secretome sample and one proteome sample, among which 385 proteins

were detected in all secretomes and proteomes studied, and 74 proteins were detected

in all secretomes and in at least one proteome. All the proteins that were shared

between secretome and proteome samples had a functional annotation, except for 36

proteins that were previously detected in the proteome as *S. solidus*-specific and that

were also secreted (2.3% of putative *S. solidus*-specific proteins in the secretome,

compared to 68.6% of putative *S. solidus*-specific genes in the new genome and 1.7%

of putative *S. solidus*-specific proteins in the proteome) (Additional file 8).

We found that the secretome was composed of proteins enriched for biological

processes, cellular components and molecular functions that were similar to the

proteome. For the proteins detected in all secretomes, almost half of the proteins

participated in the regulation of cell division and energy production. These proteins were

enriched in functions such as "microtubule based process" [GO:0007017] involved in

the regulation of cell division, "glycolytic process" [GO:0006096] and "gluconeogenesis"

[GO:0006094], these two processes being important source of energy production

(p<0.001 hypergeometric test corrected with Bonferroni method) (Figure 2A). Proteins

were also enriched in domains with GTP binding [GO:0005525] and GTPase activity

[GO:0003924] (Figure 2B), as previously reported in the proteome (p<0.001).

Furthermore, 37.9% of the proteins detected in all the secretome samples were

predicted to be localized in the cytoplasm (p<0.001), similar to proteins in the proteome.

Notably, we found that a significant number of proteins (6.1% of the proteins detected in

all secretomes p<0.001) were reported to be specifically localized into the extracellular

space, an enrichment that was specific to the secretome.

Proteins unique to the secretome

We found eight proteins that were detected in all secretome samples, but in none of the

proteomes (Table 3). Among them, three proteins had fibronectin type-III domains. The

first (protein ID: A0A0X3PH69) was a "Neogenin", which is a protein involved in neural

development with two fibronectin type-III domains. The second (protein ID:

A0A0X3Q1B7; A0A0X3PKA1; A0A0X3Q8R6) was a Receptor-type tyrosine-protein

phosphatase eta that included three fibronectin type-III domains. The third (protein ID:

A0A0V0JBL5) included two fibronectin type-III domains, but no additional information on

the function of this protein was available. Furthermore, we detected an uncharacterized

protein with a predicted molecular function corresponding to "Neurotransmitter: sodium

symporter activity (NSS)", which is involved in transmembrane transport (protein ID:

A0A183S8K9; A0A0X3PDV9; A0A0X3PNC8; A0A0V0J682; A0A183T7R5), and a

protein potentially acting at the cell membrane as a Phospholipid scramblase (PLSCR)

(protein ID: A0A0X3P711; A0A183SGM7). Lastly, three other uncharacterized proteins

did not have information on function using Uniprot database, Blast tools and protein

domain identification. It seems that these three proteins are specific to *S. solidus*, with

the last protein (protein ID: A0A0X3Q756) having a secretory signal peptide.

 Table 3. Proteins that are excreted/secreted by _S. solidus_, detected in five

Uniprot ID	Information from Uniprot and Blast tools	Information from sequence, structure and phylogeny
A0A0X3PH69	Neogenin	Fibronectin type-III domains
A0A0X3Q1B7;A0A0X3PKA1;A0A0X3Q8R6	Receptor-type tyrosine-protein phosphatase eta	Fibronectin type-III domains
A0A0V0JBL5	Unknown	Fibronectin type-II domains
A0A183S8K9;A0A0X3PDV9;A0A0X3PNC8; A0A0V0J682;A0A183T7R5	Neurotransmitter: sodium symporter (NSS)	Unknown
A0A0X3P711;A0A183SGM7	Phospholipid scramblase	Unknown
A0A0V0J8W2	Unknown	Gene specific to _S. solidus_ (orphan)
A0A0X3P740	Unknown	Gene specific to _S. solidus_ (orphan)
A0A0X3Q756	Unknown	Gene specific to _S. solidus_ (orphan) Signal peptide (secreted)

secretomes and not in proteomes.

Protein IDs were taken from Uniprot. When several protein IDs were assigned to one

protein, these protein IDs corresponded to isoforms with identical functions. For each

protein, functional annotation was first retrieved from searches with Uniprot and Blast

tools. Complementary analyses based on sequence, structure and phylogeny were

used to obtain information.

Twenty-two proteins were detected in at least one of the five secretome samples, but in

none of the proteomes (Table 4). Three proteins were composed of fibronectin type-III

domains. The first (protein ID: A0A0X3NX35) detected in four out of five secretomes

was described as a Receptor-type tyrosine-protein phosphatase H. The second (protein

ID: A0A183TLI3;   A0A0X3PWG6;   A0A0X3PW04;   A0A0V0J316;   A0A0X3PN23;

A0A0X3PJY8), which was detected in three secretomes, was either a Tenascin (an

extracellular matrix glycoprotein) or a Receptor-type tyrosine-protein phosphatase F.

The last protein of this type (protein ID: A0A0V0JA38), also detected in three

secretomes, was, according to the blast result, a collagen-like protein, which is an

important component of cuticle.

Two proteins appeared to be involved in immunity processes. The first (protein ID:

A0A183TPG4;A0A0X3P3D7;A0A0X3PTB8), which was detected in three secretomes,

was uncharacterized and had a cystatin domain which may be involved in immuno-

modulatory   functions.   The   second   (protein   ID: A0A0V0JBV1;   A0A0V0J795;

A0A0X3NVP3; A0A0X3P0M4), which was detected in the secretome of the largest

worm only (worm 4), was according to the blast results an antigen similar to the

diagnostic antigen gp50 commonly used to detect parasitic diseases.

Two   proteins   were   associated   with   transport   functions.   The   first   (protein   ID:

A0A183TIR8; A0A0X3PCX3; A0A183TT84; A0A0X3PT59) was detected in four out of

five   secretomes   and   was   a   sodium/glucose   cotransporter   involved   in   glucose

homeostasis. The second (protein ID: A0A0X3NT74), which was only detected in the

second largest worm (worm 2), was an intraflagellar transport protein required for

ciliogenesis.

One protein (protein ID: A0A183TDP7), detected in four secretomes, had a knottin fold

similar to a domain found in the *Schistosoma* parasitic trematodes, but for which the

function is not characterized. It is a cysteine-rich protein and interestingly, blastP search

against UniProt database revealed that it is only found in platyhelminthes and amino

acid multiple sequence alignment showed that it presents a new motif, C-x(6)-C-x(7)-

CC-x(4)-C-x(9)-C-x(2)-C-x(6)-C-x(5)-CC-x(3)-C-x(4)-C. This type of protein is resistant

to proteases.

Nine proteins had functions which could not be clearly determined by blast in Uniprot

and Non-redundant protein sequences (GenBank) databases. Five other proteins were

only detected using the new genome of *S. solidus* as a database reference. We further

investigated the functions of these last 14 proteins with sequence, structure and

phylogenetic analysis: nine proteins were specific to *S. solidus*. Among them, two

proteins (protein ID: A0A0V0J2U1 - protein ID: A0A0X3PIM2) had secretion signal

peptides and one protein (protein ID: A0A0X3PXG6) had a TMH (Trans-Membrane

Helix) domain found in transmembrane proteins. Furthermore, one protein was

identified as a peptidase M28B (glutamate carboxypeptidase 2) (protein ID: g1854.t1;

g12541.t1) and one protein had fibronectin type-III domains (protein ID: g17644.t1).

DISCUSSION

Understanding the molecular mechanisms behind behavioural changes in a vertebrate

host infected by a non-cerebral parasite is a fascinating challenge. If these changes are

the result of parasite manipulation of its host, one possible mechanism hinges on

molecules secreted by the parasite (*i.e.* manipulation factors) that impact the host

physiological, immunological and central nervous systems, and ultimately the behaviour

of the host. Here, we describe for the first time the protein component of the secretome

of a parasite commonly referred as manipulative, using mass spectrometry (LC-

MS/MS), with the objective to identify potential manipulation factors that could explain

host behavioural changes. As expected, we found that the proteins that are

excreted/secreted by *S. solidus* are mostly a subset of the proteins being expressed in

the whole worm, and that both included proteases. We also found that 30 secretome

proteins were not detected in the proteome and were involved in neural, immune, and

cell communication functions, therefore having the potential to interfere with the host

physiological systems and behaviour. Finally, we highlighted that the secretome of *S.*

solidus included *S. solidus*-specific proteins that could play important roles in the tight

interaction of the parasite with its fish host. All together, these proteins represent

promising candidates to explain physiological and behavioural changes in the

stickleback host.

The secretome of *S. solidus* is mostly a subset of its proteome with similar

functions

The global protein composition of the proteomes and the secretomes of *S. solidus*

highlights functions that are crucial for its parasitic lifestyle and for its interactions with

the external environment. As it is generally reported (39), we found that the proteins that

are excreted/secreted by *S. solidus* (*i.e.* the secretome) are mostly a subset of the

proteins being expressed in the whole worm (*i.e.* the proteome), both in terms of protein

number and function. We found that both the proteome and the secretome of *S. solidus*

at the infective stage (*i.e.* when behavioural changes appear in the fish host) were

enriched in proteins involved in cell division, which was also found in the functional

analysis of the genome we sequenced, as well as energy production. Concerning cell

growth, our results may seem counter intuitive because a previous transcriptome

analysis revealed that the non-infective stage of *S. solidus* shows higher expression

levels of genes involved in growth and cell regulatory functions compared to the

infective stage (33). However, we did not analyse the proteome and the secretome of

worms at the non-infective stage, such that we can only quantify that these processes

are enriched at the infective stage but not their relative importance compared to other

life stages of the parasite. We therefore only speculate that both the non-infective and

the infective stages of *S. solidus* could rely on biological processes involving cell

growth, but production levels would be much higher at the non-infective stage,

considering the fact that growth occurs predominantly in the first 12 weeks after

installation in the fish host (86). Concerning energy production, the results are also

surprising, as the worm was empirically described to use its glycogen reserve (its

primary source of energy), mainly when it reaches its final avian host (87). However,

transcriptomic analyses demonstrated that glycogen metabolism and energy production

in *S. solidus* are complex processes (33). Specifically, 6 steps of the glycolysis cycle are

highly expressed at the infective stage (33). Furthermore, it is important to keep in mind

that, for tissue and secretome samples collection, worms were placed in a solution of

Phosphate Buffered Saline (PBS, pH 7.4) in the dark for 2 hours. While this method is

commonly used to collect parasitic secretomes (59), we cannot dismiss the possibility

that the environmental change (from the body cavity of the stickleback to the PBS tube)

encountered by *S. solidus* at this step could have influenced the levels of proteins

detected in the proteomes and secretomes. One can speculate that this environmental

stress could have triggered cell growth and energy intake in the worm (for example to

counteract other metabolic effects), explaining the discrepancies between the

proteomics data we obtained and the previous transcriptomic work (33). To confirm our

results, future work could replace the PBS solution by a medium that would better mimic

the fish abdominal cavity, for example by a medium already used *in vitro* for breeding

(21). The global analysis of the proteome and of the secretome of *S. solidus* therefore

confirms the importance of energy use for the worm, even before it reaches the bird.

We detected both in the proteome and the secretome of *S. solidus* proteins with GTP

binding and GTPase activity. GTP binding proteins also called G-proteins are known to

regulate a variety of biological processes such as mediating signals by hormones and

light, gene expression, cytoskeletal and microtubule organization, or vesicle trafficking

(88-91). In parasites, GTPases have been demonstrated to have important roles in the

secretion of virulence factors (90,91). For instance, in *Toxoplasma gondii*, which is a

parasite known to induce behavioural changes in rodents (92), Rab GTPases regulate

the secretion of proteins essential to invade host cells, and the modification of their

expression results in aberrant transport of proteins (93).

The use of the new genome of *S. solidus* as a reference database during mass

spectrometry analysis allowed us to detect 6 peptidases or proteases in the proteome of

S. solidus, and one peptidase (peptidase M28B glutamate carboxypeptidase 2)

detected only in the secretome fraction, with proteases and peptidases being typically

described as virulence factors for many parasites (Table 1). In *S. solidus*, proteases and

peptidases may have important roles in weakening the fish immune response at the

infective stage when the worm is ready to pass into its avian host to reproduce (21).

Injecting these proteases and peptidases alone or in combination, into isolated head

kidney samples from non-infected fish as in (21), would be necessary to confirm their

potential role in disrupting the host immunity. Furthermore, this shows that the new

genome presented here is a valuable tool to identify proteins that are critical for the

parasitic lifestyle. We also found an SsFAAH-like enzyme that could potentially degrade

signalling lipids of the endocannabinoid class. Endocannabinoids have previously been

reported as an important player in host-parasite interactions, by promoting the activation

of the immune response in the host (94-95). This enzyme was not detected in the

secretome of *S. solidus.*

The secretome of *S. solidus* also contains proteins not detected in the proteome

 Neuronal and immune functions

Transcriptomic analysis demonstrated that when *S. solidus* reaches the infective stage,

genes involved in neural pathways and sensory perception are expressed by the worm

at higher levels (33). Thus, we expected the secretome at the infective stage of *S.*

solidus to be enriched with proteins involved in such neural functions (as described in

other parasitic systems, reviewed in Table 1). Furthermore, behavioural changes in the

stickleback infected by *S. solidus* generally appear in concomitance with the activation

of the immune response of the fish (21). Therefore, we expected the secretome of *S.*

solidus at the infective stage to include proteins involved in immunity (as reviewed in

Table 1). We found that three proteins were playing potential roles in neural and

immune functions. The first protein, which was detected in four secretomes but in none

of the proteomes, had a knottin fold called UPF0506 composed of cysteines and

generally found in *Schistosoma* parasites. In *Schistosoma*, the function of the proteins

with such knottin fold is not defined. However, peptides with knottin domains (*i.e.*

knottins) were described in venoms from various animals. For venomous animals,

knottins are neurotoxins having high specificity towards receptors in the nervous system

of their prey or aggressor (96). Furthermore, it is a cysteine-rich protein, and in

parasites cysteine-rich proteins play a role in invasion (97) and modulation (98) of the

immune system. Therefore, this protein could be a promising manipulation factor if it

could act as a neurotoxin in the brain of infected sticklebacks. The second protein,

which was detected in three secretomes but in none of the proteomes, had a cystatin

domain which may be involved in immuno-modulatory functions. It was demonstrated in

parasitic nematodes that cystastins are important secreted molecules that help

parasites to evade immunity of the host (99). Nematode cystatins inhibit host proteases

involved in antigen processing and presentation, leading to a reduction of the host

immune response (100). This secreted protein could thereby explain in part why the

immune response is only activated late in infected sticklebacks, which needs to be

further studied. The third protein appeared to be an antigen similar to the diagnostic

antigen gp50. This diagnostic antigen is used to detect parasitic infection, for example

by *Taenia solium* (101). However, the antigen was detected only in the secretome from

the biggest worm of our study and not in its proteome. Sampling over a longer time

frame could alleviate this type of discrepancies between samples. One protein, which

was detected in all secretomes but in none of the proteomes, had a "Neurotransmitter:

sodium symporter activity (NSS)" annotation. In humans, it is a membrane protein

involved in the termination of synaptic transmission and the recycling of

neurotransmitters at the membrane (102). How this membrane protein could be

secreted and act as a manipulation factor in the secretome is unclear. Therefore, we are

cautious in interpreting the role of this protein as a manipulation factor as it could be

solely a waste product from the membrane of the parasite.

Cell communication

S. solidus is located in the abdominal cavity of the threespine stickleback (16). As the

worm is not in direct contact with its host brain, we expected to detect proteins involved

in cell communication or cell-cell signalling. During annotation of the new genome, 2.5%

of genes were related to environment sensing functions. Furthermore, we found that 7

proteins detected only in the secretome fraction were characterized by fibronectin type-

III (FNIII) domains. FNIII domains are widely found in animal proteins and are involved

in cell-cell interactions (103). The first protein with fibronectin type-III domains found in

all secretomes but in none of the proteomes was identified as a Neogenin. In addition to

playing roles in cell-cell adhesion, neogenins are involved in neural development in

humans (104). Three additional proteins that we identified with fibronectin type-III

domains were described as Receptor-type tyrosine-protein phosphatases (type eta,

detected in all secretomes but in none of the proteomes, type H, detected in 4

secretomes but in none of the proteomes, type F, detected in 3 secretomes but in none

of the proteomes). Receptor-type tyrosine-protein phosphatases are involved in cell-cell

communication (105). It was previously shown that a baculovirus secretes a protein

tyrosine phosphatase, which acts on the neural system of its host the silkworm *Bombyx*

mori and enhances its locomotory activity, so that it ultimately increases the virus

dispersal (106) (Table 1). The set of phosphatases identified here are located in the

membrane, such that they are predicted to have different specific biological functions

that the protein tyrosine phosphatase found in the baculovirus. However, their abundant

representation in the secretome is intriguing and the importance of the process of

phosphorylation represents a future avenue of study in the context of behavioural

changes in the fish host, such as increased exploration (23). For the next two proteins

identified with fibronectin type-III domains, the first protein was found in all secretomes

but in none of the proteomes and had no clear function (protein ID: A0A0V0JBL5), as

for the second protein that was detected in 4 secretomes but in none of the proteomes

(protein ID: g17644.t1). The last protein appeared to be a collagen-like protein found in

3 secretomes. In *Caenorhabditis elegans*, the cuticle is an extracellular matrix made of

small collagen☐like proteins (107). Thus, the role of this protein in cell communication is

uncertain and it may rather be a waste product excreted passively from the cuticle of *S.*

solidus. In summary, fibronectin type-III proteins secreted by *S. solidus* appear to be

good candidates for manipulation factors because of their roles in cell-cell signalling, but

also in potential neural functions.

Additionally, we expected to detect proteins with transport functions in the secretome of

S. solidus to mediate the communication between the worm localized in the host

abdominal cavity and the host brain. We detected three proteins involved in transport or

related functions at the membrane, including a protein that was detected in all

secretomes but in none of the proteomes, a Phospholipid scramblase. It is a

transmembrane protein that is known in human to bind to the 5'-promoter region of the

inositol 1,4,5-triphosphate receptor type 1 gene (*IP3R1*) so that it enhances expression

of the receptor (108). Since these proteins detected here could be simply waste

products coming from the membranes of the parasite, we thus have to be careful when

speculating about the roles of these proteins, and functional analyses will be required.

The secretome includes *S. solidus*-specific proteins that could play important

roles in parasite-host interactions

During annotation of the new genome, we found that 68.6% of the predicted genes did

not have any sequences or domain similarities with other known species that would

allow accurate annotation, therefore representing putative *S. solidus*-specific genes.

Part of these genes are effectively translated, as 1.7% of the proteome and 2.3% of the

secretome included putative proteins specific to the worm. Specifically, we detected a

total of 12 proteins only in the secretome fraction for which functions could not be

clearly determined, but that were specific to *S. solidus.* Three proteins were detected to

be secreted signal peptides (protein ID: A0A0X3Q756 - protein ID: A0A0V0J2U1 -

protein ID: A0A0X3PIM2). We found that the proteins that are specifically enriched in

the secretome are more likely to have secreted signal peptides (10% of the proteins

detected only in the secretome fraction had secreted signal peptides, while 0.17% of the

proteins detected in the proteome had such signals). One protein had a TMH (Trans

Membrane Helix) domain found in transmembrane proteins (protein ID: A0A0X3PXG6).

Proteins with TMH domains are used to diagnose parasitic infection as they are highly

specific to the parasite of interest (110,111). Interestingly, analyses conducted with the

transcriptome of *S. solidus* previously suggested that 19% of all the protein coding

genes could be *S. solidus*-specific (32). In this study, producing a novel genome

sequence and assembly of *S. solidus* led us to increase this estimation by three times.

S. solidus-specific secreted proteins represent promising candidates to explain

physiological and behavioural changes in the stickleback host. The threespine

stickleback is an obligatory host of S*. solidus* (112). Because of the potential co-

evolution between the worm and the fish (16), *S. solidus* may have developed highly

specific molecular mechanisms targeting the threespine stickleback physiological

machinery to insure it can establish and grow in this fish only, until it is ready to pass

into its final avian host. The *S. solidus*-specific proteins found here are therefore likely to

be part of this unique set of molecules crucial for the survival of the worm and we can

assign them this ecological annotation (113). It will be very interesting in the future to

produce recombinant *S. solidus*-specific proteins and test their effects on the phenotype

of the threespine stickleback by functional analysis. These proteins could also serve to

obtain structural information by X-ray crystallography or NMR spectroscopy, which

could help us to discover their function. Finally, studies of how these proteins evolve

between *Schistocephalus* populations and between species of *Schistocephalus*

infecting different fish species (123) will shed additional light on the specificity and

potential function of these proteins.

921   **CONCLUSIONS**

The secretome of *S. solidus* appears to be an important component of the molecular

interaction between the parasite and its threespine stickleback host. In accordance with

our predictions, we detected in the secretome of *S. solidus* at the infective stage (*i.e.*

when behavioural changes appear in the stickleback) proteases, proteins involved in

neural and immune functions, as well as proteins involved in cell communication. We

also highlighted in the secretome the presence of *S. solidus*-specific proteins. In the

future, comparative studies could be conducted to validate that the proteins detected in

the secretome of *S. solidus* act as manipulation factors in the behaviour of its fish host:

at the organism level, the analysis of the secretome of the worm at the non-infective

stage will confirm if the putative manipulative proteins reported in this study are

effectively detected solely at the infective stage (or at highest levels) when the

behavioural changes appear in the host. It would also be of special interest to study the

secretome of *S. solidus* at the stage when it is able to reproduce, *i.e.* in the bird host. At

the species level, comparing the secretome of *Schistocephalus solidus* with the

secretome of a close related species *Schistocephalus pungitii* will be of great interest as

S. pungitii does not have known effects on the behaviour of the nine-spine stickleback,

its specific intermediate host species (114). Finally, functional studies testing effects of

presence in non-infected sticklebacks of interesting proteins identified in the secretome,

such as Receptor-type tyrosine-protein phosphatases or Phospholipid scramblase,

would allow to better understand the contribution of these proteins in the behavioural

changes. To conclude, we hope that the genomic and proteomic resources we provide

will help other researchers to investigate general questions on host-parasite interactions

in nature.

DECLARATIONS

Ethics approval and consent to participate

This study was approved by the Comité de Protection des Animaux de l'Université Laval (CPAUL 2014069-3).

Consent for publication

Not applicable

Availability of data and materials

The datasets supporting the conclusions of this article are included within the article and its additional files. Furthermore, the dataset of the new genome assembly is available from the NCBI BioProject with accession PRJNA576252. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD016844.

Competing interests

The authors declare that they have no competing interests.

Funding

Funding for the project was provided by the Fonds de Recherche du Québec – Nature &

Technologies (FRQNT), Projet en Équipe grant to N.A.H. and C.R.L.; a fellowship

(international internship program) from FRQNT obtained through the Ressources

Aquatiques Québec (RAQ) strategic cluster and a Student/PDF Research Grant award

from the Canadian Society of Zoologists (CSZ) to C.S.B. Mass spectrometry

infrastructure used here was supported by the Canada Foundation for Innovation, the

BC Knowledge Development fund and the BC Proteomics Network. Work in L.J.F.'s

group was supported by Genome Canada and Genome British Columbia via the Pan-

Canadian Proteomics Initiative (264PRO).). C.R.L. holds the Canada Research Chair in

Cellular Systems and Synthetic Biology.

Authors' contributions

C.S.B. designed the study with input from N.A.-H., C.R.L., L.J.F. and K-M.M. C.S.B.

performed the experiments to prepare the worms and their secretomes for mass

spectrometry. K-M.M. performed the experiments with the mass spectrophotometer and

the MaxQuant software. H. Martin prepared the worm samples for genome sequencing

and performed the MinION sequencing. J.L. analyzed the genome and prepared the

files used for searches with the MaxQuant software. J.L. and H. Maaroufi assembled

and annotated the new genome and proteins. C.S.B. analyzed the data with input from

N.A.-H., C.R.L. and L.J.F. C.S.B. and N.A-H. wrote the manuscript with input from all

authors. All authors read and approved the final manuscript.

Acknowledgments

We thank Karina Nielsen, Alexandra Sébastien and Tom Clark from the Foster lab for

their advice during the preparation of the samples for mass spectrometry analysis. The

authors thank the personnel of the Laboratoire Aquatique de Recherche en Sciences

Environnementales et Médicales at Université Laval for their help with fish rearing. We

thank Eric Normandeau from Ressources Aquatiques Québec (RAQ) for his help to

identify *S. solidus* specific proteins.

REFERENCES

1. Krist AC. Effect of the digenean parasite *Proterometra macrostoma* on host

morphology in the freshwater snail *Elimia livescens*. Journal of Parasitology.

2000;86(2):262–7.

2. Jones D, Jones G, Rudnicka M, Click A, Reck-Malleczewen V, Iwaya M.

Pseudoparasitism of host *Trichoplusia ni* by *Chelonus* spp. as a new model system

for parasite regulation of host physiology. Journal of Insect Physiology. 1986;32(1):

315–28.

3. Watson DW, Mullens BA, Petersen JJ. Behavioral fever response of *Musca*

domestica (Diptera: Muscidae) to infection by *Entomophthora muscae*

(Zygomycetes: Entomophthorales). Journal of Invertebrate Pathology. 1993;61(1):

10–6.

4. Lafferty KD, Shaw JC. Comparing mechanisms of host manipulation across host

and parasite taxa. Journal of Experimental Biology. 2013;216(1):56–66.

5. Adamo SA. Parasites: evolution's neurobiologists. Journal of Experimental Biology.

2013;216(1):3–10.

6. Biron DG, Loxdale HD. Host-parasite molecular cross-talk during the manipulative

process of a host by its parasite. Journal of Experimental Biology. 2013;216(1):148–

60.

7. Herbison R, Lagrue C, Poulin R. The missing link in parasite manipulation of host

behaviour. Parasites & Vectors. 2018;11(1):222-27.

8.   Coppens I, Vielemeyer O. Insights into unique physiological features of neutral
lipids in *Apicomplexa*: from storage to potential mediation in parasite metabolic
activities. International Journal for Parasitology. 2005;35(1):597–615.

9.   Fromm B, Ovchinnikov V, Høye E, Bernal D, Hackenberg M, Marcilla A. On the
presence and immunoregulatory functions of extracellular microRNAs in the
trematode *Fasciola hepatica*. Parasite Immunology. 2017;39(1):e12399.

10. Grimwood J, Smith JE. *Toxoplasma gondii*: the role of parasite surface and
secreted proteins in host cell invasion. International Journal for Parasitology.
1996;26(1):169–73.

11. Coakley G, Maizels RM, Buck AH. Exosomes and other extracellular vesicles: The
new communicators in parasite infections. Trends in Parasitology. 2015;31(1):477–
89.

12. Knudsen GM, Medzihradszky KF, Lim K-C, Hansell E, McKerrow JH. Proteomic
analysis of *Schistosoma mansoni* cercarial secretions. Molecular & Cellular
Proteomics. 2005;4(12):1862–75.

13. Gourbal BEF, Guillou F, Mitta G, Sibille P, Thèron A, Pointier J-P, et al. Excretory–
secretory products of larval *Fasciola hepatica* investigated using a two-dimensional
proteomic approach. Molecular and Biochemical Parasitology. 2008;161(1):63–6.

14. Ditgen D, Anandarajah EM, Meissner KA, Brattig N, Wrenger C, Liebau E.
Harnessing the helminth secretome for therapeutic immunomodulators. BioMed

Research International. 2014;15(1):964350-63.

15. Garg G, Ranganathan S. Helminth secretome database (HSD): a collection of
helminth excretory/secretory proteins predicted from expressed sequence tags
(ESTs). BMC Genomics. 2012;13(13):1471-2164.

16. Barber I, Scharsack JP. The three-spined stickleback-*Schistocephalus solidus*
system: an experimental model for investigating host-parasite interactions in fish.
Parasitology. 2010;137(3):411–24.

17. Clarke AS. Studies on the life cycle of the pseudophyllidean cestode
Schistocephalus solidus. Proceedings of the Zoological Society of London.
2010;124(1):257–302.

18. Hammerschmidt K, Kurtz J. Schistocephalus solidus: establishment of tapeworms
in sticklebacks--fast food or fast lane? Experimental Parasitology. 2007;116(2):142–
9.

19. Dingemanse NJ, Oosterhof C, Van Der Plas F, Barber I. Variation in stickleback
head morphology associated with parasite infection. Biological Journal of the
Linnean Society. 2009;96(1):759–68.

20. Heins DC, Baker JA, Toups MA, Birden EL. Evolutionary significance of fecundity
reduction in threespine stickleback infected by the diphyllobothriidean cestode
Schistocephalus solidus. Biological Journal of the Linnean Society. 2010;100(1):
835–46.

21. Scharsack JP, Koch K, Hammerschmidt K. Who is in control of the stickleback

immune system: interactions between *Schistocephalus solidus* and its specific vertebrate host. Proceedings of the Royal Society B. 2007;274(1629):3151–8.

22. Barber I, Huntingford FA. The effect of *Schistocephalus solidus* (Cestoda: pseudophyllidea) on the foraging and shoaling behaviour of three-spined sticklebacks, *Gasterosteus aculeatus*. Behaviour. 1995;132(1):1223–40.

23. Barber I. The effect of hunger and cestode parasitism on the shoaling decisions of small freshwater fish. Journal of Fish Biology. 1995;47(1):524–36.

24. Grécias L, Valentin J, Aubin-Horth N. Testing the parasite mass burden effect on alteration of host behaviour in the -stickleback system. Journal of Experimental Biology. 2018;221(1): jeb174748.

25. Giles N. Behavioural effects of the parasite *Schistocephalus solidus* (Cestoda) on an intermediate host, the three-spined stickleback, *Gasterosteus aculeatus* L. Animal Behaviour. 1983;131(1):1192–4.

26. Tierney JF, Crompton DWT. Infectivity of plerocercoids of *Schistocephalus solidus* (Cestoda: Ligulidae) and fecundity of the adults in an experimental definitive host, *Gallus gallus*. The Journal of Parasitology. 1992;78(1):1049-54.

27. Barber I, Svensson PA, Walker P. Behavioural responses to simulated avian predation in female three spined sticklebacks: The effect of experimental *Schistocephalus solidus* infections. Behaviour. 2004;141(1):1425–40.

28. Poulin R. Parasite manipulation of host behavior-Chapter 5: An update and frequently asked questions. Advances in the Study of Behavior. 2010;41(1):151–86.

29. Polzer M, Conradt U. Identification and partial characterization of the proteases

from different developmental stages of *Schistocephalus solidus* (Cestoda:

Pseudophyllidae). International Journal of Parasitolology. 1994;24(7):967–73.

30. Brophy PM, Papadopoulos A, Touraki M, Coles B, Körting W, Barrett J. Purification

of cytosolic glutathione transferases from *Schistocephalus solidus* (plerocercoid):

interaction with anthelmintics and products of lipid peroxidation. Molecular &

Biochemical Parasitolology. 1989;36(2):187–96.

31. International Helminth Genomes Consortium. Comparative genomics of the major

parasitic worms. Nature Genetics. 2019;51(1):163–74.

32. Hébert FO, Grambauer S, Barber I, Landry CR, Aubin-Horth N. Transcriptome

sequences spanning key developmental states as a resource for the study of the

cestode *Schistocephalus solidus*, a threespine stickleback parasite. GigaScience.

2016;5(1):1-19.

33. Hébert FO, Grambauer S, Barber I, Landry CR, Aubin-Horth N. Major host

transitions are modulated through transcriptome-wide reprogramming events in

Schistocephalus solidus, a threespine stickleback parasite. Molecular Ecololgy.

2017;26(4):1118–30.

34. Yoneva A, Scholz T, Kuchta R. Comparative morphology of surface ultrastructure of

diphyllobothriidean tapeworms (Cestoda: Diphyllobothriidea). Invertebrate Biology.

2018;137(1):38–48.

35. Berger CS, Aubin-Horth N. The secretome of a parasite alters its host's behaviour

but does not recapitulate the behavioural response to infection. Proceedings of the

Royal Society B 287, 20200412.

36. Peichel CL, Sullivan ST, Liachko I, White MA. Improvement of the threespine

stickleback genome using a Hi-C-based proximity-guided assembly. Journal of

Heredity. 2017;108(6):693–700.

37. Barrett J, Jefferies JR, Brophy PM. Parasite proteomics. Parasitology Today.

2000;16(9):400–3.

38. Harnett W. Secretory products of helminth parasites as immunomodulators.

Molecular and Biochemical Parasitology. 2014;195(1):130–6.

39. Greenbaum D, Luscombe NM, Jansen R, Qian J, Gerstein M. Interrelating different

types of genomic data, from proteome to secretome: 'oming in on function. Genome

Research. 2001;11(9):1463–8.

40. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads

using repeat graphs. Nature Biotechnology. 2019;37(5):540–6.

41. Lin Y, Yuan J, Kolmogorov M, Shen MW, Pevzner PA. Assembly of long error-

prone reads using de Bruijn graphs. Proceedings of the National Academy of

Sciences. 2016;113(52):E8396-E8405.

42. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: an

integrated tool for comprehensive microbial variant detection and genome assembly

improvement. PLoS One. 2014;9(11):e112963.

43. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO:

assessing genome assembly and annotation completeness with single-copy

orthologs. Bioinformatics. 2015;31(1):3210–2.

44. Smit A, Hubley R. RepeatModeler Open-1.0. Institute of Systems Biology, Seattle

USA. <http://www.repeatmasker.org>.

45. Hancock JM. RepeatMasker. Dictionary of Bioinformatics and Computational

Biology. 2004.

46. Hoff KJ, Lomsadze A, Borodovsky M, Stanke M. Whole-Genome annotation with

BRAKER. Methods in Molecular Biology. 2019;1962(1):65–95.

47. Hoff KJ, Lange S, Lomsadze A, Borodovsky M, Stanke M. BRAKER1:

Unsupervised RNA-Seq-Based genome annotation with GeneMark-ET and

AUGUSTUS. Bioinformatics. 2016;32(5):767–9.

48. Stanke M, Diekhans M, Baertsch R, Haussler D. Using native and syntenically

mapped cDNA alignments to improve *de novo* gene finding. Bioinformatics.

2008;24(5):637–44.

49. Stanke M, Schöffmann O, Morgenstern B, Waack S. Gene prediction in eukaryotes

with a generalized hidden Markov model that uses hints from external sources.

BMC Bioinformatics. 2006;7(1):62-73.

50. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078–9.

51. Barnett DW, Garrison EK, Quinlan AR, Stromberg MP, Marth GT. BamTools: a C

API and toolkit for analyzing and managing BAM files. Bioinformatics.

2011;27(1):1691–2.

52. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search

tool. Journal of Molecular Biology. 1990;215(3):403–10.

53. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al.

BLAST+: architecture and applications. BMC Bioinformatics. 2009;10(1):421-30.

54. Anderson I, Brass A. Searching DNA databases for similarities to DNA sequences:

when is a match significant? Bioinformatics. 1998;14(4):349–56.

55. Eddy SR. Accelerated Profile HMM Searches. PLoS Computational Biology.

2011;7(10):e1002195.

56. Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, et al. The Pfam

protein families database: towards a more sustainable future. Nucleic Acids

Research. 2016;44(1):279–85.

57. Huerta-Cepas J, Forslund K, Coelho LP, Szklarczyk D, Jensen LJ, von Mering C, et

al. Fast Genome-Wide functional annotation through orthology assignment by

eggNOG-Mapper. Molecular Biology and Evolution. 2017;34(8):2115–22.

58. Berger CS, Aubin-Horth N. An eDNA-qPCR assay to detect the presence of the

parasite *Schistocephalus solidus* inside its threespine stickleback host. Journal of

Experimental Biology. 2018;221(1): jeb178137.

59. Cao X, Fu Z, Zhang M, Han Y, Han Q, Lu K, et al. Excretory/secretory proteome of

14-day *schistosomula, Schistosoma japonicum.* Journal of Proteomics.

2016;130(1):221–30.

60. Link AJ, LaBaer J. Trichloroacetic acid (TCA) precipitation of proteins. Cold Spring

Harbor Protocol. 2011;2011(8):993–4.

61. Shevchenko A, Wilm M, Vorm O, Mann M. Mass spectrometric sequencing of

proteins silver-stained polyacrylamide gels. Analytical Chemistry. 1996;68(5):850–8.

62. Rappsilber J, Ishihama Y, Mann M. Stop and go extraction tips for matrix-assisted

laser desorption/ionization, nanoelectrospray, and LC/MS sample pretreatment in

proteomics. Analytical Chemistry. 2003;75(3):663–70.

63. Rappsilber J, Mann M, Ishihama Y. Protocol for micro-purification, enrichment, pre-

fractionation and storage of peptides for proteomics using StageTips. Nature

Protocols. 2007;2(1):1896–906.

64. Gibbs MR, Moon K-M, Chen M, Balakrishnan R, Foster LJ, Fredrick K. Conserved

GTPase LepA (Elongation Factor 4) functions in biogenesis of the 30S subunit of

the 70S ribosome. Proceedings of the National Academy of Sciences.

2017;114(5):980–5.

65. Cox J, Mann M. MaxQuant enables high peptide identification rates, individualized

p.p.b.-range mass accuracies and proteome-wide protein quantification. Nature

Biotechnology. 2008;26(1):1367–72.

66. Pathan M, Keerthikumar S, Ang C-S, Gangoda L, Quek CYJ, Williamson NA, et al.

FunRich: An open access standalone functional enrichment and interaction network

analysis tool. Proteomics. 2015;15(15):2597–601.

67. Almagro Armenteros JJ, Tsirigos KD, Sønderby CK, Petersen TN, Winther O,

Brunak S, et al. SignalP 5.0 improves signal peptide predictions using deep neural

networks. Nature Biotechnology. 2019;37(4):420–3.

68. Bendtsen JD, Jensen LJ, Blom N, Von Heijne G, Brunak S. Feature-based

prediction of non-classical and leaderless protein secretion. Protein Engineering,

Design and Selection. 2004;17(4):349–56.

69. Krogh A, Larsson B, von Heijne G, Sonnhammer EL. Predicting transmembrane

protein topology with a hidden Markov model: application to complete genomes.

Journal of Molecular Biology. 2001;305(3):567–80.

70. Potter SC, Luciani A, Eddy SR, Park Y, Lopez R, Finn RD. HMMER web server:

2018 update. Nucleic Acids Research. 2018;46(1):200–4.

71. Ishida T, Kinoshita K. PrDOS: prediction of disordered protein regions from amino

acid sequence. Nucleic Acids Research. 2007;35(1):W460–4.

72. Eswar N, Webb B, Marti-Renom MA, Madhusudhan MS, Eramian D, Shen M-Y, et

al. Comparative protein structure modeling using MODELLER. Current Protocols in

Protein Science. 2007;2(1):1-30.

73. Laskowski RA, Rullmannn JA, MacArthur MW, Kaptein R, Thornton JM. AQUA and

PROCHECK-NMR: programs for checking the quality of protein structures solved

by NMR. Journal of Biomolecular NMR. 1996;8(4):477–86.

74. DeLano WL, Others. Pymol: An open-source molecular graphics tool. CCP4

Newsletter on protein crystallography. 2002;40(1):82–92.

75. Sievers F, Higgins DG. Clustal Omega for making accurate alignments of many

protein sequences. Protein Science. 2018;27(1):135–45.

76. Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. New

algorithms and methods to estimate maximum-likelihood phylogenies: assessing

the performance of PhyML 3.0. Systematic Biology. 2010;59(3):307–21.

77. Gouet P, Courcelle E, Stuart DI, Métoz F. ESPript: analysis of multiple sequence

alignments in PostScript. Bioinformatics. 1999;15(4):305–8.

78. Rambaut A. FigTree 1.4. 2 software. Institute of Evolutionary Biology, University of

Edinburgh. 2014.

79. Luo F, Yin M, Mo X, Sun C, Wu Q, Zhu B, et al. An improved genome assembly of

the fluke *Schistosoma japonicum*. PLOS Neglected Tropical Diseases.

2019;13(8):e0007612.

80. Stroehlein AJ, Korhonen PK, Chong TM, Lim YL, Chan KG, Webster B, et al. High-

quality *Schistosoma haematobium* genome achieved by single-molecule and long-

range sequencing. Gigascience. 2019;8(9): 1-24.

81. Maule AG, Marks NJ. Parasitic flatworms: Molecular biology, biochemistry,

immunology and physiology. Chapter 1. CABI Publishing. 2006. 448 p.

82. Azadi S, Molday LL, Molday RS. RD3, the protein associated with Leber congenital

amaurosis type 12, is required for guanylate cyclase trafficking in photoreceptor

cells. Proceedings of the National Academy of Sciences. 2010;107(49):21158–63.

83. Shen Y, Peterson AS. Atrophins' emerging roles in development and

neurodegenerative disease. Cellular and Molecular Life Sciences. 2009;66(1):437–

46.

84. Vyas S, Chesarone-Cataldo M, Todorova T, Huang Y-H, Chang P. A systematic

analysis of the PARP protein family identifies new functions critical for cell

physiology. Nature Communication. 2013;4(1):2240-53.

85. Parra G, Bradnam K, Korf I. CEGMA: a pipeline to accurately annotate core genes

in eukaryotic genomes. Bioinformatics. 2007;23(9):1061–7.

86. Barber I, Svensson PA. Effects of experimental *Schistocephalus solidus* infections

on growth, morphology and sexual development of female three-spined

sticklebacks, *Gasterosteus aculeatus*. Parasitology. 2003;126(4):359–67.

87. Hopkins CA. Studies on cestode metabolism. I. Glycogen metabolism in

Schistocephalus solidus in vivo. Journal of Parasitology. 1950;36(4):384–90.

88. Bourne HR, Sanders DA, McCormick F. The GTPase superfamily: a conserved

switch for diverse cell functions. Nature. 1990;348(6297):125–32.

89. Takai Y, Sasaki T, Matozaki T. Small GTP-binding proteins. Physiological Review.

2001;81(1):153–208.

90. Mitra BN, Saito-Nakano Y, Nakada-Tsukui K, Sato D, Nozaki T. Rab11B small

GTPase regulates secretion of cysteine proteases in the enteric protozoan parasite

Entamoeba histolytica. Cellular Microbiology. 2007;9(9):2112–25.

91. Ward GE, Tilney LG, Langsley G. Rab GTPases and the unusual secretory

pathway of plasmodium. Parasitology Today. 1997;13(2):57–62.

92. Vyas A, Kim S-K, Giacomini N, Boothroyd JC, Sapolsky RM. Behavioral changes

induced by *Toxoplasma* infection of rodents are highly specific to aversion of cat

odors. Proceedings of the National Academy of Sciences. 2007;104(15):6442–7.

93. Kremer K, Kamin D, Rittweger E, Wilkes J, Flammer H, Mahler S, et al. An

overexpression screen of *Toxoplasma gondii* Rab-GTPases reveals distinct

transport routes to the micronemes. PLoS Pathogens. 2013;9(3):e1003213.

94. Batugedara HM, Argueta D, Jang JC, Lu D, Macchietto M, Kaur J, et al. Host- and

helminth-derived endocannabinoids that have effects on host immunity are

generated during infection. Infection and Immunity. 2018;86(1): 1-26.

95. Struck TH, Wey-Fabrizius AR, Golombek A, Hering L, Weigert A, Bleidorn C, et al.

Platyzoan paraphyly based on phylogenomic data supports a noncoelomate

ancestry of spiralia. Molecular Biology and Evolution. 2014;31(7):1833–49.

96. Sachkova MY, Slavokhotova AA, Grishin EV, Vassilevski AA. Structure of the

yellow sac spider *Cheiracanthium punctorium* genes provides clues to evolution of

insecticidal two-domain knottin toxins. Insect Molecular Biology. 2014;23(4):527–38.

97. Michon P, Stevens JR, Kaneko O, Adams JH. Evolutionary relationships of

conserved cysteine-rich motifs in adhesive molecules of malaria parasites.

Molecular Biology and Evolution. 2002;19(7):1128–42.

98. Anandarajah EM, Ditgen D, Hansmann J, Erttmann KD, Liebau E, Brattig NW.

SPARC (secreted protein acidic and rich in cysteine) of the intestinal nematode

Strongyloides ratti is involved in mucosa-associated parasite-host interaction.

Molecular and Biochemical Parasitology. 2016;207(1):75–83.

99. Robinson MW, Massie DH, Connolly B. Secretion and processing of a novel multi-

domain cystatin-like protein by intracellular stages of *Trichinella spiralis*. Molecular

and Biochemical Parasitology. 2007;151(1):9–17.

100.Hartmann S, Lucius R. Modulation of host immune responses by nematode

cystatins. International Journal of Parasitology. 2003;33(11):1291–302.

101.Hancock K, Pattabhi S, Greene RM, Yushak ML, Williams F, Khan A, et al.

Characterization and cloning of GP50, a *Taenia solium* antigen diagnostic for

cysticercosis. Molecular and Biochemical Parasitology. 2004;133(1):115–24.

102.Shi L, Quick M, Zhao Y, Weinstein H, Javitch JA. The Mechanism of a

neurotransmitter:sodium symporter—Inward release of Na and substrate is

triggered by substrate in a second binding site. Molecular Cell. 2008;30(1):667–77.

103.Koide A, Bailey CW, Huang X, Koide S. The fibronectin type III domain as a

scaffold for novel binding proteins. Journal of Molecular Biology. 1998;284(4):1141–

51.

104.De Vries M, Cooper HM. Emerging roles for neogenin and its ligands in CNS

development. Journal of Neurochemistry. 2008;106(4):1483–92.

105.Tonks NK. Protein tyrosine phosphatases: from genes, to function, to disease.
Nature reviews Molecular cell biology. 2006;7(11):833-46.

106.Kamita SG, Nagasaka K, Chua JW, Shimada T, Mita K, Kobayashi M, et al. A
baculovirus-encoded protein tyrosine phosphatase gene induces enhanced
locomotory activity in a lepidopteran host. Proceedings of the National Academy of
Sciences. 2005;102(1):2584–9.

107.Johnstone IL. The cuticle of the nematode *Caenorhabditis elegans*: a complex
collagen structure. Bioessays. 1994;16(3):171–8.

108.Zhou Q, Ben-Efraim I, Bigcas J-L, Junqueira D, Wiedmer T, Sims PJ. Phospholipid
scramblase 1 binds to the promoter region of the inositol 1,4,5-triphosphate
receptor type 1 gene to enhance its expression. Journal of Biological Chemistry.
2005;280(41):35062–8.

109.Grécias L. Étude des causes proximales des changements de comportement de
l'épinoche à trois épines (*Gasterosteus aculeatus*) par son parasite
Schistocephalus solidus. Thesis Université Laval 2017; Available from:
https://corpus.ulaval.ca/jspui/handle/20.500.11794/27886

110.Ohya K, Takahara Y, Kuroda E, Koyasu S, Hagiwara S, Sakamoto M, et al.
Chlamydophila felis CF0218 is a novel TMH family protein with potential as a
diagnostic antigen for diagnosis of *C. felis* infection. Clinical and Vaccine
Immunology. 2008;15(10):1606–15.

111.Wu H, Wang C, Jiang C, Xie Y, Liu L, Song Y, et al. Localization and

characterization of two putative TMH family proteins in *Chlamydia psittaci*.

Microbiological Research. 2016;183(1):19–25.

112.Orr TSC, Hopkins CA, Charles GH. Host specificity and rejection of

Schistocephalus solidus. Parasitology. 1969;59(1):683–90.

113.Pavey SA, Bernatchez L, Aubin-Horth N, Landry CR. What is needed for next-

generation ecological and evolutionary genomics? Trends in Ecology & Evolution.

2012;27(12):673–8.

114.Heins DC, Ulinski B, Johnson J, Baker JA. Effect of the cestode macroparasite

Schistocephalus pungitii on the reproductive success of ninespine stickleback,

Pungitius pungitius. Canadian Journal of Zoology. 2004;82(1):1731–7.

115.Cwiklinski K, Dalton JP. Advances in *Fasciola hepatica* research using "omics"

technologies. International Journal for Parasitology. 2018;48(1):321–31.

116.Kim D-W, Yoo WG, Lee M-R, Yang H-W, Kim Y-J, Cho S-H, et al. Transcriptome

sequencing and analysis of the zoonotic parasite *Spirometra erinacei spargana*

(plerocercoids). Parasites & Vectors. 2014;7(1):368-387.

117.Bień J, Sałamatin R, Sulima A, Savijoki K, Conn DB, Näreaho A, et al. Mass

spectrometry analysis of the excretory-secretory (E-S) products of the model

cestode *Hymenolepis diminuta* reveals their immunogenic properties and the

presence of new E-S proteins in cestodes. Acta Parasitologica. 2016;61(1): 429-42.

118.Biron DG, Marché L, Ponton F, Loxdale HD, Galéotti N, Renault L, et al.

Behavioural manipulation in a grasshopper harbouring hairworm: a proteomics

approach. Proceedings of the Royal Society B: Biological Sciences. 2005;272(1):2117–26.

119. Biron DG, Ponton F, Marché L, Galeotti N, Renault L, Demey-Thomas E, et al. "Suicide" of crickets harbouring hairworms: a proteomics investigation. Insect Molecular Biology. 2006;15(6):731–42.

120. Prandovszky E, Gaskell E, Martin H, Dubey JP, Webster JP, McConkey GA. The neurotropic parasite *Toxoplasma gondii* increases dopamine metabolism. PLoS One. 2011;6(9):e23866.

121. De Bekker C, Quevillon LE, Smith PB, Fleming KR, Ghosh D, Patterson AD, et al. Species-specific ant brain manipulation by a specialized fungal parasite. BMC Evolutionary Biology. 2014;14(1):166-178.

122. Adamo SA, Kovalko I, Turnbull KF, Easy RH, Miles CI. The parasitic wasp *Cotesia congregata* uses multiple mechanisms to control host (*Manduca sexta*) behaviour. Journal of Experimental Biology. 2016;219(23):3750–8.

123. Nishimura N, Heins DC, Andersen RO, Barber I, Cresko WA. Distinct lineages of *Schistocephalus* parasites in threespine and ninespine stickleback hosts revealed by DNA sequence analysis. PLoS One. 2011;6(7):e22505.

124. Libersat F. Wasp uses venom cocktail to manipulate the behavior of its cockroach prey. Journal of Comparative Physiology A. 2003;189(7): 497-508.

125. Korgaonkar A, Han C, Lemire AL, Siwanowicz I, Bennouna D, Kopec R et al. A novel family of secreted proteins linked to plant gall development. *BioRxiv*. 2020:

1344   https://doi.org/10.1101/2020.10.28.359562.

1345   126. Weber JN, Steinel NC, Shim KC, Bolnick DI. Recent evolution of extreme cestode

1346   growth suppression by a vertebrate host. Proceedings of the National Academy of

1347   Sciences. 2017;114(25):6575-80.

LARGER TABLES

1349 **Table 1. Proteomic content (directly measured or inferred) of the secretomes of**

1350 **different parasite species.**

1351

Parasitic species	Host species	Manipulation factor	Host system targeted	Reference
78 helminth species: 64 nematodes, 7 trematodes, 7 cestodes (1)	Humans, animals and plants	Peptidases (61% of species)	Immune system	Helminth secretome database (HSD) Garg and Ranganathan 2012 (15)
Blood fluke *Schistosoma mansoni* (1)	Humans	Proteins involved in calcium binding and regulation	Neural system (affects cell signalization)	Knudsen et al., 2005 (12)
Blood fluke *Schistosoma mansoni* (1)	Humans	Paramyosin and SPO-1	Immune system (evasion)	Knudsen et al., 2005 (12)
Blood fluke *Schistosoma mansoni* (1)	Humans	Proteases	Immune system (degradation of skin barriers)	Knudsen et al., 2005 (12)
Liver fluke *Fasciola hepatica* (1)	Molluscs (*e.g. Galba truncatula*)	Proteases and Antioxidative enzymes: Cu/Zn-superoxide dismutase + thioredoxin	Immune system (evasion)	Cwiklinski and Dalton, 2018 (115) Gourbal et al., 2008 (13)
Cestode *Spirometra erinacei* (1)	Snakes *Rhabdophis tigrinus*	Specific proteins with no identified homologs*	Unknown	Kim et al., 2014 (116)
Cestode *Hymenolepis diminuta* (1)	Rats *Rattus rattus*	Antigens	Immune system (evasion and modulation)	Bień et al., 2016 (117)
Baculovirus *Bombyx mori* NPV (2)	Silkworm *Bombyx mori*	Protein tyrosine phosphatase*	Neural system (enhanced locomotory activity)	Kamita et al., 2005 (106)

Hairworm *Spinochordodes tellinii* + *Paragordius tricuspidatus* (2)	Grass-hopper *Meconema thalassinum* + Cricket *Nemobius sylvestris*	Wnt*	Neural system (modifications of monoamine levels)	Biron et al., 2005 (118) Biron et al., 2006 (119)
Protozoan *Toxoplasma gondii* (2)	Rat *Rattus rattus*	Tyrosine hydroxylase*	Neural system (increases dopamine levels)	Prandovszky et al., 2011 (120)
Fungus *Ophiocordyceps unilateralis* (2)	Ant *Camponotus pennsylvanicus* and *Formica dolosa*	Guanobutyric acid (GBA) and sphingosine	Neural system (action not determined)	De Bekker et al., 2014 (121)
Wasp *Cotesia congregata* (2)	Caterpillar *Manduca sexta*	Cytokine*	Immune system (activation that results in feeding reduction)	Adamo et al., 2016 (122)
Wasp *Ampulex compressa* (2)	Cockroach *Periplaneta americana*	Dopamine	Neural system (represses the activity of neurons)	Libersat et al., 2003 (124)
Aphid *Hormaphis cornu* (2)	Witch hazel *Hamamelis virginiana*	BICYCLE protein	Development (induces galls - novel plant 'organs')	Korgaonkar et al., 2020 (125)

1352

1353

1354   (1) Species of human importance (2) Species studied in the context of behavioural

1355   manipulation. (*) stands for proteins inferred from experimental or bioinformatic

1356   evidence, *i.e.* proteins not directly measured in the secretome.

 Table 4. Proteins that are excreted/secreted by *S. solidus*, detected in at least one

 secretome and not in proteomes.

Protein ID	N° of secretomes for which the protein was detected	Information from Uniprot and Blast tools	Information from sequence, structure and phylogeny
A0A0X3NX35	4/5	Receptor-type tyrosine-protein phosphatase H	Fibronectin type-III domains
A0A183TLI3;A0A0X3PWG6;A0A0X3PW04;A0A0V0J316;A0A0X3PN23;A0A0X3PJY8	3/5	-Tenascin? -Receptor-type tyrosine-protein phosphatase F?	Fibronectin type-III domains
A0A0V0JA38	3/5	Collagen-like protein	Fibronectin type-III domains
A0A183TPG4;A0A0X3P3D7;A0A0X3PTB8	3/5	Unknown	Cystatin domain
A0A0V0JBV1;A0A0V0J795;A0A0X3NVP3;A0A0X3P0M4	1/5 (worm 4)	Antigen	Intrinsically disordered
A0A183TIR8;A0A0X3PCX3;A0A183TT84;A0A0X3PT59	4/5	Sodium/glucose cotransporter	Unknown
A0A0X3NT74	1/5 (worm 2)	Intraflagellar transport protein 81 homolog	Unknown
A0A183TDP7	4/5	Protein with a knottin fold	Cysteine-rich protein Specific to Platyhelminthes

A0A0V0J2U1	4/5	Unknown	Gene specific to *S. solidus* (orphan) Signal peptide (secreted)
A0A0X3NRK5	4/5	Unknown	Gene specific to *S. solidus* (orphan)
A0A0X3PMK5;A0A0X3PDL5;A0A0X3PEZ1	4/5	Unknown	Gene specific to *S. solidus* (orphan)
A0A0X3PRL9	4/5	Unknown	Gene specific to *S. solidus* (orphan)
A0A0X3NVW4	3/5	Unknown	Gene specific to *S. solidus* (orphan)
A0A0X3PXG6	3/5	Unknown	Gene specific to *S. solidus* (orphan) TMH (membrane)
A0A0X3PIM2	2/5	Unknown	Gene specific to *S. solidus* (orphan) - Signal peptide (secreted)
A0A0X3PNW2	2/5	Unknown	Gene specific to *S. solidus* (orphan)
A0A183TE24;A0A0X3PRC0	1/5 (worm 4)	Unknown	Gene specific to *S. solidus* (orphan)
g11241.t1	3/5	Unknown	Intrinsically disordered

g1854.t1;g12541.t1	2/5	Unknown	Peptidase M28B (glutamate carboxypeptidase 2)
g17644.t1	4/5	Unknown	Fibronectin type-III domains
g2.t1	3/5	Unknown	Unknown
g6226.t1	1/5	Unknown	Unknown

1359   Protein IDs were taken from Uniprot or from the new genome assembly. When several

1360   protein IDs were assigned to one protein, these protein IDs corresponded to isoforms

1361   with identical functions. For each protein, functional annotation was first retrieved from

1362   searches with Uniprot and Blast tools. Complementary analyses based on sequence,

1363   structure and phylogeny were used to obtain information.

ADDITIONAL INFORMATION

Additional file 1. Visualization on SDS-PAGE gels of the protein content of the proteome and of the secretome for 5 *S. solidus* worms. A**.** Proteome (*i.e.* tissues) extracts from 5 distinct worms. Note that for worm 1 and worm 2, the sample was loaded on gel either in duplicate or triplicate but only one lane was used for LC-MS/MS. **B.** Secretome extracts from 5 distinct worms. On each gel, a BenchMark protein ladder (Invitrogen) and a negative control (SDS-PAGE loading buffer and water) were also loaded.

Additional file 2. Methods information for LC/MS analysis.

Additional file 3. Template custom script for analysis of the proteome and secretome LC-MS/MS data using Python.

Additional file 4. List of the proteins that were solely attributed to the threespine stickleback host, and that were consequently discarded from analysis. All the protein IDs that were specific to the threespine stickleback *Gasterosteus aculeatus* are reported. For a proteome or a secretome sample, a value above 0 indicates protein detection. The Gene Ontology (GO) annotation (biological process, cellular component and molecular function) of each protein ID is reported.

1384

1385 **Additional file 5.** List of the repeated elements that were identified in the new genome

1386 of *S. solidus*. For each class of repeated elements are indicated: the number of

1387 elements, their total length (pb) as well as their proportion (%) in the new genome.

1388

Additional file 6. Annotation results of 16 307 ORFs predicted from the new expanded

Schistocephalus solidus genome. Functional annotation was performed using PFAM

(domain database) and eggNOG (orthology database). For each ORF match in each

database: the accession number, the E-value and the functional description of the

match are indicated. The other ORFs (68.6%) predicted from the new genome did not

match with these two databases, and are considered *S. solidus*-specific.

1395

1396 **Additional file 7.** List of the proteins detected by LC-MS/MS in the proteome and the

1397 secretome of *S. solidus.* All the protein IDs that were detected in each of the proteome

1398 samples and/or secretome samples are reported. For a sample, a value above 0

1399 indicates protein detection. The Gene Ontology (GO) annotation (biological process,

1400 cellular component and molecular function) of each protein ID is reported.

1401

1402 **Additional file 8.** List of the *S. solidus*-specific proteins detected in the proteome and/or

1403 in the secretome of *S. solidus*. For each protein, the ID is reported from Uniprot. If the

1404   protein is "not secreted", then it was only detected in the proteome and not in the

1405   secretome. The Gene Ontology (GO) annotation (biological process, cellular component

1406   and molecular function) of each protein ID is reported.

1407

1408   **Additional file 9.** List of the 141 proteins that were assigned both to *S. solidus* and to

1409   the fish host during LC-MS/MS analysis. Each protein is assigned several protein IDs

1410   from the worm and the fish. The Gene Ontology (GO) annotation (biological process,

1411   cellular component and molecular function) of each protein ID is reported.