Profiling of Human Gut Virome with Oxford Nanopore Technology

Jiabao Cao[1,2,#], Yuqing Zhang[1,2,#], Min Dai[3], Jiayue Xu[1], Liang Chen[1], Faming

Zhang[3,4], Na Zhao[1,*], Jun Wang[1,*]

Author Affiliations:

1. CAS Key Laboratory of Pathogenic Microbiology and Immunology, Institute of

Microbiology, Chinese Academy of Sciences, Beijing, China.

2. University of Chinese Academy of Sciences, Beijing 100049, China.

3. Medical Center for Digestive Diseases, the Second Affiliated Hospital of Nanjing

Medical University, Nanjing 210011, China.

4. Key Lab of Holistic Integrative Enterology, Nanjing Medical University, Nanjing

210011, China.

[#] Jiabao Cao, and Yuqing Zhang contributed equally to this manuscript.

* Correspondence to:

Professor Jun Wang, CAS Key Laboratory of Pathogenic Microbiology and

Immunology, Institute of Microbiology, Chinese Academy of Sciences, No. 1-3

Beichenxi Road, Chaoyang District, Beijing, China; junwang@im.ac.cn; Dr. Na Zhao,

CAS Key Laboratory of Pathogenic Microbiology and Immunology, Institute of

Microbiology, Chinese Academy of Sciences, No. 1-3 Beichenxi Road, Chaoyang

District, Beijing, China; zhaona@im.ac.cn.

Abstract

Human gut virome play critical roles in maintaining gut microbial composition and

functionality, as well as host physiology and immunology. Yet, there are insufficient

amount of studies on this topic mainly due to methodological limitations, including

enrichment of viruses (phages and host viruses) as well as short read-length from

current sequencing technology. Here we developed a full working protocol for

analyzing human gut virome using physical enrichment, reverse transcription and

random amplification, and eventually the state-of-art single-molecule real-time

sequencing (SMRT) platform of Oxford Nanopore Technology (ONT). We

demonstrate that sequencing viral DNA directly, or viral DNA/cDNA after

amplification using ONT achieves much longer reads and provides more information

regarding virome diversity, many of the virome sequences do not have match in

current databases. Moreover, direct DNA sequencing of virome provides first insights

into the epigenetic modifications on phages, where signals of methylations can be

directly detected. Our study demonstrates that progressing sequencing technology and

bioinformatic improvements will bring more knowledge into virome composition,

diversity and potentially their important functions.

Highlights:

1. Virus-like particles were enriched from human stool samples;

2. Viral nucleotides were sequenced with Oxford Nanopore Technology with and

without amplification;

3. Gut virome in humans showed highly individualized composition;

4. Novel sequences and contigs were found to be the majority in the resulted

sequences;

5. Epigenetic modifications were detected directly on virus genomes.

Keywords:

Introduction

The human gut is home to tremendous amount of microbes [1]. They inhabit

different ecological niches in the gut, forming complex interaction networks between

themselves and with the human cells, and the dynamic balance between gut

microbiome and host is required for human health [2-6]. Studies in human cohort and

mouse models, among others, have confirmed that gut microbial communities are

associated with increasing number of some of metabolism diseases and infectious

diseases, providing insights as well as potential targets for future monitoring and

therapies [3, 7-14].

The gut microbiome contains bacteria, archaea, fungi, protozoa, and, lastly but

not leastly virus. The most abundant cellular members of the microbiome are bacteria

and archaea (account for more than 99% of biomass), and have received most

attention in human microbiome studies over the years [15-17]. Yet, the advances in

next-generation sequencing (NGS) technology and bioinformatic tools have also

facilitated the development of human virome studies. Metagenomic analysis suggests

that the gut of healthy humans harbors commensal virus, including phages, DNA

virus and RNA virus [18-22]. Virome (phages and other host viruses) play roles in

intestinal physiology, enteric immune system, host health and disease [23, 24]. The

dynamic balance between the virome and the intestinal immune system is finely

regulated by cytokines secreted by immune cells [20, 25]. For instance, virome

changes in inflammatory bowel disease (IBD) (Crohn's disease and ulcerative colitis)

are disease specific [26]. Phages residing in mucosal surfaces can influence the host

by providing nonhost-derived immunity against bacterial infections [26, 27]. By

inducing interferons (IFNs), commensal virus can protect from gut inflammation

during tissue damage [28, 29]. However, using current short read sequencing

technologies, such as Illumina, can only offer knowledge on gut virome that is both

biased and fragmentary.

Oxford Nanopore Technologies (ONT) as one of the emerging single-molecule

real-time sequencing technology (SMRT) has the advantage of rapid library

preparation, ultra-long reads and real-time data acquisition [30-32]. For virome, ONT

sequencing has the potential to acquire virus genome by producing genome-length

reads that cover all of mutation within a single virus particle. In addition, biological

nanopores are able to discriminate not only the genome but also single base

modifications such as 5-methylation of cytosine (5mC for DNA and m5C for RNA)

and 6-methylation of adenine (6mA for DNA and m6A for RNA) in native DNA/RNA

[32]. Increasing evidence in last years suggests that DNA/RNA methylation can

influence biological function, including regulation of DNA/RNA replication and

repair, and gene expression [33-36]. Recently, *Oliveira* et al. reported that DNA

methyltransterase in *Clostridioides difficile* has able to mediate sporulation, *C.*

difficile disease transmission and pathogenesis [37]. *Xue* et al. reported viral

N^6-methyladenosine could upregulate replication and pathogenesis of human

respiratory syncytial virus [38]. These findings suggest that epigenetic regulation is

also important for the pathogenesis of important pathogens.

To profile the gut virome in healthy adults, including identity as well as potential

epigenetic information in the virome, we developed a protocol combining physical

enrichment, optional reverse transcription and amplification of nucleotides, and

bioinformatic analytical pipelines, and firstly characterized the virome in five healthy

humans using the ONT PromethION platform. We were able to generate long reads

for virome up to tens of kilobases, resulting in many novel contigs that do not have

matches in the available databases, and also for abundant virus we could detect

epigenetic signals. These discoveries are instructive to future investigations into the

genomics, epigenomics and potential function of human gut virome.

Material and Methods

1. Enrichment and purification of virus-like particles (VLPs)

Each frozen faecel samples (approximately 1.5 g) from five individuals who

provided written informed consent were resuspended in 15 ml sterile Phosphate

Buffered Saline and homogenized thoroughly. The suspension was centrifuged at

4,500 rpm for 10 min at the 4 ℃ to remove large food residues (Beckman Coulter

Allegra^{TM} X-22R). Transfer the supernatant to fresh tubes and centrifuged at 4,500

rpm for 10 min at the 4 ℃ again. The suspernatant was filtered through 0.45 μm

PVDF membrane (Millipore) to remove eukaryotic and bacterial cell-sized particles

before ultracentrifugating at 180,000× g for 3 hours at the 4 ℃ (Beckman Coulter

XP-100). The pellets were resuspended in 400 μl sterile Phosphate Buffered Saline

and treated with 8 U of TURBO DNase Ⅰ (Ambion) and 20 U of RNase A (Fermentas)

at 37 ℃ for 30 min. The viral nucleic acids (DNA and RNA) were extracted by using

QIAamp MinElute Virus Spin Kit (Qiagen) following the manufacturer's instructions

and eluted into RNase-free water. [39, 40]

2. Reverse transcription and random amplification

Viral first strand cDNA was synthesized in a 20 μl reaction mixture with 13 μl of

purified vial nucleic acids from each sample and 100 pmol of primer Rrm

(5'-GACCATCTAGCGACCTCCAC - NNNNNN-3'), as previously described

[39-41]. For the double-strand cDNA synthesis, 100 pmol of primer Rrm and Klenow

fragment (3.5 U/μl; Takara) were added. Random amplification was conducted with

8μl of the double-strand cDNA template in a final reaction volume of 200μl, which

contained 4μM primer Rm (5'-GCCGGAGCTCTGCAGAATTC-3'), 90 μM dNTPs

each, 80 μM Mg^{2+}, 10x Buffer and 1 U of KOD-Plus DNA polymerase (Toyobo). The

amplification product was purified by agarose gel electrophoresis and QIAquick Gel

Extraction Kit (Qiagen).

3. PromethION library preparation and sequencing

PromethION library preparation was performed according to the manufacturer's

instructions for the barcoding cDNA/DNA and native DNA (SQK-LSK109 and

EXP-NBD104). When multiplexing, all the samples were pooled together. ONT

MinKNOW software (v.19.10.1) was used to collect raw sequencing data, and Guppy

(v.3.2.4) was used for local base-calling of the raw data after sequencing runs were

completed. The PromethION was run for up to 96 h.

4. ONT sequence analysis and assembly

Qcat (Oxford Nanopore Technology), python command-line tool for

demultiplexing ONT reads from FASTQ files, was used to trim adaptor and barcode

sequences. With genomeSize = 2k and default parameters, trimmed raw reads were

analysis using Canu v1.9 [42] for virome genome de novo assembly, which includes

read correction, read trimming and contig construction.

5. Matching to current database

Raw reads qcat trimmed were analysis using mimimap2 [43] to identify gut

virome composition, which aligned reads to the reference genome in the National

Center for Biotechnology Information (NCBI) virus genome database, including all

known viruse genome sequences. To improve the accuracy of the viral taxonomy, two

following criteria were adapted: (1) the depth of coverage of reference viral

genome >= 5X; (2) the breadth of coverage of the reference viral genome >= 50%.

To assign the taxonomy of assembled contigs by Canu, two approaches and three

databases were applied, including minimap2, blastn, NCBI virus genome database,

The human gut virome database (GVD) and NCBI nucleotide database. The filter

criteria of alignment results of contigs by minimap2 was the same with raw reads,

while contig matched length >= 1000 bp with nucleic similarity >= 98% and e-value

$<= 10^{-5}$ was adapted as the identified criteria of viruses by blastn.

6. Identification and annotation of bacteriophage ORFs

Seeker [44], a new prediction tool via deep learning framework, was used to

identify putative phages from contigs in amplified cDNA/DNA group with default

parameters. According to the multiPhATE pipeline [45], ORFs in phages were

identified by PHANOTATE [46], a tool to annotate phage genomes. Consequently,

amino sequences of ORFs were aligned to Phantome (http://www.phantome.org) and

pVOGs [47] databases by blastp with parameters "percent of identity >= 60, e-value

<= 0.01" and hmm searched to pVOGs by jackhmmer [48] with default parameters,

respectly. ORFs repeated in two or more samples were clusterd by usearch [49] with

percent of indentity >= 99. To validate these ORFs, we mapped our in-house NGS

data using bowtie2 [50] with default parameters. Coverage of ORFs was calculated by

weeSAM (https://github.com/centre-for-virus-research/weeSAM) and only ORFs

with mapped reads >= 10 were counted.

7. Methylation analysis

Tombo v1.5 was used to detect the methylation states of nucleic acids from raw

DNA samples [51]. A log-likelihood threshold of 2.5 was used to call methylation and

the filter cutoff of methylation sites was defined by the estimated fraction of

significantly modified reads >= 0.7 and coverage depth >= 10X. Different

methylation sites of 5-methylcytosine (5mC) and N6-Methyladenine DNA

Modification (6mA) were visualized by Integrative Genomics Viewer (IGV Version

2.5.3) [52] with default parameters and the putative methyltransferase recognition

motifs were identified by MEME (Version 5.0.5) [53] with the following parameters:

"-dna –mod zoops", and webLogo (https://weblogo.berkeley.edu/logo.cgi) was used

to plot the logo of the motifs we identified.

8. Accession number

The sequencing data were deposited at GSA (Genome Sequence Archive) under

BioProject accession no. PRJCA002499. The full protocol is available at

https://github.com/caojiabao/VirPipeline.

Results

1. Virome separation, enrichment and sequencing

Since metagenomic sequencing using fecal DNA usually results in only minor

fraction of virome sequences, and most of reads will be either from bacteria or

archaea, enrichment of viruses are necessary for studying virome in human gut

samples. Thus, we have combined a series of enrichment methods including filtration

and super centrifugation, to enrich for virus-like-particles (VLPs) in the fecal samples

(Figure 1). After VLPs were isolated, additional DNase/RNase treatment were used to

remove any potential free-DNA that were not virus-originated. The left-over

DNA/RNA were quantified, and half of the nucleotides were directly subjected to

ONT DNA library preparation, to profile the DNA virus abundances as well as

methylation; and the other half were first subjected to RNA reverse transcription and

then amplificaiton with short random primers, to have RNA virus sequenced as cDNA,

and improve the chance of low-abundance DNA viruses to be detected in the

sequencing results.

In our study as a primary investigation, we have first profiled five healthy

volunteers' fecal samples with our protocol. Virus-like particle (VLP) fractions of five

individuals were enriched, and raw DNA, as well as enriched cDNA/DNA were

sequenced using ONT PromthION platform. With one flowcell, the ONT PromthION

yielded a total of 8.2 Gb raw data, with a median of 1.7 Gb per sample in amplified

group; and 452 Mb raw data, with a median of 67 Mb per sample in raw DNA group

(Table S1).

2. Virome composition in healthy individuals revealed by ONT sequencing

With sequencing reads from amplified cDNA/DNA results, we have mapped the

ONT reads onto NCBI virus genomes and then analyzed the composition across five

individuals. Consistent with other studies, our result from amplified cDNA/DNA of

virome showed that bacteriophage families were the most frequently detected and

accounted for the majority of the intestinal virome in number. The final catalogue of

bacteriophages included the *Caudovirals* order (families *siphovirdae*, *podoviridae*),

family *Inoviridae* and family *Microviridae*. Meanwhile, the eukaryotic CRESS-DNA

viruses (family *genomoviridae*) was also detected (Figure 2; Table S2). Of special

note was the presence of the RNA plant viruses including family *Virgaviridae* and

Alphaflexiviridae, which showed good agreement with the findings of Shkoporov *et*

al [54]. We observed that individual specificity is probably a feature of the faecal viral

communities (Figure 2), which had been already demonstrated by several previous

studies [55-57]. However, uncultured phage WW-nAnB strain 3 belonging to family

inoviridae was detected to be presented in amplified cDNA/DNA among five

individuals.

To characterize the potential biases regarding virome composition caused by

amplification, we have compared the results from amplified cDNA/DNA and raw

virome DNA. We could not compare raw RNA due to the fact that it is still not yet

possible to multiplex RNA samples on ONT platforms. The relative abundance of

virus each virome was defined using relative proportion of each virus in terms of

breadth of coverage on the assembled genome, similar to the definition of bacterial

abundances in metagenomic studies. As expected, numbers of viruses are higher in

amplified cDNA/DNA results, except for individual 2 and individual 5 who remained

the same in terms of virus diversity. Further, abundances of the common existing

viruses between raw DNA and amplified cDNA/DNA showed essential shifts in all

five individuals, demonstrated detectable bias of reverse transcription as well as

random amplification approach we adapted.

3. Virome assembly using ONT sequences

We next focused on the assembly of nanopore sequencing reads in five

individuals. Assembler Canu was used to assemble the virome sequences separately

from raw DNA and amplified cDNA/DNA groups into contigs, which yielded a total

of 1564 contigs, with a median of 15 and 347 contigs per sample for raw DNA and

amplified cDNA/DNA group. Average length of contigs from raw DNA group was

longer than those from amplified cDNA/DNA group (Figure S1). The contigs vary

largely in length, ranging from 1kb to 53kb (Figure S2). Consequently, we obtained

the identity of certain contigs by matching with NCBI virus genomes, human gut

virus database (GVD) [58] and NCBI nucleotide databases, however all with very low

matching rate, suggesting a large collection of potentially novel genomes in our

results (Table S3). Thus, Seeker was used to identify bacteriophages in amplified

cDNA/DNA group. As a result, more than 50% of contigs per sample were identified

as phages in amplified cDNA/DNA group. To characterize these bacteriophages, we

performed phages ORFs prediction and functional annotation, which yielded average

7 ORFs per contig, ranging from 6 to 10, and average 3 ORFs per kb in length. In

addition, ORFs repeated in two or more samples were validated by mapping to

additional Illumina Hiseq sequencing data of metagenomics from same samples. We

founded that over half ORFs (59.3%, 35/59) can be matched, indicating that these

phages stably exist in our data (Table S4).

To examine the potential of covering full virus genome of long sequence

produced from the oxford nanopore technology sequencing, we mapped raw reads to

the contigs by canu separated from amplified cDNA/DNA and raw DNA. We

aggregated and counted the length of raw reads aligned to the longest contigs of our

choice from these two groups, and calculated the proportion of raw reads to contigs in

length (Figure 3). In amplified VLP cDNA/DNA group, the max value of reads length

in each sample was all more than 15% of contigs in length, the highest achieving 40%

of contigs' full length in individual 4 (Figure S3). In raw DNA group, this proportion

was higher in general , understandably due to the fact random amplification usually

can not reach full length available in the raw DNA. A small number of reads were

longer than the final contigs (Figure 3), for which the possible reason for this result

was that some long reads were trimmed by canu during assembly, due to the lower

quality of part of the sequences being abandoned by Canu.

4. Viral epigenomic detection using ONT

In addition, methylation states of bases in DNA from reads can be detected

directly by the Oxford Nanopore sequencing without extra laboratory techniques. In

this study, we have analyzed methylation signals on a few contigs that reached >10X

coverage in raw DNA data in any of individuals. Methylation detection can not be

carried out on cDNA and amplified DNA samples, for they will lose all the

methylation signals. For the only one of the contigs (contig00000015) with known

identity (Uncultured crAssphage), we detected in total 17 5mC and 120 6mA

methylation sites covered in the 8kb genome (Figure 4). For 5mC and 6mA

methylation, the nucleotide motifs YCHYTTA*C*TWMRECT (e-value = 1.2 x 10^{-2})

and motif MADWDTW*A*NADYYWW (e-value = 2.5 x 10^{-4}) were identified

respectively, with the methylated nucleotide highlighted in bold italic.

We have also found another 4 contigs with >10X coverage, but they do not have

detectable relative in the databases we have searched against. They are potentially

novel viruses and very unlikely to be bacteria or archaea, as our protocol has removed

most of the none VLPs, and bacteria/archaea contigs would most likely have matches

in the databases we have searched. They have 5mC methylation sites ranging from 0.3

to 1.8 (% of genome) and 6mA methylation sites ranging from 0.7 to 2.5 (% of

genome). The motifs for methylation are also extremely diverse, including

NHHYYKG*C*DHNNN, WDWADDW*C*DWYNDDW, MNNNNTR*C*GBNNNND and

HDNBYDV*C*VVVVNNH   for   5mC,   and   WHWHNYD*A*HNNYYHTT,

VNWWDWH*A*YBYNNNT,   DRNVRKK*A*BBNDNNN   and

NRNARND*A*SYAHHNH for 6mA (Figure S4). Since most of the methylation studies

are performed in eukaryotes, and only starting in bacteria with limited information

available, it is yet difficult to compare the methylation profiles to understand its

underlying mechanisms.

Discussion

Our study combined the physical enrichment of VLPs in fecal samples,

nucleotide amplification with the latest sequencing technology to establish a complete

workflow of human gut virome profiling. With longer reads as well as richer

information of additional epigenetic modifications, developments in sequencing

technology could bring another round of revolution in metagenome as well as virome

investigations. More importantly, despite the complex steps before sequencing were

designed for maximum enrichment of VLPs, as well as removal of any DNA/RNA

that were not of virus origin, and thus inevitably makes it relatively time-consuming,

ONT could carry out sequencing and produce reads nearly simultaneously, making it

possible to finish data generation from samples with five days of working time, and

potentially even shorter if PromethION was run for less time, or data were analyzed

during the course of being generated (real-time); comparatively, Illumina based

platforms usually takes longer to generate enough read length for downstream

analyses. There are several studies who have already utilized this property for fast

pathogen detection in infectious diseases [59-61], there are cases of virome analysis

that might require such time efficiency as well, and our protocol provides a feasible

choice for virome studies that also requires short turn-around time.

We also compared the effect of reverse transcription and consequent

amplifications on virome analysis with ONT. Such steps were used for several reasons.

Firstly, despite ONT is capable of sequencing DNA or RNA directly, multiplexing is

still only possible for DNA libraries, and directly sequencing RNA is not yet

cost-effective for virome analysis, while cDNA is a better alternative. Also, to utilize

the capacity of sequencing on ONT, very high concentrations of DNA or RNA

libraries are required to generate enough reads, yet this is also very difficult for

virome DNA/RNA, who usually only reaches 10% of the required input from 1.5

grams of fecal samples. Amplification does lead to higher amount of reads and

enables detection of low abundance viruses to be detected, as revealed in our study;

but it also leads to on average shorter reads and certain biases in the estimations of

virus abundances, resulting from both affinity to random primers as well as PCR

produced artefacts. Lastly, amplified cDNA/DNA loses all the methylation

modifications on the viral nucleotides and prevents investigations into this potentially

vital epigenetic information. Thus, future investigator needed to balance the pros and

cons of amplification processes, and could take advantage of our approach of using

both raw DNA (and/or RNA) and amplified cDNA/DNA for sequencing, gaining

complementary information within the same sequencing run.

The profiles of virome in our studied individuals suggest a highly diverse virome,

with only small amount of "core" viruses shared in between. This core could shrink

even further with increasing number of individuals while the total diversity of viruses

increase, which we plan to investigate in the future. We found phages making the

majority part of the virome in the human gut, plus a few host viruses; while the

mystery of plant RNA viruses is again present with ONT data, many reads achieving >

1kb in length, whether they are left-overs from our plant-based food, or rather human

viruses with their closest relatives in the plant-associated viruses, still call for more

investigations, especially within functional experiments. Our data suggest that there

are potentially high number of unknown, novel viruses in the human gut, as our

assembled contigs have very low rate of matches in the current databases; we consider

those not likely to be contaminations due to our vigorous depletion of any non-VLPs

and non-viral nucleotides, plus the fact that they do not have any match in nucleotide

collection of NCBI either.

It's also the first time we demonstrate that the phage genomes are methylated via

direct sequencing. RSV viruses and influenza viruses are known to have m6A

methylation on their RNA genome [38, 62], detected with more complex methods

with low throughput, while DNA phages (or other DNA viruses) are not yet studied to

our knowledge. In *E. coli* and *C. difficile* it is shown that 6mA is the main form of

DNA methylation, while eukaryotes usually lack this form, and in our results phages

also have 6mA as the main form of DNA methylation [37, 63, 64]. Since DNA

methylations play an important role in bacterial defense against phages, how phage

genome becomes methylated, and the consequent impact on phage life cycle and

interactions with bacterial hosts remain to be explored with dedicated studies. Besides,

it remains possible the motifs of methylation between phage and bacteria are

intrinsically linked, and provide additional information to determine the host range of

phages; this would require increasing the current knowledge on epigenetics of both

bacteria and phages in the future.

Conclusions

To summarize, we developed and pilot-tested a thorough protocol for human gut virome analysis using the lastest ONT sequencing platform, and generated novel insights into the individuality, diversity of gut virome with new sequencing data. Our approach of course can be applied for other virome studies, including animal gut, soil and water virome etc., and accumulating both sequences as well as epigenetic information on those samples, have the long potential of opening up new directions in metagenomic, microbiological and medical researches.

Acknowledgements

This work was supported by the National Key Research and Development Program of China (grant number 2018YFC2000500), the Strategic Priority Research Program of the Chinese Academy of Sciences (grant number XDB29020000), and the National Natural Science Foundation of China (grant number 31771481 and 91857101).

References

[1] Sender R, Fuchs S. Milo R. Are We Really Vastly Outnumbered? Revisiting the Ratio of Bacterial to Host Cells in Humans. Cell 2016;164(3):337-40. https://doi.org/10.1016/j.cell.2016.01.013.

[2] Falony G, Joossens M, Vieira-Silva S, Wang J, Darzi Y, Faust K, et al. Population-level

analysis of gut microbiome variation. Science 2016;352(6285):560-4. https:// doi.org/10.1126/science.aad3503.

[3] Moschen AR, Gerner RR, Wang J, Klepsch V, Adolph TE, Reider SJ, et al. Lipocalin 2 Protects from Inflammation and Tumorigenesis Associated with Gut Microbiota Alterations. Cell Host Microbe 2016;19(4):455-69. http://doi.org/10.1016/j.chom.2016.03.007.

[4] Wang J, Thingholm LB, Skieceviciene J, Rausch P, Kummen M, Hov JR, et al. Genome-wide association analysis identifies variation in vitamin D receptor and other host factors influencing the gut microbiota. Nat Genet 2016;48(11):1396-406. https:// doi.org/10.1038/ng.3695.

[5] Tschurtschenthaler M, Wang J, Fricke C, Fritz TMJ, Niederreiter L, Adolph TE, et al. Type I interferon signalling in the intestinal epithelium affects Paneth cells, microbial ecology and epithelial regeneration. Gut 2014;63(12):1921-31. https:// doi.org/10.1136/gutjnl-2013-305863.

[6] Wang J, Chen L, Zhao N, Xu XZ, Xu YK. Zhu BL. Of genes and microbes: solving the intricacies in host genomes. Protein Cell 2018;9(5):446-61. https:// doi.org/10.1007/s13238-018-0532-9.

[7] Belkaid Y. Hand TW. Role of the Microbiota in Immunity and Inflammation. Cell 2014;157(1):121-41. https://doi.org/10.1016/j.cell.2014.03.011.

[8] Wang ZN, Klipfell E, Bennett BJ, Koeth R, Levison BS, Dugar B, et al. Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. Nature 2011;472(7341):57-U82. https://doi.org/10.1038/nature09922.

[9] Ridaura VK, Faith JJ, Rey FE, Cheng JY, Duncan AE, Kau AL, et al. Gut Microbiota from Twins Discordant for Obesity Modulate Metabolism in Mice. Science 2013;341(6150):1079-U49. https://doi.org/10.1126/science.1241214.

[10] Jostins L, Ripke S, Weersma RK, Duerr RH, McGovern DP, Hui KY, et al. Host-microbe

interactions have shaped the genetic architecture of inflammatory bowel disease. Nature

2012;491(7422):119-24. https://doi.org/10.1038/nature11582.

[11] Giongo A, Gano KA, Crabb DB, Mukherjee N, Novelo LL, Casella G, et al. Toward defining

the   autoimmune   microbiome   for   type   1   diabetes.   Isme   J   2011;5(1):82-91.

https://doi.org/10.1038/ismej.2010.92.

[12] Cox LM. Blaser MJ. Pathways in Microbe-Induced Obesity. Cell Metab 2013;17(6):883-94.

https://doi.org/10.1016/j.cmet.2013.05.004.

[13] Zhu WF, Gregory JC, Org E, Buffa JA, Gupta N, Wang ZN, et al. Gut Microbial Metabolite

TMAO Enhances Platelet Hyperreactivity and Thrombosis Risk. Cell 2016;165(1):111-24.

https://doi.org/10.1016/j.cell.2016.02.011.

[14] Wang ZN, Roberts AB, Buffa JA, Levison BS, Zhu WF, Org E, et al. Non-lethal Inhibition of

Gut Microbial Trimethylamine Production for the Treatment of Atherosclerosis. Cell

2015;163(7):1585-95. https://doi.org/10.1016/j.cell.2015.11.055.

[15] Zarate S, Taboada B, Yocupicio-Monroy M. Arias CF. Human Virome. Arch Med Res

2017;48(8):701-16. http://doi.org/10.1016/j.arcmed.2018.01.005.

[16] Zou SM, Caler L, Colombini-Hatch S, Glynn S. Srinivas P. Research on the human virome:

where are we and what is next. Microbiome 2016;4. http://doi.org/10.1186/s40168-016-0177-y.

[17] Scarpellini E, Ianiro G, Attili F, Bassanelli C, De Santis A. Gasbarrini A. The human gut

microbiota and virome: Potential therapeutic implications. Digest Liver Dis 2015;47(12):1007-12.

http://doi.org/10.1016/j.dld.2015.07.008.

[18] Liu L, Gong T, Tao WY, Lin BL, Li C, Zheng XS, et al. Commensal viruses maintain

intestinal intraepithelial lymphocytes via noncanonical RIG-I signaling. Nat Immunol

2019;20(12):1681-1691. https://doi.org/10.1038/s41590-019-0513-z.

[19] Reyes A, Haynes M, Hanson N, Angly FE, Heath AC, Rohwer F, et al. Viruses in the faecal

microbiota of monozygotic twins and their mothers. Nature 2010;466(7304):334-U81.

https://doi.org/10.1038/nature09199.

[20] Virgin HW. The Virome in Mammalian Physiology and Disease. Cell 2014;157(1):142-50.

http://doi.org/10.1016/j.cell.2014.02.032.

[21] Zhang T, Breitbart M, Lee WH, Run JQ, Wei CL, Soh SWL, et al. RNA viral community in

human feces: Prevalence of plant pathogenic viruses. Plos Biol 2006;4(1):108-18.

https://doi.org/ARTN e310.1371/journal.pbio.0040003.

[22] Shi Y. Mu LB. An expanding stage for commensal microbes in host immune regulation. Cell

Mol Immunol 2017;14(4):339-48. https://doi.org/10.1038/cmi.2016.64.

[23] Handley SA. The virome: a missing component of biological interaction networks in health

and disease. Genome Med 2016;8. http://doi.org/10.1186/s13073-016-0287-y.

[24] Mukhopadhy I, Segal JP, Carding SR, Hart AL. Hold GL. The gut virome: the 'missing link'

between gut bacteria and host immunity? Ther Adv Gastroenter 2019;12.

http://doi.org/10.1177/1756284819836620.

[25] Foca A, Liberto MC, Quirino A, Marascio N, Zicca E. Pavia G. Gut Inflammation and

Immunity: What Is the Role of the Human Gut Virome? Mediat Inflamm 2015.

http://doi.org/10.1155/2015/326032.

[26] Norman JM, Handley SA, Baldridge MT, Droit L, Liu CY, Keller BC, et al. Disease-Specific

Alterations in the Enteric Virome in Inflammatory Bowel Disease. Cell 2015;160(3):447-60.

http://doi.org/10.1016/j.cell.2015.01.002.

[27] Manrique P, Dills M. Young MJ. The Human Gut Phage Community and Its Implications for

Health and Disease. Viruses-Basel 2017;9(6). http://doi.org/10.3390/v9060141.

[28] Yang JY, Kim MS, Kim E, Cheon JH, Lee YS, Kim Y, et al. Enteric Viruses Ameliorate Gut

Inflammation via Toll-like Receptor 3 and Toll-like Receptor 7-Mediated Interferon-beta

Production. Immunity 2016;44(4):889-900. https://doi.org/10.1016/j.immuni.2016.03.009.

[29] Broggi A, Tan Y, Granucci F. Zanoni I. IFN-lambda suppresses intestinal inflammation by

non-translational regulation of neutrophil function. Nat Immunol 2017;18(10):1084-1093.

https://doi.org/10.1038/ni.3821.

[30] Deamer D, Akeson M. Branton D. Three decades of nanopore sequencing. Nat Biotechnol

2016;34(5):518-24. https://doi.org/10.1038/nbt.3423.

[31] Laszlo AH, Derrington IM, Ross BC, Brinkerhoff H, Adey A, Nova IC, et al. Decoding long

nanopore sequencing reads of natural DNA. Nat Biotechnol 2014;32(8):829-33.

https://doi.org/10.1038/nbt.2950.

[32] Schatz MC. Nanopore sequencing meets epigenetics. Nat Methods 2017;14(4):347-8.

https://doi.org/10.1038/nmeth.4240.

[33] Low DA, Weyand NJ. Mahan MJ. Roles of DNA adenine methylation in regulating bacterial

gene expression and virulence. Infect Immun 2001;69(12):7197-204. https://doi.org/Doi

10.1128/Iai.69.12.7197-7204.2001.

[34] Casadesus J. Low D. Epigenetic gene regulation in the bacterial world. Microbiol Mol Biol R

2006;70(3):830-56. https://doi.org/10.1128/Mmbr.00016-06.

[35] Oliveira PH, Touchon M. Rocha EPC. Regulation of genetic flux between bacteria by

restriction-modification systems. P Natl Acad Sci USA 2016;113(20):5658-63.

https://doi.org/10.1073/pnas.1603257113.

[36] Cohen NR, Ross CA, Jain S, Shapiro RS, Gutierrez A, Belenky P, et al. A role for the

bacterial GATC methylome in antibiotic stress survival. Nat Genet 2016;48(5):581-6.

https://doi.org/10.1038/ng.3530.

[37] Oliveira PH, Ribis JW, Garrett EM, Trzilova D, Kim A, Sekulovic O, et al. Epigenomic

characterization of Clostridioides difficile finds a conserved DNA methyltransferase that mediates

sporulation and pathogenesis. Nat Microbiol 2019; https://doi.org/10.1038/s41564-019-0613-4.

[38] Xue MG, Zhao BS, Zhang ZJ, Lu MJ, Harder O, Chen P, et al. Viral N-6-methyladenosine

upregulates replication and pathogenesis of human respiratory syncytial virus. Nat Commun

2019;10 https://doi.org/ARTN 459510.1038/s41467-019-12504-y.

[39] Thurber RV, Haynes M, Breitbart M, Wegley L. Rohwer F. Laboratory procedures to generate

viral metagenomes. Nat Protoc 2009;4(4):470-83. https://doi.org/10.1038/nprot.2009.10.

[40] Ge XY, Li Y, Yang XL, Zhang HJ, Zhou P, Zhang YZ, et al. Metagenomic Analysis of Viruses

from Bat Fecal Samples Reveals Many Novel Viruses in Insectivorous Bats in China. J Virol

2012;86(8):4620-30. https://doi.org/10.1128/Jvi.06671-11.

[41] Froussard P. rPCR: a powerful tool for random amplification of whole RNA sequences.

Genome Research 1993;2185-90. https://doi.org/10.1101/gr.2.3.185.

[42] Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH. Phillippy AM. Canu: scalable and

accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res

2017;27(5):722-36. https://doi.org/10.1101/gr.215087.116.

[43] Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics

2018;34(18):3094-100. https://doi.org/10.1093/bioinformatics/bty191.

[44] Auslander N, Gussow AB, Benler S, Wolf YI. Koonin EV. Seeker: Alignment-free

identification of bacteriophage genomes by deep learning. bioRxiv 2020;2020.04.04.025783.

https://doi.org/10.1101/2020.04.04.025783.

[45] Ecale Zhou CL, Malfatti S, Kimbrel J, Philipson C, McNair K, Hamilton T, et al.

multiPhATE: bioinformatics pipeline for functional annotation of phage isolates. Bioinformatics

(Oxford, England) 2019;35(21):4402-4. https://doi.org/10.1093/bioinformatics/btz258.

[46] McNair K, Zhou C, Dinsdale EA, Souza B. Edwards RA. PHANOTATE: a novel approach to

gene identification in phage genomes. Bioinformatics (Oxford, England) 2019;35(22):4537-42.

https://doi.org/10.1093/bioinformatics/btz265.

[47] Grazziotin AL, Koonin EV. Kristensen DM. Prokaryotic Virus Orthologous Groups (pVOGs):

a resource for comparative genomics and protein family annotation. Nucleic acids research

2017;45(D1):D491-d8. https://doi.org/10.1093/nar/gkw975.

[48] Johnson LS, Eddy SR. Portugaly E. Hidden Markov model speed heuristic and iterative

HMM   search   procedure.   BMC   Bioinformatics   2010;11(1):431.

https://doi.org/10.1186/1471-2105-11-431.

[49] Edgar R. Taxonomy annotation and guide tree errors in 16S rRNA databases. PeerJ

2018;6e5030. https://doi.org/10.7717/peerj.5030.

[50] Langmead B, Wilks C, Antonescu V. Charles R. Scaling read aligners to hundreds of threads

on   general-purpose   processors.   Bioinformatics   (Oxford,   England)   2019;35(3):421-32.

https://doi.org/10.1093/bioinformatics/bty648.

[51] Marcus Stoiber JQ, Rob Egan, Ji Eun Lee, Susan Celniker, Robert K. Neely, Nicholas Loman,

Len A Pennacchio, James Brown. De novo Identification of DNA Modifications Enabled by

Genome-Guided Nanopore Signal Processing. bioRxiv 2017; https://doi.org/10.1101/094672.

[52] Robinson JT, Thorvaldsdottir H, Wenger AM, Zehir A. Mesirov JP. Variant Review with the

Integrative   Genomics   Viewer.   Cancer   Res   2017;77(21):e31-e4.

https://doi.org/10.1158/0008-5472.CAN-17-0337.

[53] Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, et al. MEME SUITE: tools

for motif discovery and searching. Nucleic Acids Res 2009;37(Web Server issue):W202-8.

https://doi.org/10.1093/nar/gkp335.

[54] Shkoporov AN, Clooney AG, Sutton TDS, Ryan FJ, Daly KM, Nolan JA, et al. The human

gut virome is highly diverse, stable and individual-specific. Cell Host & Microbe 2019;

https://doi.org/10.1101/657528.

[55] Moreno-Gallego JL, Chou S-P, Di Rienzi SC, Goodrich JK, Spector TD, Bell JT, et al.

Virome Diversity Correlates with Intestinal Microbiome Diversity in Adult Monozygotic Twins.

Cell Host & Microbe 2019;25(2):261-72.e5. https://doi.org/10.1016/j.chom.2019.01.019.

[56] Garmaeva S, Sinha T, Kurilshikov A, Fu J, Wijmenga C. Zhernakova A. Studying the gut

virome in the metagenomic era: challenges and perspectives. BMC Biol 2019;17(1):84.

https://doi.org/10.1186/s12915-019-0704-y.

[57] Minot S, Sinha R, Chen J, Li H, Keilbaugh SA, Wu GD, et al. The human gut virome:

inter-individual variation and dynamic response to diet. Genome Res 2011;21(10):1616-25.

https://doi.org/10.1101/gr.122705.111.

[58] Gregory AC ZO, Howell A, Bolduc B. Sullivan MB. The human gut virome database.

bioRxiv 2019; https://doi.org/10.1101/655910.

[59] Gardy JL. Loman NJ. Towards a genomics-informed, real-time, global pathogen surveillance

system. Nat Rev Genet 2018;19(1):9-20. https://doi.org/10.1038/nrg.2017.88.

[60] Depledge DP, Srinivas KP, Sadaoka T, Bready D, Mori Y, Placantonakis DG, et al. Direct

RNA sequencing on nanopore arrays redefines the transcriptional complexity of a viral pathogen.

Nat Commun 2019;10 https://doi.org/10.1038/s41467-019-08734-9.

[61] Charalampous T, Kay GL, Richardson H, Aydin A, Baldan R, Jeanes C, et al. Nanopore

metagenomics enables rapid clinical diagnosis of bacterial lower respiratory infection. Nat

Biotechnol 2019;37(7):783-792. https://doi.org/10.1038/s41587-019-0156-5.

[62] Courtney DG, Kennedy EM, Dumm RE, Bogerd HP, Tsai K, Heaton NS, et al.

Epitranscriptomic Enhancement of Influenza A Virus Gene Expression and Replication. Cell Host

Microbe 2017;22(3):377-386. https://doi.org/10.1016/j.chom.2017.08.004.

[63] Beaulaurier J, Schadt EE. Fang G. Deciphering bacterial epigenomes using modern

sequencing   technologies.   Nat   Rev   Genet   2019;20(3):157-72.

https://doi.org/10.1038/s41576-018-0081-3.

[64] Fang G, Munera D, Friedman DI, Mandlik A, Chao MC, Banerjee O, et al. Genome-wide

mapping of methylated adenine residues in pathogenic Escherichia coli using single-molecule

real-time   sequencing   (vol 30,   pg   1232,   2012).   Nat   Biotechnol   2013;31(6):566-9.

https://doi.org/10.1038/nbt0613-565f.

Figure Legends:

Figure 1. An integrated novel workflow for enrichment of virus-like particles

(VLPs), extraction of nucleic acids and ONT sequencing. The complete workflow

consists of four fragments: (1) Washing and filtration of faecal samples using

PBS/Sterile and PVDF membrane including step 1-3; (2) Precipitation of VLPs

including step 4-5; (3) Extraction, amplification and purification of viral nucleic acids

including step 6-8; (4) Construction of library and ONT sequencing including step

9-10.

Figure 2. Composition and relative abundance of viruses in each individual.

Different color of the bar represents different viral species or strains. The line between

unamplified and amplified group in each individual represents common virus species

between two groups.

Figure 3. Proportion of raw reads to contigs in length. The proportion of raw reads

to contigs in length is shown by combination of scatter diagram, boxplot and violin

chart. Different color represents five different individuals. (A) raw DNA group, (B)

amplified DNA/cDNA group.

Figure 4. Different methylation sites identification and motif recognition of viral

contig. (A) the distribution of methylation sites (5mC and 6mA) in the

contig00000015; (B) 5mC motif; (C) 6mA motif.

A. Raw DNA group

B. Amplified DNA/cDNA group

A

Contig00000015

5mC

6mA

B

5mC motif

C

6mA motif