Association of *CDH11* with ASD revealed by matched-gene

co-expression analysis and mouse behavioral studies

Nan Wu[1], Yue Wang[2], Jing-Yan Jia[1], Yi-Hsuan Pan[1*], Xiao-Bing Yuan[1,3*]

[1]Key Laboratory of Brain Functional Genomics of Shanghai and Ministry of

Education, Institute of Brain Functional Genomics, School of Life Science and the

Collaborative Innovation Center for Brain Science, East China Normal University,

Shanghai 200062, People's Republic of China

[2]Hussman Institute for Autism, Baltimore, MD 21201, USA

[3]Department of Anatomy and Neurobiology, University of Maryland School of

Medicine, Baltimore, MD 21201, USA

*Correspondence and requests for materials should be addressed to Dr. Xiao-Bing

Yuan (email: xbyuan@brain.ecnu.edu.cn) or Dr. Yi-Hsuan Pan (email:

yihsuanp@gmail.com)

Keywords: *CDH11*, ASD, gene co-expression analysis, matched-gene co-expression

analysis, MGCA

Abstract

A large number of putative risk genes of autism spectrum disorder (ASD) have been

reported. The functions of most of these susceptibility genes in developing brains

remain unknown, and a causal relationship between their variations and autism traits

has not been established. The aim of this study is to predict putative risk genes at the

whole-genome level based on the analysis of gene co-expression with a group of high

confidence ASD risk genes (hcASDs). Results showed that three gene features,

including gene size, mRNA abundance, and guanine-cytosine content, affect

genome-wide co-expression profiles of hcASDs. To circumvent the interference of

these gene features on gene co-expression analysis (GCA), we developed a method to

determine whether a gene is significantly co-expressed with hcASDs by statistically

comparing the co-expression profile of this gene with hcASDs to that of this gene

with permuted gene sets of feature-matched genes. This method is referred to as

"matched-gene co-expression analysis" (MGCA). With MGCA, we demonstrated the

convergence in developmental expression profiles of hcASDs and improved the

efficacy of risk gene prediction. Results of analysis of two recently reported ASD

candidate genes, *CDH11* and *CDH9,* suggested the involvement of *CDH11*, but not

CDH9, in ASD. Consistent with this prediction, behavioral studies showed that

Cdh11-null mice, but not *Cdh9*-null mice, have multiple autism-like behavioral

alterations. This study highlighted the power of MGCA in revealing ASD-associated

genes and the potential role of CDH11 in ASD.

Introduction

Autism spectrum disorder (ASD) is a heterogeneous neurodevelopmental condition

with a complex genetic basis [1, 2]. A large number of putative risk genes have been

identified by genetic linkage analyses, genome-wide association studies (GWAS),

whole-exome sequencing (WES), or whole-genome sequencing (WGS) [3-5]. However,

the functions of most of these putative risk genes in developing brains remain

unknown. For some novel risk genes, the genetic evidence supporting their

association with ASD is not sufficient. Therefore, a causal relationship between the

variations of many risk genes and autism traits has not been established. In order to

prioritize the investigation of genes and signaling pathways of high relevance to ASD,

a method to determine the functional importance of a large group of putative risk

genes is vital.

The highly diverse ASD risk genes are believed to functionally converge in

several common molecular pathways closely related to ASD, such as the Wnt

signaling pathway, the mammalian target of rapamycin (mTOR) pathway, and

dendrite development and synaptic remodeling pathways [3, 6]. Consistent with the

functional convergence of ASD risk genes, several studies showed the convergence of

developmental expression profiles of a large group of risk genes [7, 8]. It is generally

believed that genes with similar expression profiles are co-regulated or have related

functions [7, 9]. The co-expression of genes within a biological pathway is a strong

indication of their shared functions [9]. Based on this concept, computational analyses

of various brain transcriptomes have been conducted to identify potential

co-expression networks of ASD risk genes and to discover brain circuits that may be

affected by risk genes [7, 8, 10, 11]. In these studies, the correlation coefficient (CC) of a

pair of genes is calculated based on their expression levels in different brain regions

or developmental stages. Genome-wide gene co-expression networks are constructed

by setting an empirically determined threshold of CC [7]. Genetic mutation and protein

interaction data have also been incorporated into gene co-expression analysis (GCA)

along with novel data analysis algorithms, such as machine learning, to predict

putative ASD risk genes and their convergent molecular pathways[12]. A major

limitation in most of these studies is the lack of ways to overcome the potential effects

of confounding factors such as the size, expression level (mRNA abundance), and

guanine-cytosine (GC) content of genes on the result of GCA [13]. Most ASD risk

genes are large genes with a higher expression level in the brain than in other tissues

[14]. It is unclear whether the size or expression level of an ASD gene affects its

co-expression with other genes. It is also unclear whether the convergent pattern of

developmental expression profiles is specific to ASD risk genes or the common

property of genes with similar features, such as large gene size and high mRNA

abundance [13].

In this study, we discovered that three gene features, including mRNA

abundance, gene size (genomic DNA length), and GC content of a gene, affect gene

co-expression profiles in the brain. We developed a novel method called

"matched-gene co-expression analysis" (MGCA) (Fig. 1) to examine whether a gene

exhibits significant co-expression with a group of high-confidence ASD risk genes

(hcASDs) independent of confounding gene features. This was accomplished by

statistically comparing the co-expression level of a gene with the hcASD gene set to

that of this gene with a large number of permuted gene sets of matched features.

MGCA effectively predicted putative ASD-associated genes and revealed functionally

important convergent molecular pathways of these genes. MGCA also revealed

"homophilic cell adhesion" as one of the most significantly converged pathways of

risk genes. Results of further analysis of *CDH11* and *CDH9*, two ASD candidate

genes belonging to the cadherin family of adhesion molecules, suggested the

involvement of *CDH11*, but not *CDH9*, in ASD. This prediction was supported by

mouse behavioral studies using *Cdh11-* and *Cdh9*-null mice.

Materials and Methods

Data filtering and computation of co-expression coefficient

The human brain transcriptome dataset from BrainSpan (www.brainspan.org)

(RNA-Seq Gencode v10) was used for GCA. This dataset contained 256

transcriptomes of 16 different brain regions. The developmental stages ranged from

post-conception week 8 (PCW8) to 40 years old (40Y). Normalized mRNA

expression values were represented by RPKM (Reads Per Kilobase Per Million

Mapped Reads). The average mRNA expression level of each gene in all tissues was

considered as the mRNA abundance level of a gene. Gene lengths were determined

based on gene annotations provided by the National Center for Biotechnology

Information (NCBI). The GC content of a gene was obtained from Ensembl Genome

Browser. Based on statistical analyses of genetic data described previously, 101 risk

genes that reached a genome-wide significance threshold (false discovery rate, FDR ≤

0.1) [15] were used as the hcASD gene set (high confidence ASD risk gene set, **Table

S1**). Genes with an abundance level lower than the lowest abundance level of hcASDs

were filtered out (**Table S1**). Perl scripts were written to conduct most calculations.

Pair-wise Pearson's correlation coefficient (CC) was used to indicate the tendency of

co-expression of a gene pair. Heatmaps were constructed with the software R based

on the CC matrix of 1/100 evenly distributed genes. The mean CC was defined as

co-expression coefficient (CEC), which indicates the tendency of co-expression of a

gene with a specific set of genes ($CEC = \frac{1}{M}\sum_{i=1}^{M}CC_i$, $i = 1, 2, \ldots, M$; where M

indicates the total gene number of a gene set) or the tendency of co-expression of two

gene sets ($CEC = \frac{1}{N*M}\sum_{k=1}^{N}\left(\sum_{i=1}^{M}CC_{k\,i}\right)$, $k = 1, 2, \ldots, N$; $i = 1, 2, \ldots, M$; where M and N

represent the total gene number of two different gene sets, respectively).

Gene set definition

After data filtering, a total of 12,250 genes with information on gene length, mRNA

level, and GC content available were identified and used for the study (**Table S1**). In

addition to the hcASD gene set, the following gene sets were also used: cASD

(combined ASD genes), mRand (matched random genes), Rand (random genes),

TriM (triple-matched genes), Top (Top-ranked genes by CEC value), TriM-only

(genes only in TriM gene set), and Top-only (genes only in Top gene set). The cASD

gene set is a combined set of ASD-associated genes containing 514 non-redundant

genes from nine different sets of previously reported ASD-associated genes (**Table S5**)

[7, 15-21]. Each mRand gene set contained 101 genes, with one of the three gene features

of each gene matched with that of the corresponding hcASD gene in the hcASD gene

set. To generate the mRand gene sets, each feature-matched gene was randomly

selected within the ±50 range of the corresponding hcASD gene in the ranked gene

list using a Perl Script. Each Rand gene set contained 101 genes randomly selected

from the whole gene list using a Perl Script without considering matched gene

features. TriM was the set of MGCA-revealed genes that exhibited significant

co-expression with the hcASD gene set under all three matched conditions (see **Figs.**

1, 4A). The Top gene set contained top-ranked genes with the highest CEC values

with hcASD. TriM-only and Top-only gene sets contained non-overlapped genes

present only in the TriM or the Top gene set.

Gene ontology (GO) analysis

GO analysis was performed using DAVID v6.8 (http://david.ncifcrf.gov/tools.jsp),

and the human whole-genome genes provided by DAVID were used as the

background list. A corrected P-value of 0.05 (Benjamini-Hochberg method) was used.

Pathway enrichment analysis

Metascape (http://metascape.org) was used to perform pathway enrichment analysis and draw heatmaps. Pairwise similarities between any two significant terms were computed based on Kappa-test scores. The enriched terms were then hierarchically clustered into a tree with a kappa score of 0.3 as the threshold. Boxes were colored according to their *P*-values. Gray boxes indicate a lack of enrichment for a specific GO term.

Ethics approval

Animal care and handling were performed according to the guidelines for the Care and Use of Laboratory Animals of the National Institutes of Health. All animal experiments were approved by the Animal Care and Use Committees of Hussman Institute for Autism (06012015D), University of Maryland School of Medicine (0515017), and East China Normal University (m20190236).

Animals

Cdh9-null mice (*Cdh9*^Laz, C57BL/6-ICR mixed background) [22] were provided by Dr. Joshua R. Sanes at Harvard University. *Cdh11*-null mice [23] were from the Jackson Labs (*Cdh11^tm1Mta*/HensJ, https://www.jax.org/strain/023494, C57BL/6-129Sv-CD-1 mixed background). All mice were housed in groups of five with free access to food and water and kept on a 12-hour light/dark cycle. Mice were tattooed on the tail using

fluorescent ink for identification. A UV flashlight was used to visualize the tattooed

identification numbers. All behavioral tests were conducted during the daytime on

mice 2-5 months of age. The experimenter was blind to the genotype of the animal

during behavioral experiments. The surface of the apparatus for behavioral tests was

cleaned with 50% ethanol between tests. At least 5 min between cleaning and the

next test was allowed for ethanol evaporation and odor dissipation.

Genotyping

Genotyping of *Cdh9*-null mice was done by PCR as previously described [22]. The

PCR product for the wildtype (WT) *Cdh9* allele was 550 bp amplified with the primer

pair *Cdh9*-P1 (CCA CTA CAG GAA ACC TTT GGG TT) and *Cdh9*-P3 (ATG CAA

ACC ATC AGG TAT ACC AAC C), and that of the mutant allele was 430 bp

amplified with the primer pair *Cdh9*-P1 and *Cdh9*-P2 (CGT GGT ATC GTT ATG

CGC CT). The annealing temperature for *Cdh9* PCRs was 63°C. For genotyping of

Cdh11-null mice, the primer pair *Cdh11*-P1 (CGC CTT CTT GAC GAG TTC) and

Cdh11-P2 (CAC CAT AAT TTG CCA GCT CA) were used for amplification of the

mutant allele, and the primer pair *Cdh11*-P3 (GTT CAG TCG GCA GAA GCA G)

and *Cdh11*-P2 were used for the WT allele. The annealing temperatures for PCR were

63.1°C and 56°C for the mutant and WT alleles, respectively. The sizes of the PCR

products for the mutant and WT alleles were 500 bp and 400 bp, respectively.

Behavioral tests

Mice of 3-5 months old were used for these behavioral tests. Animals were handled

before the test (10 min/day for 3 days). The general order of behavioral tests was the

open field test, elevated plus maze, sociability test, rotarod test, and gripping force

test. Animals rested for at least 3 days after finishing one behavioral experiment.

During all behavioral experiments, the experimenter was blind to mouse genotypes.

Three batches of mice were analyzed, and data were pooled for analysis.

Open field test. The test mouse was allowed to freely explore the open field arena (50

cm $\times$ 50 cm) for 30 min. The mouse's motion was videoed and tracked by an

automated tracking system (EthoVision XT 11.5), which also recorded rearing,

hopping, turning, self-grooming, moving time, total moving distance, and time spent

in the center of the arena (1/2 of total size).

Elevated plus maze test. The standard elevated plus maze (EPM) apparatus consisted

of two open and two closed arms, 30×5 cm each, connected by a central platform (5

cm $\times$ 5 cm). The maze was 30 cm off the ground. The test mouse was gently placed

on the central platform with its head facing one closed arm and was allowed to

explore for 10 min freely. The mouse's time stayed in the two open arms and the

frequency of open arm entry were recorded.

Grip strength test. The test mouse was placed on a metal grid on top of a transparent

chamber. The grid was quickly inverted, and the time for the mouse to drop off the

grid was determined. Five consecutive trials were carried out, and the average

hanging time for each mouse was calculated. The maximum hanging time was set for

1 min. After 1 min of hanging, the trial was stopped, and the hanging time was

recorded as 1 min.

Horizontal bar test. The mouse was gently placed on a metal wire, with the two

forepaws gripping the wire. The length of time which the mouse hung on the wire was

measured. The maximum hanging time was set for 1 min. The average hanging time

was calculated from 5 consecutive trials.

Rotarod test. Mice were habituated to the rotarod apparatus (Harvard Apparatus

760770) by leaving them on the low-speed rotating rod (4 rpm) for 5 min each day for

3 days and tested on the fourth day on the accelerating rod. The time and the

maximum rotation speed that the test mouse maintained the balance on the rotating

rod were measured. Five consecutive trials were done for each mouse.

Social preference test. A modified three-chamber apparatus was used. The apparatus

comprised 3 rectangular (25 cm × 38 cm) chambers made of white Plexiglas with a 13

cm gate connecting the two side chambers to the middle chamber. A 3-sided (13 cm

wide for each side) fence made of transparent Plexiglas was placed inside each side

chamber facing the door of the side chambers, creating a 13 cm × 13 cm square area

separated from the side chambers but connected to the middle chamber through the

door (**Fig. 7A**). The two side chambers were covered by transparent Plexiglas to

minimize odorant diffusion. The test mouse was placed inside the middle chamber

and freely explored the middle chamber and the square zone in each side chamber for

10 min. Three social partner mice were then placed into the fenced area in one side

chamber, and the test mouse was allowed to explore for another 10 min freely.

Another 3 social partner mice were then placed in the other side chamber, and the

behavior of the test mouse was tracked for 10 min. The time that the test mouse spent

in each chamber was measured.

Experimental Design and Statistical Analysis

The effects of three different gene features on gene co-expression profiles were first

analyzed. MGCA (**Fig. 1**) was then performed to determine the convergence in

developmental expression profiles of hcASDs and detect genes that were significantly

co-expressed with hcASDs in the whole genome. The effectiveness of MGCA in

predicting putative ASD-associated genes was analyzed by comparing its results with

those of GCA, which does not consider. *CDH11* and *CDH9* were selected as example

genes, and mouse behavioral experiments were conducted in gene knockout mice to

verify the findings of MGCA.

Data are presented as mean ± standard error of the mean (SEM). The upper fence

test and Grubbs' test were performed to evaluate whether the hcASD-hcASD

expression level (CEC value) was significantly higher than the CEC values of

feature-matched (mRand-mRand; hcASD-mRand) or non-matched non-hcASD gene

sets (Rand-Rand; hcASD-Rand). Grubb's test was done using the "grubbs.test" script

in the R software package. The false discovery rate (FDR) of a gene was determined

by the frequency of this gene significantly co-expressed ($P < 0.001$) with 5,000

mRand gene sets determined by MGCA. Gene enrichment P-value was determined

with the Chi-Square test. Pathway enrichment P-values were determined with

Metascape. Behavioral data were analyzed by Student t-test and by one-way ANOVA

followed by Dunnett's t-test as post hoc analysis using SPSS (IBM, Armonk, USA) or

GraphPad Prism (GraphPad Software, La Jolla, CA, USA).

Availability of data and materials

Perl scripts for data analysis are available on GitHub

(https://github.com/wunan124/MGCA)

Results

Effects of gene features on gene co-expression profiles

The potential effect of the three gene features, including mRNA abundance, gDNA

size, and GC content, on gene co-expression profiles was first analyzed. The

BrainSpan human brain transcriptome dataset was used for this analysis. This dataset

contains transcriptomes of human (both sex) brain tissues from 16 different brain

regions of various developmental stages and ages (from PCW8 to 40Y). A total of

12,250 genes with information on all 3 gene features were used for analyses. These

genes were placed in ascending order of mRNA abundance, gDNA size, or GC

content as ranked gene lists (**Table S1**). The correlation coefficient (CC) of each gene

pair was calculated to reflect the co-expression level of the two genes, and results

were displayed in pseudo- color-coded matrices. In each of the CC matrices (**Fig. 2A**),

these 12,250 genes were placed in ascending order on both x and y axes. All three CC

matrices exhibited variable color intensity in different areas with higher intensity

corresponding to higher CC values. The overall color intensity was the highest in

areas corresponding to medium mRNA abundance, medium to large gDNA size, and

low GC content (**Fig. 2A**). This result suggests that all three gene features affect gene

co-expression profiles.

Most hcASDs are large genes with medium to large mRNA abundance but with

no apparent bias in GC content (**Fig. S1**). To determine whether each of these three

gene features affects the co-expression of a gene with the hcASD gene set as a whole,

the co-expression coefficient (CEC, mean CC between a gene and each of the hcASD

genes) of each of the 12,250 genes with the entire hcASD gene set was calculated

(blue dots in **Fig. 2B; Table S1**). In each of the 3 panels (**Fig. 2B**), the 12,250 genes

were placed in ascending order (x-axis). A noise-reduced (by data averaging) CEC

distribution curve was then generated by plotting the average CEC of a gene with its

neighboring 20 (10 above and 10 below; ±10), 50 (±25), 100 (±50), or 200 (±100)

genes on the gene lists under each gene ranking condition. Results showed a

bell-shaped curve when genes were ranked by mRNA abundance, suggesting that

genes with medium expression levels are more likely to co-express with the hcASD

gene set (**Fig. 2B**, left panel). There was an overall positive correlation between the

gDNA size of a gene and its CEC with the hcASD gene set (**Fig. 2B**, middle panel).

The CEC maintained a relatively high level (0.28 - 0.35) when the GC content ranged

from low to medium level (approximately < 45%, x-axis < 6000) and then gradually

declined with increasing GC content (**Fig. 2B**, right panel; **Table S1**).

With cubic regression, each noise-reduced CEC distribution curve was found to

have an R^2 value > 0.88 (**Fig. S2; Table S2**), indicating a significant correlation

between each of these gene features and the tendency of co-expression of a gene with

the hcASD gene set. When the 12,250 genes were placed in stochastic (random)

orders, CECs were evenly distributed, and the noise-reduced CEC distribution curves

were largely flat (**Fig. 2C**).

Similar genome-wide gene co-expression profiles of the hcASD gene set were

observed in transcriptomes of early (8PCW - 2Y) and late (4Y - 40Y) stages (**Fig.**

S3A), both sex, and different brain regions (**Fig. S3B, C**). These findings suggest that

the co-expression profile of hcASD genes is affected by all three gene features,

regardless of developmental stages, sex, and brain areas. Similar effects of these three

gene features on the co-expression profiles of hcASD genes were observed when a set

of 64 high susceptibility genes [24] were used as the hcASD gene set (**Fig. S4**).

Similar co-expression profiles of feature-matched gene sets

The genome-wide gene co-expression profile of the hcASD gene set was then

compared to the profiles of 200 feature-matched non-hcASD gene sets. Each gene set

comprised an equal number (101) of randomly selected and feature-matched

non-hcASD genes under the three different gene ranking conditions (**Fig. 3A**). These

gene sets were named "matched random" (mRand) gene sets (see methods). In

general, the genome-wide CEC distribution of hcASDs was similar to that of each of

the 200 mRand gene sets under all three gene ranking conditions. These findings

suggest that gene sets with matched gene features have a similar genome-wide

co-expression profile as the hcASD gene set. However, genes with low to moderate

mRNA abundance (approximately 1.2 - 30 RPMK, 1 - 10500 on x-axis) had higher

noise-reduced CECs with the hcASD gene set than with any of the 200 mRand gene

sets. In contrast, high mRNA abundance genes (> 30 RPKM; 10500 - 12250 on the

x-axis) had lower noise-reduced CECs with the hcASD gene set than with most

mRand gene sets. Moreover, genes with medium to large sizes had higher

noise-reduced CECs with the hcASD gene set than with most size-matched mRand

gene sets. Except those with the highest GC content, most genes had higher

noise-reduced CECs with the hcASD gene set than with most GC content-matched

mRand gene sets.

Co-expression of ASD risk genes

To determine whether hcASDs exhibit a significant tendency of co-expression with

each other, the mean CEC of each of the 101 hcASDs with the hcASD gene set as a

whole (hcASD-hcASD, see method) was compared to that of a large number of

permuted gene sets, each comprised equal number of feature-matched non-hcASD

genes (mRand-mRand) or randomly selected non-hcASD genes (Rand-Rand), and to

the CEC between hcASD and mRand (hcASD-mRand) or Rand (hcASD-Rand) gene

sets. Two hundred each of mRand and Rand gene sets were first analyzed. Results

showed that feature-matched gene sets (mRand) had overall higher CECs than

random gene sets (Rand) under all three matched conditions (@@@ in **Fig. 3B**),

suggesting that genes with similar features tend to co-express with each other. The

CEC of hcASD-hcASD (dashed line in **Fig. 3B**) was beyond the 3 times interquartile

range [Q3 + 3 × (Q3 - Q1), 3x upper fence] of the CECs of mRand-mRand,

Rand-Rand, hcASD-mRand, and hcASD-Rand gene sets. Results of Grubbs' test

confirmed this tendency (*** in **Fig. 3B**). These results suggest that hcASDs have a

significantly greater co-expression tendency with each other than with other

feature-matched non-hcASD genes or randomly selected genes. To corroborate this

finding, a permutation test was conducted with 100,000 permuted sets of genes with

matched or non-matched features. The CEC of hcASD-hcASD was again found to be

significantly larger (permutation *P*-value < 0.00001) than that of hcASD-mRand,

mRand-mRand, hcASD-Rand, or Rand-Rand (### in **Fig. 3B**), indicating a significant

co-expression tendency of hcASDs.

Significant co-expression of hcASDs was also observed in transcriptomes of

brain tissues from both early (8PCW – 2Y) and late (4Y – 40Y) stages (**Fig. S5A, B**),

both sex (**Fig. S5C, D**), and different brain regions (**Fig. S6A-C**). These results

indicate a highly conserved co-expression profile of hcASDs. Combined ranking of

-log10 *P*-values of Grubbs' test under all three different matched conditions was then

performed to determine the relative significance level of co-expression of hcASDs

with each other in different brain regions (**Fig. S6D**). The top four brain regions with

the highest significance levels were cerebellum (CB), striatum (STR), orbital frontal

cortex (OFC), and dorsal frontal cortex (DFC); these are the brain regions previously

implicated in ASD [25-32]. These results suggest that hcASDs play important roles in

the development and function of these ASD-relevant brain regions.

ASD-associated genes and pathways identified by MGCA

Genes whose CECs with hcASDs were significantly higher than with permuted gene

sets composed of feature-matched genes under each of the three matched conditions

were considered significantly co-expressed with hcASDs (estimated FDR of each

gene below 1.25×10^{-4}). These genes were named TriM (triple-matched) genes (**Figs.**

1, 4A; Table S3). TriM genes were then compared with a gene set containing 514

non-redundant genes from nine different sets of previously reported ASD-associated

genes (cASD, see methods) and with a set of "True negative" ASD-associated genes,

which were genes associated with non-mental health diseases but not with ASD [12].

When the permutation *P*-value was either below 0.0001 or below 0.00001, the TriM

gene set showed a significant enrichment of cASD genes ($P = 0.0027$ and $P < 0.0001$,

respectively; chi-square test, **Fig. 4B**). This result suggests that at a permutation

P-value below 0.0001, TriM genes have a high rate of positive prediction of being

ASD-associated genes. When the permutation *P*-value was below 0.00001, TriM

genes exhibited a significant negative enrichment of "True negative" genes ($P =$

0.0068; chi-square test, **Fig. 4B**), suggesting a significantly low false-positive rate in

the prediction of ASD-associated genes.

Altogether, MGCA revealed 2370 TriM genes with a permutation *P*-value <

0.0001. These genes (TriM-2370) were compared with an equal number (2370) of

genes that had the highest CECs with the hcASD gene set (referred to as Top-2370

gene set, **Table S3**). TriM-2370 and Top-2370 gene sets had 1414 genes in common

(overlapped), and each had 956 non-overlapped genes. These two non-overlapped

gene sets were named TriM-only and Top-only, respectively (**Fig. 4C**; **Table S3**).

Most Top-only genes had a medium mRNA abundance level, a large gene size, and a

high CEC value (> 0.46), whereas TriM-only genes had a broad range of mRNA

abundance, gene size, and CEC values (**Fig. 4C**). Gene ontology (GO) enrichment

analysis of the TriM-2370 gene set showed significant over-representation of genes in

molecular pathways closely related to ASD, including gene transcription regulation,

homophilic cell adhesion, axon guidance and axon extension, synapse assembly, Wnt

signaling pathway, neuron migration, covalent chromatin modification, and protein

polyubiquitination. Fewer pathways relevant to ASD were revealed in the Top-2370

gene set by GO analysis, including covalent chromatin modification, transcription

regulation, protein polyubiquitination, Wnt signaling pathway, and negative

regulation of autophagy (**Fig. 4D; Table S4**). To investigate the functional

relationship between cASD and TriM-2370 or Top-2370 gene sets, the enriched

molecular pathways of cASD, TriM-only, Top-only, and overlapped gene sets were

subject to pathway enrichment analysis [33] (**Fig. S7**). Results showed that TriM-only

and Top-only gene sets converged in different but complementary molecular

pathways. The molecular pathways of the cASD gene set were found to cluster closer

to those of the TriM-only set than to those of the Top-only set, suggesting that

TriM-only genes have a closer functional relationship with cASD genes than with

Top-only genes. Some well-established ASD risk genes, such as *FOXP1*, *TBR1*,

SHANK2, *SYNGAP1*, and *PCDH9*, were found in the TriM-only gene set, suggesting

a better performance of MGCA than conventional GCA in revealing ASD-relevant

molecular pathways.

The effectiveness of MGCA in predicting ASD-relevant genes was compared

with that of GCA and two other risk gene prediction algorithms. One was DAWN

(Detecting Association With Networks), which has been used to analyze the

association between rare genetic variations and gene co-expression in the mid-fetal

prefrontal and somatosensory cortex [34]. The other algorithm was EWML

(Evidence-Weighted Machine Learning) that has been used to predict the probability

of ASD association with whole-genome genes based on data of gene co-expression,

genetic mutations, and protein-protein interaction [12]. The combined ASD risk gene

set (cASD) and the "true negative" gene set were used to conduct the

cross-comparison between different algorisms. At a permutation *P*-value of 0.0001 or

0.00001, TriM genes had higher enrichment of cASD genes than an equal number of

genes with highest CEC values (Top), an equal number of top-ranked ASD-linked

genes predicted by EWML, and network ASD genes (nASD) identified by DAWN

(**Fig. 4E**). Furthermore, fewer TriM genes overlapped with "True negative" genes

than genes predicted by DAWN and EWML, which had significant enrichment of

"true negative" genes (**Fig. 4E**), suggesting a lower rate of false-positive prediction by

MGCA than that by EWML and DAWN (**Fig. 4E**). Thus, MGCA performs better

than GCA by a higher positive prediction rate and performs better than the EWML

and DAWN algorisms by both a higher positive prediction rate and a lower prediction

error.

Co-expression of cadherin genes with hcASDs

Consistent with previous findings [8], we found that homophilic cell adhesion is the

most significantly over-represented pathway of TriM-2370 genes (**Fig. 4D; Table S4**).

Some cadherin family members in the TriM-2370 gene set, such as *PCDH19*, are

known to be high-risk ASD genes (**Table S6**) that play important roles in brain circuit

development [35, 36]. Several cadherin family members were also found in the

TriM-2370 gene set, including many members of the protocadherins β gene cluster

and Dachsous Cadherin-related 1 (*DCHS1*), suggesting that these genes also

participate in the development and function of ASD-relevant brain circuits. Some

cadherin genes were not significantly co-expressed with hcASDs under any of the

matched conditions; these genes were referred to as tri-negative genes (TriN; **Table**

S6). Several recent genetic studies have implicated two type II cadherins, *CDH11* and

CDH9, in ASD and other psychiatric diseases [37-41]. The CEC values of *CDH11* and

CDH9 with hcASDs were ranked at 5244 and 9581, respectively, in whole-genome

genes. Therefore, neither of them belonged top-ranked genes based on traditional

GCA. Using MGCA, we found that *CDH11* and *CDH9* belonged to TriM and TriN

gene sets, respectively. We thus hypothesized that *CDH11*, but not *CDH9*, is more

likely to be associated with ASD.

Autism-like traits of *Cdh11*-null mice

To assess the functional relevance of *CDH11* and *CDH9* to ASD, the behaviors of

Cdh11- and *Cdh9*-null mice were investigated. In the open field test (OFT), both male

and female *Cdh11*-null mice spent a longer time exploring the central area of the open

field arena than wild type (WT) littermates (**Fig. 5A, D**). Heterozygous littermates

showed a similar but less significant pattern. Total locomotion distance and average

moving speed of *Cdh11*-null mice were slightly reduced compared to WT littermates

(**Fig. 5B, C**). Both male and female *Cdh9*-null mice were largely normal in the OFT

(**Fig. 5E-G**).

In the elevated plus maze test, female *Cdh11*-null mice visited the open arm

more frequently and spent a significantly longer time there. Heterozygous females

spent a slightly but not statistically significantly more time in the open arm (**Fig. 5H,**

I). The increased time and frequency of open arm exploration by female *Cdh11*-null

mice is consistent with the results of a previous study using the same mouse line of

mixed sex [42]. Male *Cdh9*-null mice showed longer exploration of the open arm, but

female *Cdh9*-null mice did not, although female heterozygotes showed an increased

frequency of open arm entry (**Fig. 5J, K**).

Individuals with ASD often have a weaker grip strength than age-matched

controls [43]. The gripping strength test and the horizontal bar test showed that both

male and female *Cdh11*-null mice exhibited significantly shorter hanging duration

than WT littermates (**Fig. 6A, B**), indicating reduced gripping strength or impaired

motor coordination. The gripping strength of *Cdh9*-null mice was normal (**Fig. 6C**).

The rotarod test was conducted to evaluate motor-related functions of null mice.

Since female and male mutant mice displayed similar behaviors in most of the above

behavioral tests, only female mice were analyzed in this test. Compared to WT

littermates, *Cdh11*-null mice, but not *Cdh9*-null mice, stayed longer on the rotarod

and endured a higher rotation speed in the initial trial (**Fig. 6D-G**). In subsequent

trials, *Cdh11*-null mice did not display significant performance improvement (**Fig. 6D,**

E), indicating impaired motor learning. The enhanced performance of *Cdh11*-null

mice in the initial trial was very similar to the phenotype of several other

well-characterized ASD mouse models and suggested increased repetitive motion of

these mutant mice [44].

Repetitive behaviors were then evaluated by measuring the duration and

frequency of self-grooming within 10 minutes, during which mice were placed in a

novel or a relatively familiar environment. As shown in **Fig. 6H-I**, during the first 10

minutes of exploring a novel chamber, *Cdh11*-null mice exhibited a significantly

greater frequency of self-grooming than WT littermates, indicating elevated repetitive

behavior in a novel environment. *Cdh11*-null mice also showed a significantly higher

frequency of self-grooming than WT littermates during the second 10-minute period

(**Fig. 6J, K**), indicating elevated repetitive behavior even in a relatively familiar

environment. No such behavioral alteration was observed in *Cdh9*-null mice (**Fig. 6L,**

M).

The modified three-chamber social preference test was conducted to evaluate the

sociability of mutant mice. One main modification was an enlargement of the area for

housing social partner mice to reduce their potential stress and anxiety. Another major

modification to the protocol was using three mice instead of a single mouse as social

partners. This was done to increase the availability of social cues and reduce the

variability of test results caused by differences in the sociability of individual social

partners (**Fig. 7A**). In addition, the two side chambers were covered on the top to slow

the diffusion and mixing of odorant cues. Results showed that female *Cdh11*-null

mice exhibited a significant preference to social partner mice than to an object and a

significant preference to novel partners than to familiar ones (**Fig. 7B, C**). However,

compared to WT littermates, mutant mice spent a significantly longer time in the

middle chamber but significantly shorter time interacting with partner mice (**Fig. 7B,**

C), indicating reduced sociability. In contrast, *Cdh9*-null mice did not show any

abnormality in this test (**Fig. 7D, E**).

Discussion

Gene co-expression analysis (GCA) is a powerful tool to find functionally convergent

genes. Several previous GCA studies had considered the potential effect of gene size

and GC content on the co-expression of ASD risk genes [7]. In the present study, we

discovered that three gene features, including mRNA abundance, gDNA size, and GC

content, affected the genome-wide gene co-expression profiles in the brain. Although

the mechanisms by which different gene features affect gene co-expression are

unknown, our findings suggest the importance of considering the effect of these gene

features in gene co-expression analysis (GCA). As an example of the potential

influence of confounding gene features on GCA, genes that are stably expressed in the

brain may have high CCs with each other. These aleatory high CCs of

high-abundance genes do not mean a real co-expression relationship of genes of

common molecular pathways. One possible influence of gene size on GCA is that

large-size genes may have long noncoding regions that could be potential regulatory

elements. Thus, compared to small genes, large genes may have more shared

regulatory elements, which means a higher chance of being co-regulated by common

transcription factors. Considering that most hcASD genes are large-size genes, they

may have a higher tendency of co-expression with large genes. Genes with high GC

content have higher mRNA stability and thus have higher chances of co-existence

with each other. However, since most hcASD genes have low to medium GC content

(**Fig 2B**), it may explain an overall negative correlation between the GC content of

genes and their co-expression with hcASDs.

Almost all previous studies ignored these important confounding factors and just

selected high-CC gene pairs to construct the gene co-expression network. Without

correcting the effect of this confounding factor, the effectiveness of GCA would be

compromised. Instead of setting a CC threshold for GCA as in most other studies, we

screened for significant co-expression relationships by comparing the co-expression

coefficient (CEC) of a gene with the hcASD gene set to that with permuted gene sets

of matched gene features. Only genes that had a CEC with the hcASD gene set

significantly higher than its CECs with permuted sets of feature-matched genes were

considered being co-expressed with hcASDs. This matched-gene co-expression

analysis (MGCA) paradigm **(Fig. 1)** allowed the demonstration of significant

co-expression of hcASDs and avoided the potential bias caused by an empirically

determined threshold for CC of gene pairs in GCA. Our results revealed that MGCA

is more efficient in predicting gene association than the pre-existing methods DAWN,

an algorism integrating genetic variants and gene co-expression data, and EWML, a

sophisticated machine learning algorithm with the integration of gene co-expression,

gene mutation database, and PPI network. We believe that the high performance of

MGCA could be attributed to the correction of three confounding gene features in the

determination of functionally relevant gene co-expression. Although correlations of

these confounding gene features with the co-expression with hcASDs have been

considered [7], the potential interference of these gene features on the construction of

gene co-expression networks was not considered in previous studies. Therefore,

MGCA will be an important complement to current gene association prediction

algorithms (**Fig. 1**). As MGCA is based solely on the gene co-expression data, future

algorithms combining MGCA with genetic mutation data and machine learning will

further improve its efficacy.

An important finding in this study is the plausible association of *CDH11* with

ASD determined by MGCA. Cadherins have been shown to accumulate in synaptic

junctions and regulate dendrite development and synapse maturation [45-48]. Several

cadherin family members, such as some protocadherins in the *FAT* cadherin

subfamily, have been implicated in ASD [49-58]. A genetic association study of a large

cohort of ASD individuals and matched controls revealed genes in the protocadherin

α gene cluster (*PCDHA*) as ASD risk genes [51]. Mutations in the *PCDH19* gene have

been shown to cause early-onset epilepsy, and many individuals with these mutations

also display autistic features [52-54]. Mutations in the cadherin EGF LAG seven-pass

G-type receptor 2 gene (*CELSR2*) were speculated to be responsible for the Joubert

syndrome, a disease with a high degree of autistic features [59, 60]. It is uncertain

whether other cadherins are also high-risk factors. Using MGCA, we found that a

group of cadherin superfamily members exhibited a high co-expression with hcASDs,

suggesting shared functions with hcASDs and a role in ASD etiology. Among them,

several protocadherins, mainly *PCDHB*s, exhibited significant co-expression with

hcASDs (**Table S6**). The functions of these putative ASD-associated cadherins in the

brain remain to be determined. One of such cadherins identified by MGCA is CDH11.

In this study, we found that *Cdh11*-null mice had significantly increased repetitive

behaviors. The brain regions, including the neocortex, CB, and STR, are known to be

involved in the control of repetitive behaviors [61]. It is likely that cadherins, *Cdh11* in

particular, play important roles in mediating synapse formation during the wiring of

circuits in these brain areas. Consistent with this postulation, our recent work showed

Cdh11 expression in ASD-associated sub-regions in the CB of developing mouse

brain [62].

In human studies, partial deletion of *CDH11* was observed in a sporadic case of

non-syndromic ASD, mild intellectual disability, and attention deficit hyperactivity

disorder (ADHD) [37]. A case-control association study revealed a high prevalence of

a homozygous single nucleotide variant rs7187376C/C of *CDH11* in patients with

ASD [37]. Several other coding variants of *CDH11* were also discovered in ASD

individuals [37]. Behavioral changes that we have observed in *Cdh11*-null mice,

including reduced anxiety, increased repetitive behavior, and reduced sociability, are

highly consistent with the non-syndromic ASD case with partial deletion of *CDH11*

[37]. This observation supports the notion that loss-of-function of a single gene, such as

CDH11, is sufficient to cause major autism traits. Recessive mutations have been

implicated in ASD, and biallelic disruption of recessive neurodevelopmental genes in

ASD has been reported [62]. We found that homozygous *Cdh11*-null mice, but not

heterozygous *Cdh11*-null mice, displayed autism-like behavioral deficits. This

suggests that *CDH11* may be a recessive gene of ASD. However, clinical data from

more families with *CDH11* mutations are needed to determine whether it is true in the

future. Behavioral phenotypes of ASD are highly heterogeneous. Some individuals

with ASD are hypoactive with elevated anxiety, and some have attention deficit

hyperactivity disorder (ADHD) but with reduced anxiety [64-67]. The genetic and

neurobiological mechanisms underlying this behavioral heterogeneity have not been

fully determined. Further investigation with a larger cohort of patient families is

needed to determine whether loss-of-function mutations of *CDH11* are associated

with ADHD.

Most genetic variants found in patients with ASD are heterozygous. In some

behavioral tests, heterozygous *Cdh11*-null mice showed a similar trend of behavioral

alterations as homozygous null mice, but not at a statistically significant level (**Figs.**

6J, K; 7C). As ASD has a complex genetic basis and is affected by environmental

factors, it is conceivable that the haplodeficiency of a single risk gene causes a

relatively mild behavioral phenotype in mice. More severe behavioral deficits may

result if the haplodeficiency of *Cdh11* is combined with other genetic or

environmental factors. Our findings suggest that *CDH11* is significantly co-expressed

with hcASDs and that its mutations may exert a causal effect in autism traits.

Cdh11-null mice would be very useful in dissecting the circuit mechanisms

underlying a subgroup of ASD and in screening drugs targeting this subgroup of

ASD.

CDH9 plays a vital role in establishing specific synaptic wiring in both the

hippocampus and the retina [22, 68]. Its association with ASD has been suggested by

several studies on exome sequencing [49, 69]. The primary evidence linking *CDH9* to

ASD is the strong association of the single nucleotide polymorphism rs4307059

located in the intergenic region between *CDH10* and *CDH9* with ASD [70]. However,

this rs4307059 genotype was not correlated with the expression of either *CDH9* or

CDH10 in adult brains [70, 71], and whether a correlation exists in fetal brains is

unknown. Recently, an antisense noncoding RNA of the moesin pseudogene 1

(MSNP1AS) was shown to be transcribed from the locus harboring rs4307059.

Alterations in this pseudogene were postulated to contribute to ASD [71-73]. Whether

CDH9 deficiency is a causal factor for ASD remains undetermined. Our MGCA

showed that, unlike *CDH11*, *CDH9* was not co-expressed with hcASDs. This is an

indication that *CDH9* may not play an essential role in the wiring of ASD-relevant

circuits. Consistent with this notion, behavioral tests showed that *Cdh9*-null mice

exhibited a very mild behavioral abnormality only in the elevated plus maze test but

not in any other tests. With recent findings by other researchers [74], our results

suggest that *CDH9* deficiency may not have a significant effect on autism traits.

In conclusion, this study revealed the importance of considering matched gene

features in the analysis of gene co-expression and demonstrated the effectiveness of

MGCA in the identification of putative ASD-associated genes and their convergent

signaling pathways. The application of MGCA led to the determination of *CDH11* as

a putative ASD-associated gene. Our results also showed that *Cdh11*-null mice could

be used to study circuit mechanisms of a subgroup of ASD and explore therapeutic

strategies for ASD.

Acknowledgments: We thank Elizabeth Benevides for editing the manuscript. This

work was funded by the National Science Foundation of China (NSFC-31871501,

NSFC-ISF-32061143016) and Hussman Foundation (HIAS15006). X.Y. was

supported by the Simons Foundation (296143).

Authors' contributions: N.W., Y.W., and Y.P. analyzed data and generated figures.

J.J and X.Y. conducted behavioral tests. Y.P. and X.Y. designed the study and wrote

the manuscript.

Conflict of interest: The authors declare that they have no competing financial

interests.

References

[1] Berg JM, Geschwind DH. Autism genetics: searching for specificity and

convergence. Genome Biol 2012, 13: 247.

[2] Jeste SS, Geschwind DH. Disentangling the heterogeneity of autism spectrum

disorder through genetic findings. Nature Reviews Neurology 2014, 10: 74-81.

[3] de la Torre-Ubieta L, Won H, Stein JL, Geschwind DH. Advancing the

understanding of autism disease mechanisms through genetics. Nat Med 2016, 22:

345-361.

[4] Sullivan PF, Geschwind DH. Defining the Genetic, Genomic, Cellular, and

Diagnostic Architectures of Psychiatric Disorders. Cell 2019, 177: 162-183.

[5] Ronemus M, Iossifov I, Levy D, Wigler M. The role of de novo mutations in the

genetics of autism spectrum disorders. Nat Rev Genet 2014, 15: 133-141.

[6] Courchesne E, Pramparo T, Gazestani VH, Lombardo MV, Pierce K, Lewis NE.

The ASD Living Biology: from cell proliferation to clinical phenotype. Mol

Psychiatry 2019, 24: 88-107.

[7] Willsey AJ, Sanders SJ, Li M, Dong S, Tebbenkamp AT, Muhle RA, *et al.*

Coexpression networks implicate human midfetal deep cortical projection neurons in

the pathogenesis of autism. Cell 2013, 155: 997-1007.

[8] Parikshak NN, Luo R, Zhang A, Won H, Lowe JK, Chandran V, *et al.* Integrative

functional genomic analyses implicate specific molecular pathways and circuits in

autism. Cell 2013, 155: 1008-1021.

[9] Stuart JM, Segal E, Koller D, Kim SK. A gene-coexpression network for global

discovery of conserved genetic modules. Science 2003, 302: 249-255.

[10] Hormozdiari F, Penn O, Borenstein E, Eichler EE. The discovery of integrated

gene networks for autism and related disorders. Genome Res 2015, 25: 142-154.

[11] Mahfouz A, Ziats MN, Rennert OM, Lelieveldt BP, Reinders MJ. Shared

Pathways Among Autism Candidate Genes Determined by Co-expression Network

Analysis of the Developing Human Brain Transcriptome. J Mol Neurosci 2015, 57:

580-594.

[12] Krishnan A, Zhang R, Yao V, Theesfeld CL, Wong AK, Tadych A, *et al.*

Genome-wide prediction and functional characterization of the genetic basis of autism

spectrum disorder. Nat Neurosci 2016, 19: 1454-1462.

[13] Kopp N, Climer S, Dougherty JD. Moving from capstones toward cornerstones:

successes and challenges in applying systems biology to identify mechanisms of

autism spectrum disorders. Front Genet 2015, 6: 301.

[14] Gabel HW, Kinde B, Stroud H, Gilbert CS, Harmin DA, Kastan NR, *et al.*

Disruption of DNA-methylation-dependent long gene repression in Rett syndrome.

Nature 2015, 522: 89-93.

[15] Satterstrom FK, Kosmicki JA, Wang JB, Breen MS, De Rubeis S, An JY, *et al.*

Large-Scale Exome Sequencing Study Implicates Both Developmental and Functional

Changes in the Neurobiology of Autism. Cell 2020, 180: 568-584.

[16] Abrahams BS, Arking DE, Campbell DB, Mefford HC, Morrow EM, Weiss LA,

et al. SFARI Gene 2.0: a community-driven knowledgebase for the autism spectrum

disorders (ASDs). Mol Autism 2013, 4: 36.

[17] pfeliciano@simonsfoundation.org SCEa, Consortium S. SPARK: A US Cohort

of 50,000 Families to Accelerate Autism Research. Neuron 2018, 97: 488-493.

[18] Basu SN, Kollu R, Banerjee-Basu S. AutDB: a gene reference resource for

autism research. Nucleic Acids Res 2009, 37: D832-836.

[19] RK CY, Merico D, Bookman M, J LH, Thiruvahindrapuram B, Patel RV, *et al.*

Whole genome sequencing resource identifies 18 new candidate genes for autism

spectrum disorder. Nat Neurosci 2017, 20: 602-611.

[20] Iossifov I, O'Roak BJ, Sanders SJ, Ronemus M, Krumm N, Levy D, *et al.* The

contribution of de novo coding mutations to autism spectrum disorder. Nature 2014,

515: 216-221.

[21] Lin Y, Afshar S, Rajadhyaksha AM, Potash JB, Han S. A Machine Learning

Approach to Predicting Autism Risk Genes: Validation of Known Genes and

Discovery of New Candidates. Front Genet 2020, 11: 500064.

[22] Duan X, Krishnaswamy A, De la Huerta I, Sanes JR. Type II cadherins guide

assembly of a direction-selective retinal circuit. Cell 2014, 158: 793-807.

[23] Horikawa K, Radice G, Takeichi M, Chisaka O. Adhesive subdivisions intrinsic

to the epithelial somites. Dev Biol 1999, 215: 182-189.

[24] Sanders SJ, Xin H, Willsey AJ, Ercan-Sencicek AG, Samocha KE, Cicek AE*, et*

al. Insights into Autism Spectrum Disorder Genomic Architecture and Biology from

71 Risk Loci. Neuron 2015, 87: 1215-1233.

[25] Tsai PT, Hull C, Chu Y, Greene-Colozzi E, Sadowski AR, Leech JM*, et al.*

Autistic-like behaviour and cerebellar dysfunction in Purkinje cell Tsc1 mutant mice.

Nature 2012, 488: 647-651.

[26] Khan AJ, Nair A, Keown CL, Datko MC, Lincoln AJ, Muller RA.

Cerebro-cerebellar Resting-State Functional Connectivity in Children and

Adolescents with Autism Spectrum Disorder. Biol Psychiatry 2015, 78: 625-634.

[27] Buxhoeveden DP, Semendeferi K, Buckwalter J, Schenker N, Switzer R,

Courchesne E. Reduced minicolumns in the frontal cortex of patients with autism.

Neuropathol Appl Neurobiol 2006, 32: 483-491.

[28] Green SA, Hernandez L, Tottenham N, Krasileva K, Bookheimer SY, Dapretto

M. Neurobiology of Sensory Overresponsivity in Youth With Autism Spectrum

Disorders. JAMA Psychiatry 2015, 72: 778-786.

[29] Green SA, Rudie JD, Colich NL, Wood JJ, Shirinyan D, Hernandez L*, et al.*

Overreactive brain responses to sensory stimuli in youth with autism spectrum

disorders. J Am Acad Child Adolesc Psychiatry 2013, 52: 1158-1172.

[30] Di Martino A, Kelly C, Grzadzinski R, Zuo XN, Mennes M, Mairena MA*, et al.*

Aberrant striatal functional connectivity in children with autism. Biol Psychiatry 2011,

69: 847-856.

[31] Wang X, Bey AL, Katz BM, Badea A, Kim N, David LK, *et al.* Altered

mGluR5-Homer scaffolds and corticostriatal connectivity in a Shank3 complete

knockout model of autism. Nat Commun 2016, 7: 11459.

[32] Li X, Zhang K, He X, Zhou J, Jin C, Shen L, et al. Structural, Functional, and

Molecular Imaging of Autism Spectrum Disorder. Neurosci Bull 2021.

[33] Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, *et al.*

Metascape provides a biologist-oriented resource for the analysis of systems-level

datasets. Nat Commun 2019, 10: 1523.

[34] Liu L, Lei J, Sanders SJ, Willsey AJ, Kou Y, Cicek AE, *et al.* DAWN: a

framework to identify autism genes and subnetworks using gene expression and

genetics. Molecular Autism 2014, 5(1): 22.

[35] Yin J, Chun CA, Zavadenko NN, Pechatnikova NL, Naumova OY, Doddapaneni

HV, *et al.* Next Generation Sequencing of 134 Children with Autism Spectrum

Disorder and Regression. Genes (Basel) 2020, 11(8): 853.

[36] Emond MR, Biswas S, Jontes JD. Protocadherin-19 is essential for early steps in

brain morphogenesis. Developmental Biology 2009, 334: 72-83.

[37] Crepel A, De Wolf V, Brison N, Ceulemans B, Walleghem D, Peuteman G, *et al.*

Association of CDH11 with non-syndromic ASD. Am J Med Genet B Neuropsychiatr

Genet 2014, 165B: 391-398.

[38] Gasso P, Ortiz AE, Mas S, Morer A, Calvo A, Bargallo N, *et al.* Association

between genetic variants related to glutamatergic, dopaminergic and

neurodevelopment pathways and white matter microstructure in child and adolescent

patients with obsessive-compulsive disorder. J Affect Disord 2015, 186: 284-292.

[39] Sokolowski M, Wasserman J, Wasserman D. Polygenic associations of

neurodevelopmental genes in suicide attempt. Mol Psychiatry 2016, 21: 1381-1390.

[40] Wang K, Zhang H, Bloss CS, Duvvuri V, Kaye W, Schork NJ, *et al.* A

genome-wide association study on common SNPs and rare CNVs in anorexia nervosa.

Mol Psychiatry 2011, 16: 949-959.

[41] Anazi S, Maddirevula S, Faqeih E, Alsedairy H, Alzahrani F, Shamseldin HE*, et*

al. Clinical genomics expands the morbid genome of intellectual disability and offers

a high diagnostic yield. Mol Psychiatry 2017, 22: 615-624.

[42] Manabe T, Togashi H, Uchida N, Suzuki SC, Hayakawa Y, Yamamoto M*, et al.*

Loss of cadherin-11 adhesion receptor enhances plastic changes in hippocampal

synapses and modifies behavioral responses. Mol Cell Neurosci 2000, 15: 534-546.

[43] Travers BG, Bigler ED, Tromp do PM, Adluru N, Destiche D, Samsin D*, et al.*

Brainstem White Matter Predicts Individual Differences in Manual Motor Difficulties

and Symptom Severity in Autism. J Autism Dev Disord 2015, 45: 3030-3040.

[44] Rothwell PE, Fuccillo MV, Maxeiner S, Hayton SJ, Gokce O, Lim BK*, et al.*

Autism-associated neuroligin-3 mutations commonly impair striatal circuits to boost

repetitive behaviors. Cell 2014, 158: 198-212.

[45] Bekirov IH, Nagy V, Svoronos A, Huntley GW, Benson DL. Cadherin-8 and

N-cadherin differentially regulate pre- and postsynaptic development of the

hippocampal mossy fiber pathway. Hippocampus 2008, 18: 349-363.

[46] Friedman LG, Riemslagh FW, Sullivan JM, Mesias R, Williams FM, Huntley

GW*, et al.* Cadherin-8 expression, synaptic localization, and molecular control of

neuronal form in prefrontal corticostriatal circuits. J Comp Neurol 2015, 523: 75-92.

[47] Bian WJ, Miao WY, He SJ, Qiu Z, Yu X. Coordinated Spine Pruning and

Maturation Mediated by Inter-Spine Competition for Cadherin/Catenin Complexes.

Cell 2015, 162: 808-822.

[48] Tsai NP, Wilkerson JR, Guo W, Maksimova MA, DeMartino GN, Cowan CW*,*

et al. Multiple autism-linked genes mediate synapse elimination via proteasomal

degradation of a synaptic scaffold PSD-95. Cell 2012, 151: 1581-1594.

[49] Cukier HN, Dueker ND, Slifer SH, Lee JM, Whitehead PL, Lalanne E*, et al.*

Exome sequencing of extended families with autism reveals genes shared across

neurodevelopmental and neuropsychiatric disorders. Mol Autism 2014, 5: 1.

[50] Butler MG, Rafi SK, Hossain W, Stephan DA, Manzardo AM. Whole exome

sequencing in females with autism implicates novel and candidate genes. Int J Mol

Sci 2015, 16: 1312-1335.

[51] Anitha A, Thanseem I, Nakamura K, Yamada K, Iwayama Y, Toyota T*, et al.*

Protocadherin alpha (PCDHA) as a novel susceptibility gene for autism. J Psychiatry

Neurosci 2013, 38: 192-198.

[52] Cappelletti S, Specchio N, Moavero R, Terracciano A, Trivisano M, Pontrelli G,

et al. Cognitive development in females with PCDH19 gene-related epilepsy.

Epilepsy Behav 2015, 42: 36-40.

[53] van Harssel JJ, Weckhuysen S, van Kempen MJ, Hardies K, Verbeek NE, de

Kovel CG, *et al.* Clinical and genetic aspects of PCDH19-related epilepsy syndromes

and the possible role of PCDH19 mutations in males with autism spectrum disorders.

Neurogenetics 2013, 14: 23-34.

[54] Camacho A, Simon R, Sanz R, Vinuela A, Martinez-Salio A, Mateos F.

Cognitive and behavioral profile in females with epilepsy with PDCH19 mutation:

two novel mutations and review of the literature. Epilepsy Behav 2012, 24: 134-137.

[55] Sanders SJ, Ercan-Sencicek AG, Hus V, Luo R, Murtha MT, Moreno-De-Luca D,

et al. Multiple recurrent de novo CNVs, including duplications of the 7q11.23

Williams syndrome region, are strongly associated with autism. Neuron 2011, 70:

863-885.

[56] Pagnamenta AT, Khan H, Walker S, Gerrelli D, Wing K, Bonaglia MC, *et al.*

Rare familial 16q21 microdeletions under a linkage peak implicate cadherin 8 (CDH8)

in susceptibility to autism and learning disability. J Med Genet 2011, 48: 48-54.

[57] Moya PR, Dodman NH, Timpano KR, Rubenstein LM, Rana Z, Fried RL, *et al.*

Rare   missense   neuronal   cadherin   gene   (CDH2)   variants   in   specific

obsessive-compulsive disorder and Tourette disorder phenotypes. Eur J Hum Genet

2013, 21: 850-854.

[58] Redies C, Hertel N, Hubner CA. Cadherins and neuropsychiatric disorders. Brain

Res 2012, 1470: 130-144.

[59] Holroyd S, Reiss AL, Bryan RN. Autistic features in Joubert syndrome: a genetic

disorder with agenesis of the cerebellar vermis. Biol Psychiatry 1991, 29: 287-294.

[60] Feng J, Han Q, Zhou L. Planar cell polarity genes, Celsr1-3, in neural

development. Neurosci Bull 2012, 28: 309-315.

[61] Kim H, Lim CS, Kaang BK. Neuronal mechanisms and circuits underlying

repetitive behaviors in mouse models of autism spectrum disorder. Behav Brain Funct

2016, 12: 3.

[62] Wang C, Pan YH, Wang Y, Blatt G, Yuan XB. Segregated expressions of autism

risk genes *Cdh11* and *Cdh9* in autism-relevant regions of developing cerebellum. Mol

Brain 2019, 12: 40.

[63] Doan RN, Lim ET, De Rubeis S, Betancur C, Cutler DJ, Chiocchetti AG, *et al.*

Recessive gene disruptions in autism spectrum disorder. Nat Genet 2019, 51:

1092-1098.

[64] Gadow KD, Devincent CJ, Pomeroy J, Azizian A. Comparison of DSM-IV

symptoms in elementary school-age children with PDD versus clinic and community

samples. Autism 2005, 9: 392-415.

[65] Simonoff E, Pickles A, Charman T, Chandler S, Loucas T, Baird G. Psychiatric

disorders in children with autism spectrum disorders: prevalence, comorbidity, and

associated factors in a population-derived sample. J Am Acad Child Adolesc Psychiatry 2008, 47: 921-929.

[66] Greene RW, Biederman J, Faraone SV, Ouellette CA, Penn C, Griffin SM. Toward a new psychometric definition of social disability in children with attention-deficit hyperactivity disorder. J Am Acad Child Adolesc Psychiatry 1996, 35: 571-578.

[67] Grzadzinski R, Dick C, Lord C, Bishop S. Parent-reported and clinician-observed autism spectrum disorder (ASD) symptoms in children with attention deficit/hyperactivity disorder (ADHD): implications for practice under DSM-5. Mol Autism 2016, 7: 7.

[68] Williams ME, Wilke SA, Daggett A, Davis E, Otto S, Ravi D*, et al.* Cadherin-9 regulates synapse-specific differentiation in the developing hippocampus. Neuron 2011, 71: 640-655.

[69] Deciphering Developmental Disorders S. Prevalence and architecture of de novo mutations in developmental disorders. Nature 2017, 542: 433-438.

[70] Wang K, Zhang H, Ma D, Bucan M, Glessner JT, Abrahams BS*, et al.* Common genetic variants on 5p14.1 associate with autism spectrum disorders. Nature 2009, 459: 528-533.

[71] Kerin T, Ramanathan A, Rivas K, Grepo N, Coetzee GA, Campbell DB. A Noncoding RNA Antisense to Moesin at 5p14.1 in Autism. Science Translational Medicine 2012, 4(128): 128ra40.

[72] DeWitt JJ, Grepo N, Wilkinson B, Evgrafov OV, Knowles JA, Campbell DB.

Impact of the Autism-Associated Long Noncoding RNA MSNP1AS on Neuronal

Architecture and Gene Expression in Human Neural Progenitor Cells. Genes (Basel)

2016, 7(10): 76.

[73] DeWitt JJ, Hecht PM, Grepo N, Wilkinson B, Evgrafov OV, Morris KV, *et al.*

Transcriptional Gene Silencing of the Autism-Associated Long Noncoding RNA

MSNP1AS in Human Neural Progenitor Cells. Dev Neurosci 2016, 38: 375-383.

[74] Grove J, Ripke S, Als TD, Mattheisen M, Walters RK, Won H, *et al.*

Identification of common genetic risk variants for autism spectrum disorder. Nat

Genet 2019, 51: 431-444.

Legends:

Figure Legends:

Figure 1. Flowchart of the matched gene co-expression analysis (MGCA).

Whole-genome genes were ranked by gene features, creating various gene lists ①.

For each gene in the hcASD gene set (hcASD-1, hcASD-2, ⋯ to hcASD-101), a

feature-matched gene was randomly selected from the range of 50 genes above and 50

genes below (±50 range) this gene in the ranked gene lists ② to generate a matched

random gene set (mRand) ③. For each gene in the whole genome, its co-expression

coefficient (CEC) with each of the mRand gene set (CEC_i) was computed ④, and

100,000 permutations were conducted (i=1,2,…100000). Each CEC_i was compared to

the CEC of the same gene with the hcASD gene set (CEC_{hcASD}). Count the number of

permutations with $CEC_i < CEC_{hcASD}$ (n started from 0 for each gene evaluated, n+1 if

$CEC_i < CEC_{hcASD}$) ⑤. At the end of 100000 permutations, genes with n > 99990,

which corresponded to a permutation *P*-value < 0.0001, were considered significantly

co-expressed with hcASDs under a feature-matched condition ⑥. Genes significantly

co-expressed with hcASDs under all three feature-matched conditions were defined as

Triple-matched genes (TriM) ⑦.

Figure 2. Effect of gene features on genome-wide gene co-expression profiles. A total

of 12,250 genes with information on all 3 gene features were identified, placed in

Table S1 as gene lists, and used for analyses. **A**: Heatmaps of correlation coefficients

(CCs) of genome-wide gene pairs. Genes were ranked according to mRNA abundance,

gDNA size, or GC content. The correlation coefficient (CC) of each gene with all

genes was plotted and displayed in pseudo-color-coded matrices. **B**: Genome-wide

distribution of co-expression coefficients (CECs) of each gene with the hcASD gene

set under three different gene ranking conditions. In each of the 3 matrix panels, the

12,250 genes were placed in ascending order on the x-axis, with 1 being the lowest

mRNA abundance and GC content or shortest in gDNA size and 12,250 being the

highest in mRNA abundance and GC content or the longest in gDNA size. Each blue

dot represents the CEC of a gene with the hcASD gene set. Purple, yellow, green, and

red dots represent noise-reduced (average) CEC of a gene with its neighboring 20 (10

above and 10 below; ± 10), 50 (± 25), 100 (± 50), or 200 (± 100) genes on the gene

lists, respectively. Rods at the bottom of each panel show locations of hcASD genes

on the ranked gene lists. **C**: Genome-wide distribution of CECs of each gene with the

hcASD gene set when genes are placed in stochastic (random) orders.

Figure 3. Convergent expression of hcASD genes determined by MGCA. **A**:

Comparison of noise-reduced CEC distribution curves between the hcASD gene set

and 200 matched random gene sets (mRand) under different gene ranking conditions.

The X-axis represents gene ranks. **B**: CECs of hcASD-hcASD, hcASD-mRand,

mRand-mRand, hcASD-Rand, and Rand-Rand gene set pairs. 200 each of mRand and

Rand gene sets were analyzed. Box plots show ranges of CECs of hcASD-mRand,

mRand-mRand, hcASD-Rand, and Rand-Rand gene set pairs. In each box plot, the

central rectangles span the first quartile to the third quartile of 200 ranked CEC values.

The white bar inside the rectangle shows the median CEC value, and whiskers above

and below the box show the maximum and minimum values, respectively. The dotted

line represents the CEC among hcASDs (hcASD-hcASD) in each panel. Three

statistical methods were used to determine whether the CEC of hcASD-hcASD is

significantly higher than that of hcASD-mRand, mRand-mRand, hcASD-Rand, and

Rand-Rand. Upper fences test: red triangles stand for the boundaries of significant

difference (3 x fence). Grubbs' test: ***$P < 0.001$. Permutation test: ### $P < 0.001$.

The student's *t*-test was used to determine whether the CECs of hcASD-mRand and

mRand-mRand are significantly greater than those of hcASD-Rand and Rand-Rand,

respectively. @ $P < 0.05$; @@@ $P < 0.001$.

Figure 4. Evaluation of TriM genes identified by MGCA. **A**: Schematic of the TriM

gene identification. TriM genes were significantly co-expressed with hcASDs under

all three matched conditions determined by MGCA. **B**: Enrichment of

ASD-associated genes and negative genes in TriM gene sets at different ranges of

P-values. Fold enrichment ($\log_2$) of cASD (left) and "True negative" (right) genes in

different groups of TriM genes are shown. ** $P < 0.01$, *** $P < 0.001$, chi-square test.

C: Three-dimensional distribution of TriM genes ($P < 0.0001$) and an equal number

(2370) of genes with highest CEC with hcASDs (Top). The three axes are CEC value,

gDNA size (bp), and mRNA abundance (RPKM). Each dot in the graph represents a

gene. The TriM-specific genes, Top-specific genes, and overlapped genes are shown

in purple, blue, and black. **D**: Gene Ontology (GO) analysis of TriM ($P < 0.0001$) and

Top (2370) genes. Significantly enriched GO terms are shown ($P < 0.05$,

Benjamini-Hochberg correction). **E**: Comparison of the enrichment of cASD genes

and "True negative" genes in TriM, Top, EWML-identified genes, and

DAWN-identified genes. *** $P < 0.001$, chi-square test.

Figure 5. Open field and elevated plus maze tests of *Cdh11-* and *Cdh9*-null mice. **A**:

Heatmaps showing cumulated frequency of locations visited by *Cdh11*-null mice,

heterozygote (Het), and WT mice in the open field arena. **B, C,** and **D**: Moving

distance, moving speed, and center exploration time of *Cdh11*-null mice. **E, F,** and **G**:

Moving distance, moving speed, and center exploration time of *Cdh9*-null mice. (male

Cdh11-null: n=21, Het: n=22, WT: n=14; female *Cdh11*-null: n=21, Het: n=22, WT:

n=21; male *Cdh9*-null: n=14, Het: n=15, WT: n=12; female *Cdh9*-null: n=8, Het:

n=15, WT: n=12). **J** and **K**: Time spent in open arm and open arm entry frequency of

Cdh11-null mice. **H** and **I**: Time spent in open arm and open arm entry frequency of

Cdh9-null mice (male *Cdh11*-null: n=14, Het: n=14, WT: n=8; female *Cdh11*-null:

n=15, Het n=14, WT n=17; male *Cdh9*-null: n=13, Het: n=15, WT: n=10; female

Cdh9-null: n=9, Het: n=9, WT: n=10). Data are mean ± SEM. Statistical difference

was determined by Student *t*-test ('$P < 0.05$) and by one-way ANOVA followed by

Dunnett's *t*-test (*$P < 0.05$, **$P < 0.01$).

Figure 6. Gripping strength and repetitive behaviors of *Cdh11*- and *Cdh9*-null mice.

A and **B**: Results of gripping test and horizontal bar test for *Cdh11*-null mice (male

Cdh11-null: n=21, Het: n=23, WT: n=14; female *Cdh11*-null: n=12, Het: n=14, WT:

n=12). **C**: Results of gripping test for *Cdh9*-null mice (male *Cdh9*-null: n=11, Het:

n=11, WT: n=7, female *Cdh9*-null: n=5, Het: n=5, WT mice n=4). **D-G**: Latency to

fall (**D, F**) and maximum durable speed (**E, G**) in rotarod test for female *Cdh11*- and

Cdh9-null mice (*Cdh11*-null: n=14, Het: n=14, WT: n=17; *Cdh9*-null: n=10, Het:

n=12; WT: n=9). Numbers below the X-axis (1-5) represent different trials of tests.

H-K: Frequency and duration of self-grooming of female *Cdh11*-null mice during the

first (stage 1, **H, I**) and the second (stage 2, **J, K**) 10 min in the open field arena

(*Cdh11*-null: n=8, Het: n=7, WT n=7). **L** and **M,** Frequency and duration of

self-grooming of female *Cdh9*-null mice during the first (stage 1) and second (stage 2)

10 min in the open filed arena (*Cdh9*-null: n=11, Het: n=17, WT: n=14). Data are

mean ± SEM. Statistical difference was determined by Student *t*-test ('$P < 0.05$) and

by one-way ANOVA followed by Dunnett's *t*-test (*$P < 0.05$, **$P < 0.01$, ***$P <$

0.001, compared to first trial. # $P < 0.05$, ## $P < 0.01$, compared to mice of the WT

littermates).

Figure 7. Modified three-chamber test of female *Cdh11*- and *Cdh9*-null mice. **a**:

Schematics of standard and modified three-chamber tests. **b** and **c**: Results of

sociability and social novelty preference tests of *Cdh11*-null mice (*Cdh11*-null: n=9,

Het: n=8; WT: n=9). **d** and **e**: Results of sociability and novelty preference tests of

Cdh9-null mice (*Cdh9*-null: n=13, Het: n=5, WT: n=10). Data are Mean ± SEM.

Statistical difference was determined by Student *t*-test ('$P < 0.05$) and by one-way

ANOVA followed by Dunnett's *t*-test (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$,

compared to the duration spent in the other side chamber. #$P < 0.05$, compared to WT

littermates).

Supplementary Information

Supplemental_Fig. S1

Distribution of hcASDs in gene rank matrices. Genes in the whole genome were

ranked by gDNA size, GC content (horizontal axis), and mRNA abundance level

(vertical axis) and plotted in matrices. Each blue dot represents a single gene, and

each red dot represents an hcASD gene. Values of both horizontal and vertical axes

are ranks (orders) of the 12,250 genes under 3 different gene ranking conditions.

Supplemental_Fig. S2

Fitting (regression) of CEC distribution curves under three different gene ranking

conditions.

Supplemental_Fig. S3

CEC distribution curves for different developmental stages, sex, and brain regions.

Supplemental_Fig. S4

Genome-wide distribution of CECs of each gene with the hcASD (64) gene set under

three different gene ranking conditions.

Supplemental_Fig. S5

Co-expression of hcASDs in the brain of different developmental stages and sex. **A**

and **B** show data from early (8PCW - 2Y) and late (3Y - 40Y) developmental stages,

respectively. **c** and **d** show data from male and female brain tissues, respectively.

1004

Supplemental_Fig. S6

Co-expression of hcASDs in different brain regions. **A-C:** Box plots show the range

of CECs of hcASD-mRand in different brain regions. **D:** Significant scores of

co-expression of hcASDs in different brain regions. Score = N - R. N: number of

brain regions. R: integrated ranking (average of 3 conditions) of -Log(P-value) in

Grubbs' test under four different gene ranking conditions.

1011

1012 **Supplemental_Fig. S7**

1013 Clustering analysis of –Log(P) values in GO analysis of TriM-only, Top-only,

1014 Overlapped, and combined ASD gene sets (cASDs). –Log(P) values (0 - 20) are

1015 color-coded and shown in the heatmap.

1016

1017 **Supplemental_Table_S1**

1018 List of total genes and hcASDs included in this study.

1019

1020 **Supplemental_Table_S2**

1021 Parameters of fitted curves (regression curves) in supplemental_Fig_S2.

1022

Supplemental_Table_S3

P-value and FDR of each gene under different gene ranking conditions and lists of

TriM, Top, TriM-only, Top-only, and Overlap genes.

1026

Supplemental_Table_S4

GO entries of TriM ($P < 0.0001$) and Top (2370) genes.

1029

Supplemental_Table_S5

Lists of cASDs, True negative genes, EWML genes, and DAWN genes.

1032

Supplemental_Table_S6

List of TriM- and TriN-cadherins

Fig. 1 Flowchart of the matched gene co-expression analysis (MGCA).

Fig. 2 Effect of gene features on genome-wide gene co-expression profiles.

Fig. 3 Convergent expression of hcASD genes determined by MGCA.

Fig. 4 Evaluation of TriM genes identified by MGCA.

Fig. 5 Open field and elevated plus maze tests of *Cdh11*- and *Cdh9*-null mice.

Fig. 6 Gripping strength and repetitive behaviors of *Cdh11*- and *Cdh9*-null mice.

Fig. 7 Modified three-chamber test of female *Cdh11*- and *Cdh9*-null mice.