Running Head: PRX01, PRX44 and PRX73 are extensin-related peroxidases active in root hair growth

Authors for Correspondence:

José M. Estevez

Fundación Instituto Leloir, Av. Patricias Argentinas 435, Buenos Aires C1405BWE, Argentina. TE: 54-

115238-7500 EXT. 3206

Centro de Biotecnología Vegetal, Facultad de Ciencias de la Vida, Universidad Andrés Bello and

Millennium Institute for Integrative Biology (iBio), Santiago CP 8370146, Chile.

Email: jestevez@leloir.org.ar / jose.estevez@unab.cl

10
11
12 **Research area most appropriate for paper**: Plant Biology

13
14
Text Word count: 3,565

Figures 1-5

Experimental procedures

References: 68

RESEARCH REPORT

PRX01, PRX44, and PRX73 are Class-III extensin-related peroxidases that modulates root hair

growth in *Arabidopsis thaliana*

Eliana Marzol[1,#], Cecilia Borassi[1,#], Philippe Ranocha[2], Alejandro Aptekman[3,4], Mauro Brindas[5],

Janice Pennington[6], Julio Paez-Valencia[6], Javier Martínez Pacheco[1], Diana Rosa Rodríguez Garcia[1],

Yossmayer Rondon[1], Mariana Carignani[1], Silvina Mangano[1], Margaret Fleming[7], Francisca Blanco-

Herrera[8,9], Patricia Bedinger[7], Christophe Dunand[2], Luciana Capece[5], Alejandro Nadra[3,4], Michael

Held[10], Marisa Otegui[6,11] & José M. Estevez[1,8,†]

[1]Fundación Instituto Leloir and IIBBA-CONICET. Av. Patricias Argentinas 435, Buenos Aires

C1405BWE, Argentina.

[2]Université de Toulouse, UPS, UMR 5546, Laboratoire de Recherche en Sciences Végétales, F-31326

CNRS, UMR 5546 Castanet-Tolosan, France.

[3]Departamento de Fisiología, Biología Molecular y Celular, Facultad de Ciencias Exactas y Naturales,

Universidad de Buenos Aires, Ciudad Universitaria, Buenos Aires C1428EGA, Argentina.

[4]Departamento de Química Biológica, Facultad de Ciencias Exactas y Naturales, Universidad de

Buenos Aires (IQUIBICEN-CONICET), Ciudad Universitaria, Buenos Aires C1428EGA, Argentina.

[5]Departamento de Química Inorgánica, Analítica y Química Física, Facultad de Ciencias Exactas y

Naturales, Universidad de Buenos Aires (INQUIMAE-CONICET), Buenos Aires, CP. C1428EGA,

Argentina.

[6]Laboratory of Cell and Molecular Biology, University of Wisconsin, Madison, WI, USA.

[7]Department of Biology, Colorado State University, Fort Collins, Colorado 80523-1878, USA.

[8]Centro de Biotecnología Vegetal, Facultad de Ciencias de la Vida, Universidad Andrés Bello and

Millennium Institute for Integrative Biology (iBio), Santiago, Chile.

[9]Center of Applied Ecology and Sustainability (CAPES), Chile.

[10]Department of Chemistry and Biochemistry, Ohio University, Athens, OH 45701, USA.

[11]Departments of Botany and Genetics, University of Wisconsin, Madison, WI, USA.

[#]co-first authors

[†] Correspondence should be addressed. Email: jestevez@leloir.org.ar / jose.estevez@unab.cl (J.M.E).

Abstract

Root hair cells acts at the root surface as important sensors of soil conditions. Root hairs are able to expand several hundred times their original size in a couple of hours to reach and uptake water-soluble nutrients. This fast and oscillating growth requires a continuous remodelling of the cell wall to allow for the root hair to expand in a tightly controlled manner. Root hair cell walls contain polysaccharides and hydroxyproline-rich glycoproteins (HRGPs) including extensins (EXTs). Class-III peroxidases (PRXs) are secreted into the apoplastic space and are thought to trigger either cell wall loosening mediated by oxygen radical species, or polymerization of cell wall components including the Tyr-mediated assembly of an EXT network (EXT-PRXs). The precise role of each of these EXT-PRXs remains to be discovered. By using genetic, biochemical and modeling approaches we have identified and characterized three root hair specific EXT-PRXs, PRX01, PRX44, and PRX73. The triple mutant *prx01,44,73* as well as the overexpressors for PRX44 and PRX73 showed opposite effect on root hair growth, peroxidase activity, ROS-production, cell wall thickness, and Tyr-crosslinks in EXTs. These results suggest a key role of these three EXT-PRXs in the control of root hair growth linked to cell wall properties during polar cell expansion. Finally, modeling and docking calculations have suggested these three EXT-PRXs might be able to interact with non-*O*-glycosylated sections of EXT peptides that reduced Tyr-to-Tyr intra-chain distances in EXT aggregates and might enhance Tyr-crosslinks.

Word count: 236

Introduction

Primary cell walls composed by a diverse network of polysaccharides and structural glycoproteins, regulate cell elongation, which is crucial for several plant growth and developmental processes. Extensins (EXTs) belong to hydroxyproline (Hyp)-rich glycoprotein (HRGP) superfamily and broadly include related glycoproteins such as proline-rich proteins (PRPs) and leucine-rich repeat extensins (LRXs) with multiple Ser-(Pro)$_{3-5}$ repeats that may be *O*-glycosylated and contain Tyr (Y)-based motifs (Marzol et al. 2018). EXTs require several modifications before they become functional (Marzol et al. 2018). First, they are hydroxylated and *O*-glycosylated in the secretory pathway and then, secreted *O*-glycosylated EXTs are crosslinked and insolubilized in the plant cell wall by the activity of secreted class-III peroxidases (PRXs) on the Tyr-based motifs (Lamport et al., 2011; Marzol et al. 2018). PRXs are thought to facilitate both intra and inter-molecular covalent Tyr–Tyr crosslinks in EXT networks possibly thought the assembly of triple helices (Velasquez et al. 2015) by generating *iso*dityrosine units (IDT) and pulcherosine, or di-*iso*dityrosine (Di-IDT), respectively (Brady et al., 1996; 1998). However, the underlying molecular mechanisms of EXT cross-linking and assembly have not yet been determined. The presence of Tyr-mediated crosslinking in EXT and related glycoproteins allow them to form a dendritic glycoprotein network in the cell wall. This network affects *de novo* cell wall formation during embryo development (Cannon *et al.*, 2008), in polar cell expansion processes in root hairs (Baumberger 2001, 2003; Ringli 2010; Velasquez et al. 2011, 2015a,b) and pollen tubes (Sede et al. 2017; Ringli et al. 2017; Wang et al. 2017).

Apoplastic class-III PRXs are heme-iron-dependent proteins, member of a large multigenic family in land plants, with 73 members in *Arabidopsis thaliana* (Passardi et al. 2004; Weng and Chapple, 2010). These PRXs catalyzed several different Classs of reactions. PRXs activities coupled to $_{apo}$ROS molecules ($_{apo}$H$_2$O$_2$) directly affect the degree of cell wall crosslinking (Dunand et al. 2007) by oxidizing cell wall compounds and leading to stiffening of the cell wall through a peroxidative cycle (PC) (Lamport et al. 2011). On the other hand, $_{apo}$ROS coupled to PRX activity enhances non-enzymatic cell wall-loosening by producing oxygen radical species (e.g., $^{\bullet}$OH) and promoting growth in the hydroxylic cycle (HC). It is unclear how opposite effects on cell wall polymers are coordinated during plant growth (Passardi et al. 2004, Cosio & Dunand 2009). Finally, PRXs also contribute to the superoxide radical (O$_2$$^{\bullet-}$) pool by oxidizing singlet oxygen in the oxidative cycle (OC), thereby affecting $_{apo}$H$_2$O$_2$ levels. Thus, several PRXs are involved in the oxidative polymerization of monolignols in the apoplast of the lignifying cells in xylem (e.g. PRX017; Cosio et al 2017), in the root endodermis (e.g. PRX64; Lee et al. 2013; Ropollo et al. 2011), and in petal detachment (Lee et al 2018). In addition, PRXs are also able to polymerize other components of the plant cell wall such as suberin, pectins, and EXTs (Schnabelrauch *et al.*, 1996; Bernards et al., 1999; Jackson et al., 2001; Francoz et al. 2019). Although several candidates of PRXs have been associated specifically to EXT-crosslinking (EXT-PRXs) by *in vitro* studies (Schnabelrauch et al., 1996; Wojtaszek et al., 1997; Jackson et al., 2001; Price et al., 2003; Pereira et al. 2011; Dong et al., 2015; Jacobowitz et al. 2019),

the *in vivo* characterization and mode of action of these EXT-PRXs remain largely unknown. In this work, using a combination of reverse genetics, molecular and cell biology, computational molecular modeling, and biochemistry, to identify three apoplastic PRXs, PRX01, PRX44 and PRX73, as key enzymes for Tyr-crosslinking of cell wall EXTs in growing root hair cells. In addition, we propose a model where *O*-glycosylation levels on the triple helixes of EXTs may regulate the degree of Tyr-crosslinking affecting the expansion properties of cell walls as suggested before (Velasquez et al. 2015a). Our results open the way for the discovery of future similar interactions during root hair development and in response to the environment changes such oscillating nutrients availability in the soil.

Results and Discussion

In this work, we have chosen to analyze root hair cells because they are an excellent model to track cell elongation and to identify PRXs involved in the EXT assembly. Previously, it was suggested that single mutants for *pxr1*, *prx44* and *prx73* were involved in root hair growth and ROS homeostasis (Mangano et al. 2017). All three PRXs are under the transcriptional regulation of the root hair specific transcription factor RSL4 (Menand et al. 2007) although their mechanisms of action remained to be characterized (Mangano et al. 2017). As expected, these three PRXs are also highly co-expressed with other root hair specific genes encoding cell wall EXTs (e.g. EXT6-7, EXT12-14 and EXT18) and EXT-related glycoproteins (e.g. LRX1 and LRX2), which are important for cell expansion (Ringli 2010; Velasquez et al. 2011; Velasquez et al. 2015b) (**Figure S1**). Based on this evidences, we hypothesized that these three PRXs might be EXT-PRXs and perform the Tyr-crosslinks to catalyze the EXT assembly in root hair cell walls.

We first made transcriptional GFP-tagged reporters of PRX01, PRX44 and PRX73 promoter regions to confirm their specific expression in root hairs. In agreement with the *in silico* database (Mangano et al. 2017 and **Figure S1**), all of them were strongly expressed in root hair cells during cell elongation (**Figure 1A**). Single and double combinations of mutants for these three PRXs did not show abnormal root hair growth (Mangano et al. 2017) suggesting a high degree of functional redundancy. Then, we obtained the triple null mutant *PRX01,44,73* (**Figure 1B**) as well as two independent lines overexpressing each of the PRXs fused to GFP and under the strong *35SCaMV* promoter (PRX[OE]) to test whether the expression levels of PRXs affect the normal root hair growth. Whereas the *prx01,44,73* mutant showed much shorter root hair cells (**Figure 1B**) than the previously reported for each of the individual *prx* mutants (Mangano et al. 2017), both independent lines of PRX44[OE] and PRX73[OE] showed significant longer root hairs than the Wt Col-0 control (**Figure 2A-B**). PRX01[OE] lines, however, failed to enhance root hair growth when compared to Wt Col-0 (**Figure 2A-B**). We reasoned that the lack of enhanced root hair expansion in PRX1[OE] could be due to reduced expression compared to the PRX44 [OE] and PRX73 [OE] lines. However, both by detection of GFP signal in intact roots (**Figure 2C**) and immunoblot analysis of protein extracts with anti-GFP

antibodies (**Figure 2D**), we found that PRX1-GFP and PRX44-GFP are strongly expressed whereas
PRX73-GFP showed a more moderate expression; the three PRX GFP-fusion proteins were detected
the expected molecular weight (**Figure 2D**), indicating that the tagged proteins are stable. These
results together highlight partially redundant but not identical roles of PRX1, PRX44 and PRX73 as
positive regulators of polar growth. This is in agreement with the negative effect of SHAM, which
blocks the peroxidase activities of most peroxidases, on root hair growth (Mangano et al. 2017).
Therefore, we measured *in vitro* total peroxidase activity using a guaiacol oxidation-based assay and
found that *prx01,44,73* roots showed reduced peroxidase activity (close to 50% reduction) (**Figure
1C**) while PRXs^{OE} lines showed enhanced activity, with a 40-50% increase in PRX73^{OE} and
approximately 20% increase in PRX44^{OE} (**Figure 2E**). Consistent with our root hair growth analysis
(**Figure 2A**), the PRX01^{OE} lines showed no or moderate increased peroxidase activity (**Figure 2E**). We
then measured cytoplasmic ROS ($_{cyt}$ROS) levels by oxidation of H$_2$DCF-DA and apoplastic ROS
($_{apo}$ROS) levels with the Amplex Ultra Red (AUR) probe in root hair tips. The *prx01,44,73* root hair
tips showed lower levels of $_{cyt}$ROS (**Figure 1D**) but increased $_{apo}$ROS accumulation (**Figure 1E**)
compared to Wt Col-0. The $_{apo}$ROS levels were similar in PRX01,44,73^{OE} lines when compared to Wt
Col-0 (**Figure 2F**). These results suggest a possible role of PRX01, PRX44 and PRX73, as apoplastic
regulators of ROS-linked root hair cell elongation, possibly related to EXT assembly in the cell walls.
These three PRXs consume ROS, most probably H$_2$O$_2$, for its catalytic functions in the cell wall-
apoplast.
168
Next, we designed an EXT reporter to track changes between EXT secretion and insolubilization in
the cell walls mediated by PRXs during root hair cell elongation. The secreted EXT-reporter carries a
Tomato tag (SS-TOM-Long-EXT) that is resistant to acidic pH (Shen et al. 2014) typical of plant cell
walls and apoplastic spaces (Stoddard & Rolland 2018). A secreted Tomato tag (ss-TOM) was used as
a control (**Figure S3A**). EXT expression was first monitored in onion cells (**Figure S3**), and then was
stably expressed in *Arabidopsis* root hairs (**Figure S2C**). In both cases, plasmolysis was used to
retract the plasma membrane from the cell surface showing that the EXT-reporter was localized in
the cell walls. By performing western blot analysis, we detected the full length EXT-Tomato fusion
protein with possible *O*-glycans modifications running as higher molecular weight bands; the control
SS-TOM was detected at the expected size (**Figure S2B**). More importantly, the EXT-reporter did not
interfere with the polar growth of root hairs (**Figure S2D**), and therefore, was used it to track
changes in the *in situ* arrangement of cell wall EXTs. Then, we tested if SHAM, a peroxidase inhibitor
known to impair root hair growth (Mangano et al. 2017), might affect the secretion and assembly of
SS-TOM-Long-EXT into the new formed apical root hair cell wall (**Figure 3A-B**). SHAM treatment not
only disrupted polar cell expansion in root hairs as previously shown (Mangano et al. 2017) but
more importantly, triggered an abnormal EXT deposition in the apical cell walls suggesting that Tyr-
crosslinks might be affected (**Figure ?**). To further analyze the ultrastructure of the cell wall in
growing root hairs, we analyzed Wt Col, PRX44^{OE}, and *prx01,44,73* triple mutant roots treated with

or without SHAM by transmission electron microscopy (**Figure 3A**). Whereas there was no statistically significant difference between cell wall thickness at the root hair tips of PRX44[OE] and Wt Col-0 plants (1.2 ±0.3 µm s for Wt and 1.05 ±0.11 µm for PRX44[OE]), the *prx01,44,73* root hairs had much thinner cell walls (0.4 ±0.4 µm) (**Figure 3B**). In the three genotypes, SHAM treatment caused a slight increase in cell wall thickness that was only statistically significant in the PRX44[OE] root hairs (**Figure 3B**).

We then measured the levels of EXT cross-links at Tyr units in isolated whole roots by measuring peptidyl-tyrosine (Tyr) and iso-dityrosine (IDT) (dimer of Tyr). We did detect significant differences in the peptidyl-Tyr between Wt Col-0 and *prx01,44,73* triple mutant while slightly higher levels on IDT in extracted EXTs from PRX73[OE] line (**Table S1**). On the other hand, strong down-regulation of Tyr- and IDT-levels were found in miss-glycosylated EXTs mutants like *p4h5 sergt1-1* and *sergt1-1 rra3* (**Table S1**) suggesting a strong impact of *O*-glycans on EXT Tyr crosslinking as previously suggested based on their drastic reduced root hair growth (Velasquez et al 2015a). We hypothesize that absent or low *O*-glycosylation levels on EXTs or a change in PRX levels may trigger a reduction in the amount of peptidyl-Tyr and IDT on EXTs with a putative concomitant increase in the amounts of higher Tyr-crosslink units (trimers and tetramers) thus impacting on root hair growth. Further research is needed to understand the *in vivo* regulation of Tyr-crosslinking of EXTs by PRXs in plant cells.

One of the major limitations in our understanding of how EXT function in the plant cell walls is the lack of a realistic full-length EXT protein model. Previously, we have generated preliminary versions of a short EXT triple helix model (Velasquez et al. 2015a). To have a more detailed understanding of how EXT molecules might behave in the plant cell wall, we have built a larger model of a triple helix EXT sequence that includes 10 conserved repeats (SPPPPYVYSSPPPPYYSPSPKVYYK) with a total length of 750 amino acids (250 in each polypeptide chain) by using a coarse grained molecular dynamics approach (**Figure S4A-B**). Parameters for the EXT *O*-glycosylated form were developed in this work and we were able to simulate the EXT *O*-glycosylated system (**Figure S5**). The EXT molecules were modeled in two different states: as a non-glycosylated trimeric helical conformation similar to animal collagen as performed before (Velasquez et al., 2015a) and in the fully *O*-glycosylated state. In addition, we generated molecular dynamics simulations restraining both ends of the polypeptide chains, to model a fully extended helix (consistent with an "indefinitely long EXT"), and without that restriction, to evaluate the conformation that an isolated 10 repeats triple helix would adopt. The results obtained in these simulations indicate the importance of the triple helix conformation in the overall protein stability and especially the conservation of the fibril-like structure, in agreement with 10 repeat single helix simulations performed previously (Marzol et al. 2018). Total volume of the extended 10 repeats triple helix was measured in both glycosylation states (**Table 1**), differentiating EXT-protein + *O*-glycans and EXT-protein-only volumes for the fully

O-glycosylated EXT state. We observe that EXT-protein-only volume is significantly augmented by
the presence of the oligosaccharide moieties, indicating that *O*-glycans increase the separation
between peptide chains in the EXT triple helix. We also report average diameter for those systems,
which are consistent with previously reported atomic force microscopy images (Cannon et al. 2008).
Additionally, *O*-glycosylation contributes to increase the average distance between the side chains
of tyrosine residues, decreasing the population of tyrosine side chains in close proximity that could
lead to cross-linked EXT chains (**Figure S4C**). Current experimental and modeling lines of evidence
are in agreement with the proposed role of the proline-hydroxylation and carbohydrate moieties in
keeping the EXT molecule in an extended helical polyproline-II conformation state (Stafstrom &
Staehelin 1986; Owen et al., 2010; Ishiwata et al., 2014). This extended conformation might allow
EXTs to interact properly with themselves and with the surrounding apoplast environment, including
PRXs and pectins, to form a proper cell wall network (Nuñez et al., 2009; Valentin et al., 2010).
237
To test if the three identified PRXs (PRX01, PRX44 and PRX73) might be able to interact directly with
single chain of EXTs, an analysis by homology modeling with a known EXT-PRX GvP1 (from *Vitis*
vinífera) that is able to crosslink EXTs *in vitro*, was performed (Jackson et al., 2001; Pereira et al.
2011). As additional controls, we selected PRX64 described as necessary for the lignin
polymerization of capsarian strips in the *Arabidopsis* root endodermis (Lee et al.2013) and PRX36
able to bind homogalacturonan pectin in the outer seed coat layer(Francoz et al. 2019). By *docking*
analysis, the interaction energy (Kcal/mol) obtained with the three PRXs (PRX01, PRX44, and PRX73)
was compared to EXT-PRX GvP1 and with PRX64/PRX36 using 4 different short EXT peptides. A non-
hydroxylated EXT peptide (SPPPYVY)$_3$, a hydroxylated but not *O*-glycosylated EXT peptide
[(SOOOYVY)$_3$; O=hydroxyproline], only arabinosylated EXT-peptide [(SOOOYVY)$_3$-A] and arabino-
galactosylated EXT peptide [(SOOOYVY)$_3$-AG] were analyzed. Previously, it was shown that mutants
carrying under-*O*-glycosylated EXTs have severe defect in root hair growth (Velasquez et al. 2011;
Velasquez et al. 2015a), and at least for PROLYL 4-HYDROXYLASE 5 mutant (*p4h5*), a modified cell
wall structure in the root hair tip was detected (Velasquez et al. 2015b). Docking results for different
PRXs tested with various degrees of *O*-glycosylation on EXT peptides show a consistent interaction
energy difference depending on the glycosylation state, being higher for non-*O*-glycosylated species
(both as proline or hydroxyproline). In addition, EXT *O*-glycosylated variants dock in a rather
disperse way while those non-*O*-glycosylated do preferentially in a groove area (**Figure 4A-C**).
Furthermore, in **Figure 4A** it is shown how a non-*O*-glycosylated peptide binds through a groove
leaving one tyrosine docked in a cavity open to the solvent and very close to the heme iron (5Å),
with a second tyrosine a few Angstroms away. This disposition and distances suggest that this could
be an active site where the Tyr cross-linking takes place. This statement is reinforced by the fact that
almost every substrate docked to the protein (even variably *O*-glycosylated peptides) bind to this
pocket. Direct comparison of interaction energies derived from docking experiments with different
species would not be rigorous due to the different size and degree of freedom (depending on the *O*-

glycosylation) among docking runs (**Figure 4B**). Despite of the fine values, a trend can be observed in
Figure 4C. When interaction energies are compared among different PRXs interacting with EXT-
substrates with the same degree of *O*-glycosylation, it is observed that interaction follows this
decreasing order PRX73>PRX01>PRX44 for the non-hydroxylated (Ser-Pro-Pro-Pro) substrate while
PRX44>PRX73>PRX01 for the hydroxylated, non-*O*-cosylatied variant (Ser-Hyp-Hyp-Hyp). In
general, it is observed a higher interaction energy (more negative values) for hydroxylated species,
followed by non-hydroxylated, and then by *O*-glycosylated variants. Among the non-hydroxylated
species, the protein displaying relatively higher interaction energy is PRX73. Among the *O*-
glycosylated species, the protein displaying relatively higher interaction energy is PRX44. All
together, these results are consistent with a constitutive root hair growth effect for PRX44[OE] and
PRX73[OE] and also with the non-glycosylated species being the substrates of peroxidation. Overall,
this may indicate that PRX44 and PRX73 might interact with EXT substrates, and possibly catalyze
Tyr-crosslinking in open regions of the EXT backbones that are naked or with low levels of *O*-glycans.
This is in agreement with previous findings suggesting that high levels of *O*-glycosylation in certain
EXT segments were found to physically restrict the EXT lateral alignments, possibly by acting as a
branching point (Velasquez et al., 2015a). Overall, taking together all the approaches used in this
study, these results show that PRX01, PRX44 and PRX73 might be able to crosslink EXTs at Tyr-sites.
280
To examine the evolution of these PRXs, PRX01, PRX44, and PRX73, we performed comprehensive
phylogenetic analyses of class-III peroxidases across diverse organisms from land plant lineages.
Under low selective pressure to maintain substrate specificity, EXT-PRXs activities might have
evolved multiple times in parallel during land plant evolution through gene duplication followed by
neofunctionalization or subfunctionalization. PRX01, PRX44, and PRX73 belonged to three
independent orthologous groups (**Figure S6**) and orthologs genes for each *A. thaliana* PRX have
been detected in available Brassicaceae genomes and in various Angiosperms and Gymnosperms
families but they were missing from Lycophytes and from non-vascular land plants. It means that
these three PRX sequences were the result of ancestral duplications before the divergence between
Gymnosperms and Angiosperms but after the emergence of the Tracheophytes (**Figure S6**). Then,
orthologs of the three PRX genes have only been detected in true-root containing organisms and
these three PRXs are expressed in roots and root hairs, as most of the orthologous sequences (when
expression data are available) (**Figure S7**). This strongly supports conserved functions in roots
among the three independent orthogroups. Except for PRX73 which belonged to a cluster containing
the EXT-PRX from tomato (LePRX38), the other 2 PRXs sequences did not cluster with sequences
already described as putative EXT-PRXs from the literature such as PRX09 and PRX40 (Jacobowitz et
al. 2019). Indeed, the other known EXT-PRXs (mostly by *in vitro* evidences) are not clustered
together but are largely distributed in the tree (**Figure S6**). This analysis suggests that plant EXT-PRXs
might have evolved several times in parallel during Tracheophytes evolution.
300

Based on the results show in this work, we proposed a working model where these three PRXs, PRX01, PRX44, and PRX73 (and possibly others PRXs) might control root hair growth by channelling H_2O_2 consumption into EXT-crosslinking and impacting on cell wall hardening process. In addition, high levels of ROS in the apical zone of the cytoplasm might activate many other processes also related to polar growth (**Figure S8**). On the contrary, when low levels of PRX proteins linked to a reduced peroxidase activity (as in the triple mutant *prx01,44,73* or SHAM-treated roots), high levels of ROS (as H_2O_2) might accumulate in the apoplast triggering through the oxidative cycle a cell wall loosening effect inhibiting cell growth (**Figure S8**). Concomitantly, deficient PRX activity in the apoplast also triggers lower ROS levels in the cytoplasm of growing root hairs impact on root hair growth. It is hypothesized that high levels of apoplastic ROS may inhibit putative PIP-mediated channelling of ROS (as H_2O_2) to the cytoplasm but this remains to be tested in future experiments.

Conclusions

Currently, several of the 73 apoplastic class-III PRXs in *Arabidopsis thaliana* have no assigned biological functions. In this work, we have characterized three EXT-PRXs, PRX01, PRX44 and PRX73 that are important for root hair growth linked to ROS-homeostasis and EXT-assembly. These PRXs might control Tyr-crosslinking in EXTs and related glycoproteins and modify its secretion and assembly in the nascent tip cell walls. By using modeling and docking approaches, it was possible to measure the interactions of these PRXs with single chain EXT substrates. All these evidences highlight that PRX01, PRX44 and PRX73 are important enzymes in the EXTs assembly during root hair growth. From the evolutionary point of view, all the putative EXT-PRXs (mostly by *in vitro* evidences) do not clustered together in the phylogenetic tree of class-III PRXs suggesting that plant EXT-PRXs might have evolved several times in parallel during Traqueophytes evolution. Interestingly, as a convergent evolutionary extracellular assembly, proline-rich collagen Class-IV similar to the green EXTs linage and related glycoproteins, is also cross-linked by the activity of a specific class of animal heme-peroxidases (named peroxidasin or PXDN) to form insoluble extracellular networks (Vanacore et al. 2009; Bhave et al. 2012). While collagen IV biophysical properties allows the correct development and function of multicellular tissues in all animal phyla (Brown et al. 2017), EXTs assemblies are also key in several processes of plant cell expansion and morphogenesis (Cannon *et al.*, 2008; Velasquez et al. 2015b; Marzol et al. 2018). This might imply that cross-linked extracellular matrices based on proline-rich polymers (e.g. collagens and EXTs) have evolved more than once along eukaryotic evolution to provide mechanical support to single and multiple cellular tissues. Further analyses are required in the future to understand how these EXT-PRXs catalyze at the molecular level the Tyr-crosslinks on EXTs and how this assembly process is regulated during polar cell expansion. In this study we characterized changes in ROS homeostasis and EXT-crosslinking although these are two aspects of a much more complex and boarder balance in these fast growing cells.

Experimental Procedures

Plant and growth conditions. *Arabidopsis thaliana* Columbia-0 (Col-0) was used as the wild Class (Wt) genoClass in all experiments. All mutants and transgenic lines tested are in this ecoClass. Seedlings were germinated on agar plates in a Percival incubator at 22°C in a growth room with 16h light/8h dark cycles for 10 days at 140 µmol m^{-2}s^{-1} light intensity. Plants were transferred to soil for growth under the same conditions. For identification of T-DNA knockout lines, genomic DNA was extracted from rosette leaves. Confirmation by PCR of a single and multiple T-DNA insertions in the target PRX genes were performed using an insertion-specific LBb1 or LBb1.3 (for SALK lines) primer in addition to one gene-specific primer. To ensure gene disruptions, PCR was also run using two gene-specific primers, expecting bands corresponding to fragments larger than in WT. We isolated homozygous lines for PRX01 (AT1G05240, *prx01-2*, Salk_103597), PRX44 (AT4G26010, *prx44-2*, Salk_057222) and PRX73 (AT5G67400, *prx73-3*, Salk_009296).

SS-TOM and SS-TOM-Long-EXT constructs. The binary vector pART27136 encoding the secreted tdTomato using the secretory signal sequence from tomato polygalacturonase, under the control of the CaMV 35S promoter, were kind gifts of Dr. Jocelyn Rose, Cornell University (named pART-SS-TOM). The entire reporter protein construct was excised from pART-SS-TOM by digesting with *Not*I. The resulting fragments were gel-purified with the QIAquick Gel Extraction Kit and ligated using T4 DNA Ligase (New England Biolabs) into dephosphorylated pBlueScript KS+ that had also been digested with *Not*I and gel-purified to make pBS-SS-TOM. The plasmids were confirmed by sequencing with primers 35S-FP (5'-CCTTCGCAAGACCCTTCCTC-3') and OCS-RP (5'-CGTGCACAACAGAATTGAAAGC-3'). The Long-EXT domain sequences from *SlPEX1* (NCBI accession AF159296128) was then removed from pUC57-high and subcloned into pUC57-less by digestion of both pUC57-high and pUC57-less with *Pst*I and *Nsi*I followed by gel purification and ligation of the two gel-purified fragments to make pUC57-full length. The plasmid pBS-SS-TOM-Long EXT was made by digestion of pUC57-full length and pBS-SS-TOM with *Nde*I and *SgrA*I, followed by gel purification band from pUC57-full length and the 5545 bp band from pBS-SS-TOM, and ligation of the two gel purified fragments. pBS-SS-TOM-Long-EXT plasmid were confirmed by sequencing. All pBS plasmids were also confirmed by digestion. Binary vectors for plant transformation were made by gel purifying the *Not*I insert fragments from the pBS-sec-tdT-long plasmids and ligating them with pART27-SS-TOM backbone that had been digested with *Not*I, gel purified, and dephosphorylated, to obtain pART27-SS-TOM and pART27-SS-TOM-Long-EXT.

Root hair phenoClass. For quantitative analysis of root hair phenoClasss in *PRX01,44,73* mutants, 35S:PRX-GFP lines and Wt Col-0, 200 fully elongated root hairs were measured (n roots= 20-30) from seedlings grown on vertical plates for 10 days. Values are reported as the mean ±SD using the Image

J software. Measurements were made after 7 days. Images were captured with an Olympus SZX7 Zoom microscope equipped with a Q-Colors digital camera.

Confocal imaging. Root hairs were ratio imaged with the Zeiss LSM 710 laser scanning confocal microscope (Carl Zeiss) using a 40X oil-immersion, 1.2 numerical aperture. EGFP (473–505nm) emission was collected using a 458-nm primary dichroic mirror and the Meta-detector of the microscope. Bright-field images were acquired simultaneously using the transmission detector of the microscope. For time-lapse analysis, images were collected every 3 s. Image sequences were analyzed using the Template Matching and Slice Alignment plug-in for ImageJ. Fluorescence intensity was measured in 7 µm ROI at the root hair apex.

Peroxidase activity. Soluble proteins were extracted from roots grown on vertical plates for 10 days by grinding in 20mM HEPES, pH 7.0, containing 1 mM EGTA, 10mM ascorbic acid, and PVP PolyclarAT (100mg g^{-1} fresh material; Sigma, Buchs, Switzerland). The extract was centrifuged twice for 10 min at 10,000 g. Each extract was assayed for protein levels with the Bio-Rad assay (Bio-Rad) and for total peroxidase activity using guaïacol/H_2O_2. Enzyme activity (expressed in nkatal/mg protein) was determined by a guaiacol oxidation-based assay. Values are the mean of three replicates ± SD. P-value of one-way anova, (**) P<0.001. PRX activity was measured at 25°C by following the oxidation of 8 mM guaiacol (Fluka) at 470 nm in the presence of 2 mM H_2O_2 (Carlo Erba) in a phosphate buffer (200 mM, pH6.0).

Cytoplasmic ROS ($_{cyt}$ROS) measurements in root hairs. Growth of Arabidopsis seeds on a plate was done with 1% sterile agar for 8 d in a chamber at 22°C with continuous light. These seedlings were incubated in darkness on a slide for 10 min with 50 µM 2',7'-dichlorodihydrofluorescein diacetate (H_2DCF-DA) at room temperature. Samples were observed with a confocal microscope equipped with a 488-nm argon laser and BA510IF filter sets. A 10× objective was used, 0.30 N.A., 4.7 laser intensity, 1.1 offset, 440 photomultiplier (PMT) (for highest ROS levels), 480 PMT (for ROS media), and gain 3. Images were taken scanning XZY with 2 µm between focal planes. Images were analyzed using ImageJ. To measure ROS highest levels, a circular region of interest (ROI) (r = 2.5) was chosen in the zone of the root hair with the highest intensities. To measure ROS mean, the total area of the root hair was taken.

Apoplastic ROS ($_{apo}$ROS) measurements in root hairs. Levels of apoplastic H_2O_2 in root hair cells were measured using a solution of Amplex™ UltraRed Reagent (AUR, Molecular Probes). Briefly, Arabidopsis seedlings were incubated with 50 µM AUR for 20 min in dark conditions. Stained root hair cells were imaged with a Zeiss LSM5 Pascal laser scanning confocal microscope. The fluorescence emission of oxidized AUR in the apoplast of root hair cells was observed between 585

and 610 nm using 543 nm argon laser excitation, 40X objective, N/A= 1.2. Quantification of the AUR
probing fluorescence signal was restricted to apoplastic spaces at the root hair tip.

Phylogenetic analysis. 73 class-III PRX protein sequences from *A. thaliana*, two putative lignin class-
III PRXs from *Zinnia elegans* and 4 putative Extensin class-III PRXs from *Lupinus album*, *Lycopersicum*
esculentum, *Phaseolus vulgaris* and *Vitis vinifera*, have been aligned with ClustalW and the tree
constructed using the Neighbor-Joining method (Saitou and Nei, 1987). The analyses were
conducted in MEGA7 (Kumar, 2016). All protein sequences are available using their ID number
(http://peroxibase.toulouse.inra.fr (Savelli et al., 2019).

Co-expression analysis network. Co-expression networks for *RSL4* root hair genes were identified
from PlaNet (http://aranet.mpimp-golm.mpg.de) and trimmed to facilitate readability (Mutwill et al.
2011). Each co-expression of interest was confirmed independently using the expression angler tool
from Botany Array Resource BAR (http://bar.utoronto.ca/ntools/cgi-
bin/ntools_expression_angler.cgi) and ATTED-II (http://atted.jp). Only those genes that are
connected with genes of interest are included.

Tyr-crosslinking analysis. Root tissues obtained from *PRX01,44,73* mutants, Col-0 and 35S:PRX-GFP
lines were hydrolyzed in 6 N HCl (aqueous) with 10 mM phenol (2 mg ml^{-1}; 110 °C; 20 h).
Hydrolysates were dried under a steady stream of nitrogen (gas) and then re-dissolved at 10 mg ml^{-1}
in water. The hydrolysates were fractionated by gel permeation chromatography on a
polyhydroxyethyl A column (inner diameter, 9.4 x 200 mm, 10 nm pore size, Poly LC Inc., Columbia,
MD) equilibrated in 50 mM formic acid and eluted isocratically at a flow rate of 0.8 ml min^{-1}. UV
absorbance was monitored at 280 nm. Phenylisothiocyanate derivatives of IDT were quantified in
hydrolysates after fractionation by gel permeation chromatography. The amounts of Tyr and IDT in
the hydrolysates were determined by comparison with authentic Tyr and IDT standards. Response
factors were determined from three level calibrations with the Tyr and IDT standards.

Western-Blot Analysis. Plant material (100 mg of root from 15 days old seedlings) was collected in a
microfuge tube and ground in liquid nitrogen with 400 mL of protein extracction buffer (125 mM
Tris-Cl, pH. 4.4, 2% [w/v] SDS, 10% [v/v] glycerol, 6M UREA, 1% [v/v] b-mercaptoethanol, 1mM
PMSF). Samples were immediately transferred to ice. After 4° centrifugation at 13000 rpm for 20
minutes, supernatant was move to a new 1.5 ml tube and equal volumes of Laemmli buffer (125
mM Tris-Cl, pH. 7.4, 4% [w/v] SDS, 40% [v/v] glycerol, 10% [v/v] b-mercaptoethanol, 0.002% [w/v]
bromphenol blue) were added. The samples (0.5–1.0 mg/mL of protein) were boiled for 5 min and
30 mL were loaded on 10% SDS-PAGE. The proteins were separated by electrophoresis and
transferred to nitrocellulose membranes. Anti-GFP mouse IgG (clones 7.1 and 13.1; Roche Applied
Science) was used at a dilution of 1:2,000 and it was visualized by incubation with goat anti-mouse

IgG secondary antibodies conjugated to horseradish peroxidase (1:2,000) followed by a
chemiluminescence reaction (Clarity Western ECL Substrate; Bio-rad).

Transmission electron microscopy of root hair cell walls. Seeds were germinated on 0.2x MS, 1%
sucrose, 0.8% agar. Seven days after germination, seedlings were transferred to new 0.2x MS, 1%
sucrose, 0.8% agar plates with or without 100 µM SHAM. After 4 additional days, 1-mm root
segments with root hairs were fix in 2% glutaraldehyde in 0.1M cacodylate buffer pH7.4. Samples
were rinsed in cacodylate buffer and post-fixed in 2% OsO4. After dehydration in ethanol and
acetone, samples were infiltrated in Epon resin (Ted Pella, Redding, CA). Polymerization was
performed at 60°C. Sections were stained with 2% uranyl acetate in 70% methanol followed by
Reynold's lead citrate (2.6% lead nitrate and 3.5% sodium citrate [pH 12.0]) and observed in a Tecnai
12 electron microscope. Quantitative analysis of cell wall thickness was performed using FIJI.

Modeling and molecular docking between PRXs and EXTs. Modeling and molecular docking: cDNA
sequences of PRXs were retrieved from TAIR (PRX01: AT1G05240, PRX36: AT3G50990, PRX44:
AT4G26010, PRX64: AT5G42180, PRX73: AT5G67400) and NCBI Nucleotide DB (PRX24Gv:Vitis
vinifera peroxidase 24, LOC100254434). Homology modeling was performed for all PRXs using
modeller 9.14 (Sali et al. 1993), using the crystal structures 1PA2, 3HDL, 1QO4 and 1HCH as
templates, available at the protein data bank. 100 structures where generated for each protein and
the best scoring one (according to DOPE score) was picked. The *receptor* for the docking runs was
generated by the prepare_receptor4 script from autodock suite, adding hydrogens and constructing
bonds. Peptides based on the sequence PYYSPSPKVYYPPPSSYVYPPPPS were used, replacing proline
by hydroxyproline, and/or adding *O*-Hyp glycosylation with up to four arabinoses per hydroxyproline
in the fully glycosylated peptide and a galactose on the serine, as it is usual in plant *O*-Hyp
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5045529/. Ligand starting structure was generated
as the most stable structure by molecular dynamics by Hugo Verli (Velasquez et al. 2015a). All ligand
bonds were set to be able to rotate. Docking was performed in two steps, using Autodock vina (Trott
et al. 2010). First, an exploratory search over the whole protein surface (exhaustiveness 4) was
done, followed by a more exhaustive one (exhaustiveness 8), reducing the search space to a
75x75x75 box centered over the most frequent binding site found in the former run.

EXT conformational coarse-grained model. The use of coarse-grained (CG) molecular dynamics
(MD) allowed collection of long timescale trajectories. System reduction is significant when
compared to all atom models, approximately reducing on order of magnitude in particle number. In
addition, a longer integration time step can be used. Protein residues and coarse grained solvent
parameters correspond to the SIRAH model (Darre et al. 2014), while ad hoc specific glycan
parameters were developed (**Figure S5**). The CG force field parameters developed correspond to
arabinofuranose and galactopyranose. Triple helix systems were simulated both, in the non-

glycosylated and fully *O*-glycosylated states, where all the hydroxyprolines are bound to a tetrasaccharide of arabinofuranoses, and specific serine residues contain one galactopyranose molecule. They were immersed in WT4 GC solvent box that was constructed to be 2 nm apart from the extensin fiber, and periodic boundary conditions were employed. Coarse grained ions were also included to achieve electroneutrality and 0.15 M ionic strength. All simulations were performed using the GROMACS MD package at constant temperature and pressure, using the Berendsen thermostat (respectively) and Parrinello-Rahman barostat (Parrinello and Rahman 1981), and a 10 fs time step. The obtained trajectories were analysed using the Mdtraj python package (McGibbon et al, 2015) and visualized with Visual Molecular Dynamics (VMD) 1.9.1 (Humphrey et al. 1996). Volume measurements were performed using a Convex Hull algorithm implemented in NumPy (Oliphant 2006), and average diameter calculations were derived from this quantity using simple geometric arguments.

Acknowledgements

We would like to thank to Andres Rossi from the microscopy facility of FIL for his assistance. We thank ABRC (Ohio State University) for providing T-DNA lines seed lines. J.M.E. is Principal Investigator of the National Research Council (CONICET) from Argentina. This work was supported by grants from ANPCyT (PICT2016-0132 and PICT2017-0066) and ICGEB CRP/ARG16-03 and a grant from ICGEB CRP/ARG16-03 and ANPCyT (PICT2016-0132 and PICT2017-0066) to J.M.E. In addition, this research is also funded by Instituto Milenio iBio – Iniciativa Científica Milenio, MINECON to J.M.E. and NSF MCB grant 1614965 to M.S.O.

Author Contribution

E.M and C.B performed most of the experiments and analysed the data. P.R. and C.D. analysed the peroxidase activity and performed phylogenetic analysis. M.H. analysed the Tyr-crosslinking on EXTs. A.A. and A.D.N performed the docking experiments and analysed this data. M.B. and L.C. perform the EXT modelling and analysed this data. M.F. and P.B generated the EXT reporter lines and performed the western blots. J.M.P., D.R.R.M., Y.R., S.M., and F.B.H. analysed the data. J.P., J.P-V., and M.S.O. performed the transmission electron microscopy analysis. J.M.E. designed research, analysed the data, supervised the project, and wrote the paper. All authors commented on the results and the manuscript. This manuscript has not been published and is not under consideration for publication elsewhere. All the authors have read the manuscript and have approved this submission.

Competing financial interest

The authors declare no competing financial interests. Correspondence and requests for materials should be addressed to J.M.E. (Email: jestevez@leloir.org.ar / jose.estevez@unab.cl).

REFERENCES

Abraham, M.J., Murtola, T., Schulz, R., Páll S., Smith, J.C., Hess, B., and Lindahl E. (2015). GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *Software X*, 1–2,19-25. ISSN 2352-7110.

Baumberger N, Ringli C, Keller B. (2001). The chimeric leucine rich repeat/extensin cell wall protein LRX1 is required for root hair morphogenesis in *Arabidopsis thaliana*. *Genes & Development* 15, 1128–1139.

Baumberger N, Steiner M, Ryser U, Keller B, Ringli C. (2003). Synergistic interaction of the two paralogous Arabidopsis genes LRX1 and LRX2 in cell wall formation during root hair development. *Plant J.* 35, 71–81.

Berardini T.Z., Reiser L., Li D., Mezheritsky Y., Muller R., Strait E., Huala E. (2015). The arabidopsis information resource: making and mining the "gold standard" annotated reference plant genome. *Genesis* 53(8):474–485.

Berendsen, H. J. C., Postma, J.P.M., van Gunsteren, W.F., DiNola, A., and Haak, J.R. (1984) Molecular dynamics with coupling to an external bath. *J. Chem. Phys.* 81, 3684. doi: 10.1063/1.448118.

Berman HM, John Westbrook, Zukang Feng, Gary Gilliland, Talapady N Bhat, Helge Weissig, Ilya N Shindyalov, and Philip E Bourne. The protein data bank. *Nucleic Acids Research*, 28(1):235–242, 2000.

Bernards, M.A., Fleming, W.D., Llewellyn, D.B., Priefer, R., Yang, X., Sabatino, A., and Plourde, G.L. (1999). Biochemical characterization of the suberization-associated anionic peroxidase of potato. *Plant Physiol.* 121: 135–146.

Bhave G, Cummings CF, Vanacore RM, Kumagai-Cresse C, Ero-Tolliver IA, Rafi M, Kang J-S, Pedchenko V, Fessler LI, Fessler JH, Hudson BG (2012) Peroxidasin forms sulfilimine chemical bonds using hypohalous acids in tissue genesis. Nat Chem Biol 8:784–790.

Brady, J.D., Sadler, I.H., and Fry, S.C. (1996). Di-isodityrosine, a novel tetrametric derivative of tyrosine in plant cell wall proteins: a new potential cross-link. *Biochem. J* 315: 323–327.

BrownK.L, C.F. Cummings, R.M. Vanacore and B.G. Hudson. (2017). Building collagen IV smart scaffolds on the outside of cells. *Protein Sci.* 26:2151-2161.

Brownleader, M.D., Ahmed, N., Trevan, M., Chaplin, M.F., and Dey, P.M. (1995). Purification and Partial Characterization of Tomato Extensin Peroxidase. *Plant Physiol.* 109: 1115–1123

Cannon, M.C., Terneus, K., Hall, Q., Tan, L., Wang, Y., Wegenhart, B.L., Chen, L., Lamport, D.T., Chen, Y., and Kieliszewski, M.J. (2008). Self-assembly of the plant cell wall requires an extensin scaffold. *Proc. Natl. Acad. Sci. USA.* 105: 2226–2231.

Case, D.A., Cerutti. D.S., Cheatham, T.E.III., Darden, T.A., Duke, R.E., Giese, T.J., Gohlke, H., Goetz, A.W., Greene, D., Homeyer, N., et al. (2017), AMBER 2017, University of California, San Francisco.

Cosio C. & Dunand C. (2009). Specific functions of individual classIII peroxidases genes. *J. Exp. Bot.* 62:391–408.

Cosio C., Ranocha Ph., Francoz E., Burlat V., Zheng Y., Perry S.E., Ripoll J.-J., Yanofsky M., Dunand C. (2017) The class III peroxidase PRX017 is a direct target of the MADS-box transcription factor 1 AGL15 and participates in lignified tissue formation. *New Phytol*. 213: 250-263.

Darré, L., Machado, M.R., Brandner, A.F., González, H.C., Ferreira, S., and Pantano, S. (2015). SIRAH:
A Structurally Unbiased Coarse-Grained Force Field for Proteins with Aqueous Solvation and
Long-Range Electrostatics. *J. Chem. Theory Comput.* 11(2):723–739

Darré, L., Tek, A., Baaden, M., and Pantano, S. (2012). Mixing atomistic and coarse grain solvation
models for MD simulations: let WT4 handle the bulk. *J. Chem. Theory Comput.* 8:3880–3894.

Dong, W., Kieliszewski, M., and Held, M.A. (2015). Identification of the pI 4.6 extensin peroxidase
from *Lycopersicon esculentum* using proteomics and reverse-genomics. *Phytochem.* 112:151–
159.

Dunand, C., Crevecoeur, M., and Penel, C. (2007). Distribution of superoxide and hydrogen peroxide
in Arabidopsis root and their influence on root development: possible interaction with
peroxidases. *New Phytol*. 174:332–341.

Fabrice, T., Vogler, H., Draeger, C., Munglani, G., Gupta, S., Herger, A.G., Knox, P., Grossniklaus, U.,
and Ringli, C. (2018). LRX proteins play a crucial role in pollen grain and pollen tube cell wall
development. Plant Physiol. 176, 1981–1992.

Francoz et al. (2019) Pectin demethylesterification generates platforms that anchor peroxidases to
remodel plant cell wall domains. *Developmental Cell* 48, 1-16.

Fry, S.C. (1982). Isodityrosine, a new cross-linking amino acid from plant cell-wall glycoprotein.
Biochem. J 204: 449–455.

Hall, Q. and Cannon, M.C. (2002). The cell wall hydroxyproline-rich glycoprotein RSH is essential for
normal embryo development in Arabidopsis. *Plant Cell* 14: 1161–1172.

Humphrey, W., Dalke, A., and Schulten, K. (1996). VMD: Visual molecular dynamics. *J. Molec.*
Graphics 14:33-38

Jackson, P.A., Galinha, C.I., Pereira, C.S., Fortunato, A., Soares, N.C., Amâncio, S.B., and Pinto
Ricardo, C.P. (2001). Rapid deposition of extensin during the elicitation of grapevine callus
cultures is specifically catalyzed by a 40-kilodalton peroxidase. *Plant Physiol.* 127: 1065–1076.

Jacobowitz, J.R, Doyle W.C., and Weng, J-K. (2019). PRX9 and PRX40 are Extensin Peroxidases
Essential for Maintaining Tapetum and Microspore Cell Wall Integrity during Arabidopsis Anther
Development. *Plant Cell* 31: 848–86.

Kim D Pruitt, Tatiana Tatusova, and Donna R Maglott. Ncbi reference sequences (refseq): a curated
non-redundant sequence database of genomes, transcripts and proteins. *Nucleic Acids Research*,
35(suppl 1):D61–D65, 2006.

Kumar, S., Stecher G., Tamura K. (2016) MEGA7: molecular evolutionaary genetic analysis version
7.0 for bigger datasets. *Mol. Biol. Evol.* 33(7): 1870-1874.

Kunieda, T., Shimada, T., Kondo, M., Nishimura, M., Nishitani, K., and Hara-Nishimura, I. (2013).
Spatiotemporal secretion of PEROXIDASE36 is required for seed coat mucilage extrusion in
Arabidopsis. *Plant Cell* 25: 1355–1367.

Lamport, D.T.A., Kieliszewski, M.J., Chen, Y., and Cannon, M.C. (2011). Role of the extensin
superfamily in primary cell wall architecture. *Plant Physiol*. 156: 11–19.

Lee, Y., Rubio, M.C., Alassimone, J., and Geldner, N. (2013). A mechanism for localized lignin
deposition in the endodermis. *Cell* 153: 402–412.

Liszkay, A., van der Zalm, E., and Schopfer, P. (2004). Production of reactive oxygen intermediates
(O(2)(.-), H(2)O(2), and (.)OH) by maize roots and their role in wall loosening and elongation
growth. *Plant Physiol.* 136: 3114–23.

Machado, M.R. and Pantano, S. (2016) SIRAH tools: mapping, backmapping and visualization of
coarse-grained models. *Bioinformatics* 32(10):1568–1570

Mangano S. *et al.* (2017). The molecular link between auxin and ROS-controlled root hair growth.
Proc. Natl. Acad. Sci. U.S.A. 114(20):5289-5294.

Marzol E, Borassi C, Bringas M, Sede A, Rodríguez Garcia DR, Capece L, Estevez JM. (2018). Filling the
Gaps to Solve the Extensin Puzzle. *Mol Plant*. 11(5):645-658.

McGibbon R.T, Beauchamp K.A., Harrigan M.P., Klein C, Swails, J.M., Hernández C.X., Schwantes
C.R., Wang, L.P., Lane T.J., Pande V.S. (2015) MDTraj: A Modern Open Library for the Analysis of
Molecular Dynamics Trajectories. *Biophysical Journal*, 109(8): 1528-1532

Menand B, Yi K, Jouannic S, Hoffmann L, Ryan E, Linstead P, Schaefer DG, Dolan L. (2007). An ancient
mechanism controls the development of cells with a rooting function in land plants. Science
316:1477-1480.

Møller, S.R. et al. (2017). Identification and evolution of a plant cell wall specific glycoprotein
glycosyl transferase, ExAD. *Sci. Rep.* 7: 45341.

Mutwil, M., Klie, S., Tohge, T., Giorgi, F.M., Wilkins, O., Campbell, M.M., Fernie, A.R., Usadel, B.,
Nikoloski, Z. and Persson, S. (2011). PlaNet: combined sequence and expression comparisons
across plant networks derived from seven species. *Plant Cell*, 23, 895– 910.

Oliphant, TE. USA: Trelgol Publishing, (2006). A guide to NumPy. 376 pp.

Parrinello, M. and Rahman, A. *(1981).* Polymorphic transitions in single crystals: A new molecular
dynamics method. *J. of Applied Physics* 52:7182. doi.org/10.1063/1.328693

Passardi, F., Penel, C., and Dunand, C. (2004). Performing the paradoxical: how plant peroxidases
modify the cell wall. *Trends Plant Sci*. 9:534–540.

Pereira S.P., J.M.L. Ribeiro, A.D. Vatulescu, K. Findlay, A.J. MacDougall and P.A.P. Jackson. (2011)
Extensin network formation in *Vitis vinifera* callus cells is an essential and causal event in rapid
and H_2O_2-induced reduction in primary cell wall hydration. *BMC Plant Biology* 111:106-121.

Price, N.J., Pinheiro, C., Soares, C.M., Ashford, D.A., Ricardo, C.P., and Jackson, P.A. (2003). A
biochemical and molecular characterization of LEP1, an extensin peroxidase from lupin. *J. Biol.*
Chem. 278:41389–41399.

Ringli C. (2010). The hydroxyproline-rich glycoprotein domain of the Arabidopsis LRX1 requires Tyr
for function but not for insolubilization in the cell wall. *Plant J.* 63, 662–669.

Roppolo, D., Boeckmann, B., Pfister, A., Boutet, E., Rubio, M.C., Denervaud-Tendon, V., Vermeer,
J.E.M., Gheyselinck, J., Xenarios, I., and Geldner, N. (2014). Functional and evolutionary analysis
of the CASPARIAN STRIP MEMBRANE DOMAIN PROTEIN family. Plant Physiol. 165, 1709–1722.

Saitou N, Nei M. 1987. The neighbor-joining method: a newmethod for reconstruction of
phylogenetic trees. Mol Biol Evol4:406–25.

Saito, F., Suyama, A., Oka, T., Yoko-O, T., Matsuoka, K., Jigami, Y., and Shimma, Y.-I. (2014).
Identification of Novel Peptidyl Serine α-Galactosyltransferase Gene Family in Plants. *J. Biol.*
Chem. 289: 20405–20420.

Šali A., and Tom L Blundell. Comparative protein modelling by satisfaction of spatial restraints. *J. of*
Molecular biology, 234(3):779–815, 1993.

Savelli B., Li Q., Webber M., Jemmat A.M., Robitaille A., Zamocky M., Mathe C., Dunand C. (2019).
RedoxiBase a database for ROS homeostasis regulated proteins. *Redox Biol*.
doi.org/10.1016/j.redox.2019.101247.

Schnabelrauch, L.S., Kieliszewski, M., Upham, B.L., Alizedeh, H., and Lamport, D. (1996). Isolation of
pI 4.6 extensin peroxidase from tomato cell suspension cultures and identification of Val-Tyr-Lys
as putative intermolecular cross-link site. *Plant J.* 9:477–489.
Sede, A.R., Borassi, C., Wengier, D.L., Mecchia, M.A., Estevez, J.M., and Muschietti, J.P. (2018).
Arabidopsis pollen extensins LRX are required for cell wall integrity during pollen tube growth.
FEBS Lett. 592, 233–243.
Shen Y., Rosendale M., Campbell R.E., D. Perrais (2014). pHuji, apH-sensitive red fluorescent protein
for imaging of exo- and endocytosis. *J.Cell Biol.* 207 (3):419-432.
Shigeto, J. and Tsutsumi, Y. (2016). Diverse functions and reactions of class III peroxidases. *New
Phytol*. 209: 1395–1402.
Showalter AM, Keppler B, Lichtenberg J, Gu D, Welch LR. 2010. A bioinformatics approach to the
identification, classification, and analysis of hydroxyproline-rich glycoproteins. *Plant Physiology*
153, 485–513.
Smith, A.T., Santama, N., Dacey, S., Edwards, M., Bray, R.C., Thorneley, R.N., and Burke, J.F. (1990).
Expression of a synthetic gene for horseradish peroxidase C in *Escherichia coli* and folding and
activation of the recombinant enzyme with Ca2+ and heme. *J. Biol. Chem.* 265: 13335–13343.
Stoddard, A & Rolland, V. (2018). I see the light!. Fluorescent proteins suitable for the cell
wall/apoplat targeeting in *Nicotiana benthamiana* leaves. *Plant Direct.* doi.org/10.1002/pld3.112
Tognolli, M., Penel, C., Greppin, H., and Simon, P. (2002). Analysis and expression of the class III
peroxidase large gene family in *Arabidopsis thaliana*. *Gene* 288: 129–138.
Trott O., and Arthur J Olson. (2010). Autodock vina: improving the speed and accuracy of docking
with a new scoring function, efficient optimization, and multithreading. *Journal of
Computational Chemistry*, 31(2):455–461.
Vanacore R, Ham AJ, Voehler M, Sanders CR, Conrads TP, Veenstra TD, Sharpless KB, Dawson PE,
Hudson BG (2009) A sulfilimine bond identified in collagen IV. Science 325:1230–1234.
Velasquez, S.M., Ricardi, M.M., Dorosz, J.G., Fernandez, P.V., Nadra, A.D., Pol-Fachin, L., Egelund, J.,
Gille, S., Ciancia, M., Verli, H., et al. (2011). O-glycosylated cell wall extensins are essential in root
hair growth. *Science* 332:1401–1403.
Velasquez M, Salter JS, Dorosz JG, Petersen BL, Estevez JM. (2012). Recent advances on the
posttranslational modifications of EXTs and their roles in plant cell walls. *Frontiers in plant
science* 3, 93.
Velasquez SM, Marzol E, Borassi C, Pol-Fachin L, Ricardi MM, Mangano S, Denita JS, Salgado SJ,
Gloazzo DJ, Marcus SE. (2015a). Low sugar is not always good: Impact of specific O-glycan
defects on tip growth in Arabidopsis. *Plant Physiology* 168, 808–813.
Velasquez, S.M., Ricardi, M.M., Poulsen, C.P., Oikawa, A., Dilokpimol, A., Halim, A., Mangano, S.,
Denita-Juarez, S.P., Marzol, E., Salgoda Salter, J.D., et al. (2015b). Complex regulation of prolyl-4-
hydroxylases impacts root hair expansion. *Mol. Plant* 8:734–746.
Wang, X., Wang, K., Yin, G., Liu, X., Liu, M., Cao, N., Duan, Y., Gao, H., Wang, W., Ge, W., et al.
(2018). Pollen-expressed leucine-rich repeat extensins are essential for pollen germination and
growth. Plant Physiol. 176, 1993–2006.
Welinder, K.G., Justesen, A.F., Kjaersgard, I.V., Jensen, R.B., Rasmussen, S.K., Jespersen, H.M., and
Duroux, L. (2002). Structural diversity and transcription of class III peroxidases from *Arabidopsis
thaliana*. *Eur. J. Biochem*. 269: 6063–6081.

Wojtaszek, P., Trethowan, J., and Bolwell, G.P. (1997). Reconstitution in vitro of the components and
conditions required for the oxidative cross-linking of extracellular proteins in French bean
(Phaseolus vulgaris L.). *FEBS Lett.* 405:95–98.

696
697
Figure 1. Characterization of root hair specific PRX01, PRX44 and PRX73 expression and mutant analysis.

(**A**) Transcriptional reporters of PRX01, PRX44 and PRX73 GFP-tagged in the root elongation zone
and specifically in root hairs (bottom). Scale bar = 20 μm. (*) means lack of expression in
atrichoblast cell layers.

(**B**) Root hair phenoClass of wild-Class (Columbia, Col-0) and *prx01,44,73* triple mutant. Box-plot of
root hair length. Horizontal lines show the means. (***) P<0.001 determined by one-way ANOVA.
Scale bars, 1 mm.

(**C**) Peroxidase activity in wild-Class (Columbia, Col-0) and *prx01,44,73* triple mutant roots. Enzyme
activity values (expressed as nkatal/mg protein) are the mean of three replicates ± SD. P-value of
one-way anova, (***) P < 0.001.

(**D**) Cytoplasmic ROS levels measured with H_2DCF-DA in Wild-Class (Columbia Col-0) and in the triple
mutant *prx01,44,73*. P-value of one-way anova, (***) P<0.001 and (**) P<0.01.

(**E**) Apoplastic ROS levels measured with Amplex™ UltraRed (AUR) in wild-Class (Columbia Col-0) and
in the triple mutant *prx01,44,73*. ROS signal was quantified from the root hair cell tip. P-value of
one-way anova, (***) P<0.001. Scale bar= 10 μm.

714
715
Figure 2. High level of expression of PRX44 and PRX73 trigger ectopic growth linked to higher
peroxidase activity.
(**A**) Root hair phenoClass of wild-Class (Columbia, Col-0) and PRXs^OE lines (in Wt Col-0). Box-plot of
root hair length. Horizontal lines show the means. ** P<0.001 determined by one-way ANOVA. Scale
bars, 1 mm. (NS) not significantly different.
(**B**) Selected pictures of root hair phenoClass analyzed in Figure 2A. Scale bar= 0.5mm.

(**C**) Expression of GFP-tagged 35S:PRX01, 35S:PRX44 and 35S:PRX73 in root hair cells.

(**D**) Western Blot of PRX01-GFP, PRX44-GFP and PRX73-GFP. Soluble GFP (sGFP) was used as control.

The predicted molecular weights are 62.6 KDa for PRX01-GFP, 60.8 KDa for PRX44-GFP, 62.9 KDa for

PRX73 and 27 KDa for sGFP.

(**E**) Assays of total peroxidase activity in wild-Class (Columbia Col-0) and PRXs^{OE} lines (in Wt Col-0).

Enzyme activity (expressed in nkatal/mg protein) was determined by a guaiacol oxidation-based

assay. Values are the mean of three replicates ± SD. P-value of one-way anova, (***) P<0.001, (**)

P<0.01, (*) P<0.1.

(**F**) Apoplastic ROS levels measured with Amplex™ UltraRed (AUR) in wild-Class (Columbia Col-0) and

PRXs^{OE} lines (in Wt Col-0). ROS signal was quantified from the root hair cell tip. P-value of one-way

anova, (NS) not significantly different. Scale bar= 10 μm.

733
734
Figure 3. Impact of deficient PRXs on the cell wall thickness in the root hair tips.
(A) Transmission electron micrographs of root hair tips from Wt Col-0, *prx1,44,73* triple mutant, and PRX44[OE] with (+) and without (-) SHAM. For each genotype and treatment, a representative overview of a root hair (RH) and a detail of the cell wall at the root hair tip (CW) is shown. (B) Whisker plot showing the cell wall thickness at the root hair tip of the three genotypes with or without SHAM treatment. ** P<0.001 determined by one-way ANOVA. (NS) not significantly different. Scale bars = 1μm.

Figure 4. Interaction by an *in silico* docking approach of PRX01, PRX44 and PRX73 with EXT peptides.

(A) PRX44 is shown as a protein surface in gray, with 10 docking results superimposed for each *O*-glycosylation state, to evaluate the consistency of docking sites. Model of a PRX44 complexed to a

non-*O*-glycosylated EXT substrate (SPPPYVY)$_3$ (in green). Detail of a non-glycosylated EXT substrate
depicted as sticks, docked to the protein represented as surface in gray. Heme is depicted as thin
sticks while iron is as a red sphere. On the bottom, two close tyrosine residues dock close to the
active site of PRX44.
(**B**) Model of a PRX44 complexed to an *O*-glycosylated-EXT substrate (in magenta the protein
backbone and in light blue the *O*-Glycans). Detail of an *O*-glycosylated EXT substrate depicted as
sticks, docked to the protein represented as surface in gray. Heme is depicted as thin sticks while
iron is as a red sphere. Arabino-galactosylated EXT peptide = [(SOOOYVY)$_3$-AG].
(**C**) A non-hydroxylated EXT peptide (SPPPYVY)$_3$, a hydroxylated but not *O*-glycosylated EXT peptide
[(SOOOYVY)$_3$; O=hydroxyproline], only arabinosylated EXT-peptide [(SOOOYVY)$_3$-A] and arabino-
galactosylated EXT peptide [(SOOOYVY)$_3$-AG] were analyzed. Comparison of the binding energy of
different peroxidases to its EXT substrates with different degrees of *O*-glycosylation.

Supplementary Materials
760

PRX01, PRX44, and PRX73 are class-III extensin-related peroxidases that modulates root hair
growth in *Arabidopsis thaliana*

Eliana Marzol[1,#], Cecilia Borassi[1,#], Philippe Ranocha[2], Alejandro Aptekman[3,4], Mauro Brindas[5],
Janice Pennington[6], Julio Paez-Valencia[6], Javier Martínez Pacheco[1], Diana Rosa Rodríguez Garcia[1],
Yossmayer Rondon[1], Mariana Carignani[1], Silvina Mangano[1], Margaret Fleming[7], Francisca Blanco-
Herrera[8,9], Patricia Bedinger[7], Christophe Dunand[2], Luciana Capece[5], Alejandro Nadra[3,4], Michael
Held[10], Marisa Otegui[6,11] & José M. Estevez[1,8,†]
769
770
[#]co-first authors
[†]Correspondence should be addressed. Email: jestevez@leloir.org.ar / jose.estevez@unab.cl (J.M.E).

Supplementary Materials

Table S1. Peptidyl-Tyr and iso-dityrosine (IDT) contents in cell walls isolated from Wt Col-0, PRX[OE]
lines and under *O*-glycosylated EXTs lines. *** P<0.001, ** P<0.01 and * P<0.1 determined by one-
way ANOVA. STD=Standard Deviation. In blue higher values and in light blue lower values than Wt
Col-0 are highlighted.

	ng Tyr/µg CW (STD)	ng IDT/µg CW (STD)
Wt Col-0	7.799 ± 0.26	0.853 ± 0.08
prx01 prx44 prx73	9.588 ± 0.31**	0.963 ± 0.02
PRX44[OE]	8.649 ± 0.07	0.953 ± 0.04
PRX73[OE]	8.700 ± 0.12	1.042 ± 0.02*
under O-glycosylated EXTs		
sergt1-1 rra3	3.530 ± 0.08***	0.235 ± 0.01***
p4h5 sergt1-1	3.766 ± 0.06***	0.225 ± 0.02***

785 **Table S2**. Volume and average diameter measurements on EXTs from molecular dynamics
786 simulations. These magnitudes were measured in the fully extended EXT system.

	Non-glycosylated EXT	*O*-glycosylated EXT	*O*-glycosylated EXT (protein only)
Volume /nm^3	280 ± 30	1080 ± 50	450 ± 30
EXT length /nm	65 ± 2	70 ± 2	70 ± 2
Average diameter /nm	2.4	4.5	2.9
Distance Tyr-Tyr			
0-5 Å	20%	25%	-
5-7 Å	55%	65%	-

789
790

Figure S1. **Root hair transcriptional co-expression network: PRXs, EXTs, and the transcriptional**
regulator RSL4 are highlighted. RSL4 was used as gene bait to narrow down the number of co-
expressed genes. Transcriptional connections between RSL4 and PRXs genes (in red) and bait genes
RSL4 (in black) are highlighted. The co-expression network was identified from PLaNet
(http://aranet.mpimp-golm.mpg.de/aranet) (Mutwil et al., 2011) and trimmed to facilitate
readability. Chromatin immunoprecipitation assay (ChiP) showed that RSL4 binds and controls the
expression of PRX01,44,73 (Mangano et al. 2017).

Figure S2. EXT reporter is expressed and targeted to root hair cell walls.

(**A**) Schematic diagrams of SS-TOM and SS-TOM-Long-EXT constructs expressed under the control of the 35S promoter in Arabidopsis plants. SS, Tomato polygalacturonase signal sequence. AS, Ala-spacer, 6 alanines between tdTomato (TOM) and Long-EXT domain. Long-EXT, C-terminal 290 amino acids of *SlPEX1*.

(**B**) Expression and detection of EXT-reporters in homozygous *Arabidopsis* lines. Lines expressing detectable levels of SS-TOM (lines #2 and #5) and SS-TOM-Long-EXT (lines #2 and #7) showed similarly unexpectedly large sizes of protein on the Western blot, with bands much larger than 94 kDa (which is the predicted size of the unmodified EXT protein). All the gel blots shown are part of the same run.

(**C**) Expression of SS-TOM and SS-TOM-Long-EXT reporters in *Arabidopsis* root hair cells. (On the bottom) Bright-field images (On the bottom left) cells were plasmolyzed in 1 M NaCl. Cell wall location is denoted with solid arrowheads while plasma membrane is indicated with an empty arrowhead. Scale bar= 10 µm.

(**D**) Root hair phenoClass of Wild-Class (Columbia, Col-0), SS-TOM (#2 and #5) and SS-TOM-Long-EXT (in Wt Col-0) as box-plot. Horizontal lines show the means. NS= not significant different, determined by one-way ANOVA.

816
817
Figure S3. **Expression of EXT-reporter in transiently transformed onion epidermal cells.**

(**A**) Onion epidermis expressing SS-TOM. (On the right) Bright field images. (On the bottom) cells were plasmolyzed in 1 M NaCl. Cell wall location is denoted with solid arrowheads while plasma membrane is indicated with an empty arrowhead. Scale bar= 100 μm. SS=Signal Peptide. TOM=Tomato fluorescent protein.

(**B**) Onion epidermis expressing SS-TOM-Long-EXT. (On the right) Bright fields images. (On the bottom) Cells were plasmolyzed in 1 M NaCl. Cell wall location is denoted with solid arrowheads while plasma membrane is indicated with an empty arrowhead. Scale bar= 100 μm. SS=Signal Peptide. TOM=Tomato fluorescent protein. Long-EXT= C-terminal 290 amino acids of *Solanum lycopersicum* (Sl) *SlPEX1*.

Figure S4. Effect of the _O_-glycosylation status on the triple helix EXT assembly and Tyr-Tyr distances.

Selected snapshots of the O-glycosylated (**A**) and non-glycosylated (**B**) triple EXT chain MD (molecular dynamics) trajectory, showing the stability of the triple strand along 1 us molecular dynamics simulation. The results obtained in these simulations highlight the importance of the triple-helix EXT in overall protein stability, and especially the maintenance of the fibril-like structure. Peptide chains are depicted in red, black and blue and glycans in green. In these simulations the peptide chains are allowed to move freely inside the simulation box, without any restrictions. Insets: a snapshot of a 25-aminoacid portion of the triple helix taken from the CG MD trajectory.

(**C**) Distribution of the Tyr-Tyr distances along the MD simulations for the glycosylated (red line) and non-glycosylated (black line) states. Distances are measured between the C-_ortho_ of the Tyrosine residues (as depicted in the inset of the figure).

-(Ara$_f$)$_4$ -Gal$_p$

Figure S5. **Coarse-grained models for tetra-arabinofuranose and galactopyranose moieties (diffuse green spheres) superimposed on all atom description (solid spheres connected by tubes)**. Interaction points or beads are located in the position of the heavy atoms and hydrogens corresponding to hydroxyl groups. The latter were included due to the importance of directional hydrogen bond-like interactions in this Class of molecules. Mass and Lennard-Jones parameters of CG beads and spring constants for intramolecular interactions were chosen to be compatible with the SIRAH coarse-grained model (parameters are available upon request).

Figure S6. Evolutionary relationships of class-III PRXs with putative EXT cross-linking activity.

The 73 PRXs protein sequences from *A. thaliana*, 2 putative Lignin-PRX from *Zinnia elegans* and 4

putative EXT-PRX from *Lupinus album*, *Lycopersicum esculentum*, *Phaseolus vulgaris* and *Vitis*

vinifera, have been aligned with ClustalW and the tree constructed using the Neighbor-Joining

method (Saitou and Nei, 1987). The analysis was conducted in MEGA7 (Kumar, 2016). Previous

identified PRXs with putative EXT-PRXs are indicated in blue boxes. The EXT-PRXs studied in the

present work are highlighted in green boxes. All protein sequences are available using their ID

number (http://peroxibase.toulouse.inra.fr; Savelli et al., 2019).

Figure S7. Evolutionary and root tissue expression relationships of putative EXT-PRXs in Arabidopsis.

The evolutionary history of apoplastic class-III PRXs in root hairs associated has been associated to 15 PRXs protein sequences of *A. thaliana*. They have been aligned with ClustalW and the tree constructed using the Neighbor-Joining method (Saitou and Nei, 1987). The analyses were conducted in MEGA7 (Kumar 2016). The sequences of AtPRX01/PRX2 and AtPRX50/51 are either identical or very close, leading to only two independent expression profiles from Affymetrix data. A brown (max value)-to-cream (min value) heat-map was drawn from 105 anatomical parts from data selection available from AT_AFFY_ATH1-0 and plotted in front of each branch and using Genevestigator. The class-III PRXs studied in the present work were surrounded in green. Green and blue circles correspond to duplication posterior to Brassicaceae and to Dicotyledons emergences respectively. All protein sequences are available using their ID number (http://peroxibase.toulouse.inra.fr; Savelli et al., 2019).

Figure S8. Current model of PRX01, PRX44, and PRX73 function on the ROS homeostasis linked to

EXT-crosslinking in the root hair cell walls. ROS homeostasis in the apoplast includes ROS

production, ROS consumption, and ROS transport to the cytoplasm. ROS as superoxide ion (O_2^-) is

primarily produced by RBOHC and by RBOHH,J (based on Mangano et al. (2017) and, possibly, by

unknown PRXs in the root hair tip, and then dismutated to hydrogen peroxide (H_2O_2). (**A**) It is

proposed that under high levels of PRX01,44,73 (in OE lines), this might trigger during the

peroxidative cycle (PC) a high consumption of apoplastic H_2O_2 and the crosslinking of EXT-

monomers at Tyr-units to allow the formation of the EXT-network and a concomitant cell wall

hardening. A portion of the apoplastic H_2O_2 might be transported into the cytoplasm by aquaporins

(PIPs). (**B**) In the triple mutant *prx01,44,73*, most of the H_2O_2 produced is accumulated in the

apoplast-cell walls and most of the EXT monomers are not cross-linked. The balance between cell

wall hardening and cell wall loosening processes might be compromised impacting on polar root

hair growth. CW= cell wall, PM= plasma membrane; HC= Hydroxylic Cycle; OC= Oxidative Cycle.