Cooperative policing behavior regulates reproductive division of labor in a termite

Qian Sun [1, 2], Jordan D. Hampton[2], Kenneth F. Haynes[2], Austin Merchant[2], Xuguo Zhou[2*]

1. Department of Entomology, Louisiana State University Agricultural Center, Baton Rouge, LA

  70803, USA

2. Department of Entomology, University of Kentucky, Lexington, KY 40546-0091, USA

Q.S. and J.D.H. contributed equally to this work.

[*]**Corresponding Author:**

Dr. Xuguo "Joe" Zhou

Department of Entomology

University of Kentucky

S-225 Agricultural Science Center North

Lexington, KY 40546-0091

Phone: 859-257-3125; Fax: 859-323-1120

Email: xuguozhou@uky.edu

Short Title: Policing behavior in termites

Abstract

Reproductive conflicts are common in insect societies where helping castes retain reproductive

potential. One of the mechanisms regulating the conflicts is policing, a coercive behavior that

reduces direct reproduction by other individuals. In eusocial Hymenoptera (ants, bees, and

wasps), workers or the queen act aggressively toward fertile workers, or destroy their eggs. In

many termite species (order Blattodea), upon the death of primary queen and king, workers or

nymphs can differentiate into neotenic reproductives and inherit the breeding position. During

this process, competition among neotenics is inevitable, but how this conflict is resolved remains

unclear. Here, we report a policing behavior that regulates reproductive division of labor in the

eastern subterranean termite, *Reticulitermes flavipes*. Our results demonstrate that the policing

behavior is a cooperative effort performed sequentially by successful neotenics and workers. A

neotenic reproductive initiates the attack of the fellow neotenic by biting and displays alarm

behavior. Workers are then recruited to cannibalize the injured neotenic. Furthermore, the

initiation of policing is age-dependent, with older reproductives attacking younger ones, thereby

inheriting the reproductive position. This study provides empirical evidence of policing behavior

in termites, which represents a convergent trait shared between eusocial Hymenoptera and

Blattodea.

Keywords: policing behavior, reproductive conflict, neotenic reproduction, termites, eusociality

1. Introduction

Eusocial insects exhibit reproductive division of labor between a few reproductive individuals and numerous sterile workers. However, potential conflicts arise in species where workers are capable of reproduction [1-4]. In addition to queen pheromones that suppress worker reproduction in many species [5-7], policing behavior is an important solution to maintain reproductive harmony [2, 8]. Queen policing, a term coined by Oster and Wilson in 1978, describes behaviors carried out by the queen to retain her reproductive dominance over workers [9]. Worker policing, which was named by Ratnieks in 1988, was used to describe the actions of workers that reduce worker-produced males in favor of sons of the queen in honey bee [8]. The concept of "policing behavior" was expanded by Monnin and Ratnieks in 2001 to include all "coercive actions that reduce direct reproduction by other individuals", which accommodates various forms of behavioral regulation observed in diverse social insects [10]. In eusocial Hymenoptera (ants, bees, and wasps), policing behavior is performed via egg-eating or different forms of aggression, such as immobilization, biting, and stinging [10-13]. Policing interactions may occur among workers [11], among reproductives [14, 15], or between reproductives and workers [12, 16-18]. However, the vast majority of investigations on policing behavior have been focused on Hymenoptera, with little is known about its occurrence or nature in termites (order Blattodea), a group of eusocial insects evolved 50 million years earlier than eusocial Hymenoptera [19].

Policing behavior serves two functions that are not mutually exclusive, which are regulating genetic conflicts and improving colony efficiency [8, 20, 21]. Genetic conflicts arise between colony members due to relatedness asymmetries. For example, in social Hymenoptera with haplodiploid sex determination, workers are often capable of laying unfertilized male eggs,

and they are more closely related to their own sons than males produced by other individuals [2,

8, 22]. Moreover, policing behavior contributes to colony efficiency even when little or no

genetic conflict is present, as it optimizes the allocation of colony resources to reproduction [8,

23], or maintains an adaptive colony-level phenotype [24].

Unlike social Hymenoptera, termite colonies are usually founded and dominated by a pair

of primary reproductives (queen and king). Upon their death, neotenic reproductives of both

sexes can differentiate either from workers, which become ergatoid reproductives, or from

nymphs, which become nymphoid reproductives [25]. Reproductive succession by neotenics has

been reported in at least 13.4% of "higher" termite genera (Termitidae) and 61.7% of "lower"

termite genera (all other termite families) [26]. In termites with diplodiploid sex-determination,

individuals are more closely related to their own offspring than that of their siblings, regardless

of outbreeding or inbreeding. Therefore, genetic conflicts are potential and competition for

reproduction between nestmates is expected [27]. Moreover, given the fact that workers are

reproductively totipotent in many species, colony-level efficiency can be compromised if excess

neotenics differentiate, as it results in reduced labor force and increased resource demand by the

reproductives and their brood.

In the presence of fertile reproductives, neotenic formation is inhibited through

pheromones [6, 28, 29], and policing behavior through overt aggression was considered rare in

termites [28, 30]. However, during the process of reproductive succession when colonies are

orphaned and inhibitory pheromones are temporarily absent, production of excessive neotenic

reproductives in the colony is expected. In addition, cannibalism of neotenics was observed in

several termite families including Termopsidae [31], Kalotermitidae [30, 32-34] and

Rhinotermitidae [35, 36], suggesting the presence of policing behavior that directly regulates

reproduction in termites. Empirical studies of the process and causes of policing behavior,

however, are lacking. In this study, we conducted a series of experiments to understand whether

the number of ergatoid neotenics is regulated behaviorally during reproductive succession, and

how policing behavior is performed in the eastern subterranean termite, *Reticulitermes flavipes*.

Furthermore, we investigated a proximate factor that determines the succession of reproductives.

2. Methods

(a) Insect collection and maintenance

Field and laboratory colonies of *R. flavipes* were used in this study. Field colonies were collected

from Red River Gorge area, Daniel Boone National Forest (Slade, Kentucky, USA) and the

University of Kentucky Arboretum (Lexington, Kentucky, USA). These colonies were obtained

in summer using trapping stations filled with dampened cardboard rolls. Once captured termites

were extracted from traps, placed in Petri dishes (14.5 cm × 2.0 cm) with moistened unbleached

paper towel as their only food source for 7- 10 days before they were used in experiments. Only

workers and soldiers were collected from the field. Laboratory colonies were established in 2010

by pairs of sibling alates from a dispersal flight in Lexington, Kentucky, and they were kept in

closed plastic boxes filled with moistened wood mulch and pinewood blocks in laboratory for 5

years before use. All colonies were maintained in complete darkness (L:D = 0:24), at 27 ± 1ºC,

80%- 99% RH.

(b) Orphaning assay to test ergatoid number restriction

This assay was used to simulate the reproductive replacement process after the death of primary

reproductives. Groups of 100 workers were kept in petri dishes (35-mm-diameter) with

moistened paper towel placed at the bottom. Two treatments including "removal" and "non-

removal" of ergatoids were conducted. This experiment was specifically designed in a way to

compare between the numbers of ergatoids that can potentially differentiate and survive

("removal") and that actually survived ("non-removal"). All termites were maintained at 27 ±

1℃ and in complete darkness for 90 days. Each dish was checked by identifying the sex and

counting the number of newly differentiated ergatoids. Each ergatoid was removed and replaced

with a worker in the "removal" treatment, but returned to the dishes in the "non-removal"

treatment. Dishes in "removal" treatment were checked every day, while dishes in "non-

removal" treatment were checked every 10 days to reduce stress to reproductives resulting from

manipulations needed for sex identification. The number of remaining termites in each group

was counted every 30 days, and mortality was calculated based on the difference between the

numbers of initial and remaining individuals. Injured ergatoids were not counted. A total of 20

replications were made with five replications in each of the four colonies. Two field and two

laboratory colonies were used in this experiment.

(c) Orphaning assay for observation of policing behavior

This assay was designed to observe policing behavior under orphaning condition, which

resembled the "non-removal" treatment above. The dishes were incubated at 27 ± 1℃ for a total

of 90 days. Once a week, the dishes were checked for dryness, and water was added if the paper

towels at the bottom appeared to be dry. Between 60 and 90 days, each dish was checked for the

presence of ergatoids, and the dishes with ergatoids were selected to be video recorded for 6 days.

Video cameras (Canon Vixia HF G20, Canon Inc., Tokyo, Japan) were used for recording and

yield high quality images. The dishes and the cameras were shaded under a piece of cardboard

(1.5 m × 0.8 m) during recording. Ergatoids were identified as males or females and color

marked using enamel paint (Testor Corporation, Rockford, IL, USA) on their head capsule prior

to recording. During recording, the dishes were checked every day for missing ergatoids (which

were cannibalized), and newly formed ergatoids were color marked. When a marked ergatoid

was missing, the video of the previous 24 hours was quantitatively analyzed.

We define the "victim" as the ergatoid that was eventually cannibalized, the "attacker" as

the individual who attacked the victim, and "bystander" as the ergatoids who did not perform the

first major attack. A major attack was recognized when the attacker visibly injured the victim

such that the researcher could see the abdomen was torn, hemolymph was leaking, and the victim

quickly fled. The first major attack was designated as time "0", and the frequency of vibration

and number of workers surrounding the victim were documented. A one-minute sample (30

seconds before and after the time point) was analyzed for all these behaviors, with the sample

selected for the time midpoints 5, 15, 20, 25, 30, 45, and 60 minutes before and after time "0".

Worker density near the victim was an indicator of cannibalism. The density was quantified by

counting the number of workers and soldiers within 4 mm radius from the center of the victim

(the radius approximately equals to body length of a worker). Only field-collected colonies were

used in this experiment.

(d) Policing assay to test the effect of ergatoid age

This assay was used to determine if policing behavior in *R. flavipes* is associated with the age of

ergatoids. Each group of 50 workers was kept in a petri dish (55-mm-diameter) lined with 6

layers of moistened paper towel. A pair of virgin ergatoids, female and male, was added to each

dish on day 1. Female and male treatments were conducted, with a younger female or male

ergatoid being added to the initial group every day. When added to the dish, the initial pair of

ergatoids was 7 days post differentiation (7-day old) and became older over the course of the

experiment, and subsequent ergatoids were no more than 7-day old. All of the ergatoids were

color marked as previously described, and no new ergatoids differentiated during this assay. The

dishes were recorded until one of the ergatoids was missing. The video was then analyzed to

identify who attacked that ergatoid. A total of 9 and 10 replications were made for female and

male treatments, respectively; aggressively interacting pairs that were the same sex were

analyzed for their age differences. This experiment did not attempt to address sex-specificity, but

it was designed for increased chance of sex-specific aggression, and only same-sex aggressions

were analyzed to eliminate confounding factors associated with sex. Only field-collected

colonies were used in this experiment.

(e) Statistical analyses

Data were analyzed using R (https://www.r-project.org/) and Statistix 10 (Analytical Software,

Tallahassee, FL, USA), and graphs were generated using SigmaPlot 13 (Systat Software Inc,

Chicago, IL, USA). Data for ergatoid number and mortality in reproductive "removal" and "non-

removal" assays were fitted to Poisson family generalized linear mixed models using R's *glmer*

function. In the model for ergatoid number, ergatoid sex and treatment group (removal vs. non-

removal) were coded as fixed effects, while colony of origin was coded as a random effect. An

observation-level random effect was introduced to avoid overdispersion. In the model for

mortality, treatment group was coded as a fixed effect, while colony of origin was coded as a

random effect. In both cases, data were analyzed separately for each 10-day interval. Data testing

the influence of ergatoid age on aggressive interactions in the policing assay were analyzed in

Statistix using a Wilcoxon signed-rank test.

3. Results

(a) The number of ergatoids is restricted

In "removal" treatment, there were significantly more female and male ergatoids differentiated

than that remained in "non-removal" treatment within 90 days (Figure 1; $x^2 = 137.14$, df = 1, $P <$

0.001; GLMM, Poisson family; n = 20). At the end of day 90, there were 11.80 ± 2.59 (mean ±

SE) female and 4.75 ± 1.06 male ergatoids differentiated when they were removed daily,

compared to only 2.60 ± 0.34 female and 1.05 ± 0.05 male ergatoids if they were not removed.

In addition, there was a significantly higher overall mortality in "non-removal" than that in

"removal" treatment (Figure S1; $x^2 = 30.36$, df = 1, $P < 0.001$; GLMM, Poisson family; n = 20).

In "non-removal" groups, we frequently observed injured ergatoids partially consumed by

workers, along with other intact ergatoids. Between the two treatments, the mortality difference

(8.6 individuals at day 60 and 10.2 at day 90) closely matched the difference in the number of

ergatoids that differentiated and that survived (6.2 at day 60 and 12.9 at day 90), suggesting

cannibalism of ergatoids in "non-removal" treatment is primarily responsible for the difference

in mortality.

(b) Ergatoids and workers cooperate in elimination of excessive ergatoids

A total of seven events were captured with full behavioral process, which started with one

ergatoid attacking another ergatoid and ended with the injured individual being cannibalized by

workers (Figure 2, Table S1, Movies S1-S4). In this behavior, the attacker antennated the victim

first, and used mouthparts to hold the abdomen or thorax of the victim before biting. The bite

always caused the victim to leak hemolymph and quickly flee. Right after the attack, the attacker

displayed alarm behavior by vigorously vibrating the body toward multiple directions;

interestingly, the ergatoids who did not participate in the aggression (i.e., bystanders) also

performed alarm behavior after the attack, while the victim rarely engaged in vibration (Figure

3a). Workers, on average, displayed little vibration (Figure 3a). With the alarm of ergatoids,

workers rapidly began to surround the injured victim, biting and consuming it while it was still

alive, and the cannibalism reached a peak 30 minutes after the attack (Figure 3b). As the

cannibalism began to decline due to little of the remaining body to consume, the ergatoid

attacker and bystanders began to decline in the frequency of alarm (45 to 60 minutes after the

attack) (Figure 3a). Soldiers did not participate in either alarm behavior or cannibalism (Figure

3a, 3b).

(c) Ergatoid elimination is age-dependent

The majority of the aggressive interactions involved an ergatoid attacking the same-sex ergatoid

(15 out of 19), while the others involved either a male ergatoid attacking a female (1 out 19), or a

group of workers attacking and consuming an ergatoid (1 and 2 cases in female and male

treatments, respectively). In the same-sex aggressive interactions, the attacker was always older

than the victim in both female and male treatments (Figure 4; female: $Z = -2.3664$, $P < 0.01$, n =

7; male: $Z = -2.5205$, $P < 0.01$, n = 8; Wilcoxon signed-rank test, one-tailed). The median age

differences between the pair of attacker and victim were 5 and 2 days for females and males,

respectively.

4. Discussion

(a) Cooperative effort and justification of "policing behavior"

Overall, this study reveals a behavioral mechanism regulating reproductive division of labor

during reproductive succession in termites. Unlike butting behavior that is an indicator of

reproductive dominance reported in a drywood termite *Cryptotermes secundus* and a dampwood

termite *Zootermopsis nevadensis* [37, 38], this behavior directly acts to eliminate reproductive

individuals, thus fulfilling the definition of policing [10].

Policing in *R. flavipes* is a sequential behavior cooperatively performed by the older

reproductives in concert with workers. This is comparable to the sting smearing behavior in a

queenless ant, *Dinoponera quadriceps*, a textbook example of policing behavior [39]. In this ant,

the alpha female chemically marks a low-ranking challenger using stinger, causing workers to

immobilize the marked individual [40]. Such a cooperative effort allows the alpha to inflict

punishment indirectly and maintain her dominant status without fighting. Similarly, in *R. flavipes*,

the ergatoid attacker does not kill its rival directly; rather, it induces hemolymph exposure of the

victim by biting, and proceeds with an alarm behavior. Workers eventually eliminate the injured

individual. The workers are likely recruited by the vibrational signals, and the cannibalistic

behavior is potentially caused by the chemical cues in the hemolymph, which remains to be

investigated.

(b) Caste fate conflict

In the absence of reproductives and their inhibitory pheromones, the excess ergatoid production

in *R. flavipes* is an expression of caste fate conflict, which also occurs in social Hymenoptera.

For example, in the *Melipona* stingless bees, caste is self-determined, and immature females

selfishly develop into queens to maximize direct reproduction [41]. Such excess queen

production causes depletion of workforce and poses a cost to the colony, leading to a situation

known as "tragedy of the commons" [42]. The policing behavior in *R. flavipes* provides an

effective solution that prevents unsustainable reproduction upon the loss of former reproductives,

and supports colony efficiency. Uncontrolled reproduction causes increased resource demand by

the reproductive individuals and their brood, and a balanced ratio of reproductives and workers is

often optimal to the colony. For example, in a parthenogenic ant *Platythyrea punctate*, worker to

brood ratio is limited, and colonies are incapable of rearing brood produced by additional

reproductives [23]. Similarly, an optimal allocation to reproduction is required in the Japanese

subterranean termite *R. speratus*, and increased number of queens does not lead to additional

reproductive output [43].

In social insects, policing often inhibits reproduction of the focal individual without

killing it. In Hymenoptera, policing behavior often enforces workers to stop reproducing and

cooperate in brood care. In subterranean termites, however, worker-reproductive differentiation

is an irreversible process achieved through at least one molt [25]. Ergatoid reproductives are a

non-foraging caste that depends on workers to provide food [44], and the presence of excess

ergatoids and the subsequent brood are costly for the colony. Elimination of additional ergatoids

avoids future colony investment; cannibalism, in addition, allows the colony to recycle nutrients

from the policed individuals and partially rescue the cost that has already occurred upon their

differentiation.

(c) Age-dependent elimination

The age of ergatoids plays an important role in the policing behavior of *R. flavipes*. The result is

consistent with a dampwood termite *Porotermes adamsoni*, in which neotenic reproductives that

develop earlier have higher survivorship than that differentiate later [45]. Age is also positively

correlated with dominance rank in the naked mole-rat *Heterocephalus glaber* [46], in which

younger individuals receive more aggression than older ones [47]. In termites, older ergatoids

have the first chance to utilize resources from the colony, such as food provided by workers and

mating opportunities with existing reproductives. These factors may contribute to the maturity of

ergatoids in terms of gonad development [48], body weight, and mandible sclerotization, thus

allowing them to outcompete the younger ones. In *C. secundus*, the surviving neotenics

performed increased interactions with workers through proctodeal trophallaxis after their

differentiation, while the ones eventually eliminated did not [30]. This behavior is possibly

associated with age, but it is yet to be tested if neotenics increase trophallaxis as they become

older.

The elimination of younger ergatoids in termites is also similar to the selective

elimination of small queens in a stingless bee *Schwarziana quadripunctata*, where large fecund

queens are favored and dwarf queens tend to be killed by workers [49]. With a policing behavior

that strongly acts against small queens, small females should be less tempted to develop into

queens, a situation indicated by a theoretical study based on inclusive fitness theory [49]. In *R.*

flavipes, while a few ergatoids are formed and eliminated, the majority of workers do not molt

into ergatoids. This suggests that the same theory may apply to termites: in the presence of older

ergatoid, rather than developing into reproductives and being killed, workers can gain indirect

fitness benefits by not differentiating.

(d) Reproductive competition

The aggressive interaction between ergatoids in termites reflects competition among colony

members in inheriting the breeding position after the death of primary reproductives. This form

of cooperative policing might be common in termite species where workers or nymphs retain

reproductive potential and compete for breeding positions. Indeed, a similar behavior has been

observed in a drywood termite, *Kalotermes flavicollis*, where neotenic reproductives attack each

other and injured individuals are cannibalized by workers and nymphs [34]. In addition,

reproductive competition also occurs upon fusion of neighboring conspecific colonies. In *Z.*

nevadensis, reproductives of encountering colonies engage in agonistic behavior, leading to a

reduction in their numbers [50]. Moreover, reproductive conflict among unrelated queens

happens in species where colonies are founded by pleometrosis. In a fungus-growing termite,

Macrotermes michaelseni, the mutilation of queen antennae indicates aggression between

primary reproductives that are co-founders, and this behavior may influence queen number [51].

These findings suggest that aggressive interaction between reproductives is widespread in

termites under diverse social contexts. Aggression represents a conserved component of policing

behavior, which, at colony level, regulates reproductive division of labor and serve a collective

interest.

(e) Future directions

Several open questions remain to be addressed in regard to the pheromonal mechanisms

underlying policing behavior, including the dynamic change of reproductive pheromones with

age, and the chemical cues inducing cannibalism of ergatoids by workers. Importantly, policing

behavior, in a broad sense, may be more widespread in termites than previously considered. In

addition to a number of convergent traits in social Hymenoptera and Blattodea, such as suicidal

colony defense [52], collective foraging using trail pheromones [53], and undertaking behavior

[54], policing behavior represents another important behavior that independently evolved in both

eusocial groups. Comparative studies on proximate and ultimate aspects of policing behavior

among social insect taxa will provide important insights into the evolution of eusociality in

insects.

Supplementary material. Supplementary material includes 1 table, 1 figure, and 4 movies.

Data accessibility. Raw data can be accessed as supplementary material.

Authors' contributions. Q.S., J.D.H. contributed equally to this study. Q.S., J.D.H., K.F.H. and

X.Z. designed the study. Q.S. and J.D.H. performed the experiments. Q.S., J.D.H and A.M

analyzed the data. Q.S., J.D.H., K.F.H. and X.Z. wrote the manuscript. All authors read and

approved the final manuscript.

Competing interests. The authors declare no competing interests.

Acknowledgments. We thank members of Zhou laboratory for discussion and comments on the

study. This work was supported by William L. and Ruth D. Nutting Student Research Grant from

the International Union for the Study of Social Insects (North American Section) to Q.S., and a

Hatch fund (Accession Number: 1004654; Project Number: KY008071) from the USDA

National Institute of Food and Agriculture to X.Z.

References

1.   Robinson, G.E. (1992). Regulation of division of labor in insect societies. Annu. Rev.

305   Entomol. *37*, 637-665.

306   2.   Ratnieks, F.L., Foster, K.R., and Wenseleers, T. (2006). Conflict resolution in insect

307   societies. Annu. Rev. Entomol. *51*, 581-608.

308   3.   Hart, A.G., and Monnin, T. (2006). Conflict over the timing of breeder replacement in

309   vertebrate and invertebrate societies. Insectes Soc. *53*, 375-389.

310   4.   Korb, J. (2018). Chemical fertility signaling in termites: idiosyncrasies and

311   commonalities in comparison with ants. J. Chem. Ecol. *44*, 818-826.

312   5.   Conte, Y.L., and Hefetz, A. (2008). Primer pheromones in social Hymenoptera. Annu.

313   Rev. Entomol. *53*, 523-542.

314   6.   Matsuura, K., Himuro, C., Yokoi, T., Yamamoto, Y., Vargo, E.L., and Keller, L. (2010).

315   Identification of a pheromone regulating caste differentiation in termites. Proc. Natl.

316   Acad. Sci. U. S. A. *107*, 12963-12968.

317   7.   Holman, L., Jørgensen, C.G., Nielsen, J., and d'Ettorre, P. (2010). Identification of an ant

318   queen pheromone regulating worker sterility. Proc. Royal Soc. B *277*, 3793-3800.

319   8.   Ratnieks, F.L.W. (1988). Reproductive harmony via mutual policing by workers in

320   eusocial Hymenoptera. Am. Nat. *132*, 217-236.

321   9.   Oster, G.F., and Wilson, E.O. (1978). Caste and ecology in the social insects (Princeton

322   University Press).

323   10.   Monnin, T., and Ratnieks, F.L. (2001). Policing in queenless ponerine ants. Behav. Ecol.

324   Sociobiol. *50*, 97-108.

325   11.   Ratnieks, F.L.W., and Visscher, P.K. (1989). Worker policing in the honeybee. Nature

326   *342*, 796-797.

327   12.   Wenseleers, T., Tofilski, A., and Ratnieks, F.L.W. (2005). Queen and worker policing in

328   the tree wasp *Dolichovespula sylvestris*. Behav. Ecol. Sociobiol. *58*, 80-86.

329   13.   Foster, K.R., and Ratnieks, F.L.W. (2001). Convergent evolution of worker policing by

330   egg eating in the honeybee and common wasp. Proc. Royal Soc. B *268*, 169-174.

331   14.   Bourke, A.F.G. (1991). Queen behaviour, reproduction and egg cannibalism in multiple-

332   queen colonies of the ant *Leptothorax acervorum*. Anim. Behav. *42*, 295-310.

333   15.   Medeiros, F.N.S., Lopes, L.E., Moutinho, P.R.S., Oliveira, P.S., and Hölldobler, B.

334   (1992). Functional polygyny, agonistic interactions and reproductive dominance in the

335   Neotropical ant *Odontomachus chelifer* (Hymenoptera, Formicidae, Ponerinae). Ethology

336   *91*, 134-146.

337   16.   Adams, E.S., and Balas, M.T. (1999). Worker discrimination among queens in newly

338   founded colonies of the fire ant *Solenopsis invicta*. Behav. Ecol. Sociobiol. *45*, 330-338.

339   17.   Loope, K. (2015). Queen killing is linked to high worker-worker relatedness in a social

340   wasp. Curr. Biol. *25*, 2976-2979.

341   18.   Monnin, T., and Peeters, C. (1997). Cannibalism of subordinates' eggs in the

342   monogynous queenless ant *Dinoponera quadriceps*. Naturwissenschaften *84*, 499-502.

343   19.   Misof, B., Liu, S., Meusemann, K., Peters, R.S., Donath, A., Mayer, C., Frandsen, P.B.,

344   Ware, J., Flouri, T., Beutel, R.G., et al. (2014). Phylogenomics resolves the timing and

345   pattern of insect evolution. Science *346*, 763-767.

346   20.   Tom Wenseleers, and Francis L. W. Ratnieks (2006). Comparative analysis of worker

347   reproduction and policing in eusocial Hymenoptera supports relatedness theory. Am. Nat.

348   *168*, E163-E179.

349   21.   Walter, B., Brunner, E., and Heinze, J. (2011). Policing effectiveness depends on

350   relatedness and group size. Am. Nat. *177*, 368-376.

351   22.   Ratnieks, F.L.W., and Reeve, H.K. (1992). Conflict in single-queen hymenopteran

352   societies: the structure of conflict and processes that reduce conflict in advanced eusocial

353   species. J. Theor. Biol. *158*, 33-65.

354   23.   Hartmann, A., Wantia, J., Torres, J.A., and Heinze, J. (2003). Worker policing without

355   genetic conflicts in a clonal ant. Proc. Natl. Acad. Sci. U. S. A. *100*, 12836-12840.

356   24.   Teseo, S., Kronauer, D.J.C., Jaisson, P., and Châline, N. (2013). Enforcement of

357   reproductive synchrony via policing in a clonal ant. Curr. Biol. *23*, 328-332.

358   25.   Korb, J., and Hartfelder, K. (2008). Life history and development - a framework for

359   understanding developmental plasticity in lower termites. Biol. Rev. *83*, 295-313.

360   26.   Myles, T.G. (1999). Review of secondary reproduction in termites (Insecta: Isoptera)

361   with comments on its role in termite ecology and social evolution. Sociobiology *33*, 1-43.

362   27.   Bartz, S.H. (1979). Evolution of eusociality in termites. Proc. Natl. Acad. Sci. U. S. A. *76*,

363   5764-5768.

364   28.   Matsuura, K. (2012). Multifunctional queen pheromone and maintenance of reproductive

365   harmony in termite colonies. J. Chem. Ecol. *38*, 746-754.

366   29.   Sun, Q., Haynes, K.F., Hampton, J.D., and Zhou, X. (2017). Sex-specific inhibition and

367   stimulation of worker-reproductive transition in a termite. Sci. Nat. *104*, 79.

30. Hoffmann, K., and Korb, J. (2011). Is there conflict over direct reproduction in lower

termite colonies? Anim. Behav. *81*, 265-274.

31. Lenz, M. (1985). Is inter-and intraspecific variability of lower termite neotenic numbers

due to adaptive thresholds for neotenic elimination?—Considerations from studies on

Porotermes adamsoni (Froggatt)(Isoptera: Termopsidae). In Caste differentiation in

social insects, Volume 3, J.A.L. Watson, B.M. Okot-Kotber and C. Noirot, eds. (Oxford:

Pergamon Press), p. 125.

32. Lenz, M., McMahan, E.A., and Williams, E.R. (1982). Neotenic production in

Cryptotermes brevis (Walker): Influence of geographical origin, group composition, and

maintenance conditions (Isoptera: Kalotermitidae). Insectes Soc. *29*, 148-163.

33. Lüscher, M. (1952). Die produktion und elimination von ersatzgeschlechtstieren bei der

termite *Kalotermes flavicollis* Fabr. J. Comp. Physiol. A *34*, 123-141.

34. Ruppli, E. (1969). Die Elimination Überzähliger Ersatzgeschlechtstiere bei der Termite

Kalotermes Flavicollis (Fabr.). Insectes Soc. *16*, 235-248.

35. Costa-Leonardo, A.M., Arab, A., and Casarin, F.E. (2004). Neotenic formation in

laboratory colonies of the termite *Coptotermes gestroi* after orphaning. J. Insect Sci. *4*, 10.

36. Miyata, H., Furuichi, H., and Kitade, O. (2004). Patterns of neotenic differentiation in a

subterranean termite, *Reticulitermes speratus* (Isoptera: Rhinotermitidae). Entomol. Sci. *7*,

309-314.

37. Korb, J., Weil, T., Hoffmann, K., Foster, K.R., and Rehli, M. (2009). A gene necessary

for reproductive suppression in termites. Science *324*, 758-758.

38.   Penick, C.A., Trobaugh, B., Brent, C.S., and Liebig, J. (2013). Head-butting as an early

390   indicator of reproductive disinhibition in the termite *Zootermopsis nevadensis*. J. Insect

391   Behav. *26*, 23-34.

392   39.   Rubenstein, D.R., and Alcock, J. (2019). Animal Behavior, (Sinauer Associates).

393   40.   Monnin, T., Ratnieks, F.L.W., Jones, G.R., and Beard, R. (2002). Pretender punishment

394   induced by chemical signalling in a queenless ant. Nature *419*, 61.

395   41.   Wenseleers, T., Ratnieks, F.L., and Billen, J. (2003). Caste fate conflict in swarm-

396   founding social Hymenoptera: an inclusive fitness analysis. J. Evol. Biol. *16*, 647-658.

397   42.   Wenseleers, T., and Ratnieks, F.L. (2004). Tragedy of the commons in *Melipona* bees.

398   Proc. Royal Soc. B *271*, S310-S312.

399   43.   Yamamoto, Y., and Matsuura, K. (2011). Queen pheromone regulates egg production in a

400   termite. Biol. Lett. *7*, 727-729.

401   44.   Eggleton, P. (2011). An introduction to termites: biology, taxonomy and functional

402   morphology. In Biology of Termites: a Modern Synthesis, D.E. Bignell, Y. Roisin and N.

403   Lo, eds. (Dordrecht: Springer Netherlands), pp. 1-26.

404   45.   Mensa-Bonsu, A. (1976). The production and elimination of supplementary reproductives

405   in *Porotermes adamsoni* (Froggatt) (Isoptera, Hodotermitidae). Insectes Soc. *23*, 133-153.

406   46.   Clarke, F.M., and Faulkes, C.G. (1997). Dominance and queen succession in captive

407   colonies of the eusocial naked mole-rat, *Heterocephalus glaber*. Proc. Royal Soc. B *264*,

408   993-1000.

409   47.   O'Riain, J. (1996). Pup ontogeny and factors influencing behavioural and morphological

410   variation in naked mole-rats, *Heterocephalus glaber* (Rodentia, Bathyergidae). In

411   Zoology, Volume Ph.D. (University of Cape Town).

48. Brent, C.S., and Traniello, J.F.A. (2001). Social influence of larvae on ovarian maturation

in primary and secondary reproductives of the dampwood termite *Zootermopsis*

angusticollis. Physiol. Entomol. *26*, 78-85.

49. Wenseleers, T., Hart, A.G., and Ratnieks, F.L. (2004). When resistance is useless:

policing and the evolution of reproductive acquiescence in insect societies. Am. Nat. *164*,

E154-E167.

50. Thorne, B.L., Breisch, N.L., and Muscedere, M.L. (2003). Evolution of eusociality and

the soldier caste in termites: Influence of intraspecific competition and accelerated

inheritance. Proc. Natl. Acad. Sci. U. S. A. *100*, 12808-12813.

51. Brandl, R., Hacker, M., Bagine, R., and Kaib, M. (2001). Geographic variation of

polygyny in the termite *Macrotermes michaelseni* (Sjöstedt). Insectes Soc. *48*, 134-137.

52. Shorter, J., and Rueppell, O. (2012). A review on self-destructive defense behaviors in

social insects. Insectes Soc. *59*, 1-10.

53. Czaczkes, T.J., Grüter, C., and Ratnieks, F.L. (2015). Trail pheromones: an integrative

view of their role in social insect colony organization. Annu. Rev. Entomol. *60*, 581-599.

54. Sun, Q., Haynes, K.F., and Zhou, X. (2018). Managing the risks and rewards of death in

eusocial insects. Philos. Trans. R. Soc. B *373*, 20170258.

Figure legends

Figure 1. Restriction of ergatoid numbers in orphaning assay. Female and male ergatoids that

formed from groups of 100 workers were documented for 90 days post establishment of

experiment. Cumulative number of differentiated ergatoids (removal treatment) and surviving

number of ergatoids (non-removal treatment) are presented (mean ± SE) and compared within

the same observation day. n.s., not significant; ***, $P < 0.001$; GLMM, Poisson family; n = 20

per treatment per observation day.

Figure 2. Behavioral process of cooperative policing. (a) An ergatoid attacker holds the victim using mouthparts (Time = -10 sec). (b) The attacker performs a major attack by biting the victim on its abdomen (Time = 0). (c) The major attack causes the victim to quickly flee (Time = 0). (d) The attacker displays alarm behavior by vigorously vibrating its body toward multiple directions (Time = 1 min). (e) Bystanders also display alarm behavior in the same manner (Time = 5 min). (f) Workers then surround and nibble the victim (Time = 30 min). (g) Victim is partially cannibalized (Time = 2 hour). Attacker: ergatoid who performed the first major attack. Victim: ergatoid who received the attack and was later cannibalized. Bystander: other ergatoids.

Figure 3. Participation of different castes in policing behavior. (a) Figure shows the number of

vibrations per minute performed by each individual (mean ± SE, n = 7). Time "0" was the time

of the first major attack. A major attack was recognized when an ergatoid visibly injured another

ergatoid. A one-minute sample (30 seconds before and after) was analyzed for each time point.

(b) The number of individuals surrounding the victim before and after the attack (mean ± SE, n =

7). The increased density of workers reflects cannibalism.

Figure 4. Age of interacting ergatoids in policing behavior. Each dot represents an individual attacker or victim, and lines connect the attacker to the corresponding victim. The negative slope of the lines in both graphs shows that an older ergatoid attacked a younger ergatoid in every replication, and the age differences are significant (female: $Z = -2.3664$, $P < 0.01$, n = 7; male: Z $= -2.5205$, P < 0.01, n = 8; Wilcoxon signed-rank test, one-tailed; **, $P < 0.01$).