# Genetic Disruption of Serine Biosynthesis is a Key Driver of Macular Telangiectasia Type 2 Etiology and Progression

Roberto Bonelli,[1,2] Brendan R E Ansell,[1,2] Luca Lotta,[3] Thomas Scerri,[1,2] Traci E Clemons,[4] Irene Leung,[5] The MacTel Consortium,[6] Tunde Peto,[7] Alan C Bird,[8] Ferenc Sallo,[9] Claudia Langenberg,[3] and Melanie Bahlo*[1,2].

Department of Medical Biology, The University of Melbourne, 3052, Parkville, Victoria, Australia.
Population Health and Immunity Division, Walter and Eliza Hall Institute of Medical Research, 3052, Parkville, Victoria, Australia.
MRC Epidemiology Unit, University of Cambridge, CB2 0SL, Cambridge, UK.
The EMMES Corporation, Rockville, 20850, Maryland, United States.
Department of Research and Development, Moorfields Eye Hospital NHS Foundation Trust, EC1V 2PD, London, United Kingdom.
A list of members and affiliations is provided in **Table S8**.
Department of Ophthalmology, Queen's University, Belfast, BT7 1NN, United Kingdom.
Inherited Eye Disease, Moorfields Eye Hospital NHS Foundation Trust, EC1V 2PD, London, United Kingdom.
Department of Ophthalmology, University of Lausanne, Hôpital Ophtalmique Jules-Gonin, Fondation Asile des aveugles, Switzerland.
* Correspondence: bahlo@wehi.edu.au, +61 03 9345 2555

# Abstract

**Purpose**

Macular telangiectasia type 2 (MacTel) is a rare, heritable and largely untreatable retinal disorder, often comorbid with diabetes. Genetic risk loci subtend retinal vascular calibre, and glycine/serine/threonine metabolism genes. Serine deficiency may contribute to MacTel via neurotoxic deoxysphingolipid production, however an independent vascular contribution is also suspected. Here we use statistical genetics to dissect the causal mechanisms underpinning this complex disease.

**Methods**

We integrated genetic markers for MacTel, vascular, and metabolic traits, and applied Mendelian randomization, MTAG, and conditional/interaction genome-wide association analysis to discover causal contributors to both disease, and spatial retinal imaging sub-phenotypes.

**Results**

Serine was a key causal driver of disease occurrence and progression, with a lesser contribution to type 2 diabetes risk. Conversely, glycine, threonine and retinal vascular traits are unlikely to be causal for MacTel. Conditional regression analysis resolved three novel disease loci independent of endogenous serine biosynthetic capacity. By aggregating retinal phenotypes into endophenotypes, we demonstrate that SNPs constituting independent risk loci act via related endophenotypes.

**Discussion**

Our findings will aid in early diagnosis and accurate prognosis of MacTel, and improve prospects for effective therapeutic intervention. Our integrative genetics approach also serves as a useful template for post-GWAS analyses in other disorders.

Retinal Disease | Mendelian Randomization | Metabolomics | GWAS | Serine

# Introduction

## MacTel Disease and previous GWAS study

Macular telangiectasia type II (MacTel [1]), is a rare degenerative eye disease affecting the macula [2,3]. MacTel is bilateral and progressively affects visual acuity [4,5], reading ability [6], and vision-related quality of life [7,8]. A successful phase II clinical trial has recently been reported for an intravitreal encapsulated cell therapy implant, showing efficacy in slowing disease progression [9]. However, less invasive, non-surgical and more economical therapies are lacking. Given the rarity of MacTel and its subtle clinical signs, often requiring several ophthalmological diagnostic methods, the disease has been largely under/misdiagnosed [4]. Hence further insight into the genetic basis of MacTel is key for accurate diagnosis, identifying future therapies, developing predictive models for the disease, and increasing prognostic accuracy.

We previously published the first genome-wide association study on 476 MacTel patients and 1733 controls [10], identifying and replicating five loci. A single nucleotide polymorphism (SNP) at locus 5q14.3 (rs73171800) showed the strongest association with the disease and was previously identified to be associated with the quantitative traits of retinal venular and arterial calibre [11,12]. The other four loci, 1p12 (rs477992), 2q34 (rs715), 7p11.2 (rs4948102), and 3q21.3 (rs9820286) were implicated in glycine and serine metabolism [13,14]. Importantly, loci 3q21.3 and 7p11.2 did not reach genome-wide significance, and the SNPs at locus 3q21.3 were only in proximity, but not in linkage disequilibrium (LD) with those associated with MacTel. In this GWAS we measured and found significant depletion of serum serine, glycine and threonine in MacTel patients compared to controls. These data provided the first insight into the genetic complexity underpinning MacTel, and highlighted the potential involvement of metabolic and vascular trait disturbances in disease aetiology.

Glycine and serine are essential amino acids involved in many fundamental biochemical reactions. Some of the most well-known biochemical pathways relying on glycine and serine are the sphingolipid metabolism pathway (KEGG:map00600) and the glycerophospholipid pathway (KEGG:map00564). Pathogenic variants in genes involved in serine and glycine synthesis lead to severe congenital disorders such as phosphoglycerate dehydrogenase deficiency (*PHGDH*, 601815) and glycine encephalopathy (*GLDC*, 605899). Glycine and serine can be synthesized from one another as well as from other metabolic compounds such as sarcosine and hydroxypyruvate. Additionally, they can both be obtained through dietary intake.

A recent study reported the co-occurrence of MacTel in patients with a rare neuropathy, hereditary sensory and autonomic systemic neuropathy type I,,orHSAN1 (162400) [15]. This neuropathy results from loss of function mutations in the enzyme serine palmityltransferase, leading to accumulation of neurotoxic deoxysphingolipids. A similar accumulation of deoxysphingolipids was demonstrated in MacTel patients, but unlike HSAN1, was attributed to systemic serine deficiency.

MacTel is often comorbid with type 2 diabetes (T2D) [16]. It is unknown whether traits such as
metabolite abundances, T2D, or retinal vascular calibre, now known to be associated with
MacTel, are causative or merely associated with the disease for other reasons. Causation
hypotheses are now routinely investigated using Mendelian Randomization (MR). MR borrows
from Instrumental Variable (IV) analysis and assumes that if a disease is caused by a particular
intermediate phenotype, and the latter is caused by genetic variants, then the same genetic
variants should be also associated with the disease [17–19]. Genetic variants involved in retinal
vascular calibre traits and T2D have been previously identified [11,12,20] and the largest
metabolomic GWAS meta-analysis to date recently identified almost 500 loci that are
associated with differences in 142 serum metabolite levels in humans [21,22]. The results from
these studies can be used to investigate causal drivers of MacTel susceptibility via the MR
approach.
112
GWA studies on rare complex traits such as MacTel are inevitably underpowered due to the
difficulty in recruiting a large number of samples. However, well-powered GWAS are now
collated in resources such as the European Bioinformatics Institute GWAS catalogue.
Methods leveraging such resources to aid underpowered GWA studies have now been
developed. An example is Multi-Trait Association GWAS (MTAG) [23] which exploits the genetic
correlation between traits to identify new loci.
119
Progress in MacTel diagnosis and treatment is complicated by a lack of clear indicators of
early stages of the disease and by the substantial inter-patient and interocular heterogeneity
of retinal phenotypes. Apportioning genetic variation to separate retinal malformations has the
potential to shed light on the different biological mechanisms via which each locus contributes
to the disease. Performing joint genetic and retinal phenotypic data can investigate whether
different retinal abnormalities are consequences of common or largely independent biological
perturbations.
127
In this study, we test for the causality of several traits on MacTel disease (**Figure 1A**). We
exploit independent studies on MacTel-related traits to increase discovery power in our
disease cohort, and both identify new disease loci while further resolving previously identified
loci (**Figure 1B**). We use extensive retinal phenotypic data from MacTel patients to identify
key genetic drivers of specific subphenotypes (**Figure 1C**). We collate genetic loci into
functional groups by testing their combined effects on ocular spatial phenotypes (**Figure 1D**).
This study represents the first integrated analysis of MacTel genotypic data with MacTel-
related traits as well as retinal imaging phenotypic data. Our study serves as a model for post-
GWAS studies with phenotyping datasets.

**Figure1:** Study conceptual map. Each panel represents a study aim and depicts the data and the analysis technique used. Data is presented as pastel color pieces with black typing while analysis techniques are presented as darker panels with white writing. MacTel genetics refers to individual level SNP data available from our previous GWAS study. MacTel GWAS refers to the MacTel GWAS summary statistics. Vasculature GWAS refers to the summary statistics data available from two previous retinal vasculature calibre genetic studies [11,12]. Metabolomics GWAS refers to summary statistics data available from a recently published metabolomics GWAS [21,22]. T2D GWAS refers to summary statistics data available from GWAS study on T2D [20]. MTAG: Multi-Trait Association GWAS; MR: Mendelian Randomization.

# Methods

Unless otherwise stated, all statistical and computational analyses were performed using the R statistical software, version 3.5.1. Corrected p-values less than 0.05 were considered statistically significant. For genome-wide association analyses, uncorrected p-values less than 5e-8 were considered genome-wide significant.

## DNA samples and SNP genotyping

Genotypic data was available for 476 MacTel patients and 1733 controls. SNPs were genotyped using the Illumina 5.0M chip as described previously [10]. Retinal phenotypic data was available for 455 patients MacTel patients. Genetic predictors (betas or odds ratios (ORs), with corresponding SNP ID and relevant allele) for metabolites and retinal vascular calibre, were provided by the corresponding authors of each study [11,12,21,22].

## Mendelian Randomization Procedures

To perform Mendelian randomization analysis on MacTel with metabolites, T2D and retinal vasculature we used the allele-score method for individual-level genetic data as described by Burgess et al. 2016 [24]. Using a p-value threshold of 5e-8 we extracted SNPs significantly associated with serum metabolite concentrations or other traits of interest (retinal venular calibre, retinal arteriolar calibre, and T2D). We estimated Genetically Predicted Metabolites (GPMs) and Genetically Predicted Traits (GPTs), for T2D and retinal vasculature (**Table S1**) for each subject by using the magnitudes of the significant SNPs as weights, in the manner of a polygenic risk score (**Supplementary Methods**). We used logistic regression models to test for association between GPMs, or GPTs and MacTel susceptibility. This analysis was corrected for genetically determined sex at birth and the first principal component to account for batch effects including population stratification,  as in our previous publication [10]. We used Benjamini-Hochberg multiple testing correction to control the false discovery rate threshold of 5%. A sequential conditional modelling approach was used to identify GPMs independently associated with the disease. Specifically, the most significant GPMs were iteratively added as covariates in the logistic regression model and re-tested until no GPM was nominally significant after FDR correction.

## Conditional and Interaction GWAS and MTAG analysis

Conditional GWAS were performed by including GPMs or GPTs in the logistic regression models as covariates. For the interaction GWAS, an interaction term was included for each SNP and the GPM or GPT combination of interest. Consistent with the original GWAS on this data, each model also included genetically determined sex at birth and the first principal component [10]. Analyses were performed using Plink v1.9 [25]. Results from the conditional and interaction GWAS were analysed using FUMA [26]. LocusZoom plots were produced using the LocusZoom software [27]. MTAG analysis was performed by integrating the available summary statistics for each trait of interest using the MTAG software [23].

## Retinal phenotype clustering

As part of the Natural History Observation and Registry studies of Macular Telangiectasia [28], longitudinal retinal phenotypic data (colour fundus images, optical coherence tomography and fluorescein angiographic images) was collected 2005-2015 from 1,716 patients (3,410 eyes). The data consisted of 143 spatial measurements of retinal phenotypes (**Table S2)**. A detailed

195  description of the corresponding methods can be found elsewhere [28]. Each phenotype was
196  measured in 9 different sub-fields of the retina (defined by the ETDRS grid [29] **Figure S1**). The
197  phenotype data was cleaned by performing missing data imputation (**Supplementary**
198  **Methods**). The cleaned dataset contained 119 phenotypes that were collapsed into 30
199  biologically relevant endophenotypes using factor analysis (**Supplementary Methods**).

## 200 Investigating the relationship of retinal endophenotypes with
## 201 genetic drivers

We tested for association between retinal endophenotypes and significant disease-associated
GPMs, GPTs and prioritised SNPs. The dataset containing retinal endophenotypes and
genetic information included 3,280 observations from 455 MacTel patients (907 eyes in total)
with an average of 3.6 observations per eye over 10 years. Association testing was performed
using a linear mixed model approach, assuming an additive effect of SNP alleles that
increased MacTel risk. P-values were corrected using an adaptive Benjamini-Hochberg
procedure [30]. Further details are provided in **Supplementary Methods**.
209

# 210 Results

## 212 Discovering key drivers of MacTel disease: Metabolites

The MR procedure (**Figure 1A**) applied to the metabolite panel highlighted 14 GP metabolites
significantly associated with MacTel **Table 1**. A list of all metabolite associations is provided
in **Table S3**. The two most significant metabolites were GP serine and GP glycine In contrast
to the metabolomics results from our previous metabolomics study with directly measured
metabolites, GP threonine was not significant. Other significant metabolites included GP
arginine, GP phenylalanine and GP phosphatidylcholine species. GP alanine was borderline
significant.
220

The significant GPMs correlated with each other in our sample (**Figure S2**), mirroring
correlations observed with directly measured metabolite abundances. When all GPMs were
re-tested in a model including GP serine as a covariate, GP glycine was the only associated
metabolite (**Table S3**) suggesting that the effect of GP glycine on MacTel is largely due to
biochemical interactions with serine. GP serine remained significant when included alongside
GP glycine. No other GP metabolites remained significant once GP serine and GP glycine had
been included (**Table S3**).
228
229

| GP Metabolite / Trait | Beta | OR | 95% C.I. | P-value | Biochemical Family |
|---|---|---|---|---|---|
| Serine | -0.65 | 0.52 | 0.46 - 0.58 | 8.38E-28 | Amino Acids and Biogenic Amines |
| Glycine | -0.58 | 0.56 | 0.50 - 0.63 | 5.44E-19 | Amino Acids and Biogenic Amines |
| lysoPhosphatidylcholine acyl C14:0 | 0.21 | 1.23 | 1.11 - 1.37 | 2.48E-03 | Glycerophospholipids |
| Phosphatidylcholine diacyl C32:1 | 0.21 | 1.23 | 1.11 - 1.37 | 2.48E-03 | Glycerophospholipids |
| Phosphatidylcholine diacyl C34:1 | 0.21 | 1.23 | 1.11 - 1.37 | 2.48E-03 | Glycerophospholipids |
| Phosphatidylcholine with acyl-alkyl residue C38:1 | 0.20 | 1.22 | 1.10 - 1.36 | 2.97E-03 | Glycerophospholipids |
| Arginine | 0.17 | 1.18 | 1.07 - 1.31 | 2.83E-02 | Amino Acids and Biogenic Amines |
| Phenylalanine | -0.17 | 0.85 | 0.76 - 0.94 | 2.85E-02 | Amino Acids and Biogenic Amines |
| Arteriolar Retinal Calibre | 0.32 | 1.38 | 1.25 - 1.52 | 2.22E-10 | Vasculature Trait |
| Venular Retinal Calibre | 0.18 | 1.19 | 1.08 - 1.32 | 7.00E-04 | Vasculature Trait |
| T2D | 0.17 | 1.19 | 1.07 - 1.33 | 7.72E-04 | Comorbidity |

**Table 1**: Significant associations between GPMs and GPT and MacTel. Regression coefficients are presented in the "Beta" column. Odds Ratios (OR) are relative to a single standard deviation increase on the GPM/GPT scale. Odds Ratios 95% confidence intervals are shown in the column 95% C.. Benjamini-Hochberg adjusted P-values are presented under the column P-value.

The GWAS used to predict the genetic complement of serine was underpowered compared to the glycine GWAS[22] (**Supplementary Materials**). However, biochemically-based genetic correlation exists between the two. Hence, we attempted to boost the power of the serine GWAS results by combining them with the glycine GWAS results in an MTAG analysis (**Figure 1A**). The newly identified MTAG serine loci were used to construct a new and more powerful genetic predictor of serine (GP MTAG serine, **Table S1**). We found that GP MTAG serine encompassed the entire metabolic predictive signal for MacTel, as the original GP glycine was no longer required in the model once GP MTAG serine was included. Further MR testing of a broader set of 248 GP metabolite abundances using results from Shin et al [14,31] (**Table S1**) found a few additional significant univariate contribution from other metabolites

248  (**Supplementary Materials** and **Table S4**). However, none remained significant after
249  inclusion of GP serine as a covariate.
250

## Discovering key drivers of MacTel disease: T2D and retinal vasculature

253
Genetically predicted T2D was significantly associated with MacTel, as were both GP
arteriolar and venular calibre when tested separately. However, the vasculature trait results
were driven by a single SNPs: rs2194025 (for arteriolar calibre) and rs17421627 (for venular
calibre), both in strong LD with SNP rs73171800 at 5q14.3 previously identified in our original
GWAS [10]. When including both vascular traits in the same model, only arteriolar calibre
remained significant, due to the reduced number of SNPs (N = 2) used to construct that GPT
(**Figure S3**). When GPTs for retinal arteriolar calibre and T2D were included in a model with
GP glycine and serine, all remained significant.
262
As a quality control analysis, we compared the effect sizes for each instrumental SNP on both
MacTel and the metabolites or traits to inspect for potential SNP outlier effects of pleiotropy
(**Figure 2**). A negative, linear relationship was detected between the effect on metabolic
abundance and MacTel risk for the SNPs used to define glycine and serine. A positive
relationship was observed for T2D and vasculature traits, while no relationship was detected
for threonine which was used as a control trait, where no significant association was observed.
We additionally observed that the regression line for all significant traits except the vasculature
related traits, did not have intercept terms that were significantly different from 0 ($b_{0,serine}$= −
0.07, p=0.2 ; $b_{0,glycine}$=0.070, p=0.131 ; $b_{0,T2D}$=0.002, p=0.88) suggesting a non-pleiotropic
effect of the instruments used to test such traits[32]. The same conclusion was drawn when
Glycine and T2D SNPs considered to have potential outlier effects were removed.
274
275

**Figure 2:** Effect size comparisons between MacTel and specific trait of SNPs used to define GPM and GPT. Each dot represents a SNP. Each panel represents a traits and contains the SNPs that were found to have a genome-wide significant effect on that trait. The x-axis captures the effect sizes of the SNPs on the trait while the y-axis captures the effect of the same SNPs on MacTel risk. The blue line represents a simple regression line between the two. Metabolites or traits causally affecting the disease are expected to have correlated effect sizes. Colour of the dots represents chromosomal positions where different chromosomes are represented by different colours.

# Using causal traits to discover new disease susceptibility loci for MacTel

## MTAG results

To discover new loci putatively involved in MacTel, we performed MTAG analysis by including GWAS summary statistics for glycine, serine, retinal venular calibre, and retinal arteriolar calibre along-side MacTel (**Figure 1B**). T2D was not included as genome-wide summary statistics were not available. The number of SNPs shared across all datasets was 799,436 (maximal FDR = 0.397). This analysis confirmed the three previously identified MacTel risk loci [10] and revealed 5 novel loci attaining genome-wide significance (**Table 2, Table S5** for details).

| rsid | Locus | Candidate Gene | MTAG P-val MacTel | New GW. | P-val Art | P-val Ven | P-val Gly | P-val Ser | P-val MacTel |
|------|-------|----------------|-------------------|---------|-----------|-----------|-----------|-----------|--------------|
| | | | | | Sim et al 2013 | Ikram et al 2010 | Lotta et al 2019 | Lotta et al 2019 | Scerri et al 2017 |
| rs3815630 | 2q34 | *CPS1* | **7.62E-143** | No | *1.51E-01* | *6.23E-01* | **2.23E-308** | **6.50E-31** | **4.43E-08** |
| rs1035387 | 5q14.3 | *TMEM161B-AS1* | **1.16E-16** | No | **1.50E-09** | **1.15E-08** | *5.43E-01* | *7.53E-01* | **1.16E-16** |
| rs477992 | 1p12 | *PHGDH* | **2.21E-15** | No | *7.44E-01* | *7.95E-04* | **6.09E-07** | **2.01E-80** | **2.60E-12** |
| rs4543497 | 7p11.2 | *PSPH* | **2.22E-15** | Yes | *3.56E-01* | *2.07E-01* | **9.01E-28** | **3.29E-54** | 8.11E-06 |
| rs4841132 | 8p23.1 | *LOC157273* | **8.43E-14** | Yes | *5.55E-02* | *2.95E-02* | **1.73E-34** | 1.23E-04 | 4.54E-02 |
| rs1563075 | 16q23 | *GCSH* | **1.79E-13** | Yes | *6.86E-01* | *3.77E-01* | **2.57E-28** | 1.19E-01 | 4.66E-03 |
| rs2954021 | 8q24.1 | *TRIB1* | **1.53E-12** | Yes | *3.89E-01* | *6.33E-01* | **2.35E-21** | **2.40E-10** | 8.70E-05 |
| rs4742212 | 9p24 | *GLDC* | **1.19E-09** | Yes | *4.73E-01* | *6.88E-01* | **3.06E-34** | 1.69E-04 | 2.82E-01 |

**Table 2**: Genome-wide significant loci for MacTel from MTAG analysis using retinal venular (Ven) and arteriolar (Art) calibre, serine (Ser) and glycine (Gly). Novel genome-wide significant loci are identified by the column New GW. Bold text indicates genome-wide significant P-values on respective original studies. Grey italicised p-values are associations expected to be null *a priori*.

Among the five new genome-wide significant loci was locus 7p11.2 which did not reach genome-wide significance in the original GWAS but was deemed important given its established relationship with glycine and serine [10]. We did not observe any significant MTAG results for locus 3q21.3, originally proposed to affect MacTel though a connection with glycine and serine [10]. This SNP was in fact not genome-wide significant for either glycine ($p = 0.041$) or serine ($p = 0.65$) and is likely to present either a false positive or a GWAS signal not contributed to by glycine or serine.

## Conditional and Interaction GWAS

To uncover genetic correlates of MacTel independent of glycine and serine, we performed a GWAS conditioning on these genetically predicted traits (**Figure 1B**). This analysis identified four genome-wide significant peaks (**Figure S4 a)**.

As expected, the original signal on locus 5q14.3 (rs73171800) related to vascular calibre, remained significant, confirming the independence of this locus from genetic drivers of serine and glycine. A second genome-wide significant signal on locus 3p24.1 (rs35356316, $p = 3.10\text{e-}08$) was identified and was situated in a 'gene desert' proximal to the genes *EOMES*

322   and *SLC4A4*. Another single SNP in very close proximity to rs35356316 reached genome-
323   wide significance as in the original MacTel GWAS study [10] but was believed to be a false
324   positive, given the lack of LD with any other significant SNP (**Figure S4 b**).
325
The remaining two conditionally significant SNPs tagged independent signals in locus 19p13.2
(**Figure S4 c-d**). SNP rs36259 is an exonic non-synonymous SNP located in the *CERS4* gene
which achieved close to genome-wide significance (p=6.270e-08) and was nominally
significant in our original study (p = 1.69e-7). In close proximity, we found an independent
intergenic SNP rs4804075 (p = 3.72e-07) which lacked evidence of an eQTL effect on any
neighbouring genes, and which did not reach genome-wide significance. These results remain
significant when conditioning on SNP rs73171800 (locus 5q14.3) and GP T2D.
333
334   We performed additional GWAS analyses testing for interactions between all SNPs with GP
335   serine, glycine, T2D, and SNP rs73171800 (**Figure 1B**). However, no further significant
336   interacting loci were found (**Figure S5 a-d**).
337

# Effect of key drivers on retinal endophenotypes

Given the small sample size of the endophenotypic data (N = 455), we decided to only test for
association between endophenotypes and GP glycine, serine and T2D, as these bore the
clearest association with disease etiology (**Figure 1C, Table S6**). No significant association
between GP glycine or T2D and endophenotypes was found after correcting for GP serine,
and we, therefore, excluded them from additional testing. High GP serine levels were found to
be protective against loss of retinal transparency, leakage at the outer capillary network and
the retinal pigment epithelium (RPE) in MacTel progression areas, and leakage at the RPE in
the MacTel area. We also found suggestive significance for protection from perivascular
pigment clustering in the MacTel area and progression area. Higher GP serine levels were
associated with protection from macular thinning in the MacTel area, inferior inner area, and
suggestive protection on nasal inner area, and foveal area. Together these results highlight
that GP serine is associated not only with MacTel disease risk but also with those retinal
endophenotypes characterising disease progression.
352
353

**Figure 3:** Association plot summarising all nominally significant association between MacTel SNPs and the
endophenotypes (**Table S6**). SNPs are divided into categories based on MR results. Each dot represents the
nominal p-value for each association between all SNPs and endophenotypes. Red dots indicate that MacTel risk
alleles for that SNP also increases the risk of the retinal phenotype, while blue dots indicate a reduced risk. The
dashed line represents the nominal significance of p=0.05. The top right corner displays the ETDRS grid for the
right eye (OD) [29] used to divide the endophenotypes into 9 retinal areas; 1: foveal area, 2: superior inner area, 3:
nasal inner area, 4: inferior inner area, 5: temporal inner area, 6: superior outer area, 7: nasal outer area, 8: inferior
outer area, and 9: temporal outer area. Although definitions vary across the literature, for simplicity in this
manuscript we define the pink area as the 'MacTel [onset] area' while the green areas are MacTel 'progression
areas'.

## Determining the effects of GWAS loci on endophenotypes

To determine the likely functional impacts of the original GWAS loci we tested for associations with retinal endophenotypes (**Figure 1D, Figure 3, Table S6**).In No associations remained significant after correction for multiple testing, likely due to the modest sample and effect sizes of individual SNPs on disease endophenotypes. However, using a nominally (uncorrected) significant p-value threshold denoted p* < 0.05 in an exploratory approach, we observed that locus 1p12 (rs477992) was associated with an increased risk of perivascular pigment clustering in the progression area, but protected against risk of blunted vessels. Furthermore, locus 2q34 (rs715) increased the risk of macular thinning in the MacTel area and nasal inner area and the risk of perivascular pigment clustering in the MacTel area. Finally, the MacTel risk allele at locus 7p11.2 (rs4948102) increased the risk of retinal transparency, leakage at the RPE in both the MacTel area and progression area, as well as presence of inner empty spaces and EZ break. This locus however protected against leakage in the outer layer of the foveal area.

These three loci, which converge on glycine/serine metabolism (1p12/rs477992/*PHGDH*, 2q34/rs715/*CPS1*, and 7p11.2/rs4948102/*PSPH*) all demonstrated an overall susceptibility increment to RPE endophenotypes and other macular phenotype risks, lending further support for their role, despite not attaining significance after p-value adjustment.

Interestingly the MacTel risk allele at locus 3q21.3 (rs9820286) was only associated with protection from retinal disease endophenotypes. Specifically, locus 3q21.3 was associated with protection against retinal holes, retinal detachment, perivascular pigment clustering in the MacTel area and progression areas; leakage at the RPE level in the MacTel area and progression area; and foveal slope flattening in MacTel area, inferior inner area, and nasal inner area.

Locus 5q14.3 (rs73171800) corresponded to protection against macular thinning in the foveal area and nasal-inferior outer areas and increased risk of foveal flattening in the nasal inner area. The effect of SNP rs73171800 on macular thickness was consistent with recent findings [33]. We attempted to cluster the loci using hierarchical clustering of their regression coefficients (**Figure S7**). This analysis highlighted the unique endophenotype effect profiles of the 5q14.3 (rs73171800) and 3q21.3 (rs9820286) loci, and their distinctiveness in comparison to the other three loci, known to act on the serine/glycine pathway.

# Discussion

This study exploited well-powered publicly available GWAS data from traits of interest as well as deep retinal phenotyping data to further investigate the genetic aetiology of MacTel, a moderately rare retinal disorder for which the first GWAS was only recently performed [10]. We report causal genetic drivers for MacTel, delineating independent genetically-determined

412 metabolic contributors to the disease, and quantifying the contribution of MacTel-related traits
413 and genetic risk loci to retinal degeneration.
414

415 Mendelian randomization and publicly available data were used to "genetically predict"
416 abundances of different metabolites. Even though all GPMs in this study were generated using
417 estimated SNP allele dosage effects from blood-based metabolomic studies rather than retina,
418 we believe that our results represent a useful approach for interrogating other retinal disorders.
419

We found genetically-predicted serine depletion as the strongest causal driver of MacTel risk,
with an association magnitude much greater than those observed when using single SNPs.
This pronounced effect reflects the power of combining multiple genetic variants associated
with the disease through a shared biological mechanism; and further emphasizes the
causative role of serine in this disease. The direction of association between GP serine and
MacTel agrees with results from direct serum serine measurements reported previously [10].
This provides compelling additional evidence for a causal association between serine
deficiency and MacTel. Indeed, the MacTel disease odds doubled for every standard unit
decrease in GP serine, and individuals in the lowest quintile of GP serine presented an OR of
6.34 for MacTel compared to individuals in the top quintile. Our study results also confirm the
pronounced role that serine plays in disease progression. As this analysis was applied only to
MacTel patients, the heterogeneity of retinal phenotypes at early disease stages may be
under-represented. GP serine was nevertheless able to partly discriminate between subjects
with more advanced retinal abnormalities and those without. For example, lower GP serine
had a clear association with retinal greying in all retinal areas, which is not observed in all
MacTel patients [4]. GP serine also affected retinal thinning in the temporal parafoveal area, a
marker of photoreceptor degeneration. Although our results suggest a highly plausible
biochemical explanation for differences in disease heterogeneity and progression, we
acknowledge that our study used only aggregated retinal phenotypes and proxy measures of
progression. Longitudinal data documenting progression is now required to confirm these
findings. High concentrations of deoxy-sphingolipids, a byproduct of serine deficiency, have
recently been shown to cause MacTel [15]. Our results provide evidence that genetically-
encoded serine depletion is causal for MacTel, which likely contributes to the disease by
promoting deoxy-sphingolipid biosynthesis [15].
444

Our results indicate that the association between glycine and MacTel is likely an artefact of
the shared genetic signal between this metabolite and serine. This may relate to the
considerably greater sample size used to construct the glycine GP, and/or its metabolic
dependence on serine. Similarly, the previously observed significant association of threonine
to MacTel disease risk [10] is likely due to its metabolic dependence on glycine and serine and
thus a bystander effect.
451

Multi-Trait Association GWAS (MTAG) confirmed the central role of locus 7p11.2 (rs4543497),
which was previously found to be only nominally significant, and identified a new locus, 8q24.1
(rs2954021). The latter locus is of great interest since rs2954021 is predictive of endogenous
serum serine concentrations [22] and thus might confer MacTel risk by contributing to the same
serine pathway as previously identified SNPs. Conversely, we did not find any additional signal
arising from shared genetic correlations with retinal vascular calibre traits.
459

Although not significant after correction for multiple testing, we observed a pronounced effect
of SNP rs73171800 on macular thickness, specifically, a protective effect of the C allele
against macular thinning, which is tantamount to an increase in macular thickness.
Interestingly, a large GWAS study previously found that the G allele of SNP rs17421627 -
corresponding to the C allele of SNP rs73171800 (LD with rs73171800 $r^2$=0.67), was
significantly associated with increased macular thickness [33]. Further, a targeted study of SNP
rs17421627, leveraged the deep conservation of this locus to demonstrate enhancer activity
which modifies the retinal vasculature. Madelaine et al substituted the homologous zebrafish
locus with a construct containing rs17421627, and showed enhancer activity and changed
expression of a proximal microRNA, mir-9-2 (homologous with human mir-9-5) [34]. Both mir-9-
2 knock-down and endogenous enhancer knock-out animals showed dysmorphic retinal
vasculature, indicating that rs17421627 may act on miRNA expression to modify the formation
of the retinal vasculature in humans[33] Although our MR analysis did not find a causal
relationship between retinal vasculature calibre and MacTel, it may be that other features of
the vasculature, for example, leakage, branching or integrity, are modified in rs17421627
carriers, and account for the gross differences in macular thickness. Further deep phenotyping
of the retinal vasculature in these individuals may yield the physiological basis for this
phenotype and its impact on MacTel.
478
By conditioning on GP serine, glycine and T2D we revealed independent genetic contributors
to MacTel. The locus 3p24.1, tagged by SNP rs35356316, lies between *EOMES* (604615) and
*SLC4A4* genes (603345). Interestingly, *EOMES* encodes a transcriptional activator which is
shown in mouse studies to interact with the retinal transcription factor *Pou4f2* [35]. The
downstream gene *SLC4A4* encodes a sodium-coupled bicarbonate transporter which is
expressed in Müller glia and the RPE, and functions to balance pH in the subretinal space [36].
485
486 Conditional analysis further revealed SNP rs36259 at locus 19p13.2 which tags an exonic,
487 non-synonymous SNP located within the gene *CERS4* (615334), encoding a dihydroceramide
488 synthase involved in sphingolipid biosynthesis [37]. Just as serine depletion is associated with
489 defective sphingolipid synthesis, this locus may reduce sphingolipid production and thus
490 contribute to MacTel.
491
Type 2 diabetics are over-represented among MacTel patients [16]. We found a weak positive
association between GP T2D and MacTel risk. A possible explanation for this is that T2D
involves major perturbations of patient metabolism. Specifically, a recent meta-analysis of
metabolite abundances associated with pre-diabetes and/or T2D, found that glycine depletion
tends to occur in this disease [38]. We show that glycine depletion is not likely to be causative
for MacTel and is likely to be a consequence of metabolomic co-regularisation
counterbalancing the genetically induced serine depletion. We speculate that diabetes may
be a consequence rather than a cause of the metabolic phenotype underlying MacTel. Future
MR studies may further dissect this interaction but will require a more highly powered MacTel
GWAS. Indeed, T2D risk remained significant even after controlling for GP serine and GP
glycine, which indicates the possibility of a separate mechanism unrelated to glycine/serine
metabolism.
504
505 Our analysis revealed that locus 3q21.3, which was previously believed to act on the disease
506 by modifying serum glycine and serine, is instead likely disconnected from these metabolites,

507 as is the case for loci 3p24.1 and 5q14.3. Furthermore, these three loci do not cluster with
508 each other, implying that additional potential disease mechanisms exist.
509
To conclude, by integrating different sources of genetic and phenotypic data related to macular
telangiectasia type II with advanced statistical methods, we find evidence that genetically-
encoded serine depletion causes disease onset and likely also progression. Further targeted
studies of disease progression and experimental validation of serine-independent
contributions will resolve the complex aetiology of MacTel and to enhance treatment efficacy.
The methods applied here should also serve as a model for dissecting GWAS results,
especially for rarer traits.
517
518

## 519 Acknowledgments:

We would like to acknowledge the funding support from the Lowy Medical Research Institute.
This work was also made possible through the Victorian State Government Operational
Infrastructure Support and Australian Government National Health and Medical Research
Council (NHMRC) independent research Institute Infrastructure Support Scheme (IRIISS). RB
was supported by the Melbourne International Research Scholarship. BREA was supported
by an NHMRC early career Fellowship (1157776). MB was supported by an NHMRC Senior
Research Fellowship (1102971) and Program Grant (1054618). We are grateful to Dr Saskia
Freytag, Dr Anna Quaglieri, Prof Terry Speed, Prof Gordon Smyth, Dr Mari Gantner, Dr Kevin
Eade, Dr Martina Wallace, A/Prof Christian Metallo, Prof Martin Friedlander, Dr Tjebo Heeren,
Dr Mali Okada, Dr Sasha Woods, Prof Marcus Fruttiger, and Dr Catherine Egan for their
extremely helpful contributions that greatly improved the quality of this manuscript. We are
especially grateful to Dr Xueling Sim and Prof Wong Tien Yin for kindly sharing their summary
data on retinal vascular calibre traits.
533

## 534 Declaration of Interests

535 The authors declare no conflict of interest.
536

## 537 Web Resources

538 FUMA: http://fuma.ctglab.nl/
539 Locus zoom: http://locuszoom.org/
540 Metabolomics GWAS server: http://metabolomics.helmholtz-muenchen.de/gwas/
541

## 542 References

1.   Gass J. Some problems in the diagnosis of macular diseases. In: *Symposium on*
*Retinal Diseases*. ; 1977:268-270.

2.   Klein R, Blodi BA, Meuer SM, Myers CE, Chew EY, Klein BEK. The prevalence of
macular telangiectasia type 2 in the Beaver Dam eye study. *Am J Ophthalmol*.
2010;150(1):55-62.e2. doi:10.1016/j.ajo.2010.02.013

3.   Aung KZ, Wickremasinghe SS, Makeyeva G, Robman L, Guymer RH. The prevalence
estimates of macular telangiectasia type 2: the Melbourne Collaborative Cohort Study.
*Retina*. 2010;30(3):473-478. doi:10.1097/IAE.0b013e3181bd2c71

4.   Charbel Issa P, Gillies MC, Chew EY, et al. Macular telangiectasia type 2. *Prog Retin*
*Eye Res*. 2013/5;34:49-77. doi:10.1016/j.preteyeres.2012.11.002

5.   Gass JD, Blodi BA. Idiopathic juxtafoveolar retinal telangiectasis. Update of
classification and follow-up study. *Ophthalmology*. 1993;100(10):1536-1546.
https://www.ncbi.nlm.nih.gov/pubmed/8414413.

6.   Finger RP, Charbel Issa P, Fimmers R, Holz FG, Rubin GS, Scholl HPN. Reading
performance is reduced by parafoveal scotomas in patients with macular telangiectasia
type 2. *Invest Ophthalmol Vis Sci*. 2009;50(3):1366-1370. doi:10.1167/iovs.08-2032

7.   Lamoureux EL, Maxwell RM, Marella M, Dirani M, Fenwick E, Guymer RH. The
longitudinal impact of macular telangiectasia (MacTel) type 2 on vision-related quality of
life. *Invest Ophthalmol Vis Sci*. 2011;52(5):2520-2524. doi:10.1167/iovs.10-6568

8.   Clemons TE, Gillies MC, Chew EY, et al. The National Eye Institute Visual Function
Questionnaire in the Macular Telangiectasia (MacTel) Project. *Invest Ophthalmol Vis*
*Sci*. 2008;49(10):4340-4346. doi:10.1167/iovs.08-1749

9.   Chew EY, Clemons TE, Jaffe GJ, et al. Effect of Ciliary Neurotrophic Factor on Retinal
Neurodegeneration in Patients with Macular Telangiectasia Type 2: A Randomized
Clinical Trial. *Ophthalmology*. October 2018. doi:10.1016/j.ophtha.2018.09.041

10.  Scerri TS, Quaglieri A, Cai C, et al. Genome-wide analyses identify common variants
associated with macular telangiectasia type 2. *Nat Genet*. February 2017.
doi:10.1038/ng.3799

11.  Ikram MK, Sim X, Xueling S, et al. Four novel Loci (19q13, 6q24, 12q24, and 5q14)
influence the microcirculation in vivo. *PLoS Genet*. 2010;6(10):e1001184.
doi:10.1371/journal.pgen.1001184

12.  Sim X, Jensen RA, Ikram MK, et al. Genetic loci for retinal arteriolar microcirculation.
*PLoS One*. 2013;8(6):e65804. doi:10.1371/journal.pone.0065804

13.  Xie W, Wood AR, Lyssenko V, et al. Genetic variants associated with glycine
metabolism and their role in insulin sensitivity and type 2 diabetes. *Diabetes*.
2013;62(6):2141-2150. doi:10.2337/db12-0876

14.  Shin S-Y, Fauman EB, Petersen A-K, et al. An atlas of genetic influences on human
blood metabolites. *Nat Genet*. 2014;46(6):543-550. doi:10.1038/ng.2982

15.  Gantner ML, Eade K, Wallace M, et al. Serine and Lipid Metabolism in Macular Disease
and Peripheral Neuropathy. *N Engl J Med*. September 2019.
doi:10.1056/NEJMoa1815111

16.  Clemons TE, Gillies MC, Chew EY, et al. Medical characteristics of patients with
macular telangiectasia type 2 (MacTel Type 2) MacTel project report no. 3. *Ophthalmic*

*Epidemiol*. 2013;20(2):109-113. doi:10.3109/09286586.2013.766757

17.   Davey Smith G, Hemani G. Mendelian randomization: genetic anchors for causal
inference in epidemiological studies. *Hum Mol Genet*. 2014;23(R1):R89-R98.
doi:10.1093/hmg/ddu328

18.   Ebrahim S, Davey Smith G. Mendelian randomization: can genetic epidemiology help
redress the failures of observational epidemiology? *Hum Genet*. 2008;123(1):15-33.
doi:10.1007/s00439-007-0448-6

19.   Evans DM, Davey Smith G. Mendelian Randomization: New Applications in the Coming
Age of Hypothesis-Free Causality. *Annu Rev Genomics Hum Genet*. 2015;16:327-350.
doi:10.1146/annurev-genom-090314-050016

20.   Morris AP, Voight BF, Teslovich TM, et al. Large-scale association analysis provides
insights into the genetic architecture and pathophysiology of type 2 diabetes. *Nat*
*Genet*. 2012;44(9):981-990. doi:10.1038/ng.2383

21.   Wittemans LBL, Lotta LA, Oliver-Williams C, et al. Assessing the causal association of
glycine with risk of cardio-metabolic diseases. *Nat Commun*. 2019;10(1):1060.
doi:10.1038/s41467-019-08936-1

22.   Lotta LA, Pietzner M, Stewart ID, et al. Cross-platform genetic discovery of small
molecule products of metabolism and application to clinical outcomes. *bioRxiv*.
February 2020:2020.02.03.932541. doi:10.1101/2020.02.03.932541

23.   Turley P, Walters RK, Maghzian O, et al. Multi-trait analysis of genome-wide
association summary statistics using MTAG. *Nat Genet*. 2018;50(2):229-237.
doi:10.1038/s41588-017-0009-4

24.   Burgess S, Dudbridge F, Thompson SG. Combining information on multiple
instrumental variables in Mendelian randomization: comparison of allele score and
summarized data methods. *Stat Med*. 2016;35(11):1880-1906. doi:10.1002/sim.6835

25.   Purcell S, Neale B, Todd-Brown K, et al. PLINK: a tool set for whole-genome
association and population-based linkage analyses. *Am J Hum Genet*. 2007;81(3):559-
575. doi:10.1086/519795

26.   Watanabe K, Taskesen E, van Bochoven A, Posthuma D. Functional mapping and
annotation of genetic associations with FUMA. *Nat Commun*. 2017;8(1):1826.
doi:10.1038/s41467-017-01261-5

27.   Pruim RJ, Welch RP, Sanna S, et al. LocusZoom: regional visualization of genome-
wide association scan results. *Bioinformatics*. 2010;26(18):2336-2337.
doi:10.1093/bioinformatics/btq419

28.   Clemons TE, Gillies MC, Chew EY, et al. Baseline characteristics of participants in the
natural history study of macular telangiectasia (MacTel) MacTel Project Report No. 2.
*Ophthalmic Epidemiol*. 2010;17(1):66-73. doi:10.3109/09286580903450361

29.   Mathew R, Sivaprasad S, Florea D, et al. Agreement between time-domain and
spectral-domain optical coherence tomography in the assessment of macular thickness
in patients with idiopathic macular telangiectasia type 2. *Ophthalmologica*.
2013;230(3):144-150. doi:10.1159/000353455

30.    Benjamini Y, Krieger AM, Yekutieli D. Adaptive linear step-up procedures that control
628           the false discovery rate. *Biometrika*. 2006;93(3):491-507. doi:10.1093/biomet/93.3.491

629    31.    metabolomics gwas server. http://metabolomics.helmholtz-muenchen.de/gwas/.
630           Accessed December 4, 2018.

631    32.    Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid
632           instruments: effect estimation and bias detection through Egger regression. *Int J*
633           *Epidemiol*. 2015;44(2):512-525. doi:10.1093/ije/dyv080

634    33.    Gao XR, Huang H, Kim H. Genome-wide association analyses identify 139 loci
635           associated with macular thickness in the UK Biobank cohort. *Hum Mol Genet*.
636           December 2018. doi:10.1093/hmg/ddy422

637    34.    Madelaine R, Notwell JH, Skariah G, et al. A screen for deeply conserved non-coding
638           GWAS SNPs uncovers a MIR-9-2 functional mutation associated to retinal vasculature
639           defects in human. *Nucleic Acids Res*. March 2018. doi:10.1093/nar/gky166

640    35.    Mao C-A, Kiyama T, Pan P, Furuta Y, Hadjantonakis A-K, Klein WH. Eomesodermin, a
641           target gene of Pou4f2, is required for retinal ganglion cell and optic nerve development
642           in the mouse. *Development*. 2008;135(2):271-280. doi:10.1242/dev.009688

643    36.    Pushkin A, Abuladze N, Lee I, Newman D, Hwang J, Kurtz I. Mapping of the human
644           NBC3 (SLC4A7) gene to chromosome 3p22. *Genomics*. 1999;57(2):321-322.
645           doi:10.1006/geno.1999.5781

646    37.    Riebeling C, Allegood JC, Wang E, Merrill AH Jr, Futerman AH. Two mammalian
647           longevity assurance gene (LAG1) family members, trh1 and trh4, regulate
648           dihydroceramide synthesis using different fatty acyl-CoA donors. *J Biol Chem*.
649           2003;278(44):43452-43459. doi:10.1074/jbc.M307104200

650    38.    Guasch-Ferré M, Hruby A, Toledo E, et al. Metabolomics in Prediabetes and Diabetes:
651           A Systematic Review and Meta-analysis. *Diabetes Care*. 2016;39(5):833-846.
652           doi:10.2337/dc15-2251

653
654
655

# Supplementary Material: Genetic Disruption of Serine Biosynthesis is a Key Driver of Macular Telangiectasia Type 2 Etiology and Progression

## Supplementary Images:

**Figure S1:** Right (OD) and left (OS) eyes macula, were divided into 9 areas according to the ETDRS grid [1]. 1: foveal area, 2: superior inner area (progression area), 3: nasal inner area (progression area), 4: inferior inner area (progression area), 5: temporal inner area (MacTel area), 6: superior outer area, 7: nasal outer area, 8: inferior outer area, and 9: temporal outer area.

678
679
680 **Figure S2:** Correlation between GPM scores for all GPM scores that achieved multiple testing
681 corrected significance (p < 0.05). Bigger dots and saturated colour indicate stronger
682 correlations. Blue indicates a positive correlation while red indicates a negative one.

688
**Figure S3:** Comparison between SNPs effect on vasculature traits and MacTel. Each dot is a
SNP which was found to be significantly associated with a vasculature trait. The x-axis
represents the effect that the SNP has on the particular trait while the y-axis displays the effect
that the SNP has on MacTel. Most SNPs have no effect on MacTel. The only two SNPs that
have an effect are the SNPs rs2194025 and rs17421627 which are very close to each other
and in strong LD with SNP rs73171800 ($r^2$=0.94 and $r^2$=0.67).

**Figure S4:** MacTel GWAS results conditioned by GP serine and GP glycine. A) Manhattan plot. B) Locus zoom plot of the significant signal on locus 3p24.1. C) Locus zoom plot of the significant signal by SNP rs36259 on locus 19p13.2. D) Locus zoom plot of the significant signal by SNP rs4804075 on locus 19p13.2.

715
716 **Figure S5**: MacTel interaction GWAS results. A) Manhattan plot showing SNP interactions
717 with GP serine, B) Manhattan plot showing SNP interactions with GP glycine, C) Manhattan
718 plot showing SNPs interaction with SNP rs73171800 in locus 5q14.3, D) Manhattan plot
719 showing SNP interactions with GP T2D.

726　**Figure S6:** Visual representation of the endophenotypes. A: Endophenotypes created from
727　OCT scans. B: Endophenotypes created from fundus reflectance colour scans. C:
728　Endophenotypes created from fundus fluorescein angiography. In this figure each
729　endophenotype has a unique colour, the same colour in different images modality indicate
730　different endophenotypes.

738
739　**Figure S7:** Loci clustering heatmap. This clustering heatmap visualises the regression
740　coefficients of each SNP on the endophenotypes. Rows and columns of the heatmaps were
741　ordered according to the hierarchical clustering to give a visual representation of the
742　similarities. SNP rs73171800 (5q14.3) and SNP rs9820286 (3q21.3) have very different
743　effects from the other four SNPs which show a more consistent, shared pattern.

# Supplementary Tables:

# Supplementary Methods:

## GPM score calculation

We used the formula $GP(M/T)_i = \sum_{j \in i} m_j * SNP_j$, where $GP(M/T)_i$ is the genetically predicted metabolite ($M$) or trait, $Ti$. $SNP_j$ is the $jth$ SNP that was significantly associated with the metabolite or trait $i$. $m_j$ is the magnitude (weight) that the $SNP_j$ has on the metabolite or trait. The weight $m_j$ was obtained as $m_j = \beta_j$ where $\beta_j$ is the linear coefficient of the regression of $SNP_j$ on the metabolite or trait $i$. We used a total of 499 SNPs to predict the abundances of 142 metabolites in blood serum, 37 SNPs to construct a genetically predicted T2D risk, 5 for retinal venular calibre and 2 for the retinal arteriolar calibre. GPMs and GPTs were normalized to have a mean of zero and unit standard deviation.

# Imputation of phenotypic data

The phenotyping dataset contained multiple missing data points. Most missingness was related to imperfections in the ophthalmological images. By assuming missingness at random we imputed the missing data using predictive mean matching through chained equations, available in the package MICE [2].

# Generation of Endophenotypes with Factorial Analysis

The phenotypic data was summarised into endophenotypes. Endophenotypes have been defined as "measurable components unseen by the unaided eye along the pathway between disease and distal genotype"[3]. To identify and construct these underlying endophenotypes while ensuring that each only contained phenotypes strongly related to each other, we used an iterative factorial analysis algorithm. Given the very different nature of the phenotypes observed we performed two separate analyses for the creation of the endophenotypes one to capture macular thickness endophenotypes, and the other to capture macular abnormality phenotypes.

Starting with 43 phenotypic measurements for macular thickness and 76 for macular abnormality phenotypes we reduced these two data sets further, by only retaining correlated phenotypes within each dataset.
We discarded all those phenotypes/variables which:

- Had an absolute value of Spearman correlation with all other variables less than 0.3.
- Had an absolute value of Spearman correlation with the overall dataset score of less than 0.3.
- Would decrease the Cronbach's alpha of the dataset

This procedure retained 33 phenotypes for macular thickness and 52 phenotypes for macular abnormalities. Variables that were retained after this step underwent an iterative factorial analysis procedure:

1. Perform a *parallel analysis* on the dataset to discover the number of factors $F$ needed to summarise the phenotypic data
2. Run a *factorial analysis* with $F$ number of factors
3. Rotate the factors using the *oblimin* rotation to allow for correlation between endophenotypes and obtain a loading value for each phenotype in each factor.
4. Create a sequence of loading thresholds $T$ from 0.2 to 0.5 by 0.01
5. For each loading threshold $t \in T$ and for each factor $f \in F$, create a dataset that contains all the phenotype that have a loading greater than $t$. This results in F datasets.
6. Calculate the Cronbach's alpha in each dataset assessing the reliability of each factor.
7. Calculate the average of all factor's alphas
8. Select the loading threshold $t*$ that maximize the average alpha.

841   9. Eliminate from the dataset all the variables that do not have any loading greater than
842     $t*$.
843   10. Eliminate all variables which create factors by themselves.
844   11. Repeat 1-10 the entire process until no variables are removed.

846 This procedure was followed by an additional step, where we took the phenotypes that were
847 discarded from the previous step and re-introduced into the endophenotype creation
848 process, by themselves. Those phenotypes that were discarded a second time,  were
849 manually collapsed into meaningful new endophenotypes from previous clinical knowledge.
850 These endophenotypes were obtained by simply summing the phenotypes values together.

852 Each factor created by this process was considered an endophenotype in further analysis. All
853 endophenotypes resulted in continuous variables for which higher values represented
854 negative impacts on patient's eye health. Specifically, higher values of macular thickness
855 endophenotypes represented a risk of macular thinning, higher values for foveal slopes
856 endophenotypes represented increase risk of foveal flattening, and higher values for macular
857 abnormalities endophenotypes represented an increased risk of observing that particular
858 endophenotype. Most of the analyses to obtain the endophenotypes were performed using
859 the R package Psych[4].

861 Following the method used by [5], we also created foveal slope measurements from each
862 parafoveal area using the macular thickness endophenotypes. The formula used is was
863 $Slope_j = tan^{-1}((Thick_j - Thick_1)/500)$. Where $Slope_j$ indicates the slope from the fovea to
864 the parafoveal area $j$ and $Thick_j$ is the endophenotype of macular thickness in area $j$. Since the
865 foveal slope is defined as the slope from the fovea to the parafoveal areas the only areas used
866 were area 2, area 3, area 4, and area 5 of the ETDRS grid.

869 The factorial analysis results in 9 endophenotypes describing macular thickness and 19
870 describing macular abnormalities. We discarded one macular thickness endophenotype and
871 one macular abnormalities endophenotype since they were believed to describe only a
872 residual correlation between phenotypes and have no real clinical relevance. Additionally, we
873 obtained 4 endophenotypes describing macular slope leading to a total of 30 endophenotypes
874 to be used in the analysis.

# Testing the identified disease key drivers and SNP loci against the endophenotypes

880 Endophenotypes were normalized to have zero average and unit standard deviation. Given
881 the longitudinal nature of the data, we chose a linear mixed model approach to take into
882 account the relationships within the subject and eye observation. Each model included all
883 factors of interest as well as time defined as the number of days from the initial visit date, a
884 quadratic function of visit year, gender, and age as covariates. Each model also contained a

random intercept and random time slope for each subject as well as a random intercept and
random time slope for each eye within each subject. We included all these random effects
since the log-likelihood of models that contained them and models which did not were
significantly different (results not shown. Models that compared fixed effects were estimated
using restricted maximum likelihood estimation. Given the large sample size, the p-values
were obtained by assuming the asymptotic normality property of the t-values for each
association. To take into account false discovery rate we used an adaptive Benjamini-
Hochberg procedure [6]. We adopted this procedure rather than strict Bonferroni correction
because of the very conservative nature of the linear mixed model and the small magnitude
of SNP effect sizes.
895
896
897

# Supplementary Results:

## Using MTAG to increase discovery power on GP serine

The much higher powered discovery GWAS for glycine and serine were used to predict glycine
and serine GPMs. The samples sizes for these two traits were different: (nGlycine = 79,293
and nSerine=30,68). For this reason, we tried to increase the discovery power of the serine
GWAS by using MTAG and combining serine GWAS results with the glycine GWAS results,
leveraging the genetic correlation between them (**Figure 1A**). Although MTAG estimated the
genetic correlation between the two traits to be high (0.413), the maximal FDR for serine was
also high (0.246). This was most likely caused by the difference in sample sizes of the two
studies. By using the results from this analysis, we constructed GP MTAG serine and tested
it for significance alongside GP glycine on MacTel. This resulted in a slight decrease of
significance for GP MTAG serine (p=6.24e-13) and a complete loss in significance for the
glycine original GPM (p=0.907).
912

## MR using Shin et al 2014 metabolomics study:

We also tried to perform MR testing on all metabolites available from one of the first studies
on the genetic impact on metabolomics abundances by Shin et al 2014 [7,8] which contained
many more metabolites than those initially available to us. Of the 248 GPMs available we
found few additional metabolites to be significant (**Table S4**), None of them was as strongly
associated as GP glycine and GP serine and none retained significance after their inclusion
as covariates.

## Composition of the endophenotypes

An illustrative image of the endophenotypes is presented in **Figure S6** while a detailed list of
phenotypes and respective loadings values used to estimate each of the endophenotypes is
available in **Supplementary Materials** and **Table S7**. Interestingly, among the 30 retained

endophenotypes, we noticed that some distinguished between abnormalities occurring in the
initial MacTel region (area 5 in the ETDRS grid) and abnormalities expanding to the
surrounding areas of the retina, which is usually denoted to indicate progression of the disease
(areas 2, 3 and 4 on the ETDRS grid). For this reason, we defined these endophenotypes as
"MacTel area" and "Progression" respectively.

929
930
931
932

## Supplementary Table 8:

| **The Emmes Corporation:** *Coordinating Center* 401 N. Washington Street, Suite 700, Rockville, Maryland 20850 **Tel: 301.251.1161** **MacTel@emmes.com** | | |
|---|---|---|
| Sarah Duwel, RN MA CCRA *Principal Investigator / Project Director* Tel: x10874 | Traci Clemons, PhD *Principal Investigator / Statistician* Tel: x10212 | Clay Styer *Data Manager* *Advantage eClinical* Tel: x 17345 |
| Alexa Irazabal *Administrative Coordinator – Genetic Shipping Supplies* Tel: x10860 | Lena Bradley *Project Manager – Lead CRA* Tel: x10865 | |

| **Lowy Medical Research Institute** *Executive and Fiscal Sponsor* L31, 85 Castlereagh Street Sydney, New South Wales, AU | | |
|---|---|---|
| **Primary Contact:** Jennifer Trombley RN MSN CCRC *Director, Clinical Operations* 3366 N. Torrey Pines Ct., Suite 300 La Jolla, CA 92037 858.249.7109 office (US) 760.666.0452 mobile (US) JTrombley@LMRI.net | Martin Friedlander MD PhD *President – LMRI* 858.249.7100 office (US) 760.666.0452 mobile (US) MFriedlander@LMRI.net | Anton Lever *Group Manager, Finance* AUS Tel: +612 9295 6015 AUS Mobile: +614 0307 2143 alever@lfg.com.au |

| moorfields Reading Centre | Columbia University, Genetics Lab |
|---|---|
| *Moorfields Eye Hospital NHS*<br>*162 City Road*<br>*London EC1V 2PD*<br>MacTel@moorfields.nhs.uk | *Eye Institute Research Annex, Rm. 202*<br>*635 West 165th Street*<br>*New York, NY 10032*<br>*Tel: 212-305-8989* |
| Main contact:<br><br>**Konstantinos Balaskas**<br>**Consultant Ophthalmic Surgeon**<br>Konstantinos.Balaskas@moorfields.nhs.uk<br><br>**Administration:**<br><br>**Norma Jack**<br>**Tel: 020 7566 2815**<br>Norma.Jack@moorfields.nhs.uk | Main Contact:<br>**Carolyn Cai, Research Assistant**<br><br>Tel: 212-305-8501<br>Fax 212-305-7014<br>Cc3529@cumc.columbia.edu |

| Sites | | |
|---|---|---|
| Site Name | Principal investigator | Coordinator |
| Centre for Eye Research (002)<br>Australia Macula Research Unit<br>Level 1, 32 Gisborne St.<br>East Melbourne VIC 3002 | Robyn Guymer, PhD<br>Tel: 61 3 9929 8393<br>Fax: 61 3 9662 3859<br>Email:<br>rhg@unimelb.edu.au | Melinda Cain<br>Tel: 61 3 9929 8396<br>Email:<br>mcain@unimelb.edu.au<br><br>**EMILY CARUSO**<br>Tel: 61 3 9929  8113<br>Email:<br>emily.caruso@unimelb.edu.au |

| | | |
|---|---|---|
| Hôpital Lariboisière (004)<br>  Service d'Ophtalmologie<br>2, rue<br>Ambroise<br>Paré,<br>Cedex 10<br>Paris<br>75475 | Alain Gaudric, MD<br>  Tel: 33 1 4995 6480<br>  Fax: 33 1 4995 6484<br>Email: alain.gaudric@aphp.fr | Abir Zureik<br>  Tel: 33 1 49 95<br>64 86<br>Fax: 33 1 49<br>95 64 83<br>Email: abir.zureik@aphp.fr |
| Jules Stein Eye Institute,<br>UCLA (005)<br>  100 Stein Plaza, Room 2-565<br>Los Angeles CA,<br>90095 | Jean-Pierre Hubschman, MD<br>  Tel: 310-206-5004<br>  Fax: 310-825-3350<br>Email: hubschman@jsei.ucla.e | Rosaleen Ostrick<br>Tel: 310-794-5595<br>Fax: 310-794-5591<br>Email: ostrick@jsei.ucla.edu<br><br>Veronika Sadykov<br>Tel: 310-794-5596<br>Fax: 310-794-5591<br>Email:<br>Sadykov@jsei.ucla.edu<br>Mary Crowley<br>Tel: 310-267-5278<br>Email:<br>MECrowley@mednet.ucla.e<br>du |
| Lions Eye Institute (006)<br>  2 Verdun Street<br>  Nedlands WA 6009 | Ian Constable, MD<br>  Tel: 61 8 9381 0882<br>  Fax: 61 8 9381 0883<br>Email:<br>ian.constable@lei.org.au | Gareth Cooper<br>Email: GarethCooper@lei.org.au<br><br>Diana Bowman<br>Tel: +61 8 9381 0777<br>Fax: +61 8 9381 0700<br>Email: DianaBowman@lei.org.au |

| | | |
|---|---|---|
| Vitreous, Retina, MACULA Consultants of New York  (007)<br>460 Park Avenue<br>5<sup>th</sup> Floor<br>New York, NY 10022 | Lawrence A. Yannuzzi, MD<br>Tel: 212-605-3777<br>Fax: 212-607-3795<br>Email:<br>layannuzzi@gmail.com | MAria Scolara<br>Email:<br>mscolaro@vrmny.com<br><br>Samira Shirwa<br>Tel: 212-452-6929<br>Email: sshirwa@vrmny.com<br>Sariah R. Persaud<br>Tel: 212-452-6929<br>Email:<br>sariahpersaud@vrmny.com |
| Moorfields Eye Hospital (008)<br>Clinical Trials Unit<br>162 City Road<br>London EC1V 2PD | Cathy Egan, MD<br>Tel: 44 20 7566 2262<br>Fax: 44 20 7608 6925<br>Email:<br>catherine.egan@moorfields.nhs.uk | Supeetha david<br>Tel:  0207 566 2835<br>Email:<br>Supeetha.David@moorfields.nhs.uk<br><br>Tjebo Heeren<br>Email:<br>Tjebo.Heeren@moorfields.nhs.uk |
| Retina Associates of Cleveland, Inc. (010)<br>3401 Enterprise Parkway, Suite 300<br>Beachwood, OH 44122 | Lawrence Singerman, MD<br>Tel: 216-831-5700<br>Fax: 216-360-2196<br>Email: lsingerman@retina-assoc.com | Diane Weiss, RN<br>Tel: 216-831-5704  Ext. 244<br>Fax: 216-360-2196<br>Email: dew@retina-assoc.com |

| | | |
|---|---|---|
| Save Sight Institute (011) | Mark Gillies, MD, PhD<br>  GPO Box 4337 (use 2001 zip)<br>  8 MacQuarie Street<br>  Sydney NSW 2000<br>  Tel: 61 4 1206 0313<br>  Fax: 61 2 9382 7318<br>  Email: mark.gillies@sydney.edu.au | Stella Xu<br>  Tel +61 2 9382 7309<br>  Fax +61 2 93827278<br>  Email: stella.xu@sydney.edu.au<br>Sharon McKenzie<br>  Tel +61 2 9382 7309<br>  Fax +61 2 9382  7278<br>  Email: sharon.mckenzie@sydney.edu.au<br><br>Maria Williams<br>  Tel +61 2 9382 7286<br>  Fax +61 2 9382  7278<br>  Email : maria.williams@sydney.edu.au<br><br>Liza Shilpakar<br>  Liza.shilpakar@sydney.edu.au |
| Scripps HEalth (012)<br>  Scripps Clinical Research<br>  10666 N. Torrey Pines Rd<br>  La Jolla, CA 92037 | Martin Friedlander, MD, PhD<br>  Tel: 858-784-9138<br>  Fax: 858-784-9135<br>  Email: friedlan@scripps.edu | Jennifer Trombley, RN<br>  Tel: 858-554-6678<br>  Fax: 858-554-5016<br>  Mobile: 760-492-6600<br>  Email: Trombley.jennifer@scrippshealth.org |
| St. Franziskus Hospital (013)<br>  St. Franziskus Hospital Hohenzollernring 74<br>  Muenster D-48145 | Daniel Pauleikhoff, Prof. Dr<br>  Tel: 49 251 933080<br>  Email: dapauleikhoff@muenster.de | elke attenberger<br>  Email: elke.attenberger@uveitis-zentrum.de<br>  Tel: 0251-9352758<br>Martin Ziegler<br>  Email: mpz@alex-ziegler.de |

| | | |
|---|---|---|
| The Goldschleger Eye Institute (014)<br>The Chaim Sheba Medical Center<br>Tel Hashomer 52621 | Joseph Moisseiev, MD<br>Tel: 972 3 534 3462<br>Email: Joseph.moisseiev@sheba.health.gov.il | Rinat Rashty<br>Tel: 972-3-530 2877<br>Email: Rinat.rashty@sheba.health.gov.il |
| The New York Eye and Ear Infirmary (015)<br>Department of Ophthalmology<br>310 East 14th Street<br>Suite 319 South Building<br>New York NY, 10003 | Richard Rosen, MD<br>Tel: 212-979-4284<br>Fax: 212-979-4512<br>Email: rrosen@nyee.edu | **MELIZA UNSON**<br>Tel: 212-979-4672<br>Email: munson@nyee.edu |
| University of BONN (017)<br>Dept. of Ophthalmology<br>Ernst-Abbe-Str. 2<br>Bonn, 53127 | Frank Holz, MD<br>Tel: 49 228 287 5647<br>Fax: 49 228 287 5603<br>Email: Frank.Holz@ukbonn.de | Simone Müller, Assistenzärztin<br>Email: Simone.Mueller@ukbonn.de<br><br>**Kerstin Bartsch**<br><br>Email: Kerstin.Bartsch@ukbonn.de |
| University of Michigan, Kellogg Eye Center (019)<br>1000 Wall Street<br>Ann Arbor MI, 48105 | Grant Comer, MD<br>Room 714<br>Tel: 734-763-5906<br>Fax: 734-936-2340<br>Email: gcomer@umich.edu | Pam Campbell, COT, CCRP<br>Room 1415<br>Tel: 734-936-0138<br>Email: pamtitus@med.umich.edu |
| Scheie Eye Institute (022)<br>51 North 39th Street<br>Philadelphia PA, 19104 | Alexander Brucker, MD<br>Tel: 215-662-8675<br>Fax: 215-662-8025<br>Email: ajbrucke@mail.med.upenn.edu | Sheri Drossner<br>Tel: 215-662-8177<br>Fax: 215-662-8025<br>Email: Sheri.Grand@uphs.upenn.edu |
| University of Bern (024)<br>Universitätsklinik für Augenheilkunde<br>Inselspital<br>Bern<br>, CH<br>3010 | Sebastian Wolf, MD, PhD<br>Tel: 41 (0)31 632 8503<br>Fax: 41 (0)31 632 4382<br>Email: Sebastian.wolf@insel.ch | Corinne Stockli<br>Tel: 41 (0)31 632 1197<br>Fax: 41 (0)31 632 0114<br>Email: Corinne.Stoeckli@insel.ch |

| | | |
|---|---|---|
| University of Utah Health Care (025)<br><br>Moran Eye Center<br>University of Utah Health Care<br>65 Mario Capecchi Drive<br>Salt Lake City, UT 84132 | Paul Bernstein, MD, PhD<br>Tel: 801-581-6078<br>Fax: 801-581-3357<br>Mobile: 801-560-3093<br>Email: paul.bernstein@hsc.utah.edu | Barbara Hart<br>Email: Barbara.hart@hsc.utah.edu<br><br>Kara Halsey<br>Tel: 801-213-2034<br>Email: Kara.halsey@utah.edu<br><br>Kimberly Wegner<br>Tel: 801-584-6265<br>Email: Kimberley.wegner@hsc.utah.edu<br><br>Sam Taylor<br>Email: samclarktaylor@gmail.com<br><br>**ELIZABETH NUTTALL – Regulatory Documents**<br><br>Email: Elizabeth.nuttall@hsc.utah.edu |
| Bascom Palmer (Miami) (026) | Philip Rosenfeld, MD, PhD<br>900 NW 17th Street<br>Miami FL, 33136<br>Tel: 305-326-6148<br>Fax: 305-326-6538<br>Email: prosenfeld@med.miami.edu | Cristina Lage-Rodriguez, MS, CCRC<br>1638 NW 10th Avenue, Suite 505B<br>Miami, FL 33136<br>Tel: 305-326-6117<br>Fax: 305-547-3662<br>Email: clage@med.miami.edu |
| Massachusetts Eye and Ear (027)<br><br>Retina Service<br><br>243 Charles Street<br>Boston, MA 02114 | Joan Miller, MD<br>Tel: 617-573-3915<br>Fax: 617-573-3364<br>Email: joan_miller@meei.harvard.edu | Patricia Houlihan<br>Tel: 617-573-3646<br>Fax: 617-573-3514<br>Email: patricia_houlihan@meei.harvard.edu<br><br>Alex Leite<br>Email: Alex_Leite@meei.harvard.edu |
| Emory University (028)<br><br>Emory Eye Center<br><br>1365B Clifton RD NE<br>Atlanta, GA 30322 | Jiong Yan, MD<br>Tel: 617-573-3915<br>Email: jyan2@emory.edu | Jayne Brown<br>Tel: 404-778-4430<br>Email: jmbrown@emory.edu |

| | | |
|---|---|---|
| UNiversity of California – San francisco (029)<br>UCSF Department of Ophthalmology<br>10 Koret Way Box 0730, Room K209<br>San Francisco, CA 94143 | JaCque Duncan, MD<br>Tel: 415-514-4241<br>Email: jacque.duncan@ucsf.edu | Arshia MiaN<br>Email: Arshia.Mian@ucsf.edu<br><br>MARY LEW<br>Email: Mary.Lew@ucsf.edu |
| University of Rochester (030)<br>Flaum Eye Institute<br>601 Elmwood Avenue<br>Box 659<br>Rochester, NY 14642 | Mina Chung, MD<br>Tel: 585-275-4141<br>Email: mina_chung@urmc.rochester.edu | Katherine Nedrow<br>Tel: 585-275-9485<br>Email:katherine_nedrow@urmc.rochester.edu |
| Medical College of Wisconsin (031)<br>The Eye Institute<br>925 N. 87th Street, Room 807<br>Milwaukee, WI 53226 | David Weinberg, MD<br>Tel: 414-955-2026<br>Email: dweinber@mcw.edu | Ellie Dorsey<br>Tel: 414-955-7854<br>Email: edorsey@mcw.edu<br><br>Shay Bourgeois<br>Tel: 414-955-2001<br>Email: sbourgeois@mcw.edu<br><br>**Brittany Rego**<br>Tel: 14-955-2031<br>Email: brego@mcw.edu |
| Radboud (033)<br>Radboud University Medical Center<br>Departments of Ophthalmology & Human Genetics<br>P.O. Box 9101, 6500 HB Nijmegen (409) | CB Hoyng, MD<br>Email: trialcentrum.ohk@radboudumc.nl<br>Tel: +31 (0)24 36 15255 | Asha Kalisingh<br>Email: asha.kalisingh@radboudumc.nl<br>Tel: + 0243613212<br><br>chantal jumelet vanast<br>Email: Chantal.Jumelet-vanAst@radboudumc.nl<br><br>Clasien Oomen – regulatory<br>Email: clasien.oomenvanast@radboudumc.nl<br>Tel: + 31-24-36 15255 |
| Oxford eye hospital (034)<br>John Radcliffe Hospital West Wing | **Peter Charbel Issa MD, DPhil, FEBO**<br><br>T: 01865 227678 | Alexina Fantano<br>T: 01865 231053<br>F: 01865 231160<br>Email: alexina.fantato@ouh.nhs.uk |

| | | |
|---|---|---|
| Headley Way, Headington, Oxford, OX3 9DU | Email: peter.charbelissa@ndcn.ox.ac.uk | |
| Eye Center, Medical Center, Faculty of Medicine, University of Freiburg (035) German: Klinik für Augenheilkunde, Universitätsklinikum Freiburg<br><br>UNIVERSITÄTSKLINIKUM FREIBURG Klinik für Augenheilkunde Studienteam der Sektion Retinologie Dagmar Rommel Studienassistentin / Studienkoordination Killianstraße 5 · 79106 Freiburg /Germany Telefon: 0761 270-40231 Telefax: 0761 270-86210 | Felicitas Bucher, MD Email:   bucherf@scripps.edu | Dagmar Rommel Email: dagmar.rommel@uniklinik-freiburg.de |
| Retina Research CEnter (036)<br>  3705 Medical Parkway<br>  Suite 420<br>  Austin, Texas 78705 | Chirag Jhaveri, Md Email: cjhaveri@e-retina.net Tel: 512-279-1251 | ivana Gunderson Email: igunderson@e-retina.net Tel: 512-279-1251<br><br>Daniela Rodridguez Email: drodriguez@e-retina.net |

| | | |
|---|---|---|
| Retina Consultants of Southern Colorado (037)<br>2770 N. Union Blvd. Suite 140<br>Colorado Springs, CO 80909 | Ryan Rich, MD<br>Email: researchmdrich@coloradoretina.com<br>Tel: 719-473-9595 | Jordan Townsend<br>719-473-9595 Ext. 123<br>Email: jtownsend@coloradoretina.com<br><br>Dragana Kuga<br>Email: dkuga@coloradoretina.com<br>Victoria Lee<br>Email vlee@coloradoretina.com<br><br>Kari Winger<br>Email kwinger@coloradoretina.com |
| Cincinnati Eye Institute – CEI Physicians PSC, Inc (038)<br>1945 CEI Drive<br>Cincinnati, OH 45242 | Daniel Miller, MD<br>Email: dmiller@cincinnatieye.com<br>Tel: 513-569-3669 | Megan Kingdon, RN, BSM, CCRC<br>Email: mkingdon@cincinnatieye.com<br>Tel: 513-569-3669<br><br>Kelsi Dwyer<br>Email: kdwyer@cincinnatieye.com |
| University of Washington Eye Institute (039)<br><br>908 Jefferson St., 7th Floor, Seattle, WA 98104<br>Mailing Address: Box 359607, 325 Ninth Avenue, Seattle, WA 98104 | Cecilia Lee, MD<br>Email: jung.cecilia@gmail.com | Ian Luttrell<br>Email: luttri@uw.edu<br><br>Francy Moses<br>Email: mosesf@uw.edumary |
| Byers Eye Institute - Stanford University (040)<br>2370 Watson Court<br>Palto Alto, California 94303 | Diana Do, MD<br>Email: dianado@stanford.edu<br>Tel: 650-723-6995 | Lisa Greer<br>Email: lgreer7@stanford.edu<br>Tel: 650-725-9184 |

| | | |
|---|---|---|
| MAyo Clinic (041)<br>200 First St SW<br>Rochester, MN 55905 | Sophie Bakri, MD<br>Email:<br>bakri.sophie@mayo.edu | Heidi Rubin<br>Email:<br>Rubin.heidi@mayo.edu |
| Retina Center of New Jersey, LLC (042)<br>1255 Broad Street, Suite 104<br>Bloomfield, NJ 07003 | Patrick Higgins, MD<br>  Email:<br>Drhiggins@Retinacenternj.Com<br>  Tel: 973-707-5632 | Diane Deininger<br>  Email:<br>ddeininger@retinacenternj.com<br>  Tel: 973-707-5632<br>  Fax :973-337-5865<br>Susan Anderson<br>  Email:<br>sanderson@retinacenternj.com<br>  Tel: 973-707-5632<br><br>Valerie Benavides<br>  Email:<br>vbenavides@retinacenternj.com<br>  Tel: 973-707-5632<br>KAthleen Rodriguez<br>  Email:krodriguez@retinacenternj.com<br>  Tel: 973-707-5632 |
| Retina Associates New Orleans (043)<br>4315 Houma Boulevard Suite 201<br>Metairie, LA 70006 | Stanislav A. Zhuk, MD, FACS<br>  Email:stanislav.zhuk@retinaassociates.org<br>  Tel: 504-456-9061<br>  Fax: 504-895-8094 | MElinda Mitchell<br>  Email:<br>Melinda.mitchell@retinaassociates.org |

| | | |
|---|---|---|
| Royal Oak (044)<br><br>Associated Retinal Consultants, P.C. Neuroscience Center Building<br>3555 West Thirteen Mile Road Ste. LL-20<br>Royal Oak, MI 48073 | Sandeep Randhawa | Kendra Mellert<br>E-mail: kmellert@arcpc.net<br>Tel: 248-288-9132 ext. 1313<br>Fax: 248-288-2265 |
| Grand Rapids (144)<br><br>Associated Retinal Consultants, P.C.<br>1179 East Paris, S.E.; suite: 250<br>Grand Rapids, MI 49546 | Paul V. Raphaelian, M.D. | Deb Markus<br>Email: dmarkus@arcpc.net |
| Traverse City (244)<br><br>Associated Retinal Consultants, P.C.<br>860 E. Front St.<br>Traverse City, Mi 49686 | Scott Sneed | Serena Neal<br>Email: sneal@arcpc.net |
| Sierra Eye Associates (045)<br>950 Ryland Street, Reno, NV, 89502 | Arshad Khanani, MD<br>Email: arshad.khanani@gmail.com<br>Tel: 775-329-0286 | Deanna Dorman<br>Email:ddorman@sierraeyeassociates.com<br>Tel: 775-329-0286 x 141<br><br>**Heather Southwick**<br>Email: hpopovich@sierraeyeassociates.com |

| | | |
|---|---|---|
| Retina Northwest – PORTLAND (046) Portland 5440 SW Westgate Drive Suite 217 Portland, OR 97221 | Michael Lee, MD Email: MLee@retinanorthwest.com <br><br> Tel: 503-274-2121 | Brian Puckett Email: bpuckett@retinanorthwest.com <br> Tel: 503-274-2121 Fax: 503-229-8472 |
| Eramus - Netherlands (047) Department of Ophthalmology Erasmus Medical Center Rotterdam BA-130 postvak 'Trials' 3000 CA Rotterdam The Netherlands | J. Vingerling, MD Email: j.vingerling@erasmusmc.nl | MARjo-Lan jansen Email: m.jansen-vandrecht@erasmusmc.nl |
| New England Retina Consultants, PC (048) 3640 Main Street, Suite 201 Springfield, MA. 01107 | David LaLLY, MD Email: david.lally@gmail.com | Leah Persky Email: persky.leah@gmail.com Tel: 413-788-6044 Fax: 413-746-9715 |
| CUMBERLAND VALLEY RETINA CONSULTANTS (049) 1150 Opal Court Hagerstown, MD 21740 | DAVID WARROW, MD Email: Davidw@retinacare.net | Paula SMith Email: paulas@retinacare.net Tel: 301-665-1712 Fax: 301-655-1714 <br> Katie Alvord Email: katiea@retinacare.net Tel: 301-665-1712 x 1024 |

| | | |
|---|---|---|
| Northwestern University (050)<br>259 E. Erie, Suite 15-20,<br>Chicago, IL 60611 | Amani Fawzi, MD<br>Email: amani.fawzi@northwestern.edu | Regulatory:<br>Michell Kim<br>Email: michell.kim@northwestern.edu<br>Tel:312-503-2229<br><br>Nicole Seddon<br>Email: nreinhol@nm.org<br>Tel: 312.695-0252<br><br>Priya Thakkar<br>Email: Priya.Thakkar@nm.org<br>Tel: 312.695.2573 |
| ZNA Middelheim Hospital (051)<br>Lindendreef 1<br>2020 Anterwerp, Belguim | Joke Ruys, MD<br>Email: joke.ruys@zna.be<br>Tel : +0032 3 280 3614 | LiEsBeth VanPeene<br>Email: liesbeth.vanpeene@zna.be<br>Tel. +0032 3 280 3614 |
| Bay Area Retina Associates (052)<br>122 La Casa Via Suite 223<br>Walnut Creek, CA 94598 | Roger Goldberg, MD<br>Email:rgoldberg@bayarearetina.com | Farhan AhmeD<br>Email:fahmed@bayarearetina.com |
| Southeastern Retina Associates, P.C. (053)<br>1124 E. Weisgarber Rd. | Suite 207<br>Knoxville, TN 37909 | Joseph M. Googe Jr, MD<br>Email: ctgooge@seretina.com<br>Tel: 865-588-0811 | Kristina Oliver<br>Email: koliver@seretina.com |
| Mayo Clinic, Scottsdale Arizona (054)<br>13400 East Shea Boulevard<br>Scottsdale, AZ 85259 | Odette Houghton, MD<br>Email: Houghton.Odette@mayo.edu | Michaele Menghini<br>Email: Menghini.Michaele@mayo.edu |

| | | |
|---|---|---|
| Elman Retina Group, P.A. (055)<br>9114 Philadelphia Rd.<br>Suite 310<br>Baltimore, MD 21237 | Michael J. Elman, M.D.<br>Email:<br>elman@elmanretina.com<br>Phone: 410- 686-3000<br>Fax: 410- 686-3690<br>Cell: 410-812-6440 | JoAnn Starr<br>Email:<br>Starr@elmanretina.com<br>Phone: **410-686-3000**<br>FAX: **410-686-3690** |
| Retina Consultants of Houston (056)<br>6560 Fannin Street<br>Suite 750<br>Houston, TX 77030 | Charles Wykoff, MD, PhD<br>Email :<br>ccwmd@houstonretina.com<br>Phone: 713-524-3434 | Kirsten Poldervart<br>Email:<br>kirsten.poldervart@houstonretina.com<br>Phone: 713-524-3434<br><br>Jose Munoz<br>Email:<br>jose.munoz@houstonretina.com<br>Phone: 713-524-3434<br>Cassandra 'Cassie' Cone<br>Email:<br>Cassandra.cone@houstonretina.com<br>Phone: 713-524-3434<br><br>MArie Chin (Regulatory)<br>Email:<br>marie.chin@houstonretina.com<br>Phone: 713-524-3434 |
| Charles Retina (057)<br>1432 Kimbrough Road<br>Germantown, TN 38138 | Steve Charles, MD<br>Email: scharles@att.net | Melanie Woods<br>Email:<br>mwoods@charlesretina.com |
| Tulsa Retina Consulants (059)<br>2424 E 21st Street,<br>Suite 200<br>Tulsa, OK 74114<br>918-794-6089 | Lars Freisberg, MD<br>Email<br>Larsfreisberg@mac.com | Scott Hewitt<br>Email:<br>shewitt@retinatulsa.com<br>Phone: 918 -734-6344<br><br>Debbie Hardison<br>Email:<br>dhardison@retinatulsa.com |

| | | |
|---|---|---|
| Palmetto Retina Center, LLC (061)<br>124 Sunset Court<br>West Columbia, SC 29169 | JOHN A. Wells, III, MD<br>Email: jackwells@palmettoretina.com | Peggy Miller<br>Email: pmcdougal@palmettoretina.com<br>Phone: 803-404-5038<br>Fax: 803-744-0145<br><br>Mallie Taylor<br>Email: mtaylor@palmettoretina.com |
| Texas Retina Associates (062)<br>9600 N. Central Expressway, Suite 100<br>Dallas, TX 75231 | Gary Fish, MD<br>Email: gefgef@ix.netcom.com | Karen Duignan<br>Email : kduignan@texasretina.com<br>Phone: 214-692-6885 x1<br>Fax: 214-265-0935<br><br>Tiffanie Keaton<br>Email: tkeaton@texasretina.com |
| Ophthalmic Consultants of Long Island (063)<br>360 Merrick Rd<br>3rd Floor<br>Lynbrook, NY 11563 | Glenn Stoller, MD<br>Email : glstoller@gmail.com<br>Phone: 516-593-4026 | Brennon Taylor<br>Email: btaylor@ocli.net<br>Phone: 516-593-4026<br><br>Susan Jones<br>Email: sjones@ocli.net<br>Phone: 516-593-4026<br><br>Carla Del Castillo<br>Email : cdelcastillo@ocli.net<br>Phone: 516-593-4026 |

| | | |
|---|---|---|
| Center for Retina and Macular Disease - Central Florida (065)<br><br>**250 Avenue K, SW**<br><br>**Suite 200**<br><br>**Winter Haven, FL 33880** | Richard Hamilton<br>Email: rich@crmd.net | Research Supervisor<br>Toni Collins<br>Email: tocol@crmd.net<br>Phone: 863-297-5400 Ext. 2046<br><br>Tammy<br>Email: tamc@crmd.net |
| **Retina-Vitreous Associates Medical Group (066)**<br><br>**9001 Wilshire Boulevard, Suite 301**<br><br>**Beverly Hills, CA 90211** | David Boyer M.D.<br>Email: vitdoc@aol.com | Janet Kurokouchi<br>Email: JKurokouchi@laretina.com<br>Phone: (310) 289-2478 x 3 (main study line) |
| **Global Research Foundation  (067)**<br><br>**7447 N Figueroa Street**<br><br>**Los Angeles, CA 90041** | **Khaled Tawansy**<br><br>Email: ktawansy@globalresearchgroup.net | Research Supervisor<br>Monica Torres<br>Email: mtorres@globalresearchgroup.net<br>Phone:  323-257-3300 ext 120 |

# References

1.    Mathew R, Sivaprasad S, Florea D, et al. Agreement between time-domain and
spectral-domain optical coherence tomography in the assessment of macular thickness
in patients with idiopathic macular telangiectasia type 2. *Ophthalmologica*.
2013;230(3):144-150. doi:10.1159/000353455

2.    Buuren S van, Groothuis-Oudshoorn K. mice: Multivariate Imputation by Chained
Equations in R. 2011. http://doc.utwente.nl/78938/. Accessed March 10, 2017.

3.    Gottesman II, Gould TD. The endophenotype concept in psychiatry: etymology and
strategic intentions. *Am J Psychiatry*. 2003;160(4):636-645.
doi:10.1176/appi.ajp.160.4.636

4.    Revelle W. psych: Procedures for personality and psychological research. *Northwestern*
*University, Evanston R package version*. 2014;1(1).

5.    Jaisankar D, Raman R, Gondhale H, Chhablani J, Agarwal M. VISUAL FUNCTION
CORRELATES OF FOVEAL SLOPE CHANGES ON OPTICAL COHERENCE
TOMOGRAPHY IN MACULAR TELANGIECTASIA TYPE 2. *Retina*. December 2016.
doi:10.1097/IAE.0000000000001416

6.    Benjamini Y, Krieger AM, Yekutieli D. Adaptive linear step-up procedures that control
the false discovery rate. *Biometrika*. 2006;93(3):491-507. doi:10.1093/biomet/93.3.491

7.    metabolomics gwas server. http://metabolomics.helmholtz-muenchen.de/gwas/.
Accessed December 4, 2018.

8.    Shin S-Y, Fauman EB, Petersen A-K, et al. An atlas of genetic influences on human
blood metabolites. *Nat Genet*. 2014;46(6):543-550. doi:10.1038/ng.2982

979