Joint Inference of Clonal Structure using Single-cell Genome and Transcriptome Sequencing Data

Xiangqi Bai,[1] Zhana Duren,[2] Lin Wan,[3,4,*] Li C. Xia[5,*]

[1] Division of Oncology, Department of Medicine,
Stanford School of Medicine, Stanford, CA, 94305, USA
[2] Center for Human Genetics and Department of Genetics and Biochemistry,
Clemson University, Greenwood, 29646, SC, USA
[3] NCMIS, LSC, Academy of Mathematics and Systems Science,
Chinese Academy of Sciences, Beijing, 100190, China
[4] University of Chinese Academy of Sciences, Beijing, 100049, China
[5] Department of Statistics and Finance, School of Mathematics,
South China University of Technology, Guangzhou, 510006, China
[*] Corresponding author. lwan@amss.ac.cn (Lin Wan), lcxia@scut.edu.cn (Li C. Xia)

Latest advancements in high-throughput single-cell genome (scDNA) and transcriptome (scRNA) sequencing technologies enabled cell-resolved investigation of tissue clones. However, it remains challenging to cluster and couple single cells for heterogeneous scRNA and scDNA data generated from the same specimen. In this study, we present a computational framework called CCNMF, which employs a novel Coupled-Clone Non-negative Matrix Factorization technique to jointly infer clonal structure for matched scDNA and scRNA data. CCNMF couples multi-omics single cells by linking copy number and gene expression profiles through their general concordance. We validated CCNMF using both simulated benchmarks and real-world applications, demon-

strating its robustness and accuracy. We analyzed scRNA and scDNA data from an ovarian cancer cell lines mixture, a gastric cancer cell line, as well as a primary gastric cancer, successfully resolving underlying clonal structures and identifying high correlations of coexisting clones between genome and transcriptome. Overall, CCNMF is a coherent computational framework that simultaneously resolves genomic and transcriptomic clonal architecture, facilitating understanding of how cellular gene expression changes along with clonal genome alternations.

Introduction

Understanding of how genomic content changes impact gene expression in individual cells is essential to further understand the cell clone development in normal and diseased tissues. In particular, characterizing the clone-wise gene dosage effect, i.e., the sensitivity of cellular gene expression to the copy number profile shared by the group of cells, is critical to elucidate the functional consequence of diseases-associated genomic copy number variants (CNVs), a significant challenge in current structural variant research (*1–3*).

However, there is no technology available that can efficiently and accurately measure both DNA copy number and gene expression profiles of individual cells simultaneously. Although several technologies (*4–6*), like scTrio-seq had made an attempt to measure genomic and transcriptomic content of up to a few cells per batch, it remains a low-throughput technique. High-throughput single-cell sequencing technologies that are currently available can only measure either the transcriptome (*7–10*) or the genome (*11–13*) content of individual cells, but not both simultaneously.

For examples, droplet-based single-cell RNA sequencing (scRNA) technology is routinely employed to measure cellular expression so as to assess the clonal development states of various

tissues and cell systems (*12, 14*). Recently, droplet-based single-cell DNA sequencing (scDNA)

technology enabled cell-wise and genome-wide measurement of genomic alternations, such as

copy number variants, in thousands of cells (*12, 15*). High quality single-cell copy number vari-

ants combined with single-cell gene expression profiles promise to further reveal the clonal het-

erogeneity in complex tissues and cell systems (*16–20*). Realizing the potential, however, will

require high fidelity co-clustering of heterogeneous single cells measured scRNA and scDNA

sequencing technologies.

Addressing this difficulty, we developed an efficient computational method – **C**oupled-

Clone **N**on-negative **M**atrix **F**actorization, termed as CCNMF. It reasonably models the shared

underlying clonal structure and the general concordance between cellular expression level and

copy number states (*12, 21*). CCNMF then employs machine learning algorithms to infer the

most likely multi-modal integration solution. Briefly, CCNMF took two matrices as inputs:

single-cell gene expression matrix obtained through scRNA-seq technology and single-cell

copy number variants matrix obtained from scDNA-seq technology, both derived from the same

biological specimen.

CCNMF was established on a model-based approach which couples single cells across

scDNA and scRNA data by maximizing their global concordance between gene expression and

copy number variants. CCNMF optimizes an objective function that simultaneously maximizes

intra-clone compactness and inter-clone structure coherence, this coupled-clone nonnegative

matrix factorization framework followed the co-clustering concept as introduced in ref. (*22*). A

coherent underlying clonal structure, i.e. the identity-linked cell clusters between scDNA and

scRNA data, is thus inferred as the weight matrix optimally assigning all cells to their most

likely cluster identity. Based on that, CCNMF accurately estimates the dosage effect per gene

and cell cluster.

Before CCNMF, only a few methods were available for analyzing the combined scDNA

and scRNA data. These methods mostly operate in a map-to-reference mechanism, i.e., data from one technology is mapped to the reference clonal structure derived from another technology (*21, 23–27*). For examples, *clonealign*, an early attempt of integrative modeling gene dosage effect of DNA copy number and gene expression, statistically assigns scRNA gene expression states to a reference phylogenetic tree representing scDNA-derived clones, in a Bayesian way (*21*); *Seurat*, which mainly integrates multiple scRNA datasets, can also project other types of single cell data to the scRNA-derived clusters using the mutual nearest neighbor search (*23*); *DENDRO* infers single cell copy numbers from scRNA data and validates the result using the scDNA data (*24*). The model-free single-cell integrative nonnegative matrix factorization (iNMF) method (*25*) identified the shared cell types across multi samples and species from scRNA-seq data. However, these map-to-reference inference methods risk systematic bias because the choice of reference technology was largely arbitrary, and different choices significantly influence downstream analysis.

Instead, CCNMF takes a data-driven approach, unbiased toward data and technology sources. CCNMF utilizes the coherence of the underlying clonal structure shared within the biological specimen to maximize the inference for true cell clonal identity and gene expression effects. Using both simulated and real cancer datasets, we validated that CCNMF can faithfully recover the underlying clonal structure, accurately identify clonal identity for all single cells, and statistically infer gene-wise dosage and expression changes that differentiate each clone. We applied CCNMF to characterize an ovarian cancer cell mixture, a gastric cancer cell line and a primary gastric cancer, and the results showed CCNMF is capable of identifying clonal structure and dosage effect in real cell systems. Thus, scDNA- and scRNA-seq combined with CCNMF analysis offers a new way to study the functional consequence of clonal gene dosage change and how it contributes to clonal development.

Results

The CCNMF Toolkit

The CCNMF analytical framework has been implemented as an *R* package, with a workflow illustrated in Fig. 1. CCNMF toolkit can accept scRNA and scDNA data in standard formats, including those generated by 10X Genomics scRNA-/scDNA-seq and DLP scDNA technologies. The toolkit executes the statistical framework and analytical steps as follows:

1. Aligns scRNA genes and scDNA genome bins to the same genome reference.

2. One-to-multiple mapping of scDNA bins to genes using location overlapping (> 1bp) to reduce the scDNA data to the same gene coordinates as the scRNA data.

3. Initializes the coupled term between scRNA (E) and scDNA data (O) using a coupling matrix (A), which is either an identity matrix or a user provided matrix with prior information.

4. Iteratively optimizes the objective function using a gradient descent algorithm until convergence.

5. Identifies the most coherent clonal structure by finding the maximum weights of the H matrices that represented the most likely cell clonality membership.

The outputs of CCNMF include: W matrices that represent the expression or copy number profile centroids of scRNA or scDNA clones; H matrices that represent the cell-wise membership weights toward clone for each of scRNA and scDNA cells. The toolkit is platform-independent and works with general R installation. It is made available as an open source software on Github with detailed readme, manual and examples.

CCNMF recovers the underlying clonal structures in Simulation

We firstly evaluated the performance of CCNMF using simulated scDNA and scRNA datasets – *Sim*. The evaluation was based on Adjusted Rand Index (ARI), and the results were presented in **Supplementary tables S1-S3**. *Sim* included two different scenarios, each with three parameters. In each scenario, ARI was assessed by varying the parameter of interest over its range while keeping the others as default. The default parameters for copy number fraction, outlier percentage and dropout percentage were 50%, 0 and 0, respectively.

Supplementary table S1 showed the simulation results for the linear and bifurcate clonal structure scenarios, with various copy number fractions ranging from 10% to 50%. It was defined as the percentage of genome undergoing copy number changes. As shown in the table, CCNMF was able to recover the exact underlying clonal structure with the highest accuracy (ARI=1) for all cases under both scenarios, except for one case with an ARI of 98%. With the decreasing of the copy number fraction, the clonal copy number difference becomes smaller, making it harder for CCNMF to resolve the clones accurately. The results demonstrated that such effect is only modest, as with only 10% of genome having copy number difference between the clones, CCNMF was still able to correctly uncover the underlying structure.

Supplementary table S2 showed the simulation results with varying dropout percentages from 10% to 90%. Dropout percentage was defined as the percentage of cells with zero values for gene expression and copy number. The reason behind it could be the limited sensitivity of a technology or the gene was not present or expressed. Dropouts are very common in scRNA and scDNA data because of amplification bias and other random effects. In *Sim*, a dropout percentage at 10% means that 10% of all simulated gene expression or copy numbers were perturbed to be zeros. As shown in the table, CCNMF achieved high accuracy in recovering the underlying clonal structure for all cases under both scenarios. All resulted ARIs were $> 98\%$, except for one case with ARI of 81%.

Supplementary table S3 showed the simulation results for outlier percentage ranging from

10% to 90%. It was defined as the percent of cells with non-realistic copy numbers or expression

values. In practice, these data points are typically deemed technical errors and are excluded

from downstream analysis. In *Sim*, outlier percentage stands for the percent of all simulated

scDNA and scRNA cells were perturbed to be outliers. It was obvious from the table that

CCNMF was robust to the presence of outliers. When the outlier percentage was less than 60%,

CCNMF achieved high accuracy with all ARIs greater than 92% for both scenarios. In summary,

our comprehensive simulation study demonstrated that CCNMF achieved good performance for

resolving the coherent underlying clonal structures in scDNA and scRNA data with practical

noise considerations.

CCNMF detected cell origins for ovarian cancer mixture cell lines *OV* data

We conducted CCNMF analysis on the *OV* benchmark dataset consisting of a mixture of ovarian

cancer cell lines. It was composed of two cell lines, OV-2295(R) and TOV-2295(R) from the

same patient. OV-2295(R) was an ascites site cell line that is abnormal adjacent tissue but

not cancerous, while TOV-2295(R) was a high-grade serous ovarian cancer cell line (*28*). The

ground truth for individual cells were obtained from the original publication (*28*).

CCNMF successfully characterized two subclones in the *OV* mixture cells lines. The scDNA

cells were classified into subclones C1 and C2, which comprised of 394 cells from TOV-

2295(R) and 371 cells from OV-2295(R), respectively. Correspondingly, the scRNA cells were

categorized into subclones R1 and R2, consisting of 4918 cells from TOV-2295(R) and 1717

cells from OV-2295(R), respectively. Notably, C1 and R1 were from the same cell line TOV-

2295(R), while C2 and R2 were from cell line OV-2295(R), indicating corresponding relation-

ship with scDNA subclones with scRNA subclones. The tSNE plots of Fig. 2A and 2B demon-

strated the clear separation of the two mixture cells lines in scDNA and scRNA, respectively.

We compared the identified clones with the ground truth, resulting in an adjusted Rand index

(ARI) of 1, indicating that CCNMF accurately recovered the underlying clonal structure of *OV*.

159　CCNMF identified coherent clonal structure in gastric cancer cell line $NCI - N87$
160　data

To investigate the performance of CCNMF on a cancer cell line, we analyzed the $NCI - N87$

gastric caner cell line to determine whether the clone structure can be detected by CCNMF.

The large-scale scDNA data of $NCI - N87$ was composed of 1105 single cells. A substantial

proportion of replicating cells existed in the scDNA which will cause analytical difficulty for

clone reconstruction (*12*). It is because the most copy number changes in replicating cells were

driven by ongoing DNA replication, which overwhelm the true clonal copy number variants.

To address this challenge, we filtered out replicating cells before joint clustering analysis. We

identified the replicating cells by calculating the variance of copy number for each cell because

the higher replication activity gives rise to a higher variance of observed intra-cell segments.

We utilized the Expectation-Maximization algorithm to fit a two-component mixture normal

distribution on the obtained copy number variance across all cells (**Supplementary Fig. S4**).

The cells were identified as replicating cells if they were assigned to a normal distribution with

a larger mean, and were excluded from further analysis. Using this efficient filtering procedure,

we successfully identified and removed the group of replicating cells with highly fluctuating

copy numbers and retained 724 cells (**Supplementary Fig. S5**).

We utilized CCNMF to analyze single-cell copy number and gene expression matrices with

selected and shared genes/features. We successfully identified and characterized three sub-

clones (C1, C2 and C3) in the $NCI - N87$ cell line for matched scDNA and scRNA, as shown

in the tSNE plots (see Fig. 3A and 3B). Out of the initial 1005 cells of scDNA, 281 cells

were identified as replicating cells and were discarded from downstream clonal reconstruction.

The remaining 724 cells were classified into three subclones, with 456, 91 and 177 cells in

C1, C2 and C3, respectively. Notably, all the cells in C1 exhibited a consistent large CNVs

amplification on the chromosome 19, while all cells in C3 shared a focal amplification event

on the chromosome 3q arm. However, C3 was involved in a smaller proportion cells without

amplifications on chromosomes 3 and 19, and instead, shared lesser amplification events on the

chromosome 11 (see Fig. 3C). As an independent validation, our analysis successfully resolved

the two major subclones and one minor subclones characterized in another study (*12*), with C1

and C3 corresponding to the two major subclones and C2 corresponding the minor one.

To increase comparability between scRNA and scDNA, we also filtered out the replicating

cells in scRNA by calculating cell cycle scores using the "CellCycleScoring" function in Seu-

rat. Ultimately, we selected 2168 scRNA cells in G0/G1 phase out of a total of 3246 cells and

coupled them with 724 scDNA cells in G0/G1 phase as well. We identified three subclones (R1,

R2 and R3) from 2168 scRNA cells, with 1337, 128, and 703 cells in each, respectively. To es-

timate the clonal large-scale CNVs, we applied the inferCNV package on gene expression with

clone identities. We then visualized the inferred CNVs changes among the clones of scRNA

by heatmap (Fig. 3D). Notably, the three scRNA subclones (R1, R2 and R3) corresponded to

the scDNA clones (C1, C2 and C3). The inferred CNVs in R1 were consistent with C1, which

shared a amplification event on chromosome 20. The focal amplification event on chromosome

3q arm was shared by R3 and C3. The cells of R2 shared lesser amplification events on chro-

mosome 11, similar to C2. It is worth noting that the amplification-like events on chromosome

1 and 4 observed in R3 represented the relatively higher gene expressions in those cells than

others in R1 and R2, which were potentially corresponding cells in C2 with diploids. We calcu-

lated Pearson's correlation between pair-wise cells from scDNA and scRNA. It depicted a clear

block pattern where cells in matched clones between two modalities had high correlations than

cells in different clones, due to the clone-wise gene dosage effect associated with CNVs (Fig.

3E).

CCNMF characterized cohort clone structure for primary gastric cancer *P5931*

Besides analyzing tumor cell lines, we performed CCNMF on real-world application for pri-

mary gastric adenocarcinoma. The scRNA and scDNA datasets were derived from a primary

gastric cancer patient *P5931* (*29*). Primary tumors usually had higher degree of intracellular

heterogeneity and more complex tumor microenvironment than cancer cell lines. Notably, we

identified the tumor cells in G0/G1 phase for *P5931* scDNA after filtering out the replicating

and normal cells (**Supplementary Fig. S6**). Moreover, we also detected the epithelial cells in

G0/G1 phase in scRNA (*29*). Then we applied the CCNMF to find the underlying clone struc-

ture between tumor G0/G1 cells in scDNA and epithelial G0/G1 cells in scRNA for *P5931*.

Due to the heterogeneity in primary cancer specimen, there were no ground truth available

of cell clone identities. Therefore, we interpreted the clonal structure based on the result of

CCNMF. As shown in Fig. 4A and 4B, two clones overlaid with cell identities on tSNE plots

for scDNA and scRNA. The two clones were clearly separated in scDNA through several cells

in C1 were closer to C2 group. The clonal structure of scDNA of this primary gastric cancer

sample was dominated by these significant copy number alternations affecting chromosomes

7 and 21 (Fig. 4C). The somatic events shared by cells of C1 were chromosome 8 and 21

amplifications. Clearly, the chromosome 7 amplification and chromosome 21 deletion were the

major defining somatic events that separated the C2 from C1.

We visualized the underlying large-scale CNVs pattern of scRNA clones with cell identi-

ties from CCNMF (Fig. 4D). Here we applied the inferCNV package to infer the large-scale

CNVs on the clones of scRNA. The inferred CNVs were estimated based on the relative gene

expression levels among different clusters. In Fig. 4D, the two clones R1 and R2 of scRNA

were corresponding with C1 and C2 of scDNA, respectively. Here, the inferred CNVs in R1 were consistent with C1, which shared chromosome 8 and 20 amplifications. Also chromosome 7 amplification and chromosome 21 deletion were shared by R2 and C2. The Pearson's correlation was calculated between pair-wise cells between scDNA and scRNA. The correlation heatmap demonstrated that the cells in matched clones between two modalities had high correlations due to the clone-wise gene dosage effect associated with CNVs (Fig. 4E).

Discussion

Single-cell multi-omics technology enables the identification of cellular and genomic characteristics of cancer cells. The advancements in scDNA-seq technology allow for characterizing tumor subclone architectures by providing genomic DNA variation such as CNVs. Importantly, each subclone has distinct genomic alternations and cellular properties. Understanding the biological features and phenotypes of subclones is essential for precision cancer treatment. However, scDNA-seq data cannot be used to identify cellular types or phenotypes. Single-cell RNA-seq provides gene expression informations that elucidates the biology of individual cells, but it is limited in defining specific cancer subclones. Inferring CNVs from scRNA-seq remains as a potential area for obtaining biological insights of subclones (*20, 30, 31*). However, the inferred CNVs are based on changes in read depth or gene expression across the genome, and only provide limited reliability of copy number information of individual cells. Therefore, the matched scDNA-seq data is still needed as a ground truth of genome changes.

It is important to note that currently, there is a lack of technologies that are capable of simultaneously measuring copy number variants and gene expression of the same cell with high throughput. Although G&T-seq (*4*), DR-seq (*5*) and scTrio-seq (*6*) can measure genomic and transcriptome in up to a few cells per batch, there are low throughput manner. High throughput scRNA-seq can only measure transcriptome, and scDNA-seq is only for genome content, but not

for both modalities in a single cell. To overcome these limitations, the integration of matched

scDNA-seq and scRNA-seq of the same specimen is a promising approach.

To facilitate the understanding of tumor clonal structure and the associated biological char-

acteristics like clonal-wise gene dosage effect, we proposed a joint-clustering approach CC-

NMF for the integration of matched scRNA- and scDNA-seq data from the same specimen.

CCNMF optimizes an objective function that simultaneously maximizes for intra-technology

clonal compactness, inter-technology clonal coherence and expected dosage effect consistence.

We demonstrated the utility of CCNMF by achieving high accuracy for resolving coherent

clonal structure in simulation datasets with various noise levels. In real-world applications, CC-

NMF has been validated by identifying the underlying clonal structure in mixtures of ovarian

cancer cell lines, a gastric caner cell line and a primary gastric cancer patient. The consis-

tent CNV change patterns between corresponding subclones between scDNA and scRNA were

observed in results of CCNMF.

CCNMF uncover the coherent clonal architecture from the matched tumor single-cell genome

and transcriptome data. The integration analysis showcased heterogeneous and clonal genetic

nature of pathological tissues, which provides crucial dosage effect information for elucidating

genetic cause and etiology of diseases. Furthermore, CCNMF can serve as a bioinformatics tool

for performing single-cell level clonal dosage effect analysis for the community.

Materials and Methods

Coupled factorization of scDNA and scRNA data

We utilized the coupled-clone nonnegative matrix factorization framework to identify the un-

derlying clonal structure of scDNA and scRNA data from the same biological specimen. The

input can be any matched scDNA and scRNA metrics generated by various technologies. Input

matrix $O \in R^{p \times n_1}$ is the copy numbers of p genes and n_1 cells from the scDNA; while the input

matrix $E \in R^{p \times n_2}$ is the gene expression of p genes and n_2 cells from the scRNA (Fig. 1).

CCNMF was established on a powerful approach–Nonnegative Matrix Factorization (NMF),

which uncovers the latent low-dimensional representation for a given matrix (*32, 33*). Briefly,

NMF factorizes the given features by samples matrix into two non-negative matrices W and

H, where W represents the latent structure of features (i.e. genes and CNVs), while H de-

scribes the weight of those features among samples (i.e. cells). Besides the NMF, the most

important concept in CCNMF is to coupling the nonnegative factorization of matrices O and

E. We additionally defined $A \in R^{p \times p}$ to represent the linked sensitivity of gene expression

to copy number. The matrix A serves as a bridge to enforce the link between changes in copy

number and gene expression level for corresponding genes. A can be estimated priorly either

by a linear regression model using public bulk RNA and DNA sequencing data (*22*) or by an

identity matrix. The diagonal elements of the identity matrix represent the full informative links

between copy number and gene expression on the same genes, but not for other genes. Hence,

we simultaneously co-factorize the single-cell datasets O and E by minimizing the following

objective function:

$$
\begin{aligned}
\mathcal{F}(W, H) = \min_{W_1, H_1, W_2, H_2 \geq 0} & \frac{1}{2} \left\| O - W_1 H_1 \right\|_F^2 \\
& + \frac{\lambda_1}{2} \left\| E - W_2 H_2 \right\|_F^2 - \lambda_2 tr(W_2^T A W_1) \\
& + \mu(\left\| W_1 \right\|_F^2 + \left\| W_2 \right\|_F^2)
\end{aligned}
\tag{1}
$$

$$
\text{subject to} : \left\| W_1 \right\|_F^2 = 1, \left\| W_2 \right\|_F^2 = 1, W_1, W_2, H_1, H_2 \geq 0.
$$

where we denoted $W_1 \in R^{p \times k}, W_2 \in R^{p \times k}$ and $H_1 \in R^{k \times n_1}, H_2 \in R^{k \times n_2}$ by shorthands W

and H, $tr()$ represents the trace of a matrix.

By minimizing the first two terms of the objective function in Equation (1), we ensured

the respective NMF decompositions of O and E as $O = W_1 H_1$ and $E = W_2 H_2$. Notably,

W_i $(i = 1, 2)$ represents the mean matrix of clusters for the n_i cells, while H_i is the weight

matrices that softly assign n_i single cells to the underlying identity linked cell clusters. Upon

convergence, the weight matrix H_i provides the inferred cluster identities for all single cells

by the maximum weight. Additionally, minimizing the cross term $-tr(W_2^T A W_1)$ ensured the

coherence of the inferred clone structure between the scRNA and scDNA data.

Optimization of the objective function

303 For optimization, we applied the alternating direction methods of multipliers (ADMM) (*34,*

304 *35*) to the objective function. Let μ_1 and μ_2 be the Lagrangian multipliers for W_1 and W_2,

305 respectively, thus the transformed objective function was written as follows:

$$L(W, H, \mu_1, \mu_2) = \mathcal{F}(W, H) + \sum_{n=1}^{2} \mu_n tr(W_n^T W_n) \tag{2}$$

306   To solve the transformed objective function, we first obtained required gradients by setting

307 its first order derivatives to zeros. Then, we used the gradient descent algorithm to iteratively

308 update and optimize the objective function until convergence by following those steps (for proof

309 see **Supplementary Methods**):

$$
\begin{aligned}
h_{ij}^1 &\leftarrow h_{ij}^1 \frac{(W_1^T O)_{ij}}{(W_1^T W_1 H_1)_{ij}}, \\
w_{ij}^1 &\leftarrow w_{ij}^1 \frac{(O H_1^T + \lambda_2 A^T W_2 + W_1 m_{11})_{ij}}{(W_1 H_1 H_1^T + W_1 m_{12})_{ij}}, \\
h_{ij}^2 &\leftarrow h_{ij}^2 \frac{(W_2^T E)_{ij}}{(W_2^T W_2 H_2)_{ij}}, \\
w_{ij}^2 &\leftarrow w_{ij}^2 \frac{(E H_2^T + \frac{\lambda_2}{\lambda_1} A W_1 + W_2 m_{21})_{ij}}{(W_2 H_2 H_2^T + W_2 m_{22})_{ij}},
\end{aligned}
\tag{3}
$$

310   where

$$m_{11} \leftarrow tr(W_1^T(W_1 H_1 H_1^T)),$$

$$m_{12} \leftarrow tr(W_1^T(O H_1^T + \lambda_2 A^T W_2)),$$

$$m_{21} \leftarrow tr(W_2^T(W_2 H_2 H_2^T)),$$

$$m_{22} \leftarrow tr(W_2^T(E H_2^T + \frac{\lambda_2}{\lambda_1} A W_1)).$$

(4)

Model inputs

The model had two hyper-parameters inputs as λ_1 and λ_2, which were used to initialize the iter-

ative computation. Our experiences were that λ_1 and λ_2 are data-dependent. Nonetheless, they

can be empirically determined by the input data. In practice, we used an automatic balancing

strategy to determine the parameters, which ensured the initial values of the four terms of the

objective function are within the same order.

The coupling matrix A is also expected as input, for which an identify matrix can be sup-

plied. The non-zero diagonal elements represent the strengths of linked copy number and gene

expression on the same gene, while the zero non-diagonal elements mean uninformative rela-

tionships between copy number and genes on different genes. To incorporate a real informative

prior for A, we can estimate it from known associations between copy number and gene expres-

sion using bulk sequencing data of the same tissue source. It was well known that DNA copy

number is highly positively correlated with the gene expression levels for the most ($> 99\%$)

of expressed human genes (36). We thus estimated a diagonal coupling matrix ($A \in R^{p \times p}$)

where each diagonal element was calculated by the ratio of gene-wise mean expression to mean

copy number using the bulk RNA and DNA data downloaded from The Cancer Genome Atlas

(TCGA) (https://www.cancer.gov/tcga). Moreover, this obtained empirical matrix

A also was used in simulation to generate realistic-like matched scRNA and scDNA datasets.

Model Selection

When the number of cluster k is unknown, CCNMF runs specified range of models with different k. The optimal number of k was selected based on the minimum coverage value of the optimization function across various k sets.

Preprocessing matched scRNA & scDNA data

We preprocessed the scRNA data with several steps, including filtering outlier genes and cells, normalizing for sequencing depth and log transformation. We also performed chromosome-level smoothing procedure for each cell, which the expression of order genes along each chromosome were smoothed the average of 101 gene window.

It is noted that scDNA features were located on the genome segmental bins, while scRNA features were presented across genes. To ensure the consistency of between scDNA and scRNA features, we associated multiple neighbor genome bins from scDNA with its corresponding gene from scRNA using the following preprocessing steps. First, we aligned both scRNA and scDNA into the same human genome assembly (GRCh37 or GRCh38). Then, we identified the one-to-multiple mapping between each gene and bins using the IRanges R package (*37*). Finally, the bin-level copy number variants of scDNA matrix was converted into the mapped gene by taking the average of CNVs on corresponding bins.

To perform feature selection on scDNA-seq copy number matrix, we developed a novel statistical approach due to the limited availability of methods in this task. After converting segmental bin-level scDNA data into gene-level CNVs, we selected the most highly variable genes across tumor cells while excluding replicating cells. Specifically, we modeled the variances of CNVs for all genes using a mixture normal distribution, and iteratively selected genes located at the normal distribution with a higher mean until at least 20% genes were chosen. The resulting genes were considered potentially informative and used as input features for CCNMF. Finally,

353 we extracted the properly formed both scRNA and scDNA matrices $E \in R^{p \times n_2}$ and $O \in R^{p \times n_1}$

354 for CCNMF analysis (see Fig. 1), where p genes were the remaining selected features from

355 scDNA.

Simulation procedures

We generated the matched scRNA and scDNA data from the same clonal structure by presetting

the ground truth genetic copy number (**GCN**) changes (as illustrated in **Supplementary Fig.**

S1). Notably, the ground truth GCN profile represented a specific clonal structure. First, we

specifically set the first clone as normal cells with GCN vector $V_1 = [2, \cdots, 2] \in R^m$, where

m enumerates over all genome segmental bins. The second tumor clone's associated GCN

vector as $V_2 \in R^m$, in which a fraction of V_2 were randomly sampled from $\{0, 1, 3, 4\}$ with

various probability parameters. Similarly, V_3 of the third clone were also simulated by the

above procedure. Finally, we obtained a ground truth GCN profile $V = [V_1, V_2, V_3, \cdots]$ as the

baseline genomic change profile for the underlying clonal structure.

To simulate the scDNA data, we estimated their specific parameters and noise from bulk

CNV datasets downloaded from cBioPortal (*38, 39*). Specifically, (i) we estimated the probabil-

ity transition matrix $P(C|G = k)$ for observed integer copy number when given genetic changes

using bulk CNV data, where C is the observed CNV, G is the genetic CNV, k $(0, 1, 2, 3, \cdots)$

is the value of copy number (**Supplementary Fig. S2**); (ii) We simulated copy number for per

gene and per cell $D_{ij} \sim multinomial(P(C|G) * P(V))$ (**Supplementary Fig. S3**) when given

the clonal GCN matrix V; (iii) We then added outlier and dropout events (**Supplementary**

Methods).

We extended the associated genetic copy number (GCN) profile V to genes$\times$cells matrix in

which GCN determines the clone-wise genes expression and copies. We next used the Splatter

pipeline (*40*) to simulate library effects, dropout, outlier events in scRNA-seq data based on

the above extended clonal gene-wise matrix (**Supplementary Methods**). Splatter parameters

were estimated from the same tumor tissue with RNA-seq and bulk CNV data downloaded from

cBioPortal.

Simulated and real datasets

As the first benchmark, we simulated 46 scDNA- and scRNA-seq datasets. They were referred

as the **Sim** data. The linear and bifurcated clone structure scenarios with 3 clones were sim-

ulated in this study. We varied common experimental parameters, such as the percentages of

outliers, dropouts, and CN dosage effect. For each simulated dataset, we randomly generated

cell-wise scDNA and scRNA data according to the specified scenario and parameters using

the procedure as detailed previously (also see **Supplementary Fig. S1**). Each of the obtained

dataset in *Sim*, has 1000 cells and 2000 genes/CNV bins, and the three composing clones have

200, 400 and 400 cells each. The first clone was designed as normal cells, and the second and

third clones represented by deletion and amplification events, respectively. We set the percent-

ages of differentially acquired deletions and amplifications to affect 10% to 50% chromosome

regions. We deposited the *Sim* data into GitHub.

For the real data benchmark, we obtained a mixture of high grade serous carcinoma (HGSC)

cell lines for matched scDNA and scRNA, referred to as the **OV** data. It was sequenced by

DLP scDNA-seq and the 10X Genomics scRNA-seq technologies and was downloaded from

European Genome-Phenome archive with accession EGAD00001004553 (*21*). The mixture

cells were made up from ascites (OV2295R) and solid tumors (TOV2295R). The scRNA subset

included 1717 cells from ascites (OV2295R) and 4918 cells from solid tumors (TOV2295R),

while the scDNA subset had 371 cells from OV2295R and 394 cells from TOV2295R.

As another real data application, we also downloaded the matched scRNA and scDNA data

for $NCI - N87$ gastric cancer cell line from Gene Expression Omnibus (GSE142750) and

National Institute of Health's SRA (PRJNA498809) (*12*). We firstly processed the scDNA-seq data using Cellranger-DNA pipeline with reference genome GRCh38 according to described procedures in (*12*). The scDNA data includes 1005 single cells, each contains 154423 copy number bins across the whole genome. The scRNA-seq data has 3246 cells with 13513 genes per cell.

Finally, we also collected a primary gastric adenocarcinoma patient data, called *P5931* (*29*), which was sequenced using 10X Genomics technologies for both scRNA and scDNA. The scDNA consists of 796 cells with 154423 copy number bins across the whole genome. The scRNA of *P*5931 has in total 11217 cells with 19129 genes per cell. The data was downloaded from dbGAP repositories with accession numbers phs001711 (*29*).

References

1. L. C. Xia, J. M. Bell, C. Wood-Bouwens, J. J. Chen, N. R. Zhang, H. P. Ji, Identification of large rearrangements in cancer genomes with barcode linked reads. *Nucleic acids research* **46**, e19 (2018).

2. L. C. Xia, S. Sakshuwong, E. Hopmans, J. Bell, S. Grimes, D. Siegmund, H. Ji, N. Zhang, A genome-wide approach for detecting novel insertion-deletion variants of mid-range size. *Nucleic Acids Research* **44**, e126 (2016).

3. L. C. Xia, D. Ai, H. Lee, N. Andor, C. Li, N. R. Zhang, H. P. Ji, Svengine: an efficient and versatile simulator of genome structural variations with features of cancer clonal evolution. *GigaScience* **7**, giy081 (2018).

4. I. C. Macaulay, W. Haerty, P. Kumar, Y. I. Li, T. X. Hu, M. J. Teng, M. Goolam, N. Saurat, P. Coupland, L. M. Shirley, M. Smith, N. Van der Aa, R. Banerjee, P. D. Ellis, M. A.

Quail, H. P. Swerdlow, M. Zernicka-Goetz, F. J. Livesey, C. P. Ponting, T. Voet, G&T-seq: parallel sequencing of single-cell genomes and transcriptomes. *Nature Methods* **12**, 519-522 (2015).

5. S. S. Dey, L. Kester, B. Spanjaard, M. Bienko, A. van Oudenaarden, Integrated genome and transcriptome sequencing of the same cell. *Nature Biotechnology* **33**, 285-289 (2015).

6. Y. Hou, H. Guo, C. Cao, X. Li, B. Hu, P. Zhu, X. Wu, L. Wen, F. Tang, Y. Huang, J. Peng, Single-cell triple omics sequencing reveals genetic, epigenetic, and transcriptomic heterogeneity in hepatocellular carcinomas. *Cell Research* **26**, 304–319 (2016).

7. F. Tang, C. Barbacioru, Y. Wang, E. Nordman, C. Lee, N. Xu, X. Wang, J. Bodeau, B. B. Tuch, A. Siddiqui, mrna-seq whole-transcriptome analysis of a single cell. *Nature Methods* **6**, 377-382 (2009).

8. G. X. Y. Zheng, J. M. Terry, P. Belgrader, P. Ryvkin, Z. W. Bent, R. Wilson, S. B. Ziraldo, T. D. Wheeler, G. P. McDermott, J. Zhu, Massively parallel digital transcriptional profiling of single cells. *Nature Communications* **8**, 14049 (2017).

9. E. Z. Macosko, A. Basu, R. Satija, J. Nemesh, K. Shekhar, M. Goldman, I. Tirosh, A. R. Bialas, N. Kamitaki, E. M. Martersteck, J. J. Trombetta, D. A. Weitz, J. R. Sanes, A. K. Shalek, A. Regev, S. A. McCarroll, Highly parallel genome-wide expression profiling of individual cells using nanoliter droplets. *Cell* **161**, 1202-1214 (2015).

10. J. N. Campbell, E. Z. Macosko, H. Fenselau, T. H. Pers, A. Lyubetskaya, D. Tenen, M. Goldman, A. M. J. Verstegen, J. M. Resch, S. A. McCarroll, E. D. Rosen, B. B. Lowell, L. T. Tsai, A molecular census of arcuate hypothalamus and median eminence cell types. *Nature Neuroscience* **20**, 484-496 (2017).

11. H. Zahn, A. Steif, E. Laks, P. Eirew, M. VanInsberghe, S. P. Shah, S. Aparicio, C. L.

Hansen, Scalable whole-genome single-cell library preparation without preamplification.

Nature Methods **14**, 167–173 (2017).

12. N. Andor, B. T. Lau, C. Catalanotti, A. Sathe, M. Kubit, J. Chen, C. Blaj, A. Cherry,

C. D. Bangs, S. M. Grimes, C. J. Suarez, H. P. Ji, Joint single cell DNA-seq and RNA-

seq of gastric cancer cell lines reveals rules of in vitro evolution. *NAR Genomics and*

Bioinformatics **2** (2020). Lqaa016.

13. M. Efremova, S. A. Teichmann, Computational methods for single-cell omics across

modalities. *Nature Methods* **17**, 14-17 (2020).

14. D. J. McCarthy, *et al.*, Cardelino: computational integration of somatic clonal substructure

and single-cell transcriptomes. *Nature Methods* **17**, 414-421 (2020).

15. E. I. Velazquez-Villarreal, S. Maheshwari, J. Sorenson, I. T. Fiddes, V. Kumar, Y. Yin,

M. G. Webb, C. Catalanotti, M. Grigorova, P. A. Edwards, J. D. Carpten, D. W. Craig,

Single-cell sequencing of genomic dna resolves sub-clonal heterogeneity in a melanoma

cell line. *Communications Biology* **3**, 318 (2020).

16. Y. Wang, J. Waters, M. L. Leung, A. Unruh, W. Roh, X. Shi, K. Chen, P. Scheet, S. Vattathil,

H. Liang, A. Multani, H. Zhang, R. Zhao, F. Michor, F. Meric-Bernstam, N. E. Navin,

Clonal evolution in breast cancer revealed by single nucleus genome sequencing. *Nature*

512, 155-160 (2014).

17. M. Markowska, T. Cakała, B. Miasojedow, B. Aybey, D. Juraeva, J. Mazur, E. Ross,

E. Staub, E. Szczurek, Conet: copy number event tree model of evolutionary tumor his-

tory for single-cell data. *Genome Biology* **23**, 128 (2022).

18. S. Zaccaria, B. J. Raphael, Characterizing allele- and haplotype-specific copy numbers in
single cells with chisel. *Nature Biotechnology* **39**, 207-214 (2021).

19. C.-Y. Wu, B. T. Lau, H. S. Kim, A. Sathe, S. M. Grimes, H. P. Ji, N. R. Zhang, Integrative
single-cell analysis of allele-specific copy number alterations and chromatin accessibility
in cancer. *Nature Biotechnology* **39**, 1259-1269 (2021).

20. C.-Y. Wu, A. Sathe, J. Rong, P. R. Hess, B. T. Lau, S. M. Grimes, H. P. Ji, N. R. Zhang,
Cancer subclone detection based on dna copy number in single cell and spatial omic se-
quencing data. *bioRxiv* (2022).

21. K. R. Campbell, A. Steif, E. Laks, H. Zahn, D. Lai, A. McPherson, H. Farahani, F. Kabeer,
C. O'Flanagan, J. Biele, J. Brimhall, B. Wang, P. Walters, I. Consortium, A. Bouchard-
Côté, S. Aparicio, S. P. Shah, clonealign: statistical integration of independent single-cell
rna and dna sequencing data from human cancers. *Genome Biology* **20**, 54 (2019).

22. Z. Duren, X. Chen, R. Jiang, Y. Wang, W. H. Wong, Modeling gene regulation from paired
expression and chromatin accessibility data. *Proceedings of the National Academy of Sci-
ences* **114**, E4914-E4923 (2017).

23. T. Stuart, A. Butler, P. Hoffman, C. Hafemeister, E. Papalexi, W. M. Mauck, Y. Hao,
M. Stoeckius, P. Smibert, R. Satija, Comprehensive integration of single-cell data. *Cell*
177, 1888-1902.e21 (2019).

24. Z. Zhou, B. Xu, A. Minn, N. R. Zhang, Dendro: genetic heterogeneity profiling and sub-
clone detection by single-cell rna sequencing. *Genome Biology* **21**, 10 (2020).

25. J. D. Welch, V. Kozareva, A. Ferreira, C. Vanderburg, C. Martin, E. Z. Macosko, Single-
cell multi-omic integration compares and contrasts features of brain cell identity. *Cell* **177**,
1873-1887.e17 (2019).

26. S. Milite, R. Bergamin, L. Patruno, N. Calonaci, G. Caravagna, A Bayesian method to

cluster single-cell RNA sequencing data using copy number alterations. *Bioinformatics* **38**,

2512-2518 (2022).

27. M. Edrisi, X. Huang, H. A. Ogilvie, L. Nakhleh, Macrodna: Accurate integration of single-

cell dna and rna data for a deeper understanding of tumor heterogeneity. *bioRxiv* (2022).

28. I. J. Letourneau, M. C. Quinn, L. L. Wang, L. Portelance, K. Y. Caceres, L. Cyr, N. Delvoye,

L. Meunier, M. de Ladurantaye, Z. Shen, S. L. Arcand, P. N. Tonin, D. M. Provencher, A.-

M. Mes-Masson, Derivation and characterization of matched cell lines from primary and

recurrent serous ovarian cancer. *BMC Cancer* **12**, 379 (2012).

29. X. Bai, B. Lau, S. M. Grimes, A. Sathe, H. P. Ji, Single cell multi-omic mapping of sub-

clonal architecture and pathway phenotype in primary gastric and metastatic colon cancers.

bioRxiv (2022).

30. T. Tickle, I. Tirosh, C. Georgescu, M. Brown, B. Haas, infercnv of the trinity ctat project,

`https://github.com/broadinstitute/inferCNV` (2019).

31. R. Gao, S. Bai, Y. C. Henderson, Y. Lin, A. Schalck, Y. Yan, T. Kumar, M. Hu, E. Sei,

A. Davis, F. Wang, S. F. Shaitelman, J. R. Wang, K. Chen, S. Moulder, S. Y. Lai, N. E.

Navin, Delineating copy number and clonal substructure in human tumors from single-cell

transcriptomes. *Nature Biotechnology* **39**, 599-608 (2021).

32. D. D. Lee, H. S. Seung, Learning the parts of objects by non-negative matrix factorization.

Nature **401**, 788-791 (1999).

33. D. D. Lee, H. S. Seung, *Advances in Neural Information Processing Systems 13* (MIT

Press, 2001), pp. 556–562.

34. S. Boyd, N. Parikh, E. Chu, B. Peleato, J. Eckstein, Distributed optimization and statistical
learning via the alternating direction method of multipliers. *Foundations and Trends in*
Machine Learning **3**, 1-122 (2010).

35. Z. Duren, X. Chen, M. Zamanighomi, W. Zeng, A. T. Satpathy, H. Y. Chang, Y. Wang,
W. H. Wong, Integrative analysis of single-cell genomics data by coupled nonnegative
matrix factorizations. *Proceedings of the National Academy of Sciences* **115**, 7723-7728
(2018).

36. R. S. N. Fehrmann, J. M. Karjalainen, M. Krajewska, H.-J. Westra, D. Maloney, A. Sime-
onov, T. H. Pers, J. N. Hirschhorn, R. C. Jansen, E. A. Schultes, H. H. H. B. M. van Haagen,
E. G. E. de Vries, G. J. te Meerman, C. Wijmenga, M. A. T. M. van Vugt, L. Franke, Gene
expression analysis identifies global gene dosage sensitivity in cancer. *Nature Genetics* **47**,
115 (2015).

37. M. Lawrence, W. Huber, H. Pagès, P. Aboyoun, M. Carlson, R. Gentleman, M. Morgan,
V. Carey, Software for computing and annotating genomic ranges. *PLoS Computational*
Biology **9** (2013).

38. E. Cerami, J. Gao, U. Dogrusoz, B. E. Gross, S. O. Sumer, B. A. Aksoy, A. Jacobsen, C. J.
Byrne, M. L. Heuer, E. Larsson, Y. Antipin, B. Reva, A. P. Goldberg, C. Sander, N. Schultz,
The cbio cancer genomics portal: An open platform for exploring multidimensional cancer
genomics data. *Cancer Discovery* **2**, 401–404 (2012).

39. J. Gao, B. A. Aksoy, U. Dogrusoz, G. Dresdner, B. Gross, S. O. Sumer, Y. Sun, A. Jacobsen,
R. Sinha, E. Larsson, E. Cerami, C. Sander, N. Schultz, Integrative analysis of complex
cancer genomics and clinical profiles using the cbioportal. *Science Signaling* **6**, pl1–pl1
(2013).

40. L. Zappia, B. Phipson, A. Oshlack, Splatter: simulation of single-cell rna sequencing data.

Genome Biology **18**, 174 (2017).

Acknowledgments: XB gratefully acknowledges the financial support from China Scholarship

Council (No.201904910816).

Funding: LW was supported by the National Key Research and Development Program of China

under Grant 2022YFA1004801 and the National Natural Science Foundation of China (No.

12071466). LCX was supported by GuangDong Basic and Applied Basic Research Foundation

2022A1515-011426.

Competing interests: No competing interest is declared.

Author contributions: XB, LW and LCX designed the study; XB, ZD, LW and LCX were

involved in algorithm development; XB performed data analysis; LW and LCX supervised this

study; XB, LW and LCX wrote the manuscript. All authors reviewed and approved the final

manuscript.

Data and materials availability: The R package of CCNMF is available at

https://github.com/labxscut/CCNMF.

Figures

Fig. 1. The workflow of Coupled-Clone Nonnegative Matrix Factorization (CCNMF). It

took scDNA matrix $O \in R^{p \times n_1}$ and scRNA matrix $E \in R^{p \times n_2}$ as input for inferring shared

clonal structure between two modalities from the same bio-specimen. We applied a coupling

matrix $A \in R^{p \times p}$ to link matrices $O \in R^{p \times n_1}$ and $E \in R^{p \times n_2}$, which represented the global

concordance between DNA copy number and gene expression.

Fig. 2. The coherent clonal structure between scDNA and scRNA of the mixture ovarian

cancer cell lines *OV***.** (A) tSNE plot of scDNA clones. (B) tSNE plot of scRNA clones. (C)

Heatmap shows CNVs changes across scDNA's clones. (D) Heatmap shows inferred CNVs

changes estimated by gene expression level across scRNA's clones. (E) Heatmap of Pearson's

Correlation between single cells in coherent clones from scDNA and scRNA. Cells composing

the same clone were coded in the same color. Each row represents a single cell and each column

represents a genomic region for (C) and (D). The color in each dot of the heatmap represents

the CNV status for (C) and (D). In panel (E), each row represents a single cell of scDNA, while

each column represents a single cell of scRNA, the color in the heatmap represents the Pearson's

correlation of cells between scDNA and scRNA.

Fig. 3. The coherent clonal structure between scDNA and scRNA of the gastric cancer

cell line *NCI-N87*. (A) tSNE plot of scDNA clones. (B) tSNE plot of scRNA clones. (C)

Heatmap shows CNVs changes across scDNA's clones. (D) Heatmap shows inferred CNVs

changes estimated by gene expression level across scRNA's clones. (E) Heatmap of Pearson's

Correlation between single cells in coherent clones from scDNA and scRNA.

Fig. 4. The coherent clonal structure between scDNA and scRNA of the primary gastric

cancer patient *P5931*. (A) tSNE plot of scDNA clones. (B) tSNE plot of scRNA clones. (C)

Heatmap shows CNVs changes across scDNA's clones. (D) Heatmap shows inferred CNVs

changes estimated by gene expression level across scRNA's clones. (E) Heatmap of Pearson's

Correlation between single cells in coherent clones from scDNA and scRNA.

Supplementary materials

577　Supplementary Methods

578　Supplementary Figs. S1 to S6

579　Supplementary Tables S1 to S3

Fig. 1. The workflow of Coupled-Clone Nonnegative Matrix Factorization (CC-NMF).

Fig. 2. The coherent clonal structure between scDNA and scRNA of the mixture ovarian cancer cell lines *OV*.

Fig. 3. **The coherent clonal structure between scDNA and scRNA of the gastric cancer cell line** *NCI-N87.*

Fig. 4. **The coherent clonal structure between scDNA and scRNA of the primary gastric cancer patient** *P5931.*