Segregation, connectivity, and gradients of deactivation in neural correlates of evidence in social decision making

Roberto Viviani+*, Lisa Dommes*, Julia E. Bosch*, Karin Labek+

+ Institute of Psychology, University of Innsbruck, Austria

*** Department of Psychiatry and Psychotherapy III
University of Ulm, Germany**

Correspondence: roberto.viviani@uibk.ac.at

Abstract

Functional imaging studies of sensory decision making have detected a signal associated with evidence for decisions that is consistent with data from single-cell recordings in laboratory animals. However, the generality of this finding and its implications on our understanding of the organization of the fMRI signal are not clear. In the present functional imaging study, we investigated decisions in an elementary social cognition domain to seek evidence for neural correlates of evidence, their segregation, connectivity, and their relationship to task deactivations. Participants were asked to decide which of two depicted individuals was saddest on the basis of information rich in sensory features (facial expressions) or through contextual cues suggesting the mental state of others (stylized drawings of mourning individuals). The evidence signal was located in two distinct sparse networks differentially recruited depending on the two image types. Using the main evidence hubs as seeds in a database of connectivity data, these two networks were wholly and selectively retrieved. Furthermore, all hubs were located near or along a ribbon of cortex located between task activations and deactivations between areas affected by perceptual priming and the deactivated areas of the default network system. In associative cortex, these findings suggest gradients of progressive deactivation as a possible neural correlate of the cortical organization envisaged by predictive coding theories of cortical function.

Segregation, connectivity, and gradients of deactivation in neural correlates of evidence in social decision making

Introduction

Single-cell recording studies or perceptual decision making in laboratory animals have demonstrated the existence of neurons in several regions of the brain whose firing rates increase with the evidence for a decision (Gold and Shadlen 2002; Cisek and Kalaska 2010; Shadlen and Kiani 2013; Gold and Heekeren 2014). The firing rates may reflect the gradual accumulation of the evidence for choosing among the options on offer, until a decision threshold is reached (Gold and Shadlen 2002). It has also been argued that this mechanism may help to understand how the brain processes information more generally, providing a "window on cognition" (Shadlen and Kiani 2013) and revealing that brain function may differ in important respects from conventional views in cognitive science (Cisek and Kalaska 2010; Hunt and Hayden 2017).

In its purest version, the conventional view envisages informational flow as a sequence of processing steps through specialized units (Fodor 1983), whose activation may be univocally interpreted as the recruitment of an identifiable process, and that terminates with the specification of motor output (Petersen et al. 1989). In functional imaging studies, subtracting activation between carefully chosen experimental conditions allows identifying the mapping between the processing units and cortical regions (cognitive subtraction, Posner and Raichle 1994). An 'alternative view' is that the areas in which firing rates are associated with the evidence for a decision are part of a dynamically configured network of connected areas in which the information coming from lower neurons and goal representations are integrated, computing evidence for decisions through the competition of representations of alternatives (Heekeren et al. 2008; Cisek 2012). This view integrates segregation (the areas where evidence for a decision is accumulated are anatomically distinct, presumably reflecting information of different types) and connectivity (these areas must talk to each other if all aspects contributing to the decision criterion are to be considered) differently than in the conventional view. The discrepancy is best exemplified by motor control. Because accumulation of evidence also takes place in areas deputed to motor control, it is simultaneously available as a motor plan corresponding to the available options (Figure 1).

Figure 1. Simplified schemas of two views of cortical organization, highlighting the differences in connectivity and hierarchical organization that are relevant to the present study. a: in its purest form, the conventional view envisages information processing as a sequence of distinct steps with action planning following input analysis. b: a possible alternative architecture is characterized by accumulation of evidence up to a sparse interconnected network where the decision is elaborated (modified from Cisek 2012). Choice is implemented by a distributed network whose activity increases with evidence until a threshold is reached.

The centerpiece of this view is the computation of a signal reflecting evidence for a decision. Several neuroimaging studies have sought a signal with these properties. Studies of the neural correlates of perceptual decision making (for review, see #Error in retrieving data.#) have shown the existence of such as signal in a sparse network of small areas (Heekeren et al. 2004; Tosoni et al. 2008). Because when the evidence is large the decision is easy, these studies detect areas that are more active in easy than in difficult decisions (Heekeren et al. 2008). Proponents of the generality of the decision making mechanism have also suggested that it may also apply to preference-based choice (Krajbich and Rangel 2011; Cisek 2012; Glimcher 2015), as well as to choices in social cognition (Shadlen and Kiani 2013).

Whatever the merits of what we have called here the 'alternative view' may be in explaining the computational principles underlying a large set of findings, it may be noted that the strategy of contrasting easy with difficult decisions in a neuroimaging study is itself non-standard, because difficult decisions are not a control condition for easy decisions. Standard reasoning, based on the cognitive subtraction logic (Posner and Raichle 1994), would suggest that the neural correlates of decision making are those that increase with the difficulty of decisions (as in studies of cognition, Rypma et al. 1999), not those associated with their easiness (Kayser et al. 2010; for a discussion, see Heekeren et al. 2008). We readily acknowledge that the present work alone cannot resolve these conceptual discrepancies, and also that the activations associated with task difficulty (always present in these studies, including the present one) are certain to play a role in the computation of decisions. However, in the present work we intentionally left aside the investigation of the effect of task difficulty, which follows the logic of cognitive subtraction with which all neuroimagers are familiar. Instead, the purpose of the present study was to explore the extent to which the data support an alternative modelling strategy, see how far the argument in its favour can be pushed, and highlight the points where its conclusions may depart from those of the conventional approach.

Our experiment was designed to address three related issues. The first was the detection of the neural correlates of evidence for a decision in an elementary social

cognition task, thus exploring the applicability of the modelling strategy outside domains of sensory and preference-based decision making. We asked participants to indicate the saddest individual from those shown in two images, and identified a sparse network of evidence-related areas by contrasting easy to difficult decisions. Second, we looked for evidence for segregation and connectivity in this sparse network. Participants were divided into two groups, who viewed either facial expressions of varying degrees of sadness or stylized drawings of individuals without facial features depicted in scenes of mourning or in neutral scenes (Labek et al. 2017). When deciding between faces, participants can rely on purely sensory information to make their choice (Vuilleumier et al. 2003). In contrast, the scenes of mourning individuals provided minimal visual detail, and require considering the scenic context or inference about the mental state of the depicted individual to judge between degrees of sadness (Labek et al. 2017). We tested the existence of specific hubs of the decision network selectively recruited when comparing the effect of easy decisions in these two types of stimuli. Decisions based on facial expressions recruited a sparse specific cortical network in posterior associative cortex in decisions from facial expressions that was not recruited when deciding between mourning scenes, while a second network was active in both tasks. To assess connectivity, we selected the location of the main evidence hubs in these two networks and used it as seed in a database of connectivity data. The seeds recovered these two networks selectively and completely.

The third issue we wished to address concerns the location of evidence-related areas in the context of the global organization of the fMRI signal. We mapped the location of the evidence-related network with respect of task activations and deactivations, which in the posterior part of the brain form a gradient from activated perceptual cortex to the deactivations of the default network system. Areas associated with subjective value in preference-based decision making, such as vmPFC and the postsplenial cortex, are usually deactivated by the task, a fact that has spurred interest in the role of neural inhibition in the constitution of the value-associated fMRI signal (Jocham et al. 2012; Hunt et al. 2015). If preference-based and other types of choice making are related, we would expect this relation to deactivations to occur in both. Indeed, neuroimaging of negative emotions often deliver results in vmPFC and in associative cortical areas that combine deactivation and control such as to challenge conventional interpretations (Benelli et al. 2012; Viviani 2014).

We found that none of the hubs of the evidence network were located in strongly activated areas. Instead, they were typically located in proximity of task deactivations, at a point along a gradient from activation to deactivation. In the sensory-selective network recruited by deciding about faces, its hubs spanned the transition zone between task activations and deactivations, which formed a gradient of activity across the hubs. Furthermore, we observed that task deactivations in the cortical areas occupied by the hubs were modulated by the emergence of the hubs themselves. This analysis will provide suggestive evidence that these areas are embedded in a global organization of information processing whose functional imaging correlate is a gradient of decreasing activation.

Finally, we turned to the gross temporal dynamics of the signal. Using the number of previous presentations of the stimuli, we first retrieved the task signal decreases that are associated with adaptation and neural priming, thought to be a neural correlate of semantic encoding (Grill-Spector et al. 2006; Barron et al. 2016). We found that the neural priming signal, located in areas of task activation, included the cortex adjacent to the decision network hubs, without engulfing the hubs themselves. While the faces-specific decision network remained stable during the task, we found traces of an

evidence network recruited by mourning scenes that evolved during time. In the initial phase of the experiment, deciding between mourning scenes activated temporo-parietal and medial prefrontal areas that coincided with task deactivations.

Taken together, the results of the analytic strategy we followed here raise the possibility that, in so far as an fMRI signal referring to a global organization of information processing may be detected in the data of this study, this signal consists of a gradient of progressive deactivation. A consequence of this would be that lack of activation cannot be equated to lack of recruitment, contradicting a basic tenet of the cognitive subtraction logic. While a neuroimaging study cannot provide direct evidence on neural activity, these findings raise the question of whether deactivation has any plausibility at all as a neural correlate of information processing. In the discussion, we will mention some independent proposals on the organization of cortical processing that are relevant to this issue. Cortical inhibitory mechanisms arise naturally in neural models that implement competitive weighting of evidence, or in models of cortical function more generally. This suggests that the progressive task deactivation reported here, while usually a neglected aspect of neuroimaging analyses, is not a finding at odds with current reasoning about cortical function.

Methods

Task and modelling

Two groups of participants were asked to indicate the saddest individual depicted in two separate images. In the first group, participants decided on the basis of two pictures of faces displaying varying degrees of sadness (Viviani et al. 2018). In the second group, participants decided on the basis of stylized drawings of individuals in situations ranging from neutral to extreme sadness (mourning the loss of a loved one, Labek et al. 2017). Using two separate groups of participants ensured that the strategy used was not biased by the fact that the task was the same in both. The faces were displayed in frontal view from the Karolinska Directed Emotional Faces inventory (KDEF, Lundqvist et al. 1998), and were chosen so as to display varying degrees of sadness range from the neutral to the desperate. The scenes of mourning individuals and neutral scenes were taken from Labek et al. (2017).

In each trial, two pictures were displayed side by side on a screen on the back of the scanner visible through mirrors mounted on the coil. All possible combinations of the pictures from a total pool of seven, taken two at a time, were shown (all combinations of the faces pictures in the faces group and all combinations of the mourning scenes pictures in the mourning group). Button presses indicating decisions in the trials could be made only after 2.5 sec of display, signaled by the appearance on the screen of two blue circles symbolizing the buttons under the depicted items. Participants had 1.5 sec to make their choice, after which the trial was declared as a miss. Trials were presented at variable intervals (10.5 sec on average) according to a random exponential schedule bounded at 9 and 12 sec, for a total task duration of 8 min 50 sec. Participants had the opportunity to familiarize with the task in a brief session prior to scanning. The presentation of trials was programmed in standard software (Presentation 14, Neurobehavioral Systems Inc., Albany, CA).

Several strategies have been used in the literature to investigate the neural correlates of evidence. The procedure we adopted here follows the approach of sensory-based decision making in contrasting easy to difficult decisions (Heekeren et al. 2004), but a similar approach may also be found in several neuroeconomic studies of

preference-based decisions (Serences 2008; Boorman et al. 2009; FitzGerald et al. 2009; Hunt et al. 2012). From the neuroeconomic studies we borrowed the notion that the possibly subjective degree of sadness participants saw in the pictures could be inferred from their choices. After scanning, the score of the times each picture was selected as the saddest of the pair by the participant was computed. The difference and the sum of the sadness scores of the two pictures on display in the trial were computed. The difference was an approximate estimate of the evidence participants had to make their decision. The sum was an estimate of the amount of the properties that determine the decision was present on display. After centering, this participant-specific score was then used as the 'parametric modulation' of a regressor sampling the modeling of the BOLD response at each trial. This means that the model included an effect of trials relative to fixation, modeling the effects of perception and encoding of stimuli as well as the generation of responses common to all trials, and two the parametric modulations, modeling the interaction of this effect with the differences between the scores and their sum. Additional parametric modulations were given by a centered score of the number of times the pictures of the trials had been previously presented, and the interaction of the score with the difference and sum of the sadness scores. The parametric modulations captured variance not explained by the main regressor, which modeled the effect of trials. Trials in which participants gave no response (misses) were modelled separately and considered as a nuisance effect.

This modelling strategy was chosen on the basis of its capacity to control for sensory properties of the stimuli associated with the criterion for the choices, thus minimizing confounding of the criterion on the basis of which decisions are made from the effect of evidence. For example, sadness in facial expression may be confounded with salience (or any other property of the visual stimuli that correlates with sad expressions). However, the effect of evidence is not confounded with salience for two reasons. First, in so far as correlated with the degree of sadness, the salience of the images in the trial is summarized by the sum of the scores of these images. The inclusion of the sum of scores in the model therefore adjusts for salience of the trial or any other property that determines choice. Second, when participants make consistent choices, and there are no misses, then the difference and sum of scores regressors are orthogonal. This fact is difficult to prove in general, but may easily be checked directly for a given number of trials (the Matlab code to do this is provided in Viviani et al. 2017). The orthogonality follows from the fact that each picture is presented in association with all other pictures. Two very sad pictures give the same low difference score as two neutral pictures, while the same pictures, in a different combination, give high difference scores. Because the effect of the stimuli themselves is subtracted away in the regression on the score differences that detects the signal associated with evidence for the decision, this regression detects only evidence-related areas (i.e. degree of evidence for sadness), not areas differentially recruited by the sensory encoding of the stimuli.

An alternative modelling strategy consists of obtaining separate coefficients for the regression on the highest and lowest scores of the two items presented in each trial, and taking the contrast high score vs. low score to the second level. This modelling strategy differs from the one adopted here in that an additional degree of freedom is spent to model the slopes of the effects of the low and high score items independently. Using this alternative modelling strategy we obtained results that were qualitatively identical to those reported below. We used here the score difference strategy because of its better control for the confounding of the choice criterion.

Recruitment and image acquisition

The study was conducted at the Psychiatry and Psychotherapy Clinic of the University of Ulm, Germany, after approval by the Ethical Review Board. Healthy participants (N = 63) were recruited through local announcements and admitted to the study after verifying that they met the inclusion criteria and had provided written informed consent. To limit the correlation between the difference and sum of scores regressors, we set a threshold of max 4 missed trials to consider the data as valid, leading to the exclusion of 3 participants in each group. The final sample comprised 27 and 30 participants in the faces and mourning groups, respectively.

Data were collected in a Prisma 3T Siemens Scanner using a T2*-sensitive echo-planar imaging sequence (TR/TE: 2460/30 msec, flip angle 82°, image size 64×64×39 obtained from transversal slices of 2.5 mm with a gap of 0.5 mm, giving an isotropic voxel size of 3 mm). A 64-channels head coil was used with foam padding to minimize head motion.

Data analysis

Data were analyzed with the freely available software SPM12 (www.fil.ion.ucl.ac.uk/spm). After realignment, normalization, smoothing (FWHM 8 mm), and high-pass filtering (512 sec cutoff), trials were modeled by a boxcar function with fixed trial duration 2.5 sec, convolved with a standard haemodynamic function. Realignment parameters were also included in the model as nuisance covariates. The model was estimated separately in each voxel and included a first-order autoregressive term to model autocorrelation in the residuals. Estimates of the contrasts of interest were brought to the second level to account for the random effect of subjects. At the second level, we report significance tests corrected at the peak and cluster level computed by permutation (8000 re-sampling). Peak-level corrections (also known as voxel-level) achieve strong control of false rejection errors. For cluster-level corrections, clusters were defined *a priori* by the uncorrected threshold $p < 0.001$. In both corrections, the testing family was defined by a mask of gray and white matter determined by majority voting on the segmentation computed in each individual separately as part of the normalization procedure.

Overlay from Figures 2 and 7c were obtained using the SPM software. All other overlays were obtained with the freely available software MriCroN (Chris Rorden, https://people.cas.sc.edu/rorden/mricron/install.html) from the statistical parametric maps of *t* values. Figures were then annotated with text and legends in Adobe Illustrator. The hatch pattern of Figures 3 and 7a, b was obtained by drawing a grid and using it as a clipping mask on the original overlay from the statistical parametric maps. The signal course of Figures 2 and 7c was obtained by fitting a Fourier series of 5 basis functions for the 6 time-points in the interval of the plot (software package fda, MATLAB version, Ramsay and Silverman 2005) to partial residuals from the fit, with adjustment for the intercept, for the regressor for the misses (if present), and for the movement covariates. The trials were defined by an interval of 15 seconds after each onset. Ninety percent confidence intervals for the fitted curves were computed point-wise, as customary in functional data analysis (Ramsay and Silverman 2005). Histological area classifications were determined with the Anatomical toolbox for SPM (Eickhoff et al. 2005).

Results

Participants gave consistent answers, with most subjects identifying the saddest stimulus in both tasks (for a maximal classification score of 8, in the sad faces task only

two individuals obtained a maximal classification score of 6, and six a maximal classification score of 7; in the mourning scenes task, there were one and four individuals with these scores). The difference in the maximal classification scores in the two groups was not significant (logistic regression with overdispersion, t = 0.90, p = 0.37). However, there were more misses in the mourning scenes task at trend significance levels (t = 2.03, p = 0.05), possibly indicating a slightly more difficult task.

In the neuroimaging data, we first looked at the effect of the evidence for the decision (easy vs. difficult decision) in both groups taken as a whole, and at the existence of an interaction between the groups that would support segregation in this system. The evidence for the decision was associated with a signal in a sparse network of areas that included the sensory association cortex (SII) in the anterior portion of the parietal lobe (areas PFt and PFop in the histological classification of Caspers et al. 2006), the postcentral gyrus (Area 1, Geyer et al. 1999) and the adjacent precentral gyrus, and the middle insula (Figure 2a, in brown colour). This network was recruited bilaterally but with a slight left lateralization (see Table A1 in the Appendix).

Figure 2. a: the top row shows the main effect of the evidence for decision (in brown) and the interaction with the stimulus type (in blue-yellow). The interactions are highlighted by ovals and arrows (overlays thresholded at *p* < 0.01, uncorrected, for illustration purposes). The middle and bottom rows show the effect of evidence in the sad faces and mourning scenes separately (threshold at *p* < 0.05). b: The signal course at the somatosensory and insular peaks (left), and at the iPC/sTS peak (right), where it is shown separately for the faces and the mourning groups. The curves show the signal course and 90% confidence intervals, fitted by a Fourier series, in the tertiles of the trials where the evidence for the decision were highest (easy decisions, red/orange color) and lowest (difficult decisions, light blue color). Coordinates in MNI standard space. SII: sensory association cortex; MI: middle insula; iPC: inferior parietal cortex (posterior portion).

In the interaction evidence x group, there were bilateral effects in the most posterior portion of inferior parietal cortex/superior temporal gyrus (iPC/sTS), also encroaching into the occipital lobe (area PGp in Caspers et al. 2006) and in the cuneus-precuneus/retrosplenial cortex (Figure 2a, top row, in blue to yellow colours; Table A2). Inspection of the effect of evidence in the two groups taken separately (Figure 2a, middle rows, in brown) revealed this interaction to arise because these areas were recruited during the sad faces task, but not in the mourning scenes task. For this reason, we will refer to this effect as the effect of evidence in faces. Of note are also bilateral effects of evidence in the faces task in the anterior portion of the fusiform gyrus (visible at z = −16 in Figure 2a, area FG4 according to Lorenz et al. 2017). These loci failed to reach significance in the interaction, but we draw attention to them here because they will reappear later in the analysis. There were no significant effects in the other direction, i.e. areas that were associated with evidence only in the mourning scenes task, even if in Figure 2 the insular peak was stronger in the mourning scenes group. As shown in Figure 2b in these areas the signal showed a more or less marked brief deactivation at the presentation of the stimuli, followed by a rise that was steeper in the trials where the evidence for the decision was stronger.

Figure 3. a: connectivity from the left iPC/sTS at *z* = 28, overlaid on the effect of evidence for sad faces (interaction evidence x faces). The connectivity is shown as an orange hatch pattern, which largely overlaps with the effect of evidence for sad faces, which is the same effect shown in the top row of Figure 2a (blue to yellow colours). b: connectivity from the left middle insular seed, overlaid on the main effect for evidence (in brown, the same as in the top row of Figure 2a). The positive connectivity is shown as an orange hatch pattern, as in panel a. Negative connectivity is in shades of grey. All connectivity values were thresholded at 0.20, as in the default display from neurosynth.org.

In the alternative view, the areas where evidence for a decision is accumulated must be talking to each other to integrate different properties of the options and reach a consistent decision. We therefore looked for data on the connectivity of the evidence in faces network in the public database neurosynth.org (Yarkoni et al. 2011). We looked at the connectivity spanned by the left iPC/sTS, as shown in Figure 3a. In this and the following figures, we show the same data as in Figure 2a (effect of evidence in faces in blue-yellow, effect for evidence in the whole sample in brown) in combination with other effects. In Figure 3a, one can see that the connectivity of this seed, shown in the hatched areas, closely overlaps with the effect of evidence for faces. This overlap also includes the anterior portion of the fusiform gyrus that failed to reach significance in the previous analysis. The only partial mismatch between this connectivity and the effect of evidence for faces concerns the medial face of the brain, where the connectivity extends toward areas that are weakly detected by the effect for evidence for the whole group

(brain areas in Figure 1a, $x = -8$). There was no substantial negative connectivity in the data from neurosynth.org for this seed.

To look for evidence for connectivity in the region associated with evidence in the whole group, we examined the data for the left insular region. We chose this seed because of its importance in the encoding of pain (Lamm et al. 2011; Peyron and Fauchon 2019), and because it had a robust activation in the mourning group. The results, shown in Figure 3b, show again a substantial overlap with the areas associated with the amount of evidence, again with the possible exception of the medial aspect of the brain. Specifically, this connectivity retrieved the activation in the somatosensory cortex detected in the effect for evidence in the sample.

The data on the connectivity for the insular seed also included areas of negative association. These areas are shown as grey hatched areas in Figure 3b. One can see that, in the posterior portion of the brain, these areas of negative connectivity retrieved the areas associated with evidence for faces. In summary, the connectivity analysis showed existence of connections between the sparse areas activated in the decision task, and reproduced the dissociation between the system activated by the information from faces and a more general system active in both tasks. The data also suggest the possibility of a third network centered in the posterior cingulus/retrosplenial cortex and vmPFC. While the effects of evidence obtained in this third network were too weak to survive the correction for multiple comparisons, we draw attention to them here because these regions will resurface in the analysis later.

Figure 4. a: overlays for the effect of evidence for sad faces, drawn with overlays of task activation and deactivation (yellow and light blue colors). b: overlays for the common effect of evidence for both groups, together with the same overlays of task activation and deactivation (threshold for illustration purposes *p* < 0.01, uncorrected).

As noted in the Introduction, an issue we wished to address was the relationship between the hubs of the evidence network and task deactivations. In Figure 4, we drew the same effects of decision evidence of Figure 2 together with the activations and deactivations of the task (relative to the implicit baseline of the intertrial intervals). These activations and deactivations are very strong relative to noise levels, giving rise to high test statistics that are incompatible with the null hypothesis. Figure 4 shows that both evidence networks were located toward the end of the areas of activation, in some cases extending beyond it and encroaching the adjacent deactivated areas (in the inferior parietal region visible at $x = 22$ and $y = -27$ of Figure 4). This was especially the case for the faces network, whose hubs were loci located on the ribbon of the transition between activations and deactivations (Figure 4a). The evidence hubs in the medial

aspect of the brain were located entirely in deactivated areas. The correspondence of evidence networks and lack of strong activation is consistent with the course of the signal shown in Figure 2b.

Even in the areas in which the evidence networks were located in an area activated by the task, the activation effect was modest relative to the main activations. Figure 5a shows the contrasts for task activations and deactivations masked by the areas detected for the evidence in faces network. One can see that the location of the evidence network in the transitional zone between activations and deactivations more clearly, and the existence of a centrifugal gradient relative to a point in the occipital pole.

Figure 5. a: Overlays of task activations and deactivations (in warm and cold colors), thresholded at the uncorrected significance level _p_ < 0.001 and masked for the effect of evidence for faces (shown in Figure 4a). At _z_ = 28, the right retrosplenial hub is too weakly activated to appear in the figure. The figure shows the gradient of progressive task deactivation in the evidence hubs. b: signal course in the progression from activation to deactivation in the iPC/sTS hub in the faces task (at the points from _α_ to _δ_ marked in panel a). In red, signal for trials with high evidence for the decision; in light blue, trials with low evidence.

Figure 5b shows the progressive modulation of the task activation signal through the progression from activated to deactivated cortex in and around the iPC/sTS hub. One can see the emergence of a signal associated with high evidence for the decision to emerge when the task activation is weak or there is deactivation (loci _β_ and _γ_ in Figure 5). The locus _δ_ corresponds to the peak deactivation in the posterior iPC. Here, this signal is much weaker, if present at all.

The location of the evidence hubs at the transition to task deactivations raises the issue of whether the type of input (faces or mourning) changed the level of activity in the task vs. baseline contrast itself. If the evidence hubs are related to a decrement of activation, one would expect activation to be lower in the faces group, as these hubs are present only in this group. This is also the contrast we would adopt in a conventional analysis (even if using faces and mourning scenes as control for each other would not be entirely appropriate), except that we would look for a neural correlate in the form of higher instead of lower activation. It turns out that in the hubs and the adjacent cortex, activity was lower in the faces than in the mourning group, the opposite of what one would expect if activation in these areas reflected recruitment for the purposes of encoding sensory features (Figure 6; iPC/sTS hub, x, y, z: −34, −76, 42, t = 4.97, p = 0.04; −36, −78, 16, t = 5.25, p = 0.015; retrosplenial cortex, −14, −56, 18, t = 7.10, p < 0.001; 16, −50, 20, t = 6.20, p = 0.001, all peak-level corrected). In the faces group, activation was lower than in the mourning group also in the anterior fusiform gyrus hub (−30, −36, −16,

t = 6.75, *p* < 0.001; 32, −32, −14, *p* = 7.21, *p* < 0.001, peak-level corrected). These results do not contradict known findings about the effects of faces obtained with the cognitive subtraction approach. In more posterior portions of the fusiform gyrus, more active during the task, the activation was higher in the faces group, as expected (10, −72, −2, *t* = 5.60, *p* = 0.005, peak-level corrected). Likewise, the amygdala was more active in the faces than in the mourning group (26, 2, −26, *t* = 4.66, *p* = 0.09).

Figure 6. Effect of the contrast mourning vs. faces, thresholded for illustration purposes at p < 0.01, uncorrected, shown in the same locations as in Figure 5, but without masking.

In the final analyses of this study, we considered the change of the signal over time. The task signal has been shown to decrease over the experiment as an effect of practice, and sensory and semantic association areas show signal decrements consistent with their recruitment during encoding (sensory priming, Schacter and Buckner 1998; see also Grill-Spector et al. 2006; Barron et al. 2016). Figure 7a simultaneously shows the evidence for faces network, the pattern of task activations and deactivations of Figure 4, and the decrements of this task activation elicited by regressing the data on the number of times the items of the trial had been previously displayed (green hatch pattern; Table A3).

Figure 7. a: overlays for the effect of evidence in sad faces (blue-yellow), drawn with overlays of the effect of repetition (decreases, hatched areas in green) and of the third order interaction evidence x faces x repetition (hatched areas in violet). Task activations and deactivations are in yellow and light blue colors. b: overlays for the common effect of evidence for both groups (brown colors), together with the same other overlays as in panel a (threshold for illustration purposes *p* < 0.01, uncorrected). c: signal course and 90% confidence intervals at the iPS/sTS peak in the tertile of novel and most repeated images trials in the mourning group (left). On the right, the same plots for the posterior cingular locus (PC). The curves show the signal fitted by a Fourier series in the tertiles of the trials where the evidence for the decision were highest (easy decisions, red/orange color) and lowest (difficult decisions, blue color).

In the posterior parietal/occipital region and in the fusiform gyrus, decrements consistent with a priming effect were visible in the activated areas (Henson et al. 2002),

adjacent to and partially overlapping the inferior parietal hub of the evidence for faces network. Decrements over time were also observed in activated areas in the prefrontal cortex and in the ventral striatum/brainstem. These latter are known effect respectively of practice or conceptual priming (Schacter and Buckner 1998) and novelty (Bunzeck and Düzel 2006) and will not be commented further here.

When looking at the interaction between the effect of evidence and repetition of stimuli, we found a third order interaction evidence x group x repetition (Table A4), shown in Figure 7a and b as a violet hatch pattern. In the parietal region, this interaction can be seen to involve the deactivated areas adjacent to but distinct from the evidence for the faces hubs (PGa and PFm, Geyer et al. 1999). In the retrosplenial hub, the interaction appeared as a distinct locus in the posterior cingulus (Figure 7b), which however failed to reach significance after correction. The most extensive areas affected by this third order interaction were in the medial aspect of the brain, where it modulated task deactivations that hosted hubs of the common decision network. The strongest effects here were in the vmPFC and dmPFC, which we had seen involved by connectivity patterns in Figure 3.

Separate inspection of the interaction evidence x repetition within the groups revealed the third order interaction to arise entirely from the mourning scenes task, which showed a much more complex encoding of the evidence for sadness. In the faces group, the effect of evidence was stable across the whole experiment, giving no interaction with the effect of repetition. In contrast to faces, in the mourning scenes task the hubs were encoding evidence for the decision only briefly at the beginning of the experiment (Figure 7c). In the last part of the experiment, encoding was characterized by activity for neutral scenes, which precluded any simple mapping of high and low evidence trials to activity levels.

Discussion

In both groups, regressing the signal on the evidence for the decision revealed a network of areas centered in the somatosensory association cortex and the middle insula. Deciding on the basis of facial expressions, which offer rich sensory information for this task, additionally recruited the inferior parietal and postsplenial cortices, supporting the notion that recruitment of evidence hubs is to some extent selective. Furthermore, this selectivity was consistent with connectivity data suggesting the existence of at least two partially segregated networks accumulating evidence for the decision. This concordance represents an argument for the validity of the modelling strategy of contrasting easy with difficult decisions, since it retrieved areas that were shown to be preferentially connected in independent data. These findings are consistent with a generalized mechanism of evidence-accumulation, active also when deciding in an elementary social cognition task.

The hubs detected by modelling evidence for the decision are also consistent with data from cortical lesions. The main hub of the decision network recruited in both groups was located in the somatosensory association cortex, an area also implicated in studies of patients with impairments in classifying emotions (even if in lesions studies the effects are prevalently right-lateralized, Adolphs et al. 1996, 2000). The same studies also show a locus associated with impairment in the anterior fusiform gyrus (Adolphs et al. 1996), as in our data. The effect of lesions in the posterior iPC/sTS, the site of the faces evidence hubs and an area involved in the appraisal of emotional stimuli in imaging studies (Viviani 2013), are impairments in the computation of "a mental representation of extrapersonal events in terms of their motivational salience"

(Mesulam 1999). Also worth noting is the fact that, as apparent from Figure 4b, the precentral gyrus was not very active in the task, even if participants indicated their choice with a button press. However, in line with the alternative view, a cluster of the common decision network included the left precentral gyrus, which is consistent with the possibility that participants were accumulating evidence for button pressing (Cisek and Kalaska 2010). This finding is difficult to reconcile with the cognitive subtraction modelling strategy, as is the fact that none of the hubs was strongly activated by the task.

The course of the signal showed that the lack of activation was due to a brief deactivation of variable intensity at the presentation of the stimuli in the evidence hubs. This deactivation then progressed to activation at the time of the decision with a slope that was steeper in trials where the evidence of the decision was large. This pattern is visible in other decision making studies that report the signal course (Boorman et al. 2009; Viviani et al. 2017; in some of the areas reported by Tosoni et al. 2008). In contrast, in the adjacent constantly task-deactivated areas, the deactivation remained stable. In a sensory-based decision making study, Tosoni et al. (2008) showed that stably task-deactivated areas, attributed to the default network system, were distinct from those modulated by decision making. Their findings are consistent to those obtained here in so far as the parietal hubs, while contributing to modulating the activation in the faces vs. mourning contrast, were adjacent to but did not completely engulf the task deactivations. The initially activated evidence hubs in the mourning group, however, were located within the deactivations of the default network. The relationship of areas such as vmPFC and the inferior parietal cortex, often active in tasks of social cognition and emotion processing, and the default network has been noted before (Mars et al. 2012; Viviani 2014).

As whole, the fMRI signal in posterior part of the brain appeared to be organized in large portions of cortex characterized by a gradient of progressive deactivation. Sensory areas, strongly activated by the task, were succeeded by associative areas modulated by sensory priming. The sparse evidence network selectively recruited by decision about faces was located at the transition area to deactivations. The progression ended in the deactivated areas of the default network system. Areas associated with differential task activation identified in numerous previous studies that followed the cognitive subtraction approach were present also in our data. This was the case for the activation of posterior portion of the fusiform gyrus and of the amygdala in the faces task. However, these activations did not completely characterize the pattern of the fMRI signal.

If the organization of information processing in the posterior part of the brain has a neural correlate consisting of a gradient of progressive deactivation, questions arise about why this may be the case. While several related models have been developed in the psychological literature to explain empirical findings in decision making, it has been shown that all these models share a common fundamental computational step (Bogacz et al. 2006). Among the proposals for the implementation of this mechanism are several variations of competition between representations of the choice alternatives via mutual inhibition (Usher and McClelland 2001; Shadlen and Newsome 2001; Wang 2002; Cisek 2006). For this reason, evidence for decrements of the fMRI signal in preference-based choice tasks has been interpreted as evidence for this inhibition (Hunt et al. 2015), possibly mediated by GABA transmission (Jocham et al. 2012; for discussion, see Hunt and Hayden 2017). Since competition via mutual inhibition is thought to be a characteristic feature of neural circuits, one would expect decrements due to this mechanism to be present also in other choice tasks.

A similar conclusion may be reached when considering the theoretical literature on more general models of information processing. According the predictive coding model of cortical organization (Rao and Ballard 1999; Friston 2005; Kilner et al. 2007; Friston et al. 2013), the decision evidence signal arises by matching the predicted distribution of the signal of interest in the external world (in the present case, the inferred degree of sadness of the displayed individuals). The same distribution in the pre-motor areas provides the match at the level of the response. According to this model, the sensory and associative areas would be tuned by feedback so as to provide the signal in support of this prediction (referred to as a 'prediction error' relative to the invariant signal common to all stimuli). As a result, only the information that is relevant for the target classification of input is present in the activity of the associative areas. These tuned cortical areas may correspond to those where neural priming is observed (Auksztulewicz and Friston 2016), and that were here located between the early sensory processing areas and the evidence hubs.

In algorithmic implementations of predictive coding, where the relevant information in the input channel is represented by different pools of hidden units in a network, training is constrained by minimizing the amount of information in these units (Hinton and Zemel 1994). This increases the redundancy of the codes represented by the activation states of decision nodes. This account draws attention to the fact that in the evidence hubs the amount of information may be minimal, encoding only a position between two possible points (the two alternative decisions, here indexed by the relative degree of sadness). This makes the units that encode the relative degree of sadness to have similar activity, explaining why they may be all silent when the evidence for prediction is low. In contrast, early in the sensory processing channel the amount of information reflects relevant details of the visual input, requiring larger amounts of information to be coded. Also in this model, neural implementations are thought to make use of competitive inhibition to enforce progressive reduction in the amount of information (Hinton 2007).

In both models, deactivation (increased inhibition) cannot be equated with lack of recruitment, as in the cognitive subtraction approach. This is exemplified by the findings in the occipito-temporal cortex near the iPC/sTS hub and in the fusiform gyrus. In the posterior portion of the fusiform gyrus, where more basic features of the visual input may be expected to be processed, task activity was strong and increased in the faces task, which may be expected to make more intensive use of this type of visual information. Anteriorly, however, the evidence hub, where the evidence for the decision represents a high-level type of information, was located adjacent to the transition area to task deactivations, consistently with a stronger inhibitory influence.

If the progressive deactivation is related to inhibitory processes that constrain the complexity of representations of the encoded stimuli, the possibility arises that deactivated areas may host a parallel system that, operating side by side the activated centres of the executive dorsal network, organizes encoding of information. Whether this conjecture is correct or not, much appears to take place in the weakly active or deactivated ventral associative areas of the brain that is not captured by traditional analysis approaches, and has received comparatively little attention as an imaging phenotype.

Acknowledgements

This work was supported in part by a collaborative grant of the Federal Institute for Drugs and Medical Devices (BfArM, Bonn, Grant No. V-17568/68502/2017-2020) to Roberto Viviani. The authors declare no conflict of interest.

Appendix

Table A1 Evidence

Cl. #	Brain area	MNI coord.	k	p (cl.)	t	p (peak)
1	Left Inferior Parietal/Supramarginal (BA 40)	-56 -26 26	1055	0.007	6.16	<0.001
	Left Postcentral Gyrus (BA 1)	-54 -20 56			4.95	0.027
	Left Inferior Parietal Lobule (BA 40)	-60 -26 48			3.98	0.262
	Left Postcentral/Somatosensory (BA 1/43)	-62 -20 38			3.90	0.300
	Left Postcentral Gyrus (BA 3)	-44 -24 46			3.71	0.418
	Left Inferior Parietal Lobule (BA 40)	-58 -40 46			3.59	0.490
	Left Postcentral Gyrus (BA 3)	-42 -30 62			3.27	0.688
2	Left Insula (BA 13)	-38 0 14	258	0.044	5.33	0.007
	Left Primary Somatosensory Cortex (BA 43)	-54 -4 10			4.17	0.183
3	Right Supramarginal Gyrus (BA 40)	64 -36 42	466	0.019	4.98	0.024
	Right Supramarginal Gyrus (BA 40)	66 -26 30			4.62	0.062
	Right Rolandic Operculum (BA 43)/Supra-marginal Gyrus (BA 48)	52 -22 24			4.07	0.222
4	Right Insula (BA 13)	38 2 14	101	0.125	4.59	0.066
5	Left Precentral Gyrus (BA 6)	-32 -18 66	89	0.140	4.21	0.166

Legend. MNI coordinates: Montreal Neurological Institute coordinates, in mm; *k*: cluster size, in 2x2x2 mm voxels; *p* (cl.): significance level, cluster-level correction; *t*: peak *t*; *p* (peak): significance level, peak-level correction. The table lists up to 8 peaks per cluster, at minimum distance 8mm.

Table A2 Evidence x faces

Cl. #	Brain area	MNI coord.	k	p (cl.)	t	p (peak)
1	Left Parieto-occipital Gyrus (BA 39)	-42 -72 28	475	0.018	5.54	0.004
	Left Parieto-occipital Gyrus (BA 39)	-50 -72 22			5.19	0.012
2	Right Precuneus (BA 23)	16 -58 24	306	0.036	5.54	0.004
3	Left Precuneus (BA 31)	-10 -62 24	306	0.036	4.59	0.070
	Left Retrosplenial Cortex (BA 29)	-8 -48 4			3.61	0.482
4	Left Parahippocampal Gyrus (BA 27)	-30 -42 -8	93	0.137	4.24	0.164
	Left Fusiform Gyrus (BA 36/37)	-34 -36 -18			3.82	0.356
5	Right Angular/Parieto-occipital (BA 39/19)	48 -70 22	285	0.040	4.23	0.168
	Right Angular Gyrus (BA 39)	42 -70 32			4.05	0.242
6	Right Fusiform Gyrus (BA 37)	34 -36 -16	42	0.252	3.92	0.308

Legend. MNI coordinates: Montreal Neurological Institute coordinates, in mm; *k*: cluster size, in 2x2x2 mm voxels; *p* (cl.): significance level, cluster-level correction; *t*: peak *t*; *p* (peak): significance level, peak-level correction. The table lists up to 8 peaks per cluster, at minimum distance 8mm.

Table A3 Repetition (decreases)

Cl. #	Brain area	MNI coord.	k	p (cl.)	t	p (peak)

1	Right Inferior Frontal Gyrus (BA 45)	46 8 26	6197	0.001	8.72	<0.001
	Right Anterior Insula (BA 44/47)	32 24 0			7.68	<0.001
	Right Nucleus Accumbens	12 6 -2			7.25	<0.001
	Left Premotor Cortex/Supplementary Motor Cortex (BA 6)	-6 20 46			7.00	<0.001
	Right Inferior Frontal Gyrus (BA 45)	50 28 16			6.84	<0.001
	Right Supplementary Motor Cortex (BA 6)	8 12 52			5.72	0.003
	Right Precentral Gyrus (BA 6)	40 2 42			4.97	0.027
2	Left Anterior Insula (BA 44/47)	-30 22 2	4835	0.001	8.38	<0.001
	Left Premotor Cortex/Supplementary Motor Cortex (BA 6)	-48 10 26			6.95	<0.001
	Left Inferior Frontal Gyrus (BA44/45)	-44 18 24			6.83	<0.001
	Left Inferior Frontal Gyrus (BA 44)	-40 6 32			6.77	<0.001
	Left Nucleus Accumbens	-10 6 -4			6.70	<0.001
	Left Precentral Gyrus (BA 6)	-38 -2 56			6.28	<0.001
	Left Inferior Frontal Gyrus (BA 45)	-46 24 14			5.82	0.002
	Left Subgenual Ventromedial Prefrontal Cortex (BA 25)	-14 0 -10			5.54	0.005
3	Right Lingual Gyrus (BA 19)	24 -60 -6	3556	0.001	7.71	<0.001
	Right Middle Temporal Gyrus (BA 21)	48 -62 12			6.00	0.001
	Right Fusiform Gyrus (BA 37)	30 -38 -14			5.57	0.004
	Right Fusiform Gyrus (BA 37)	42 -46 -16			5.38	0.009
	Right Inferior/Middle Temporal Gyrus (BA 20/21)	48 -64 -2			5.37	0.009
	Right Superior Occipital Gyrus (BA 19)	26 -74 36			5.29	0.011
	Right Lingual Gyrus (BA 19)	24 -46 -8			5.12	0.017
	Right Middle Occipital Gyrus (BA 19)	36 -82 24			4.87	0.034
4	Left Fusiform Gyrus (BA37)	-26 -44 -10	4469	0.001	7.14	<0.001
	Left Fusiform Gyrus (BA37)	-28 -56 -10			7.08	<0.001
	Left Fusiform Gyrus (BA37)	-32 -72 -8			5.99	0.001
	Left Middle Temporal Gyrus (BA21)	-48 -66 4			5.87	0.002
	Left Middle Occipital Gyrus (BA 19)	-24 -74 32			5.71	0.003
	Left Middle Occipital Gyrus (BA 19)	-30 -84 30			5.53	0.005
	Left Middle Temporal Gyrus (BA 39)	-44 -66 16			5.42	0.008
	Left Fusiform Gyrus (BA 37)	-42 -48 -18			5.07	0.020
5	Left Middle Temporal Gyrus (BA 21)	-54 -6 -14	106	0.117	4.64	0.062
6	Right Postcentral Gyrus (BA 3)	44 -26 42	234	0.048	4.30	0.142
	Right Postcentral Gyrus (BA 3)	56 -20 40			3.73	0.405
	Right Postcentral Gyrus (BA 3)	36 -30 48			3.70	0.420
	Right Postcentral Gyrus (BA3)	64 -22 34			3.64	0.455
7	Left Postcentral Gyrus (BA 3)	-46 -26 38	176	0.070	4.20	0.180
1	Left Supramarginal Gyrus (BA 40)	-32 -34 40			4.09	0.224

| 8 | Right Primary Somatosensory Cortex (BA7) | 26 -48 54 | 53 | 0.209 | 3.85 | 0.331 |

Legend. MNI coordinates: Montreal Neurological Institute coordinates, in mm; *k*: cluster size, in 2x2x2 mm voxels; *p* (cl.): significance level, cluster-level correction; *t*: peak *t*; *p* (peak): significance level, peak-level correction. The table lists up to 8 peaks per cluster, at minimum distance 8mm.

Table A4. Interaction evidence x repetition (decreases) x mourning

Cl. #	Brain area	MNI coord.	*k*	*p* (cl.)	*t*	*p* (peak)
1	Left Superior Temporal Gyrus (BA 22)/Angular Gyrus (BA 39)	-54 -54 20	1380	0.005	6.12	<0.001
	Left Supramarginal Gyrus (BA 40)	-52 -58 30			5.87	<0.001
	Left Inferior Parietal Lobule/Supramarginal Gyrus (BA 40)	-48 -50 40			4.75	0.041
2	Left Middle Cingulate Cortex (BA 8)	-2 44 46	1587	0.004	5.61	0.003
	Left Middle Cingulate Cortex (BA 8)	-10 36 38			5.12	0.015
	Left Anterior Cingulate Cortex (BA 24)	-4 48 16			4.68	0.051
	Left Dorsomedial Prefrontal Cortex/Frontal Eye Field (BA 8)	-14 26 52			4.46	0.089
	Right Dorsolateral Prefrontal (BA 9)	10 48 32			4.37	0.111
	Left Middle Frontal Gyrus (BA 46)	-10 36 50			4.22	0.159
	Left Middle Frontal Gyrus (BA 46)	-18 58 24			3.93	0.272
	Left Middle Frontal Gyrus (BA 46)	-4 54 30			3.88	0.294
3	Right Supramarginal Gyrus (BA 40)	-38 18 44	296	0.038	4.55	0.069
	Right Supramarginal Gyrus (BA 40)	-48 18 36			3.72	0.383
4	Left Anterior Cingulate (BA 24)	-46 48 -4	149	0.072	4.27	0.141
5	Right Inferior Frontal Gyrus (BA 45)	54 -56 32	355	0.031	4.13	0.188
	Right Superior Temporal Gyrus (BA 22)/Angular Gyrus (BA 39)	58 -64 30			3.72	0.387
6	Left Supramarginal Gyrus (BA 40)	-6 36 -4	75	0.145	4.03	0.224
7	Left Inferior Parietal Lobule/Supramarginal Gyrus (BA 40)	-50 18 20	51	0.200	3.71	0.389

Legend. MNI coordinates: Montreal Neurological Institute coordinates, in mm; *k*: cluster size, in 2x2x2 mm voxels; *p* (cl.): significance level, cluster-level correction; *t*: peak *t*; *p* (peak): significance level, peak-level correction. The table lists up to 8 peaks per cluster, at minimum distance 8mm.

References

Adolphs, R., Damasio, H., Tranel, D., Cooper, G., Damasio, A.R., 2000. A role for somatosensory cortices in the visual recognition of emotion as revealed by three-dimensional lesion mapping. J. Neurosci. 20, 2683-2690.

Adolphs, R., Damasio, H., Tranel, D., Damasio, A.R., 1996. Cortical systems for the recognition of emotion in facial expression. J. Neurosci. 16, 7678-7687.

Auksztulewicz, R., Friston, K., 2016. Repetition suppression and ist contextual determinants in predictive coding. Cortex 80, 125-140.

Barron, H.C., Garvert, M.M., Behrens, T.E.J., 2016. Repetition suppression: A means to index neural representations using BOLD?. Phil. Trans. R. Soc. B 371, 20150355.

Benelli, E., Mergenthaler, E., Walter, S., Messina, I., Sambin, M., Buchheim, A., Sim, E.J., Viviani, R., 2012. Emotional and cognitive processing of narratives and individual

appraisal styles: Recruitment of cognitive control networks vs. modulation of deactivations. Front. Hum. Neurosci. doi, 10.3389/fnhum.2012.00239.

Bogacz, R., Brown, E., Moehlis, J., Holmes, P., Cohen, J.D., 2006. The physics of optimal decision making: A formal analysis of models of performance in two-alternative forced-choice tasks. Psychol. Rev. 113, 700-765.

Boorman, E.D., Behrens, T.E.J., Wollrich, M.W., Rushworth, M.F.S., 2009. How green is the grass on the other side? Frontopolar cortex and the evidence in favour of alternative courses of action. Neuron 62, 733-743.

Bunzeck, N., Düzel, E., 2006. Absolute coding of stimulus novelty in the human substantia nigra/VTA. Neuron 51, 369-379.

Caspers, S., Geyer, S., Schleicher, A., Mohlberg, H., Amunts, K., Zilles, K., 2006. The human inferior parietal cortex: Cytoarchitectonic parcellation and interindividual variability. NeuroImage 33, 430-448.

Cisek, P., 2006. Integrated neural processes for defining potential actions and deciding between them: A computational model. J. Neurosci. 26, 9770-9761.

Cisek, P., 2012. Making decisions through a distributed consensus. Curr. Opin. Neurobiol. 22, 927-936.

Cisek, P., Kalaska, J.F., 2010. Interacting with a world full of action choices. Annu. Rev. Neurosci. 33, 269-298.

Eickhoff, S.B., Stephan, K.E., Mohlberg, H., Grefkes, C., Fink, G.R., Amunts, K., Zilles, K., 2005. A new SPM toolbox for combining probabilistic cytoarchitectonic maps and functional imaging data. NeuroImage 25, 1325-1335.

FitzGerald, T.H.B., Seymour, B., Dolan, R.J., 2009. The role of human orbitofrontal cortex in value comparison for incommensurable objects. J. Neurosci. 29, 8388-8395.

Fodor, J.A., 1983. The Modularity of Mind. An Essay on Faculty Psychology. The MIT Press, Cambridge (Mass.).

Friston, K., 2005. A theory of cortical responses. Proc. R. Soc. Lond. B 360, 815-836.

Friston, K., Schwartenbeck, P., FitzGerald, T., Moutoussis, M., Behrens, T., Dolan, R.J., 2013. The anatomy of choice: Active inference and agency. Front. Hum. Neurosci. doi, 10.3389/fnhum.2013.00598.

Geyer, S., Schleicher, A., Zilles, K., 1999. Areas 3a, 3b, and 1 of human primary somatosensory cortex: 1. Mictrostructural organization and interindividual variability. NeuroImage 10, 63-83.

Glimcher, P., 2015. Understanding the hows and whys of decision making: From expected utility to divisive normalization. Cold Spring Harb. Symp. Quant. Biol. doi, 10.1101/sqb.2014.79.024778.

Gold, J.I., Heekeren, H.R., 2014. Neural mechanisms for perceptual decision making. In: Glimcher, P.W., Fehr, E. (Eds). Neuroeconomics: Decision Making and the Brain, pp. 355-372.

Gold, J.I., Shadlen, M.N., 2002. Banburism and the brain: Decoding the relationship between sensory stimuli, decisions, and reward. Neuron 36, 299-308.

Grill-Spector, K., Henson, R., Martin, A., 2006. Repetition and the brain: Neural models of stimulus-specific effects. Trends Cogn. Sci. 10, 14-23.

Heekeren, H.R., Marrett, S., Bandettini, P.A., Ungerleider, L.G., 2004. A general mechanism for perceptual decision-making in the human brain. Nature 431, 859-862.

Heekeren, H.R., Marrett, S., Ungerleider, L.G., 2008. The neural systems that mediate human perceptual decision making. Nature Rev. Neurosci. 9, 467-479.

Henson, R.N.A., Shallice, T., Gorno-Tempini, M.L., Dolan, R.J., 2002. Face repetition effects in implicit and explicit memory tests as measured by fMRI. Cereb. Cortex 12, 178-186.

Hinton, G.E., 2007. Learning multiple layers of representation. Trends Cogn. Sci. 11, 428-434.

Hinton, G.E., Zemel, R.S., 1994. Autoencoders, minimum description length and Helmholtz free energy. Advances in Neural Information Processing Systems 1994, 3-10.

Hunt, L.T., Dolan, R.J., Behrens, T.E.J., 2015. Hierarchical competitions subserving multi-attribute choice. Nat. Neurosci. 17, 1613-1622.

Hunt, L.T., Hayden, B.Y., 2017. A distributed, hierarchical and recurrent framework for reward-based choice. Nature Rev. Neurosci. 16, 172-182.

Hunt, L.T., Kolling, N., Soltani, A., Woolrich, M.W., Rushworth, M.F.S., Behrens, T.E.J., 2012. Mechanisms underlying cortical activity during value-guided choice. Nat. Neurosci. 15, 470-476.

Jocham, G., Hunt, L.T., Near, J., Behrens, T.E.J., 2012. A mechanism for value-guided choice based on the excitation-inhibition balance in prefrontal cortex. Nat. Neurosci. 15, 960-961.

Kayser, A.W., Buchsbaum, B.R., Erickson, D.T., D'Esposito, M., 2010. The functional anatomy of a perceptual decision in the human brain. J. Neurophysiol. 103, 1179-1194.

Kilner, J.M., Friston, K.J., Frith, C.D., 2007. Predictive coding: An account of the mirror neuron system. Cogn. Process. 8, 159-166.

Krajbich, I., Rangel, A., 2011. Multialternative drift-diffusion model predicts the relationship between visual fixations and choice in value-based decisions. Proc. Natl Acad. Sci. USA 108, 13852-13857.

Labek, K., Berger, S., Buchheim, A., Bosch, J., Spohrs, J., Dommes, L., Beschoner, P., Stingl, J.C., Viviani, R., 2017. The iconography of mourning and its neural correlates: A functional neuroimaging study. Soc. Cogn. Aff. Neurosci. 12, 1303-1313.

Lamm, C., Decety, J., Singer, T., 2011. Meta-analytic evidence for common and distinct neural networks associated with directly experienced pain and empathy for pain. NeuroImage 54, 2492-2502.

Lorenz, S., Weiner, K.S., Caspers, J., Mohlberg, H., Schleicher, A., Bludau, S., Eickhoff, S.B., Grill-Spector, K., Zilles, K., Amunts, K., 2017. Two new cytoarchitectonic areas on the human mid-fusiform gyrus. Cereb. Cortex 27, 373-385.

Lundqvist, D., Flykt, A., Öhman, A., 1998. The Karolinska Directed Emotional Faces - KDEF. CD ROM from Deparment of Clinical Neuroscience Psychology section, Karolinska Institutet, .

Mars, R.B., Neubert, F.X., Noonan, M.A.P., Sallet, J., Toni, I., Rushworth, M.F.S., 2012. On the relationship between the 'default mode network' and the 'social brain'. Front. Hum. Neurosci. doi, 10.3389/fnhum.2012.00189.

Mesulam, M.-M., 1999. Spatial attention and neglect: Parietal, frontal and cingulate contributions to the mental representation and attentional targeting of salient extrapersonal events. Phil. Trans. R. Soc. B 354, 1325-1346.

Petersen, S.E., Fox, P.T., Posner, M.I., Mintun, M., Raichle, M.E., 1989. Positron emission tomographic studies of the processing of single words. J. Cogn. Neurosci. 2, 153-170.

Peyron, R., Fauchon, C., 2019. The posterior insular-opercular cortex: An access to the brain networks of thermosensory and nociceptive processes? Neurosci. Lett. 702, 34.39.

Posner, M.I., Raichle, M.E., 1994. Images of Mind. Freeman, New York.

Rao, R.P.N., Ballard, D.H., 1999. Predictive coding in the visual cortex: A functional interpretation of some extra-classical receptive-field effects. Nat. Neurosci. 2, 79-87.

Rypma, B., Prabhakaran, V., Desmond, J.E., Glover, G.H., Gabrieli, J.D.E., 1999. Load-dependent roles of frontal brain regions in the maintenance of working memory. NeuroImage 9, 216-226.

Schacter, D.L., Buckner, R.L., 1998. Priming and the brain. Neuron 20, 185-195.

Serences, J.T., 2008. Value-based modulations in human visual cortex. Neuron 60, 1169-1181.

Shadlen, M.N., Kiani, R., 2013. Decision making as a windon on cognition. Neuron 30, 791-806.

Shadlen, M.N., Newsome, W.T., 2001. Neural basis of a perceptual decision in the parietal cortex (Area LIP) of the rhesus monkey. J. Neurophysiol. 86, 1916-1936.

Tosoni, A., Galati, G., Romani, G.G., Corbetta, M., 2008. Sensory-motor mechanisms in human parietal cortex underlie arbitrary visual decisions. Nat. Neurosci. 11, 1446-1453.

Usher, M., McClelland, J.L., 2001. On the time course of perceptual choice: The leaky competing accumulator model. Psychol. Rev. 108, 550-592.

Viviani, R., 2013. Emotion regulation, attention to emotion, and the ventral attentional network. Front. Hum. Neurosci. doi, 10.3389/fnhum.2013.00746.

Viviani, R., 2014. Neural correlates of emotion regulation in the ventral prefrontal cortex and the encoding of subjective value and economic utility. Frontiers Psychiatry doi, 10.3389/fpsyt.2014.00123.

Viviani, R., Dommes, L., Bosch, J., Beschoner, P., Stingl, J.C., Schnell, T., 2017. Choosing between human values. bioRxiv, doi: https://doi.org/10.1101/856294.

Viviani, R., Dommes, L., Bosch, J., Stingl, J.C., Beschoner, P., 2018. The neural correlates of decisions about sadness in facial expressions. J. Neurosci. Psychol. Econ. 11, 93-105.

Vuilleumier, P., Armony, J.L., Driver, J., Dolan, R.J., 2003. Distinct spatial frequency sensitivities for processing faces and emotional expressions. Nat. Neurosci. 6, 624-631.

Wang, X.J., 2002. Probabilistic decision making by slow reverberation in cortical circuits. Neuron 36, 1-20.

Yarkoni, T., Poldrack, R.A., Nichols, T.N., Van, Essen, D.C., Wager, T.D., 2011. Large-scale automated synthesis of human functional neuroimaging data. Nat. Meth. 8, 665-670.