RIG-I and PKR, but not stress granules, mediate the pro-inflammatory response to

Yellow fever virus

Guillaume Beauclair[1], Felix Streicher[1], Daniela Bruni[1], Ségolène Gracias[1], Salomé Bourgeau[1], Laura

Sinigaglia[1], Takashi Fujita[2], Eliane F. Meurs[1], Frédéric Tangy[1] and Nolwenn Jouvenet[1]*

[1] Viral Genomics and Vaccination Unit, Institut Pasteur, UMR3569 CNRS, 75015 Paris, France

[2] Department of Virus Research, Institute for Frontier Life and Medical Sciences, Kyoto University,

Kyoto, Japan.

Short title: Innate response against YFV in liver cells

*Corresponding author: nolwenn.jouvenet@pasteur.fr

Abstract

Yellow fever virus (YFV) is a single-stranded RNA virus primarily targeting the liver. Severe YF cases are responsible for viral hemorrhagic fever, plausibly precipitated by excessive pro-inflammatory cytokine response, resulting in up to 50% fatality. Pathogen recognition receptors (PRRs), such as the ubiquitously expressed cytoplasmic RIG-I-like receptors (RLRs), as well as the viral RNA sensor PKR are known to initiate a pro-inflammatory response upon recognition of viral genomes and replication intermediates. The PRRs responsible for cytokine production in cells infected with YFV strains are not known.

Here, we sought to reveal the main determinants responsible for the acute cytokine expression occurring in human hepatocytes following YFV infection. Using a RIG-I-defective human hepatoma cell line, we found that RIG-I largely contributes to cytokine secretion upon infection with the Asibi strain. In infected RIG-I-proficient hepatoma cells, RIG-I was localized in stress granules. These granules are large aggregates of stalled translation preinitiation complexes known to concentrate RLRs and PKR, and are so far recognized as hubs orchestrating RNA virus sensing. Using PKR-deficient hepatoma cells, we found that PKR contributes to both stress granule formation and cytokine induction upon Asibi infection. However, we showed that stress granules disruption did not affect the cytokine response to Asibi infection, as assessed by siRNA-knockdown-mediated inhibition of stress granule assembly and an automated determination of granule size and number. Finally, no viral RNA was detected in stress granules using a fluorescence *in situ* hybridization approach coupled with immunofluorescence.

Collectively, our data strongly suggest that both RIG-I and PKR mediate pro-inflammatory cytokine induction in YFV-infected hepatocytes, in a stress granule-independent manner. Therefore, by showing the uncoupling of the early cytokine response from the stress granules formation, our model challenges the current view by which stress granules are required for the mounting of the acute antiviral response.

Author Summary

Yellow fever is a mosquito-borne acute hemorrhagic disease caused by yellow fever virus (YFV). The

mechanisms responsible for its pathogenesis remain largely unknown, although increased inflammation has

been linked to worsened outcome. In the early phases of infection, YFV targets the liver, where it primarily

infects hepatocytes. Pathogen-recognition receptors (PRRs) are the first immune sensors detecting microbial

infections. The PRR(s) responsible for the recognition of the YFV in human hepatocytes remains to be

established. Here, we first show that YFV's sensing by two PRRs, RIG-I and PKR, elicits the induction and

production of pro-inflammatory mediators in human hepatocytes. We show that YFV infection promotes the

formation of cytoplasmic structures, termed stress granules, in a PKR-, but not RIG-I-dependent manner, and

that these structures are dispensable for downstream signaling via these receptors. Whilst stress granules were

previously postulated to be essential platforms for PRRs activation, here we first uncovered that they are not

required for pro-inflammatory mediators' production upon YFV infection. Collectively, our work uncovered

the molecular events triggered by YFV, which could prove instrumental in clarifying the pathogenesis of the

disease, with possible repercussions on disease management.

Introduction

Flaviviruses are a group of more than seventy enveloped RNA viruses that can cause serious diseases in

humans and animals [1,2]. They have provided some of the most important examples of emerging or

resurging diseases of global significance. Most flaviviruses, such as dengue virus (DENV), yellow fever

virus (YFV) and Zika virus (ZIKV), are arthropod-borne viruses transmitted to vertebrate hosts by

mosquitoes or ticks [1]. YFV is the prototype and eponymous virus of the Flavivirus genus. It is a small

(40–60 nm), enveloped virus harboring a single positive-strand RNA genome of 11 kb. The genome

encodes a polyprotein that is cleaved co- and post-translationally into three structural proteins: capsid

(C), membrane precursor (prM) and envelope (E), and seven non-structural (NS) proteins (NS1, NS2a,

NS2b, NS3, NS4a, NS4b, and NS5). The C, prM, and E proteins are incorporated into the virions, while

NS proteins are found only in infected cells [3]. NS proteins coordinate RNA replication, viral assembly

and modulate innate immune responses, while the structural proteins constitute the virion.

YFV is endemic in the tropical regions of sub-Saharan Africa and South America. The reference

strain Asibi was isolated in 1927 in West Africa from the blood of a human patient. The vaccine strain

17D was developed empirically in the 1930s, by passaging the Asibi strain in rhesus macaques, mouse

and chicken embryonic tissues [4]. 17D is one of the most effective vaccines ever generated. It has been

used safely and effectively on more than 600 million individuals over the past 70 years [4]. However,

due to poor vaccine coverage and vaccine shortages, the virus continues to cause disease in endemic areas,

as illustrated by recent outbreaks in Angola and Brazil [5,6].

Yellow fever (YF) pathogenesis is viscerotropic in humans, with viral replication in the liver critical

to the establishment of the disease [3]. Severe YF is responsible for multi-system organ failure and viral

hemorrhagic fever resulting in up to 50% fatality. Similar to Ebola hemorrhagic fever, cytokine dysregulation

is thought to result in endothelial damage, disseminated intravascular coagulation and circulatory shock

observed in the terminal stage of the disease. Viral hemorrhagic fever is considered as an illness

precipitated by excessive pro-inflammatory cytokine response ('cytokine storm') [7,8]. Almost any cell

type in the body can produce pro-inflammatory cytokines in response to various stimuli, such as a viral

infection [9]. The pattern of cytokine expression in response to infection with pathogenic strains of YFV

has been described both in humans and in rhesus macaques. Studies performed on patients during a YF

outbreak in the Republic of Guinea in 2000 showed that excessive production of pro-inflammatory

cytokines, such as IL6 and TNFα, correlates with high viremia and disease outcome [10]. The levels of

these two cytokines were statistically higher in patients with fatal hemorrhagic fever than in patients

who survived. Similar findings were obtained with rhesus macaques infected with the strain Dakar-

HD1279 [11]. The animals died of hemorrhagic fever within five days. These studies also suggest that

pro-inflammatory cytokines might be produced by injured organs, especially the liver, and not by cells

circulating in the blood [11].

Upon viral infection, the cytokine response is initiated by the recognition of viral genomes and viral

replication intermediates by pathogen recognition receptors (PRRs) [12], such as the ubiquitously

expressed cytoplasmic RIG-I-like receptors (RLRs). Three RLRs have been identified in vertebrates:

melanoma differentiation-associated gene 5 (MDA5), laboratory of genetics and physiology 2 (LGP2)

and retinoic acid inducible gene 1 (RIG-I) [13]. Upon binding to viral nucleic acids, MDA5 and RIG-I

undergo a conformational change, which allows interaction with the adaptor protein MAVS and

subsequent recruitment of signaling complexes that comprises protein kinases such as TBK1, IKKα,

IKKβ and IKKε [14]. These events lead to the activation and nuclear translocation of the transcription

factors IRF3 and NF-κB, which stimulate the rapid expression of pro-inflammatory cytokines and type

I interferons (IFNs).

RIG-I plays an important role in initiating the IFN-mediated antiviral response against DENV and

ZIKV [15–18]. Induction of cytokines upon infection with the vaccine strain of YFV is also mediated

by RIG-I in plasmacytoid dendritic cells [19]. Whether RIG-I contributes to cytokine production in cells

infected with wild-type YFV strains is not known.

The subcellular localization where RIG-I interacts with its ligands is not well described. Upon

infection with influenza A virus lacking its non-structural protein 1 (IAVΔNS1), Newcastle disease

virus or Sendai virus [20–23], RIG-I accumulates in cytoplasmic foci. Disruption of these foci, which

have been identified as stress granules, reduced the abundance of *IFNB* mRNAs in IAVΔNS1-infected

cells [20]. Thus, stress granules may be important for RIG-I signaling and could serve as a platform for

viral RNA sensing. Moreover, other viral nucleic sensors, such as MDA5 [24], LGP2 [25] and cGAS

[26], are also concentrated in stress granules. Finally, numerous viruses interfere with stress granule

formation, *via*, for instance, the cleavage of one of their core components by a viral protease. Thus,

accumulation of viral sensors in stress granules and viral counteraction mechanisms suggest that these

granules act as hubs that coordinate foreign nucleic acid sensing [27,28].

Stress granules are non-membranous cytoplasmic structures that concentrate translationally stalled

mRNAs and proteins [29]. Their assembly is triggered by sensing distinct types of stress, including viral

infection, by one of four kinases, resulting in phosphorylation of the eukaryotic initiation factor 2α

(eIF2α). These kinases are: the heme-regulated inhibitor (HRI), which is activated by heat shock or

oxidative stress (*e.g.* by reactive oxygen species (ROS) produced upon treatment with sodium arsenite);

general control non-depressible protein 2 (GCN2), which reacts to UV-induced damage or amino acid

deprivation; dsRNA-sensing protein kinase R (PKR); and PKR-like endoplasmatic reticulum (ER)

kinase (PERK), which is activated upon ER stress [29]. Stress granule assembly does not only arrest the host translation machinery, but also heavily disturb viral translation, which is further evidence for their antiviral nature [27,28,30]. Here, we investigated the role of RIG-I, PKR and stress granules in the induction inflammatory cytokines upon infection of hepatoma cells with YFV.

Results

The Asibi strain of YFV triggers the production of pro-inflammatory cytokines in hepatoma cells through RIG-I, IRF3 and NF-κB

We examined whether the YFV reference strain Asibi could trigger the expression of pro-inflammatory cytokines in human hepatoma cells through the RIG-I signaling pathway. Huh7 and Huh7.5 cells, a clone derived from Huh7 that expresses an inactive form of the viral sensor RIG-I [31], were infected with the Asibi strain for 24 or 48 hours. The abundances of *TNFA*, *IL6* and *IFNB* mRNAs increased over time with infection in Huh7 cells, while being significantly lower in Huh7.5 cells at 48 hours post-infection (hpi) (**Fig 1A-C**). Addition of BAY11-7085 (BAY), an irreversible inhibitor of IκBα phosphorylation that prevents activation of NF-κB [32], reduced induction of *TNFA*, *IL6* and *IFNB* mRNAs in Huh7 cells infected with Asibi (**Fig 1A-C**). However, the mRNA levels of these three cytokines in BAY-treated Huh7 cells were higher compared to those in Huh7.5 cells (**Fig 1A-C**). Infection of the Huh7 cells, either treated with BAY or left untreated, produced similar amounts of viral RNA (**Fig 1D)**, which suggests that the chemical treatment did not affect viral replication. Furthermore, we noticed that Huh7.5 cells infected for 48 hours produced approximately three-fold more viral RNA than Huh7 cells infected for the same amount of time (**Fig 1D)**, suggesting that absence of RIG-I signaling favors viral replication. Untreated Huh7 cells secreted around 10 to 20 pg/ml of IL6 and TNFα protein after 48 hours of infection, but not after 24 hpi (**Fig 1E, F**). By contrast, the amount of IL6 and TNFα protein secreted by cells treated with the inhibitor of IκBα phosphorylation or expressing the inactive form of RIG-I was below the detection limit of the ELISA (**Fig 1E, F**). Together, these data suggest that Asibi infection triggers the induction and secretion of pro-inflammatory cytokines in a RIG-I- and NF-κB- dependent manner in human hepatoma cells.

By Western blot analysis, we found that IRF3 was activated in Huh7 cells infected with Asibi for

24 or 48 hours, but not in non-infected cells (**Fig 1G**). The level of IRF3 phosphorylation was higher at

48 than at 24 hpi, despite the total protein abundance being similar at these two points in time (**Fig 1G**).

IRF3 was not phosphorylated in Huh7.5 cells infected with Asibi (**Fig 1G**). In Huh7 cells treated with

BAY, IRF3 was not phosphorylated at 24 hpi (**Fig 1G**). Phosphorylation was present at 48 hpi, tough in

small levels than in non-treated cells (**Fig 1G**). IκBα is a negative regulator of the NF-κB pathway: its

degradation triggers the nuclear translocation of the NF-κB subunits p65 and p50. IκBα was degraded

in Huh7 cells infected for 48 hours but less so in cells treated with BAY for 48 hours (**Fig 1G**). Slightly

more of the viral protein NS4b was produced in Huh7.5 cells than in Huh7 cells upon 48 hours of Asibi

infection (**Fig 1G**), which is consistent with the RT-qPCR analysis (**Fig 1D**). These experiments indicate

that both IRF3 and NF-κB are activated during Asibi infection of human hepatoma cells and that RIG-

I contributes to IRF3 activation.

Immunofluorescence analysis revealed that IRF3 was detectable exclusively in the cytosol of 100%

of uninfected cells, whereas it was found both in the cytosol and the nucleus of around 50% of cells

infected for 24 hours (**Fig 1H**). Infected cells were identified with antibodies recognizing dsRNA

structures, including probably viral replicative intermediates (**Fig 1H**). Similar quantification showed

that around 3% of uninfected cells exhibited nuclei positive for the NF-κB subunit p65 whereas around

65% of Asibi-infected cells showed nuclei positive for p65 (**Fig 1I**).

Together, these results suggest that both the NF-κB and the IRF3 pathways are involved in cytokine

secretion in human hepatoma cells upon Asibi infection and that their initiation is dependent on RIG-I.

RIG-I localizes in stress granules in human and monkey cells infected with YFV

Having determined that RIG-I is important for mediating a cytokine response in Huh7 cells infected

with Asibi (**Fig 1**), we then investigated its subcellular localization. The distribution of the RIG-I signal

was found to be weak and diffuse in the cytosol of uninfected Huh7 cells (**Fig 2A**), which is consistent

with data reported for unstimulated HeLa cells [20]. After infection with Asibi for 24 hours, RIG-I

accumulated in cytosolic foci (**Fig 2A**). Since RIG-I was previously shown to localize in stress granules

in HeLa cells infected with several types of viruses [20–23], the Huh7 cells were also stained with the

stress granule marker TIA-1 related protein (TIAR). In non-infected cells, the TIAR signal was weak

and diffused in the cytoplasm and in the nucleus of the cells. In infected cells, TIAR was redistributed

into cytoplasmic foci (**Fig 2A**). The RIG-I structures present in infected cells co-localized with TIAR

foci (**Fig 2A**, zoom images for details), suggesting that RIG-I is recruited to stress granules upon

infection. The intensity profiles of individual foci confirmed that the RIG-I signal and the TIAR signal

overlapped (**Fig 2A**, right panels). These profiles also revealed that dsRNAs seemed to be excluded

from stress granules (**Fig 2A**).

To follow the formation of stress granules in infected cells, the localization of TIAR and RAS

GTPase-activating protein SH3 domain-binding protein 1 and 2 (G3BP), which are canonical stress

granule markers [33], were examined in Huh7 cells infected with Asibi for over time (**Fig 2B**). Infected

cells were identified by the presence of the viral protein NS4b (**Fig 2B**). TIAR and G3BP were diffuse

in non-infected cells. They remained diffuse until 12 hpi, where 16% of cells were determined positive

for NS4b. Fifteen hours after infection, around 80% of cells were infected and around 40% of the total

number of cells exhibited TIAR- and G3BP-positive foci (**Fig 2B**). At 24 hpi, all cells were infected and

around 70% of them were positive for TIAR- and G3BP- containing foci. At 37 hpi, all cells remained

infected and around 20% of them exhibited TIAR- and G3BP positive foci. These foci resembled the

ones detected in Huh7 cells treated with sodium arsenite (NaArs), a chemical commonly used for

induction of oxidative stress in eukaryotic cells, for 30 minutes (**Fig 2B**). These results indicate that

upon Asibi infection, TIAR- and G3BP- positive foci appeared between 12 and 15 hpi, peak around 24

hpi and persist in around 20% of cells until later times. Moreover, these data suggest that the formation

of the stress granules occur throughout infection with Asibi.

To further validate the nature of the TIAR- and G3BP- positive foci observed in infected cells, we

assessed their sensitivity to cycloheximide. Cycloheximide is a compound that inhibits stress granule

formation and fosters their disassembly by stabilizing mRNAs on polysomes [34]. Huh7 cells treated

with NaArs for 30 min and subsequently incubated with cycloheximide for 1 hour or left untreated were

used as positive controls (**Fig 2C**). Alternatively, cells were infected with Asibi for 48 hours and then

treated, or not, with cycloheximide for 1 hour. The TIAR and the G3BP signals overlapped, both in cells

infected by Asibi for 48 hours and in NaArs-treated cells (**Fig 2C**). Cycloheximide treatment led to a

complete loss of TIAR- and G3BP- positive foci, both in NaArs -treated cells and in Asibi-infected cells

(**Fig 2C**, compare non-treated and treated cells). These results suggest that, under both conditions, most

mRNAs trapped in these cytoplasmic foci can be released within 1 hour and undergo some translation

on polysomes. Most importantly, these experiments confirmed that the cytoplasmic foci containing

TIAR, G3BP and RIG-I identified in infected cells also contained mRNAs, and as such, can be defined

as canonical stress granules. Furthermore, these experiments also showed that the stress granule proteins

TIAR and G3BP are not recruited to Asibi replication complexes identified by NS4b,in contrast to

observations reported for ZIKV, DENV and WNV [35,36].

We next monitored the formation of stress granules upon Asibi infection in different cell models

than Huh7. Like Huh7 cells, HepG2 cells are hepatocyte-derived cells and are thus relevant for infection

with YFV, a hepatotropic virus. TIAR- and G3BP-positive foci were detected in HepG2 cells infected

for 48 hours (**Fig 3A**). Stress granules were also detected in CMMT cells, a *Macaca mulatta* epithelial

cell line, upon 24 hours of infection with Asibi (**Fig 3B**). Macaques are natural hosts of YFV during the

jungle cycle of transmission [37]. Similarly, TIAR- and G3BP-positive foci were observed in Huh7 cells

infected with the clinical isolate HD1279 (HD Dakar) for 24 hours (**Fig 3C**). These data indicate that

infection with two YFV strains triggers the formation of stress granules in several mammalian cell lines,

suggesting that the stress response is neither viral strain nor cell type specific.

PKR contributes to stress granule formation and innate response in Asibi-infected Huh7 cells

PKR is known to be activated upon infection with several flaviviruses, including ZIKV, WNV and

Japanese Encephalitis virus (JEV) [38–40] and to play an important role in stress granule nucleation

through its ability to phosphorylate the eukaryotic initiation factor eIF2α [41]. We first assessed whether

PKR was activated upon Asibi infection of Huh7 cells. Western blot analysis showed that PKR was

phosphorylated in Huh7 cells infected for 24 and 48 hours, but not in non-infected cells (**Fig 4A**). Thus,

PKR is activated upon Asibi infection of Huh7 cells. To evaluate the role of PKR in the formation of

stress granules in Asibi-infected human hepatic cells, we generated a Huh7 short hairpin (sh)-PKR cell

line by transduction with lentiviruses expressing short hairpin RNAs (shRNAs) specific for PKR. While

PKR expression was detectable in Huh7 cells transduced with lentivirus expressing control shRNA

targeting luciferase (Huh7-shLuc cells), it was not detectable in Huh7-shPKR cells (**Fig 4B**). As a

control experiment, we treated the cells with NaArs, which induces stress granule formation *via* HRI-

mediated eIF2α phosphorylation [42]. TIAR- and G3BP- positive stress granules were detected in both

Huh7-shLuc and Huh7-shPKR cells treated with NaArs for 30 minutes (**Fig 3C**). This confirms that the

absence of PKR does not disrupt the formation of NaArs-dependent stress granule induction.

Automated microscopic quantification revealed that around 10% of Huh7-shLuc cells infected at a

MOI of 2 for 24 or 34 hours contained at least two stress granules (**Fig 4 D-E**). At a MOI of 20, around

25% of Huh7-shLuc cells were positive for stress granules at 24 and 34 hpi (**Fig 4 D-E**). No TIAR- nor

G3BP- positive foci were detected in Huh7-shPKR cells infected at a MOI of 2 for 24 or 34 hours. At a

MOI of 20, a low percentage (around 1%) of Huh7-shPKR cells exhibited detectable stress granules

(**Fig 4 D-E**). These stress granules possibly formed in a PKR-independent manner, *via* the activation of

other members of the eIF2α kinase family. Alternatively, a minute quantity of PKR still being expressed

in shPKR cells may initiate the formation of these granules. A similar proportion of Huh7-shLuc and

Huh7-shPKR cells were found positive for NS4b at 24 or 34 hpi at the two tested MOIs (**Fig 4F**).

Together, we conclude that PKR plays a key role in stress granule formation in Asibi-infected human

hepatic cells and that the absence of stress granules has no effect on viral infection.

We next tested whether other members of the eIF2α kinase family are involved in stress granule

formation in Huh7 cells infected with Asibi. Therefore, cells infected for 24 hours were treated for four

hours with either GSK2606414 (GSK) or GCN2-IN-1 (iGCN2), which are inhibitors of PERK [43] and

GCN2, respectively. Around 78% of control DMSO-treated cells were positive for TIAR- and G3BP-

foci at 24 hpi (**Fig 5**). Only 2% and 4% of GSK- and iGCN2- treated cells, respectively, exhibited stress

granules (**Fig 5**), indicating that both inhibitors suppressed Asibi-induced stress granule formation.

These data suggest that at least three of the four eIF2α kinases (GCN2, PERK and PKR) contribute to

stress granule formation in human hepatoma cells infected with Asibi. Thus, Asibi infection induces an

integrated stress response in Huh7 cells.

Since no stress granules were detected in cells depleted of PKR in three of four conditions tested

(**Fig 4E**), we used Huh7-shPKR cells to determine whether absence of granules affected cytokine

induction. Consistent with our Western blot data (**Fig 4B**), RT-qPCR analysis showed that *PKR* mRNAs

abundance was reduced by more than 90% in Huh7-shPKR cells in comparison to Huh7-shLuc cells

(**Fig 6A**). These RT-qPCR analyses also showed that the viral RNA yield in Huh7-shPKR cells was not

markedly different from that in Huh7-shLuc cells 24 hours or 34 hours after infection either at a MOI

of 2 or 20 (**Fig 6B**), which is consistent with our microscopic analysis (**Fig 4F**). *IL6* mRNA abundance

was significantly lower in Huh7-shPKR cells than that of Huh7-shLuc when cells were infected at a

MOI of 20 (**Fig 6C**). The abundances of *TNFA* and *IFNB* mRNAs were significantly lower in Huh7-

shPKR cells than those in infected Huh7-shLuc under the same conditions of infection (**Fig 6D-E**). This

is consistent with our preceding observation of cytokine induction in Asibi infected cells being

dependent on NF-κB (**Fig 1**) and with a previous report showing that PKR is involved in the induction

of the *IFN* through the activation of NF-κB [44]. Thus, at least two nucleic acid sensors, namely RIG-I

(**Fig 1**) and PKR (**Fig 4**), contribute to the antiviral response against Asibi in Huh7 cells.

Taken together, our data show that PKR contributes to both stress granule formation and induction

of cytokines in Huh7 cells infected with Asibi. However, since PKR does not require its kinase activity

to stimulate NF-κB [44,45], it might contribute to the cytokine induction in Asibi-infected cells in a

eIF2α- (and thus stress granule-) independent manner.

Inhibition of stress granule formation has a limited effect on antiviral response

To further investigate the putative link between stress granules and the induction of antiviral

cytokines, we used a siRNA-knockdown approach to inhibit stress granule assembly. We tested

numerous combinations of siRNAs targeting core granule components in infected cells. The most

efficient strategy consisted in transfecting Huh7 cells twice with four pools of siRNAs targeting TIAR,

TIA-1, G3BP1 and G3BP2 (siSG). In these cells, we monitored stress granule formation using

antibodies against eIF3b and eIF4G, two core components of stress granules. Around 30% of Huh7 cells

transfected with scrambled control siRNAs and infected at a MOI of 2 or 20 for 24 hours contained at

least two eIF4G- and eIF3b-positive foci (**Fig 7 A-B**). When cells were transfected with the 4 siRNA

pools and infected under the same conditions and for the same amount of time, less than 5% of cells

exhibited stress granules (**Fig 7 A-B**). The impact of the siRNA against TIAR, TIA-1, G3BP1 and

G3BP2 on the number of cells positive for stress granules following 34 hours of infection was also

significant at the two tested MOIs (**Fig 7B**). Under the four tested conditions, the surface area occupied

by stress granules per cell was significantly reduced in cells transfected with the mixture of four siRNA

pools compared to the controls (**Fig 7C**). These experiments validate our silencing approach as an

efficient way to significantly reduce both the number of cells positive for stress granules and the surface

occupied by stress granules per cell. Transfection with the 4 siRNA pools had no effect on viral

infection, when compared to transfection with control siRNAs (**Fig 7D**). RT-qPCR analysis showed that

TIAR, TIA-1, G3BP1 and G3BP2 mRNA abundances were significantly reduced in cells transfected

with the 4 siRNA pools as compared to cells transfected with control siRNAs (**Fig 8 A-D**). Of note,

TIAR and TIA-1 expression was induced three- to four-fold by infection, in an MOI-dependent

response. This may be a consequence of virally-induced stress. Transfection of the four siRNA pools

has no effect on *TNFA, IL6* and *IFNB* mRNA abundances nor on viral RNA yield (**Fig 8 E-H**). Together,

these data show that significant reduction of stress granule numbers and size did not suppress the

cytokine response to Asibi infection. Thus, in our model, stress granules have a limited contribution to

the initiation of the antiviral response.

RIG-I signaling is not involved in stress granule assembly

In Huh7 cells, RIG-I signaling is crucial for cytokine induction and secretion upon Asibi infection

(**Fig 1**). Therefore, we wondered whether RIG-I signaling would participate in stress granule assembly

and/or maintenance. We quantified the proportion of Huh7 and Huh7.5 cells that were positive for stress

granules at 24 and 34 hpi following infection at two MOIs. A similar proportion of Huh7 and Huh7.5

cells were found positive for stress granule markers in all tested conditions (**Fig 9 A-B**). No difference

in the number of cells positive for NS4b was observed between Huh7 and Huh7.5 cells (**Fig 9C**). We

conclude that the absence of RIG-I signaling had no effect on the number of granules that assemble in

response to infection. It is thus unlikely that RIG-I mediates a feed-forward loop to maintain stress

granule assembly.

YFV RNA is excluded from stress granules

Our previous analysis showed that neither dsRNA (**Fig 2A**) nor the viral proteins NS4b and NS1 (**Fig 2B, 3, 4D, 5, 7A** and **9A**) seem to be recruited to stress granules. To further investigate the putative association between stress granules containing RIG-I (**Fig 2A**) and viral RNA, we used an RNA fluorescence *in situ* hybridization (FISH) approach coupled with immunofluorescence and confocal microscopy. The Asibi-specific FISH probe produced no signal in non-infected Huh7 control cells (**Fig 10A**) and a bright punctate signal in the cytoplasm of infected cells (**Fig 10A**), confirming the specificity of the probe [46]. Asibi RNA was never found to be associated with TIAR-foci (**Fig 10A,** enlarged panels). This was unexpected since RIG-I signaling is key to cytokine induction and production (**Fig.1**) and massively concentrate in stress granules in Asibi-infected cells (**Fig 2A**). These results suggest that an interaction between viral RNA and RIG-I might not happen in stress granules in Asibi-infected Huh7 cells. A minute quantity of RIG-I might however interact with viral RNA in ER-derived replication sites and initiates a stress granule-independent IFN response. In line with this, RIG-I was recruited to TIAR-positive granules in Huh7 cells treated with NaArs (**Fig 10B**), which is consistent with previous observations on NaArs-treated HeLa cells [20]. These data further indicate that RIG-I recruitment to stress granules can occur in absence of viral RNA.

Discussion

The identity of the PRR(s) responsible for the initiation of a pro-inflammatory response upon direct infection of human hepatocytes with the wild-type strain of YFV has not been uncovered thus far. In this study, we first identified RIG-I and PKR as main actors responsible for the induction and production of pro-inflammatory cytokines upon infection of human hepatocytes with the Asibi strain of YFV. We demonstrated the crucial involvement of RIG-I in this process by directly comparing the dysfunctional RIG-I-expressing human hepatoma cell line Huh7.5. with its RIG-I -competent counterpart (Huh7 cells). The level of IL6 and TNFα secreted in the first 48 hours of infection by Huh7.5 cells upon Asibi infection was under the limit of detection, whilst both cytokines were produced in Huh7 cells in the same experimental conditions. This is in accordance with our previous studies showing that human plasmacytoid dendritic cells infected with the YFV vaccine strain 17D induced an IFN response in a

RIG-I dependent manner [19]. Furthermore, the increased viral load found in the absence of a functional

RIG-I demonstrates the direct anti-viral role of the effector mechanisms initiated by this sensor. Other

members of the RLR family may contribute to YFV sensing in other cell types or at later time of

infection. Such distinct temporal activity of RIG-I and MDA5 during infection with the related West

Nile Virus has been reported in mice [47].

In Huh7 cells silenced for PKR and infected with Asibi, cytokine expression was, in most of the

tested conditions, below the detection limit. This suggests that PKR, like RIG-I, largely contributes to

the initiation of the antiviral response. This is consistent with the previously reported role of PKR in the

production of IFN and other pro-inflammatory cytokines upon viral infection, possibly *via* its interaction

with the IKK complex and subsequent activation of NF-κB [44,48]. Moreover, PKR was previously

described as an important effector protein against infection with WNV in mice [40].

The fact that abolishing RIG-I signaling or silencing PKR drastically reduces the pro-inflammatory

response in Asibi infected cells shows the non-redundant nature of these two dsRNA sensors in

maintaining cytokine production. We speculate that RIG-I and PKR cooperate to induce an anti-YFV

response. Indeed, PKR interacts with RIG-I and MDA5 in HEK293T cells transfected with FLAG-

tagged versions of RIG-I or MDA5 and infected with a mutant vaccinia virus lacking a gene known to

inhibit PKR [49]. However, in this context, PKR was only required for MDA5, but not RIG-I mediated,

IFN production [49]. Consistent with the hypothesis that PKR and RIG-I cooperate to orchestrate

antiviral response, PKR is known to interact with several members of the TRAF family, including

TRAF2 and TRAF6 [50,51], two proteins involved in MAVS signaling. Finally, MAVS, which acts

downstream of RIG-I and MDA5, interacts with PKR [52] and is essential for PKR-induced IFN

production [49].

Stress granules have been demonstrated to contain and concentrate viral nucleic sensors; hence,

they are generally regarded as a functional platform for ligand recognition and subsequent induction of

an antiviral response. We report here that Asibi replication triggers the formation of stress granules in

human hepatoma cells and epithelial monkey cells. This is consistent with previous studies showing that

infection with the vaccine strain 17D induces the formation of stress granules in human dendritic cells

and hamster kidney cells [53]. Thus, formation of stress granules seems to be a general feature of

flavivirus infection [35,36,54,55]. However, unlike WNV, DENV, ZIKV and the related tick-borne

encephalitis virus [35,36,38,54–57], YFV does not seem to efficiently antagonize stress granule

biogenesis. YFV may not express proteins that inhibit granule formation. Alternatively, stress granules

may not have an antiviral function against YFV and thus the virus has no pressure to evolve an evasion

strategy.

Stress granules were not detected in Huh7 cells depleted of PKR upon 24 hours of Asibi infection,

strongly suggesting a PKR-dependent granule establishment early in infection. Of note, RIG-I signaling

was not involved in stress granule assembly, indicating that, whilst the two sensors are both critically

implicated in the pro-inflammatory cytokine production, only PKR is able to activate stress granule

formation. At 34 hours, very few cells silenced for PKR exhibited stress granules. Small quantities of

remaining PKR, below the detection limit of Western blot and qPCR analysis, may be responsible for

the formation of these foci. Alternatively, they may form subsequent to the phosphorylation of eIF2α

by one of the other three eIF2α kinases. YFV replication and assembly, which takes place within ER

membranes [3], likely causes an ER stress and might thus trigger the activation of the ER resident kinase

PERK. Since GCN2 is phosphorylated in mouse and human dendritic cells upon infection with 17D

[53], this kinase could also potentially contribute to granule formation in Asibi infected cells. This would

explain the effect of the GCN2 inhibitor on stress granule formation in our experimental setting.

Similar to what was reported in the context of infection with numerous viruses [27,28,58], we

showed that RIG-I accumulates in stress granules upon Asibi infection. PKR, MDA5, LGP2 and the

DNA sensor cGAS were also shown to concentrate in stress granules in cells infected with a large panel

of RNA and DNA viruses [20,22–26]. However, we were unable to detect neither Asibi RNA nor

dsRNA structures in stress granules of infected cells. Thus, the interaction between YFV RNA by RIG-

I and PKR is unlikely to occur in stress granules. Similarly, in cells infected with Orthohantaviruses, no

viral RNA was found associated with stress granules [59]. Conversely, viral RNA was detected in stress

granules induced by infection with IAVΔNS1 [20]. We propose that, at least in Asibi-infected cells, the

large accumulation of PKR and RIG-I in stress granules masks the detection of small quantities of PKR

and RIG-I which are recruited to perinuclear ER-derived replication sites, where viral RNA sensing

occurs. Importantly, inhibition of stress granule formation didn't affect the pro-inflammatory cytokine

expression by infected hepatocytes, indicating that the anti-viral response and stress granules formation

are independent events in the early phases of YFV infection. Whether the formation of stress granules

and their composition affect later stages of infection remains to be established.

Collectively, our data strongly suggest that, in cells infected with YFV, pro-inflammatory cytokine

induction is mediated by RIG-I and PKR, independent of their localization in stress granules. Thus,

contrary to the widely believed consensus, we demonstrated that the stress granule formation can be

uncoupled from the initiation of a pro-inflammatory response upon viral infection.

Materials and Methods

Cell lines, virus, antibodies and inhibitors

Huh7 human hepatocellular carcinoma cells (kindly provided by A. Martin, Institut Pasteur Paris)

and Huh7.5 cells [60] were maintained in Dulbecco's modified eagle's medium (DMEM)(Gibco)

containing GlutaMAX I and sodium pyruvate (Invitrogen) supplemented with 10% heat-inactivated

foetal bovine serum (FBS, Dominique Dutscher) and 1% penicillin and streptomycin (P/S, 10.000 U/ml,

Thermofischer). HepG2 cells (kindly provided by C. Neuveut, Institut Pasteur Paris) were maintained

in DMEM-F12 complemented with 10% FBS, 3.5×10^{-7} M hydrocortisone, and 5 µg/ml insulin. Human

embryonic kidney (HEK) 293FT cells (American Type Culture Collection; ATCC CRL-1573), *Macaca*

mulatta mammary gland epithelial CMMT cells (kindly provided by O. Schwartz, Institut Pasteur Paris)

and African green monkey kidney epithelial Vero cells (ATCC) were maintained in DMEM 10% FBS.

The Asibi strain, which is the YFV reference strain was provided by the Biological Resource Center

of Institut Pasteur. The YFV-DAK strain (YFV-Dakar HD1279) was provided by the World Reference

Center for Emerging Viruses and Arboviruses (WRCEVA), through the University of Texas Medical

Branch at Galveston, USA. Virus stocks were propagated on Vero cells. Viruses were concentrated by

polyethylene glycol 6000 precipitation and purified by centrifugation in a discontinued gradient of

sucrose. Infectious virus titers were determined by plaque assays on Vero cells. Cells were infected

with 10-fold serial dilutions of viral stocks and incubated for 4 days in DMEM containing 2% FBS and

carboxymethyl cellulose. Cells were then washed with PBS, fixed with 3% formaldehyde crystal violet solution for 20 min at room temperature, rinsed with water and plaques were counted.   Cell infections were carried out at the indicated MOIs. The viral inoculum was replaced with fresh culture medium two hours post-infection.

The following antibodies were used: pIRF3 S386 (clone EPR2346, Abcam), IRF3 (FL-425, Santa Cruz), NF-κB p65 (D14E12, Cell Signalling), IκBα (L35A5, Cell Signalling), β-Actin (clone AC-74, Sigma-Aldrich), dsRNA J2 (English & Scientific Consulting Kft), G3BP (BD Biosciences), TIAR (C-18, Santa-Cruz), eIF4G (sc-11373, Santa-Cruz), eIF3b (N-20, sc-16377), YFV-NS4b (kind gift from C.M. Rice, Rockefeller University), Alexa Fluor® 488 donkey Anti-Goat IgG (H+L), Alexa Fluor® 594 donkey Anti-Mouse IgG (H+L), Alexa Fluor® 647 donkey Anti-Rabbit IgG (H+L) (Life Technologies), anti-mouse 680 (LI-COR Bioscience), anti-rabbit 800 (Thermo Fisher Scientific). The RIG-I antibody was previously described [20].

BAY11-7085 (Abcam), GSK2606414 (Selleck Chemicals), GCN2-IN-1 (A-92) (Selleck Chemicals) were resuspended in DMSO (Pan Bioteck) and used at the indicated concentration. Sodium arsenite (NaArs, Sigma) and Cycloheximide (CHX, Sigma) were diluted in water.

Generation of shRNA-expressing Huh7 cells by lentiviral transduction

pALPS-shPKR1 and pALPS-shPKR3 were generated by cloning the following sequences in pALPS [61]: (TGC TGT TGA CAG TGA GCG CCA GCA GAT ACA TCA GAG ATA ATA GTG AAG CCA CAG ATG TAT TAT CTC TGA TGT ATC TGC TGA TGC CTA CTG CCT CGG A) and (TGC TGT TGA CAG TGA GCG ACA GAA GAA AAG ACT AAC TGT ATA GTG AAG CCA CAG ATG TAT ACA GTT AGT CTT TTC TTC TGG TGC CTA CTG CCT CGG A). HEK293FT cells were transfected with pCMV-VSV-G codon-optimized, pCMVR8.74 (both plasmids were given by P. Charneau, Institut Pasteur) and lentiviral vectors (pALPS-shPKR1/3 or pALPS-shLuc) at a mass ratio of 1:4:4. Huh7 cells were transduced with supernatant from transfected HEK293FT and selected with 10 μg/μL puromycin (Sigma Aldrich) four days later.

Small Interfering RNA (siRNA) transfection

Huh7 cells were transfected using Lipofectamine RNAiMax (Life Technologies) at 30 nM final of

nontargeting scrambled negative control (siScr) or G3BP1, G3BP2 and TIAR siRNAs. Alternatively,

cells were transfected at 40 nM final of siRNA-TIA1. All siRNAs were obtained from Dharmacon (ON-

TARGETplus SMARTpool). Transfected cells were harvested or used for infection assays 48 hours

after transfection.

Enzyme-linked immunosorbent assay (ELISA)

The amount of TNFα and IL6 in cell-culture supernatants was quantified by ELISA using the Human

TNFα or IL6 ELISA kit (eBiosciences), according to manufacturer's instructions. The detection limits

were 4 pg/ml for TNFα and 2 pg/ml for IL6.

Western blot analysis

Cells were lysed in radioimmunoprecipitation assay (RIPA) buffer (Sigma) containing a protease and

phosphatase inhibitor mixture (Roche). Cell lysates were normalized for protein content, boiled in

NuPAGE LDS sample buffer (Thermo Fisher Scientific) in non-reducing conditions and the proteins

were separated by SDS-PAGE (NuPAGE 4-12% Bis-Tris Gel, Life Technologies). Proteins were

transferred to nitrocellulose membranes (Bio-Rad) using a Trans-Blot Turbo Transfer system (Bio-Rad).

After blocking with PBS-Tween-20 0,1% (PBST) containing 5% milk or BSA for 1 hour at RT, the

membrane was incubated overnight at 4 °C with primary antibodies diluted in a blocking buffer. Finally,

the membrane was incubated for 45 min at RT with secondary antibodies (Anti-rabbit/mouse IgG (H+L)

DyLight 680/800) diluted in a blocking buffer, washed, and scanned using an Odyssey CLx infrared

imaging system (LI-COR Bioscience).

RNA extraction and RT-qPCR analysis

Total RNAs were extracted from cell-lysates using the NucleoSpin RNA II Kit (Macherey-Nagel)

following the manufacturer's protocol and were eluted in nuclease-free water. First-strand

complementary DNA synthesis was performed with the RevertAid H Minus M-MuLV Reverse

Transcriptase (Thermo Fisher Scientific). Quantitative real-time PCR was performed on a real-time PCR

system (QuantStudio 6 Flex, Applied Biosystems) with SYBR Green PCR Master Mix (Life

Technologies). Data were analyzed with the $\Delta\Delta$CT method, with all samples normalized to GAPDH.

All experiments were performed in technical triplicate. Primers used in this study are: GAPDH (GGT

CGG AG TCA ACG GAT TTG and ACT CCA CGA CGT ACT CAG CG), YF-NS3 (GCG TAA GGC

TGG AAA GAG TG and CTT CCT CCC TTC ATC CAC AA), TNFA (GCC CAT GTT GTA GCA

AAC CC and GGA GGT TGA CCT TGG TCT GG), IL6 (TGC AAT AAC CAC CCC TGA CC and

GTG CCC ATG CTA CAT TTG CC), IFNB (AAG CAA TTG TCC AGT CCC A and TGC ATT ACC

TGA AGG CCA AG), PKR (CAG AAT TGA CGG AAA GAC TTA CG and CTC TCA AGA GAA

TCA TCA CTG GT), G3BP1 (CAG TTA TAT AGA CCG GCG GC and TAT GTC CAA ACC TAC

GCG GC), G3BP2 (CAA TCT CAG CCA CCT CGT GT and CCA CAA CGT TTC CAA AAC TCA

T), TIAR (CGT CTG GGT TAA CAG ATC AGC and CAT GGG CTG CAC TTT CAT GG), TIA1

(AAT CCC GTG CAA CAG CAG GA and AGG CGG TTG CAC TCC ATA AT). Viral genome

equivalents concentrations (GE/ml) were determined by extrapolation from a standard curve generated

from serial dilutions of the plasmid encoding the YF-R.luc2A-RP [62].

Immunofluorescence, fluorescence *in situ* hybridization (FISH) and confocal analysis.

Cells were fixed with 4% paraformaldehyde (PFA; Sigma-Aldrich) for 30 min at RT, permeabilized

with 0.5% Triton X-100 in PBS and then blocked with 0,05% Tween-, 5% BSA-containing PBS before

incubation with the indicated primary antibodies. Antibody labelling was revealed with Alexa Fluor®

488-, 594 or 647-conjugated antibodies. Coverslips were mounted on slides using ProLong® Gold

Antifade Reagent with NucBlue solution (Invitrogen). Images were acquired with a Zeiss LSM 700

inverted confocal microscope. Fluorescence in situ hybridization was performed after immunostaining.

The (+) RNA strands of YFV were detected following the manufacturer's protocol (ViewRNA ISH

Cells Assays) using probe sets designed by Affymetrix. The Alexa-Fluor 546-conjugated (+)-strand

probe set targets a region between position nucleotides 4567 and 5539 of the YFV genome. Intensity

plot data were generated using Zen 2 lite (Zeiss) and plotted using GraphPad Prism 6.

Stress granule quantification

Nucleus and stress granule quantifications were performed using CellProfiler (v3.0.0) using custom pipeline (available on request). In a nutshell: (i) nuclei were detected using the NucBlue signal, (ii) cellular border were estimated using TIAR or eiF3b signal, (iii) stress granule borders were estimated using G3BP, TIAR, eIF4G or eIF3b signals. Further analyses were performed using R and R studio [63]. Cells were considered positive for stress granules if they exhibited at least two of them.

Statistical analysis

Data are presented as means ± SD and were analyzed using GraphPad Prism 6. Statistical analysis of percentage values or fold enrichment values were performed on logit or log-transformed values, respectively. Statistical analysis was performed with two tailed unpaired t-test with Welch correction for pairwise comparisons or by one- or two-way analysis of variance (ANOVA) with Tukey's or Dunnet's multiple comparisons test when comparing more than two sets of values. Each experiment was performed at least twice, unless otherwise stated. Statistically significant differences are indicated as follows: *: $p < 0.05$, **: $p < 0.01$ and ***: $p < 0.001$; ns, not significant.

Acknowledgments. We thank C.M Rice for generously providing the anti-YFV-NS4b antibodies; R. Kuhn for the YFV replicon construct, M. Flamand for anti-NS1 antibodies (17A12), C. Neuveut for the HepG2 cells, O. Schwartz for the CMMT cells, A. Martin for Huh7 cells, P. Schnupf for eIF3b and eIF4G antibodies, P. Charneau for the plasmids pCMV-VSV-G and pCMVR8.74, A. Cimarelli for the pALPS plasmid, Olivier Disson for immuno-histochemistry advice and Léa Richard for cloning shPKR into pALPS. We are grateful to the members of our laboratory for helpful discussions and technical advice. We thank Audrey Salles and Julien Fernandes (UTechS Photonic Bioimaging platform, Institut Pasteur) for their help in confocal acquisition. We are grateful to Stéphane Rigaud (Image Analysis Hub, Institut Pasteur) for his advice in analyzing confocal images and to Vincent Guillemot (Bioinformatics and Biostatistics HUB, Institut Pasteur) for his help in statistical analysis. Finally, we thank Stephanie Straubel for critical reading of the manuscript.

References

1.   Gould EA, Solomon T. Pathogenic flaviviruses. Lancet. 2008;371: 500–9.
doi:10.1016/S0140-6736(08)60238-X
2.   Gubler DJ, Kuno G, Markoff L. Flaviviruses. 5th Edition. Fields Virology. 5th
Edition. 2007. pp. 1153–1252.
3.   Douam F, Ploss A. Yellow Fever Virus: Knowledge Gaps Impeding the Fight Against
an Old Foe. Trends Microbiol. 2018;26: 913–928. doi:10.1016/j.tim.2018.05.012
4.   Pulendran B. Learning immunology from the yellow fever vaccine: innate immunity to
systems vaccinology. Nat Rev Immunol. 2009;9: 741–7. doi:nri2629 [pii] 10.1038/nri2629
5.   Barrett AD. Yellow Fever in Angola and Beyond--The Problem of Vaccine Supply
and Demand. N Engl J Med. 2016;375: 301–3. doi:10.1056/NEJMp1606997
6.   Barrett ADT. The reemergence of yellow fever. Science. 2018;361: 847–848.
doi:10.1126/science.aau8225
7.   Messaoudi I, Basler CF. Immunological features underlying viral hemorrhagic fevers.
Curr Opin Immunol. 2015;36: 38–46. doi:10.1016/j.coi.2015.06.003
8.   Paessler S, Walker DH. Pathogenesis of the viral hemorrhagic fevers. Annu Rev
Pathol. 2013;8: 411–40. doi:10.1146/annurev-pathol-020712-164041
9.   Chen IY, Ichinohe T. Response of host inflammasomes to viral infection. Trends
Microbiol. 2015;23: 55–63. doi:10.1016/j.tim.2014.09.007
10.   ter Meulen J, Sakho M, Koulemou K, Magassouba N, Bah A, Preiser W, et al.
Activation of the cytokine network and unfavorable outcome in patients with yellow fever.
The Journal of infectious diseases. 2004;190: 1821–7. doi:10.1086/425016
11.   Engelmann F, Josset L, Girke T, Park B, Barron A, Dewane J, et al. Pathophysiologic
and transcriptomic analyses of viscerotropic yellow fever in a rhesus macaque model. PLoS
Negl Trop Dis. 2014;8: e3295. doi:10.1371/journal.pntd.0003295
12.   Streicher F, Jouvenet N. Stimulation of Innate Immunity by Host and Viral RNAs.
Trends in Immunology. 2019;40: 1134–1148. doi:10.1016/j.it.2019.10.009
13.   Chow KT, Gale M, Loo Y-M. RIG-I and Other RNA Sensors in Antiviral Immunity.
Annual Review of Immunology. 2018;36: 667–694. doi:10.1146/annurev-immunol-042617-
053309
14.   Nan Y, Nan G, Zhang YJ. Interferon induction by RNA viruses and antagonism by
viral pathogens. Viruses. 2014;6: 4999–5027. doi:10.3390/v6124999
15.   Chazal M, Beauclair G, Gracias S, Najburg V, Simon-Loriere E, Tangy F, et al. RIG-I
Recognizes the 5' Region of Dengue and Zika Virus Genomes. Cell Rep. 2018;24: 320–328.
doi:10.1016/j.celrep.2018.06.047
16.   Sprokholt JK, Kaptein TM, van Hamme JL, Overmars RJ, Gringhuis SI, Geijtenbeek
TBH. RIG-I-like Receptor Triggering by Dengue Virus Drives Dendritic Cell Immune
Activation and TH1 Differentiation. J Immunol. 2017;198: 4764–4771.
doi:10.4049/jimmunol.1602121
17.   Sprokholt JK, Kaptein TM, van Hamme JL, Overmars RJ, Gringhuis SI, Geijtenbeek
TBH. RIG-I-like receptor activation by dengue virus drives follicular T helper cell formation
and antibody production. PLoS Pathog. 2017;13: e1006738.
doi:10.1371/journal.ppat.1006738
18.   Hertzog J, Junior AGD, Rigby RE, Donald CL, Mayer A, Sezgin E, et al. Infection
with a Brazilian isolate of Zika virus generates RIG-I stimulatory RNA and the viral NS5
protein blocks type I IFN induction and signaling. European Journal of Immunology.
2018;48: 1120–1136. doi:10.1002/eji.201847483
19.   Bruni D, Chazal M, Sinigaglia L, Chauveau L, Schwartz O, Despres P, et al. Viral

entry route determines how human plasmacytoid dendritic cells produce type I interferons.
Sci Signal. 2015;8: ra25. doi:10.1126/scisignal.aaa1552

20.   Onomoto K, Jogi M, Yoo J-S, Narita R, Morimoto S, Takemura A, et al. Critical Role
of an Antiviral Stress Granule Containing RIG-I and PKR in Viral Detection and Innate
Immunity. PLOS ONE. 2012;7: e43031. doi:10.1371/journal.pone.0043031

21.   Oh S-W, Onomoto K, Wakimoto M, Onoguchi K, Ishidate F, Fujiwara T, et al.
Leader-Containing Uncapped Viral Transcript Activates RIG-I in Antiviral Stress Granules.
PLOS Pathogens. 2016;12: e1005444. doi:10.1371/journal.ppat.1005444

22.   Yoo J-S, Takahasi K, Ng CS, Ouda R, Onomoto K, Yoneyama M, et al. DHX36
enhances RIG-I signaling by facilitating PKR-mediated antiviral stress granule formation.
PLoS Pathog. 2014;10: e1004012. doi:10.1371/journal.ppat.1004012

23.   Sánchez-Aparicio MT, Ayllón J, Leo-Macias A, Wolff T, García-Sastre A. Subcellular
Localizations of RIG-I, TRIM25, and MAVS Complexes. J Virol. 2017;91.
doi:10.1128/JVI.01155-16

24.   Langereis MA, Feng Q, Kuppeveld FJ van. MDA5 Localizes to Stress Granules, but
This Localization Is Not Required for the Induction of Type I Interferon. Journal of Virology.
2013;87: 6314–6325. doi:10.1128/JVI.03213-12

25.   Narita R, Takahasi K, Murakami E, Hirano E, Yamamoto SP, Yoneyama M, et al. A
Novel Function of Human Pumilio Proteins in Cytoplasmic Sensing of Viral Infection. PLOS
Pathogens. 2014;10: e1004417. doi:10.1371/journal.ppat.1004417

26.   Hu S, Sun H, Yin L, Li J, Mei S, Xu F, et al. PKR-dependent cytosolic cGAS foci are
necessary for intracellular DNA sensing. Sci Signal. 2019;12. doi:10.1126/scisignal.aav7934

27.   Onomoto K, Yoneyama M, Fung G, Kato H, Fujita T. Antiviral innate immunity and
stress granule responses. Trends Immunol. 2014;35: 420–428. doi:10.1016/j.it.2014.07.006

28.   McCormick C, Khaperskyy DA. Translation inhibition and stress granules in the
antiviral immune response. Nat Rev Immunol. 2017;17: 647–660. doi:10.1038/nri.2017.63

29.   Ivanov P, Kedersha N, Anderson P. Stress Granules and Processing Bodies in
Translational Control. Cold Spring Harb Perspect Biol. 2019;11: a032813.
doi:10.1101/cshperspect.a032813

30.   Reineke LC, Lloyd RE. Diversion of stress granules and P-bodies during viral
infection. Virology. 2013;436: 255–267. doi:10.1016/j.virol.2012.11.017

31.   Sumpter R, Loo YM, Foy E, Li K, Yoneyama M, Fujita T, et al. Regulating
intracellular antiviral defense and permissiveness to hepatitis C virus RNA replication
through a cellular RNA helicase, RIG-I. J Virol. 2005;79: 2689–99.
doi:10.1128/JVI.79.5.2689-2699.2005

32.   Pierce JW, Schoenleber R, Jesmok G, Best J, Moore SA, Collins T, et al. Novel
inhibitors of cytokine-induced IkappaBalpha phosphorylation and endothelial cell adhesion
molecule expression show anti-inflammatory effects in vivo. J Biol Chem. 1997;272: 21096–
103.

33.   Ivanov P, Kedersha N, Anderson P. Stress Granules and Processing Bodies in
Translational Control. [cited 17 Jan 2020]. doi:10.1101/cshperspect.a032813

34.   Kedersha N, Cho MR, Li W, Yacono PW, Chen S, Gilks N, et al. Dynamic Shuttling
of Tia-1 Accompanies the Recruitment of mRNA to Mammalian Stress Granules. J Cell Biol.
2000;151: 1257–1268. doi:10.1083/jcb.151.6.1257

35.   Bonenfant G, Williams N, Netzband R, Schwarz MC, Evans MJ, Pager CT. Zika
Virus Subverts Stress Granules To Promote and Restrict Viral Gene Expression. Journal of
Virology. 2019;93. doi:10.1128/JVI.00520-19

36.   Emara MM, Brinton MA. Interaction of TIA-1/TIAR with West Nile and dengue virus
products in infected cells interferes with stress granule formation and processing body
assembly. Proceedings of the National Academy of Sciences. 2007;104: 9041–9046.

doi:10.1073/pnas.0703348104

37.   Monath TP. Yellow fever vaccine. Expert review of vaccines. 2005;4: 553–74.
doi:10.1586/14760584.4.4.553

38.   Hou S, Kumar A, Xu Z, Airo AM, Stryapunina I, Wong CP, et al. Zika Virus Hijacks
Stress Granule Proteins and Modulates the Host Stress Response. Journal of Virology.
2017;91. doi:10.1128/JVI.00474-17

39.   Tu Y-C, Yu C-Y, Liang J-J, Lin E, Liao C-L, Lin Y-L. Blocking Double-Stranded
RNA-Activated Protein Kinase PKR by Japanese Encephalitis Virus Nonstructural Protein
2A. Journal of Virology. 2012;86: 10347–10358. doi:10.1128/JVI.00525-12

40.   Samuel MA, Whitby K, Keller BC, Marri A, Barchet W, Williams BRG, et al. PKR
and RNase L Contribute to Protection against Lethal West Nile Virus Infection by Controlling
Early Viral Spread in the Periphery and Replication in Neurons. J Virol. 2006;80: 7009–7019.
doi:10.1128/JVI.00489-06

41.   Dabo S, Meurs EF. dsRNA-Dependent Protein Kinase PKR and its Role in Stress,
Signaling and HCV Infection. Viruses. 2012;4: 2598–2635. doi:10.3390/v4112598

42.   Lu L, Han AP, Chen JJ. Translation initiation control by heme-regulated eukaryotic
initiation factor 2alpha kinase in erythroid cells under cytoplasmic stresses. Mol Cell Biol.
2001;21: 7971–7980. doi:10.1128/MCB.21.23.7971-7980.2001

43.   Axten JM, Medina JR, Feng Y, Shu A, Romeril SP, Grant SW, et al. Discovery of 7-
methyl-5-(1-{[3-(trifluoromethyl)phenyl]acetyl}-2,3-dihydro-1H-indol-5-yl)-7H-pyrrolo[2,3-
d]pyrimidin-4-amine (GSK2606414), a potent and selective first-in-class inhibitor of protein
kinase R (PKR)-like endoplasmic reticulum kinase (PERK). J Med Chem. 2012;55: 7193–
7207. doi:10.1021/jm300713s

44.   Bonnet MC, Weil R, Dam E, Hovanessian AG, Meurs EF. PKR stimulates NF-kappaB
irrespective of its kinase function by interacting with the IkappaB kinase complex. Mol Cell
Biol. 2000;20: 4532–4542. doi:10.1128/mcb.20.13.4532-4542.2000

45.   Bonnet MC, Daurat C, Ottone C, Meurs EF. The N-terminus of PKR is responsible for
the activation of the NF-κB signaling pathway by interacting with the IKK complex. Cellular
Signalling. 2006;18: 1865–1875. doi:10.1016/j.cellsig.2006.02.010

46.   Sinigaglia L, Gracias S, Decembre E, Fritz M, Bruni D, Smith N, et al. Immature
particles and capsid-free viral RNA produced by Yellow fever virus-infected cells stimulate
plasmacytoid dendritic cells to secrete interferons. Sci Rep. 2018;8: 10889.
doi:10.1038/s41598-018-29235-7

47.   Errett JS, Suthar MS, McMillan A, Diamond MS, Gale M. The essential,
nonredundant roles of RIG-I and MDA5 in detecting and controlling West Nile virus
infection. J Virol. 2013;87: 11416–25. doi:10.1128/JVI.01488-13

48.   Zamanian-Daryoush M, Mogensen TH, DiDonato JA, Williams BR. NF-kappaB
activation by double-stranded-RNA-activated protein kinase (PKR) is mediated through NF-
kappaB-inducing kinase and IkappaB kinase. Mol Cell Biol. 2000;20: 1278–1290.
doi:10.1128/mcb.20.4.1278-1290.2000

49.   Pham AM, Santa Maria FG, Lahiri T, Friedman E, Marié IJ, Levy DE. PKR
Transduces MDA5-Dependent Signals for Type I IFN Induction. PLoS Pathog. 2016;12.
doi:10.1371/journal.ppat.1005489

50.   Gil J, García MA, Gomez-Puertas P, Guerra S, Rullas J, Nakano H, et al. TRAF
Family Proteins Link PKR with NF-κB Activation. Molecular and Cellular Biology. 2004;24:
4502–4512. doi:10.1128/MCB.24.10.4502-4512.2004

51.   Arnaud N, Dabo S, Akazawa D, Fukasawa M, Shinkai-Ouchi F, Hugon J, et al.
Hepatitis C virus reveals a novel early control in acute immune response. PLoS Pathog.
2011;7: e1002289. doi:10.1371/journal.ppat.1002289 PPATHOGENS-D-11-00682 [pii]

52.   Zhang P, Li Y, Xia J, He J, Pu J, Xie J, et al. IPS-1 plays an essential role in dsRNA-

induced stress granule formation by interacting with PKR and promoting its activation. J Cell
Sci. 2014;127: 2471–2482. doi:10.1242/jcs.139626
53.   Ravindran R, Khan N, Nakaya HI, Li S, Loebbermann J, Maddur MS, et al. Vaccine
activation of the nutrient sensor GCN2 in dendritic cells enhances antigen presentation.
Science. 2014;343: 313–317. doi:10.1126/science.1246829
54.   Albornoz A, Carletti T, Corazza G, Marcello A. The stress granule component TIA-1
binds tick-borne encephalitis virus RNA and is recruited to perinuclear sites of viral
replication to inhibit viral translation. J Virol. 2014;88: 6611–6622. doi:10.1128/JVI.03736-
13
55.   Roth H, Magg V, Uch F, Mutz P, Klein P, Haneke K, et al. Flavivirus Infection
Uncouples Translation Suppression from Cellular Stress Responses. mBio. 2017;8.
doi:10.1128/mBio.02150-16
56.   Basu M, Courtney SC, Brinton MA. Arsenite-induced stress granule formation is
inhibited by elevated levels of reduced glutathione in West Nile virus-infected cells. PLOS
Pathogens. 2017;13: e1006240. doi:10.1371/journal.ppat.1006240
57.   Amorim R, Temzi A, Griffin BD, Mouland AJ. Zika virus inhibits eIF2α-dependent
stress granule assembly. PLOS Neglected Tropical Diseases. 2017;11: e0005775.
doi:10.1371/journal.pntd.0005775
58.   Zhang Q, Sharma NR, Zheng Z-M, Chen M. Viral Regulation of RNA Granules in
Infected Cells. Virol Sin. 2019;34: 175–191. doi:10.1007/s12250-019-00122-3
59.   Christ W, Tynell J, Klingström J. Puumala and Andes Orthohantaviruses Cause
Transient Protein Kinase R-Dependent Formation of Stress Granules. J Virol. 2020;94.
doi:10.1128/JVI.01168-19
60.   Blight KJ, McKeating JA, Rice CM. Highly permissive cell lines for subgenomic and
genomic hepatitis C virus RNA replication. J Virol. 2002;76: 13001–14.
61.   Pertel T, Hausmann S, Morger D, Zuger S, Guerra J, Lascano J, et al. TRIM5 is an
innate immune sensor for the retrovirus capsid lattice. Nature. 2011;472: 361–5.
doi:10.1038/nature09976
62.   Jones CT, Patkar CG, Kuhn RJ. Construction and applications of yellow fever virus
replicons. Virology. 2005;331: 247–59. doi:10.1016/j.virol.2004.10.034
63.   RStudio Team. RStudio: Integrated Development Environment for R. 2016.
733
734

 Figure legends

Fig 1: The Asibi strain of YFV triggers the production of pro-inflammatory cytokines in hepatoma

cells through RIG-I, IRF3 and NF-κB

Huh7 and Huh7.5 cells were left uninfected or infected with Asibi at an MOI of 1 for 24 or 48 hours in

the absence or presence of 10 µM of BAY11-7085 (BAY). The relative amounts of *IL6* (**A**), *TNFA* (**B**)

and *IFNB* (**C**) mRNAs were determined by RT-qPCR analysis and normalized to GAPDH mRNA and

non-infected samples. The relative amounts of cell-associated viral RNA were determined by RT-qPCR

analysis (**D**) and expressed as genome equivalents (YFV-GE) per µg of total cellular RNA. The data are

the means ± SD of two (Huh7.5) or three (Huh7, Huh7+BAY) independent experiments. ns: non-

significant, *: p<0.05, **: p<0.01. (**E**, **F**) Culture media of the indicated cells were analysed by ELISA

to determine the amounts of secreted IL6 (**E**) and TNFα (**F**). The data are the means ± SD of two

(Huh7.5) or three (Huh7, Huh7+BAY). The dashed lines indicate the limit of detection of the IL6 and

TNFα ELISA (2 and 4 pg/ml, respectively). (**G**) Whole-cell lysates of the indicated cells were analysed

by Western blotting with antibodies against the indicated proteins. (**H**, **I**) Huh7 cells were left uninfected

(NI) or infected with YFV-Asibi for 48 hours at an MOI of 20. Cells were then stained with IRF3 (**H**)

or p65 (**I**) (red), dsRNA (green) antibodies and NucBlue® (blue). Around 200 cells per independent

experiment were scored for nuclear translocation of IRF3 or p65 (**H** and **I**, respectively). ns: non-

significant, *: p<0.05.

Fig 2: RIG-I localizes to stress granules in Huh7 cells infected with YFV

(**A**) Huh7 cells were left uninfected (NI) or infected with Asibi at MOI of 20 for 24 hours. Cells were

then stained with dsRNA (red), RIG-I (purple), TIAR (green) antibodies and NucBlue® (blue). (**B**)

Huh7 cells were left uninfected (NI), treated with 0.5 mM NaArs for 30 min or infected with Asibi at

an MOI of 20 for the indicated time. Cells were then stained with NS4b (purple), G3BP (red), TIAR

(green) antibodies and NucBlue® (blue). Percentage of NS4b-positive and G3BP-positive cells were

estimated and shown in purple and yellow respectively. (**C**) Huh7 cells were treated with 0.5 mM NaArs

for 30 min or infected with YFV-Asibi at MOI of 20 for 48 hours. Cells were then left untreated (NT)

or treated with 100 μg/mL Cycloheximide (CHX) for 1 hour and stained with antibodies against NS4b

(purple), TIAR (green) and G3BP (red), as well as with NucBlue® (blue).

Fig 3: YFV infection induces the formation of stress granules in several mammalian cells

HepG2 or CMMT cells were infected with Asibi at MOI of 20 for, respectively, 48 and 24 hours. Huh7

cells were infected with YFV-HD Dakar at MOI of 10 for 24 hours. Cells were then stained with NS4b

(purple), TIAR (green), G3BP (red) antibodies and NucBlue® (blue).

Fig 4: PKR contributes to stress granule formation in Asibi-infected Huh7 cells

(**A**) Huh7 cells were left uninfected (NI) or infected with Asibi at an MOI of 20 for 24 or 48 hours.

Whole-cell lysates were analysed by Western blotting with antibodies against the indicated proteins.

Samples were loaded on the same gel. *: unspecific bands. (**B**) Whole-cell lysates of Huh7-shLuc and

Huh7-shPKR cells were analysed by Western blotting with antibodies against the indicated proteins.

(**C**) Huh7-shLuc and Huh7-shPKR cells were treated with 0.5 mM NaArs for 30 min and stained with

TIAR (green) and G3BP (red) antibodies, as well as with NucBlue® (blue). (**D**) Huh7 cells were left

uninfected (NI) or infected with YFV-Asibi at MOI of 2 or 20 for 24 or 34 hours. Cells were then stained

with NS4b (purple), TIAR (green), G3BP (red) antibodies and NucBlue® (blue). Images are

representative of two independent experiments. Percentage of G3BP-positive (**E**) or NS4b-positive cells

(**F**) were estimated by analysing at least 200 cells per condition. ns: non-significant, *: $p<0.05$, **:

$p<0.01$, ***: $p<0.001$.

Fig 5: PERK and GCN2 contribute to stress granule formation in Asibi-infected cells

Huh7 cells were left uninfected (NI) or infected with Asibi at an MOI of 20 for 24 hours. Cells were

then treated for 4 hours with either DMSO, 10 μM GSK2606414 (GSK) or 10 μM GCN2-IN-1 (A-92)

(iGCN2) and stained with NS4b (purple), TIAR (green), G3BP (red) antibodies and NucBlue® (blue).

Fig 6: PKR contributes to innate response in Asibi-infected Huh7 cells

Huh7-shLuc or -shPKR cells were left uninfected (NI) or infected with Asibi at MOI of 2 or 20 for 24 or 34 hours. The relative amounts of (A) *PKR* mRNA, (B) YFV RNA, (C) *IL6* mRNA, (D) *TNFA* mRNA and (E) *IFNB* mRNA were determined by qPCR analysis and normalized to that of GAPDH mRNA and non-infected shLuc samples. Mean ± SD is represented. One-way ANOVA on log transformed data with Tukey corrected multiple comparisons tests were performed. shPKR samples were compared to shLuc ones. ns: non-significant, *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$.

Fig 7: Knock-down of G3BP1, G3BP2, TIAR and TIA1 inhibits stress granules formation

(**A**) Huh7 cells were transfected with scramble siRNA (siScr) or with siRNA targeting G3BP1, G3BP2, TIAR and TIA1 (siSG) 48 and 24 hours prior to Asibi infection. Transfected cells were left uninfected (NI), treated with 0.5 mM NaArs for 30 min (NaArs) or infected with Asibi at MOI of 2 or 20 for 24 or 34 hours. Cells were then stained with NS1 (purple), eIF3b (green), eIF4G (red) antibodies and NucBlue® (blue). Images are representative of two independent confocal microscopy analyses. Percentages of eIF4G- and eIF3b-positive (**B**) or NS1-positive cells (**C**) were estimated. (**D**) Stress granule area was determined for each cell as the sum of the whole area occupied by stress granules divided by the total cell surface. (**B-D**) Around 200 cells per independent experiment were scored. ns: non-significant, *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$.

Fig 8: Inhibition of stress granule formation has limited effect on antiviral response

Huh7 cells were transfected with scramble siRNA (siScr) or with four pools of siRNA targeting G3BP1, G3BP2, TIAR and TIA1 (siSG) 48 and 24 hours prior to infection. Transfected cells were left uninfected (NI) or infected with Asibi at an MOI of 2 or 20. Cells were harvested prior to infection (0 h), 24 or 34 hours post-infection. The relative amount of (A) *G3BP1* mRNA, (B) *G3BP2* mRNA, (C) *TIAR* mRNA, (D) *TIA1* mRNA, (E) YFV RNA, (F) *IL6* mRNA, (G) *TNFA* mRNA and (H) *IFNB* mRNA was determined by qPCR analysis and normalized to that of GAPDH mRNA and non-infected sh-Scr samples. Mean ± SD is represented. One-way ANOVA on log transformed data with Tukey corrected multiple comparisons test were performed. Samples treated with siSG-RNA were compared to siScr samples. ns: non-significant, *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$.

Fig 9: RIG-I signaling is not involved in stress granule assembly

(**A**) Huh7 and Huh7.5 cells were left uninfected (NI), treated with 0.5 mM NaArs for 30 min (NaArs) or infected with Asibi at MOI of 2 or 20 for 24 or 34 hours. Cells were then stained with NS4b (purple), TIAR (green), G3BP (red) antibodies and NucBlue® (blue). (**B-C**) Images are representative of two independent confocal microscopy analyses. Around 200 cells per independent experiment were scored. Percentages of G3BP-positive (**B**) or NS4b-positive cells (**C**) were estimated. ns: non-significant, *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$.

Fig 10: YFN RNA is excluded form stress granules

(**A**) Huh7 cells were left uninfected (NI) or infected with Asibi at MOI of 20 for 24 hours. Cells were then stained with TIAR (green) and NucBlue® (blue), prior to be processed for FISH using a probe specific for YFV (+) strand RNA (red). (**B**) Huh7 cells were left untreated (NT) or treated with 0.5 mM NaArs for 30 min. Cells were then stained with RIG-I (purple), TIAR (green) antibodies and NucBlue® (blue).

A *PKR mRNA*

B *YFV RNA*

C *IL6 mRNA*

D *TNFA mRNA*

E *IFNB mRNA*