Mapping the immunogenic landscape of near-native HIV-1 envelope trimers in non-human

primates

Christopher A. Cottrell[1,2,3], Jelle van Schooten[4], Charles A. Bowman[1], Meng Yuan[1], David Oyen[1],

Mia Shin[1], Robert Morpurgo[4], Patricia van der Woude[4], Marielle van Breemen[4], Jonathan L.

Torres[1], Raj Patel[1], Justin Gross[1], Leigh M. Sewall[1], Jeffrey Copps[1], Gabriel Ozorowski[1,2,3], Bartek

Nogal[1,2,3], Devin Sok[2,3,5], Eva G. Rakasz[6], Celia Labranche[7], Vladimir Vigdorovich[12], Scott

Christley[13], Diane G. Carnathan[3,14], D. Noah Sather[12], David Montefiori[7,8], Guido Silvestri[3,14],

Dennis R. Burton[2,3,5,9], John P. Moore[10], Ian A. Wilson[1,2,3,11], Rogier W. Sanders[1,10,#], Andrew B.

Ward[1,2,3,#], Marit J. van Gils[4,#]

[1]Department of Integrative Structural and Computational Biology, The Scripps Research Institute,

La Jolla, CA 92037, USA.

[2]IAVI Neutralizing Antibody Center, The Scripps Research Institute, La Jolla, CA 92037, USA.

[3]Consortium for HIV/AIDS Vaccine Development, The Scripps Research Institute, La Jolla, CA

92037, USA.

[4]Department of Medical Microbiology, Amsterdam UMC, University of Amsterdam, 1105 AZ

Amsterdam, The Netherlands.

[5]Department of Immunology and Microbiology, The Scripps Research Institute, La Jolla, CA

92037, USA.

[6]Wisconsin National Primate Research Center, University of Wisconsin, Madison, WI 53715, USA.

[7]Department of Surgery, Duke University Medical Center, Durham, NC 27710, USA.

[8]Duke Human Vaccine Institute, Duke University Medical Center, Durham, NC 27710, USA.

[9]The Ragon Institute of Massachusetts General Hospital, Massachusetts Institute of Technology

and Harvard University, Cambridge, MA 02139, USA.

[10]Department of Microbiology and Immunology, Weill Medical College of Cornell University, New

York, NY 10021,USA

[11]The Skaggs Institute for Chemical Biology, The Scripps Institute, La Jolla, CA 92037, USA

[12]Center for Global Infectious Disease Research, Seattle Children's Research Institute, Seattle, WA

98109, USA.

[13]Department of Population and Data Sciences, UT Southwestern Medical Center, Dallas, TX

75390, USA

[14]Yerkes National Primate Research Center, Emory University, Atlanta, GA 30322, USA;

[#]corresponding author; m.j.vangils@amc.uva.nl; andrew@scripps.edu; r.w.sanders@amc.uva.nl

Abstract

The induction of broad and potent immunity by vaccines is the key focus of research efforts

aimed at protecting against HIV-1 infection. Soluble native-like HIV-1 envelope glycoproteins

have shown promise as vaccine candidates as they can induce potent autologous neutralizing

responses in rabbits and non-human primates. In this study, monoclonal antibodies were isolated

and characterized from rhesus macaques immunized with the BG505 SOSIP.664 trimer to better

understand vaccine-induced antibody responses. Our studies reveal a diverse landscape of

antibodies recognizing immunodominant strain-specific epitopes and non-neutralizing neo-

epitopes. Additionally, we isolated a subset of mAbs against an epitope cluster at the gp120-gp41

interface that recognize the highly conserved fusion peptide and the glycan at position 88 and

have characteristics akin to several human-derived broadly neutralizing antibodies.

Introduction

HIV-1 continues to cause significant morbidity and mortality around the world with an estimated

1.7 million new infections in 2018[1], which emphasizes the need for an effective prophylactic

vaccine. The HIV-1 envelope glycoprotein (Env) is the sole target for neutralizing antibody (NAb)

responses. Studies of infected patients have led to the isolation of NAbs against multiple different

epitopes on the Env surface that are capable of both neutralizing most circulating strains and

providing passive protection against repeated viral challenges in non-human primates (NHPs)[2-5].

Extensive research efforts, including structure-based engineering of Env immunogens, are

currently directed towards developing vaccine strategies to successfully elicit broadly

neutralizing antibodies (bNAbs) against specific Env epitopes[6-12]. The development and structural

determination of soluble native-like Env trimer mimics, particularly ones based on the SOSIP

technology, has provided a platform for structure-based immunogen design[13-16]. NAbs induced

by SOSIP trimers in NHPs can protect against challenge with an autologous Simian-Human

Immunodeficiency virus (SHIV)[17,18]. However, as NAbs with the required breadth of activity have

not yet been induced in trimer-immunized animals, improvements to current vaccine design and

delivery strategies are clearly needed.

Characterizing the antibody response to SOSIP trimers may provide useful information for

guiding immunogen design. Initial analyses of rabbits immunized with BG505 SOSIP.664 trimers,

including studies of isolated monoclonal antibodies (mAbs), showed that autologous NAbs

targeted a large hole in the glycan shield of the BG505 virus caused by the absence of

glycosylation sites at positions 241 and 289[19,20]. The 241 glycan is highly conserved (97%) among

HIV-1 strains and while the 289 glycan is less conserved, it is still present in 79% of viruses. The

required absence of typically conserved glycans explains why the NAbs isolated from the trimer-

immunized rabbits lack breadth[19]. Later studies involving BG505 trimer-immunized rabbits,

guinea pigs, and NHPs have identified additional narrow-specificity neutralizing serum responses

that recognize epitopes in the C3/V4, C3/V5, and V1 regions, with mAbs isolated from guinea pigs

targeting the C3/V4 epitope[17,20-23].

Here, we describe a detailed characterization of neutralizing and non-neutralizing mAbs

isolated from two rhesus macaques (RMs) previously immunized with the BG505 SOSIP.664

trimer [24]. We identified multiple mAbs targeting the 289-glycan hole on the BG505 SOSIP.664

trimer or a neo-epitope cluster at the base of the trimer. The most potent NAb isolated targeted

the gp120/gp41 interface at an epitope that significantly overlaps with the epitope of human

bNAb VRC34[25]. Insights from the induction of these NAbs through vaccination can be further used

to develop immunogens and immunization strategies to induce cross-reactive antibody

responses.

Results

Indian origin rhesus macaque BCR germline database

The majority of bNAbs isolated from HIV-infected patients have exceedingly high levels of

somatic hypermutation (SHM)[5,26]. Accurately measuring levels of SHM elicited during

immunization experiments is a critical component to ensuring the elicited antibodies are

acquiring the level of mutations associated with neutralization breadth. To accurately measure

the extent of SHM, the mAb sequences that we obtained from BG505 SOSIP.664 trimer-

immunized RMs required comparison to a germline B-cell receptor (BCR) reference database.

The IMGT reference database for RMs is incomplete and contains a mixture of genes/alleles from

both Chinese and Indian origin animals. Given the high levels of genetic diversity in the

Immunoglobulin (Ig) loci among RMs from different origins[27] and the general use of Indian origin

RMs in most HIV-1 immunization experiments conducted in the United States, we constructed a

germline database containing gene/alleles from only Indian origin RMs. The gene/alleles from

the published database[28] were aligned to the Mmul_8.0.1 Indian origin RM genome assembly

using BLAST[29,30]. Sequences that were not identical to the reference genome were eliminated.

Additional genes/alleles from publicly available Indian origin RM genomic DNA sequencing

datasets[21,31] were added to the database. After duplicates were removed, the resulting new

database contained 189 IGHV, 70 IGHD, 9 IGHJ, 188 IGKV, 5 IGKJ, 147 IGLV, and 13 IGLJ

genes/alleles (Table S1).

Recent advances in BCR repertoire sequencing and analysis have enabled the use of next-

generation sequencing (NGS) datasets for inferring novel genes/alleles[27,32,33]. Using the database

described above as an initial database, we performed IgDiscover analysis[27] on IgM BCR sequences

derived from five Indian origin RMs. Additionally, we performed IgDiscover analysis on previously

obtained Indian origin RM BCR NGS datasets that were downloaded from the NCBI sequence read

archive[27,34] or obtained directly from the study authors[28]. Inferred genes/alleles were kept if they

were detected in more than one animal or if they were identical to RM genes/alleles that were

previously deposited in the NCBI database. We added 113 IGHV, 18 IGKV, and 18 IGLV

genes/alleles to our germline database (http://ward.scripps.edu/gld/) resulting in a total of 302

IGHV, 206 IGKV, and 165 IGLV genes/alleles. This updated database was then converted into a

custom IgBLAST database and subsequently used to analyze our BG505-specific mAbs sequences

with IgBLAST[35].

Antigen-specific mAbs isolated from BG505 SOSIP.664 trimer-immunized RMs

To better understand the immunogenicity of the BG505 SOSIP.664 trimers in previously

immunized Indian origin RMs[24] we selected the two animals (rh1987 and rh2011) with the highest

serum neutralization activity against the autologous BG505.T332N pseudovirus for in-depth mAb

analysis. Peripheral blood mononuclear cells (PBMCs) from the following time points were

selected for BG505 SOSIP.664 trimer-specific IgG-positive single memory B-cell sorting: (i) two

weeks prior to the fourth immunization (week 22), (ii) 1 week after the fourth immunization

(week 25) and (iii) 1 week after the sixth immunization (week 53) (Fig 1A). In total, 25 and 17

mAbs were cloned from RMs rh1987 and rh2011, respectively (Fig 1B).

The BG505-specific mAb sequences were analysed using our germline database and

shown to be evenly distributed between kappa and lambda light chains (KC and LC) usage (Fig

1B). For animal rh1987, 11 KC and 14 LC mAbs were isolated. Their average heavy chain (HC) SHM

(nucleotide level) was 6.4% (range: 2.1%-10.2%) with an average HC complementarity-

determining region 3 (CDR-H3) length of 15 amino acids (aa) (range: 7-23) (Table S2). The rh1987

KC mAbs utilized HC variable genes from the IGHV3 and IGHV4 families and predominantly used

KC V genes from the IGKV1 family (Table S2). All of the rh1987 KC mAbs had a CDR-L3 length of 9

aa and their average KC SHM (nucleotide level) was 4.7% (range: 2.6%-6.0%) (Table S2). A single

clonal family with 4 members (RM19A) was detected among rh1987 KC mAbs with members

isolated from both week 22 and week 25 samples (Table S2). The rh1987 LC mAbs used HC V

genes from the IGHV1, IGHV3 and IGHV4 families and LC V genes mainly from the IGLV2 gene

family (Table S2). The rh1987 LC mAbs had an average CDR-L3 length of 10 aa (range: 9-11) with

an average LC SHM (nucleotide level) of 3.8% (range: 0.9%-10.6%) (Table S2). Two clonal families

(RM19B [2 members] and RM19C [4 members]) were identified among the rh1987 LC mAbs

isolated from weeks 22 and 25 (Table S2).

For animal rh2011, 8 KC and 9 LC mAbs were isolated. Their average HC SHM rate

(nucleotide level) was 6.1% (range: 3.0%-9.1%) and they had an average CDR-H3 length of 17 aa

(range 10-20) (Table S2). Half of the rh2011 KC mAbs belonged to the RM20E clonal family,

isolated from the week 53 sample. The RM20E clonal family utilized the HC V gene IGHV5-

ABI*01_S2502 and the KC V gene LJI.Rh_IGKV2.71 (Table S2). Overall, the rh2011 KC mAbs had

an average KC SHM rate (nucleotide level) of 4.1% (range: 3.0%-5.3%) and a CDR-L3 length of 9

aa (Table S2). Among the rh2011 LC mAbs, two clonal families (RM20A [4 members] and RM20B

[2 members]) were isolated from weeks 22 and 25 samples (Table S2). Overall, the rh2011 mAbs

had an average LC SHM rate (nucleotide level) of 4.4% (range: 2.1%-6.4%) and an average CDR

L3 length of 10.6 aa (range: 9-11) (Table S2).

BG505-specific mAbs recognize multiple Env regions

All 42 mAbs bound to the BG505 SOSIP.664 trimer in ELISA, but only 11 of the 25 mAbs from

rh1987 and 9 of the 17 from rh2011 bound the corresponding gp120 monomer (Fig S1). The mAbs

were tested for neutralization activity against the autologous BG505 clade A Tier 2 virus, its

glycan-611 knockout variant (N611A), and the heterologous SF162 clade B Tier 1A virus. Only a

few mAbs, 4 from rh1987 and 2 from rh2011, neutralized the BG505.T332N pseudovirus but one

of them, RM20F from rh2011, did so potently (Fig 1C). Two mAbs from rh1987 and 4 from rh2011

were able to potently neutralize the N611A-variant despite lacking activity against the autologous

BG505.T332N pseudovirus (Fig 1C and D). None of the 42 mAbs neutralized the easy-to-neutralize

heterologous SF162 virus (Fig 1C and D).

EM-based epitope mapping revealed mAbs isolated from both animals target 4 distinct, but somewhat overlapping epitopes

We used low resolution, negative stain, single particle electron microscopy (EM) to visualize

where a representative subset of the isolated mAbs bound on the surface of the BG505 SOSIP

trimer. The majority (55%) of mAbs isolated were non-neutralizing antibodies that bound to the

base of the BG505 SOSIP trimer (Figs 2A, S2, and Table S2) at a neo-epitope cluster that is

occluded by the viral membrane on HIV-1 virions. Fabs bound to the base of the soluble trimer

via multiple angles of approach and utilized a variety of heavy and light chain genes/alleles to do

so (Fig 2A and Table S2). The extent of SHM in the base-targeting mAbs ranged from 2-10% in the

HC and 1-11% in the light chain (Table S2). Previous studies examining the polyclonal antibody

responses elicited by the BG505 SOSIP trimers in rabbits and RMs have shown that epitopes at

the base of the soluble trimer were targeted in every single animal analyzed [21,23,36]. Taken

together with our new data, it is clear that the base of SOSIP trimers contain an immunodominant

non-neutralizing neo-epitope cluster that is easily targeted by a variety of precursor BCRs in

different species.

A subset of mAbs from both animals bound to an epitope near the N611 glycan (Figs 2B

and S2). These mAbs were not capable of neutralizing the autologous BG505.T332N pseudovirus

but neutralized the BG505 N611A variant (Fig 1C and D). Two additional mAbs isolated from

rh2011 (RM20F and RM20H) bound to an epitope near the fusion peptide (FP) and were capable

of neutralizing both the autologous and N611A BG505 pseudoviruses, but the latter virus more

potently (Figs 2C and S2). MAbs from both animals targeted the 289-glycan hole epitope on

BG505, with some, isolated from rh1987, showing weak neutralization of the autologous

BG505.T332N pseudovirus (Figs 1C and 2D; Table S2). Multiple germline genes/alleles were used

to target the same 289-glycan hole epitope (Table S2).

To further assess the epitopes targeted following immunization with the BG505

SOSIP.664 trimer and verify that we isolated mAbs representative of the full serum antibody

response, we performed electron microscopy polyclonal epitope mapping (EMPEM)[36] using week

28 serum. EMPEM revealed that similar epitopes were targeted in both animals and that the

epitope assignments correlated well with the epitopes ascribed to mAbs generated by antigen-

specific B-cell sorting (Fig 2E versus S2).

A previous analysis of purified serum IgGs from RMs rh1987 and rh2011 identified the

C3/V5 epitope as a major target for neutralization activity[20]; however, none of the mAbs isolated

here targeted the C3/V5 epitope nor were they detected by EMPEM. While low resolution,

negative stain EM provides valuable information on where mAbs bind on the surface of HIV-1

Env, the molecular detail necessary to guide structure-based immunogen design requires high-

resolution structural data obtained by cryoEM and x-ray crystallography. We therefore selected

three Fabs (RM20J, RM20F, and RM20E1) that bound to different epitopes for high-resolution

structural determination.

MAb RM20J binds to the α2 helix of gp120 and exploits a hole in the glycan shield of BG505 at

position 289

We solved a 2.3 Å crystal structure of unliganded RM20J Fab and a 3.9 Å cryoEM structure

of RM20J Fab bound to the BG505 SOSIP Env trimer (Figs 3A and S3; Tables S4 and S5.). Together

these structures revealed the RM20J Fab binds to an epitope on a single gp120 protomer with

982 Å² of buried surface area (BSA). The CDR-H1 and CDR-H2 make contact with the C2 region of

gp120 including residues N289 and T290 (Fig 3B). A glycan at position N289 would directly clash

with both the CDR-H1 and CDR-H2 of RM20J (Fig 3B). CDR-L2 makes contact with the first N-

acetyl glucosamine sugar of the N355 glycan (Fig 3C). Additional contacts are made to the α2

helix of gp120 by RM20J CDR-H3 and CDR-L2 (Fig 3C). When compared to 10A, a previously

characterized 241/289 glycan hole targeting NAb isolated from a BG505 SOSIP.664-immunized

rabbit[19,36], RM20J binds to an epitope biased more towards 289 and away from 241 in the

241/289 glycan hole, revealing subtle differences in the recognition of the epitope (Figs 3D and

3E). Despite binding to the BG505 SOSIP trimer with high affinity (Table S3), RM20J was not able

to neutralize the autologous BG505.T332N pseudovirus (Fig 1C). Although the hypervariable

region of V4 was not resolved in the trimer structure, it lies directly above the RM20J epitope

and contains two additional glycans (N406 and N411) that may affect RM20J binding.

Comparisons between the glycosylation profiles of the BG505 viral Env and the SOSIP.664 trimer

revealed differences in the glycoforms present at positions N355, N406, and N411[37,38] with more

complex glycans being found on the viral Env that could hinder the ability of RM20J to bind on

the surface of the virus and, therefore, render it incapable of neutralization. Several of the mAbs

isolated from rh1987, including the RM19A clonal family, bind to a similar epitope as RM20J (Fig

2D, Fig S2, Table S2) and either fail to neutralize the autologous BG505 virus or do so with weak

potency (Fig 1C and D).

MAb RM20F binds to a quaternary epitope at the gp120/gp41 interface that includes elements

of the fusion peptide and the N88 glycan

For a more detailed view of the mode of RM20F recognition, we solved a 2.2 Å crystal

structure of unliganded RM20F Fab and a 4.3 Å cryoEM structure of RM20F Fab bound to BG505

SOSIP trimer (Figs 4A and S3; Tables S4 and S5). RM20F recognizes an epitope spanning two gp41

protomers and a single gp120 protomer that has 1126 Å² of BSA at the interface. The RM20F LC

contributes 22% of the paratope surface area (250 Å²) and makes contact with the poorly

conserved residues H85 (8.1% prevalence among global strains) and K229 (12.5% prevalence) in

the C1 and C2 regions of gp120 respectively (Fig 4B). The RM20F HC contributes the remaining

78% of the paratope surface area (876 Å²) and uses its 20 residue CDR-H3 to wedge between the

FP of the primary gp41 protomer and the HR2 helix of the adjacent gp41 protomer (Fig 4C).

Additional contacts with the fusion peptide proximal region (FPPR) of the primary gp41 protomer

are made by residues at the tip of CDR-H2 (Fig 4B). The N88 glycan accounts for 18% (198 Å^2) of

the epitope BSA and makes contact with the CDR-H2 and FR-H3 regions of RM20F (Fig 4B). The

lack of connecting density, even at lower contour, between RM20F and the glycans at N611 and

N637 suggests these glycans do not substantially contribute to the epitope. Epitope mapping

using BG505.T332N mutant pseudoviruses showed that knocking out the N611 glycan (N611Q

mutant) substantially enhanced neutralization by RM20F, while knocking out the N637 glycan

(N637Q mutant) had no effect. (Fig 4D). Other virus mutants revealed that neutralization by

RM20F was sensitive to various sequence changes within the epitope, particularly at residues

H85 and E647 (84.3% prevalence) and N88 (N88 glycan knock out) (Fig 4D). The N88 glycan knock

out and the H85A mutation (to a lesser extent) significantly reduced neutralization activity of the

FP-targeting bNAb VRC34, but no effect on the CD4 binding site targeting bNAb VRC01 (Fig 4D).

Introducing the 241 or 289 glycans (S241N and P291T, respectively) modestly reduced the

neutralization activity of RM20F (Fig 4D). In comparison to the FP-targeting bNAbs VRC34 and

ACS202, RM20F lacked neutralization breadth when tested against a panel that included multiple

heterologous viruses (Fig S4) likely due to the dependency on poorly conserved residues.

MAb RM20E1 binds to the fusion peptide and makes contact with two adjacent protomers

253   We solved a 2.3 Å crystal structure of the unliganded RM20E1 Fab and 4.4 Å cryoEM

structure of RM20E1 Fab bound to BG505 SOSIP trimer and Fab PGT122 (Figs 5A and S3; Tables

S4 and S5). RM20E1 binds to an epitope composed of one gp120 and two gp41 protomers with

1178 Å^2 of BSA. Residues 515 to 520 of the FP in the primary gp41 protomer are sandwiched

between CDR-H3, CDR-L1, and CDR-L3 of RM20E1 (Fig 5B). CDR-H3 and FR-H1 make contact with

HR2 in the adjacent gp41 protomer (Fig 5C). Additionally, the FR-H1 makes contact near the

N611-glycan site in the adjacent gp41 protomer, but we observed no connecting density that

could be attributed to the N611-glycan itself (Fig 5C). RM20E1 avoids the N88 glycan but does

interact with residues in the C1 region of gp120, including H85, via its CDR-L1 (Fig 5B). Despite

recognition of the conserved FP, RM20E1 did not neutralize the autologous BG505.T332N

pseudovirus. The antibody did however potently neutralize the N611A glycan KO BG505

pseudovirus (Fig 1C) suggesting the epitope is shielded by the N611 glycan. The epitopes of the

FP-targeting bNAbs VRC34, ACS202, and DFPH-a.15 overlap to a large extent with the epitopes

of RM20E1 and RM20F (Fig 5D), with DFPH-a.15 and VRC34 also neutralizing more potently in

the absence of the N611 glycan[9,25,39,40]. The RM20E1-bound FP conformation is similar to the FP

conformation when bound by the bNAb VRC34 (Fig S5); however, the inability of RM20E1 to

accommodate the N611 glycan likely results in the lack of neutralization of the wild-type virus.

Discussion

A major goal of HIV-1 vaccine research is to elicit bNAbs able to neutralize the large

diversity of circulating HIV-1 strains in humans. However, how to achieve this goal remains a

critical problem. Native-like Env trimers are an important design platform for engineering

immunogens for bNAb induction[7-13,41,42]. BG505 SOSIP trimers were able to induce responses in

immunized RMs that potently neutralized the autologous Tier 2 virus[17,24]. When present at

sufficient titers, those NAbs protected against BG505-SHIV challenge[18]. Evaluating SOSIP trimers

in RMs can yield valuable information because of the close genetic relationship between RMs

and humans. Here, we isolated mAbs from two BG505 SOSIP.664 trimer-immunized RMs to

better understand how the immune system recognizes the trimers and the epitopes associated

with the potent, but limited, HIV-1 Tier 2 neutralization.

By mapping the epitopes of all of the mAbs isolated from the immunized RMs, rather than

focusing only on NAbs, we were able to identify several non-neutralizing and potentially

immunodominant epitopes that would ideally be eliminated in future immunization studies. As

shown previously with mAbs from rabbits immunized with BG505 SOSIP.664 trimers[19], the lack

of glycans at positions 241 and 289 in BG505 creates a large glycan hole which is targeted by

mAbs from both RMs. The mAbs isolated from RMs that target the 241/289 glycan hole are more

biased towards the 289-site compared to the previously characterized rabbit mAbs. This

difference may be attributed to the underlying differences in BCR repertoires between the two

animal models. In addition to the lack of glycans due to missing sequons that encode for

glycosylation, the recombinant BG505 SOSIP.664 trimer may also contain missing glycans even

when the correct sequon is present as previously observed by mass spectrometry studies of

glycopeptides[37,43]. Our study identified gp120/gp41 interface antibodies whose neutralization

was enhanced in the absence of the N611 glycan, suggesting that the BG505 SOSIP.664 trimer

may have sub-stoichiometric glycan occupancy in gp41 at this position, creating an unexpected

but immunogenic glycan hole. The elicitation of FP targeting mAbs in RMs with the BG505

SOSIP.664 trimer provides evidence that the FP bNAb epitope is accessible and immunogenic on

soluble Env trimer immunogens. Recent studies in mice, guinea pigs and RMs using synthetically

produced HIV fusion peptides covalently attached to carrier proteins as priming immunogens

followed by boosts with soluble Env trimer immunogens have also elicited FP specific antibodies

including some mAbs with neutralization breadth[9,44,45]. However, the majority of the animals

immunized in these studies do not develop neutralization breadth and instead develop potent

neutralization against the BG505 pseudovirus with the N611 glycan KO[9,44,45]. Given the

consistency across studies and animal models in eliciting potent NAbs that target the FP epitope

and require the absence of the N611 glycan, investing in strategies to quantify and enhance the

N611 glycan occupancy in soluble Env trimer immunogens, particularly for boost immunogens,

may improve the neutralization breadth elicited by FP-targeting immunization protocols.

The sorting probe used to isolate BG505 Env-specific B-cells was a C-terminally

biotinylated BG505 SOSIP.664 trimer bound to a fluorescent streptavidin tetramer. Steric

constraints between the base of the trimer and the streptavidin tetramer likely resulted in a

lower recovery of base epitope-specific B-cells. Despite this potential selection bias, 55% of the

mAbs isolated from the two immunized RMs bound to the base of the BG505 SOSIP.664 trimer,

indicating the base of the soluble trimer is the major target for antibody responses during

immunization. To reduce the immunogenicity of this epitope, glycans can be introduced to shield

this site or the native-like trimers could be constructed onto scaffolds or particles[46-49].

We were unable to construct individual germline BCR databases from RMs rh1987 and

rh2011 to precisely determine the SHM and gene/alleles usage as additional PBMC samples were

no longer available. Instead, we constructed a germline database containing BCR gene/alleles

from multiple Indian origin RMs that allowed us to measure SHM levels in the mAbs we isolated

from RMs. The new database provides web-based access (http://ward.scripps.edu/gld/) to a

curated and highly annotated general resource for examining BCR gene/alleles from Indian origin

RMs. Previous estimates of average SHM rates in mAb sequences from Env-immunized RMs were

8.9% and 6.1% for the HC and LC, respectively [50]. These apparently high levels of SHM following

repeated immunization with the exact same soluble Env immunogens were likely due to missing

germline gene/alleles from the database used to calculated SHM. Using the germline database

reported in this study, we were able to assign vaccine-elicited antibody sequences to specific

germline genes/alleles and determine levels of SHM much more accurately. The average levels

of SHM reported in this study (6.3% for HC and 4.2% for LC) are comparable to the average levels

of SHM reported in similar immunization studies where per animal germline BCR databases were

inferred using IgDiscover [51,52]. Our germline database provides a resource for assigning germline

genes/alleles and accurately calculating rates of SHM when inferring individual germline

databases for each animal is logistically impractical.

In conclusion, in this study, neutralizing and non-neutralizing mAbs with distinctive

epitopes were isolated and characterized in BG505 SOSIP.664-immunized RMs. We

demonstrated that a polyclonal response was elicited in two different RMs that target the BG505

SOSIP.664 trimer in highly similar ways. While rabbit antibody responses are dominated by the

base and 241-glycan hole epitopes, the RM mAbs target more diverse epitopes with mAbs also

targeting the FP and gp120/gp41 interface. The mAbs characterized here also provide a valuable

resource for epitope mapping and comparison to a wide array of BG505-based immunization

experiments, including the recently initiated BG505 SOSIP.664 human clinical trial

(NCT03699241). Finally, the FP-targeting mAbs in particular provide a structure-guided

opportunity to modify BG505 SOSIP and other trimers to focus the antibody response on the FP-

region, with the goal of eliciting bNAb-like antibodies.

Methods

Immunizations of rhesus macaques

Immunization samples used in this study were obtained from previously immunized RMs

described in Sanders et al. 2015 [24]. Briefly, RMs were immunized intramuscularly (i.m.) with 100

µg of BG505 SOSIP.664 trimer formulated in 75 units of ISCOMATRIX given at week 0, 4, 12, 24,

38 and 52. All immunizations and blood samplings were performed at the Wisconsin National

Primate Research Center as described previously [24].

Rhesus macaque naïve B-cell repertoire sequencing

Frozen PBMCs from five naïve Indian origin RMs were obtained from Yerkes National Primate

Research Center (IACUC approval YER2001036). The cells were rapidly thawed in a 37°C water

bath and immediately diluted into 10 mL of pre-warmed RPMI media with 10% (v/v) heat-

inactivated fetal bovine serum (FBS). Cells were pelleted at 400xg for 7 minutes and resuspended

in 0.5 mL FACS buffer (PBS + 1% (v/v) FBS and stained on ice for 1 hr with the panel of fluorescent

antibodies against IgM ([clone G20-127] BD), CD4 ([clone OKT-4] BioLegend), CD3 ([clone SP34-

2] BD), IgG ([clone G18-145] BD), CD20 ([clone 2H7] BioLegend), CD8 ([clone RPA-T8] BioLegend),

CD14 ([clone M5E2] BD), CD16 ([clone eBioCB16] ThermoFisher) and an eFluor780 viability

marker (ThermoFisher). A MoFlo Astrios cell sorter (Beckman Coulter), with gating for live

$IgM^+/CD20^+/CD3^-/CD4^-/CD8^-/CD14^-/CD16^-$ cells, was used to isolate cells that were then pelleted

at 600xg for 10 min, resuspended in RLT+BME buffer (Qiagen), snap frozen in a dry ice ethanol

bath and stored at -80°C. RNA extraction was performed using RNeasy Protect Mini kit (Qiagen)

following the manufacturer's instructions. The 5' rapid amplification of cDNA ends (5'RACE) with

template switching method was used to obtain cDNA with unique molecular identifiers (UMIs)

using a protocol modified from Turchaninova *et al.* [53]. Briefly, 300 ng of RNA was used in a 5'RACE

cDNA synthesis reaction. The first-strand cDNA synthesis was performed at 42°C for 1 hr using

the RM IgM outer reverse primer (5'-GTGATGGAGTCGGGAAGGAAG-3'), a template switch

adaptor with incorporated UMIs (5'-

AAGCAGUGGUAUCAACGCAGAGUNNNNUNNNNUNNNNUCTTrGrGrGrG-3'), and SMARTScribe

Reverse Transcriptase (Clontech). Residual template switch adaptor was removed by incubation

with 5 U of uracil DNA glycosylase (New England BioLabs) for 40 min at 37°C. The resulting cDNA

was purified using the MinElute PCR Purification Kit (Qiagen) following the manufacturer's

instructions. PCR amplification was performed using the Q5® High-Fidelity DNA Polymerase (New

England BioLabs), the forward primer (5'-NNNNAAGCAGTGGTATCAACGCA-3'), and the RM IgM

inner reverse primer (5'-NNNNNAGGGGGAAAAGGGTTG-3'). Illumina adaptors were added using

the NEBNext® Ultra™ II DNA Library Prep Kit (New England BioLabs) following the manufacturer's

instructions. Libraries were sequenced on an Illumina MiSeq using the Illumina v3, (2x 300 bp)

sequencing kit.

Indian origin RM germline BCR database

Gene/alleles published by Vigdorovich et al. [28] were aligned to the Mmul_8.0.1 Indian origin RM

genome assembly using BLAST [29,30]. Sequences that were not identical to the reference genome

were eliminated. Additional full-length genes/alleles from available Indian origin RM genomic

DNA sequencing datasets [21,31] were added to the database. Duplicates and sequences containing

ambiguous bases were removed. The resulting initial Indian origin RM germline BCR database

(Table S1) was used for running IgDiscover on additional NGS datasets that were obtained during

this study as described above, downloaded from the NCBI SRA [27,34], or obtained directly from the

study authors [28]. Paired sequence reads were aligned and filtered for length and quality using

VDJServer [54]. Novel germline BCR gene/alleles were inferred using IgDiscover v0.11 with the

germline_filter parameters "unique_cdr3s" and "unique_js" set to 10 and 4 respectively to

reduce the rate of false positives [27]. Inferred genes/alleles were kept if they were detected in

more than one animal or if they were identical to RM genes/alleles that were previously

deposited   in   NCBI.   The   resulting   germline   BCR   database   (available   at

http://ward.scripps.edu/gld/) was converted into a custom IgBLAST database and subsequently

used to analyze the BG505-specific mAb sequences [35].

Env sequence analysis

Prevalence of specific amino acids or potential N-linked glycosylation sites (PNGS) were

calculated   using   HIVAnchor   (https://github.com/chazbot72/anchor).   Pairwise   alignments

between the HxB2 Env reference sequence (K03455) and the LANL 2018 Group M super filtered

web alignment was performed using Clustal Omega [55]. The results were parsed into a database

keyed on positions relative to the reference, with gaps notated as sub positions following the last

identical residue. The database was subsequently interrogated for conservation of amino acids

or PNGS at specific positions.

Env protein production

BG505 SOSIP.664, BG505 SOSIP.664-D7324 tag, BG505 SOSIP.664-AviTag, BG505 SOSIP.v4.1, and

BG505 SOSIP.v5.2 were expressed in HEK293F cells and purified with either PGT145 or 2G12

affinity chromatograph followed by size exclusion chromatography (SEC) using a HiLoad® 16/600

Superdex® pg200 (GE Healthcare) as described previously [7,8,13]. Monomeric gp120 proteins

(AviTag or D7324 tagged) were purified using a *Galanthus nivalis* lectin (Vector Labs) column. The

Avi-tagged   proteins   were   biotinylated   using   the   BirA   enzyme   (Avidity)   according   to   the

manufacturer's protocol. The resulting biotinylated proteins are referred to using the descriptor

AviB.

Monoclonal antibody isolation

BG505 SOSIP.664-specific IgG$^+$ memory B-cells from isolated PBMCs from RMs rh1987 and

rh2011 were single cell sorted in lysis buffer in order to amplify the antigen-specific mAbs, as

previously described [56]. PBMCs were stained with primary fluorophore-conjugated antibodies to

human CD3, CD8, CD14, CD20, IgG and IgM (BD Pharmigen). For staining with Env proteins, 50

nM of BG505 SOSIP.664-AviB, BG505 SOSIP.664 7C3-AviB or gp120-AviB were coupled in

equimolar ratios to Streptavidin-PE, Streptavidin-FITC or Streptavidin-APC (Life Technologies),

respectively. Cells were stained for 1 hr at 4°C in PBS supplemented with 1 mM EDTA and 1% FBS.

In the gating strategy, we first excluded unwanted cell populations (CD3$^-$/CD8$^-$/CD14$^-$) followed

by selection of HIV Env-specific (positive for any of the 3 probes) memory B-cells

(CD20$^+$/IgG$^+$/IgM$^-$/HIV$^+$). Cells of interest were single-cell sorted using a BD FACSAria III machine,

into 96-well plates containing lysis buffer, and immediately stored at -80°C. One round of reverse-

transcription and two rounds of nested PCR were performed to amplify the antibody V(D)J genes

as previously described by Tiller *et al.*[15]. The PCR products containing the variable regions of the

heavy chain or light chain, kappa or lambda were cloned into human IgG expression vectors to

produce mAbs as described previously [56]. Fab expression vectors were made by introducing two

stop codons following residue D234 (Kabat numbering [57]) in the IgG heavy chain vectors using

the QuikChange® Lightning Site-Directed Mutagenesis kit (Agilent). Sequences were verified by

Sanger sequencing (Genewiz).

Monoclonal antibody and Fab production

MAbs and Fabs were expressed in HEK293F cells and purified using affinity chromatography.

Briefly, HEK293F cells (Invitrogen) were co-transfected with heavy and light chain plasmids (1:1

ratio) using PEImax. Transfections were performed according to the manufacturer's protocol.

Supernatants were harvested 4-6 days following transfection and passed through a 0.45 μm filter.

MAbs were purified using Protein A/G (ThermoFisher) or MAbSelect™ (GE Healthcare) affinity

chromatography. Fabs were purified using CaptureSelect™ CH1-XL (ThermoFisher) affinity

chromatography.

D7324-capture ELISA for monomeric and trimeric BG505 Env proteins

Binding ELISAs were conducted as described previously [13,40].

TZM-bl cell-based neutralization assays

Neutralization assays using the autologous BG505.T332N virus and mutants, and the

heterologous SF162 virus, were carried out as described previously [58]. Nonlinear regression

curves were determined and 50% inhibitory concentration (IC_{50}) values were calculated using a

sigmoid function in Graphpad Prism v7.03.

Bio-Layer Interferometry (BLI)

An Octet RED instrument (FortéBio) was used to determine the kinetic parameters of the

antibody–antigen interactions by Biolayer Interferometry. Monoclonal Fabs were loaded onto

anti-human Fab-CH1 (FAB2G) biosensors (FortéBio) at a concentration of 10 μg/mL in kinetics

buffer (PBS, pH 7.4, 0.01% [w/v] BSA, and 0.002% [v/v] Tween 20) until a response of 1 nanometer

shift was reached. Loaded biosensors were dipped into kinetics buffer for 1 min to acquire a

baseline and then moved to wells containing a series of 2-fold dilutions of BG505 SOSIP.v5.2 in

kinetics buffer, starting at a 4000 nM. The trimers were allowed to associate for 180 secs before

the biosensor were move back to the wells containing kinetics buffer where the baseline was

acquired. Disassociation of the trimers from the Fab-loaded biosensors was recorded for 300

secs. All BLI experiments were conducted at 37°C. Kinetic parameters were calculated using the

Octet System Data Analysis v9.0 (FortéBio).

Negative Stain Electron Microscopy

BG505 SOSIP/Fab complexes were made by mixing 10-15 μg SOSIP with a 3 to 6-fold per protomer

molar excess for monoclonal Fabs or 500 μg polyclonal Fabs and allowed to incubate for 18 to 24

hrs at room temperature (RT). Complex samples were either diluted to 0.02 mg/mL and applied

to glow discharged negative stain grids or they were SEC purified using a Superose™ 6 Increase

10/300 GL (GE Healthcare) column to remove excess Fab prior to EM grid preparation. Fractions

containing the SOSIP/Fab complexes were pooled and concentrated using 10 kDa Amicon® spin

concentrators (Millipore). Samples were diluted to 0.03 mg/mL in TBS (0.05 M Tris pH 7.4, 0.15

M NaCl) and adsorbed onto glow discharged carbon-coated Cu400 EM grids (Electron Microscopy

Sciences) and blotted after 10 seconds. The grids were then stained with 3 μL of 2% (w/v) uranyl

formate, immediately blotted, and stained again for 45 secs followed by a final blot. Image

collection and data processing was performed as described previously on either an FEI Tecnai T12

microscope (2.05 Å/pixel; 52,000× magnification) or FEI Talos microscope (1.98 Å/pixel; 72,000×

magnification) with an electron dose of ∼25 electrons/Å^2 using Leginon [59,60]. 2D classification, 3D

sorting and 3D refinement conducted using Relion v3.0 [61]. EM density maps were visualized using

UCSF Chimera and segmented using Segger [62,63].

X-ray Crystallography Data Collection and Processing

All crystals were grown using sitting drop vapor diffusion. The RM20F Fab was crystallized from

a solution containing 10 mg/mL protein in TBS with a well solution containing 0.1M MES, pH 5.0

and 2M ammonium sulfate. The crystals were cryoprotected by soaking in a well solution

supplemented with 30% ethylene glycol. The RM20J Fab was crystallized from a solution

containing 10 mg/mL protein in TBS with a well solution containing 0.1M MES, pH 6.0, 5%

PEG3000 and 40% PEG400, with no cryoprotectant supplemented. The RM20E1 Fab was

crystallized from a solution containing 6.3 mg/mL protein in TBS with a well solution containing

0.1M glycine, pH 10.5, 1.2M NaH_2PO_4, 0.8M Na_2HPO_4, and 0.2M Li_2SO_4, with 15% ethylene glycol

supplemented as cryoprotectant. All crystals were grown at 298 K. Diffraction data for RM20F

and RM20E1 were collected at the Stanford Synchrotron Radiation Lightsource (SSRL) beamline

BL12–2, and that for RM20J collected at the Advanced Photon Source (APS) beamline 23ID-B.

Data collection and processing statistics are detailed in Table S5. Data sets were indexed,

integrated, and scaled using the HKL-2000 package [64]. The structures were solved by molecular

replacement using PHASER [65] with a homology model (SWISS-MODEL; [66-68]) as a search model

and further refined using phenix.refine [69] combined with manual building cycles in Coot [70].

Cryo Electron Microscopy Sample Preparation

RM20J complex: 500 µg BG505 SOSIP.v5.2 was mixed with 656 µg RM20J Fab and incubated at

RT overnight. The complex was SEC purified using a HiLoad® 16/600 Superdex® pg200 (GE

Healthcare) column in TBS. Fractions containing the complex were concentrated to 6.1 mg/mL

using a 10 kDa Amicon® spin concentrator (Millipore). 3.5 µL of the complex was mixed with 0.57

µL of 0.04 mM lauryl maltose neopentyl glycol (LMNG) and applied to a C-Flat grid (CF-2/1-4C,

Protochips, Inc.), which had been plasma-cleaned for 5 seconds using a mixture of N_2/O_2 (Gatan

Solarus 950 Plasma system). The grid was blotted and plunged into liquid ethane using a Vitrobot

Mark IV (ThermoFisher).

RM20F complex: 500 µg BG505 SOSIP.v4.1 was mixed with approximately 1,000 µg RM20F Fab

and incubated at RT overnight. The complex was SEC purified using a Superose^{TM} 6 Increase

10/300 GL (GE Healthcare) column in TBS. Fractions containing the complex were concentrated

to 6 mg/mL using a 10 kDa Amicon® spin concentrator (Millipore). 3 µL of the complex was mixed

with 1 µL of a n-Dodecyl-β-D-Maltopyranoside (DDM) solution to a final DDM concentration of

0.06 mM and applied to a C-Flat grid (CF-2/2-4C, Protochips, Inc.), which had been plasma-

cleaned for 5 seconds using a mixture of N_2/O_2 (Gatan Solarus 950 Plasma system). The grid was

blotted and plunged into liquid Ethane using a Vitrobot Mark IV (ThermoFisher).

RM20E1 complex: 355 µg BG505 SOSIP.v5.2 was mixed with 484 µg RM20E1 Fab and 484 µg

PGT122 Fab and incubated at RT overnight. The complex was SEC purified using a HiLoad® 16/600

Superdex® pg200 (GE Healthcare) column in TBS. Fractions containing the complex were

concentrated to 4 mg/mL using a 10 kDa Amicon® spin concentrator (Millipore). 3 µL of the

complex was mixed with 1 µL of a n-Dodecyl-β-D-Maltopyranoside (DDM) solution to a final DDM

concentration of 0.06 mM and applied to a grid (Quantifoil R 1.2/1.3, 400), which had been

plasma-cleaned for 5 seconds using a mixture of N_2/O_2 (Gatan Solarus 950 Plasma system). The

grid was blotted and plunged into liquid Ethane using a Vitrobot Mark IV (ThermoFisher).

Cryo Electron Microscopy Data Collection and Processing

Samples were imaged on either FEI Titan Krios electron microscope (ThermoFisher) operating at

300 keV (RM20F dataset) or a FEI Talos Arctica electron microscope (ThermoFisher) operating at

200 keV (RM20J and RM20E1 datasets). Both microscopes were equipped with Gatan K2 Summit

direct electron directors operating in counting mode. Automated data collection was performed

using the Leginon software suite [59]. Micrograph movie frames were aligned and dose-weighted

using MotionCor2 [71], and CTF models were determined using Gctf [72]. Particle picking, 2D

classification, Ab-initio reconstruction, and 3D refinement were conducted using cryoSPARCv2 [73].

Data collection and processing parameters are reported in Table S4.

Initial molecular models of the BG505 SOSIP trimer/Fab complexes were built by docking the Env

portion of PDB: 5V8M [74] into the EM density maps along with the relevant Fab crystal structures

(PDB: 4JY5 was used for PGT122 [75]) using UCSF Chimera [62]. The Fab constant regions were

removed due to flexibility in the elbow region as commonly found in Fab structures [76], the

appropriate stabilizing mutations (v4.1 or v5.2) were introduced into the Env sequence, and N-

linked glycans were added using Coot [77]. The models were iteratively refined into the EM density

maps using RosettaRelax and Coot [70,77-80]. Glycan structures were validated using Privateer [81].

Overall structures were evaluated using EMRinger [82] and MolProbity [83]. Protein interface

calculations were performed using jsPISA [84]. Final model statistics are summarized in Table S4.

Statistical analysis

Statistical models inherent to Relion 3.0 [61] and cryoSPARC [73] were employed in image analysis to

derive 2D classes and 3D models. All ELISA and neutralization assays were conducted with at least

duplicate measurements.

Data and Software Availability

The accession numbers for Env-specific BCR sequences are DDBJ/ENA/GenBank: MT002976-

MT002992 and MT008262-MT008328. RM IgM BCR sequences are available under BioProject ID:

PRJNA604386. Atomic coordinates and structure factors of the reported crystal structure have

been deposited in the Protein Data Bank (PDB: 6VOS, 6VOR, 6VSR). Cryo-EM reconstructions have

been deposited in the Electron Microscopy Data Bank (EMDB: EMD-21246, EMD-21257, EMD-

21232), and in the Protein Data Bank (PDB: 6VN0, 6VO1, 6VLR). The accession numbers for the

negative stain 3D EM reconstructions are Electron Microscopy DataBank: EMD-21053, EMD-

21055, EMD-21056, EMD-21057, EMD-21058, EMD-21059, EMD-21061, EMD-21062, EMD-

21064, EMD-21065, EMD-21066, EMD-21075, EMD-21076, EMD-21077, EMD-21078, EMD-

21079, EMD-21080, EMD-21081, EMD-21082, EMD-21083, EMD-21084, EMD-21085, EMD-

21086, EMD-21087, EMD-21088, EMD-21089, EMD-21090, EMD-21091, EMD-21092, EMD-

21093, EMD-21272, EMD-21273, EMD-21274, EMD-21275, EMD-21276, EMD-21277, EMD-

21278.

Figure Legends

Figure 1. MAb isolation and characterization from BG505 SOSIP.664 trimer-immunized RMs. (A)

Simplified immunization scheme adapted from Sanders et al. 2015 [24]. Black arrows indicate i.m.

immunizations with 100 µg of BG505 SOSIP.664 with 75 units of ISCOMATRIX adjuvant. Red

arrows indicate blood draws. (B) Heavy and light chain genetic characteristics for mAbs isolated

from rh1987 and rh2011. (C) TZM-bl neutralization of BG505.T332N, BG505.T332N.N611A, and

SF162 pseudoviruses for mAbs isolated from RM rh1987. (D) TZM-bl neutralization of

BG505.T332N, BG505.T332N.N611A, and SF162 pseudoviruses for mAbs isolated from RM

rh2011. Assay limit of detection was at an IC_{50} of 50 µg/mL.

Figure 2. Epitope mapping by negative stain electron microscopy. (A) Representative base-

targeting mAbs for animals rh1987 and rh2011. (B) Overlapping gp120/gp41 interface epitope

targeted by mAbs from both animals. (C) Overlapping FP epitope targeted by mAbs from both

animals. (D) Overlapping N289-glycan hole epitope targeted by mAbs from both animals. (E)

EMPEM analysis for wk28 IgG from rh2011 and rh1987. All structural figures were generated with

UCSF Chimera [62]. EM density maps were segmented with the Segger extension in UCSF Chimera

[62,63].

Figure 3. RM20J binds to the N289 glycan hole region of BG505 SOSIP.v5.2 (A) Segmented 3.9 Å

cryoEM reconstruction of RM20J Fab (pink) in complex with BG505 SOSIP.v5.2 (gp120, dark blue;

gp41, light blue). (B) and (C) Zoomed-in views of the epitope/paratope interaction between

gp120 (blue, ribbon diagram) and RM20J Fab (surface representation). (D) and (E) Comparison of

the RM20J epitope with that of the BG505 SOSIP.664 elicited rabbit neutralizing NAb 10A (PDB:

6DID) [19,36].

Figure 4. RM20F binds to a quaternary epitope at the gp120/gp41 interface of BG505 SOSIP.v4.1.

(A) Segmented 4.3 Å cryoEM reconstruction of RM20F Fab (orange) in complex with BG505 SOSIP

(gp120, dark blue; gp41, light blue). (B) and (C) Zoomed-in views of the epitope/paratope

interaction between BG505 SOSIP (gp120, dark blue; gp41s, light blue and light green; ribbon

diagram) and RM20F Fab (surface representation). (D) Neutralization data for BG505 mutant

pseudoviruses against VRC01, RM20F, and VRC34.

Figure 5. RM20E1 binds to the fusion peptide of BG505 SOSIP.v5.2. (A) Segmented 4.2 Å cryoEM

reconstruction of RM20E1 Fab (yellow) and PGT122 Fab (light green) in complex with BG505

SOSIP (gp120, dark blue; gp41, light blue). (B) and (C) Zoomed-in views of the epitope/paratope

interaction between BG505 SOSIP (gp120, dark blue; gp41s, light blue and light green; ribbon

diagram) and RM20E1 Fab (surface representation). (D) Comparison of RM20E1 and RM20F

epitopes to those of FP targeting bNAbs VRC34, ACS202, and DFPH-a.15 [9,25,39,40].

Table S1. Initial germline BCR database used in IgDiscover

Acknowledgements

We thank Jean-Christophe Ducom, Hannah Turner, and Bill Anderson for assistance with

computational resources, microscope management, and data collection. We thank Alan Saluk,

Brian Seegers, Steven Head, Jessica Ledesma, Curt Wittenberg, and Nitya Bhaskaran for

assistance with naïve B-cell sorting and NGS sequencing. This work was supported by the HIV

Vaccine Research and Design (HIVRAD) program (P01 AI110657) (A.B.W., R.W.S, J.P.M., and

I.A.W.), NIH CHAVI-ID (UM1 AI100663) and CHAVD (UM1 AI44462) awards (A.B.W., I.A.W. and

D.R.B.), NIH R01 AI13082 (J.P.M.), the International AIDS Vaccine Initiative Neutralizing Antibody

Center, the Bill and Melinda Gates Foundation CAVD (OPP1115782, OPP1132237, OPP1084519,

OPP119635), and the European Union's Horizon 2020 research and innovation program under

grant agreement no. 681137 (R.W.S.). C.A.C. is supported by the NIH F31 Ruth L. Kirschstein

Predoctoral Award AI131873 and by the Achievement Rewards for College Scientists Foundation.

R.W.S. is a recipient of a Vici fellowship from the Netherlands Organization for Scientific Research

(NWO). G.O. and M.J.v.G. are supported by amfAR Mathilde Krim Fellowships in Basic Biomedical

Research grant numbers 109718-63-RKNT and 109514-61-RKVA, respectively. J.v.S. is a recipient

of a 2017 AMC Ph.D. Scholarship. Computational analyses of EM data were performed using

shared instrumentation funded by NIH grant S10OD021634. Naïve B-cell sorting and NGS

sequencing was conducted as part of the Technologies Across Scale graduate course with support

from Skaggs Graduate School of Chemical and Biological Sciences at Scripps Research. Use of the

Stanford Synchrotron Radiation Lightsource, SLAC National Accelerator Laboratory, is supported

by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences under

Contract No. DE-AC02-76SF00515. The SSRL Structural Molecular Biology Program is supported

by the DOE Office of Biological and Environmental Research, and by the National Institutes of

Health, National Institute of General Medical Sciences (including P41GM103393). The contents

of this publication are solely the responsibility of the authors and do not necessarily represent

the official views of NIGMS, NIAID, or NIH.

Author Contributions

C.A.C., J.v.S., M.Y., D.O., I.A.W., R.W.S., A.B.W., and M.J.v.G. designed the experiments. C.A.C.,

J.v.S., M.S., R.M., P.v.d.W., M.v.B., J.L.T., R.P., J.C., D.S., C.L., and M.J.v.G. performed the

experiments. C.A.C., J.v.S., C.L., D.M., and M.J.v.G. analyzed the data. C.A.C., M.Y., D.O., M.S.,

L.M.S., G.O., and B.N. performed the structural studies. C.A.C., C.A.B., J.G., and S.C. developed

the macaque germline database. E.G.R., D.G.C., and G.S. provided the macaque PBMC samples.

V.V., D.N.S., D.R.B., and J.P.M. provided resources. C.A.C., A.B.W., and M.J.v.G. wrote the

manuscript with input from all authors.

References

1   UNAIDS. Global HIV & AIDS statistics — 2018 fact sheet. (2019).

2   Moldt, B. *et al.* Neutralizing antibody affords comparable protection against vaginal and
rectal simian/human immunodeficiency virus challenge in macaques. *Aids* **30**, 1543-
1551, doi:10.1097/QAD.0000000000001102 (2016).

3   Burton, D. R. & Hangartner, L. Broadly Neutralizing Antibodies to HIV and Their Role in
Vaccine Design. *Annual review of immunology* **34**, 635-659, doi:10.1146/annurev-
immunol-041015-055515 (2016).

4   Pegu, A. *et al.* A Meta-analysis of Passive Immunization Studies Shows that Serum-
Neutralizing Antibody Titer Associates with Protection against SHIV Challenge. *Cell Host*
& Microbe **26**, 336-+, doi:10.1016/j.chom.2019.08.014 (2019).
5   Wu, X. HIV Broadly Neutralizing Antibodies: VRC01 and Beyond. *Adv Exp Med Biol* **1075**,
53-72, doi:10.1007/978-981-13-0484-2_3 (2018).
6   Jardine, J. G. *et al.* HIV-1 broadly neutralizing antibody precursor B cells revealed by
germline-targeting immunogen. *Science* **351**, 1458-1463, doi:10.1126/science.aad9195
(2016).
7   Torrents de la Pena, A. *et al.* Improving the Immunogenicity of Native-like HIV-1
Envelope Trimers by Hyperstabilization. *Cell Rep* **20**, 1805-1817,
doi:10.1016/j.celrep.2017.07.077 (2017).
8   de Taeye, S. W. *et al.* Immunogenicity of Stabilized HIV-1 Envelope Trimers with
Reduced Exposure of Non-neutralizing Epitopes. *Cell* **163**, 1702-1715,
doi:10.1016/j.cell.2015.11.056 (2015).
9   Kong, R. *et al.* Antibody Lineages with Vaccine-Induced Antigen-Binding Hotspots
Develop Broad HIV Neutralization. *Cell* **178**, 567-584 e519,
doi:10.1016/j.cell.2019.06.030 (2019).
10   Escolano, A. *et al.* Immunization expands B cells specific to HIV-1 V3 glycan in mice and
macaques. *Nature* **570**, 468-473, doi:10.1038/s41586-019-1250-z (2019).
11   Escolano, A. *et al.* Sequential Immunization Elicits Broadly Neutralizing Anti-HIV-1
Antibodies in Ig Knockin Mice. *Cell* **166**, 1445-1458 e1412,
doi:10.1016/j.cell.2016.07.030 (2016).
12   Steichen, J. M. *et al.* HIV Vaccine Design to Target Germline Precursors of Glycan-
Dependent Broadly Neutralizing Antibodies. *Immunity* **45**, 483-496,
doi:10.1016/j.immuni.2016.08.016 (2016).
13   Sanders, R. W. *et al.* A next-generation cleaved, soluble HIV-1 Env trimer, BG505
SOSIP.664 gp140, expresses multiple epitopes for broadly neutralizing but not non-
neutralizing antibodies. *PLoS pathogens* **9**, e1003618, doi:10.1371/journal.ppat.1003618
(2013).
14   Julien, J. P. *et al.* Crystal structure of a soluble cleaved HIV-1 envelope trimer. *Science*
342, 1477-1483, doi:10.1126/science.1245625 (2013).
15   Lyumkis, D. *et al.* Cryo-EM structure of a fully glycosylated soluble cleaved HIV-1
envelope trimer. *Science* **342**, 1484-1490, doi:10.1126/science.1245627 (2013).
16   Pancera, M. *et al.* Structure and immune recognition of trimeric pre-fusion HIV-1 Env.
Nature **514**, 455-461, doi:10.1038/nature13808 (2014).
17   Pauthner, M. *et al.* Elicitation of Robust Tier 2 Neutralizing Antibody Responses in
Nonhuman Primates by HIV Envelope Trimer Immunization Using Optimized
Approaches. *Immunity* **46**, 1073-1088 e1076, doi:10.1016/j.immuni.2017.05.007 (2017).
18   Pauthner, M. G. *et al.* Vaccine-Induced Protection from Homologous Tier 2 SHIV
Challenge in Nonhuman Primates Depends on Serum-Neutralizing Antibody Titers.
Immunity **50**, 241-252 e246, doi:10.1016/j.immuni.2018.11.011 (2019).
19   McCoy, L. E. *et al.* Holes in the Glycan Shield of the Native HIV Envelope Are a Target of
Trimer-Elicited Neutralizing Antibodies. *Cell Rep* **16**, 2327-2338,
doi:10.1016/j.celrep.2016.07.074 (2016).

20   Klasse, P. J. *et al.* Epitopes for neutralizing antibodies induced by HIV-1 envelope
679   glycoprotein BG505 SOSIP trimers in rabbits and macaques. *PLoS pathogens* **14**,
680   e1006913, doi:10.1371/journal.ppat.1006913 (2018).
681   21   Cirelli, K. M. *et al.* Slow Delivery Immunization Enhances HIV Neutralizing Antibody and
682   Germinal Center Responses via Modulation of Immunodominance. *Cell*,
683   doi:10.1016/j.cell.2019.04.012 (2019).
684   22   Lei, L. *et al.* The HIV-1 Envelope Glycoprotein C3/V4 Region Defines a Prevalent
685   Neutralization Epitope following Immunization. *Cell Rep* **27**, 586-598 e586,
686   doi:10.1016/j.celrep.2019.03.039 (2019).
687   23   Nogal, B. *et al.* Mapping polyclonal antibody responses in non-human primates
688   vaccinated with HIV Env trimer subunit vaccines. *bioRxiv*, 833715, doi:10.1101/833715
689   (2019).
690   24   Sanders, R. W. *et al.* HIV-1 VACCINES. HIV-1 neutralizing antibodies induced by native-
691   like envelope trimers. *Science* **349**, aac4223, doi:10.1126/science.aac4223 (2015).
692   25   Kong, R. *et al.* Fusion peptide of HIV-1 as a site of vulnerability to neutralizing antibody.
693   *Science* **352**, 828-833, doi:10.1126/science.aae0474 (2016).
694   26   Landais, E. & Moore, P. L. Development of broadly neutralizing antibodies in HIV-1
695   infected elite neutralizers. *Retrovirology* **15**, 61, doi:10.1186/s12977-018-0443-0 (2018).
696   27   Corcoran, M. M. *et al.* Production of individualized V gene databases reveals high levels
697   of immunoglobulin genetic diversity. *Nat Commun* **7**, 13642, doi:10.1038/ncomms13642
698   (2016).
699   28   Vigdorovich, V. *et al.* Repertoire comparison of the B-cell receptor-encoding loci in
700   humans and rhesus macaques by next-generation sequencing. *Clin Transl Immunology*
701   **5**, e93, doi:10.1038/cti.2016.42 (2016).
702   29   Johnson, M. *et al.* NCBI BLAST: a better web interface. *Nucleic Acids Res* **36**, W5-9,
703   doi:10.1093/nar/gkn201 (2008).
704   30   Zimin, A. V. *et al.* A new rhesus macaque assembly and annotation for next-generation
705   sequencing analyses. *Biol Direct* **9**, 20, doi:10.1186/1745-6150-9-20 (2014).
706   31   Ramesh, A. *et al.* Structure and Diversity of the Rhesus Macaque Immunoglobulin Loci
707   through Multiple De Novo Genome Assemblies. *Frontiers in immunology* **8**, 1407,
708   doi:10.3389/fimmu.2017.01407 (2017).
709   32   Ralph, D. K. & Matsen, F. A. I. Per-sample immunoglobulin germline inference from B
710   cell receptor deep sequencing data. *arXiv* (2018).
711   33   Gadala-Maria, D., Yaari, G., Uduman, M. & Kleinstein, S. H. Automated analysis of high-
712   throughput B-cell sequencing data reveals a high frequency of novel immunoglobulin V
713   gene segment alleles. *Proceedings of the National Academy of Sciences of the United*
714   *States of America* **112**, E862-870, doi:10.1073/pnas.1417683112 (2015).
715   34   Guo, K. *et al.* Immunoglobulin VH gene diversity and somatic hypermutation during SIV
716   infection of rhesus macaques. *Immunogenetics* **67**, 355-370, doi:10.1007/s00251-015-
717   0844-3 (2015).
718   35   Ye, J., Ma, N., Madden, T. L. & Ostell, J. M. IgBLAST: an immunoglobulin variable domain
719   sequence analysis tool. *Nucleic Acids Res* **41**, W34-40, doi:10.1093/nar/gkt382 (2013).

36 Bianchi, M. *et al.* Electron-Microscopy-Based Epitope Mapping Defines Specificities of
Polyclonal Antibodies Elicited during HIV-1 BG505 Envelope Trimer Immunization.
Immunity **49**, 288-300 e288, doi:10.1016/j.immuni.2018.07.009 (2018).
37 Cao, L. *et al.* Differential processing of HIV envelope glycans on the virus and soluble
recombinant trimer. *Nat Commun* **9**, 3693, doi:10.1038/s41467-018-06121-4 (2018).
38 Struwe, W. B. *et al.* Site-Specific Glycosylation of Virion-Derived HIV-1 Env Is Mimicked
by a Soluble Trimeric Immunogen. *Cell Rep* **24**, 1958-1966 e1955,
doi:10.1016/j.celrep.2018.07.080 (2018).
39 Yuan, M. *et al.* Conformational Plasticity in the HIV-1 Fusion Peptide Facilitates
Recognition by Broadly Neutralizing Antibodies. *Cell Host Microbe* **25**, 873-883 e875,
doi:10.1016/j.chom.2019.04.011 (2019).
40 van Gils, M. J. *et al.* An HIV-1 antibody from an elite neutralizer implicates the fusion
peptide as a site of vulnerability. *Nat Microbiol* **2**, 16199,
doi:10.1038/nmicrobiol.2016.199 (2016).
41 Dubrovskaya, V. *et al.* Vaccination with Glycan-Modified HIV NFL Envelope Trimer-
Liposomes Elicits Broadly Neutralizing Antibodies to Multiple Sites of Vulnerability.
Immunity **51**, 915-929 e917, doi:10.1016/j.immuni.2019.10.008 (2019).
42 Yang, L. *et al.* Structure-Guided Redesign Improves NFL HIV Env Trimer Integrity and
Identifies an Inter-Protomer Disulfide Permitting Post-Expression Cleavage. *Frontiers in
immunology* **9**, 1631, doi:10.3389/fimmu.2018.01631 (2018).
43 Cao, L. *et al.* Global site-specific N-glycosylation analysis of HIV envelope glycoprotein.
Nat Commun **8**, 14954, doi:10.1038/ncomms14954 (2017).
44 Xu, K. *et al.* Epitope-based vaccine design yields fusion peptide-directed antibodies that
neutralize diverse strains of HIV-1. *Nature medicine* **24**, 857-867, doi:10.1038/s41591-
018-0042-6 (2018).
45 Cheng, C. *et al.* Consistent elicitation of cross-clade HIV-neutralizing responses achieved
in guinea pigs after fusion peptide priming by repetitive envelope trimer boosting. *PloS
one* **14**, e0215163, doi:10.1371/journal.pone.0215163 (2019).
46 Brouwer, P. J. M. *et al.* Enhancing and shaping the immunogenicity of native-like HIV-1
envelope trimers with a two-component protein nanoparticle. *Nat Commun* **10**, 4272,
doi:10.1038/s41467-019-12080-1 (2019).
47 Kulp, D. W. *et al.* Structure-based design of native-like HIV-1 envelope trimers to silence
non-neutralizing epitopes and eliminate CD4 binding. *Nat Commun* **8**, 1655,
doi:10.1038/s41467-017-01549-6 (2017).
48 Georgiev, I. S. *et al.* Two-Component Ferritin Nanoparticles for Multimerization of
Diverse Trimeric Antigens. *ACS Infect Dis* **4**, 788-796, doi:10.1021/acsinfecdis.7b00192
(2018).
49 Ringe, R. P. *et al.* Neutralizing antibody induction by HIV-1 Envelope glycoprotein SOSIP
trimers on iron oxide nanoparticles may be impaired by mannose binding lectin. *J Virol*,
doi:10.1128/JVI.01883-19 (2019).
50 Wang, Y. *et al.* HIV-1 Cross-Reactive Primary Virus Neutralizing Antibody Response
Elicited by Immunization in Nonhuman Primates. *J Virol* **91**, doi:10.1128/JVI.00910-17
(2017).

51   Martinez-Murillo, P. *et al.* Particulate Array of Well-Ordered HIV Clade C Env Trimers
Elicits Neutralizing Antibodies that Display a Unique V2 Cap Approach. *Immunity* **46**,
804-817 e807, doi:10.1016/j.immuni.2017.04.021 (2017).

52   Phad, G. E. *et al.* Extensive dissemination and intraclonal maturation of HIV Env vaccine-
induced B cell responses. *The Journal of experimental medicine* **217**,
doi:10.1084/jem.20191155 (2020).

53   Turchaninova, M. A. *et al.* High-quality full-length immunoglobulin profiling with unique
molecular barcoding. *Nature protocols* **11**, 1599-1616, doi:10.1038/nprot.2016.093
(2016).

54   Christley, S. *et al.* VDJServer: A Cloud-Based Analysis Portal and Data Commons for
Immune Repertoire Sequences and Rearrangements. *Frontiers in immunology* **9**, 976,
doi:10.3389/fimmu.2018.00976 (2018).

55   Sievers, F. & Higgins, D. G. Clustal Omega for making accurate alignments of many
protein sequences. *Protein science : a publication of the Protein Society* **27**, 135-145,
doi:10.1002/pro.3290 (2018).

56   Sok, D. *et al.* Recombinant HIV envelope trimer selects for quaternary-dependent
antibodies targeting the trimer apex. *Proceedings of the National Academy of Sciences*
of the United States of America **111**, 17624-17629, doi:10.1073/pnas.1415789111
(2014).

57   Kabat, E. A. *Sequences of proteins of immunological interest : tabulation and analysis of*
amino acid and nucleic acid sequences of precursors, V-regions, C-regions, J-chain, T-cell
receptors for antigenm T-cell surface antigens, [beta]2-microglobulins,major
histocompatibility antigens, Thy-1, complement, C-reactive protein, thymopoietin,
integrins, post-gamme globulin, [alpha]2-macroglobulins, and other related proteins. 5th
edn, (U.S. Dept. of Health and Human Services, Public Health Service, National Institutes
of Health, 1991).

58   Derking, R. *et al.* Comprehensive antigenic map of a cleaved soluble HIV-1 envelope
trimer. *PLoS pathogens* **11**, e1004767, doi:10.1371/journal.ppat.1004767 (2015).

59   Suloway, C. *et al.* Automated molecular microscopy: the new Leginon system. *Journal of*
structural biology **151**, 41-60, doi:10.1016/j.jsb.2005.03.010 (2005).

60   Pugach, P. *et al.* A native-like SOSIP.664 trimer based on an HIV-1 subtype B env gene. *J*
Virol **89**, 3380-3395, doi:10.1128/JVI.03473-14 (2015).

61   Zivanov, J. *et al.* New tools for automated high-resolution cryo-EM structure
determination in RELION-3. *eLife* **7**, doi:10.7554/eLife.42166 (2018).

62   Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research and
analysis. *J Comput Chem* **25**, 1605-1612, doi:10.1002/jcc.20084 (2004).

63   Pintilie, G. D., Zhang, J., Goddard, T. D., Chiu, W. & Gossard, D. C. Quantitative analysis
of cryo-EM density map segmentation by watershed and scale-space filtering, and fitting
of structures by alignment to regions. *Journal of structural biology* **170**, 427-438,
doi:10.1016/j.jsb.2010.03.007 (2010).

64   Otwinowski, Z. & Minor, W. Processing of X-ray diffraction data collected in oscillation
mode. *Methods Enzymol* **276**, 307-326 (1997).

65   McCoy, A. J. *et al.* Phaser crystallographic software. *J Appl Crystallogr* **40**, 658-674,
doi:10.1107/S0021889807021206 (2007).

807 66 Arnold, K., Bordoli, L., Kopp, J. & Schwede, T. The SWISS-MODEL workspace: a web-
808   based environment for protein structure homology modelling. *Bioinformatics* **22**, 195-
809   201, doi:10.1093/bioinformatics/bti770 (2006).
810 67 Biasini, M. *et al.* SWISS-MODEL: modelling protein tertiary and quaternary structure
811   using evolutionary information. *Nucleic Acids Res* **42**, W252-258,
812   doi:10.1093/nar/gku340 (2014).
813 68 Bordoli, L. *et al.* Protein structure homology modeling using SWISS-MODEL workspace.
814   *Nature protocols* **4**, 1-13, doi:10.1038/nprot.2008.197 (2009).
815 69 Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular
816   structure solution. *Acta crystallographica. Section D, Biological crystallography* **66**, 213-
817   221, doi:10.1107/S0907444909052925 (2010).
818 70 Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot.
819   *Acta crystallographica. Section D, Biological crystallography* **66**, 486-501,
820   doi:10.1107/S0907444910007493 (2010).
821 71 Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for
822   improved cryo-electron microscopy. *Nature methods* **14**, 331-332,
823   doi:10.1038/nmeth.4193 (2017).
824 72 Zhang, K. Gctf: Real-time CTF determination and correction. *Journal of structural biology*
825   **193**, 1-12, doi:10.1016/j.jsb.2015.11.003 (2016).
826 73 Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for
827   rapid unsupervised cryo-EM structure determination. *Nature methods* **14**, 290-296,
828   doi:10.1038/nmeth.4169 (2017).
829 74 Lee, J. H. *et al.* A Broadly Neutralizing Antibody Targets the Dynamic HIV Envelope
830   Trimer Apex via a Long, Rigidified, and Anionic beta-Hairpin Structure. *Immunity* **46**,
831   690-702, doi:10.1016/j.immuni.2017.03.017 (2017).
832 75 Julien, J. P. *et al.* Broadly neutralizing antibody PGT121 allosterically modulates CD4
833   binding via recognition of the HIV-1 gp120 V3 base and multiple surrounding glycans.
834   *PLoS pathogens* **9**, e1003342, doi:10.1371/journal.ppat.1003342 (2013).
835 76 Bailey, L. J. *et al.* Locking the Elbow: Improved Antibody Fab Fragments as Chaperones
836   for Structure Determination. *Journal of molecular biology* **430**, 337-347,
837   doi:10.1016/j.jmb.2017.12.012 (2018).
838 77 van Beusekom, B. *et al.* Building and rebuilding N-glycans in protein structure models.
839   *Acta Crystallogr D Struct Biol* **75**, 416-425, doi:10.1107/S2059798319003875 (2019).
840 78 DiMaio, F., Tyka, M. D., Baker, M. L., Chiu, W. & Baker, D. Refinement of protein
841   structures into low-resolution density maps using rosetta. *Journal of molecular biology*
842   **392**, 181-190, doi:10.1016/j.jmb.2009.07.008 (2009).
843 79 Frenz, B. *et al.* Automatically Fixing Errors in Glycoprotein Structures with Rosetta.
844   *Structure*, doi:10.1016/j.str.2018.09.006 (2018).
845 80 Casanal, A., Lohkamp, B. & Emsley, P. Current Developments in Coot for
846   Macromolecular Model Building of Electron Cryo-microscopy and Crystallographic Data.
847   *Protein science : a publication of the Protein Society*, doi:10.1002/pro.3791 (2019).
848 81 Agirre, J. *et al.* Privateer: software for the conformational validation of carbohydrate
849   structures. *Nature structural & molecular biology* **22**, 833-834, doi:10.1038/nsmb.3115
850   (2015).

851   82   Barad, B. A. *et al.* EMRinger: side chain-directed model and map validation for 3D cryo-
852   electron microscopy. *Nature methods* **12**, 943-946, doi:10.1038/nmeth.3541 (2015).

853   83   Williams, C. J. *et al.* MolProbity: More and better reference data for improved all-atom
854   structure validation. *Protein science : a publication of the Protein Society* **27**, 293-315,
855   doi:10.1002/pro.3330 (2018).

856   84   Krissinel, E. Stock-based detection of protein oligomeric states in jsPISA. *Nucleic Acids
857   Res* **43**, W314-319, doi:10.1093/nar/gkv314 (2015).

A

Immunizations

Weeks 0 4 12 22 24 25 28 38 52 53

Blood Draws

B

rh1987 heavy
Avg SHM: 6.4%
Avg CDR3
length: 15aa
25 mAbs

rh2011 heavy
Avg SHM: 6.1%
Avg CDR3
length: 17aa
17 mAbs

VH1 VH3 VH4 VH5

rh1987 light
Avg SHM: 4.2%
Avg CDR3
length: 10aa
25 mAbs

rh2011 light
Avg SHM: 4.3%
Avg CDR3
length: 10aa
17 mAbs

VK1 VK2 VK3 VK6 VL1 VL2 VL4 VL8 VL11

C

rh1987

mAb	BG505 T332N	BG505 T332N+N611A	SF162
RM19A	41.8	>50	>50
RM19A1	44.9	>50	>50
RM19A2	>50	>50	>50
RM19A3	>50	>50	>50
RM19B1	>50	>50	>50
RM19C3	>50	>50	>50
RM19C4	>50	>50	>50
RM19D	>50	>50	>50
RM19J	>50	>50	>50
RM19M	12.9	5.63	>50
RM19N	>50	>50	>50
RM19O	>50	25.5	>50
RM19P	23.5	>50	>50
RM19R	>50	>50	>50
RM19S	>50	3.51	>50
RM19T	>50	>50	>50

IC_{50} (µg/mL)

D

rh2011

mAb	BG505 T332N	BG505 T332N+N611A	SF162
RM20A3	>50	>50	>50
RM20B	>50	>50	>50
RM20C	>50	>50	>50
RM20D	>50	>50	>50
RM20E	>50	<0.41	>50
RM20E1	>50	<0.41	>50
RM20E2	>50	<0.41	>50
RM20E3	>50	<0.41	>50
RM20F	1.6	<0.41	>50
RM20H	42.9	<0.41	>50
RM20I	>50	>50	>50
RM20J	>50	>50	>50

IC_{50} (µg/mL)

Figure#3

A

gp120
gp41
RM20J

90°

B

E269
K344
E268
N289
N339
T290

C

gp120
LC
CDR-L2
HC
CDR-H1
CDR-H2
CDR-H3

N355
K347
E340
disordered V4
K351
N339

D

gp120
gp41
RM20J
10A

E

disordered V4
N289
S241

A

gp120
gp41
RM20F

90°

B

gp120
gp41_1
gp41_2
HC
LC
CDR-H2
CDR-H3
CDR-L3
FR-H3

K229
H85
N88 glycan

C

gp120
gp41_1
gp41_2
fusion peptide
LC
CDR-H2
CDR-H3
CDR-L3

L523
V518
HR2

D

	VRC01	RM20F	VRC34
WT	1.0	1.0	1.0
H85A	1.2	11.9	4.7
E87A	1.2	4.3	0.8
N88A	1	>28	>24
K229A	0.6	3.1	1.2
S241N	1.1	6.6	n.d.
E267A	0.8	1.3	0.9
P291T	1.1	2.9	n.d.
N332T	0.9	0.5	n.d.
F522A	0.8	3.9	1.0
M535A	0.9	2.9	1.0
T536A	0.8	2.9	1.5
T538A	0.8	3.6	1.2
N611Q	1.5	<0.1	n.d.
N625Q	1.9	0.6	n.d.
N637Q	1.6	1.2	n.d.
G644A	1.0	4.2	1.8
E647A	1.7	13.5	0.6

fold reduction in neutralization

Figure#5

A
PGT122
gp120
gp41
RM20E1

B
gp120
gp41_1
gp41_2
fusion peptide
HC
CDR-H1
CDR-H3
LC
CDR-L1
CDR-L3

H85
N88 glycan

C
gp120
gp41_1
gp41_2
fusion peptide
HC
FR-H1
CDR-H1
CDR-H3
LC
CDR-L1
CDR-L3

F519
HR2
N611
N637

D
Env   RM20E1   RM20F   DFPH-a.15   VRC34   ACS202