The Nucleus Accumbens Core is Necessary to Scale Fear to Degree of Threat

Madelyn H. Ray, Alyssa N. Russ, Rachel A. Walker, and Michael A. McDannald

Department of Psychology and Neuroscience, Boston College, Chestnut Hill, MA 02467

Correspondence and requests for materials should be addressed to:

Michael A. McDannald
Department of Psychology and Neuroscience
Boston College
140 Commonwealth Ave.
514 McGuinn Hall
Chestnut Hill, MA 02467, USA
michael.mcdannald@bc.edu

or

Madelyn H. Ray
Department of Psychology and Neuroscience
Boston College
140 Commonwealth Ave.
Chestnut Hill, MA 02467, USA
madelyn.ray@bc.edu

Figures: 4
Tables: 1
Pages: 26
Words
Abstract: 185
Introduction: 397
Discussion: 1,429

Conflict of interest: The authors declare no competing financial interests.

Acknowledgements: We would like to thank Alexa LaBanca and Andrew Thomson for technical support. This work was supported by NIH DA034010 to M.A.M. We thank Bret Judson and the Boston College Imaging Core for infrastructure and support.

Abstract

Fear is maximally adaptive when the response level rapidly scales to the degree of threat. Using a behavioral procedure consisting of danger, uncertainty and safety cues, we have found rapid scaling of fear (within two seconds of cue presentation) in male rats. Here we examined a possible role for the nucleus accumbens core (NAcc) in the acquisition and expression of rapid fear scaling. In experiment 1, male Long Evans rats received bilateral sham or neurotoxic NAcc lesions. NAcc-lesioned rats showed general impairment in scaling fear to degree of threat, as well as impairment in a specific component of scaling: rapid uncertainty-safety discrimination. In experiment 2, male Long Evans rats received NAcc transduction with halorhodopsin or a control fluorophore. Rats were trained to the point of discrimination, and the NAcc was green-light illuminated during cue or control periods. NAcc-halorhodopsin rats receiving light illumination during cue presentation were specifically impaired in rapid uncertainty-safety discrimination. The results reveal a general role for the NAcc in scaling of fear to degree of threat, and a specific role in one component of scaling: rapid discrimination of uncertain threat and safety.

Significance Statement

Rapidly discriminating cues for threat and safety is essential for survival and impaired threat-safety discrimination is a hallmark of anxiety disorders. In the current study, we induced nucleus accumbens core (NAcc) dysfunction while rats underwent fear discrimination consisting of cues for danger, uncertainty and safety. Permanent NAcc dysfunction, induced via neurotoxic lesion, generally disrupted scaling of fear to degree of threat, and specifically impaired one component of scaling: the acquisition of rapid uncertainty-safety discrimination. Reversible NAcc dysfunction, induced via optogenetic inhibition, specifically impaired expression of rapid, uncertainty-safety discrimination. The results reveal that the NAcc is essential to scale fear to degree of threat, and is a plausible source of dysfunction in anxiety disorders.

Introduction

The ability to discriminate danger from safety is critical for survival. Individuals with stress and anxiety disorders are impaired in discrimination, showing excessive fear-related responses to safety (Jovanovic et al., 2010; Jovanovic et al., 2012; Lissek et al., 2014; Duits et al., 2015). Danger and safety represent extremes of a threat continuum, with most real-world threats involving uncertainty. Ideally, one's level of fear should scale to the degree of threat. This scaling of the fear response would be most adaptive if observed rapidly following encounter with a potential threat.

Drawing from learning theory (Rescorla, 1968), our laboratory has observed rapid scaling of fear to danger, uncertainty and safety in male rats (DiLeo et al., 2016). Present work in our laboratory is aimed at identifying brain regions that permit rapid scaling of fear. Candidate regions should be able to rapidly process valence information and should receive amygdalar input (Quirk et al., 1995; Goosens and Maren, 2001; Koo et al., 2004; McDannald and Galarce, 2011). We identified the nucleus accumbens core (NAcc) as a likely candidate, based on its ability to rapidly process reward-predictive cues (Cromwell and Schultz, 2003; Setlow et al., 2003; Ambroggi et al., 2011; McGinty et al., 2013; Saddoris and Carelli, 2014; Sugam et al., 2014; Ottenheimer et al., 2018), as well as its anatomical connectivity with the amygdala (Kita and Kitai, 1990; Petrovich et al., 1996; Wright and Groenewegen, 1996). Even more, the NAcc is implicated in a variety of fear-related processes (Haralambous and Westbrook, 1999; Thomas et al., 2002; Schwienbacher et al., 2004; Iordanova et al., 2006a; Fadok et al., 2010; Badrinarayan et al., 2012; Oleson et al., 2012; Li and McNally, 2015; Correia et al., 2016).

In two experiments, we examined roles for the NAcc in scaling of fear. In experiment 1, we permanently ablated NAcc neurons via neurotoxic lesion. Following recovery, rats received fear discrimination in which auditory cues predicted unique foot shock probabilities: danger (p=1.00), uncertainty (p=0.25), and safety (p=0.00). To isolate rapid scaling of fear, we broke the 10-s cues into five, 2-s cue intervals. This permitted analysis of rapid scaling of fear, focusing on the first 2-s cue interval, as well as overall scaling across all intervals. In experiment 2, we examined a role for the NAcc in the expression of fear scaling. Rats acquired fear scaling and once stable, NAcc neural activity was optogenetically inhibited at the time of cue presentation or a control period. Combined, the two experiments allowed us to examine roles for the NAcc in the acquisition and expression of fear scaling.

Materials and methods

Experiment 1

Subjects

Subjects were forty-five male Long Evans rats weighing 275-300 g upon arrival (Charles River Laboratories; RGD Cat# 2308852, RRID:RGD_2308852). Rats were individually housed and maintained on a 12-h dark light cycle (lights off at 6:00 PM) with water *ad libitum*. Procedures adhered to the NIH Guide for the Care and Use of Laboratory Animals and were approved by the Boston College Institutional Animal Care and Use Committee.

Behavioral apparatus

Eight sound-attenuated enclosures each housed a behavior chamber with aluminum front and back walls, clear acrylic sides and top, and a metal grid floor. Grid floors were electrically connected to a shock generator. A single external food cup and central nose poke opening equipped with infrared photocells were present on one wall. Auditory stimuli were presented through two speakers mounted on the ceiling of each behavior chamber.

Surgical procedures

Stereotaxic surgery was performed under isoflurane anesthesia (2-5%) using aseptic technique. Twenty-four rats received bilateral infusions of *N*-Methyl-D-aspartic acid (15 µg/µl in Dulbecco's PBS) aimed at the nucleus accumbens core (0.40 µl, +1.90 AP, ±1.80 ML, -6.60 DV from skull). Infusions were delivered via 2 µl syringe (Hamilton, Neuros) controlled by a microsyringe pump (World Precision Instruments, UMP3-2). Infusion rate was ~0.11 µl/min. Thirty seconds after the completion of each infusion, the syringe was raised 0.1 mm then left in place for five minutes to encourage delivery to the target site. The remaining twenty-one rats received identical surgical treatment without infusions. Rats received carprofen (5 mg/kg) for post-operative analgesia.

Nose poke acquisition

Following recovery from surgery, rats were food restricted to 85% of their initial free feeding body weight, then fed (2 - 20 g/day) to increase their target body weight by 1 g/day for the remainder of testing. Rats were shaped to nose poke for pellet (BioServ F0021 – protein/fat/carbohydrate blend) delivery using a fixed ratio 1 schedule: one nose poke yielded one pellet. Shaping sessions lasted 30 min or approx-

imately 50 nose pokes. Over the next 3, 60-min sessions, rats were placed on variable interval (VI) schedules in which nose pokes were reinforced on average every 30 s (session 1), or 60 s (sessions 2 and 3). For the remainder of testing, nose pokes were reinforced on a VI-60 schedule independent of all Pavlovian contingencies.

Pre-exposure

In two separate sessions, each rat was pre-exposed to the three cues to be used in Pavlovian fear discrimination. Cues were auditory stimuli, 10-s in duration and consisted of repeating motifs of a broadband click, phaser, or trumpet. Stimuli can be heard or downloaded at http://mcdannaldlab.org/resources/ardbark. Previous studies have found these stimuli to be equally salient, yet highly discriminable (Berg et al., 2014; Wright et al., 2015; DiLeo et al., 2016; Ray et al., 2018). The 42-min pre-exposure sessions consisted of four presentations of each cue (12 total presentations) with a mean inter-trial interval (ITI) of 3.5 min. The order of trial type presentation was randomly determined by the behavioral program and differed for each rat during each session throughout behavioral testing.

For all sessions, fear to each auditory cue was measured using a suppression ratio based on nose poke rates during the 20-s baseline period immediately preceding the 10-s cue period: suppression ratio = (baseline nose poke rate – cue nose poke rate) / (baseline nose poke rate + cue nose poke rate). A ratio of 1 indicated complete suppression of nose poking during the cue and a high level of fear; 0, no suppression and no fear. Intermediate suppression ratios reflected intermediate fear levels. The same suppression ratio formula was used to calculate fear in 2-s cue intervals.

Fear discrimination

For the next 16 sessions, each rat underwent Pavlovian fear discrimination. Each 54-min session began with a ~5-min warm-up period during which no cues or shock were presented. Each cue was associated with a unique foot shock (0.5 mA, 0.5-s) probability: danger (1.00), uncertainty (0.25), and safety (0.00). Foot shock was administered 1-s following cue offset. A single session consisted of four danger, six uncertainty-no shock, two uncertainty-shock, and four safety trials. Auditory stimulus identity was counterbalanced across rats. Mean inter-trial interval was 3.5 min.

Histology

Upon the conclusion of behavior, rats were anesthetized with an overdose of isoflurane and perfused

intracardially with 0.9% biological saline. Brains were extracted and stored in 4% (v/v) formalin and 10% (w/v) sucrose. Forty-micrometer sections were collected on a sliding microtome. Tissue was then washed with PBS, incubated in NeuroTrace (Thermo Fisher, N21479) at a 1:200 concentration, washed again, mounted, dried, and coverslipped with Vectashield Hardset mounting media (Vector Labs, H-1400). Slides were imaged within 3 weeks of processing.

Experiment 2

Subjects

Subjects were 25 male Long Evans rats weighing 275-300 g upon arrival (Charles River Laboratories; RGD Cat# 2308852, RRID:RGD_2308852). Rats were individually housed and maintained on a 12-h dark light cycle (lights off at 6:00 PM) with water *ad libitum*. Procedures adhered to the NIH Guide for the Care and Use of Laboratory Animals and were approved by the Boston College Institutional Animal Care and Use Committee.

Behavioral apparatus

Behavioral apparatus was identical to experiment 1. In addition to the standard behavior apparatus, green (532 nm, 500 mW) lasers (Shanghai Laser & Optics Century Co., Ltd.; Shanghai, China) were used to illuminate the NAcc. Lasers were connected to the behavior cables via 1X2 fiber optic rotatory joints (Doric; Quebec, Canada). A ceramic sleeve maintained contact between the ferrules on the opto-genetic cable and the head cap. The ceramic sleeve/ferrule junction was shielded with black shrink wrap to block light emission into the behavioral chamber. A PM160 light meter (Thorlabs; Newton, NJ) was used to measure light output.

Optogenetic materials

Optical ferrules were constructed using 2.5mm ceramic zirconia ferrules (Precision Fiber Products; Chu-la Vista, CA). Behavior cables were custom made for light delivery (Multimode Fiber, 0.22 NA, High-OH, Ø200 μm Core). All protocols can be downloaded at http://mcdannaldlab.org/resources/optogenetics.

Surgical procedures

Stereotaxic surgery was performed under isoflurane anesthesia (2-5%) using aseptic technique. Thir-teen rats received bilateral infusions of AAV-hSyn-eNpHR3.0-EYFP (*halorhodopsin*) aimed at the nu-

cleus accumbens core (0.50 μl, +1.90 AP, ±1.80 ML, -6.60 DV at a 0° angle) and bilateral optical ferrules (+1.70 AP, ±2.80 ML, -6.00 DV at a 10° angle). Infusions were delivered via 2 μl syringe (Hamilton, Neuros) controlled by a microsyringe pump (World Precision Instruments, UMP3-2). Infusion rate was ~0.11 μl/min. At the completion of each infusion, the syringe was raised 0.1 mm then left in place for five min to encourage delivery to the target site. The remaining 12 rats received identical surgical treatment but received a control fluorophore instead (AAV-hSyn-EYFP). Implants were secured with dental cement surrounded by a modified, 50 mL centrifuge tube. Post-surgery, rats received 2 weeks of undisturbed recovery with prophylactic antibiotic treatment (cephalexin; Henry Schein 049167) before beginning nose poke acquisition. All rats received carprofen (5 mg/kg) for post-operative analgesia.

Pre-illumination training and cable habituation

Nose poke acquisition, pre-exposure and initial fear discrimination (10 sessions) were identical to Experiment 1. We increased the delay between cue offset and shock onset to 2 s to ensure that neural activity would not be inhibited during shock delivery. Cable habituation was provided in two consecutive sessions by plugging rats into optogenetic cables and administering fear discrimination without illumination. In total, rats received twelve fear discrimination sessions prior to receiving light illumination.

NAcc illumination

Rats received eight sessions of fear discrimination plus NAcc illumination. The NAcc was illuminated via bilateral delivery of 12.5 mW of 532 nm 'green' light: DPSS laser → optogenetic cables → implanted ferrules. There were two types of illumination sessions: cue and ITI. For cue sessions, light illumination began 0.5 s prior to cue onset and ended 0.5 s following cue offset, resulting in a total illumination time of 11 s. Light illumination was given for all trial types (danger, uncertainty and safety) for a total of 16 illumination events per session. For ITI sessions, illumination occurred during the inter-trial intervals between cue presentations. Illumination was roughly equidistant from previous cue offset and subsequent cue onset (~90 s from each). Sixteen ITI illumination events were administered, each lasting 11 s, meaning that total illumination time was equal for cue and ITI sessions. The within subjects design meant that each rat received four cue illumination sessions and four ITI illumination sessions. Illumination was given in two-session blocks, with half of the subjects starting with cue illumination.

Histology

Upon conclusion of behavior, rats were anesthetized with an overdose of isoflurane and perfused intracardially with 0.9% saline and 4% paraformaldehyde in a 0.2 M potassium phosphate buffered solution. Brains

were extracted and stored in 4% (v/v) formalin and 10% (w/v) sucrose. Forty-micrometer sections were collected on a sliding microtome. Tissue was washed with PBS, incubated in NeuroTrace (Thermo Fisher, N21479) at a 1:200 concentration, washed again, mounted, dried, and coverslipped with Vectashield Hard-set (Vector Labs, H-1400). Slides were imaged within 3 weeks of processing.

Figure 1. Neurotoxic lesion experimental timeline and histology. Rats received stereotaxic surgery, recovered, were trained to nose poke and received 16 sessions of Pavlovian fear discrimination. (**A**) Representative sham with NAcc intact (left) and lesion with NAcc damage (right) is shown. Dotted lines (left) show approximate NAcc location. Arrows (right) indicate gliosis and damage restricted to the NAcc. (**B**) The extent of neurotoxic NAcc lesions across four coronal planes is shown, and the anterior distance from bregma (millimeters) indicated. (**C**) Pavlovian fear discrimination consisted of three, 10-s cues predicting unique shock probability: danger (p=1.00), red; uncertainty (p=0.25), purple; and safety (p=0.00), blue. Each 10-s cue was divided into 5, 2-s intervals for rapid fear discrimination analyses. (**D**) Mean **± SEM** baseline nose poke rates for the sixteen fear discrimination sessions are shown for sham (black) and NAcc rats (gray).

Statistical analysis

Behavioral data were acquired using Med Associates Med-PC IV software (MED PC, RRID:SCR_012156). Raw data were processed in Matlab (MATLAB, RRID:SCR_001622) to extract time stamps for nose poke and cue onset. Suppression ratios were calculated as: (baseline poke rate – cue poke rate) / (baseline poke rake + cue poke rate) and were analyzed with repeated measures ANOVA in SPSS (RRID:SCR_002865). Repeated measures ANOVA was performed with factors of group, cue, and time (experiment 1) or group, cue, time, and illumination (experiment 2). Partial eta squared (η_p^2) and observed power (op) are reported for ANOVA results for indicators of effect size. For all analyses, $p<0.05$ (or an appropriate Bonferroni correction) was considered significant.

Results

Experiment 1

Histological Results

Rats received bilateral sham or neurotoxic NAcc lesions. Neurotoxic damage (cell loss and gliosis) was quantified for the NAcc. Twenty-four NAcc rats showed damage primarily in the nucleus accumbens core (>85%) with minor damage (~10%) in the accumbens shell. Shams showed no evidence of neurotoxic damage. Representative sham (Figure 1A, right), and NAcc lesion (Figure 1A, left) sections are shown. Each subject's lesion was drawn, made transparent, and stacked (Figure 1B). Darker areas indicate regions of greater overlap and more consistent damage. Rats fully recovered from surgery before receiving fear discrimination (Figure 1C).

Baseline Nose Poking

NAcc lesions altered the progression of nose poking over discrimination sessions but did not grossly reduce nose poke rates (Figure 1D). ANOVA for baseline nose poke rate with session (16) and group (sham vs. NAcc) as factors found a main effect of session ($F_{(15,645)} = 47.14$, $p=3.77 \times 10^{-93}$, $\eta_p^2 = 0.52$, op = 1.00), a session x group interaction ($F_{(15,645)} = 2.10$, $p=0.008$, $\eta_p^2 = 0.05$, op = 0.97) but no main effect of group ($F_{(1,43)} = 0.16$, $p=0.69$).The interaction was the result of sham rats showing greater increases in baseline nose poking from the first half to second half of discrimination. So although sham and NAcc rats showed equivalent nose pokes rates when the first and second half of discrimination were compared in isolation (independent samples t-test, $t_{43} < 1$, p>0.4), shams showed greater increases in poking from the first to second half of discrimination (independent samples t-test, $t_{43} = 2.19$, p=0.034). Mean ± SEM baseline nose

pokes rates for sessions 1-8 were: sham (28.44 ± 2.96) and NAcc (28.83 ± 1.97); sessions 9-16 were: sham (38.80 ± 3.62) and NAcc (35.33 ± 2.46).

Figure 2. Fear scaling in sham and NAcc-lesioned rats. Mean ± SEM suppression ratio for danger (red), uncertainty (purple), and safety (blue) are shown for (**A**) sham and (**B**) NAcc rats. The vertical lines separate the two pre-exposure and sixteen fear discrimination sessions. The gray rectangles indicate the last six discrimination sessions. Mean + SEM suppression ratios for the 5, 2-s cue intervals are shown for (**C**) sham and (**D**) NAcc rats. Cue color scheme maintained from A. Mean difference score for danger vs. uncertainty (D-U, red bar) and uncertainty vs. safety (U-S, purple bar) is shown for the first and last 2-s cue intervals for (**E**) sham and (**F**) NAcc rats. Data points show individual difference scores. *One-sample t-test compared to zero, $p<0.00625$.

Fear Scaling

Sham rats acquired appropriate scaling of the fear response over the 16 sessions (Figure 2A). Suppression ratios for the entire 10-s cue were low in pre-exposure and initially increased to all cues. As discrimination proceeded, the suppression ratio for each cue diverged: high to danger, intermediate to uncertainty, and low to safety. NAcc rats showed a similar progression, but poorer overall scaling (Figure 2B). In support of the general emergence of scaling, ANOVA [between factor: group (sham vs. NAcc); within factors: session (16) and cue (danger, uncertainty and safety)] revealed a main effect of cue ($F_{(2,86)}$ = 115.51, p=4.34 x 10^{-25},

$\eta_p^2 = 0.73$, op = 1.00) and a cue x session interaction ($F_{(30,1290)} = 14.05$, $p=6.30 \times 10^{-60}$, $\eta_p^2 = 0.25$, op = 1.00). Revealing impaired scaling in NAcc rats, ANOVA found a cue x group interaction ($F_{(2,86)} = 5.76$, $p=0.004$, $\eta_p^2 = 0.12$, op = 0.86). The cue x group interaction was also observed when only the last six sessions were analyzed ($F_{(2,86)} = 4.50$, $p=0.014$, $\eta_p^2 = 0.10$, op = 0.76) – the period by which scaling patterns were stable. Independent samples t-tests for the last six session mean found decreased suppression ratios to danger in NAcc rats compared to shams ($t_{43} = 2.86$, $p=0.006$), while uncertainty and safety suppression ratios did not differ by group ($t < 1.1$, $p>0.3$). However, the impairment in NAcc rats was not solely attributable to decreased fear to danger. Performing ANOVA for all 16 discrimination sessions, but only for uncertainty and safety suppression ratios, returned a significant cue x group interaction ($F_{(1,43)} = 1.02$, $p=0.028$, $\eta_p^2 = 0.11$, op = 0.60). NAcc rats did not discriminate uncertainty and safety as well as shams. These results reveal a general role for the NAcc in the acquisition of fear scaling.

Rapid Fear Scaling

We were interested in examining a possible role for the NAcc in the rapid emergence of fear scaling. To do this, we examined mean suppression ratios from the last six sessions. Each cue was divided into 5, 2-s cue intervals and suppression ratios were calculated for each cue/interval. Sham rats showed scaling of the response in the first 2-s cue interval and in all subsequent intervals (Figure 2C). Scaling was reduced across all intervals in NAcc rats (Figure 2D). In support, ANOVA [between factor: group (sham vs. NAcc); within factors: interval (5, 2-s cue intervals) and cue (danger, uncertainty and safety)] found a group x cue interaction ($F_{(2,86)} = 3.88$, $p=0.024$, $\eta_p^2 = 0.08$, op = 0.69). Supporting a specific role for the NAcc in rapid scaling of fear, NAcc rats showed impaired scaling even when only the first 2-s cue interval was analyzed (main effect of group; $F_{(2,86)} = 5.08$, $p=0.008$, $\eta_p^2 = 0.11$, op = 0.81). By contrast, no main effect of group was observed when the last 2-s cue interval was analyzed.

ANOVA revealed a role for the NAcc in rapid scaling of fear. To specify the nature of the deficit, we separated scaling into its component parts: danger vs. uncertainty and uncertainty vs. safety. We calculated difference scores for each cue pair in the first and last 2-s cue intervals. Sham rats showed positive difference scores to each cue pair at each interval. That is, difference scores exceeded zero, as revealed by one-sample t-test: first 1-s cue interval: danger vs. uncertainty ($t_{20} = 10.95$, $p=6.7 \times 10^{-4}$), uncertainty vs. safety ($t_{20} = 3.55$, $p=0.002$); last 2-s cue interval: danger vs. uncertainty ($t_{23} = 4.60$, $p=1.76 \times 10^{-4}$), uncertainty vs. safety ($t_{23} = 5.73$, $p=1.30 \times 10^{-5}$) for shams (Figure 2E). NAcc rats were selectively impaired at rapid uncertainty-safety discrimination. One-sample tests found that only the NAcc uncertainty-safety

difference score from the first 2-s cue interval failed to differ from zero (Figure 2F): first interval: danger vs. uncertainty (t_{23} = 4.20, p=3.38 x 10^{-4}), uncertainty vs. safety (t_{23} = 1.31, p=0.20); Last interval: danger vs. uncertainty (t_{20} = 5.22, p=2.70 x 10^{-5}), uncertainty vs. safety (t_{20} = 4.19, p=3.53 x 10^{-4}). All significant one-sample t tests survive Bonferroni correction (0.05/8, p<0.00625). Altogether, these results reveal a specific role for the NAcc in the acquisition of rapid uncertainty-safety discrimination.

Figure 3. Optogenetic inhibition experimental timeline, histology, and initial fear scaling. Rats received stereotaxic surgery, recovered, underwent nose poke acquisition, ten fear discrimination sessions, two cable habituation sessions, and eight light illumination sessions. (**A**) Representative NAcc transduction is shown with YFP expression (yellow fluorescent protein; yellow) and neurotrace (blue). Dotted lines indicate approximate NAcc location. (**B**) The extent of viral transduction across four coronal planes is shown for Halo (green, left) and YFP rats (yellow, right), and the anterior distance from bregma (millimeters) indicated. Individual ferrule placement is indicated in black circles. (**C**) Mean ± SEM suppression ratios for danger (red), uncertainty (purple), and safety (blue) are shown for YFP (left) and Halo rats (right) during the ten fear discrimination sessions. (**D**) In the final eight sessions, rats received NAcc light illumination during cue presentation (top) or during the inter-trial interval (ITI, bottom). Green indicates light illumination, yellow indicates shock delivery and candy-striped indicates cue presentation. (**E**) Cue and ITI illumination were given in alternating, two-session blocks. Block order was counterbalanced with roughly half of the subjects first receiving the cue illumination block.

Experiment 2

Experiment 1 results implicate the NAcc in two aspects of fear acquisition: overall scaling and a specific component of scaling: rapid uncertainty/safety discrimination. However, neurotoxic lesions altered baseline nose poking behavior and permanently ablated NAcc neurons. To determine a temporally specific role for the NAcc in fear expression, Experiment 2 took a within-subjects, optogenetic approach. Rats were NAcc-transducted with halorhodopsin or a control fluorophore and trained to discriminate danger, uncertainty and safety. Once scaling was established, rats received sessions in which the NAcc was illuminated during cue presentation or during the inter-trial interval (Figure 3D, E). If the NAcc plays identical roles in acquisition and expression of fear scaling, we would expect to observe a three-way interaction (group x stimulation x cue) with only halorhodopsin rats showing impaired overall scaling during cue illumination sessions. If NAcc plays a more selective role in the expression of rapid fear scaling, we would anticipate a four-way interaction (group x interval x stimulation x cue) with only halorhodopsin rats showing impaired rapid uncertainty-safety discrimination during cue illumination sessions.

Histological Results

Rats received bilateral NAcc transduction with halorhodopsin (Halo) or a control fluorophore (YFP) and bilateral optical ferrule implantation just above the NAcc. Representative transduction is shown (Figure 3A). Each subject's total transduction area was drawn, made transparent, and stacked (Figure 3B). Darker areas indicate regions of greater overlap and more consistent transduction. Transduction centered around and above the anterior commissure, the precise NAcc location.

Baseline Nose Poking

YFP and Halo rats showed equivalent baseline nose poking rates throughout pre-exposure, discrimination, cable habituation, and light illumination (Figure S1). ANOVA for baseline nose poke rate [factors: session (20) and group (YFP vs. Halo)] demonstrated a main effect of session ($F_{(19,437)}$ = 12.60, p=4.19 x 10^{-31}, η_p^2 = 0.35, op = 1.00), but no main effect or interaction with group (Fs < 0.93, ps>0.55). Equivalent performance lessens the concern that differences in suppression ratios between groups result from differences in baseline nose poke rates.

Initial Fear Scaling

YFP and Halo rats acquired reliable fear scaling over the 10 sessions (Figure 3C). Suppression ratios were low in pre-exposure and initially increased to all cues. As discrimination proceeded, the suppression ratio

for each cue diverged: high to danger, intermediate to uncertainty, and low to safety. Demonstrating overall scaling, ANOVA [within factors: session (10) and 10-s cue (danger, uncertainty and safety); between factor: group (YFP vs. Halo)] revealed a main effect of cue ($F_{(2.46)}$ = 36.21, p=3.58 x 10^{-10}, η_p^2 = 0.61, op = 1.00), session ($F_{(9,207)}$ = 25.74, p=2.04 x 10^{-29} η_p^2 = 0.53, op = 1.00) and a cue x session interaction ($F_{(18,414)}$ = 6.26, p=1.14 x 10^{-13} η_p^2 = 0.21, op = 1.00). ANOVA found no main effect or interaction with group (Fs < 3.42, ps>0.08). Thus, YFP and Halo entered the light illumination phase (Figure 3D) showing equivalent fear scaling.

Figure 4. Optogenetic inhibition and rapid fear scaling. Mean + SEM suppression ratios are plotted for the 5, 2-s cue intervals during danger (red), uncertainty (purple), and safety (blue). **A & B** show suppression ratios for cable habituation and ITI illumination sessions for YFP and Halo rats. **C & D** show suppression ratios for cue illumination sessions. (**E**) Mean difference scores for uncertainty vs. safety suppression ratios in Cue vs. ITI illumination sessions are shown for YFP (black) and Halo rats (green). Scores were calculated for the first (left) and last (right) 2-s cue intervals. Individual data points are shown in filled circles. Outliers are shown in open circles. (**F**) Mean + SEM suppression ratios are plotted for the 5, 2-s intervals during ITI illumination (green) and post-illumination period (gray) (YFP, left; Halo, right). *(green) one-sample t-test compared to zero, p=0.0038. *(black) independent samples t-test, p=0.0041.

Fear Scaling during Light Illumination

Suppression ratios are shown in 5, 2-s intervals for each cue during habituation/ITI illumination sessions [YFP rats (Figure 4A) and Halo rats (Figure 4B)]. Identical plots were made for cue illumination sessions (Figure 4C, D). To examine a possible role for the NAcc in rapid fear scaling, we performed ANOVA with all factors [within factors: session-type (cable habituation, ITI illumination and cue illumination), cue (danger, uncertainty and safety), and interval (5, 2-s cue intervals); between factor: group (YFP vs. Halo)]. The complete ANOVA output is reported in Table 1. Consistent with general scaling across groups, ANOVA revealed a main effect of cue ($F_{(2.46)}$ = 89.04, p=1.53 x 10^{-16}, η_p^2 = 0.80, op = 1.00) as well as a cue x interval interaction ($F_{(8.184)}$ = 6.14, p=5.16 x 10^{-7}, η_p^2 = 0.21, op = 1.00). Indicative of a selective role for the NAcc in rapid fear scaling, ANOVA revealed a significant 4-way interaction [session-type x cue x interval x group ($F_{(16,368)}$ = 1.80, p=0.029, η_p^2 = 0.07, op = 0.95)], but not a significant 3-way interaction [session-type x cue x group ($F_{(4,92)}$ = 1.35, p=0.26, η_p^2 = 0.06, op = 0.41)].

The 4-way interaction indicates that YFP and Halo rats showed differing temporal scaling patterns across the different session types. To begin to clarify the differing patterns, we split YFP and Halo rats and performed identical ANOVAs [within factors: session-type (habituation, ITI illumination and cue illumination), cue (danger, uncertainty and safety), and interval (5, 2-s cue intervals)]. Indicative of reliable scaling, ANOVA for YFP rats found a main effect of cue ($F_{(2,22)}$ = 47.71, p=1.0 x 10^{-10}, η_p^2 = 0.81, op = 1.00) and a cue x interval interaction ($F_{(8,88)}$ = 2.76, p=0.009, η_p^2 = 0.20, op = 0.92). Revealing no effect of illumination on the temporal pattern of fear scaling, the 3-way interaction (session-type x cue x interval) was not significant ($F_{(16,176)}$ = 0.59, p = 0.89, η_p^2 = 0.05, op = 0.39). ANOVA for Halo rats also found a main effect of cue ($F_{(2,24)}$ = 41.39, p=1.66 x 10^{-8}, η_p^2 = 0.78, op = 1.00) and a cue x interval interaction ($F_{(8,96)}$ = 4.07, p=3.36 x 10^{-4}, η_p^2 = 0.25, op = 0.99). Only now, ANOVA revealed a significant 3-way interaction (session-type x cue x interval; $F_{(16,192)}$ = 1.92, p = 0.021, η_p^2 = 0.14, op = 0.95). NAcc illumination only disrupted the temporal scaling pattern for Halo rats.

Visual examination of the data suggests that – similar to NAcc-lesioned rats – Halo rats receiving NAcc optogenetic inhibition during cue presentation were specifically impaired in rapid uncertainty-safety discrimination (Figure 4D). We first calculated differential suppression ratios for uncertainty vs. safety for the first and last 2-s cue intervals during cue and ITI illumination sessions. We then calculated a difference score for the two session-types (cue differential suppression ratio – ITI differential suppression ratio). Difference scores around zero would indicate equivalent uncertainty-safety discrimination during cue and ITI

illumination sessions. Negative difference scores would indicate *worse* uncertainty-safety discrimination during cue illumination sessions. Two individuals (1 YFP and 1 Halo) had first interval difference scores ±2 standard deviations beyond the group mean. The data for these individuals are shown (Figure 4E, open circles), but were not included in t-test analyses.

In the first 2-s cue interval, Halo rats showed impaired uncertainty-safety discrimination in cue illumination sessions, compared to ITI illumination (Figure 4E, left). This was supported by significant, negative shift of differences scores away from zero (one-sample t-test, t_{11} = -3.65, *p*=0.004). YFP differences scores did not differ from zero, indicating equivalent uncertainty-safety discrimination in cue and ITI illumination sessions (t_{10} = 1.22, *p*=0.25). Further, YFP and Halo difference scores from the first 2-s cue interval differed from one another (independent samples t-test, t_{21} = 3.22, p=0.004). Impaired uncertainty-safety discrimination in Halo rats receiving cue illumination was specific to the first 2-s cue interval. Identical analysis of the last 2-s cue interval found that difference scores did not differ from zero for YFP (one-sample t-test, t_{10} = -0.41, *p*=0.69) and Halo rats (one-sample t-test, t_{11} = 0.27, *p*=0.80) (Figure 4E, right). Moreover, differences scores were similar between the two groups (independent samples t-test, t_{21} = 0.48, p=0.64). Altogether, the results reveal that NAcc activity at the time of cue presentation is necessary to rapidly discriminate uncertainty and safety.

Of course, it is possible that NAcc optogenetic inhibition simply suppressed rewarded nose poking. In this case, impaired rapid fear scaling would be the byproduct of a general reduction in poking. To rule out this possibility, we examined nose poke suppression during light illumination in ITI sessions (Figure 4F). No cues were present during this period, allowing us to determine the effect of light illumination alone to suppress nose poking. The middle 10 s of the 11-s light illumination was divided into 5, 2-s cue intervals – exactly as was done for the cue illumination analyses. For comparison, we also sampled 10 s of nose poking 30-s following illumination offset. This post-illumination served as a control period to which light illumination could be compared. ANOVA [within factors: period (light and post) and interval (5, 2-s cue intervals); between factor: group (YFP vs. Halo)] revealed main effects of period ($F_{(1,23)}$ = 34.53, *p*=5 x 10^{-6}, η_p^2 = 0.60, op = 1.00) and interval ($F_{(4,92)}$ = 2.49, *p*=0.049, η_p^2 = 0.10, op = 0.69). Critically, ANOVA found no main effect or interaction with group (Fs < 1.10, *p*s>0.31). So while suppression ratios were higher during light illumination, this did not differ between YFP and Halo rats and was therefore not due to inhibition of NAcc activity.

Discussion

We set out to examine a role for the nucleus accumbens core in the acquisition and expression of fear scaling. Neurotoxic lesions revealed a general role for the NAcc in fear scaling, as well as a specific role in one component of scaling: rapid uncertainty-safety discrimination. Optogenetic inhibition revealed a specific role for NAcc cue activity in the expression of rapid, uncertainty-safety discrimination. The results reveal that the NAcc is an essential component of a neural circuit permitting fear to rapidly scale to degree of threat.

Before discussing our results more broadly, we must consider several limitations of our experimental design and results. First, our experiments only used male rats. Several studies have reported sex differences in danger-safety discrimination (Day et al., 2016; Foilb et al., 2018; Greiner et al., 2019). We find only modest sex differences in our fear procedure (Walker et al., 2018; Walker et al., 2019), suggesting similar neural circuits may be utilized across sexes. Of course, females and males may achieve similar performance through differing neural mechanisms. Relatedly, the form of the fear response may differ between sexes. Males may be biased towards more passive fear responses, such as freezing (Maren et al., 1994; Gupta et al., 2001; Gruene et al., 2015). Females may be biased towards more active fear responses, such as darting (Gruene et al., 2015; Greiner et al., 2019; Odynocki and Poulos, 2019). Passive and active fear responses can both potentially contribute to conditioned suppression. So if the NAcc mediates active fear responses, and our procedure is capturing suppression via active fear, there may be an even greater role for the NAcc in female fear discrimination. By contrast, if the NAcc mediates active fear responses, and conditioned suppression is more the result of passive fear, there may be a diminished role for the NAcc in female fear discrimination.

Another important consideration is that our dependent measure of fear is derived from the rate of rewarded nose poking. Conditioned suppression is a strength because it provides an objective measure of fear on multiple time scales. It is a potential weakness because the NAcc plays a well-established role in reward-seeking. Disrupting NAcc function can attenuate reward-related behavior in many settings (Corbit et al., 2001; Hall et al., 2001; Ito et al., 2004; Blaiss and Janak, 2009; Ambroggi et al., 2011; McDannald et al., 2011; McDannald et al., 2013), though this finding is not universal (Ramirez and Savage, 2007; Corbit and Balleine, 2011). In our first experiment, NAcc lesions slowed the increase of baseline nose poking over discrimination sessions and also impaired fear scaling. However, the temporal emergence of the deficits in reward-seeking and fear scaling did not align. The fear scaling deficit was apparent across all sessions

while the nose poking deficit only emerged in the later sessions.

Our second experiment better demonstrated independent roles for the NAcc in rapid fear scaling and reward-seeking. Light illumination during the cue period effectively impaired one component of scaling in NAcc-halorhodopsin rats but not NAcc-YFP rats: rapid uncertainty-safety discrimination. By contrast, light illumination during the inter-trial interval produced modest but equivalent reductions in nose poking for both groups. Optogenetic inhibition of the NAcc was insufficient to reduce rewarded nose poking. The failure of NAcc inhibition to suppress nose poking may seem odd. Mice will readily perform actions that channelrhodopsin-excite NAcc D1 and D2 cell types (Cole et al., 2018), and rats will perform actions exciting glutamatergic inputs to the NAcc via channelrhodopsin (Stuber et al., 2011; Britt et al., 2012). However, these studies demonstrate that NAcc activity is sufficient, but not necessary, to support reward-seeking. Further, prominent theories posit that reward-seeking initially depends on medial striatal structures, such as the NAcc. With further training, lateral striatal regions, such as the dorsolateral striatum, control reward-seeking (Gerdeman et al., 2003; Belin and Everitt, 2008; Corbit et al., 2012; Burton et al., 2015). In our second experiment, rats had extensive experience with nose poking by the time the NAcc was optogenetically inhibited. By this time, reward-seeking may have no longer been under NAcc control, yet the NAcc continued to contribute to rapid fear scaling.

A role for the NAcc in fear would be expected based on immediate early gene studies. Shock-associated cues and contexts reliably upregulate NAcc c-fos and zif268 (Beck and Fibiger, 1995; Campeau et al., 1997; Thomas et al., 2002). Although the NAcc contains information about shock-associated cues and contexts, specifying the role of the NAcc in fear has presented a considerable challenge. Parkinson and colleagues found that NAcc lesions impaired cued fear, but enhanced contextual fear (Parkinson et al., 1999). Taking a similar experimental approach, Levita and colleagues found that NAcc lesions had no impact on the acquisition or expression of cued fear, but impaired retention of contextual fear (Levita et al., 2002). Contemporary work by Haralambous and Westbrook found that inhibiting accumbens activity (core + shell) specifically impaired the acquisition, but not expression of contextual fear, and had no effect on cued fear (Haralambous and Westbrook, 1999). Even considering slightly different methodologies, it is difficult to reconcile these disparate results.

These are not the only conflicts in the literature. Schwienbacher and colleagues found that blocking NAcc activity with tetrodotoxin abolished acquisition, and impaired expression, of fear-potentiated startle (Schwienbacher et al., 2004). The very next year, Josselyn and colleagues utilized a variety of methods

to manipulate the NAc during fear-potentiated startle: lesion, agonizing dopamine and blocking glutamate. NAcc manipulation had no effect on any aspect of fear-potentiated startle (Josselyn et al., 2005). Since these initial studies, the NAcc has been implicated in a variety of fear-related processes. For example, the NAcc can modulate salience, regulating the ability of cues to enter into associations with shock (Iordanova et al., 2006b; Iordanova et al., 2006a; Iordanova, 2009). Human imaging studies have observed correlates of prediction error, a theoretical signal that strengthens or weakens cue-shock associations (Seymour et al., 2004; Delgado et al., 2008; Schiller et al., 2008; Li et al., 2011), consistent with a role for the NAcc in predictive learning (Li and McNally, 2015).

What can we make of the mixed NAcc fear literature? Though dissatisfying, one answer is that the NAcc must play multiple roles in fear. Genetically- and anatomically-defined NAcc neuron types may be linked to specific fear processes. Future work dissecting the NAcc in this way, as has been done in reward settings (Kupchik et al., 2015; Francis and Lobo, 2017; Tejeda et al., 2017), is likely to be fruitful. A more fulfilling answer might be that the behavioral procedures used in early studies did not isolate an essential NAcc fear function. The NAcc may not be necessary to demonstrate fear to certain threat or to withhold fear to certain safety. For example, the NAcc is not necessary to behaviorally discriminate contexts/cues associated with certain shock and certain safety (Antoniadis and McDonald, 2006; McDannald and Galarce, 2011; Piantadosi, 2017).

We propose that a necessary role for the NAcc in fear emerges when subjects are confronted with threats on a continuum from safety to danger. The NAcc is a core component of a neural circuit permitting the level of the fear response to scale to the degree of threat. Further, our results suggest general and specific roles for the NAcc in fear scaling. During acquisition, the NAcc is generally necessary for fear scaling for the duration of the encounter – in our case for the entirety of cue presentation. At the same time, the NAcc is specifically necessary for one component of fear scaling: rapid discrimination of uncertain threat and safety. Once a scaled fear response is acquired, the general role for the NAcc diminishes. However, the NAcc continues to play a specific role in rapidly discriminating uncertain threat and safety. Importantly, we are not claiming that fear scaling is *the* function of the NAcc in fear, but rather *a* function.

Environmental threats are not absolute, but exist on a continuum from safety to danger. Using a behavioral procedure that attempts to capture this continuum, we find the NAcc is essential to scale fear to degree of threat. Our results clarify at least one role for the NAcc in fear, yet much more work remains. There is evidence of altered NAcc structure and function in anxiety disorders and post-traumatic stress disorder (Cha

et al., 2014; Felmingham et al., 2014; Manning et al., 2015; Morey et al., 2017). Disrupted threat-safety discrimination in anxiety and stress disorders may be conceptualized as maladaptive fear scaling. Detailing NAcc function, and mapping a more complete neural circuit for fear scaling, may inform strategies to promote adaptive fear in anxiety and stress disorders.

References

Ambroggi F, Ghazizadeh A, Nicola SM, Fields HL (2011) Roles of nucleus accumbens core and shell in incentive-cue responding and behavioral inhibition. J Neurosci 31:6820-6830.

Antoniadis EA, McDonald RJ (2006) Fornix, medial prefrontal cortex, nucleus accumbens, and mediodorsal thalamic nucleus: Roles in a fear-based context discrimination task. Neurobiology of Learning and Memory 85:71-85.

Badrinarayan A, Wescott SA, Vander Weele CM, Saunders BT, Couturier BE, Maren S, Aragona BJ (2012) Aversive Stimuli Differentially Modulate Real-Time Dopamine Transmission Dynamics within the Nucleus Accumbens Core and Shell. Journal of Neuroscience 32:15779-15790.

Beck CH, Fibiger HC (1995) Conditioned fear-induced changes in behavior and in the expression of the immediate early gene c-fos: with and without diazepam pretreatment. Journal of Neuroscience 15:709-720.

Belin D, Everitt BJ (2008) Cocaine seeking habits depend upon dopamine-dependent serial connectivity linking the ventral with the dorsal striatum. Neuron 57:432-441.

Berg BA, Schoenbaum G, McDannald MA (2014) The dorsal raphe nucleus is integral to negative prediction errors in Pavlovian fear. European Journal of Neuroscience 40:3096-3101.

Blaiss CA, Janak PH (2009) The nucleus accumbens core and shell are critical for the expression, but not the consolidation, of Pavlovian conditioned approach. Behav Brain Res 200:22-32.

Britt JP, Benaliouad F, McDevitt RA, Stuber GD, Wise RA, Bonci A (2012) Synaptic and behavioral profile of multiple glutamatergic inputs to the nucleus accumbens. Neuron 76:790-803.

Burton AC, Nakamura K, Roesch MR (2015) From ventral-medial to dorsal-lateral striatum: Neural correlates of reward-guided decision-making. Neurobiology of Learning and Memory 117:51-59.

Campeau S, Falls WA, Cullinan WE, Helmreich DL, Davis M, Watson SJ (1997) Elicitation and reduction of fear: Behavioural and neuroendocrine indices and brain induction of the immediate-early gene c-fos. Neuroscience 78:1087-1104.

Cha J, Carlson JM, DeDora DJ, Greenberg T, Proudfit GH, Mujica-Parodi LR (2014) Hyper-Reactive Human Ventral Tegmental Area and Aberrant Mesocorticolimbic Connectivity in Overgeneralization of Fear in Generalized Anxiety Disorder. Journal of Neuroscience 34:5855-5860.

Cole SL, Robinson MJF, Berridge KC (2018) Optogenetic self-stimulation in the nucleus accumbens: D1 reward versus D2 ambivalence. PLoS One 13:e0207694.

Corbit LH, Balleine BW (2011) The general and outcome-specific forms of pavlovian-instrumental transfer are differentially mediated by the nucleus accumbens core and shell. J Neurosci 31:11786-11794.

Corbit LH, Muir JL, Balleine BW (2001) The role of the nucleus accumbens in instrumental conditioning: Evidence of a functional dissociation between accumbens core and shell. Journal of Neuroscience 21:3251-3260.

Corbit LH, Nie H, Janak PH (2012) Habitual Alcohol Seeking: Time Course and the Contribution of Subregions of the Dorsal Striatum. Biological Psychiatry 72:389-395.

Correia SS, McGrath AG, Lee A, Graybiel AM, Goosens KA (2016) Amygdala-ventral striatum circuit activation decreases long-term fear. Elife 5.

Cromwell HC, Schultz W (2003) Effects of expectations for different reward magnitudes on neuronal activity in primate striatum. J Neurophysiol 89:2823-2838.

Day HLL, Reed MM, Stevenson CW (2016) Sex differences in discriminating between cues predicting threat and safety. Neurobiology of Learning and Memory 133:196-203.

Delgado MR, Li J, Schiller D, Phelps EA (2008) The role of the striatum in aversive learning and aversive prediction errors. Philos Trans R Soc Lond B Biol Sci 363:3787-3800.

DiLeo A, Wright KM, McDannald MA (2016) Sub-second fear discrimination in rats: Adult impairment in adolescent heavy alcohol drinkers. Learning & Memory 23:618-622.

Duits P, Cath DC, Lissek S, Hox JJ, Hamm AO, Engelhard IM, van den Hout MA, Baas JM (2015) Updated meta-analysis of classical fear conditioning in the anxiety disorders. Depress Anxiety 32:239-253.

Fadok JP, Darvas M, Dickerson TMK, Palmiter RD (2010) Long-Term Memory for Pavlovian Fear Conditioning Requires Dopamine in the Nucleus Accumbens and Basolateral Amygdala. Plos One 5.

Felmingham KL, Falconer EM, Williams L, Kemp AH, Allen A, Peduto A, Bryant RA (2014) Reduced Amygdala and Ventral Striatal Activity to Happy Faces in PTSD Is Associated with Emotional Numbing. Plos One 9.

Foilb AR, Bals J, Sarlitto MC, Christianson JP (2018) Sex differences in fear discrimination do not manifest as differences in conditioned inhibition. Learning & Memory 25:49-53.

Francis TC, Lobo MK (2017) Emerging Role for Nucleus Accumbens Medium Spiny Neuron Subtypes in Depression. Biol Psychiatry 81:645-653.

Gerdeman GL, Partridge JG, Lupica CR, Lovinger DM (2003) It could be habit forming: drugs of abuse and striatal synaptic plasticity. Trends in Neurosciences 26:184-192.

Goosens KA, Maren S (2001) Contextual and auditory fear conditioning are mediated by the lateral, basal, and central amygdaloid nuclei in rats. Learning & Memory 8:148-155.

Greiner EM, Muller I, Norris MR, Ng KH, Sangha S (2019) Sex differences in fear regulation and reward-seeking behaviors in a fear safety-reward discrimination task. Behavioural Brain Research 368:131-139.

Gruene TM, Flick K, Stefano A, Shea SD, Shansky RM (2015) Sexually divergent expression of active and passive conditioned fear responses in rats. Elife 4.

Gupta RR, Sen S, Diepenhorst LL, Rudick CN, Maren S (2001) Estrogen modulates sexually dimorphic contextual fear conditioning and hippocampal long-term potentiation (LTP) in rats. Brain Research 888:356-365.

Hall J, Parkinson JA, Connor TM, Dickinson A, Everitt BJ (2001) Involvement of the central nucleus of the amygdala and nucleus accumbens core in mediating Pavlovian influences on instrumental behaviour. European Journal of Neuroscience 13:1984-1992.

Haralambous T, Westbrook RF (1999) An infusion of bupivacaine into the nucleus accumbens disrupts the acquisition but not the expression of contextual fear conditioning. Behavioral Neuroscience 113:925-940.

Iordanova MD (2009) Dopaminergic Modulation of Appetitive and Aversive Predictive Learning. Reviews in the Neurosciences 20:383-404.

Iordanova MD, Westbrook RF, Killcross AS (2006a) Dopamine activity in the nucleus accumbens modulates blocking in fear conditioning. European Journal of Neuroscience 24:3265-3270.

Iordanova MD, McNally GP, Westbrook RF (2006b) Opioid receptors in the nucleus accumbens regulate attentional learning in the blocking paradigm. Journal of Neuroscience 26:4036-4045.

Ito R, Robbins TW, Everitt BJ (2004) Differential control over cocaine-seeking behavior by nucleus accumbens core and shell. Nature Neuroscience 7:389-397.

Josselyn SA, Falls WA, Gewirtz JC, Pistell P, Davis M (2005) The nucleus accumbens is not critically involved in mediating the effects of a safety signal on behavior. Neuropsychopharmacology 30:17-26.

Jovanovic T, Kazama A, Bachevalier J, Davis M (2012) Impaired safety signal learning may be a biomarker of PTSD. Neuropharmacology 62:695-704.

Jovanovic T, Norrholm SD, Blanding NQ, Davis M, Duncan E, Bradley B, Ressler KJ (2010) Impaired fear inhibition is a biomarker of PTSD but not depression. Depress Anxiety 27:244-251.

Kita H, Kitai ST (1990) Amygdaloid Projections to the Frontal-Cortex and the Striatum in the Rat. Journal of Comparative Neurology 298:40-49.

Koo JW, Han JS, Kim JJ (2004) Selective neurotoxic lesions of basolateral and central nuclei of the amygdala produce differential effects on fear conditioning. Journal of Neuroscience 24:7654-7662.

Kupchik YM, Brown RM, Heinsbroek JA, Lobo MK, Schwartz DJ, Kalivas PW (2015) Coding the direct/indirect pathways by D1 and D2 receptors is not valid for accumbens projections. Nature Neuroscience 18:1230-+.

Levita L, Dalley JW, Robbins TW (2002) Disruption of Pavlovian contextual conditioning by excitotoxic lesions of the nucleus accumbens core. Behavioral Neuroscience 116:539-552.

Li J, Schiller D, Schoenbaum G, Phelps EA, Daw ND (2011) Differential roles of human striatum and amygdala in associative learning. Nat Neurosci 14:1250-1252.

Li SSY, McNally GP (2015) A Role of Nucleus Accumbens Dopamine Receptors in the Nucleus Accumbens Core, but Not Shell, in Fear Prediction Error. Behavioral Neuroscience 129:450-456.

Lissek S, Kaczkurkin AN, Rabin S, Geraci M, Pine DS, Grillon C (2014) Generalized Anxiety Disorder Is Associated With Overgeneralization of Classically Conditioned Fear. Biological Psychiatry 75:909-915.

Manning J, Reynolds G, Saygin ZM, Hofmann SG, Pollack M, Gabrieli JD, Whitfield-Gabrieli S (2015) Altered resting-state functional connectivity of the frontal-striatal reward system in social anxiety disorder. PLoS One 10:e0125286.

Maren S, De Oca B, Fanselow MS (1994) Sex differences in hippocampal long-term potentiation (LTP) and Pavlovian fear conditioning in rats: positive correlation between LTP and contextual learning. Brain Res 661:25-34.

McDannald MA, Galarce EM (2011) Measuring Pavlovian fear with conditioned freezing and conditioned suppression reveals different roles for the basolateral amygdala. Brain Research 1374:82-89.

McDannald MA, Setlow B, Holland PC (2013) Effects of ventral striatal lesions on first- and second-order appetitive conditioning. Eur J Neurosci 38:2589-2599.

McDannald MA, Lucantonio F, Burke KA, Niv Y, Schoenbaum G (2011) Ventral Striatum and Orbitofrontal Cortex Are Both Required for Model-Based, But Not Model-Free, Reinforcement Learning. Journal of Neuroscience 31:2700-2705.

McGinty VB, Lardeux S, Taha SA, Kim JJ, Nicola SM (2013) Invigoration of reward seeking by cue and proximity encoding in the nucleus accumbens. Neuron 78:910-922.

Morey RA, Davis SL, Garrett ME, Haswell CC, Marx CE, Beckham JC, McCarthy G, Hauser MA, Ashley-Koch AE, Workgrp M-AM (2017) Genome-wide association study of subcortical brain volume in

PTSD cases and trauma-exposed controls. Translational Psychiatry 7.

Odynocki N, Poulos AM (2019) Delving into darting. Lab Animal 48:204-205.

Oleson EB, Gentry RN, Chioma VC, Cheer JF (2012) Subsecond dopamine release in the nucleus accumbens predicts conditioned punishment and its successful avoidance. J Neurosci 32:14804-14808.

Ottenheimer D, Richard JM, Janak PH (2018) Ventral pallidum encodes relative reward value earlier and more robustly than nucleus accumbens. Nature Communications 9.

Parkinson JA, Olmstead MC, Burns LH, Robbins TW, Everitt BJ (1999) Dissociation in effects of lesions of the nucleus accumbens core and shell on appetitive pavlovian approach behavior and the potentiation of conditioned reinforcement and locomotor activity by D-amphetamine. Journal of Neuroscience 19:2401-2411.

Petrovich GD, Risold PY, Swanson LW (1996) Organization of projections from the basomedial nucleus of the amygdala: a PHAL study in the rat. J Comp Neurol 374:387-420.

Piantadosi PT (2017) Contributions of nucleus accumbens circuitry to aspects of aversively-motivated behaviors. In: Psychology, p 235: University of British Columbia.

Quirk GJ, Repa C, LeDoux JE (1995) Fear conditioning enhances short-latency auditory responses of lateral amygdala neurons: parallel recordings in the freely behaving rat. Neuron 15:1029-1039.

Ramirez DR, Savage LM (2007) Differential involvement of the basolateral amygdala, orbitofrontal cortex, and nucleus accumbens core in the acquisition and use of reward expectancies. Behav Neurosci 121:896-906.

Ray MH, Hanlon E, McDannald MA (2018) Lateral orbitofrontal cortex partitions mechanisms for fear regulation and alcohol consumption. PLoS One 13:e0198043.

Rescorla RA (1968) Probability of shock in the presence and absence of CS in fear conditioning. Journal of Comparative and Physiological Psychology 66:1-5.

Saddoris MP, Carelli RM (2014) Cocaine self-administration abolishes associative neural encoding in the nucleus accumbens necessary for higher-order learning. Biol Psychiatry 75:156-164.

Schiller D, Levy I, Niv Y, LeDoux JE, Phelps EA (2008) From Fear to Safety and Back: Reversal of Fear in the Human Brain. Journal of Neuroscience 28:11517-11525.

Schwienbacher I, Fendt M, Richardson R, Schnitzler HU (2004) Temporary inactivation of the nucleus accumbens disrupts acquisition and expression of fear-potentiated startle in rats. Brain Research 1027:87-93.

Setlow B, Schoenbaum G, Gallagher M (2003) Neural encoding in ventral striatum during olfactory discrimination learning. Neuron 38:625-636.

Seymour B, O'Doherty JP, Dayan P, Koltzenburg M, Jones AK, Dolan RJ, Friston KJ, Frackowiak RS (2004) Temporal difference models describe higher-order learning in humans. Nature 429:664-667.

Stuber GD, Sparta DR, Stamatakis AM, van Leeuwen WA, Hardjoprajitno JE, Cho S, Tye KM, Kempadoo KA, Zhang F, Deisseroth K, Bonci A (2011) Excitatory transmission from the amygdala to nucleus accumbens facilitates reward seeking. Nature 475:377-380.

Sugam JA, Saddoris MP, Carelli RM (2014) Nucleus accumbens neurons track behavioral preferences and reward outcomes during risky decision making. Biol Psychiatry 75:807-816.

Tejeda HA, Wu J, Kornspun AR, Pignatelli M, Kashtelyan V, Krashes MJ, Lowell BB, Carlezon WA, Bonci A (2017) Pathway-and Cell-Specific Kappa-Opioid Receptor Modulation of Excitation-Inhibition Balance Differentially Gates D1 and D2 Accumbens Neuron Activity. Neuron 93:147-163.

Thomas KL, Hall J, Everitt BJ (2002) Cellular imaging with zif268 expression in the rat nucleus accumbens and frontal cortex further dissociates the neural pathways activated following the retrieval of contextual and cued fear memory. European Journal of Neuroscience 16:1789-1796.

Walker RA, Andreansky C, Ray MH, McDannald MA (2018) Early adolescent adversity inflates threat estimation in females and promotes alcohol use initiation in both sexes. Behav Neurosci 132:171-182.

Walker RA, Wright KM, Jhou TC, McDannald MA (2019) The ventrolateral periaqueductal gray updates fear via positive prediction error. Eur J Neurosci.

Wright CI, Groenewegen HJ (1996) Patterns of overlap and segregation between insular cortical, intermediodorsal thalamic and basal amygdaloid afferents in the nucleus accumbens of the rat. Neuroscience 73:359-373.

Wright KM, DiLeo A, McDannald MA (2015) Early adversity disrupts the adult use of aversive prediction errors to reduce fear in uncertainty. Front Behav Neurosci 9:227.

Table 1. Experiment 2 fear discrimination during light illumination ANOVA results

Effect	F	p	η_{p2}	op
group	1.940	0.177	0.078	0.266
session-type	10.639	**$1.59 \times 10_{-4}$**	0.316	0.985
session-type x group	2.219	0.120	0.088	0.430
cue	89.044	**$1.53 \times 10_{-16}$**	0.795	1.000
cue x group	0.571	0.569	0.024	0.139
interval	6.00	**$2.46 \times 10_{-4}$**	0.207	0.982
interval x group	1.367	0.252	0.056	0.411
session-type x cue	4.213	**0.004**	0.155	0.912
session-type x cue x group	1.346	0.259	0.055	0.405
session-type x interval	0.708	0.684	0.030	0.323
session-type x interval x group	1.463	0.174	0.060	0.649
cue x interval	6.139	**$5.16 \times 10_{-7}$**	0.211	1.000
cue x interval x group	0.639	0.744	0.027	0.291
session-type x cue x interval	0.777	0.712	0.033	0.538
session-type x cue x interval x group	1.803	**0.029**	0.073	0.945

Figure S1. Baseline nose poking.
Mean +/- SEM nose poke rate (y-axis) is shown YFP (black) and Halo rats (green) during the 10 pre-illumination, 2 cable habituation (C) and 8 illlumination (I) sessions. Repeated measures ANOVA (between factor: group; within factor: session) found neither a main effect of group nor a group x session interaction.