The nucleus accumbens core is necessary to scale fear to degree of threat

Madelyn H. Ray, Alyssa N. Russ, Rachel A. Walker, and Michael A. McDannald

Department of Psychology & Neuroscience, Boston College, Chestnut Hill, MA 02467

Correspondence and requests for materials should be addressed to:

Michael A. McDannald
Department of Psychology & Neuroscience
Boston College
140 Commonwealth Ave.
514 McGuinn Hall
Chestnut Hill, MA 02467, USA
michael.mcdannald@bc.edu

or

Madelyn H. Ray
Department of Psychology & Neuroscience
Boston College
140 Commonwealth Ave.
Chestnut Hill, MA 02467, USA
madelyn.ray@bc.edu

Figures: 5
Tables: 2
Pages: 30
Words
Abstract: 169
Introduction: 492
Discussion: 1283

Conflict of interest: The authors declare no competing financial interests.
Acknowledgements: We would like to thank Alexa LaBanca and Andrew Thomson for technical support. This work was supported by NIH DA034010 to M.A.M. We thank Bret Judson and the Boston College Imaging Core for infrastructure and support.

Abstract

Fear is adaptive when the level of the response rapidly scales to degree of threat. Using a discrimination procedure consisting of danger, uncertainty and safety cues, we have found rapid fear scaling (within two seconds of cue presentation) in male rats. Here we examined a possible role for the nucleus accumbens core (NAcc) in the acquisition and expression of fear scaling. In experiment 1, male Long Evans rats received bilateral sham or neurotoxic NAcc lesions, recovered and underwent fear discrimination. NAcc-lesioned rats were generally impaired in scaling fear to degree of threat, and specifically impaired in rapid uncertainty-safety discrimination. In experiment 2, male Long Evans rats received NAcc transduction with halorhodopsin or a control fluorophore. After fear scaling was established, the NAcc was illuminated during cue or control periods. NAcc-halorhodopsin rats receiving cue illumination were specifically impaired in rapid uncertainty-safety discrimination. The results reveal a general role for the NAcc in scaling fear to degree of threat, and a specific role in rapid discrimination of uncertain threat and safety.

Significance Statement

Rapidly discriminating cues for threat and safety is essential for survival and impaired threat-safety discrimination is a hallmark of stress and anxiety disorders. In two experiments, we induced nucleus accumbens core (NAcc) dysfunction in rats receiving fear discrimination consisting of cues for danger, uncertainty and safety. Permanent NAcc dysfunction, via neurotoxic lesion, generally disrupted the ability to scale fear to degree of threat, and specifically impaired one component of scaling: rapid discrimination of uncertain threat and safety. Reversible NAcc dysfunction, via optogenetic inhibition, specifically impaired rapid discrimination of uncertain threat and safety. The results reveal that the NAcc is essential to scale fear to degree of threat, and is a plausible source of dysfunction in stress and anxiety disorders.

Introduction

The ability to discriminate danger from safety is critical to survival. Individuals with stress and anxiety disorders are impaired in discrimination, showing excessive fear-related responses to safety (Jovanovic et al., 2010; Jovanovic et al., 2012; Lissek et al., 2014; Duits et al., 2015). Danger and safety represent extremes of a threat continuum, with most real-world threats involving uncertainty. Ideally, one's level of fear should scale to the degree of threat. A scaled fear response would be most adaptive if it was rapidly organized following encounter with a potential threat.

Drawing from learning theory (Rescorla, 1968), our laboratory devised a discrimination procedure in which distinct auditory cues predict unique foot shock probabilities: danger ($p=1.00$), uncertainty ($p=0.25$), and safety ($p=0.00$) (Berg et al., 2014). Using this procedure, we have found that fear level scales to shock probability within two seconds of cue presentation (DiLeo et al., 2016). Present work in our laboratory seeks to identify brain regions necessary for fear scaling and its rapid emergence. Candidate regions should be able to process valence and receive amygdalar input (Quirk et al., 1995; Goosens and Maren, 2001; Koo et al., 2004; McDannald and Galarce, 2011). We identified the nucleus accumbens core (NAcc) as a likely candidate, based on its ability to rapidly process reward-predictive cues (Cromwell and Schultz, 2003; Setlow et al., 2003; Ambroggi et al., 2011; McGinty et al., 2013; Saddoris and Carelli, 2014; Sugam et al., 2014; Ottenheimer et al., 2018), as well as its anatomical connectivity with the amygdala (Kita and Kitai, 1990; Petrovich et al., 1996; Wright and Groenewegen, 1996). Even more, the NAcc is implicated in a variety of fear-related processes (Haralambous and Westbrook, 1999; Thomas et al., 2002; Schwienbacher et al., 2004; Iordanova et al., 2006b; Fadok et al., 2010; Badrinarayan et al., 2012; Oleson et al., 2012; Li and McNally, 2015; Correia et al., 2016).

In two experiments, we examined roles for the NAcc in fear scaling. In experiment 1, we

permanently ablated NAcc neurons via neurotoxic lesion. Following recovery, rats received fear

discrimination consisting of danger, uncertainty, and safety cues. Fear was measured with

suppression of rewarded nose poking (Estes and Skinner, 1941; Bouton and Bolles, 1980).

Examining suppression over the entire 10-s cue permitted analysis of overall fear scaling. To

examine the temporal emergence of scaling, we divided the 10-s cues into five, 2-s cue

intervals. Focusing on suppression during the first 2-s cue interval permitted analysis of rapid

fear scaling. In experiment 2, we examined a role for the NAcc in the expression of fear scaling.

Rats were NAcc-transducted with halorhodopsin or a control fluorophore, and bilaterally

implanted with ferrules above the NAcc. Following recovery, rats received fear discrimination

until fear scaling was stable. Over the next eight sessions, the NAcc was green-light illuminated

during cue presentation or a control period, optogenetically inhibiting activity in halorhodopsin

rats. The two experiments allowed us to examine roles for the NAcc in the acquisition and

expression of fear scaling.

Materials and methods

Experiment 1

Subjects

Subjects were forty-five male Long Evans rats weighing 275-300 g upon arrival (Charles River

Laboratories; RGD Cat# 2308852, RRID:RGD_2308852). Rats were individually housed and

maintained on a 12-h dark light cycle (lights off at 6:00 PM) with water *ad libitum*. Procedures

adhered to the NIH Guide for the Care and Use of Laboratory Animals and were approved by

the Boston College Institutional Animal Care and Use Committee.

Behavioral apparatus

Eight sound-attenuated enclosures each housed a behavior chamber with aluminum front and

back walls, clear acrylic sides and top, and a metal grid floor. Grid floors were electrically

connected to a shock generator. A single external food cup and central nose poke opening

equipped with infrared photocells were present on one wall. Auditory stimuli were presented

through two speakers mounted on the ceiling of each behavior chamber.

Surgical procedures

Stereotaxic surgery was performed under isoflurane anesthesia (2-5%) using aseptic technique.

Twenty-four rats received bilateral infusions of *N*-Methyl-D-aspartic acid (15 µg/µl in Dulbecco's

PBS) aimed at the nucleus accumbens core (0.40 µl, +1.90 AP, ±1.80 ML, -6.60 DV from skull).

Infusions were delivered via 2 µl syringe (Hamilton, Neuros) controlled by a microsyringe pump

(World Precision Instruments, UMP3-2). Infusion rate was ~0.11 µl/min. Thirty seconds after the

completion of each infusion, the syringe was raised 0.1 mm then left in place for five minutes to

encourage delivery to the target site. The remaining twenty-one rats received identical surgical

treatment without infusions. Rats received carprofen (5 mg/kg) for post-operative analgesia.

Nose poke acquisition

Following recovery from surgery, rats were food restricted to 85% of their initial free feeding

body weight, then fed (2 - 20 g/day) to increase their target body weight by 1 g/day for the

remainder of testing. Rats were shaped to nose poke for pellet (BioServ F0021 –

protein/fat/carbohydrate blend) delivery using a fixed ratio 1 schedule: one nose poke yielded

one pellet. Shaping sessions lasted 30 min or approximately 50 nose pokes. Over the next 3,

60-min sessions, rats were placed on variable interval (VI) schedules in which nose pokes were

reinforced on average every 30 s (session 1), or 60 s (sessions 2 and 3). For the remainder of

testing, nose pokes were reinforced on a VI-60 schedule independent of all Pavlovian

contingencies.

Pre-exposure

In two separate sessions, each rat was pre-exposed to the three cues to be used in Pavlovian

fear discrimination. Cues were auditory stimuli, 10-s in duration and consisted of repeating

motifs of a broadband click, phaser, or trumpet. Stimuli can be heard or downloaded at

http://mcdannaldlab.org/resources/ardbark. Previous studies have found these stimuli to be

equally salient, yet highly discriminable (Berg et al., 2014; Wright et al., 2015; DiLeo et al., 2016;

Ray et al., 2018). The 42-min pre-exposure sessions consisted of four presentations of each

cue (12 total presentations) with a mean inter-trial interval (ITI) of 3.5 min. The order of trial type

presentation was randomly determined by the behavioral program and differed for each rat

during each session throughout behavioral testing.

For all sessions, fear to each auditory cue was measured using a suppression ratio based on

nose poke rates during the 20-s baseline period immediately preceding the 10-s cue period:

suppression ratio = (baseline nose poke rate – cue nose poke rate) / (baseline nose poke rate +

cue nose poke rate). A ratio of 1 indicated complete suppression of nose poking during the cue

and a high level of fear; 0, no suppression and no fear. Intermediate suppression ratios reflected

intermediate fear levels. The same suppression ratio formula was used to calculate fear in 2-s

cue intervals.

Fear discrimination

Each rat received sixteen, 54-min Pavlovian fear discrimination sessions. Sessions began with

a ~5-min warm-up period during which no cues or shock were presented. The three cues were

associated with a unique foot shock (0.5 mA, 0.5-s) probability: danger (1.00), uncertainty

(0.25), and safety (0.00). Foot shock was administered 1-s following cue offset. A single session

consisted of four danger, six uncertainty omission, two uncertainty shock, and four safety trials.

153 Auditory stimulus identity was counterbalanced across rats. Mean inter-trial interval was 3.5

154 min.

Histology

Upon the conclusion of behavior, rats were anesthetized with an overdose of isoflurane and

perfused intracardially with 0.9% biological saline. Brains were extracted and stored in 4% (v/v)

formalin and 10% (w/v) sucrose. Forty-micrometer sections were collected on a sliding

microtome. Tissue was then washed with PBS, incubated in NeuroTrace (Thermo Fisher,

N21479) at a 1:200 concentration, washed again, mounted, dried, and coverslipped with

Vectashield Hardset mounting media (Vector Labs, H-1400). Slides were imaged within 3 weeks

of processing.

Experiment 2

Subjects

Subjects were 25 male Long Evans rats weighing 275-300 g upon arrival (Charles River

Laboratories; RGD Cat# 2308852, RRID:RGD_2308852). Rats were individually housed and

maintained on a 12-h dark light cycle (lights off at 6:00 PM) with water *ad libitum*. Procedures

adhered to the NIH Guide for the Care and Use of Laboratory Animals and were approved by

the Boston College Institutional Animal Care and Use Committee.

Behavioral apparatus

Behavioral apparatus was identical to experiment 1. In addition to the standard behavior

apparatus, green lasers (532 nm, max 500 mW; Shanghai Laser & Optics Century Co., Ltd.;

Shanghai, China) were used to illuminate the NAcc. Lasers were connected to the behavior

cables via 1X2 fiber optic rotary joints (Doric; Quebec, Canada). A ceramic sleeve maintained

contact between the ferrules on the optogenetic cable and the head cap. The ferrule junction

was shielded with black shrink wrap to block light emission into the behavioral chamber. A

PM160 light meter (Thorlabs; Newton, NJ) was used to measure light output.

Optogenetic materials

Optical ferrules were constructed using 2.5mm ceramic zirconia ferrules (Precision Fiber

Products; Chula Vista, CA). Behavior cables were custom made for light delivery (Multimode

Fiber, 0.22 NA, High-OH, Ø200 µm Core). All protocols can be downloaded at

http://mcdannaldlab.org/resources/optogenetics.

Surgical procedures

Stereotaxic surgery was performed under isoflurane anesthesia (2-5%) using aseptic technique.

Thirteen rats received bilateral infusions of AAV-hSyn-eNpHR3.0-EYFP (*halorhodopsin*) aimed

at the nucleus accumbens core (0.50 µl, +1.90 AP, ±1.80 ML, -6.60 DV at a 0° angle) and

bilateral optical ferrules (+1.70 AP, ±2.80 ML, -6.00 DV at a 10° angle). Infusions were delivered

via 2 µl syringe (Hamilton, Neuros) controlled by a microsyringe pump (World Precision

Instruments, UMP3-2). Infusion rate was ~0.11 µl/min. The syringe was raised 0.1 mm after

each infusion, then left in place for five min to encourage delivery to the target site. The

remaining 12 rats received identical surgical treatment but were infused with a control

fluorophore (AAV-hSyn-EYFP). Implants were secured with dental cement surrounded by a

modified, 50 mL centrifuge tube. Post-surgery, rats received 2 weeks of undisturbed recovery

with prophylactic antibiotic treatment (cephalexin; Henry Schein 049167) before beginning nose

poke acquisition. All rats received carprofen (5 mg/kg) for post-operative analgesia.

Pre-illumination training and cable habituation

Nose poke acquisition, pre-exposure and initial fear discrimination (10 sessions) were identical

to Experiment 1. We increased the delay between cue offset and shock onset to 2 s to ensure

that neural activity would not be inhibited during shock delivery. Cable habituation was provided

in two consecutive sessions by plugging rats into optogenetic cables and administering fear

discrimination without illumination. In total, rats received twelve fear discrimination sessions

prior to receiving light illumination.

NAcc illumination

Rats received eight sessions of fear discrimination plus NAcc illumination. The NAcc was

illuminated via bilateral delivery of 12.5 mW of 532 nm 'green' light: DPSS laser → optogenetic

cables → implanted ferrules. There were two types of illumination sessions: cue and ITI. For

cue sessions, light illumination began 0.5 s prior to cue onset and ended 0.5 s following cue

offset, resulting in a total illumination time of 11 s. Light illumination was given for all trial types

(danger, uncertainty and safety) for a total of 16 illumination events per session. For ITI

sessions, illumination occurred during the inter-trial intervals between cue presentations.

Illumination was roughly equidistant from previous cue offset and subsequent cue onset (~90 s

from each). Sixteen ITI illumination events were administered, each lasting 11 s, equating total

illumination time for cue and ITI sessions. The within-subjects design meant that each rat

received four cue illumination sessions and four ITI illumination sessions. Illumination was given

in two-session blocks, with half of the subjects starting with cue illumination.

Histology

After behavioral testing ended, rats were anesthetized with an overdose of isoflurane and

perfused intracardially with 0.9% biological saline and 4% paraformaldehyde in a 0.2 M

potassium phosphate buffered solution. Brains were extracted and stored in 4% (v/v) formalin

and 10% (w/v) sucrose. Forty-micrometer sections were collected on a sliding microtome.

Tissue was rinsed, incubated in NeuroTrace (Thermo Fisher, N21479) at a 1:200 concentration,

rinsed again, mounted, dried, and coverslipped with Vectashield Hardset (Vector Labs, H-1400).

Slides were imaged within 3 weeks of processing.

Statistical analysis

Behavioral data were acquired using Med Associates Med-PC IV software (MED PC,

RRID:SCR_012156). Raw data were processed in Matlab (MATLAB, RRID:SCR_001622) to

extract time stamps for nose poke and cue onset. Suppression ratios were calculated as:

(baseline poke rate – cue poke rate) / (baseline poke rake + cue poke rate) and were analyzed

with repeated measures ANOVA in SPSS (RRID:SCR_002865). Repeated measures ANOVA

was performed with factors of group, cue, and time (experiment 1) or group, cue, time, and

illumination (experiment 2). Partial eta squared (η_{p2}) and observed power (op) are reported for

ANOVA results for indicators of effect size. For all analyses, $p<0.05$ (or an appropriate

Bonferroni correction) was considered significant.

Results

Experiment 1

Histological Results

Rats received bilateral sham or neurotoxic NAcc lesions. Neurotoxic damage (cell loss and

gliosis) was quantified. Twenty-four NAcc rats showed damage primarily in the NAcc (>85%)

with minor damage (~10% or less) in the neighboring accumbens shell. Shams showed no

evidence of neurotoxic damage. Representative sham (Figure 1A, left), and NAcc lesion (Figure

1A, right) sections are shown. Each subject's lesion was drawn, made transparent, and stacked

(Figure 1B). Darker areas indicate regions of greater overlap and more consistent damage. Rats

fully recovered from surgery before receiving fear discrimination (Figure 1C).

Baseline Nose Poking

NAcc lesions altered the progression of nose poking over discrimination sessions, but did not

grossly reduce nose poke rates (Figure 1D). ANOVA for baseline nose poke rate with session

(16) and group (sham vs. NAcc) as factors found a main effect of session ($F_{(15,645)} = 47.14$,

248 p=3.77 x 10^{-93}, η_{p2} = 0.52, op = 1.00), a session x group interaction ($F_{(15,645)}$ = 2.10, p=0.008, η_{p2}

249 = 0.05, op = 0.97) but no main effect of group ($F_{(1,43)}$ = 0.16, p=0.69, η_{p2} = 0.004, op = 0.07).

250 Dividing the 16 sessions into 2, 8-session blocks; ANOVA found a block x group interaction

251 ($F_{(1,43)}$ = 4.81, p=0.034, η_{p2} = 0.10, op = 0.57). While sham (t_{20} = 7.69, p=2.13 x 10^{-7}) and NAcc

252 rats (t_{23} = 5.63, p=1.00 x 10^{-5}) both increased poking from the first to second half of

253 discrimination, sham rats showed greater increases (Figure 1E). Mean ± SEM baseline nose

254 pokes rates for sessions 1-8: sham (28.44 ± 2.96) and NAcc (28.83 ± 1.97); sessions 9-16:

255 sham (38.80 ± 3.62) and NAcc (35.33 ± 2.46; Figure 1E).

256 **Fear Scaling**

257 Sham rats acquired appropriate scaling of the fear response over the 16 sessions (Figure 2A,

258 left). Suppression ratios for the entire 10-s cue were low in pre-exposure and initially increased

259 to all cues. As discrimination proceeded, the suppression ratio for each cue diverged: high to

260 danger, intermediate to uncertainty, and low to safety. NAcc rats showed a similar progression,

261 but poorer overall scaling (Figure 2A, right). In support of the general emergence of scaling,

262 ANOVA [between factor: group (sham vs. NAcc); within factors: session (16) and cue (danger,

263 uncertainty and safety)] revealed a main effect of cue ($F_{(2,86)}$ = 115.51, p=4.34 x 10^{-25}, η_{p2} = 0.73,

264 op = 1.00) and a cue x session interaction ($F_{(30,1290)}$ = 14.05, p=6.30 x 10^{-60}, η_{p2} = 0.25, op =

265 1.00). Revealing impaired scaling in NAcc rats, ANOVA found a cue x group interaction ($F_{(2,86)}$ =

266 5.76, p=0.004, η_{p2} = 0.12, op = 0.86). The cue x group interaction was also observed when only

267 the last six sessions were analyzed ($F_{(2,86)}$ = 4.50, p=0.014, η_{p2} = 0.10, op = 0.76), the period by

268 which scaling patterns were stable.

269 To further reveal the deficit in NAcc rats, we focused on suppression ratios from the final six

270 sessions. Difference scores were calculated for the two components of scaling: (danger –

271 uncertainty) and (uncertainty – safety). Sham (Figure 2B, left) and NAcc rats (Figure 2B, right)

discriminated each cue pair. One-sample t-tests found that difference scores exceeded zero for

each comparison: sham, danger vs. uncertainty (t_{20} = 10.25, p=2.07 x 10^{-9}), uncertainty vs.

safety (t_{20} = 6.11, p=4.17 x 10^{-8}); NAcc, danger vs. uncertainty (t_{23} = 8.01, p=0.001), uncertainty

vs. safety (t_{23} = 3.65, p=0.002). However, NAcc rats showed poorer overall discrimination.

ANOVA [between factor: group (sham vs. NAcc); within factor: discrimination (danger –

uncertainty) and (uncertainty – safety)] revealed a main effect of group ($F_{(1,43)}$ = 5.68, p=0.022,

η_{p2} = 0.12, op = 0.64). Difference scores were reduced across both components in NAcc rats.

These results reveal a general role for the NAcc in fear scaling.

Rapid Fear Scaling

We were interested in revealing a possible role for the NAcc in the rapid emergence of fear

scaling. To do this, we examined mean suppression ratios from the last six sessions. Each cue

was divided into 5, 2-s cue intervals and suppression ratios were calculated for each

cue/interval. Sham rats showed scaling of the fear response in the first 2-s cue interval and in all

subsequent intervals (Figure 2C, left). Scaling was reduced across all 2-s cue intervals in NAcc

rats (Figure 2C, right). ANOVA [between factor: group (sham vs. NAcc); within factors: interval

(5, 2-s cue intervals) and cue (danger, uncertainty and safety)] found a group x cue interaction

($F_{(2,86)}$ = 3.88, p=0.024, η_{p2} = 0.08, op = 0.69). Supporting a specific role for the NAcc in rapid

fear scaling, NAcc rats showed impaired scaling even when only the first 2-s cue interval was

analyzed (cue x group interaction; $F_{(2,86)}$ = 5.08, p=0.0008, η_{p2} = 0.11, op = 0.81). No cue x

group interaction was observed when the last 2-s cue interval was analyzed ($F_{(2,86)}$ = 1.90,

p=0.16, η_{p2} = 0.04, op = 0.39).

To specify the nature of the deficit in NAcc rats, we reduced scaling into its component parts:

(danger – uncertainty) and (uncertainty – safety). We calculated difference scores for the first

and last 2-s cue intervals. Sham rats showed positive difference scores for each cue pair at

each interval (Figure 2D, left). Difference scores exceeded zero, as revealed by one-sample t-

tests: first 2-s cue interval: danger vs. uncertainty (t_{20} = 10.95, p=6.7 x 10^{-4}), uncertainty vs.

safety (t_{20} = 3.55, p=0.002); last 2-s cue interval: danger vs. uncertainty (t_{23} = 4.60, p=1.76 x 10^{-4}),

uncertainty vs. safety (t_{23} = 5.73, p=1.30 x 10^{-5}) for shams. NAcc rats were generally impaired

at rapid scaling. ANOVA for the first 2-s cue interval differences revealed a main effect of group

($F_{(1,43)}$ = 6.50, p=0.014, η_{p2} = 0.01, op = 0.70), while ANOVA for the last 2-s cue interval

differences scores found no main effect ($F_{(1,43)}$ = 2.49, p=0.12, η_{p2} = 0.05, op = 0.34). Difference

scores also suggest that NAcc rats were more specifically impaired in rapid uncertainty-safety

discrimination (Figure 2D, right). One-sample tests found that only the NAcc uncertainty-safety

difference score from the first 2-s cue interval failed to differ from zero: first interval: danger vs.

uncertainty (t_{23} = 4.20, p=3.38 x 10^{-4}), uncertainty vs. safety (t_{23} = 1.31, p=0.20); last interval:

danger vs. uncertainty (t_{20} = 5.22, p=2.70 x 10^{-5}), uncertainty vs. safety (t_{20} = 4.19, p=3.53 x 10^{-4}).

All significant, one-sample t tests survive Bonferroni correction (0.05/8, p<0.00625).

Altogether, these results reveal a general role for the NAcc in the acquisition of rapid and overall

fear scaling, as well as a more specific role in rapid uncertainty-safety discrimination.

Experiment 2

Experiment 1 results implicate the NAcc in the acquisition of fear scaling. However, neurotoxic

lesions altered baseline nose poking behavior and permanently ablated NAcc neurons. To

determine a temporally specific role for the NAcc in expression, Experiment 2 took a within-

subjects, optogenetic approach. Rats were NAcc-transducted with halorhodopsin or a control

fluorophore, recovered, then acquired a scaled fear response to danger, uncertainty and safety.

Once scaling was established, rats received sessions in which the NAcc was illuminated during

cue presentation or during the inter-trial interval. If the NAcc plays identical roles in the

acquisition and expression of fear scaling, we would expect to observe a three-way interaction

(group x illumination x cue) with only halorhodopsin rats showing impaired overall scaling during

cue illumination sessions. If the NAcc plays a more selective role in the expression of rapid fear

scaling, we would anticipate a four-way interaction (group x interval x illumination x cue) with

only halorhodopsin rats showing impaired rapid uncertainty-safety discrimination during cue

illumination sessions.

Histological Results

Rats received bilateral NAcc transduction with halorhodopsin (Halo) or a control fluorophore

(YFP) and bilateral optical ferrule implantation just above the NAcc. Representative transduction

is shown (Figure 3A). Each subject's total transduction area was drawn, made transparent, and

stacked (Figure 3B). Darker areas indicate regions of greater overlap and more consistent

transduction. Transduction centered around and above the anterior commissure, the precise

NAcc location.

Baseline Nose Poking

YFP and Halo rats showed equivalent baseline nose poking rates throughout pre-exposure,

discrimination, cable habituation, and light illumination (Figure S1). ANOVA for baseline nose

poke rate [factors: session (20) and group (YFP vs. Halo)] demonstrated a main effect of

session ($F_{(19,437)} = 12.60$, $p=4.19 \times 10^{-31}$, $\eta_{p2} = 0.35$, op = 1.00), but no main effect or interaction

with group (Fs < 0.93, ps>0.55). Equivalent performance lessens the concern that differences in

suppression ratios between groups result from differences in baseline nose poke rates.

Initial Fear Scaling

YFP and Halo rats acquired reliable fear scaling over the 10 sessions (Figure 3C). Suppression

ratios were low in pre-exposure and initially increased to all cues. As discrimination proceeded,

the suppression ratio for each cue diverged: high to danger, intermediate to uncertainty, and low

to safety. Demonstrating overall scaling, ANOVA [within factors: session (10) and 10-s cue

(danger, uncertainty and safety); between factor: group (YFP vs. Halo)] revealed a main effect

of cue ($F_{(2,46)} = 36.21$, *p*=3.58 x 10^{-10}, $\eta_{p2} = 0.61$, op = 1.00), session ($F_{(9,207)} = 25.74$, *p*=2.04 x 10^{-}

^{29} $\eta_{p2} = 0.53$, op = 1.00) and a cue x session interaction ($F_{(18,414)} = 6.26$, *p*=1.14 x 10^{-13}, $\eta_{p2} =$

0.21, op = 1.00). ANOVA found no main effect or interaction with group (Fs < 3.42, *p*s>0.08).

Thus, YFP and Halo entered the light illumination phase (Figure 3D, E) showing equivalent fear

scaling.

Overall Fear Scaling during Light Illumination

When suppression ratios were calculated for the entire 10-s cue, YFP and Halo rats showed

scaling of the fear response over the 10 sessions of cable habituation, cue illumination and ITI

illumination (Figure 4). ANOVA [between factor: group (YFP vs. Halo); within factors: session

(10) and cue (danger, uncertainty and safety)] was separately performed for rats receiving ITI-

cue illumination order (YFP, n = 5; Halo, n = 5; Figure 4A) and cue-ITI illumination order (YFP, n

= 7; Halo, n = 8; Figure 4B). Each ANOVA returned a main effect of cue (Fs > 29, *p*s<2 x 10^{-7}),

but neither returned a main effect of group, group x cue interaction or a group x cue x session

interaction (Fs < 2.5, ps>0.1). Complete ANOVA results provided in Table 1. Next we calculated

difference scores for the two components of scaling: (danger – uncertainty) and (uncertainty –

safety) (Figure 4C). ANOVA [between factors: group (YFP vs. Halo) and order (ITI-cue vs. cue-

ITI); within factors: illumination (hab/ITI vs. cue) and discrimination (danger – uncertainty vs.

uncertainty – safety)] found main effects of illumination ($F_{(1,21)} = 8.90$, *p*=0.007, $\eta_{p2} = 0.30$, op =

0.81) and discrimination ($F_{(1,21)} = 14.29$, *p*=0.001, $\eta_{p2} = 0.41$, op = 0.95), as well as a group x

illumination interaction ($F_{(1,21)} = 4.75$, *p*=0.041, $\eta_{p2} = 0.19$, op = 0.55). The interaction resulted

from YFP rats showing poorer overall discrimination in cue illumination sessions compared to

ITI illumination, whereas Halo rats showed equivalent discrimination in each session type. No

main effect of group ($F_{(1,21)} = 0.19$, *p*=0.67, $\eta_{p2} = 0.009$, op = 0.07) or any group interaction was

detected (Fs < 1.2, ps>0.3). These results reveal that NAcc activity is not necessary for the

expression of fear scaling when suppression is measured for the duration of cues.

Rapid Fear Scaling during Light Illumination

To examine rapid fear scaling, we divided the 10-s cue into 5, 2-s intervals. Suppression ratios

are shown for each cue/interval during habituation/ITI illumination sessions [YFP rats (Figure

5A) and Halo rats (Figure 5B)] and for cue illumination sessions [YFP rats (Figure 5C) and Halo

rats (Figure 5D)]. To examine a possible role for the NAcc in rapid fear scaling, we performed

ANOVA with all factors [within factors: session-type (cable habituation, ITI illumination and cue

illumination), cue (danger, uncertainty and safety), and interval (5, 2-s cue intervals); between

factor: group (YFP vs. Halo)]. The complete ANOVA output is reported in Table 2. Consistent

with general scaling across groups, ANOVA revealed a main effect of cue ($F_{(2.46)} = 89.04$,

$p=1.53 \times 10^{-16}$, $\eta_{p2} = 0.80$, op = 1.00) as well as a cue x interval interaction ($F_{(8.184)} = 6.14$, $p=5.16$

$\times 10^{-7}$, $\eta_{p2} = 0.21$, op = 1.00). Indicative of a selective role for the NAcc in rapid fear scaling,

ANOVA revealed a significant 4-way interaction [session-type x cue x interval x group ($F_{(16,368)} =$

1.80, $p=0.029$, $\eta_{p2} = 0.07$, op = 0.95)], but not a significant 3-way interaction [session-type x cue

x group ($F_{(4,92)} = 1.35$, $p=0.26$, $\eta_{p2} = 0.06$, op = 0.41)].

The 4-way interaction indicates that YFP and Halo rats showed differing temporal scaling

patterns across the different session types. To begin to clarify the differing patterns, we split

YFP and Halo rats and performed identical ANOVAs [within factors: session-type (habituation,

ITI illumination and cue illumination), cue (danger, uncertainty and safety), and interval (5, 2-s

cue intervals)]. Indicative of reliable scaling, ANOVA for YFP rats found a main effect of cue

($F_{(2,22)} = 47.71$, $p=1.0 \times 10^{-10}$, $\eta_{p2} = 0.81$, op = 1.00) and a cue x interval interaction ($F_{(8,88)} = 2.76$,

$p=0.009$, $\eta_{p2} = 0.20$, op = 0.92). Revealing no effect of illumination on the temporal pattern of

fear scaling, the 3-way interaction (session-type x cue x interval) was not significant ($F_{(16,176)} =$

0.59, $p = 0.89$, $\eta_{p2} = 0.05$, op = 0.39). ANOVA for Halo rats also found a main effect of cue

($F_{(2,24)} = 41.39$, $p=1.66 \times 10^{-8}$, $\eta_{p2} = 0.78$, op = 1.00) and a cue x interval interaction ($F_{(8,96)} = 4.07$,

$p=3.36 \times 10^{-4}$, $\eta_{p2} = 0.25$, op = 0.99). Only now, ANOVA revealed a significant 3-way interaction

(session-type x cue x interval; $F_{(16,192)}$ = 1.92, p = 0.021, η_{p2} = 0.14, op = 0.95). NAcc illumination

only disrupted the temporal scaling pattern for Halo rats. It appears that – similar to NAcc rats –

Halo rats receiving NAcc optogenetic inhibition during cue presentation were specifically

impaired in rapid uncertainty-safety discrimination (Figure 5D). If this were the case, then Halo

rats should show poorer uncertainty-safety discrimination in the first 2-s cue interval during cue

illumination sessions compared to ITI illumination sessions. YFP rats show would equivalent

performance during each type of illumination. Further, this deficit should not be observed in the

last 2-s cue interval.

We calculated (uncertainty – safety) difference scores for the first and last 2-s cue intervals.

Separate scores were calculated for cue and ITI illumination sessions. We then calculated a

difference score for the two session-illumination-types (cue difference score – ITI difference

score). This approach capitalized on our within-subject design; each rat was tested during cue

and ITI illumination. The approach is consistent with our ANOVA results, which found a

differential effect of cue and ITI illumination for Halo rats, but not for YFP rats. A difference

score of difference scores has the added benefit of reducing the differential illumination effects

to a single value. Values around zero would indicate equivalent uncertainty-safety discrimination

during cue and ITI illumination sessions. Negative values would indicate *worse* uncertainty-

safety discrimination during cue illumination sessions. Two individuals (1 YFP and 1 Halo) had

first interval difference scores ±2 standard deviations beyond the group mean. The data for

these individuals are shown (Figure 5E, open circles), but were not included in t-test analyses.

In the first 2-s cue interval, Halo rats showed worse uncertainty-safety discrimination during cue

illumination sessions compared to ITI illumination sessions (Figure 5E, left). This was supported

by significant, negative shift of differences scores away from zero (one-sample t-test, t_{11} = -3.65,

p=0.004). YFP rats showed equivalent uncertainty-safety discrimination during cue and ITI

illumination sessions; difference scores hovered around zero (t_{10} = 1.22, p=0.25). Further, YFP

and Halo difference scores differed from one another (independent samples t-test, t_{21} = 3.22,

p=0.004). Impaired uncertainty-safety discrimination in Halo rats receiving cue illumination was

restricted to the first 2-s cue interval. Identical analysis of the last 2-s cue interval found that

difference scores did not differ from zero for YFP (one-sample t-test, t_{10} = -0.41, p=0.69) and

Halo rats (one-sample t-test, t_{11} = 0.27, p=0.80) (Figure 5E, right). Differences scores were

similar between the two groups (independent samples t-test, t_{21} = 0.48, p=0.64). Altogether, the

results reveal that NAcc activity at the time of cue presentation is necessary to rapidly

discriminate uncertainty and safety.

Of course, it is possible that NAcc optogenetic inhibition simply suppressed rewarded nose

poking. In this case, impaired rapid fear scaling would be the byproduct of a general reduction in

poking. To rule out this possibility, we examined nose poke suppression during light illumination

in ITI sessions (Figure 5F). No cues were present during this period, allowing us to determine

the effect of light illumination alone to suppress nose poking. The middle 10 s of the 11-s light

illumination was divided into 5, 2-s cue intervals – exactly as was done for the cue illumination

analyses. For comparison, we also sampled 10 s of nose poking 30-s following illumination

offset. This post-illumination served as a control period to which light illumination could be

compared. ANOVA [within factors: period (light and post) and interval (5, 2-s cue intervals);

between factor: group (YFP vs. Halo)] revealed main effects of period ($F_{(1.23)}$ = 34.53, p=5 x 10_{-6},

η_{p2} = 0.60, op = 1.00) and interval ($F_{(4,92)}$ = 2.49, p=0.049, η_{p2} = 0.10, op = 0.69). Critically,

ANOVA found no main effect or interaction with group (Fs < 1.10, ps>0.31). So while

suppression ratios were higher during light illumination, this did not differ between YFP and Halo

rats and was therefore not due to inhibition of NAcc activity.

Discussion

We set out to examine a role for the nucleus accumbens core in fear scaling. Neurotoxic lesions

revealed a general role for the NAcc in the acquisition of fear scaling, as well as a specific role

in acquiring rapid uncertainty-safety discrimination. Optogenetic inhibition revealed a specific

role for NAcc cue activity in the expression of rapid, uncertainty-safety discrimination. The

results reveal that the NAcc is an essential component of a neural circuit permitting fear to

rapidly scale to degree of threat.

Before discussing our results more broadly, we must consider several limitations of our

experimental design and results. First, our experiments only used male rats. Several studies

have reported sex differences in danger-safety discrimination (Day et al., 2016; Foilb et al.,

2018; Greiner et al., 2019). We find only modest sex differences in our discrimination procedure

(Walker et al., 2018; Walker et al., 2019), suggesting similar neural circuits may be utilized

across sexes. Of course, females and males may achieve similar performance through differing

neural mechanisms. Another important consideration is that our dependent measure of fear is

derived from the rate of rewarded nose poking. Conditioned suppression is a strength because

it provides an objective measure of fear on multiple time scales (Estes and Skinner, 1941;

Bouton and Bolles, 1980). It is a potential weakness because the NAcc plays a well-established

role in reward-seeking. Disrupting NAcc function can attenuate reward-related behavior in many

settings (Corbit et al., 2001; Hall et al., 2001; Ito et al., 2004; Blaiss and Janak, 2009; Ambroggi

et al., 2011; McDannald et al., 2011; McDannald et al., 2013), though this finding is not

universal (Ramirez and Savage, 2007; Corbit and Balleine, 2011). In our first experiment, NAcc

lesions slowed the increase of baseline nose poking over discrimination sessions and also

impaired fear scaling. However, the temporal emergence of the deficits in reward-seeking and

fear scaling did not align. The fear scaling deficit was apparent across all sessions while the

nose poking deficit only emerged in the later sessions.

Our second experiment better demonstrated independent roles for the NAcc in rapid fear

scaling and reward-seeking. Light illumination during the cue period impaired rapid uncertainty-

safety discrimination in NAcc-halorhodopsin rats, but not NAcc-YFP rats. By contrast, light

illumination during the inter-trial interval produced equivalent and modest reductions in nose

poking for both groups. Optogenetic inhibition of the NAcc was insufficient to reduce rewarded

nose poking. The failure of NAcc inhibition to suppress nose poking may seem odd. Mice will

readily perform actions that channelrhodopsin-excite NAcc D1 and D2 cell types (Cole et al.,

2018), and rats will perform actions exciting glutamatergic inputs to the NAcc via

channelrhodopsin (Stuber et al., 2011; Britt et al., 2012). However, these studies demonstrate

that NAcc activity is sufficient, but not necessary, to support reward-seeking. Further, prominent

theories posit that reward-seeking initially depends on medial striatal structures, such as the

NAcc. With further training, lateral striatal regions, such as the dorsolateral striatum, control

reward-seeking (Gerdeman et al., 2003; Belin and Everitt, 2008; Corbit et al., 2012; Burton et

al., 2015). In our second experiment, rats had extensive experience with nose poking by the

time the NAcc was optogenetically inhibited. By this time, reward-seeking may have no longer

been under NAcc control, yet the NAcc continued to contribute to rapid fear scaling.

A role for the NAcc in fear would be expected based on immediate early gene studies. Shock-

associated cues and contexts reliably upregulate NAcc c-fos and zif268 (Beck and Fibiger,

1995; Campeau et al., 1997; Thomas et al., 2002). Despite these clear findings, specifying the

role of the NAcc in fear has presented a considerable challenge. Initial work by Parkinson and

colleagues found that NAcc lesions impaired cued fear, but enhanced contextual fear

(Parkinson et al., 1999). Taking a similar experimental approach, Levita and colleagues found

that NAcc lesions had no impact on the acquisition or expression of cued fear, but impaired

retention of contextual fear (Levita et al., 2002). Contemporary work by Haralambous and

Westbrook found that inhibiting accumbens activity (core + shell) specifically impaired the

acquisition, but not expression of contextual fear, and had no effect on cued fear (Haralambous

and Westbrook, 1999). Even considering slightly different methodologies, it is difficult to

reconcile these disparate results.

These are not the only conflicts in the literature. Schwienbacher and colleagues found that

blocking NAcc activity with tetrodotoxin abolished acquisition, and impaired expression, of fear-

potentiated startle (Schwienbacher et al., 2004). The very next year, Josselyn and colleagues

utilized a variety of methods to manipulate the NAcc during fear-potentiated startle: lesion,

agonizing dopamine and blocking glutamate. NAcc manipulation had no effect on any aspect of

fear-potentiated startle (Josselyn et al., 2005). Since these initial studies, the NAcc has been

implicated in a variety of fear-related processes. For example, the NAcc can modulate salience,

regulating the ability of cues to enter into associations with shock (Iordanova et al., 2006a;

Iordanova et al., 2006b; Iordanova, 2009). Human imaging studies have observed correlates of

prediction error, a theoretical signal that strengthens or weakens cue-shock associations

(Seymour et al., 2004; Delgado et al., 2008; Schiller et al., 2008; Li et al., 2011), consistent with

a role for the NAcc in predictive learning (Li and McNally, 2015).

What can we make of the mixed NAcc fear literature? Though dissatisfying, one answer is that

the NAcc must play multiple roles in fear. Genetically- and anatomically-defined NAcc neuron

types may be linked to specific fear processes. Future work dissecting the NAcc in this way, as

has been done in reward settings (Kupchik et al., 2015; Francis and Lobo, 2017; Tejeda et al.,

2017), is likely to be fruitful. A more fulfilling answer might be that standard cued and contextual

fear conditioning procedures do not isolate essential NAcc functions. The NAcc may not be

needed to demonstrate fear to certain threat or to withhold fear to certain safety. For example,

the NAcc is not necessary to behaviorally discriminate contexts/cues associated with certain

shock and certain safety (Antoniadis and McDonald, 2006; McDannald and Galarce, 2011;

Piantadosi, 2017).

We propose that a necessary role for the NAcc in fear emerges when subjects are confronted

with threats on a continuum from safety to danger. The NAcc is a core component of a neural

circuit permitting the level of the fear response to scale to the degree of threat. Further, our

results suggest general and specific roles for the NAcc in fear scaling. During acquisition, the

NAcc is generally necessary for fear scaling for the duration of an encounter – in our case for

the entirety of cue presentation. At the same time, the NAcc is specifically necessary for one

component of fear scaling: rapid discrimination of uncertain threat and safety. Once a scaled

fear response is acquired, the general role for the NAcc diminishes. However, the NAcc

continues to play a specific role in rapidly discriminating uncertain threat and safety. Of course,

we are not claiming that fear scaling is *the* function of the NAcc in fear, but rather *a* function.

Environmental threats are not absolute, but exist on a continuum from safety to danger. Using a

behavioral procedure that attempts to capture this continuum, we find the NAcc is essential to

scale fear to degree of threat. Our results clarify at least one role for the NAcc in fear, yet much

more work remains. There is evidence of altered NAcc structure and function in anxiety and

stress disorders (Cha et al., 2014; Felmingham et al., 2014; Manning et al., 2015; Morey et al.,

2017). Disrupted threat-safety discrimination may be conceptualized as maladaptive fear

scaling. Detailing NAcc function, and mapping a more complete neural circuit for rapid fear

scaling, may inform strategies to promote adaptive fear in anxiety and stress disorders.

Figure 1. NAcc lesion experimental outline. (**A**) Representative sham with NAcc intact (left) and lesion with NAcc damage (right) is shown. Dotted lines (left) show approximate NAcc location. Arrows (right) indicate gliosis and damage restricted to the NAcc. (**B**) The extent of neurotoxic NAcc lesions across four coronal planes is shown, and the anterior distance from bregma (millimeters) indicated. (**C**) Pavlovian fear discrimination consisted of three, 10-s cues predicting unique foot shock probabilities: danger (p=1.00), red; uncertainty (p=0.25), purple; and safety (p=0.00), blue. Cues were divided into 5, 2-s intervals for rapid analyses. (**D**) Mean ± SEM baseline nose poke rates for the sixteen fear discrimination sessions are shown for sham (black) and NAcc (gray) rats. (**E**) Mean baseline nose poke rates for sessions 1-8 and 9-16 for sham and NAcc rats. Data points show individual poke rates. *independent samples t-test, p<0.025, +block x group interaction p<0.05. Abbreviations: NAcc – nucleus accumbens core, NAs – nucleus accumbens shell.

Figure 2. NAcc lesions and fear scaling. (**A**) Mean ± SEM suppression ratio for danger (red), uncertainty (purple), and safety (blue) are shown for sham (left) and NAcc (right) rats. The vertical lines separate the two pre-exposure and sixteen fear discrimination sessions. The last six discrimination sessions are shaded. (**B**) Mean difference score for danger vs. uncertainty (D-U, red bar) and uncertainty vs. safety (U-S, purple bar) across the entire 10-s cue is shown for sham (left) and NAcc (right) rats. Data points show individual difference scores. *One-sample t-test compared to zero, p<0.0125; +main effect of group, p<0.05. (**C**) Mean + SEM suppression ratios for the 5, 2-s cue intervals are shown for sham (left) and NAcc (right) rats. Cue color scheme maintained from A. (**D**) Mean difference score for danger vs. uncertainty (D-U, red bar) and uncertainty vs. safety (U-S, purple bar) is shown for the first 2-s cue interval (left) and last 2-s cue interval (right) for sham and NAcc rats. Data points show individual difference scores. *One-sample t-test compared to zero, p<0.00625; +main effect of group, p<0.05.

Figure 3. NAcc illumination experimental outline. (**A**) Representative NAcc transduction is shown with YFP expression (yellow fluorescent protein; yellow) and neurotrace (blue). Dotted lines indicate approximate NAcc location. (**B**) The extent of viral transduction across four coronal planes is shown for Halo (green, left) and YFP rats (yellow, right), and the anterior distance from bregma (millimeters) indicated. Individual ferrule placement is indicated in black circles. (**C**) Mean ± SEM suppression ratios for danger (red), uncertainty (purple), and safety (blue) are shown for YFP (left) and Halo rats (right) during the ten initial fear discrimination sessions. (**D**) In the final eight sessions, rats received NAcc light illumination during cue presentation (top) or during the inter-trial interval (ITI, bottom). Green indicates light illumination, yellow indicates shock delivery and candy-striped indicates cue presentation. (**E**) Cue and ITI illumination were given in alternating, two-session blocks. Block order was counterbalanced with roughly half of the subjects first receiving ITI illumination. Abbreviations: NAcc – nucleus accumbens core, NAs – nucleus accumbens shell.

Figure 4. NAcc illumination and overall fear scaling. (**A**) Mean ± SEM suppression ratios over the entire 10-s cue are plotted for danger (red), uncertainty (purple), and safety (blue). Data are plotted for cable habituation (H), ITI illumination (I) and cue illumination (C) for YFP (n=5) and Halo rats (n=5) receiving ITI-cue illumination. ITI illumination sessions shaded. (**B**) YFP (n=7) and Halo rats (n=8) receiving cue-ITI illumination plotted as in A. (**C**) Difference scores for danger vs. uncertainty (D-U, red bar) and uncertainty vs. safety (U-S, purple bar) are shown for YFP (black) and Halo rats (green) during cable habituation/ITI illumination (left) and cue illumination (right). ITI-cue rats indicated by open circles, cue-ITI rats by closed circles.

Figure 5. NAcc illumination and rapid fear scaling. Mean + SEM suppression ratios are plotted for the 5, 2-s cue intervals for danger (red), uncertainty (purple), and safety (blue), for YFP and Halo rats during (**A** & **B**) cable habituation/ITI illumination, and (**C** & **D**) cue illumination. (**E**) Uncertainty – safety difference scores were separately calculated for cue and

ITI illumination, then difference of difference scores calculated (cue difference score – ITI

difference score). Mean and individual difference of difference scores are plotted for the first 2-s

cue interval (left) and last 2-s cue interval (right), for YFP (black) and Halo rats (green). Open

circles are outliers. *(green) one-sample t-test compared to zero, p=0.0038. *(black)

independent samples t-test, p=0.0041. (**F**) Mean + SEM suppression ratios are plotted for the 5,

2-s intervals during ITI illumination (light green) and for the 5, 2-s intervals during the post-

illumination period (gray) (YFP, left; Halo, right).

References

Ambroggi F, Ghazizadeh A, Nicola SM, Fields HL (2011) Roles of nucleus accumbens core and shell in incentive-cue responding and behavioral inhibition. J Neurosci 31:6820-6830.

Antoniadis EA, McDonald RJ (2006) Fornix, medial prefrontal cortex, nucleus accumbens, and mediodorsal thalamic nucleus: Roles in a fear-based context discrimination task. Neurobiology of Learning and Memory 85:71-85.

Badrinarayan A, Wescott SA, Vander Weele CM, Saunders BT, Couturier BE, Maren S, Aragona BJ (2012) Aversive Stimuli Differentially Modulate Real-Time Dopamine Transmission Dynamics within the Nucleus Accumbens Core and Shell. Journal of Neuroscience 32:15779-15790.

Beck CH, Fibiger HC (1995) Conditioned fear-induced changes in behavior and in the expression of the immediate early gene c-fos: with and without diazepam pretreatment. Journal of Neuroscience 15:709-720.

Belin D, Everitt BJ (2008) Cocaine seeking habits depend upon dopamine-dependent serial connectivity linking the ventral with the dorsal striatum. Neuron 57:432-441.

Berg BA, Schoenbaum G, McDannald MA (2014) The dorsal raphe nucleus is integral to negative prediction errors in Pavlovian fear. European Journal of Neuroscience 40:3096-3101.

Blaiss CA, Janak PH (2009) The nucleus accumbens core and shell are critical for the expression, but not the consolidation, of Pavlovian conditioned approach. Behav Brain Res 200:22-32.

Bouton ME, Bolles RC (1980) Conditioned fear assessed by freezing and by the suppression of three different baselines. Animal Learning & Behavior 8:429-434.

Britt JP, Benaliouad F, McDevitt RA, Stuber GD, Wise RA, Bonci A (2012) Synaptic and behavioral profile of multiple glutamatergic inputs to the nucleus accumbens. Neuron 76:790-803.

Burton AC, Nakamura K, Roesch MR (2015) From ventral-medial to dorsal-lateral striatum: Neural correlates of reward-guided decision-making. Neurobiology of Learning and Memory 117:51-59.

Campeau S, Falls WA, Cullinan WE, Helmreich DL, Davis M, Watson SJ (1997) Elicitation and reduction of fear: Behavioural and neuroendocrine indices and brain induction of the immediate-early gene c-fos. Neuroscience 78:1087-1104.

Cha J, Carlson JM, DeDora DJ, Greenberg T, Proudfit GH, Mujica-Parodi LR (2014) Hyper-Reactive Human Ventral Tegmental Area and Aberrant Mesocorticolimbic Connectivity in Overgeneralization of Fear in Generalized Anxiety Disorder. Journal of Neuroscience 34:5855-5860.

Cole SL, Robinson MJF, Berridge KC (2018) Optogenetic self-stimulation in the nucleus accumbens: D1 reward versus D2 ambivalence. PLoS One 13:e0207694.

Corbit LH, Balleine BW (2011) The general and outcome-specific forms of pavlovian-instrumental transfer are differentially mediated by the nucleus accumbens core and shell. J Neurosci 31:11786-11794.

Corbit LH, Muir JL, Balleine BW (2001) The role of the nucleus accumbens in instrumental conditioning: Evidence of a functional dissociation between accumbens core and shell. Journal of Neuroscience 21:3251-3260.

Corbit LH, Nie H, Janak PH (2012) Habitual Alcohol Seeking: Time Course and the Contribution of Subregions of the Dorsal Striatum. Biological Psychiatry 72:389-395.

Correia SS, McGrath AG, Lee A, Graybiel AM, Goosens KA (2016) Amygdala-ventral striatum circuit activation decreases long-term fear. Elife 5.

Cromwell HC, Schultz W (2003) Effects of expectations for different reward magnitudes on neuronal activity in primate striatum. J Neurophysiol 89:2823-2838.

Day HLL, Reed MM, Stevenson CW (2016) Sex differences in discriminating between cues
predicting threat and safety. Neurobiology of Learning and Memory 133:196-203.
Delgado MR, Li J, Schiller D, Phelps EA (2008) The role of the striatum in aversive learning and
aversive prediction errors. Philos Trans R Soc Lond B Biol Sci 363:3787-3800.
DiLeo A, Wright KM, McDannald MA (2016) Sub-second fear discrimination in rats: Adult
impairment in adolescent heavy alcohol drinkers. Learning & Memory 23:618-622.
Duits P, Cath DC, Lissek S, Hox JJ, Hamm AO, Engelhard IM, van den Hout MA, Baas JM
(2015) Updated meta-analysis of classical fear conditioning in the anxiety disorders.
Depress Anxiety 32:239-253.
Estes KW, Skinner BF (1941) Some Quantitative Properties of Anxiety. Journal of Experimental
Psychology 29:390-400.
Fadok JP, Darvas M, Dickerson TMK, Palmiter RD (2010) Long-Term Memory for Pavlovian
Fear Conditioning Requires Dopamine in the Nucleus Accumbens and Basolateral
Amygdala. Plos One 5.
Felmingham KL, Falconer EM, Williams L, Kemp AH, Allen A, Peduto A, Bryant RA (2014)
Reduced Amygdala and Ventral Striatal Activity to Happy Faces in PTSD Is Associated
with Emotional Numbing. Plos One 9.
Foilb AR, Bals J, Sarlitto MC, Christianson JP (2018) Sex differences in fear discrimination do
not manifest as differences in conditioned inhibition. Learning & Memory 25:49-53.
Francis TC, Lobo MK (2017) Emerging Role for Nucleus Accumbens Medium Spiny Neuron
Subtypes in Depression. Biol Psychiatry 81:645-653.
Gerdeman GL, Partridge JG, Lupica CR, Lovinger DM (2003) It could be habit forming: drugs of
abuse and striatal synaptic plasticity. Trends in Neurosciences 26:184-192.
Goosens KA, Maren S (2001) Contextual and auditory fear conditioning are mediated by the
lateral, basal, and central amygdaloid nuclei in rats. Learning & Memory 8:148-155.
Greiner EM, Muller I, Norris MR, Ng KH, Sangha S (2019) Sex differences in fear regulation and
reward-seeking behaviors in a fear safety-reward discrimination task. Behavioural Brain
Research 368:131-139.
Hall J, Parkinson JA, Connor TM, Dickinson A, Everitt BJ (2001) Involvement of the central
nucleus of the amygdala and nucleus accumbens core in mediating Pavlovian influences
on instrumental behaviour. European Journal of Neuroscience 13:1984-1992.
Haralambous T, Westbrook RF (1999) An infusion of bupivacaine into the nucleus accumbens
disrupts the acquisition but not the expression of contextual fear conditioning. Behavioral
Neuroscience 113:925-940.
Iordanova MD (2009) Dopaminergic Modulation of Appetitive and Aversive Predictive Learning.
Reviews in the Neurosciences 20:383-404.
Iordanova MD, McNally GP, Westbrook RF (2006a) Opioid receptors in the nucleus accumbens
regulate attentional learning in the blocking paradigm. Journal of Neuroscience 26:4036-
4045.
Iordanova MD, Westbrook RF, Killcross AS (2006b) Dopamine activity in the nucleus
accumbens modulates blocking in fear conditioning. European Journal of Neuroscience
24:3265-3270.
Ito R, Robbins TW, Everitt BJ (2004) Differential control over cocaine-seeking behavior by
nucleus accumbens core and shell. Nature Neuroscience 7:389-397.
Josselyn SA, Falls WA, Gewirtz JC, Pistell P, Davis M (2005) The nucleus accumbens is not
critically involved in mediating the effects of a safety signal on behavior.
Neuropsychopharmacology 30:17-26.
Jovanovic T, Kazama A, Bachevalier J, Davis M (2012) Impaired safety signal learning may be
a biomarker of PTSD. Neuropharmacology 62:695-704.

Jovanovic T, Norrholm SD, Blanding NQ, Davis M, Duncan E, Bradley B, Ressler KJ (2010)
Impaired fear inhibition is a biomarker of PTSD but not depression. Depress Anxiety
27:244-251.
Kita H, Kitai ST (1990) Amygdaloid Projections to the Frontal-Cortex and the Striatum in the
Rat. Journal of Comparative Neurology 298:40-49.
Koo JW, Han JS, Kim JJ (2004) Selective neurotoxic lesions of basolateral and central nuclei of
the amygdala produce differential effects on fear conditioning. Journal of Neuroscience
24:7654-7662.
Kupchik YM, Brown RM, Heinsbroek JA, Lobo MK, Schwartz DJ, Kalivas PW (2015) Coding the
direct/indirect pathways by D1 and D2 receptors is not valid for accumbens projections.
Nature Neuroscience 18:1230-+.
Levita L, Dalley JW, Robbins TW (2002) Disruption of Pavlovian contextual conditioning by
excitotoxic lesions of the nucleus accumbens core. Behavioral Neuroscience 116:539-
552.
Li J, Schiller D, Schoenbaum G, Phelps EA, Daw ND (2011) Differential roles of human striatum
and amygdala in associative learning. Nat Neurosci 14:1250-1252.
Li SSY, McNally GP (2015) A Role of Nucleus Accumbens Dopamine Receptors in the Nucleus
Accumbens Core, but Not Shell, in Fear Prediction Error. Behavioral Neuroscience
129:450-456.
Lissek S, Kaczkurkin AN, Rabin S, Geraci M, Pine DS, Grillon C (2014) Generalized Anxiety
Disorder Is Associated With Overgeneralization of Classically Conditioned Fear.
Biological Psychiatry 75:909-915.
Manning J, Reynolds G, Saygin ZM, Hofmann SG, Pollack M, Gabrieli JD, Whitfield-Gabrieli S
(2015) Altered resting-state functional connectivity of the frontal-striatal reward system in
social anxiety disorder. PLoS One 10:e0125286.
McDannald MA, Galarce EM (2011) Measuring Pavlovian fear with conditioned freezing and
conditioned suppression reveals different roles for the basolateral amygdala. Brain
Research 1374:82-89.
McDannald MA, Setlow B, Holland PC (2013) Effects of ventral striatal lesions on first- and
second-order appetitive conditioning. Eur J Neurosci 38:2589-2599.
McDannald MA, Lucantonio F, Burke KA, Niv Y, Schoenbaum G (2011) Ventral Striatum and
Orbitofrontal Cortex Are Both Required for Model-Based, But Not Model-Free,
Reinforcement Learning. Journal of Neuroscience 31:2700-2705.
McGinty VB, Lardeux S, Taha SA, Kim JJ, Nicola SM (2013) Invigoration of reward seeking by
cue and proximity encoding in the nucleus accumbens. Neuron 78:910-922.
Morey RA, Davis SL, Garrett ME, Haswell CC, Marx CE, Beckham JC, McCarthy G, Hauser
MA, Ashley-Koch AE, Workgrp M-AM (2017) Genome-wide association study of
subcortical brain volume in PTSD cases and trauma-exposed controls. Translational
Psychiatry 7.
Oleson EB, Gentry RN, Chioma VC, Cheer JF (2012) Subsecond dopamine release in the
nucleus accumbens predicts conditioned punishment and its successful avoidance. J
Neurosci 32:14804-14808.
Ottenheimer D, Richard JM, Janak PH (2018) Ventral pallidum encodes relative reward value
earlier and more robustly than nucleus accumbens. Nature Communications 9.
Parkinson JA, Olmstead MC, Burns LH, Robbins TW, Everitt BJ (1999) Dissociation in effects of
lesions of the nucleus accumbens core and shell on appetitive pavlovian approach
behavior and the potentiation of conditioned reinforcement and locomotor activity by D-
amphetamine. Journal of Neuroscience 19:2401-2411.
Petrovich GD, Risold PY, Swanson LW (1996) Organization of projections from the basomedial
nucleus of the amygdala: a PHAL study in the rat. J Comp Neurol 374:387-420.

Piantadosi PT (2017) Contributions of nucleus accumbens circuitry to aspects of aversively-
motivated behaviors. In: Psychology, p 235: University of British Columbia.
Quirk GJ, Repa C, LeDoux JE (1995) Fear conditioning enhances short-latency auditory
responses of lateral amygdala neurons: parallel recordings in the freely behaving rat.
Neuron 15:1029-1039.
Ramirez DR, Savage LM (2007) Differential involvement of the basolateral amygdala,
orbitofrontal cortex, and nucleus accumbens core in the acquisition and use of reward
expectancies. Behav Neurosci 121:896-906.
Ray MH, Hanlon E, McDannald MA (2018) Lateral orbitofrontal cortex partitions mechanisms for
fear regulation and alcohol consumption. PLoS One 13:e0198043.
Rescorla RA (1968) Probability of shock in the presence and absence of CS in fear
conditioning. Journal of Comparative and Physiological Psychology 66:1-5.
Saddoris MP, Carelli RM (2014) Cocaine self-administration abolishes associative neural
encoding in the nucleus accumbens necessary for higher-order learning. Biol Psychiatry
75:156-164.
Schiller D, Levy I, Niv Y, LeDoux JE, Phelps EA (2008) From Fear to Safety and Back: Reversal
of Fear in the Human Brain. Journal of Neuroscience 28:11517-11525.
Schwienbacher I, Fendt M, Richardson R, Schnitzler HU (2004) Temporary inactivation of the
nucleus accumbens disrupts acquisition and expression of fear-potentiated startle in
rats. Brain Research 1027:87-93.
Setlow B, Schoenbaum G, Gallagher M (2003) Neural encoding in ventral striatum during
olfactory discrimination learning. Neuron 38:625-636.
Seymour B, O'Doherty JP, Dayan P, Koltzenburg M, Jones AK, Dolan RJ, Friston KJ,
Frackowiak RS (2004) Temporal difference models describe higher-order learning in
humans. Nature 429:664-667.
Stuber GD, Sparta DR, Stamatakis AM, van Leeuwen WA, Hardjoprajitno JE, Cho S, Tye KM,
Kempadoo KA, Zhang F, Deisseroth K, Bonci A (2011) Excitatory transmission from the
amygdala to nucleus accumbens facilitates reward seeking. Nature 475:377-380.
Sugam JA, Saddoris MP, Carelli RM (2014) Nucleus accumbens neurons track behavioral
preferences and reward outcomes during risky decision making. Biol Psychiatry 75:807-
816.
Tejeda HA, Wu J, Kornspun AR, Pignatelli M, Kashtelyan V, Krashes MJ, Lowell BB, Carlezon
WA, Bonci A (2017) Pathway-and Cell-Specific Kappa-Opioid Receptor Modulation of
Excitation-Inhibition Balance Differentially Gates D1 and D2 Accumbens Neuron Activity.
Neuron 93:147-163.
Thomas KL, Hall J, Everitt BJ (2002) Cellular imaging with zif268 expression in the rat nucleus
accumbens and frontal cortex further dissociates the neural pathways activated following
the retrieval of contextual and cued fear memory. European Journal of Neuroscience
16:1789-1796.
Walker RA, Andreansky C, Ray MH, McDannald MA (2018) Early adolescent adversity inflates
threat estimation in females and promotes alcohol use initiation in both sexes. Behav
Neurosci 132:171-182.
Walker RA, Wright KM, Jhou TC, McDannald MA (2019) The ventrolateral periaqueductal gray
updates fear via positive prediction error. Eur J Neurosci.
Wright CI, Groenewegen HJ (1996) Patterns of overlap and segregation between insular
cortical, intermediodorsal thalamic and basal amygdaloid afferents in the nucleus
accumbens of the rat. Neuroscience 73:359-373.
Wright KM, DiLeo A, McDannald MA (2015) Early adversity disrupts the adult use of aversive
prediction errors to reduce fear in uncertainty. Front Behav Neurosci 9:227.

A Sham | NAcc
neurotrace

C
0   10-s Cue   10
Danger (4)
Uncertainty Shock (2)
Uncertainty Omission (6)
Safety (4)
1 2 3 4 5
2-s Cue Interval

Surgery → Nose Poke Acquisition → Fear Discrimination over Rewarded Nose Poking

B
■ 100%
■ 75%
■ 50%
■ 25%

+2.16
+1.68
+1.20
+0.72

D
Poke/min
Sham
NAcc
Session
p p 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16

E
Poke/min
*
+
*
1-8 9-16   1-8 9-16
Sham   NAcc

A

yfp
neurotrace

C

YFP   Halo

Suppression Ratio

Danger
Unc
Safety

Session   Session

Surgery → Nose Poke Acquisition → Fear Discrimination → Cable Habituation → Light Illumination

Halo   YFP

B

+2.16

+1.68

+1.20

+0.72

D Cue Illumination

0   10

ITI Illumination

0   10   ~90 s   0   10

E

| ITI | ITI | Cue | Cue | ITI | ITI | Cue | Cue |

or

| Cue | Cue | ITI | ITI | Cue | Cue | ITI | ITI |

A. YFP ITI-Cue (n=5)   Halo ITI-Cue (n=5)

B. YFP Cue-ITI (n=7)   Halo Cue-ITI (n=8)

C. Hab/ITI Illumination   Cue Illumination

A. YFP Hab/ITI Illumination

Suppression Ratio vs 2-s Cue Interval. Traces: Danger, Unc, Safety.

B. Halo Hab/ITI Illumination

Suppression Ratio vs 2-s Cue Interval. Traces: Danger, Unc, Safety.

C. YFP Cue Illumination

Suppression Ratio vs 2-s Cue Interval. Traces: Danger, Unc, Safety.

D. Halo Cue Illumination

Suppression Ratio vs 2-s Cue Interval. Traces: Danger, Unc, Safety.

E.

U-S Difference Score Cue − ITI Illumination. First Interval: YFP, Halo. Last Interval: YFP, Halo.

F. ITI Illumination

YFP and Halo. Suppression Ratio vs 2-s Interval. Traces: Light, Post.

10-s cue ANOVA results
ITI-Cue

Term	F	p	η_p^2	op
cue	**103.37**	**7.09 x 10⁻¹⁰**	**0.93**	**1.00**
cue x group	0.55	0.59	0.06	0.13
session	1.55	0.15	0.16	0.68
session x group	**2.04**	**0.047**	**0.2**	**0.82**
cue x session	1.21	0.26	0.13	0.79
cue x session x group	1.18	0.29	0.13	0.78
group	2.51	0.15	0.24	0.29

10-s cue ANOVA results
Cue-ITI

Term	F	p	η_p^2	op
cue	**29.23**	**2.23 x 10⁻⁷**	**0.69**	**1.00**
cue x group	0.15	0.86	0.01	0.07
session	**3.46**	**0.001**	**0.21**	**0.98**
session x group	0.77	0.65	0.06	0.37
cue x session	1.07	0.39	0.08	0.74
cue x session x group	1.47	0.10	0.10	0.90
group	1.19	0.30	0.08	0.17

2-s cue interval ANOVA results

Term	F	p	η_p^2	op
group	1.94	0.18	0.08	0.27
session-type	**10.64**	**1.59 x 10-4**	**0.32**	**0.99**
session-type x group	2.22	0.12	0.09	0.43
cue	**89.04**	**1.53 x 10-16**	**0.80**	**1.00**
cue x group	0.57	0.57	0.02	0.14
interval	**6.00**	**2.46 x 10-4**	**0.21**	**0.98**
interval x group	1.37	0.25	0.06	0.41
session-type x cue	**4.21**	**0.004**	**0.16**	**0.91**
session-type x cue x group	1.35	0.26	0.06	0.41
session-type x interval	0.71	0.68	0.03	0.32
session-type x interval x group	1.46	0.17	0.06	0.65
cue x interval	**6.14**	**5.16 x 10-7**	**0.21**	**1.00**
cue x interval x group	0.64	0.74	0.03	0.29
session-type x cue x interval	0.78	0.71	0.03	0.54
session-type x cue x interval x group	**1.80**	**0.029**	**0.07**	**0.95**