A robust mean and variance test with application to high-dimensional phenotypes

James R Staley[1], Frank Windmeijer[1,2], Matthew Suderman[1], Matthew S Lyon[1,3], George Davey Smith[1] and Kate Tilling[1,*].

[1]MRC Integrative Epidemiology Unit, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK.

[2]Department of Economics, University of Bristol, Bristol, UK.

[3]National Institute for Health Research Bristol Biomedical Research Centre, University of Bristol, Oakfield House, Bristol, BS8 2BN, UK.

[*]Corresponding author.

Short title: Joint mean and variance test for omics data

Correspondence:

Prof Kate Tilling

MRC Integrative Epidemiology Unit, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, BS8 2BN, UK.

Email: kate.tilling@bristol.ac.uk

Telephone: +44 (0)117 331 0019

Abstract

Most studies of high-dimensional phenotypes focus on assessing differences in mean levels (location) of the phenotype by exposure, e.g. epigenome-wide association studies of DNA methylation at CpG sites. However, identifying effects on the variability (scale) of these outcomes, and combining tests of mean and variability (location-and-scale), could provide additional insights into biological mechanisms. Here, we review variability tests, specifically an extended (for continuous exposures) version of the Brown-Forsythe test, and develop a novel joint location-and-scale score test for both categorical and continuous exposures (JLSsc). The Brown-Forsythe test and JLSsc performed well in comparison to alternative approaches in simulations. These approaches identified >7500 CpG sites that were associated with either mean or variability with gender or gestational age in cord blood methylation in ARIES (Accessible Resource for Integrated Studies). The Brown-Forsythe test and JLSsc are robust tests that can be used to detect associations not solely driven by a mean effect.

Keywords: variability test, joint location-and-scale test, DNA methylation, ARIES, ALSPAC.

1 Introduction

Most investigations into health-related phenotypes have focused on determining whether an

exposure affects the mean of a phenotype (location test). However, assessing whether an

exposure affects the variability of a phenotype (scale test) could also provide insight into the

biological mechanisms that control phenotypic variation and disease pathogenesis as well as

identify possible interactions[1, 2, 3]. Furthermore, the potential of combining a location test with

a scale test (joint location-and-scale test) has yet to be fully explored, especially in the context

of high-dimensional phenotypes where these tests could be used to improve power as well as

to identify markers involved in interactions[4]. One example where these approaches could be

particularly useful is for epigenome-wide association studies (EWAS), where DNA

methylation at CpG (cytosine followed by a guanine) sites across the genome are tested for

association with an exposure (Supplementary Text)[5, 6].

A range of statistical tests have been developed to test whether an exposure affects variability

of an outcome, specifically in the context of evaluating variability differences for a

continuous variable between groups of individuals[7]. Li *et al*.[8] compared approaches for

assessing methylation variability in the EWAS setting, and showed that the Brown-Forsythe

test[9] performed well compared to alternative approaches. Since this test can be re-formulated

in a regression framework[10, 11], it can be extended to continuous exposures. Methods for

jointly testing mean and variability have also been proposed[4, 10, 11, 12, 13, 14, 15], although these

approaches are either limited by sensitivity to distributional assumptions or are restricted to

binary exposures.

Here, we review variability tests, specifically the Brown-Forsythe test, and develop a novel

joint location-and-scale test, which can be used for both continuous and categorical

exposures. We performed a simulation study to compare these approaches to alternative tests,

and then applied these modelling approaches to investigate the effect of gender and

gestational age on cord blood DNA methylation mean and variability.

Results

Review of statistical methods

Ordinary least squares (OLS) regression is commonly used to assess mean differences in an

outcome by an exposure and is known to be relatively robust to its underlying assumptions

(Methods and Supplementary Text), so is used as the location test. Likewise, the Brown-

Forsythe approach has been found to be robust to non-normality and outliers[8], so is also a

good candidate for analysing high-dimensional phenotypes, especially since it can be

extended for use with continuous exposures (see Methods). A computationally efficient joint

location-and-scale test, which combines p-values from location and scale tests using Fisher's

method (JLSp), has been developed for genetic data[10, 11]. However, this test like most existing

joint location-and-scale tests requires the phenotype to be normally distributed, while other

tests are only designed for binary exposures. Here, we develop a novel joint location-and-

scale score test (JLSsc) that is: (i) robust to distributional assumptions, and (ii) can be used for

both categorical and continuous exposures.

Our approach essentially combines a location test and scale test, while accounting for the

correlation between these tests (see Methods). Briefly, we propose to test the joint null

hypothesis $H_0: \beta = \delta = 0$ in the model specification:

$$y_i = \alpha + x_i'\beta + \varepsilon_i$$

$$(y_i - \bar{y})^2 = \lambda + x_i'\delta + u_i ,$$

20 where y_i is the outcome for the i-th individual, x_i is the (k_x) vector of exposures for the i-th

21 individual and $\bar{y}$ is the sample average of y_i.

1   Let $\tilde{y}_i = y_i - \bar{y}$, $\tilde{x}_i = x_i - \bar{x}$ (and $\tilde{X} = [\tilde{x}_i{'}]$) and $\tilde{d}_i = \tilde{y}_i^2 - \hat{\sigma}^2$, where $\hat{\sigma}^2 = \frac{1}{n}\sum_{i=1}^{n}\tilde{y}_i^2$.

2   Further, let $\hat{\theta} = \begin{pmatrix}\hat{\beta}\\\hat{\delta}\end{pmatrix}$, where , and $\hat{\Sigma} = \frac{1}{n}\sum_{i=1}^{n}\begin{bmatrix}\tilde{y}_i^2 & \tilde{y}_i\tilde{d}_i \\ \tilde{y}_i\tilde{d}_i & \tilde{d}_i^2\end{bmatrix}$, then the score test for $H_0: \beta =$

3     $\delta = 0$ or $H_0: \theta = 0$, is given by

$$S = \hat{\theta}'\left(\hat{\Sigma}^{-1} \otimes \left(\tilde{X}'\tilde{X}\right)\right)\hat{\theta} \xrightarrow{d} \chi^2_{2k_x}\ .$$

Simulation study

We performed a simulation study to evaluate the performance of the location and scale tests

as well as the joint location-and-scale tests with both binary and continuous exposures in the

context of DNA methylation. We compared the Brown-Forsythe scale test to Bartlett's test[16]

(for simulations with a binary exposure) and a likelihood ratio test (LRT) comparing mixed

models with and without a variability effect (LRTv). JLSsc was compared to JLSp as well as

a LRT comparing mixed models with and without including both a mean and variability effect

for the exposure (LRTmv) and a double generalized linear mixed model (DGLM)[13, 14, 17]. The

simulations were performed using data from the Tsaprouni *et al.* study[18], which investigated

the relationship between smoking and DNA methylation (data accessible at NCBI GEO

database[19], accession GSE50660). Type I error simulations were performed across all CpG

sites in Tsaprouni *et al.* for varying sample sizes, while power simulations were performed for

various phenotypic distributions using 1000 samples where in each simulation replicate the

characteristics from a randomly selected CpG site from Tsaprouni *et al.* were used (see

Methods and Supplementary Text). In the type I error simulations, common data

transformations such as the M-value transformation for DNA methylation data[20] and inverse

normal rank transformation were applied to the methylation data for those approaches that

failed to control type I error due to deviations from normality.

OLS regression test of mean differences was not inflated under the null of no mean or

variability effect even in 100 samples (Figures 1a and S1). Similarly, the Brown-Forsythe

variability test accurately controlled type I error rates (Figures 1b and S2). Bartlett's test and

LRTv had extreme type I error inflation due to the deviations from normality and the

existence of outlying values in methylation levels (Figure S3). Likewise, the test statistics

from the likelihood-based approaches for joint testing the mean and variability (LRTmv and

DGLM) were also heavily inflated (Figure S3). The extreme inflated type I error rates of these

approaches were still present after transforming methylation levels using the M-value

transformation (Figure S4) but were no longer present after using an inverse normal rank

transformation (Figure S5). However, when using this transformation a mean effect can

induce a variability effect and vice versa (Figure S6), as seen previously[21]. JLSp fared better

than the aforementioned joint tests in controlling type I error rates, although the non-

independence of the p-values did lead to a small amount of type I error inflation (Figure 1c

and Figure S7). The JLSsc approach, on the other hand, correctly controlled type I error rates

(Figure 1d and Figure S7).

In the power simulations, when there was either a mean or variability effect and the

underlying data were normally distributed, the Brown-Forsythe test and JLSsc were less

powerful but still performed well in comparison to the equivalent LRT and the alternative

approaches (Figure 2). This is expected as the Brown-Forsythe test and JLSsc sacrifice a

small amount of power under the normal model for robustness to deviations from this model.

Broadly similar results were found when the residual error was heavy-tailed or skewed, when

the exposure was a categorical variable with three categories, when a squared exposure term

was added to JLSsc for a continuous exposure (although outlier removal in the outcome is

necessary here to retain type I error levels), and when there was an outlier in the dataset

(Figures S8-S12).

The computational time required to complete each approach for 100,000 CpGs with a binary

exposure were as follows: 22 minutes for the extended Brown-Forsythe test, 113 minutes for

LRTv, 16 minutes for JLSsc and 123 minutes for LRTmv. The relative computation times

between the respective variability and joint tests were even greater when the exposure was

continuous.

Application to offspring gender and gestational age on cord blood DNA

methylation

We applied the methods that demonstrated good performance in the simulation study, namely

OLS regression, the Brown-Forsythe test, JLSp and JLSsc, to identify methylation at CpG

sites that are associated with offspring gender and gestational age in the Accessible Resource

for Integrated Studies (ARIES) project. ARIES is a sub-sample of the Avon Longitudinal

Study of Parents and Children (ALSPAC) (see Methods). In ARIES, 858 children (417 male

and 441 female) were available for the analysis of gender, and after excluding offspring with

missing maternal information we were left with 708 children (345 males and 363 females) for

the analysis of gestational age (mean: 39.5 weeks, standard deviation: 1.5 weeks; Table S1).

Methylation at 8174 CpG sites were associated with gender in cord blood (through the mean,

variability or joint tests; Figure 3a and Table S2). Most of these sites were identified through

a mean difference in methylation of males and females (7642 CpGs had a mean difference

with $p < 1 \times 10^{-7}$), although 240 CpG sites were associated with a variability difference

between males and females. For instance, cg18918831 was more variable in males compared

to females (Figure 4). The joint location-and-scale tests identified 7724 of these CpG sites

(JLSp identified 7213 sites and JLSsc identified 7228 sites), including all of those with a

variability effect. Mean methylation at 5359 of these sites were associated with gender in

previous EWAS (Table S2)[22, 23, 24, 25].

Gestational age was associated with cord blood methylation at 412 CpG sites (Figure 3b and

Table S3). Most of these CpG sites were associated with a mean effect of gestational age on

methylation, and there were no CpG sites with a variability effect with $p < 1 \times 10^{-7}$. The

joint mean and variability tests identified 93.7% of the CpG site associations (JLSp identified

317 and JLSsc identified 340 CpG sites, respectively), including sites that were mostly

identified through a variability association (e.g. cg24577594; Table S3). The majority of the

CpG sites identified have been found previously in EWAS of gestational age (402 CpG sites;

Table S3)[25, 26].

9   Discussion

In this study, we have introduced a framework for testing variability using an extended

version of the Brown-Forsythe test and for jointly testing mean and variability. These

approaches were compared to the LRTs as well as other alternative methods in simulations

and were used to investigate the effect of gender and gestational age on cord blood DNA

methylation.

Without transforming the phenotype to be normally distributed, the approaches which assume

normality of the phenotype (Bartlett's test, LRTv, LRTmv and DGLM) had extremely

inflated type I error rates when faced with real methylation data. Indeed, these approaches

essentially became tests of deviations from normality and outlying values, which can have

some utility in identifying outliers caused by disease[27]. However, because of these drawbacks

these approaches are not useful for assessing variability nor joint mean and variability effects,

especially as normalizing outcome levels to overcome this problem can induce effects that

were not present prior to the transformation[21]. The extended Brown-Forsythe test and the

JLSsc approach retained correct type I error rates and performed well in comparison to the

other approaches in detecting variability and joint effects. These tests were also at least 5

times more efficient than their LRT counterparts.

Over 8000 CpG sites were associated with gender in cord blood methylation, while

methylation at 412 CpG sites were associated with gestational age. The majority of these CpG

sites were associated with effects of gender and gestational age on mean methylation.

However, 240 CpG sites were associated with differences in variability between males and

females. JLSsc identified most of the associations in both analyses, except where there was

little evidence of a mean/variability effect in the presence of a borderline effect of the other.

These methods are applicable to any area of medical research where variability and joint

effects are of interest, although they will be particularly useful for analysing high-dimensional

phenotypes where it is not possible to assess the distribution at all markers. For instance, there

has been recent interest in using variability tests to attempt to identify gene-environment

interactions, as these interactions will often cause heterogeneity in the variance across

genotypes[4, 21]. The Brown-Forsythe test has been proposed as a useful test in this scenario[21],

although the extended version presented here and elsewhere[10, 11] could be used to assess

variability trends across genotypes. Furthermore, JLSsc avoids the distributional assumptions

made by current methods proposed in the genetics literature[4, 10, 11].

The limitations of this study also warrant consideration. In the simulations and the applied

example, we only analysed DNA methylation data, although we fully expect these results to

be generalisable to all phenotypes. The application of the approaches to detect CpG sites

associated with gender and gestational age also have several limitations, especially with

regards to residual confounding. In particular, there are likely to be other important maternal

factors involved in gestation period that we have not adjusted for in our analysis. The ARIES

cohort is also not selected at random from the full ALSPAC cohort[28], and as such, the results

from this study may not generalise to the full ALSPAC cohort or the general population.

In summary, the extended Brown-Forsythe test and JLSsc are robust tests of variability and joint mean and variability effects, respectively. These tests can be used in analyses to detect associations for any type of exposure with high-dimensional phenotypes.

Methods

Modelling approaches

Location tests

Ordinary least squares (OLS) regression is commonly used to assess mean differences in methylation by an exposure. That is,

$$y_i = \alpha + x_i'\beta + \epsilon_i , i = 1, \ldots, n , \qquad [1]$$

where y_i is the outcome for the i-th individual (e.g. DNA methylation levels in epigenome-wide association studies), x_i is the exposure(s) for the i-th individual and $\epsilon_i \sim N(0, \sigma_\epsilon^2)$. OLS regression is known to be relatively robust to the underlying assumptions related to the residuals when estimating the regression coefficients (discussed further in the Supplementary Text).

Scale tests

There are several statistical tests for assessing variability differences of continuous outcome by a categorical exposure[7]. Bartlett's test[16] is perhaps the most well-known of these tests (Supplementary Text) and has been used to analyse high-dimensional phenotypes[3, 29]. However, this test is known to be very sensitive to outliers and non-normality of the outcome, which is a major cause of concern when analysing data like DNA methylation. The Brown-Forsythe test[9], on the other hand, is robust to non-normality of the outcome and outliers[8]. This test is essentially a one-way analysis of variability of the variable $z_{ij} = |y_{ij} - \tilde{y}_j|$, where y_{ij} is the methylation of the i-th individual in the j-th group and $\tilde{y}_j$ is the median of the j-th

1   group. Let $y_{ij} \sim \left(\mu_j, \sigma_j^2\right)$, where μ_j and σ_j^2 are the mean and variance of y_{ij} in the j-th group,

2   then the test statistic for $H_0: \sigma_1^2 = \sigma_2^2 = \cdots = \sigma_k^2$ is given by

$$BF = \frac{(n-k)\sum_{j=1}^k n_j \left(\bar{z}_j - \bar{z}\right)^2}{(k-1)\sum_{j=1}^k \sum_{i=1}^{n_j} \left(z_{ij} - \bar{z}_j\right)^2} \sim F_{k-1, n-k}, \qquad [2]$$

3   where k is the number of groups, n_j is the number of individuals in the j-th group and $\bar{z}_j$ and

4      $\bar{z}$ are the group mean and overall mean of z_{ij}, respectively.

5   The Brown-Forsythe test can be re-formulated as a two-stage approach[10, 11]:

6   (i)   Obtain the absolute values of the residuals from a least absolute deviation regression,

7           $d_i = |y_i - (\hat{\alpha} + x_i' \hat{\beta})|$.

8   (ii)   Test for an association between the d_i's and a function of the x_i's using a regression

9   F-test.

Since this regression framework does not depend on the exposure (x_i) being categorical, it

can also be applied to continuous exposures. Indeed, this approach has the same structure as

the Glejser and Bresuch-Pagan tests of heteroskedasticity[30, 31].

Joint location-and-scale tests

If the data are symmetrically distributed then the p-values from the location and scale tests are

independent and can be combined using Fisher's method (JLSp)[10, 11]. However, often high-

dimensional phenotypes are not all symmetrically distributed (e.g. DNA methylation at CpG

sites), which will likely lead to correlated p-values for at least some markers. Other

alternative approaches for jointly testing for mean and variability effects include likelihood-

ratio tests (LRT) comparing linear mixed models with and without including a fixed-effect

and random-effect for the exposure (LRTmv) and double generalized linear mixed models

(DGLM)[13, 14, 17] (further details in Supplementary Text). However, these tests are also

sensitive to deviations from normality and outlying values[13].

To alleviate some of the issues involved in testing for mean and variability effects

simultaneously, we have developed a joint location-and-scale score test (JLSsc). This

approach essentially combines a location test and scale test, while accounting for the

correlation between these tests. We propose to test the joint null hypothesis $H_0\colon \beta = \delta = 0$ in

the model specification:

$$y_i = \alpha + x_i'\beta + \varepsilon_i$$

$$(y_i - \bar{y})^2 = \lambda + x_i'\delta + u_i \,, \tag{3}$$

where x_i is a (k_x) vector of exposures and $\bar{y}$ is the sample average of y_i. The first part,

$H_0\colon \beta = 0$, is the null hypothesis that x does not affect the mean of y. The second part,

$H_0\colon \delta = 0$, is the null hypothesis that x does not affect the variability of y.

Let $\tilde{y}_i = y_i - \bar{y}$, $\tilde{x}_i = x_i - \bar{x}$ and $\tilde{d}_i = \tilde{y}_i^2 - \hat{\sigma}^2$, where $\hat{\sigma}^2 = \frac{1}{n}\sum_{i=1}^n \tilde{y}_i^2$. Further, let the

$n \times k_x$ matrix $\tilde{X} = [\tilde{x}_i']$ and the n vectors $\tilde{y} = (\tilde{y}_i)$ and $\tilde{d} = (\tilde{d}_i)$. Then the OLS estimators

for β and δ are given by

$$\hat{\beta} = (\tilde{X}'\tilde{X})^{-1}\tilde{X}'\tilde{y}$$

$$\hat{\delta} = (\tilde{X}'\tilde{X})^{-1}\tilde{X}'\tilde{d} \,. \tag{4}$$

14   Let $\theta = \begin{pmatrix} \beta \\ \delta \end{pmatrix}, \hat{\theta} = \begin{pmatrix} \hat{\beta} \\ \hat{\delta} \end{pmatrix}$ and $\hat{\Sigma} = \frac{1}{n}\sum_{i=1}^n \begin{bmatrix} \tilde{y}_i^2 & \tilde{y}_i\tilde{d}_i \\ \tilde{y}_i\tilde{d}_i & \tilde{d}_i^2 \end{bmatrix}$. The estimator for the variance of $\hat{\theta}$

15   under the null that $\beta = \delta = 0$ and the additional assumption that the conditional skewness

16   and kurtosis of y_i do not vary with the values of x_i, is then given by

$$V\hat{a}r(\hat{\theta}) = \hat{\Sigma} \otimes (\tilde{X}'\tilde{X})^{-1} \,. \tag{5}$$

1   Hence, the score test for $H_0: \beta = \delta = 0$ or $H_0: \theta = 0$, is given by

$$S = \hat{\theta}' \left(\hat{\Sigma}^{-1} \otimes \left(\tilde{X}' \tilde{X} \right) \right) \hat{\theta} \xrightarrow{d} \chi^2_{2k_x} \ . \tag{6}$$

Additional terms such as the square of a continuous exposure, especially useful for modelling

the relationship with outcome variability, can be added as part of x_i vector and would be

included in both parts of the test. Covariates are regressed out of both the outcome and

exposure variables by taking residuals from OLS regression prior to analysis with JLSsc.

Further details of JLSsc are discussed in the Supplementary Text.

We have developed an R package to perform these tests available at:

https://github.com/jrs95/jlst.

Simulation study

We assessed the performance of the location and scale tests as well as the joint location-and-

scale tests with both binary and continuous exposures in a simulation study based on

methylation data. We assessed the performance of OLS regression, Bartlett's test (for

simulations with a binary exposure), Brown-Forsythe test, LRT comparing mixed models

with and without a variability effect (LRTv), JLSsc, JLSp, LRTmv and DGLM. For

approaches which failed to adequately control type I error rates, we repeated the tests after

applying M-value (i.e. $\log_2(y_i/(1 - y_i)))$[20] and inverse normal rank transformations to the

methylation levels. This simulation study was performed based on data from the Tsaprouni *et*

al. study[18], which investigated the relationship between smoking and DNA methylation (data

accessible at NCBI GEO database[19], accession GSE50660).

Type I error simulations were performed by randomly generating a binary or continuous

exposure (uncorrelated with mean or variability of any of the methylation levels) and testing

the associations across all CpG sites in Tsaprouni *et al.*. To generate datasets with varying

sample size (100, 500, 1000 and 10,000 samples), samples were randomly sampled with

replacement from the Tsaprouni *et al.* dataset (Supplementary Text). The binary and

categorical exposures were randomly generated using $Ber(0.5)$ and $N(0,1)$, respectively.

Quantile-quantile (QQ) plots were used to assess deviations from normality and detect

outlying test statistics.

Power simulations were performed using the same exposure distributions as above and setting

these exposures to affect the mean and variability of methylation. In each simulation replicate,

one CpG was selected at random from the Tsaprouni *et al.* dataset, the mean and standard

deviation of this CpG site were used to set the average methylation and to generate mean and

variability effects (Supplementary Text). The mean and variability effects of the exposure on

methylation were simulated using normal distributions, while the residual error was simulated

to be either normally distributed, heavy-tailed or skewed (Supplementary Text). We also

performed simulations for a categorical exposure with three categories, adding a squared

exposure term to JLSsc for a continuous exposure, and where we generated an outlying value

(Supplementary Text). Statistical power was calculated as the proportion of simulation

replicates where either the location, scale or joint test had $p < 1 \times 10^{-7}$. For each simulation

scenario, 1000 simulation replicates were performed for a sample size of 1000 samples.

The computational time of the extended Brown-Forsythe test and JLSsc were compared to

their equivalent LRTs for 100,000 randomly selected CpGs from the Tsaprouni *et al.* dataset

for the binary and continuous exposures describe above. This analysis was performed using

one core (2.6 GHz; 4GB) on a linux server.

Application to offspring gender and gestational age on cord blood DNA

methylation

Study population

This study used DNA methylation data generated as part of the Avon Longitudinal Study of

Parents and Children (ALSPAC)[32, 33]. ALSPAC recruited 14,541 pregnant women with

expected delivery dates between April 1991 and December 1992. Of these initial pregnancies,

there were 14,062 live births and 13,988 children who were alive at 1 year of age. Please note

that the study website contains details of all the data that is available through a fully

searchable data dictionary and variable search tool

(http://www.bristol.ac.uk/alspac/researchers/our-data/). Ethical approval for the study was

obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics

Committees. Informed consent for the use of data collected via questionnaires and clinics was

obtained from participants following the recommendations of the ALSPAC Ethics and Law

Committee at the time. Consent for biological samples has been collected in accordance with

the Human Tissue Act (2004).

As part of the Accessible Resource for Integrated Studies (ARIES) project

(http://www.ariesepigenomics.org.uk)[28], a sub-sample of 1018 ALSPAC child–mother pairs

had DNA methylation measured. The ARIES participants were selected based on availability

of DNA samples at two time-points for the mother (antenatal and at follow-up when the

offspring was in adolescence) and at three time-points for the offspring (neonatal from cord

blood, childhood (age 7) and adolescence (age 17)).

Laboratory methods, quality control and pre-processing

The laboratory methods and quality control procedures used have been described elsewhere[34].

In brief, the DNA methylation wet laboratory and pre-processing analyses were performed at

1 the University of Bristol as part of the ARIES project, where the Infinium

2 HumanMethylation450 BeadChip[35] was used to measure genome-wide DNA methylation

3 levels at over 485,000 CpG sites. The methylation level at each CpG site was calculated as a

4 beta value: the ratio of the methylated probe intensity and the overall intensity. These beta

5 values range from 0 (no methylation) to 1 (complete methylation). The samples were

6 processed using functional normalization with the meffil package[36, 37]. Further quality control

7 procedures are described in the Supplementary Text.

8 *Statistical analysis*

9 To investigate the mean and variability effects of gender and gestational age (in weeks,

10 Supplementary Text) on cord blood methylation, we used the approaches which controlled

11 type I error rates without transforming methylation levels, namely OLS regression, the

12 Brown-Forsythe test, JLSp and JLSsc. All analyses were adjusted for cell counts estimated

13 using the method described by de Goede *et al.* for cord blood methylation[38]. We further

14 adjusted for 20 surrogate variables to account for residual batch effects[39]. The gestational age

15 analysis was further adjusted for offspring gender and whether the birth was by caesarean

16 section as well as for maternal characteristics: age, smoking, pre-pregnancy BMI and weight,

17 parity, education, family social class and alcohol intake during pregnancy. CpGs were

18 considered to be associated with either gender or gestational age if one of the location, scale

19 or joint tests had $p < 1 \times 10^{-7}$.

20 All analyses were performed using R (version 3.5.2).

Funding

22 This work was supported by an MRC Methodology Research Grant [grant number

23 MR/M025020/1]. Work was performed in the MRC Integrative Epidemiology Unit [grant

24 numbers MC_UU_00011/1 and MC_UU_00011/3]. The UK Medical Research Council and

Wellcome [grant number 102215/2/13/2] and the University of Bristol provide core support for ALSPAC. A comprehensive list of grants funding is available on the ALSPAC website (http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf); this research was specifically funded by BBSRC [grant numbers BBI025751/1 and BB/I025263/1], MRC [grant numbers MC_UU_00011/1, MC_UU_00011/3, MC_UU_12013/9 and MC_UU_00011/5], National Institute of Child and Human Development [grant number R01HD068437], NIH [grant number 5RO1AI121226-02] and CONTAMED EU collaborative Project [grant number 212502]. This study was supported by the NIHR Biomedical Research Centre at University Hospitals Bristol and Weston NHS Foundation Trust and the University of Bristol. The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care. This publication is the work of the authors and Kate Tilling will serve as the guarantor for the contents of this paper.

Acknowledgements

We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses.

Author contributions

J.R.S. and M.S. performed the analyses. F.W. and J.R.S. developed the methodology. All authors contributed to the conception of the study and writing the manuscript.

Conflicts of interest

J.R.S became a full-time employee of UCB while this manuscript was being written.

References

1. Feinberg AP, Irizarry RA. Evolution in health and medicine Sackler colloquium: Stochastic epigenetic variation as a driving force of development, evolutionary adaptation, and disease. *Proc Natl Acad Sci U S A* **107 Suppl 1**, 1757-1764 (2010).

2. Kitano H. Biological robustness. *Nat Rev Genet* **5**, 826-837 (2004).

3. Ek WE, Rask-Andersen M, Karlsson T, Enroth S, Gyllensten U, Johansson A. Genetic variants influencing phenotypic variance heterogeneity. *Hum Mol Genet* **27**, 799-810 (2018).

4. Young AI, Wauthier FL, Donnelly P. Identifying loci affecting trait variability and detecting interactions in genome-wide association studies. *Nat Genet* **50**, 1608-1614 (2018).

5. Flanagan JM. Epigenome-wide association studies (EWAS): past, present, and future. *Methods Mol Biol* **1238**, 51-63 (2015).

6. Paul DS, Beck S. Advances in epigenome-wide association studies for common diseases. *Trends Mol Med* **20**, 541-543 (2014).

7. Conover WJ, Johnson ME, Johnson MM. A Comparative-Study of Tests for Homogeneity of Variances, with Applications to the Outer Continental-Shelf Bidding Data. *Technometrics* **23**, 351-361 (1981).

8. Li X*, et al.* A Comparative Study of Tests for Homogeneity of Variances with Application to DNA Methylation Data. *PLoS One* **10**, e0145295 (2015).

9. Brown MB, Forsythe AB. Robust Tests for Equality of Variances. *Journal of the American Statistical Association* **69**, 364-367 (1974).

10. Soave D*, et al.* A Joint Location-Scale Test Improves Power to Detect Associated SNPs, Gene Sets, and Pathways. *Am J Hum Genet* **97**, 125-138 (2015).

11. Soave D, Sun L. A generalized Levene's scale test for variance heterogeneity in the presence of sample correlation and group uncertainty. *Biometrics* **73**, 960-971 (2017).

12. Ahn S, Wang T. A powerful statistical method for identifying differentially methylated markers in complex diseases. *Pac Symp Biocomput*, 69-79 (2013).

13. Cao Y, Wei P, Bailey M, Kauwe JSK, Maxwell TJ. A versatile omnibus test for detecting mean and variance heterogeneity. *Genet Epidemiol* **38**, 51-59 (2014).

14. Smyth GK. Generalized Linear-Models with Varying Dispersion. *Journal of the Royal Statistical Society Series B-Methodological* **51**, 47-60 (1989).

15. Li X, Fu Y, Wang X, Qiu W. Robust joint score tests in the application of DNA methylation data analysis. *BMC Bioinformatics* **19**, 174 (2018).

16. Bartlett MS. Properties of sufficiency and statistical tests. *Proceedings of the Royal Society of London Series a-Mathematical and Physical Sciences* **160**, 0268-0282 (1937).

17. Goldstein H. *Multilevel statistical models*, 4th edn. Wiley (2011).

18. Tsaprouni LG*, et al.* Cigarette smoking reduces DNA methylation levels at multiple genomic loci but the effect is partially reversible upon cessation. *Epigenetics* **9**, 1382-1396 (2014).

19. Edgar R, Domrachev M, Lash AE. Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res* **30**, 207-210 (2002).

20. Du P*, et al.* Comparison of Beta-value and M-value methods for quantifying methylation levels by microarray analysis. *BMC Bioinformatics* **11**, 587 (2010).

21. Wang H*, et al.* Genotype-by-environment interactions inferred from genetic effects on phenotypic variability in the UK Biobank. *bioRxiv*, (2019).

22. Shah S*, et al.* Genetic and environmental exposures constrain epigenetic drift over the human life course. *Genome Res* **24**, 1725-1733 (2014).

23. Yousefi P, Huen K, Dave V, Barcellos L, Eskenazi B, Holland N. Sex differences in DNA methylation assessed by 450 K BeadChip in newborns. *BMC Genomics* **16**, 911 (2015).

24. Singmann P*, et al.* Characterization of whole-genome autosomal differences of DNA methylation between men and women. *Epigenetics Chromatin* **8**, 43 (2015).

25. Spiers H*, et al.* Methylomic trajectories across human fetal brain development. *Genome Res* **25**, 338-352 (2015).

26. Bohlin J*, et al.* Prediction of gestational age based on genome-wide differentially methylated regions. *Genome Biol* **17**, 207 (2016).

27. Teschendorff AE, Jones A, Widschwendter M. Stochastic epigenetic outliers can define field defects in cancer. *BMC Bioinformatics* **17**, 178 (2016).

28. Relton CL*, et al.* Data Resource Profile: Accessible Resource for Integrated Epigenomic Studies (ARIES). *Int J Epidemiol* **44**, 1181-1190 (2015).

29. Teschendorff AE, Widschwendter M. Differential variability improves the identification of cancer risk markers in DNA methylation studies profiling precursor cancer lesions. *Bioinformatics* **28**, 1487-1494 (2012).

30. Glejser H. A New Test for Heteroskedasticity. *Journal of the American Statistical Association* **64**, 316-323 (1969).

31. Breusch TS, Pagan AR. A simple test for heteroscedasticity and random coefficient variation. *Econometrica* **47**, 1287-1294.

32. Fraser A*, et al.* Cohort Profile: the Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. *Int J Epidemiol* **42**, 97-110 (2013).

33. Boyd A*, et al.* Cohort Profile: the 'children of the 90s'--the index offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol* **42**, 111-127 (2013).

34. Richmond RC*, et al.* Prenatal exposure to maternal smoking and offspring DNA methylation across the lifecourse: findings from the Avon Longitudinal Study of Parents and Children (ALSPAC). *Hum Mol Genet* **24**, 2201-2217 (2015).

35. Dedeurwaerder S, Defrance M, Calonne E, Denis H, Sotiriou C, Fuks F. Evaluation of the Infinium Methylation 450K technology. *Epigenomics* **3**, 771-784 (2011).

36. Fortin JP*, et al.* Functional normalization of 450k methylation array data improves replication in large cancer studies. *Genome Biol* **15**, 503 (2014).

37. Min J, Hemani G, Davey Smith G, Relton CL, Suderman M. Meffil: efficient normalisation and analysis of very large DNA methylation samples. *bioRxiv*, (2017).

38. de Goede OM*, et al.* Nucleated red blood cells impact DNA methylation and expression analyses of cord blood hematopoietic cells. *Clin Epigenetics* **7**, 95 (2015).

39. Leek JT, Storey JD. Capturing heterogeneity in gene expression studies by surrogate variable analysis. *PLoS Genet* **3**, 1724-1735 (2007).

Figures

Figure 1: QQ plots for type I error simulations using a binary exposure and 1000 samples. a) OLS (mean test); b) Brown-Forsythe (variability test); c) JLSp (joint test); and d) JLSsc (joint test).

Figure 2: Power simulation results comparing approaches for identifying CpG sites associated with either a mean and/or a variance effect with the exposure at   . a) & b) are plots for a binary exposure and c) & d) are plots for a continuous exposure.

Figure 3: Miami plots for the mean (OLS) and variability (Brown-Forsythe test) associations of methylation with gender a) and gestational age b). The dark red and blue lines represent the threshold and the orange points are CpG sites that are associated with a variance effect.

Figure 4: Methylation variability plot for gender.