Visual mental imagery engages the left fusiform gyrus, but not the early visual cortex: a meta-analysis of neuroimaging evidence

Alfredo Spagna[1,2], Dounia Hajhajate[2], Jianghao Liu[2,3], Paolo Bartolomeo[2,*]

[1] Department of Psychology, Columbia University in the City of New York, NY, USA, 10027

[2] Sorbonne Université, Inserm U 1127, CNRS UMR 7225, Paris Brain Institute, ICM, Hôpital de la Pitié-Salpêtrière, F-75013 Paris, France

[3] Dassault Systèmes, Vélizy-Villacoublay, France

* Corresponding author

Paolo Bartolomeo, M.D., Ph. D.

Inserm U 1127, CNRS UMR 7225, Sorbonne Université, Institut du Cerveau et de la Moelle

épinière, ICM, Hôpital de la Pitié-Salpêtrière, 75013 Paris, France

paolo.bartolomeo@icm-institute.org

Abstract

The dominant neural model of visual mental imagery (VMI) stipulates that memories from the medial temporal lobe acquire sensory features in early visual areas. However, neurological patients with damage restricted to the occipital cortex typically show perfectly vivid VMI, while more anterior damages extending into the temporal lobe, especially in the left hemisphere, often cause VMI impairments. Here we present two major results reconciling neuroimaging findings in neurotypical subjects with the performance of brain-damaged patients: (1) a large-scale meta-analysis of 46 fMRI studies, which revealed that VMI engages fronto-parietal networks and a well-delimited region in the left fusiform gyrus. (2) A Bayesian analysis showing no evidence for imagery-related activity in early visual cortices. We propose a revised neural model of VMI that draws inspiration from recent cytoarchitectonic and lesion studies, whereby fronto-parietal networks initiate, modulate, and maintain activity in a core temporal network centered on the fusiform imagery node, a high-level visual region in the left fusiform gyrus.

Keywords: fronto-parietal networks; attention; working memory; fusiform gyrus; temporal lobe.

Introduction

Close your eyes and think of Leonardo da Vinci's *Monna Lisa*. Is she looking at you or not? Is

her hair curled or straight? Visual Mental Imagery (VMI) is the set of abilities whereby we can

"see" things that are elsewhere (or nowhere: now imagine *Monna Lisa* frowning at you). These

capacities are important for predicting the outcome of everyday tasks (Moulton & Kosslyn,

2009), for example to decide whether our car fits in a narrow parking spot. The subjective

vividness of visual mental images varies substantially across individuals (Cui, Jeter, Yang,

Montague, & Eagleman, 2007; Dijkstra, Bosch, & van Gerven, 2017; Galton, 1880; J. Pearson,

Naselaris, Holmes, & Kosslyn, 2015), with some individuals experiencing mental images as

"quasi-visual" in nature, while others having less vivid images, down to the total absence of VMI

experience in otherwise normal individuals, a condition dubbed as "aphantasia" (de Vito &

Bartolomeo, 2016; Fulford et al., 2018; Jacobs, Schwarzkopf, & Silvanto, 2018; Zeman, Dewar,

& Della Sala, 2015). The neural bases of this remarkable set of cognitive functions are the object

of intense research efforts (Dentico et al., 2014; Dijkstra, Mostert, Lange, Bosch, & van Gerven,

2018; Dijkstra, Zeidman, Ondobaka, van Gerven, & Friston, 2017; Ishai, Ungerleider, & Haxby,

2000; D. G. Pearson, Deeprose, Wallace-Hadrill, Burnett Heyes, & Holmes, 2013; J. Pearson et

al., 2015; Winlove et al., 2018). Identifying the brain circuits supporting VMI and motor imagery

is also essential for clinical reasons, because detecting their activity in neuroimaging can reveal

consciousness in non-communicating patients in apparent vegetative state (Owen et al., 2006); in

addition, uncontrolled VMI activity could contribute to the vivid recollections of traumatic

memories resulting in post-traumatic stress disorder (Mary et al., 2020).

The dominant model of VMI (Dijkstra, Bosch, & van Gerven, 2019; J. Pearson et al.,

2015) stipulates the existence of common neural substrates underlying VMI and visual

perception, spanning across the ventral cortical visual stream, with a crucial implication of early

visual areas, providing the sensory and spatial representational content of VMI (Kosslyn,

Thompson, & Ganis, 2006; J. Pearson, 2019, 2020). Neuroimaging studies in healthy volunteers

supported a strong version of the model, by demonstrating the engagement of early, occipital

visual areas in VMI (Ibáñez-Marcelo, Campioni, Phinyomark, Petri, & Santarcangelo, 2019;

Kosslyn, Ganis, & Thompson, 2001). Further, TMS interference on V1 was shown to impact

VMI (Kosslyn et al., 1999). The model also provided a principled account of inter-individual

differences in VMI, because the level of activation in low-level visual areas correlates with the

subjective experience of VMI "vividness" (Dijkstra, Bosch, et al., 2017; but see Fulford et al.,

2018; Lee, Kravitz, & Baker, 2012). However, this neuroimaging evidence is correlational, and

does not directly speak to the causal role of these structures in VMI. In fact, causal evidence

from neurological patients is sharply discordant with an implication of early visual areas in VMI

(Bartolomeo, 2002, 2008; Bartolomeo, Bourgeois, Bourlon, & Migliaccio, 2013; Bartolomeo,

Hajhajate, Liu, & Spagna, 2020). Whereas the model would predict a systematic co-occurrence

of perceptual and imaginal deficits after brain damage (Farah, Levine, & Calvanio, 1988),

patients with brain damage restricted to the occipital cortex often have spared VMI abilities

(Aglioti, Bricolo, Cantagallo, & Berlucchi, 1999; Behrmann, Winocur, & Moscovitch, 1992;

Chatterjee & Southwood, 1995; Policardi et al., 1996), with preserved subjective VMI vividness

despite damaged early visual cortex (Goldenberg, Mullbacher, & Nowak, 1995), even in the case

of bilateral cortical blindness (Chatterjee & Southwood, 1995; de Gelder, Tamietto, Pegna, &

Van den Stock, 2015; Zago et al., 2010). Instead, deficits of VMI for object form, object color,

faces or orthographic material typically arise as a consequence of more anterior damage (Basso,

Bisiach, & Luzzatti, 1980; Beschin, Cocchini, Della Sala, & Logie, 1997; Bisiach & Luzzatti,

1978; Bisiach, Luzzatti, & Perani, 1979; Farah et al., 1988; Manning, 2000; Riddoch, 1990;

Sirigu & Duhamel, 2001), often extensively including the temporal lobes, especially in the left

hemisphere (Bartolomeo, 2002, 2008; Moro, Berlucchi, Lerch, Tomaiuolo, & Aglioti, 2008).

Such a strong evidence of dissociation is at odds with models proposing a crucial implication of

early visual areas in VMI and suggests an engagement in VMI of higher-order associative areas,

especially in the left temporal lobe, rather than early visual cortex. Also, growing evidence

shows that the temporal dynamics of VMI and perception are different (Daselaar, Porat,

Huijbers, & Pennartz, 2010; Dijkstra et al., 2018; Ganis, Thompson, & Kosslyn, 2004;

Yomogida et al., 2004). VMI and perception build on distinct brain connectivity patterns

(Dijkstra, Zeidman, et al., 2017), characterized by a reversed cortical information flow (Dentico

et al., 2014), and supported by anatomical connectivity between striate, extrastriate, and parietal

areas (Whittingstall, Bernier, Houde, Fortin, & Descoteaux, 2014). In addition, studies on VMI

mainly focused on the ventral cortical visual stream, but evidence also indicates VMI-related

increases in BOLD response in fronto-parietal networks (Mazard, Laou, Joliot, & Mellet, 2005;

Mechelli, Price, Friston, & Ishai, 2004; Yomogida et al., 2004). Thus, what VMI shares with

visual perception may not be the passive, low-level perceptual aspect, but the active, exploratory

aspects of vision (Bartolomeo, 2002; Bartolomeo et al., 2013; Thomas, 1999), such as those

sustained by visual attention (Bartolomeo & Seidel Malkinson, 2019). Yet, the precise identity of

these mechanisms, and of the underlying brain networks, remains unknown.

16   To address these issues, we conducted a meta-analysis of functional magnetic resonance

imaging studies that examined the neural correlates associated with visual and motor mental

imagery. Our specific aims were to assess the role of low- and high-level visual cortex, as well as

the role of the fronto-parietal networks in mental imagery, by identifying the brain regions with

higher activation for mental imagery in healthy volunteers. Specifically, we asked whether or not

VMI recruits early visual areas, defined as retinotopically organized visual areas (i.e., visual field

maps according to e.g., Wandell & Winawer, 2011; Wang, Mruczek, Arcaro, & Kastner, 2015).

To evaluate the relationships between modality-specific and more general processes in mental

imagery, we also assessed the neuroimaging of *motor* mental imagery (MMI). The studies

included the coordinates of brain regions with a significant activation increase in the contrasts of:

VMI *versus* Control; VMI *versus* Perception; and MMI *versus* Control. We predicted the

involvement in VMI of fronto-parietal and cingulo-opercular network areas, together with high-

level areas in the visual cortical ventral stream. Based on evidence from clinical neurology, we

did not expect to find an activation of the early visual areas associated with visual mental

imagery. If confirmed, this prediction would challenge the current dominant model, postulating a

crucial implication of early visual cortex in VMI, and delineate a new, alternative model, based

on a critical role in our VMI experience of high-level visual region and of fronto-parietal

networks important for attention and visual working memory.

Method

Literature search

First, we searched for functional magnetic resonance imaging (fMRI) studies in the

domain of mental imagery using the online database NeuroSynth (http://NeuroSynth.org - RRID:

SCR_006798) (Yarkoni, Poldrack, Nichols, Van Essen, & Wager, 2011). At the time of access

(October 15, 2018), the database contained 3,884 activation peaks from 84 studies. To achieve an

exhaustive search of studies, we further expanded our literature search to other databases (i.e.,

PubMed, Google Scholar), using the following search terms: Visual mental imagery, Mental

imagery, Vividness, Mental rotation, and carefully screened the reference list of previously

conducted reviews and meta-analyses. This final step led to the discovery of 39 additional

studies.

Inclusion and Exclusion criteria

We first applied the following inclusion criteria: (1) studies published in peer-review

journals in English; (2) studies with normal healthy adults (age $\geq$ 18 years); (3) studies with

table(s) reporting brain activation foci from whole-brain analysis; (4) studies with the

coordinates of the activation reported either in the Montreal Neurological Institute (MNI) or in

the Talairach space. We then applied the following exclusion criteria: (1) single-subject studies

(excluded n = 2); (2) studies not using fMRI (e.g., voxel based morphometry studies or DTI

studies; n = 8); (3) non-first hand empirical studies such as review articles (excluded n = 7); (4)

studies of special populations of adults (e.g., with a specific personality trait or neurological

disorders), whose brain functions may deviate from neurotypical controls' (excluded n = 31); (5)

studies not reporting coordinates for contrasts of interest in this study (n = 2). (6) studies that

could not be clearly classified into one of our three subdomains, or reporting contrasts not of

interest for this study (n = 19); (7) studies using analytic approaches not of interest for this study

(e.g., ROI analyses, multivariate pattern analysis, functional or effective connectivity analysis)(n

= 15). The resulting final number of studies included in the meta-analysis was 41.

Data extraction

Within the included studies, we identified the tables that reported positive activation from

defined contrasts of (1) VMI greater than control or rest conditions; (2) VMI greater than visual

perception condition; (3) MMI greater than control or rest condition. "Control" conditions

included experimental conditions unrelated to the specific imagery task. "Visual Perception"

included all conditions in which participants viewed visual stimuli related to the items employed

in the VMI conditions. In the present work, we will use the term "experiment" as the result of

one statistical map, with a single study potentially reporting results from more than one map. In

fact, five of the 41 studies reported multiple experiments or multiple contrasts of interest for this

study (e.g., reporting coordinates for both VMI *versus* Control and VMI *versus* Perception or

VMI *versus* Control and MMI *versus* Control), which brought the total number of included

experiments to 46. Specifically, we found (1) 27 experiments (376 foci from a total of 380

subjects) reporting contrasts regarding the difference in activation between VMI and rest or

control condition; (2) 4 experiments (62 foci from a total of 52 subjects) reporting contrasts

regarding the difference in activation between visual mental imagery and perception, and (3) 15

experiments (340 foci from a total of 239 subjects) reporting contrasts regarding differences in

activation between motor mental imagery and rest. **Figure 1** illustrates the selection process

based on inclusion and exclusion criteria.

Figure 1. Flow chart of the selection process conducted in the present meta-analysis

showing the following steps: (1) identification of studies through literature searches (studies

repeated in the two searches were included only once); (2) screening for inclusion and exclusion

criteria; (3) assessment for appropriate data extraction; (4) evaluation of study eligibility based

on analysis conducted (excluding studies that did not report results from whole-brain analyses

either in the main text or in the supplementary material); (5) attribution of the included studies

to one or more contrasts (some of the studies reported coordinates contrasts that could be

included in more than one contrast).

Figure 2 shows the published activation foci from the included studies projected onto an

inflated cortical surface for the contrast of a) VMI > Control, b) VMI > Perception, c) MMI >

Control. See the supplementary videos on <u>Github</u> for a dynamical representation of the foci

included in the <u>VMI > Control</u>, <u>VMI > Perception</u>, <u>MMI > Control</u> analyses.

***Figure 2**. Activation foci reported in the studies included in the present meta-analysis for*

the contrasts a) VMI > Control, b) VMI > Perception, c) MMI > Control. For the VMI >

Control panel, coordinates are color coded to represent the three categories of contrasts

included: Object Form (cyan); Faces (yellow); Orthographic Material (orange).

Activation likelihood estimation

GingerALE calculates the overlap of activation distributions across studies by modeling

the foci as 3D Gaussian distributions centered at the reported coordinates with the full-width

half-maximum (FWHM) weighted by the sample size of each study and combining the

probabilities of activation for each voxel (Eickhoff, Bzdok, Laird, Kurth, & Fox, 2012). **Tables**

1-3 report author names, publication year, type of stimuli used in the study, type of contrast, and

number of subjects that were included in the contrasts VMI *versus* Control, VMI *versus*

Perception, and MMI *versus* Control, respectively. Further, we conducted *conjunction* and

disjunction analyses between (VMI *versus* Control) & (VMI *versus* Perception) as well as

conjunction and *disjunction* analyses between (VMI *versus* Control) & (MMI *versus* Control).

Foci included in the dataset were either reported in MNI space or converted from

Talairach space to MNI space by using the icbm2tal transform in GingerALE. The ALE map was

assessed against a null-distribution of random spatial association across studies using a non-

linear histogram integration algorithm (Eickhoff et al., 2012; Eickhoff et al., 2009). All ALE

maps were thresholded at a cluster-level corrected $p < .05$, with a cluster-forming $p < .001$ with

10,000 permutations for multiple comparison correction. Conjunction and disjunction analyses

were then conducted at a $p < .05$ threshold and 10,000 permutations.

Spatial Bayesian Latent Factor Regression for Coordinate-Based Meta-Analysis

A specific aim of this study was to assess whether or not early visual activation occurred

in the VMI vs Control condition, as a way to assess the potential contribution of early visual

areas to VMI. Because the logic of our argument rests on testing the *absence* of activation, we

conducted a spatial Bayesian latent factor regression model, where the probability that a voxel is

reported as an activated focus, is modeled via a doubly stochastic Poisson process (Montagna,

Wager, Barrett, Johnson, & Nichols, 2018). To test the activation degree of each voxel, the null

hypothesis (H_0) stated that the voxel was specifically activated as a focus during VMI vs Control

condition. We took the posterior probability $p(v)$ as the probability of null hypotheses for each

voxel. The alternative hypothesis (H_1) was that the activation of the voxel belongs to the ongoing

brain activity and was not specific. Thus, we took the mean activation probability of the whole

brain voxels to get the H1 probability (see (Montagna et al., 2018) for additional details about the

method and a general discussion of the properties of this model). We defined Bayes Factors

(Rouder, Speckman, Sun, Morey, & Iverson, 2009) for accepting and rejecting the null

hypothesis. To compute the odds hypothesis conditionally on observed data $Bayes\ Factors =$

$Pr(H0|data)/Pr(H1|data)$: BF (odds) > 3 were considered as the criteria to accept null

hypothesis. Bayes Factors (BF) (odds) < 0.33 were considered as the criteria to reject the null

hypothesis and thus to accept the alternative hypothesis. Spatial Bayesian latent factor regression

analysis was conducted on the dataset consisting of 27 studies on VMI > Control condition for a

total of 376 foci (see **Table 1** for the list of studies included in this analysis). For each study, we

also included three supplementary covariates: inference method (fixed / random effects), *p*-value

correction (corrected / uncorrected), the number of participants. We used a standard brain mask

of 91 x 109 x 91 voxels of 2mm in size. Kernels falling outside the mask were discarded. We

assigned a Gamma(1, 0.3) prior distribution with mean 1/3 to the diagonal elements of Σ^{-1}. The

sampler was run for 10,000 iterations and the algorithm converged, with the first 5,000 samples

discarded as a burn-in and collecting every 50th sample to thin the chain. We obtained the

posterior intensity values of the full brain voxels, which we used to compute the BFs.

Results

The meta-analysis conducted on all the coordinates from the 46 experiments (*all mental

imagery*) showed activation increases in areas within the fronto-parietal networks (including the

bilateral superior frontal gyrus, and the bilateral inferior parietal lobe), and the bilateral anterior

insular cortices. In addition, unilateral activation increases consistent across the studies were

found in the right precuneus and in the anterior portion of the left calcarine sulcus, in the left

fusiform and parahippocampal gyri. **Figure 3** shows the surface views of brain regions showing

consistent activation increase across all domains, while **Table 4** reports the coordinates of

common activation across all domains.

Figure 3. Brain regions showing consistently increased activation associated with mental imagery (across all the domains of VMI > Control, VMI > Perception, and MMI > Control).

The meta-analysis of the 27 experiments included in the VMI *vs* Control condition showed activation increases in bilateral superior frontal gyri, in the left anterior insular cortex, in a large cluster with a peak in the bilateral inferior parietal lobe extending towards the superior parietal lobe and to the superior portion of the middle occipital gyrus (BA19), in the left middle frontal gyrus, in the left fusiform gyrus, as well as regions lateralized to the right hemisphere, including the angular gyrus, the supplementary motor area / anterior cingulate cortex (**Figure 4** and **Table 5**).

***Figure 4.** Brain regions showing consistently increased activation associated with the contrast*

VMI > Control condition, including our hotspot in the left fusiform gyrus (which we labeled the

Fusiform Imagery Node, see Discussion), and fronto-parietal areas bilaterally.

The meta-analysis of the 4 experiments included in the VMI *vs* Perception condition

showed the activation increase in the insular cortex and in the supplementary motor area /

anterior cingulate cortex, bilaterally. **Figure 5** shows the surface views of brain regions showing

consistent activation increase in the VMI *vs* Perception condition, while **Table 6** reports the

coordinates of common activation in the VMI *vs* Perception condition.

Figure 5. *Brain regions showing consistent increased activation associated with the contrast*

VMI > Perception condition.

The meta-analysis of the 15 experiments included in the MMI *versus* Control condition showed activation increases in the superior frontal gyri bilaterally, of the cerebellum bilaterally, and of the left supplementary motor area, of the anterior portion of the left calcarine sulcus, and of the parahippocampal / fusiform gyri. **Figure 6** shows the surface views of brain regions showing consistent activation increase MMI *vs* Control condition, while **Table 7** reports the coordinates of common activation in the MMI *vs* Control condition.

Figure 6. *Brain regions showing consistent increased activation associated with the contrast*

MMI > Control condition.

To further investigate common areas of activation among the VMIs contrasts, we

conducted conjunction and disjunction analyses between the VMI *versus* Control and VMI

versus Perception contrasts (31 experiments). Results of the conjunction analysis showed

activation increases in the left anterior insular cortex and in the bilateral supplementary motor

area. Results of the both disjunction analyses showed no significant cluster of activation for the

((VMI > Control) > (VMI > Perception)) contrast. **Figure 7** shows section views of the brain

showing region with consistent activation increase in the conjunction analysis between the VMI

> Control and VMI > Perception contrasts. No significant activation foci were found from either

disjunction analysis. **Table 8** lists for the coordinates of common activation in the conjunction

analyses.

Figure 7. Brain regions showing consistent increased activation in the left anterior insular

cortex and in the bilateral supplementary motor area / anterior cingulate cortex associated with

the conjunction analysis between VMI versus Control condition and VMI versus Perception

condition. See Table 8 for the coordinates of common activation in the conjunction analysis.

The meta-analysis of the conjunction between the VMI *versus* Control and MMI *versus*

Control contrasts (42 experiments) showed activation increases in the superior frontal gyri and in

the supplementary motor area bilaterally. **Figure 8** shows the surface views of brain region

showing consistent activation increase in the conjunction and disjunction analyses between the

VMI > Control and MMI > Control contrasts. See **Table 9** for the results of the conjunction and

disjunction analyses.

Figure 8. Brain regions showing consistent increased activation in the left anterior insular

cortex and in the bilateral supplementary motor area / anterior cingulate cortex associated with

the conjunction analysis between VMI versus Control condition and MMI versus Control

condition. See Table 9 for the coordinates of common activation in the conjunction analysis.

Figure 9 shows the substantial overlap between the VMI-related activation we found in

the left fusiform gyrus and the cytoarchitectonic area FG4 (Lorenz et al., 2015), located in the

rostral fusiform gyrus.

Figure 9. The left fusiform activation we found is colored in red and circled in white, while the

mask of the probability distribution of the FG4 as shown in the coordinates reported by (Lorenz

et al., 2015) is shown in colors ranging from purple to red.

A Bayesian analysis of VMI studies

Figure 10 shows the Bayes Factor (BF) values estimated using the spatial Bayesian latent

factor regression model for the contrast VMI *vs* Control. Values in the early visual areas

bilaterally were below the lower cut-off of 0.33, indicating the absence of consistent activation

associated with VMI across the studies included in this meta-analysis.

Figure 10. *The area with color shading identifies BF values < 0.33, providing evidence for the*

absence of activation in that area. The area shaded in white indicates identifies brain regions

with BF values above 0.33.

Discussion

Lack of activation in primary sensory/motor cortices during mental imagery

The dominant model of VMI stipulates a functional and anatomical equivalence between

VMI and visual perception (J. Pearson, 2019). According to this view, VMI and visual

perception should rely upon the same cortical areas across the ventral cortical visual stream

(Dijkstra et al., 2019; J. Pearson, 2019). Progression of visual information would occur in a

bottom-up fashion during perception, and in a reversed, top-down direction during VMI. Indeed,

results from hemodynamic neuroimaging in healthy volunteers demonstrate the engagement of

early, occipital visual areas for VMI in some studies, but not in others. Notably, one study

(Kilintari, Narayana, Babajani-Feremi, Rezaie, & Papanicolaou, 2016) actually showed a relative

decrease of BOLD signal in occipital cortex during VMI. On the other hand, TMS interference

on V1 was shown to impact VMI (Kosslyn et al., 1999), but this effect might have resulted from

modulation of downstream visual areas. A TMS pulse applied over V1/V2 can stimulate not only

local neuronal assemblies, but also excitatory projecting neurons reaching through the entire

visual system, and beyond, up to FEF (Bergmann & Hartwigsen, 2020; Parkin, Ekhtiari, &

Walsh, 2015). Also, more recent investigations showed mixed results. For example, Marzi,

Mancini, and Savazzi (2009) demonstrated different reaction times to TMS-induced and

imagined phosphenes compared with those to phosphene-like dots presented on the screen. This

is a critical difference between perception and imagery, that is specific to the effects of TMS on

occipital areas. Keogh, Bergmann, and Pearson (2020) reported that imagery strength showed a

negative relationship with activation in early visual areas and a positive relationship with

activation of frontal cortex clusters. Moreover, applying tDCS to decrease excitability in visual

cortex, or to increase excitability in the superior frontal lobe, induced stronger mental images.

Thus, neurostimulation evidence is at best ambiguous concerning the role of early visual areas in

visual mental imagery, and suggests the involvement of superior frontal lobe areas, consistent

with the FEF foci we found in the present meta-analysis.

The dominant model also provides a principled account of inter-individual differences in

VMI, because the level of activation in early visual areas seem to correlate with the subjective

experience of "vividness" of visual mental images (Lee et al., 2012). Recent support to this claim

came from the finding that the content of VMI can be decoded from V1 activity (Senden,

Emmerling, Van Hoof, Frost, & Goebel, 2019). However, even the decoding studies are

correlational and not causal in nature; their findings might thus reflect nonfunctional byproducts,

instead of the true neural bases of VMI. In addition, a decoding study investigating face VMI

(VanRullen & Reddy, 2019) reported that only voxels in the temporal lobe supported above-

chance decoding of imagined faces, whereas occipital and frontoparietal regions did not perform

above chance. Our Bayesian analysis indicates that we can confidently accept the hypothesis that

VMI does not increase BOLD signal in occipital cortex when VMI is compared with control

conditions. This result is unlikely to depend on different regions of V1 being implicated in

different domains of VMI, because many of the included studies used items such as objects,

words, and faces, which are preferentially processed in foveal vision. The present evidence,

together with the extensive evidence from brain-damaged patients with intact VMI after lesions

restricted to the occipital cortex (Bartolomeo, 2002), strongly suggests that VMI does not need

activity in early visual areas. Most studies included here required participants to build mental

images of complex objects, which were often retrieved from long-term memory. Future research

should assess whether early visual areas may contribute to visual mental images of elementary

forms (e.g., lines or gratings), or to mental images evoked from short-term memory (Ishai et al.,

2000). Note, however, that some of the available evidence does not seem to support this

possibility. Shelton and Pippitt (2006), found fMRI evidence of greater parietal and prefrontal

activity in mental rotation compared to visual rotation. Broggin, Savazzi, and Marzi (2012)

found similar patterns of performance for luminance, contrast, and visual motion in visually

presented and imagined stimuli, but important differences for frequency gratings.

We also note that for the motor imagery contrast we found no evidence of involvement of the

primary motor cortex (BA4), consistent with previous meta-analytic results (Hétu et al., 2013).

The lack of primary motor cortex activation might have been related to methodological

differences in the tasks used in the studies included in the meta-analysis. Due to the

topographical organization of BA4, different subregions, controlling different parts of the body,

are meant to be activated by different tasks (e.g., drawing with the right hand should have

activated the hand-portion of M1 in the left hemisphere, while a different region might have been

activated for walking). Ultimately, this null effect may also reflect the obvious fact that actual

movement execution has to be inhibited in motor mental imagery.

Results from the contrast between VMI and perception showed that the activation of the

bilateral cingulo-opercular network (i.e., the anterior cingulate cortex and the anterior insular

cortex) was greater in VMI than in perception. This result should, however, be interpreted with

caution, given the low number of studies that fit our inclusion and exclusion criteria for the VMI

> Perception contrast (n = 4).

Left fusiform involvement in VMI

The present demonstration of robust activity in the left fusiform gyrus during VMI is

remarkable in view of its substantial overlapping with the cytoarchitectonic area FG4 (see Fig.

2), despite the wide variety of methods used in the included studies, and because of its striking

agreement with the evidence from brain-damaged patients. Impaired VMI in these patients

typically occurs after extensive damage to the temporal lobe, and especially in the left

hemisphere, at least for form and color VMI (Bartolomeo, 2002, 2008; Moro et al., 2008).

Although anatomical evidence is rarely precise in stroke patients, a recently described case report

(Thorudottir et al., 2020) provides additional, converging causal evidence on the role of the left

fusiform gyrus in VMI. After a bilateral stroke in the territory of the posterior cerebral artery, an

architect, who before the stroke could easily imagine objects and buildings, spontaneously

reported to have become virtually unable to visualize items. He had now to rely on computer-

aided design for his work, because he needed to see items that he could previously imagine. The

stroke had provoked extensive damage to the right hemisphere, including the occipital pole, the

lingual gyrus, the fusiform gyrus and the parahippocampal region. In the left hemisphere, the

lesion was smaller and affected only the medial fusiform gyrus and lingual gyrus. Comparison of

the lesion location with those of other patients with strokes in the same arterial territory, but

spared VMI, showed that the patient with impaired VMI had selective damage in the right

lingual gyrus and in the left posterior medial fusiform gyrus. The fusiform lesion was located in

close proximity to the hotspot we found in the present meta-analysis. This fusiform region might

act as an interface between semantic processing in the anterior temporal lobe (Lambon Ralph,

Jefferies, Patterson, & Rogers, 2017) and perceptual information coming from the occipital

cortex. There is, however, the possibility that VMI-related regions in the left temporal lobe are

more extended than suggested by the present fMRI results. BOLD responses in the anterior

temporal lobe suffer from large magnetic susceptibility artifacts arising from the auditory canals

(Wandell, 2011). Even when distortion-corrected sequences are used in fMRI, the lowest signal

is recorded from the region between the midfusiform region and the temporal pole. Thus,

additional, more anterior temporal regions could potentially participate in VMI-related abilities.

Fronto-parietal networks and VMI

The bilateral activation of areas within the fronto-parietal network (Corbetta, 1998; Rossi,

Pessoa, Desimone, & Ungerleider, 2009; Xuan et al., 2016) and the cingulo-opercular network

(Dosenbach, Fair, Cohen, Schlaggar, & Petersen, 2008; Dubis, Siegel, Neta, Visscher, &

Petersen, 2016; Sadaghiani & D'Esposito, 2015; Sheffield et al., 2015) suggests that mental

imagery requires the activation of task-positive neural substrates supporting high-level cognitive

functions, including those for attentional control (Bartolomeo & Seidel Malkinson, 2019;

Petersen & Posner, 2012; Wu et al., 2019; Xuan et al., 2016) and visual working memory (Awh

& Jonides, 2001; Jonikaitis & Moore, 2019). Also, the implication of fronto-parietal networks in

VMI is broadly consistent with lesion location in spatial neglect for visual mental images

(Bisiach & Luzzatti, 1978), which typically goes far beyond the occipital cortex, and implicates

the fronto-parietal networks in the right hemisphere (Bartolomeo, D'Erme, & Gainotti, 1994;

Guariglia, Padovani, Pantano, & Pizzamiglio, 1993; Rode et al., 2010). Taken together, the

results from this meta-analysis shed further light on the neural correlates of VMI, and suggest a

left-hemisphere superiority that is in line with neuropsychological evidence (Basso et al., 1980;

Goldenberg, 1992; Moro et al., 2008; Riddoch, 1990). Functional lateralization is a fundamental

organization principle of the brain, and spans across the anatomical and functional realm

(Karolis, Corbetta, & De Schotten, 2019). Besides adding additional evidence regarding the

potential lateralization of VMI, this result can be put in relation to attention processes, another

family of functions showing various types of brain lateralization (Asanowicz, Marzecova,

Jaskowski, & Wolski, 2012; Bartolomeo, 2007, 2014; Bartolomeo & Seidel Malkinson, 2019;

Chica et al., 2012; Heilman & Van Den Abell, 1980; Spagna, Kim, Wu, & Fan, 2018; Spagna,

Martella, Fuentes, Marotta, & Casagrande, 2016). Putting together the high degree of similarity

between the regions of the brain showing a consistent increase of activation associated with VMI

and those associated with visual attention, it is likely that these two functions share a common

bilateral neural substrate in the rostral frontal regions; a right hemisphere bias might emerge

when directing attention towards external stimuli (as in visual cued tasks) (Bartolomeo & Seidel

Malkinson, 2019), while a left hemisphere bias might arise when directing attention towards

internally-generated stimuli (Gazzaley & Nobre, 2012) (as in the case of mental imagery).

Given the prominent role of fronto-parietal attention networks in attention-mediated

conscious processing (Chica & Bartolomeo, 2012; Chica, Paz-Alonso, Valero-Cabre, &

Bartolomeo, 2013; Chica, Valero-Cabre, Paz-Alonso, & Bartolomeo, 2014) and in neural models

of consciousness (Dehaene & Changeux, 2011), it is tempting to speculate that individual

differences in subjective vividness of visual mental images might in part depend on different

degrees of engagement of fronto-parietal networks in VMI, or on their degree of anatomical and

functional connectivity with the ventral temporal cortex. For example, vivid re-experiencing of

traumatic memories in post-traumatic stress disorder might depend in part on poor prefrontal

control on medial temporal lobe activity (Mary et al., 2020).

A revised neural model of VMI

The present results, together with the evidence from neurological patients (Bartolomeo et al.,

2020), invite a revision of the neural model of VMI (**Figure 11**).

Figure 11. *Schematic depiction of a revised neural model of VMI. FP, fronto-parietal; WM,*

visual Working Memory; FIN, Fusiform Imagery Node; ATL, Anterior Temporal Lobe; MTL;

Medial Temporal Lobe.

We propose that a core network of VMI builds upon a region in the FG4 area (Lorenz et al.,

2015) of the left fusiform gyrus, which may be labeled as Fusiform Imagery Node (FIN), in

analogy with other domain-preferring regions in the ventral temporal cortex (Mahon &

Caramazza, 2011). This region corresponds roughly to the lateral section of left middle fusiform

region, in the vicinity of regions preferentially responding to letter strings (Cohen et al., 2002)

and to faces (although predominantly in the right hemisphere (Kanwisher, McDermott, & Chun,

1997) (but also see Bukowski, Dricot, Hanseeuw, & Rossion, 2013 for a focus on the left

fusiform gyrus). More anterior regions of the left temporal lobe (ATL) provide the FIN with

semantic information (Lambon Ralph et al., 2017). Importantly, both left and right hemisphere

ATL are implicated in conceptual knowledge (Lambon Ralph et al., 2017). This bilateral

organization might allow some patients with unilateral brain damage to compensate for deficits

of VMI (see Bartolomeo & de Schotten, 2016) As noted above, we cannot exclude the

participation of more anterior regions in the ventral temporal cortex, between the mid-fusiform

gyrus and the temporal pole, which are difficult to visualize in fMRI because of susceptibility

artifacts. During VMI, prefrontal circuits might engage activity in this core ventral temporal

network. The FIN is part of the ventral cortical visual pathway; during perception, it receives

visual information from more posterior regions. Medial temporal lobe (MTL) structures may

provide a neural substrate for recombining elements of past experiences to generate episodic

simulations in VMI (Mahr, 2020; Schacter, Addis, & Buckner, 2007). MTL structures, together

with the posterior cingulate, and perhaps with the fronto-parietal attention networks, may

contribute to the phenomenal experience of VMI as "quasi-visual" in nature (sometimes referred

to as "vividness" of visual mental images) (Fulford et al., 2018). The degree of this integration

may differ among individuals. Functional disconnection of the MTL components of the network

may lead to impaired phenomenal experience associated with VMI (aphantasia). In these cases,

some recovery of visual information (without VMI experience) may still be provided by the

ATL-FIN portion of the VMI network. Dysfunction of the ATL-FIN component may lead to

VMI deficits, as suggested by the patterns of performance of patients with extensive left

temporal damage (Bartolomeo, 2002). Domain-specific VMI might show differences in

hemispheric laterality. For example, a homologue network in the right hemisphere might

contribute to aspects of VMI for faces (Barton & Cherkasova, 2003; O'Craven & Kanwisher,

2000; Sunday, McGugin, Tamber-Rosenau, & Gauthier, 2018) (but also see Ishai et al., 2000), in

analogy with the previously mentioned right fusiform superiority in perceptual face processing

(Kanwisher et al., 1997)(but also see Bukowski et al., 2013). In our model, VMI-related activity

in the temporal lobe core network is initiated, modulated and maintained by fronto-parietal

networks subserving attention and working memory. Dysfunction of right hemisphere fronto-

parietal networks, coupled with their disconnection from the left FIN (Rode et al., 2010), but

with normal functioning of the left hemisphere portions of the network, may provoke neglect for

the left side part of visual mental images, with spared processing of their right side (imaginal or

representational neglect) (Bartolomeo et al., 1994; Bisiach & Luzzatti, 1978; Guariglia et al.,

1993). In this case, visual mental images are built by the left hemisphere VMI network, but the

attentional impairment resulting from right fronto-parietal dysfunction determines the incomplete

exploration or maintenance of the left portion of mental images (Rode et al., 2010).

Conclusions

Altogether, the long-standing neuropsychological evidence of the past 30 years, together with

more recent neuroimaging evidence analyzed here, invites a reappraisal of the dominant model

of visual mental imagery, which was based on a substantial functional and structural overlap

between visual mental imagery and visual perception. The available evidence suggests a

different scenario, where prefrontal regions initiate the imagination process by recruiting the

stored information to-be-recalled from semantic circuits involving the left temporal lobe,

together with high-level ventral stream areas of the left fusiform and parahippocampal gyri, and

then activates networks important for attention and visual working memory, which maintain the

recalled images to perform the task at hand (whether a simple inspection, a rotation, or a

judgement). This alternative model seems better equipped to deal with dramatic patterns of

double dissociation between perception and imagery abilities in brain-damaged patients. In

particular, the present finding of FIN activity in VMI is strikingly consistent with the observation

of imagery deficits occurring not after occipital damage, but after extensive left temporal damage

(Bartolomeo, 2002). Further neuroimaging and neuropsychological research is needed to

determine the brain correlates of specific domains of VMI (e.g., faces, object form or color,

places, orthographic material). A clear description of these fast-occurring dynamics can only be

achieved by employing techniques with adequate spatiotemporal resolution, such as intracerebral

recordings (see e.g., Rossion, Jacques, & Jonas, 2018), to trace feedback and feedforward sweeps

of activation in the present model.

Authors Contribution

P.B. and A.S. conceived the study. A.S., D.H., and J.L. extracted the coordinates from the source articles. A.S. and D.H. conducted the analysis on Ginger ALE. A.S. and J.L. conducted the Bayesian Analyses. All the authors contributed to the writing of the manuscript. Raw Coordinates included in each of the contrasts, the maps of the results in the .nii format as well as the raw outputs from GingerALE software, code and instructions to reproduce the Bayes analyses, instructions to reproduce the figures, and supplementary videos can be found on the [Github page of A.S]().

Acknowledgments

Supported by an ICM post-doctoral fellowship to A.S., and by funding from Dassault Systèmes. P.B. is supported by the Agence Nationale de la Rercherche through ANR-16-CE37– 0005 and ANR-10-IAIHU-06. We thank Laurent Cohen, Tal Seidel Malkinson, Silvia Montagna, Katarzyna Siuda-Krzywicka, Michel Thiebaut de Schotten, and an anonymous reviewer for helpful discussion.

Competing Interests

The authors declare no conflict of interest.

Tables

Table 1. List of studies (n = 27) showing the first author's name, publication year, manuscript title, contrast used, and number of subjects (n = 340) that were included in the ALE meta-analysis for the contrast Visual Mental Imagery > Control condition. Fixed or Random Effects and Corrected or Uncorrected variables were used as covariates in the Bayesian Analysis. A total of 372 activation foci were included in the analysis for this contrast.

Name and Year	Manuscript title	Contrast	Category	Table	N	Mean Age ±SD (min:max)	Gender	Handedness	Random / Fixed Effects	Corrected / Uncorrected
1. Anderson et al. 2019	*Visual imagery during real-time fMRI neurofeedback from occipital and superior parietal cortex*	Imagine a flickering circle in one of two visual quadrants designated by auditory instruction vs. Rest (same visual display with instruction of "stop imagining")	A	4	21	24.15 (19:31)	10M	Unknown	RFX	Corrected
2. Belardinelli et al. 2009	*An fMRI investigation on image generation in different sensory modalities: The influence of vividness*	Generate and maintain mental images on auditory instruction (e.g. "to see a candle") vs.	A	3	9	25.2 ± 3.7	9F	Right	RFX	Corrected

#	Author	Title	Paradigm								
			Rest (maintain fixation on a black cross presented at the center of a white screen).								
3.	Boccia et al. 2015	*A Penny for Your Thoughts! Patterns of fMRI Activity Reveal the Content and the Spatial Topography of Visual Mental Images*	Imagine the location designated by a visually presented word (e.g. a city in the map of Italy) vs. Baseline (unclear)	A	1	15	24.67 ± 2.16	7F	Right	RFX	Corrected
4.	Boly et al. 2007	*When thoughts become action: An fMRI paradigm to study volitional brain activity in non-communicative brain injured patients*	Imagine faces of familiar relatives on auditory instructions vs. Rest (the auditory instruction was "now please, just relax").	B	2	12	24 ± 4	12M	Unknown	RFX	Corrected
5.	D'Esposito et al. 1997	*A functional MRI study of mental image generation*	Imagine the items designated by auditorily presented words vs. Control	B	1	18-37	7M	Right	FFX	Corrected	

condition (passive

listening to abstract

words).

No.	Author	Title	Task				Age	Sex	Handedness		Correction
6.	Gardini et al. 2005	*Different neuronal pathways support the generation of general and specific mental images*	Imagine the items designated by visually presented words vs. Control condition (read silently a pseudo word).	C	1	15	25.93 ± 8.87 20:51;	6F	Right	RFX	Corrected
7.	Handy et al. 2004	*Visual imagery and memory: Do retrieval strategies affect what the mind's eye sees?*	Imagine concrete objects designated by auditorily presented words vs. Control condition (passive listening to abstract words).	A	1	15	18 ± 29	10F	Right	RFX	Uncorrected
8.	Huijbers et al. 2011	*Imagery and retrieval of auditory and visual information: Neural correlates of successful and unsuccessful performance*	Imagine items designated by auditorily presented words vs. Control	A	4	21	22	16F	Right	RFX	Uncorrected

			condition (Memory retrieval of cue-words from a previously presented list).							
9.	Ishai et al. 2000	*Distributed Neural Systems for the Generation of Visual Images*	Imagine familiar houses, faces, and chairs from long-term memory when presented with a gray square vs. Control condition (passive view of the gray square).	A/B	4	9	28 ± 5	5F	Right	RFX	Corrected
10.	Kaas et al. 2009	*Imagery of a moving object: The role of occipital cortex and human MT/V5+*	Imagine a blue ball moving back and forth in one of two quadrants vs Rest (fixation on the center cross).	A	1	12	20:34	8F	Unknown	RFX	Corrected
11.	Kilintari et	*Brain activation profiles during*	Imagine a previously	A	4	14	18:41	10F	Right	RFX	Corrected

al. 2016	*kinesthetic and visual imagery: an fMRI study*	observed videotaped action vs. Rest (avoid thinking about the movement).								
12. Kukolja et al. 2006	*Neural mechanisms underlying spatial judgements on seen and imagined visual stimuli in the left and right hemifields in men*	Imagine the position of clock hands for a given time (specified in digital numbers presented visually, e.g. 04:20 or 06:50) and judge whether their angle was <90∘ or >90∘ vs. Control condition (press both response keys when seeing "43:21").	A	3(D)	20	24.5 ± 3.2 21:32	20M	Right	RFX	Corrected
13. Lambert et al. 2002	*Neural substrates of animal mental imagery: calcarine sulcus and dorsal pathway involvement — an fMRI study*	Imagine animals upon auditory presentation of their names vs. Control condition (passive	A	2	6	24±3.5; 19 :25	4F	Right	RFX	Corrected

		listening of abstract words not conducive to MI).								
14. Logie et al. 2011	*Low and high imagers activate networks differentially in mental rotation*	Mental rotation of 3D abstract object visually presented pictures of 3-D objects vs. 0°Control condition (perceptual judgements on pairs of similar objects, either identical objects and 0°or mirror images).	A	1	21	20:35	14F	Right	RFX	Corrected
15. Mazard et al. 2005	*Neural impact of the semantic content of visual mental images and visual percepts*	Auditory instructions to imagine previously seen object drawings vs. Control condition (judge whether auditorily presented	A	1	15	18:27	4F	Right	FFX	Corrected

letters / numbers are

vowels / even).

16.	Pyke et al. 2017	*When math operations have visuospatial meanings versus purely symbolic definitions: Which solving stages and brain regions are affected?*	Solving of math problems by visuospatially-trained vs. purely symbolic solving symbolically-trained participants.	C	1	40	23.6 ± 4.9	36F	Right	FFX	Uncorrected
17.	Sack et al. 2002	*Tracking the Mind's Image in the Brain II: Transcranial Magnetic Stimulation Reveals Parietal Asymmetry in Visuospatial Imagery*	Imagine the position of clock hands in two analog clock faces on auditory instructions, and judge in which clock the hands formed a greater angle vs. rest	A	1,2	6	27.8 ± 5.7	3F	Right	FFX	Corrected
18.	Sasaoka et al. 2013	*Dynamic Parieto-premotor Network for Mental Image Transformation Revealed by Simultaneous EEG*	Imagine the Clock angle vs. Baseline Mentally rotate previously memorized clock	A	1	23	Unknown	23M	Right	RFX	Corrected

			hands while maintaining central fixation vs. Rest.								
19.	Seurinck et al. 2011	*Mental Rotation Meets the Motion Aftereffect: The Role of hV5/MT+ in Visual Mental Imagery*	Mentally rotate visually presented letters or numbers vs. Control condition (maintain central fixation).	A	1	16	24 ± 2.0; 21:28	16M	Right	RFX	Corrected
20.	Slotnick et al. 2005	*Visual Mental Imagery Induces Retinotopically Organized Activation of Early Visual Areas*	Mentally fill in imagined checkerboard wedges at the display location specified by an outer arc vs. Control condition (judge whether a red square is in the right or left hemifield)	A	1	6	21 ± 1.1	3F	Unknown	FFX	Corrected
21.	Slotnick et al. 2011	*Visual memory and visual mental imagery recruit common control and*	Imagine the items indicated by visually	A	1	12	18.6 :22.2	8F	Unknown	RFX	Corrected

	sensory regions of the brain	presented words (e.g., "ladder", "clock") vs. Control condition (press the button corresponding to the visually presented word "left," "center," or "right").								
22. Soddu et al. 2009	*Reaching across the abyss: recent advances in functional magnetic resonance imaging and their potential relevance to disorders of consciousness*	Covertly describe faces on auditory instructions vs. Rest.	A/B	1	8	Unknown	Unknown	Unknown	RFX	Corrected
23. Thompson-Schill et al, 1999	*A neural basis for category and modality specificity of semantic knowledge*	Auditorily presented questions about either visual or non-visual characteristics of living and non-living things (yes/no answers) vs. Control	C	1	5	Unknown	1F	Unknown	FFX	Corrected

			condition (listen to nonsense auditory stimuli).							
24.	Trojano et al. 2000	*Matching Two Imagined Clocks: the Functional Anatomy of Spatial Analysis in the Absence of Visual Stimulation*	Imagine two analogue clock faces based on auditorily presented times and judge which of the two clocks formed a greater visual angle vs. Control condition (judge which of two auditory presented times was numerically greater).	A	2	4	27; 23:32	2F	Right	RFX	Uncorrected
25.	Yomogida et al., 2004	*Mental Visual Synthesis is Originated in the Fronto-temporal Network of the Left Hemisphere*	Imagine common objects upon visual presentation of the corresponding word (e.g. "apple", "table"), vs. Baseline	A	1	23	18:31	Unknown	Unknown	FFX	Corrected

		condition (fixation).								
26. Zeman et al. 2010	*Loss of imagery phenomenology with intact visuo-spatial task performance: A case of 'blind imagination'*	Generate mental images of famous faces in response to their visually presented names vs. Control condition (passive view of strings of random letters).	B	4	10	63.4; 55:76	Males	Right	RFX	Corrected
27. Zvyangintsev et al. 2013	*Brain networks underlying mental imagery of auditory and visual information*	Imagine common objects upon visual presentation of the corresponding word vs. Control condition (counting backwards).	A	2	15	25.1 ± 5.7	7F	Right	RFX	Corrected

Category: A = Object Form; B = Faces; C = Orthographic Material.

Table 2. List of studies (n = 4) showing the first author's name, publication year, manuscript title, contrast used, and number of subjects (n = 52) that were included in the ALE meta-analysis for the contrast Visual Mental Imagery > Perception condition. Fixed or Random Effects and Corrected or Uncorrected variables were used as covariates in the Bayesian Analysis. A total of 62 activation foci were included in the analysis for this contrast.

Name and Year	Manuscript title	Contrast	Table	N	Mean Age ±SD (min:max)	Gender	Handedness	Random / Fixed Effects	Corrected / Uncorrected
1. Arzy et al. 2009	*Subjective mental time: the functional architecture of projecting the self to past and future*	Mental time imagery - self projection in different time points and events vs. Face discrimination	1	12	33.7 ± 4.2 (29:38)	5F	Right	FFX	Corrected
2. Daselaar et al. 2010	*Modality-specific and modality-independent components of the human imagery system*	Visual Imagery vs. Visual Perception	2	16	21	10F	Right	RFX	Uncorrected
3. Kukolia et al. 2006	*Neural mechanisms underlying spatial judgements on seen and imagined visual stimuli in the left and right hemifields in men*	Visual Imagery vs. Visual Perception	3	20	24.5 ± 3.2 (21:32)	20M	Right	RFX	Corrected
4. Trojano et al. 2000	*Matching Two Imagined Clocks: the Functional Anatomy of Spatial Analysis in the Absence of Visual Stimulation*	Visual imagery vs. spatial matching of visually presented clocks	3	4	27; (23:32)	2F	Right	FFX	Uncorrected

Table 3. List of studies (n = 16) showing the first author's name, publication year, manuscript title, stimuli used, and number of subjects (n = 282) that were included in the ALE meta-analysis for the contrast Motor Mental Imagery > Control condition. Fixed or Random Effects and Corrected or Uncorrected variables were used in the Bayesian Analysis. A total of 346 activation foci were included in the analysis for this contrast.

	Name and Year	Manuscript title	Stimuli used	Table	N	Mean Age ±SD (min:max)	Gender	Handedness	Random / Fixed Effects	Corrected / Uncorrected
1.	Boly et al. 2007	*When thoughts become action: An fMRI paradigm to study volitional brain activity in non-communicative brain injured patients*	Playing tennis imagery vs. Rest; Spatial navigation imagery vs. Rest	2	12	24 ± 4	12M	Unknown	RFX	Corrected
2.	Cremers et al. 2012	*Hemispheric Specialization During Mental Imagery of Brisk Walking*	Brisk walking imagery vs. Standing; Brisk walking imagery vs. Lying	1, 2	18	22.9 ± 2.5	7F	Right	RFX	Corrected
3.	Fernandez-Espejo et al. 2014	*The Clinical Utility of fMRI for Identifying Covert Awareness in the Vegetative State: A Comparison of Sensitivity between 3T and 1.5T*	Imagery of playing tennis vs. Rest; Spatial navigation imagery vs. Rest	2,4	15	23 ± 3	6F	Right	RFX	Corrected
4.	Harrington et al 2009	*The neural basis for simulated drawing and the semantic*	Imagery of drawing familiar and non familiar objects vs.	1	8	32.8 (20–52)	7F	Right	FFX	Corrected

		implications	Rest							
5.	Jahn et al. 2008	*Imaging human supraspinal locomotor centers in brainstem and cerebellum*	Imagine body movement: lying (rest condition), standing, walking and running	1,2,3	26	33.7; (21– 61)	12F	Right	FFX	Corrected
6.	Jahn et al. 2008	*Supraspinal locomotor control in quadrupeds and humans*	Imagery of running vs. Lying; Imagery of standing vs. Lying; Imagery of walking vs. Lying;	1	38	Unknown	Unknown	Unknown	FFX	Corrected
7.	Jancke et al. 2001	*The Role of the Inferior Parietal Cortex in Linking the Tactile Perception and Manual Construction of Object Shapes*	Motor imagery vs. actual squeezing; Motor imagery vs. actual modeling	2,3	12	36	2F	Right	FFX	Corrected
8.	Kilintari et al. 2016	*Brain activation profiles during kinesthetic and visual imagery: an fMRI study*	Kinesthetic mental imagery vs. Rest	3	14	(18 – 41)	10F	Right	RFX	Corrected
9.	La Fougèere et al. 2010	*Real versus imagined locomotion: A [18F]-FDG PET-fMRI comparison*	Imagery of walking vs. Imagery of lying	2	16	61.3 ± 7.8; (51-73)	7F	Right	RFX	Corrected
10.	Piefke et al. 2009	*Neurofunctional Modulation of Brain Regions by Distinct Forms of Motor Cognition and Movement Features*	Motor imagery vs. Observation of movements	5	12	25.18 ± 4.25	14M	Right	RFX	Corrected

11.	Sauvage et al. 2011	*Reevaluating brain networks activated during mental imagery of finger movements using probabilistic Tensorial Independent Component Analysis (TICA)*	Motor Imagery vs. Rest	2	8	24.4 ± 4	2F	Right	RFX	Corrected
12.	Soddu et al. 2009	*Reaching across the abyss: recent advances in functional magnetic resonance imaging and their potential relevance to disorders of consciousness*	Motor Imagery vs. Rest	1	14	Unknown	Unknown	Unknown	RFX	Corrected
13.	Wagner et al. 2008	*Mind the bend: cerebral activations associated with mental imagery of walking along a curved path*	Motor Imagery vs. Imagery of Standing	1,2,3, 4,5	12	33.5 (21– 51)	7F	Right	RFX	Corrected
14.	Whittingstall et al. 2013	*Structural network underlying visuospatial imagery in humans*	Motor Imagery vs. Count Backwards	1	18	(20-28)	3F	Right	FFX	Uncorrected
15.	Zu Eulenburg et al. 2013	*On the recall of vestibular sensations*	Motor Imagery vs. Rest	1	16	26.3 ± 3	8F	Right	RFX	Corrected

Table 4. Brain regions showed consistent increased activation across all the mental imagery domains (46 studies, 787 foci, 671 subjects).

Region	L/R	BA	x	y	z	Z	K
Supp. Motor Area	L	6	-4	-2	64	0.05	1020
Precentral Gyrus[a]	R	6	30	-4	54	0.04	463
Insular Cortex	L	47	-30	22	4	0.04	336
Precentral Gyrus[a]	L	6	-28	-4	56	0.04	511
Inferior Parietal Lobe	L	40	-38	-42	46	0.04	503
Insular Cortex	R	47	34	22	2	0.03	357
Inferior Parietal Lobe	R	40	40	-42	48	0.03	297
Middle Occipital Gyrus	L	19	-26	-76	38	0.03	741
Cerebellum Crus1	L	0	-32	-60	-36	0.03	189
Precentral Gyrus	L	6	-52	6	36	0.03	436
Precuneus	R	7	16	-68	50	0.03	114
Calcarine Sulcus	L	17	-12	-56	10	0.03	266
Fusiform Gyrus	L	37	-46	-58	-14	0.02	149
Parahippocampal Gyrus	L	37	-32	-44	-4	0.02	161

Note: x, y, z: coordinates of the corresponding peak in MNI space, in units of millimeter.

Abbreviations: ALE, activation likelihood; BA, Brodmann area; K, cluster size in number of voxels; L, left; R, right. [a] comprises the Frontal Eye Fields.

Table 5. Brain regions showed consistent increased activation in the VMI *versus* Control condition (27 studies, 376 foci, 380 subjects).

Region	L/R	BA	x	y	z	Z	K
Precentral Gyrus	R	6	30	-4	54	0.03	270
Inferior Parietal Lobe	L	40	-40	-44	46	0.03	910
Superior Parietal Lobe	L	7	-16	-64	52	0.02	
Middle Occipital Gyrus	L	19	-28	-74	30	0.02	
Supp. Motor Area	R	32	8	14	50	0.03	273
Precentral Gyrus	L	6	-26	-2	60	0.02	308
Precentral Gyrus	L	44	-52	10	38	0.02	270
Insular Cortex	L	47	-36	18	-2	0.02	238
Fusiform Gyrus	L	37	-42	-54	-18	0.02	166
Inferior Parietal Lobe	R	40	40	-42	48	0.02	310
Angular Gyrus	R	7	32	-66	48	0.02	

Note: x, y, z: coordinates of the corresponding peak in MNI space, in units of millimeter. Abbreviations: ALE, activation likelihood; BA, Brodmann area; K, cluster size in number of voxels; Regions are listed in a descending order by their peak Z value. For a cluster with multiple local peaks, the number of voxels in the whole cluster was only listed under the first local peak; L, left; R, right. [a] comprises the Frontal Eye Fields.

Table 6. Brain regions showed consistent increased activation in the VMI *versus*

Perception condition (4 studies, 62 foci, 52 subjects).

Region	L/R	BA	x	y	z	Z	K
Insular Cortex	R	47	34	22	0	0.01	76
Insular Cortex	L	48	-30	18	2	0.01	68
Suppl. Motor Area[a]	L	6	-4	10	52	0.01	88

Note: x, y, z: coordinates of the corresponding peak in MNI space, in units of millimeter.

Abbreviations: ALE, activation likelihood; BA, Brodmann area; K, cluster size in number of

voxels; L, left; R, right. [a] This cluster extended bilaterally.

Table 7. Brain regions showed consistent increased activation in the MMI *versus* Control condition contrast (15 studies, 340 foci, 239 subjects).

Region	L/R	BA	x	y	z	Z	K
Supp. Motor Area	L	6	-2	-2	66	0.04	604
Cerebellum	L	0	-32	-58	-32	0.03	181
Cerebellum	R	0	32	-62	-30	0.03	135
Precentral Gyrus [a]	L	6	-36	-4	56	0.02	123
Parahippocampal Gyrus	L	37	-32	-44	-4	0.02	100
Calcarine Sulcus	L	17	-16	-52	12	0.02	143
Middle Frontal Gyrus [a]	R	6	32	0	60	0.02	167

Note: x, y, z: coordinates of the corresponding peak in MNI space, in units of millimeter. Abbreviations: ALE, activation likelihood; BA, Brodmann area; K, cluster size in number of voxels; L, left; R, right. [a] comprises the Frontal Eye Fields.

Table 8. Brain regions showed consistent increased activation in the conjunction and disjunction analyses between VMI > Control condition (27 studies, 376 foci, 380 subjects) and VMI > Perception condition (4 studies, 62 foci, 52 subjects).

Conjunction (VMI > Control) and (VMI > Perception)							
Region	**L/R**	**BA**	**x**	**y**	**z**	**Z**	**K**
Anterior Insular Cortex	L	48	-30	18	2	0.0156	248
Supp. Motor Area[a]	L	6	-4	10	52	0.01	51
Disjunction (VMI > Control) > (VMI > Perception)							
Region	**L/R**	**BA**	**x**	**y**	**z**	**Z**	**K**
No cluster found							
Disjunction (VMI > Perception) > (VMI > Control)							
Region	**L/R**	**BA**	**x**	**y**	**z**	**Z**	**K**
No cluster found							

Note: x, y, z: coordinates of the corresponding peak in MNI space, in units of millimeter. Abbreviations: ALE, activation likelihood; BA, Brodmann area; K, cluster size in number of voxels; L, left; R, right. [a] extending bilaterally.

Table 9. Brain regions showed consistent increased activation in the conjunction and disjunction analyses between VMI > Control condition (27 studies, 376 foci, 380 subjects) and MMI > Control condition (15 studies, 340 foci, 242 subjects).

Conjunction (VMI > Control) and (MMI > Control)							
Region	**L/R**	**BA**	**x**	**y**	**z**	**Z**	**K**
Precentral Gyrus	L	6	-32	-4	50	3.00	54
Precentral Gyrus	R	6	34	-8	52	2	56
Supp. Motor Area [a]	L	6	-8	0	64	0.02	79
Disjunction (VMI > Control) > (MMI > Control)							
Region	**L/R**	**BA**	**x**	**y**	**z**	**Z**	**K**
Fusiform Gyrus	L	37	-46	-58	-20	3.00	60
Superior Occipital Lobe	R	7	28	-78	44	2.00	61
Inferior Parietal Lobe	L	40	-50	-46	40	1.00	110
Disjunction (MMI > Control) > (VMI > Control)							
Region	**L/R**	**BA**	**x**	**y**	**z**	**Z**	**K**
Suppl. Motor Area	L	6	-10	-8	70	13.00	450
Suppl. Motor Area	R	6	8	2	52	1.00	
Precentral Gyrus	L	6	-36	-6	48	6.00	52
Cerebellum Crus1	R	0	30	-64	-36	5.00	104
Fusiform Gyrus	L	37	-26	-40	-12	4.00	106
Cerebellum Crus1	L	0	-32	-58	-38	3.00	151
Lingual Gyrus	L	18	-14	-56	4	2.00	157

Note: x, y, z: coordinates of the corresponding peak in MNI space, in units of millimeter.

Abbreviations: ALE, activation likelihood; BA, Brodmann area; K, cluster size in number of

voxels; Regions are listed in a descending order by their peak Z value. For a cluster with multiple

local peaks, the number of voxels in the whole cluster was only listed under the first local peak;

L, left; R, right. [a] extending bilaterally.

Reference List

Aglioti, S., Bricolo, E., Cantagallo, A., & Berlucchi, G. (1999). Unconscious letter discrimination is enhanced by association with conscious color perception in visual form agnosia. *Current Biology, 9*(23), 1419-1422.

Asanowicz, D., Marzecova, A., Jaskowski, P., & Wolski, P. (2012). Hemispheric asymmetry in the efficiency of attentional networks. *Brain Cogn, 79*(2), 117-128. doi:10.1016/j.bandc.2012.02.014

Awh, E., & Jonides, J. (2001). Overlapping mechanisms of attention and spatial working memory. *Trends Cogn Sci, 5*(3), 119-126. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=11239812

Bartolomeo, P. (2002). The relationship between visual perception and visual mental imagery: a reappraisal of the neuropsychological evidence. *Cortex, 38*(3), 357-378.

Bartolomeo, P. (2007). Visual neglect. *Curr Opin Neurol, 20*(4), 381-386. doi:10.1097/WCO.0b013e32816aa3a3

Bartolomeo, P. (2008). The neural correlates of visual mental imagery: An ongoing debate. *Cortex, 44*(2), 107-108. Retrieved from http://dx.doi.org/10.1016/j.cortex.2006.07.001

Bartolomeo, P. (2014). The attention systems of the human brain. In *Attention Disorders After Right Brain Damage* (pp. 1-19): Springer London.

Bartolomeo, P., Bourgeois, A., Bourlon, C., & Migliaccio, R. (2013). Visual and motor mental imagery after brain damage. In S. Lacey & R. Lawson (Eds.), *Multisensory Imagery* (pp. 249-269). New York: Springer.

Bartolomeo, P., D'Erme, P., & Gainotti, G. (1994). The relationship between visuospatial and representational neglect. *Neurology, 44*, 1710-1714.

Bartolomeo, P., & de Schotten, M. T. (2016). Let thy left brain know what thy right brain doeth: Inter-hemispheric compensation of functional deficits after brain damage. *Neuropsychologia, 93*, 407-412.

Bartolomeo, P., Hajhajate, D., Liu, J., & Spagna, A. (2020). Assessing the causal role of early visual areas in visual mental imagery. *Nature Reviews Neuroscience, 21*(9), 517-517.

Bartolomeo, P., & Seidel Malkinson, T. (2019). Hemispheric lateralization of attention processes in the human brain. *Current opinion in psychology, 29C*, 90-96. Retrieved from https://doi.org/10.1016/j.copsyc.2018.12.023

Barton, J. J., & Cherkasova, M. (2003). Face imagery and its relation to perception and covert recognition in prosopagnosia. *Neurology, 61*(2), 220-225.

Basso, A., Bisiach, E., & Luzzatti, C. (1980). Loss of mental imagery: A case study. *Neuropsychologia, 18*(4-5), 435-442.

Behrmann, M., Winocur, G., & Moscovitch, M. (1992). Dissociation between mental imagery and object recognition in a brain-damaged patient. *Nature, 359*(6396), 636.

Bergmann, T. O., & Hartwigsen, G. (2020). Inferring Causality from Noninvasive Brain Stimulation in Cognitive Neuroscience. *Journal of Cognitive Neuroscience*, 1-29.

Beschin, N., Cocchini, G., Della Sala, S., & Logie, R. H. (1997). What the eyes perceive, the brain ignores: a case of pure unilateral representational neglect. *Cortex, 33*(1), 3-26.

Bisiach, E., & Luzzatti, C. (1978). Unilateral neglect of representational space. *Cortex, 14*(1), 129-133.

Bisiach, E., Luzzatti, C., & Perani, D. (1979). Unilateral neglect, representational schema and consciousness. *Brain, 102*(3), 609-618.

Broggin, E., Savazzi, S., & Marzi, C. A. (2012). Similar effects of visual perception and imagery on simple reaction time. *Quarterly Journal of Experimental Psychology, 65*(1), 151-164.

Bukowski, H., Dricot, L., Hanseeuw, B., & Rossion, B. (2013). Cerebral lateralization of face-sensitive areas in left-handers: only the FFA does not get it right. *Cortex, 49*(9), 2583-2589.

Chatterjee, A., & Southwood, M. H. (1995). Cortical blindness and visual imagery. *Neurology, 45*(12), 2189-2195.

Chica, A. B., & Bartolomeo, P. (2012). Attentional routes to conscious perception. *Frontiers in Psychology, 3*(1), 1-12. doi:10.3389/fpsyg.2012.00001

Chica, A. B., Paz-Alonso, P. M., Valero-Cabre, A., & Bartolomeo, P. (2013). Neural bases of the interactions between spatial attention and conscious perception. *Cerebral Cortex, 23*(6), 1269-1279. doi:10.1093/cercor/bhs087

Chica, A. B., Thiebaut de Schotten, M., Toba, M., Malhotra, P., Lupianez, J., & Bartolomeo, P. (2012). Attention networks and their interactions after right-hemisphere damage. *Cortex, 48*(6), 654-663. doi:10.1016/j.cortex.2011.01.009

Chica, A. B., Valero-Cabre, A., Paz-Alonso, P. M., & Bartolomeo, P. (2014). Causal contributions of the left frontal eye field to conscious perception. *Cerebral Cortex, 24*(3), 745-753. doi:10.1093/cercor/bhs357

Cohen, L., Lehéricy, S., Chochon, F., Lemer, C., Rivaud, S., & Dehaene, S. (2002). Language-specific tuning of visual cortex? Functional properties of the Visual Word Form Area. *Brain, 125*(5), 1054-1069.

Corbetta, M. (1998). Frontoparietal cortical networks for directing attention and the eye to visual locations: identical, independent, or overlapping neural systems? *Proc Natl Acad Sci U S A, 95*(3), 831-838.

Cui, X., Jeter, C. B., Yang, D., Montague, P. R., & Eagleman, D. M. (2007). Vividness of mental imagery: individual variability can be measured objectively. *Vision Res, 47*(4), 474-478. doi:10.1016/j.visres.2006.11.013

Daselaar, S. M., Porat, Y., Huijbers, W., & Pennartz, C. M. (2010). Modality-specific and modality-independent components of the human imagery system. *Neuroimage, 52*(2), 677-685. doi:10.1016/j.neuroimage.2010.04.239

de Gelder, B., Tamietto, M., Pegna, A. J., & Van den Stock, J. (2015). Visual imagery influences brain responses to visual stimulation in bilateral cortical blindness. *Cortex, 72*, 15-26. doi:10.1016/j.cortex.2014.11.009

de Vito, S., & Bartolomeo, P. (2016). Refusing to imagine? On the possibility of psychogenic aphantasia. A commentary on Zeman et al. (2015). *Cortex, 74*, 334-335. doi:10.1016/j.cortex.2015.06.013

Dehaene, S., & Changeux, J. P. (2011). Experimental and theoretical approaches to conscious processing. *Neuron, 70*(2), 200-227. doi:S0896-6273(11)00258-3 [pii] 10.1016/j.neuron.2011.03.018

Dentico, D., Cheung, B. L., Chang, J. Y., Guokas, J., Boly, M., Tononi, G., & Van Veen, B. (2014). Reversal of cortical information flow during visual imagery as compared to visual perception. *Neuroimage, 100*, 237-243. doi:10.1016/j.neuroimage.2014.05.081

Dijkstra, N., Bosch, S. E., & van Gerven, M. A. (2017). Vividness of Visual Imagery Depends on the Neural Overlap with Perception in Visual Areas. *J Neurosci, 37*(5), 1367-1373. doi:10.1523/jneurosci.3022-16.2016

Dijkstra, N., Bosch, S. E., & van Gerven, M. A. J. (2019). Shared Neural Mechanisms of Visual Perception and Imagery. *Trends Cogn Sci, 23*(5), 423-434. doi:https://doi.org/10.1016/j.tics.2019.02.004

Dijkstra, N., Mostert, P., Lange, F. P., Bosch, S., & van Gerven, M. A. (2018). Differential temporal dynamics during visual imagery and perception. *Elife, 7*. doi:10.7554/eLife.33904

Dijkstra, N., Zeidman, P., Ondobaka, S., van Gerven, M. A. J., & Friston, K. (2017). Distinct Top-down and Bottom-up Brain Connectivity During Visual Perception and Imagery. *Sci Rep, 7*(1), 5677. doi:10.1038/s41598-017-05888-8

Dosenbach, Fair, D. A., Cohen, A. L., Schlaggar, B. L., & Petersen, S. E. (2008). A dual-networks architecture of top-down control. *Trends Cogn Sci, 12*(3), 99-105. doi:10.1016/j.tics.2008.01.001

Dubis, J. W., Siegel, J. S., Neta, M., Visscher, K. M., & Petersen, S. E. (2016). Tasks driven by perceptual information do not recruit sustained BOLD activity in cingulo-opercular regions. *Cerebral Cortex, 26*(1), 192-201.

Eickhoff, S. B., Bzdok, D., Laird, A. R., Kurth, F., & Fox, P. T. (2012). Activation likelihood estimation meta-analysis revisited. *Neuroimage, 59*(3), 2349-2361.

Eickhoff, S. B., Laird, A. R., Grefkes, C., Wang, L. E., Zilles, K., & Fox, P. T. (2009). Coordinate-based activation likelihood estimation meta-analysis of neuroimaging data: A random-effects approach based on empirical estimates of spatial uncertainty. *Human brain mapping, 30*(9), 2907-2926.

Farah, M. J., Levine, D. N., & Calvanio, R. (1988). A case study of mental imagery deficit. *Brain and cognition, 8*(2), 147-164.

Fulford, J., Milton, F., Salas, D., Smith, A., Simler, A., Winlove, C., & Zeman, A. (2018). The neural correlates of visual imagery vividness—An fMRI study and literature review. *Cortex, 105*, 26-40.

Galton, F. (1880). Visualised numerals. *Nature, 21*, 252-256.

Ganis, G., Thompson, W. L., & Kosslyn, S. M. (2004). Brain areas underlying visual mental imagery and visual perception: an fMRI study. *Brain Res Cogn Brain Res, 20*(2), 226-241. doi:10.1016/j.cogbrainres.2004.02.012

Gazzaley, A., & Nobre, A. C. (2012). Top-down modulation: bridging selective attention and working memory. *Trends Cogn Sci, 16*(2), 129-135. doi:10.1016/j.tics.2011.11.014 PMID - 22209601

Goldenberg, G. (1992). Loss of visual imagery and loss of visual knowledge—a case study. *Neuropsychologia, 30*(12), 1081-1099.

Goldenberg, G., Mullbacher, W., & Nowak, A. (1995). Imagery without perception - A case study of anosognosia for cortical blindness. *Neuropsychologia, 33*(11), 1373-1382.

Guariglia, C., Padovani, A., Pantano, P., & Pizzamiglio, L. (1993). Unilateral neglect restricted to visual imagery. *Nature, 364*, 235-237.

Heilman, K. M., & Van Den Abell, T. (1980). Right hemisphere dominance for attention The mechanism underlying hemispheric asymmetries of inattention (neglect). *Neurology, 30*(3), 327-327.

Hétu, S., Grégoire, M., Saimpont, A., Coll, M.-P., Eugène, F., Michon, P.-E., & Jackson, P. L. (2013). The neural network of motor imagery: An ALE meta-analysis. *Neuroscience & Biobehavioral Reviews, 37*(5), 930-949. doi:https://doi.org/10.1016/j.neubiorev.2013.03.017

Ibáñez-Marcelo, E., Campioni, L., Phinyomark, A., Petri, G., & Santarcangelo, E. L. (2019). Topology highlights mesoscopic functional equivalence between imagery and perception: The case of hypnotizability. *Neuroimage, 200*, 437-449.

Ishai, A., Ungerleider, L. G., & Haxby, J. V. (2000). Distributed neural systems for the generation of visual images. *Neuron, 28*(3), 979-990.

Jacobs, C., Schwarzkopf, D. S., & Silvanto, J. (2018). Visual working memory performance in aphantasia. *Cortex, 105*, 61-73.

Jonikaitis, D., & Moore, T. (2019). The interdependence of attention, working memory and gaze control: behavior and neural circuitry. *Current opinion in psychology, 29*, 126-134. doi:https://doi.org/10.1016/j.copsyc.2019.01.012

Kanwisher, N., McDermott, J., & Chun, M. M. (1997). The fusiform face area: a module in human extrastriate cortex specialized for face perception. *Journal of Neuroscience, 17*(11), 4302-4311.

Karolis, V. R., Corbetta, M., & De Schotten, M. T. (2019). The architecture of functional lateralisation and its relationship to callosal connectivity in the human brain. *Nature communications, 10*(1), 1417.

Keogh, R., Bergmann, J., & Pearson, J. (2020). Cortical excitability controls the strength of mental imagery. *Elife, 9*, e50232.

Kilintari, M., Narayana, S., Babajani-Feremi, A., Rezaie, R., & Papanicolaou, A. C. (2016). Brain activation profiles during kinesthetic and visual imagery: An fMRI study. *Brain Research, 1646*, 249-261. doi:https://doi.org/10.1016/j.brainres.2016.06.009

Kosslyn, S. M., Ganis, G., & Thompson, W. L. (2001). Neural foundations of imagery. *Nat Rev Neurosci, 2*(9), 635-642. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11533731

Kosslyn, S. M., Pascual-Leone, A., Felician, O., Camposano, S., Keenan, J. P., Thompson, W. L., . . . Alpert, N. M. (1999). The role of area 17 in visual imagery: convergent evidence from PET and rTMS. *Science, 284*(5411), 167-170. Retrieved from http://www.ncbi.nlm.nih.gov/cgi-bin/Entrez/referer?http://www.sciencemag.org/cgi/content/full/284/5411/167

Kosslyn, S. M., Thompson, W. L., & Ganis, G. (2006). *The case for mental imagery*: Oxford University Press.

Lambon Ralph, M. A., Jefferies, E., Patterson, K., & Rogers, T. T. (2017). The neural and computational bases of semantic cognition. *Nature Reviews Neuroscience, 18*(1), 42.

Lee, S.-H., Kravitz, D. J., & Baker, C. I. (2012). Disentangling visual imagery and perception of real-world objects. *Neuroimage, 59*(4), 4064-4073.

Lorenz, S., Weiner, K. S., Caspers, J., Mohlberg, H., Schleicher, A., Bludau, S., . . . Amunts, K. (2015). Two New Cytoarchitectonic Areas on the Human Mid-Fusiform Gyrus. *Cerebral Cortex, 27*(1), 373-385. doi:10.1093/cercor/bhv225

Mahon, B. Z., & Caramazza, A. (2011). What drives the organization of object knowledge in the brain? *Trends Cogn Sci, 15*(3), 97-103. doi:10.1016/j.tics.2011.01.004

Mahr, J. B. (2020). The dimensions of episodic simulation. *Cognition, 196*, 104085. doi:https://doi.org/10.1016/j.cognition.2019.104085

Manning, L. (2000). Loss of visual imagery and defective recognition of parts of wholes in optic aphasia. *Neurocase, 6*(2), 111-128.

Mary, A., Dayan, J., Leone, G., Postel, C., Fraisse, F., Malle, C., . . . Gagnepain, P. (2020). Resilience after trauma: The role of memory suppression. *Science, 367*(6479), eaay8477. doi:10.1126/science.aay8477 PMID - 32054733

Marzi, C. A., Mancini, F., & Savazzi, S. (2009). Interhemispheric transfer of phosphenes generated by occipital versus parietal transcranial magnetic stimulation. *Exp Brain Res, 192*(3), 431-441.

Mazard, A., Laou, L., Joliot, M., & Mellet, E. (2005). Neural impact of the semantic content of visual mental images and visual percepts. *Brain Res Cogn Brain Res, 24*(3), 423-435. doi:10.1016/j.cogbrainres.2005.02.018

Mechelli, A., Price, C. J., Friston, K. J., & Ishai, A. (2004). Where bottom-up meets top-down: Neuronal interactions during perception and imagery. *Cerebral Cortex, 14*(11), 1256-1265. Retrieved from http://cercor.oxfordjournals.org/cgi/content/abstract/14/11/1256

Montagna, S., Wager, T., Barrett, L. F., Johnson, T. D., & Nichols, T. E. (2018). Spatial Bayesian latent factor regression modeling of coordinate-based meta-analysis data. *Biometrics, 74*(1), 342-353.

Moro, V., Berlucchi, G., Lerch, J., Tomaiuolo, F., & Aglioti, S. M. (2008). Selective deficit of mental visual imagery with intact primary visual cortex and visual perception. *Cortex, 44*(2), 109-118.

Moulton, S. T., & Kosslyn, S. M. (2009). Imagining predictions: mental imagery as mental emulation. *Philos Trans R Soc Lond B Biol Sci, 364*(1521), 1273-1280. doi:10.1098/rstb.2008.0314

O'Craven, K. M., & Kanwisher, N. (2000). Mental imagery of faces and places activates corresponding stimulus-specific brain regions. *Journal of Cognitive Neuroscience, 12*(6), 1013-1023.

Owen, A. M., Coleman, M. R., Boly, M., Davis, M. H., Laureys, S., & Pickard, J. D. (2006). Detecting Awareness in the Vegetative State. *Science, 313*(5792), 1402-1402. doi:10.1126/science.1130197

Parkin, B. L., Ekhtiari, H., & Walsh, V. F. (2015). Non-invasive human brain stimulation in cognitive neuroscience: a primer. *Neuron, 87*(5), 932-945.

Pearson, D. G., Deeprose, C., Wallace-Hadrill, S. M., Burnett Heyes, S., & Holmes, E. A. (2013). Assessing mental imagery in clinical psychology: a review of imagery measures and a guiding framework. *Clin Psychol Rev, 33*(1), 1-23. doi:10.1016/j.cpr.2012.09.001

Pearson, J. (2019). The human imagination: the cognitive neuroscience of visual mental imagery. *Nature Reviews Neuroscience, 20*(10), 624-634. doi:10.1038/s41583-019-0202-9 PMID - 31384033

Pearson, J. (2020). Reply to: Assessing the causal role of early visual areas in visual mental imagery. *Nature Reviews Neuroscience, 21*(9), 517-518.

Pearson, J., Naselaris, T., Holmes, E. A., & Kosslyn, S. M. (2015). Mental Imagery: Functional Mechanisms and Clinical Applications. *Trends Cogn Sci, 19*(10), 590-602. doi:10.1016/j.tics.2015.08.003

Petersen, S. E., & Posner, M. I. (2012). The attention system of the human brain: 20 years after. *Annu Rev Neurosci, 35*, 73-89. doi:10.1146/annurev-neuro-062111-150525

Policardi, E., Perani, D., Zago, S., Grassi, F., Fazio, F., & Ladavas, E. (1996). Failure to evoke visual images in a case of long-lasting cortical blindness. *Neurocase, 2*(5), 381-394.

Riddoch, M. J. (1990). Loss of visual imagery: A generation deficit. *Cognitive Neuropsychology, 7*(4), 249-273.

Rode, G., Cotton, F., Revol, P., Jacquin-Courtois, S., Rossetti, Y., & Bartolomeo, P. (2010). Representation and disconnection in imaginal neglect. *Neuropsychologia, 48*(10), 2903-2911.

Rossi, A. F., Pessoa, L., Desimone, R., & Ungerleider, L. G. (2009). The prefrontal cortex and the executive control of attention. *Exp Brain Res, 192*(3), 489-497. doi:10.1007/s00221-008-1642-z

Rossion, B., Jacques, C., & Jonas, J. (2018). Mapping face categorization in the human ventral occipitotemporal cortex with direct neural intracranial recordings. *Annals of the New York Academy of Sciences*.

Rouder, J. N., Speckman, P. L., Sun, D., Morey, R. D., & Iverson, G. (2009). Bayesian t tests for accepting and rejecting the null hypothesis. *Psychonomic bulletin & review, 16*(2), 225-237.

Sadaghiani, S., & D'Esposito, M. (2015). Functional characterization of the cingulo-opercular network in the maintenance of tonic alertness. *Cerebral Cortex, 25*(9), 2763-2773.

Schacter, D. L., Addis, D. R., & Buckner, R. L. (2007). Remembering the past to imagine the future: the prospective brain. *Nature Reviews Neuroscience, 8*(9), 657-661. doi:10.1038/nrn2213

Senden, M., Emmerling, T. C., Van Hoof, R., Frost, M. A., & Goebel, R. (2019). Reconstructing imagined letters from early visual cortex reveals tight topographic correspondence between visual mental imagery and perception. *Brain Structure and Function, 224*(3), 1167-1183.

Sheffield, J. M., Repovs, G., Harms, M. P., Carter, C. S., Gold, J. M., MacDonald III, A. W., . . . Barch, D. M. (2015). Fronto-parietal and cingulo-opercular network integrity and cognition in health and schizophrenia. *Neuropsychologia, 73*, 82-93.

Shelton, A. L., & Pippitt, H. A. (2006). Motion in the mind's eye: comparing mental and visual rotation. *Cognitive, Affective, & Behavioral Neuroscience, 6*(4), 323-332.

Sirigu, A., & Duhamel, J. (2001). Motor and visual imagery as two complementary but neurally dissociable mental processes. *Journal of Cognitive Neuroscience, 13*(7), 910-919.

Spagna, A., Kim, T. H., Wu, T., & Fan, J. (2018). Right hemisphere superiority for executive control of attention. *Cortex*. doi:10.1016/j.cortex.2018.12.012

Spagna, A., Martella, D., Fuentes, L. J., Marotta, A., & Casagrande, M. (2016). Hemispheric modulations of the attentional networks. *Brain and cognition, 108*, 73.

Sunday, M. A., McGugin, R. W., Tamber-Rosenau, B. J., & Gauthier, I. (2018). Visual imagery of faces and cars in face-selective visual areas. *PloS one, 13*(9), e0205041.

Thomas, N., J., T. (1999). Are theories of imagery theories of imagination?: An active perception approach to conscious mental content. *Cogntive Science, 23*(2), 207-245.

Thorudottir, S., Sigurdardottir, H. M., Rice, G. E., Kerry, S. J., Robotham, R. J., Leff, A. P., & Starrfelt, R. (2020). The Architect Who Lost the Ability to Imagine: The Cerebral Basis of Visual Imagery. *Brain Sciences, 10*(2), 59. Retrieved from https://www.mdpi.com/2076-3425/10/2/59

VanRullen, R., & Reddy, L. (2019). Reconstructing faces from fMRI patterns using deep generative neural networks. *Communications Biology, 2*(1), 193. doi:10.1038/s42003-019-0438-y

Wandell, B. A. (2011). The neurobiological basis of seeing words. *Annals of the New York Academy of Sciences, 1224*(1), 63.

Wandell, B. A., & Winawer, J. (2011). Imaging retinotopic maps in the human brain. *Vision Res, 51*(7), 718-737.

Wang, L., Mruczek, R. E., Arcaro, M. J., & Kastner, S. (2015). Probabilistic maps of visual topography in human cortex. *Cerebral Cortex, 25*(10), 3911-3931.

Whittingstall, K., Bernier, M., Houde, J. C., Fortin, D., & Descoteaux, M. (2014). Structural network underlying visuospatial imagery in humans. *Cortex, 56*, 85-98. doi:10.1016/j.cortex.2013.02.004

Winlove, C. I. P., Milton, F., Ranson, J., Fulford, J., MacKisack, M., Macpherson, F., & Zeman, A. (2018). The neural correlates of visual imagery: A co-ordinate-based meta-analysis. *Cortex, 105*, 4-25. doi:10.1016/j.cortex.2017.12.014

Wu, T., Wang, X., Wu, Q., Spagna, A., Yang, J., Yuan, C., . . . Fan, J. (2019). Anterior insular cortex is a bottleneck of cognitive control. *Neuroimage, 195*, 490-504. doi:10.1016/j.neuroimage.2019.02.042

Xuan, B., Mackie, M.-A., Spagna, A., Wu, T., Tian, Y., Hof, P. R., & Fan, J. (2016). The activation of interactive attentional networks. *Neuroimage, 129*, 308-319. doi:10.1016/j.neuroimage.2016.01.017

Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C., & Wager, T. D. (2011). Large-scale automated synthesis of human functional neuroimaging data. *Nature methods, 8*(8), 665.

Yomogida, Y., Sugiura, M., Watanabe, J., Akitsuki, Y., Sassa, Y., Sato, T., . . . Kawashima, R. (2004). Mental visual synthesis is originated in the fronto-temporal network of the left hemisphere. *Cereb Cortex, 14*(12), 1376-1383. doi:10.1093/cercor/bhh098

Zago, S., Corti, S., Bersano, A., Baron, P., Conti, G., Ballabio, E., . . . Cappellari, A. (2010). A cortically blind patient with preserved visual imagery. *Cognitive and behavioral neurology, 23*(1), 44-48.

Zeman, A., Dewar, M., & Della Sala, S. (2015). Lives without imagery - Congenital aphantasia. *Cortex, 73*, 378-380. doi:10.1016/j.cortex.2015.05.019