Analysis

The proliferative history shapes the DNA methylome of B-cell

tumors and predicts clinical outcome

Martí Duran-Ferrer[1,2,*], Guillem Clot[1,2], Ferran Nadeu[1,2], Renée Beekman[1,2], Tycho

Baumann[2,3], Jessica Nordlund[4], Yanara Marincevic-Zuniga[4], Gudmar Lönnerholm[5],

Alfredo Rivas-Delgado[1,3], Silvia Martin[1,2], Raquel Ordoñez[2,6], Giancarlo Castellano[1],

Marta Kulis[1], Ana Queirós[1], Lee Seung-Tae[7], Joseph Wiemels[8], Romina Royo[9],

Montserrat Puiggrós[9], Junyan Lu[10], Eva Gine[1,2,3], Sílvia Beà[1,2,13], Pedro Jares[1,2,13], Xabier

Agirre[2,6], Felipe Prosper[2,6,11], Carlos López-Otín[2,12], Xosé S.Puente[2,12], Christopher C.

Oakes[13], Thorsten Zenz[13,14,], Julio Delgado[1,2,3], Armando López-Guillermo[1,2,3], Elías

Campo[1,2,15] and José Ignacio Martin-Subero[1,2,15,16*]

[1]Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain.

[2]Centro de Investigación Biomédica en Red de Cáncer, CIBERONC, Spain.

[3]Servicio de Hematología, Hospital Clínic, IDIBAPS, Barcelona, Spain.

[4]Department of Medical Sciences, Molecular Medicine and Science for Life Laboratory, Uppsala

University, Uppsala, Sweden

[5]Department of Women's and Children's Health, Pediatrics, Uppsala University, Uppsala,

Sweden.

[6]Centro de Investigación Médica Aplicada (CIMA), IDISNA, Pamplona, Spain.

[7]Department of Laboratory Medicine, Yonsei University College of Medicine, Korea.

[8]Center for Genetic Epidemiology, University of Southern California, Los Angeles.

[9]Programa Conjunto de Biología Computacional, Barcelona Supercomputing Center (BSC),

Institut de Recerca Biomèdica (IRB), Spanish National Bioinformatics Institute, Universitat de

Barcelona, Barcelona, Spain.

[10]European Molecular Biology Laboratory (EMBL), Heidelberg, Germany.

[11]Hematology and Cell Therapy Department, Clínica Universidad de Navarra, Universidad de

Navarra, Avenida Pío XII, 36 31008 Pamplona, Spain.

[12]Departamento de Bioquímica y Biología Molecular, Instituto Universitario de Oncología

(IUOPA), Universidad de Oviedo, Oviedo, Spain.

[13]Division of Hematology, Department of Internal Medicine, The Ohio State University,

Columbus, OH

[14]Department of Medical Oncology and Hematology, University Hospital Zürich and University

of Zürich, Zürich, Switzerland

[15]Departament de Fonaments Clinics, Facultat de Medicina, Universitat de Barcelona,

Barcelona, Spain

[16]Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain.

*Correspondence: maduran@clinic.cat, imartins@clinic.cat

Abstract

We report a systematic analysis of the biological and clinical implications of DNA methylation variability in five categories of B-cell tumors derived from B cells spanning the entire maturation spectrum. We used 2056 primary samples including training and validation series and show that 88% of the human DNA methylome is dynamically modulated under normal and neoplastic conditions. B-cell tumors display both epigenetic imprints of their cellular origin and *de novo*, disease-specific epigenetic alterations that in part are related to differential transcription factor binding. These differential methylation patterns were used by a machine-learning approach to create a diagnostic algorithm that accurately classifies 14 B-cell tumor entities and subtypes with different clinical management. Beyond this, we identified extensive patient-specific epigenetic variability targeting constitutively silenced chromatin regions, a phenomenon we could relate to the proliferative history of normal and neoplastic B cells. We observed that, depending on the maturation stage of the tumor cell of origin, mitotic activity leaves different imprints into the DNA methylome. Subsequently, we constructed a novel DNA methylation-based mitotic clock called epiCMIT (epigenetically-determined Cumulative MIToses), whose lapse magnitude represents a strong independent prognostic variable within specific B-cell tumor subtypes and is associated with particular driver genetic alterations. Our findings reveal DNA methylation as a holistic tracker of B-cell tumor developmental history, with implications in the differential diagnosis and prediction of the outcome of the patients.

Introduction

The process of neoplastic transformation implies a dramatic alteration of cellular

identity [1]. However, cancer cells partially maintain molecular imprints of the cellular

lineage and maturation stage from which they originate [2]. B-cell neoplasms are a

paradigmatic model of this phenomenon, as the maturation stage of different B-cell

neoplasms is the main principle behind the World Health Organization classification of

these tumors [3]. Recent studies have focused on the analysis of the DNA methylome, a

bona fide epigenetic mark related to cellular identity and gene regulation [4,5], during

the entire B-cell maturation program [6] and in various B-cell neoplasms spanning the

entire maturation spectrum. These include B-cell acute lymphoblastic leukemia (ALL)

[7,8] derived from precursor B cells, mantle cell lymphoma (MCL) [9] and chronic

lymphocytic leukemia [10,11](CLL) derived from pre- and post-germinal center mature B

cells, diffuse large B-cell lymphoma (DLBCL) [12,13] derived from germinal center B cells,

and multiple myeloma (MM) [14,15] derived from terminally-differentiated plasma cells.

These studies have revealed a dynamic DNA methylome during B-cell maturation as

well as novel insights into the cellular origin, pathogenic mechanisms and clinical

behavior of B-cell neoplasms reviewed in [16]. However, a global analysis of the entire

normal cell differentiation program and derived neoplasms is neither available for B

cells nor for any other human cell lineage. Thus, we herein exploit both previously

generated DNA methylation datasets as well newly generated data to systematically

decipher the sources of DNA methylation variability across B-cell neoplasms. This

comprehensive approach using over 2000 samples reveals previously hidden biological

insights and clinical associations. In particular, *de novo* disease-specific

hypomethylation in active regulatory regions is associated with differential

transcription factor binding and targets genes important for disease-specific

pathogenesis. From the clinical perspective, we define a set of epigenetic biomarkers

that can accurately classify B-cell neoplasms requiring differential clinical management

and construct a DNA methylation-based mitotic clock called epiCMIT as a personalized

predictor of clinical behavior within each B-cell neoplasm.

Results

Shared DNA methylation dynamics in normal and neoplastic B cells

We analyzed previously published DNA methylation profiles of samples from normal

and neoplastic B cells spanning the entire B-cell differentiation spectrum, all generated

with the 450k microarray platform from Illumina. These included 10 normal B cell

subpopulations [6] as well as the five common categories of B-cell neoplasms, i.e. ALL [7,8],

MCL [9], CLL [10,17], DLBCL (own unpublished series) and MM [14] (Fig. 1a and Supplementary

Table 1). Following the guidelines of the TCGA Consortium, we selected samples

containing a tumor-cell content more than 60% https://www.cancer.gov/about-

nci/organization/ccg/blog/2018/bcr-tips. This proportion of tumor cells was estimated

by flow cytometry [6,9,10,14,17], genetic data [18] and/or lineage-specific DNA methylation

patterns (Supplementary Table 2). To validate that 60% was an appropriate threshold

for B-cell tumors, we analyzed 5 MCLs and 3 CLLs in which we had DNA extracted both

from sorted (minimum purity of 86%) and unsorted tumor cells (purity between 48%

and 77%). Unsupervised analyses showed that samples tightly clustered according to

patient number and not based on their tumor cell content (Extended Data Fig. 1a). In

fact, tumor cell content in these paired samples was reflected in a minor component

accounting for only 2% of the total variability, and therefore was considered negligible

(data not shown). Tumor cell content estimated by any of the three methods was in

general highly correlated (Extended Data Fig. 1b) and in most of the cases, we used

DNA methylation data to estimate tumor purity. However, MM samples showed that

DNA methylation-based estimation of tumor cell content was far lower than that

estimated by flow cytometry (Extended Data Fig. 1c). This result is congruent with

previous DNA methylation data indicating that MM losses the B cell identity [14].

Unexpectedly, some DLBCL samples also showed a similar effect (Extended Fig. 1c),

and therefore in MM and DLBCL, tumor cell content was estimated by flow cytometry

and genetic data, respectively. After all filtering criteria (Methods), we generated a

curated data matrix containing 1595 high quality samples (Fig. 1a and Supplementary

Table 1) with DNA methylation values for 437182 CpGs, which was used in all

downstream analyses.

This large comprehensive dataset offered an exceptional opportunity to step-

wise dissect the DNA methylation variability of normal and neoplastic B cells at

different magnitude levels, including cancer-specific, tumor entity-specific, tumor

subentity-specific and individual-specific variability (Fig. 1b). Before studying such DNA

methylation dynamics, we identified that only 12% of the studied CpGs show stable

DNA methylation levels in normal and neoplastic B cells (Fig. 1C and Supplementary

Table 3). We characterized these CpGs based on genetic location, CpG content and

chromatin states from primary samples from the Blueprint consortium [19]. We

identified that stably methylated regions were located at gene-bodies of actively

transcribed genes whereas stably unmethylated sites were prevalent in CpG islands of

promoter regions showing active chromatin marks (Fig. 1d, e). Stably methylated and

unmethylated CpGs mainly converged into the same genes (Fig. 1e and Extended Data

Fig. 1e), which are highly expressed in normal and neoplastic B cells (Fig. 1f) and are

involved in cellular functions such as cell cycle, RNA processing and energy metabolism

(Extended Data Fig. 1g). These results indicate, on the one hand, that the stable

fraction of the DNA methylome affects genes involved in fundamental cellular

functions showing an epigenetic signature of expressed genes (Extended Data Fig. 1g).

On the other hand, they show that the great majority of the DNA methylome (88%) is

labile during normal B cell differentiation and neoplastic transformation.

We next wondered whether all B-cell tumors, regardless the entity, share a

unifying DNA methylation signature related to their neoplastic nature. As 25% of the

DNA methylome is modulated during normal B cell differentiation [6,11], we took the

remaining stable 75% and compared the methylomes of normal and neoplastic B cells.

This analysis did not identify any consistent *de novo* DNA methylation signature that is

shared by all B-cell neoplasms under study. Instead, DNA methylation variability is

related to differences among B-cell tumor entities and subtypes as well as patient-

specific variability, as will be shown in the following sections.

Disease-specific hypomethylation targeting regulatory regions is associated with

specific transcription factor bindings and differential gene expression

We next aimed at studying whether disease-specific DNA methylation patterns may

unravel differential pathogenic mechanisms underlying each B-cell neoplasm. Overall,

a principal component analysis (PCA) showed that different B-cell neoplasms form

distinct clusters (Fig. 2a). A further analysis indicated that the first nine components

contained information regarding specific B-cell neoplasms and allowed us to distil the

main biological sources of DNA methylation variability (Extended Data Fig 2a). The first

component was related to B-cell development, separating neoplasms according to the

maturation stage of their cellular origin, i.e. ALL together with pre-germinal center B

cells and mature B cell neoplasms together with germinal center B cells, memory B

cells and plasma cells. The remaining components showed tumor-specific patterns,

such as PC3, PC7 and PC9 related to MM, DLBCL and MCL-specific variability,

respectively. Next, in order to identify DNA methylation signatures specifically

associated with malignant transformation, we focused our next analysis on the

genome fraction with stable (i.e. B-cell independent) DNA methylation levels across B-

cell differentiation [6] (Fig. 2b). We showed varying numbers of tumor-specific DNA

methylation (tsDNAm) changes, ranging from 616 in CLL to 49279 in MM (Fig. 2b,

Supplementary Tables 4 and 5). Remarkably, we observed that DNA methylation

changes manifested differently in distinct neoplasms. Overall, hypermethylation was

enriched at CpG islands and promoter related regions, whereas hypomethylation at

low CpG content regions such as open sea, shore and shelfs (Extended Data Fig. 2c).

ALL and DLBCL showed more tumor-specific DNA hypermethylation (tsDNAm-hyper)

whereas MCL, CLL and MM acquired more tumor-specific DNA hypomethylation

(tsDNAm-hypo), being this skew towards hypomethylation remarkable in MM (Fig. 2b-

c and Extended Data Fig. 2b). To shed light into the potential causes of this

phenomenon, we analyzed the expression levels of DNA methyltransferases (DNMTs)

during normal B-cell maturation [6]. We identified that DNMT1, which has been linked

both to DNA methylation maintenance and DNA hypermethylation of CpG islands

marked by polycomb [20], shows the highest expression levels in precursor B cells and

germinal center B cells, the respective cells of origin of the mostly hypermethylated

ALL and DLBCL. Conversely, bone marrow plasma cells, the cell of origin of the mostly

hypomethylated MM, showed the lowest levels of DNMT1 expression (Extended Data

Fig. 2d).   Although the precise mechanisms remain to be elucidated, the level of

expression of DNMTs of the respective cells of origin may influence how DNA

methylation changes are manifested in distinct B-cell neoplasms.

Transcription factors (TFs) have been reported to affect DNA methylation levels

of regulatory regions upon their binding to the DNA [21,22]. Therefore, we performed TFs

binding site prediction analysis in active regulatory elements (i.e. marked by H3K27ac)

containing tsDNAm-hypo CpG (Fig. 2d and Methods). Interestingly, the entities in

which tsDNAm-hypo was predominantly located in H3K27ac regions (Fig. 2c) showed

enrichments for binding sites of TFs expressed in each respective entity (Extended

Data Fig. 2e and Supplementary Table 6) and with a previously reported implication in

their pathogenesis, such as SPI1/SPIB and EBF1 in ALL, TCF/ZEB in MCL, and NFAT in

CLL (Fig. 2d) [23–25]. In the case of DLBCL and MM, their associated tsDNAm-hypo CpGs

were actually depleted of regulatory elements (Fig. 2c), suggesting that TF binding may

not be a major factor leading to their tumor-specific DNA methylation signatures.

However, focusing only on the underrepresented H3K27ac-containing tsDNAm-hypo

CpGs, we could also detect significant relationships with TFs potentially involved in the

respective diseases, such as FOX family in DLBCL [26], and NRL (a member of the

oncogenic MAF family), ISL1, TEAD, and YY1 in MM [27–30].

Beyond the potential role of TFs in shaping tumor-specific DNA methylation

signatures, we also investigated the downstream transcriptional consequences of

tsDNAm-hypo signatures. An analysis of transcriptional profiles of cases from all five

diseases revealed a total of 94 genes associated with tsDNAm-hypo genes expressed in

a disease-specific manner (Fig. 2e). Although some of the identified genes have been

shown to be specifically expressed in a particular disease, such as *CTLA4* and *KSR2* in

CLL [31], this comprehensive analysis provides a rich resource of novel epigenetically-

regulated candidate oncogenes for mechanistic studies in each B-cell neoplasm entity.

Accurate classification of 14 clinico-biological subtypes of B cell neoplasms using

epigenetic biomarkers

The B-cell neoplasms shown in Fig. 1a represent broad categories which are further

classified into subtypes with different clinico-biological features based on genetic,

transcriptional or epigenetic features [3]. These include high-hyperdiploid (HeH),

11q23/MLL, t(12;21), t(1;19), t(9;22) and dic(9;20) ALLs [7]; C1 (conventional; germinal

center-inexperienced) and C2 (leukemic non-nodal; germinal center-experienced)

MCLs [9]; naïve-like/low-programmed, intermediate/intermediate-programmed and

memory-like/high-programmed CLLs [10,11], and finally germinal center B cell (GCB) and

activated B cell (ABC) DLBCLs [32]. In MM, a previous report did not show methylation

differences among the distinct cytogenetic subtypes [14] and thus MM subgrouping was

not included in our analyses. Here, we focused on the identification of epigenetic

biomarkers that may allow a comprehensive diagnosis of B-cell tumors entities and

subtypes. We devised a strategy to construct a classifier algorithm that yielded 56

CpGs as the optimal number distributed along 5 predictors (Extended Data Fig. 3a-e

and Supplementary Table 7, Methods) to accurately discriminate the main B-cell tumor

entities as a first step (predictor 1), and subsequently B-cell tumor subtypes as a

second step (predictors 2, 3, 4 or 5) (Fig. 3a and Methods). The accuracy of the five

predictors was evaluated using nested 10-fold stratified cross-validation in the training

series (n=1345) and with external validation series (n=711) (Fig. 3b). Overall, we

obtained very high accuracies in the predictions in both main B-cell tumor entities

(mean sensitivities for training series = 0.97 and validation series =0.99) and B-cell

tumor subtypes (mean sensitivities for training series = 0.9 and validation series =

0.97). In some tumor subtypes, we obtained lower accuracy in the predictions mostly

due to small sample sizes. The script to easily implement the classifier is provided as

supplementary information (SI1). This epigenetic classifier may represent the basis for

a simple and accurate diagnostic tool for main B-cell tumors and B-cell tumor subtypes

with more complicated diagnosis such as subtypes of MCL or CLL.

Patient-specific DNA methylation changes are associated with silent chromatin

without an impact on gene expression

After having characterized entity-based sources of DNA methylation variability, we

next aimed at studying patient-specific changes within each tumor subtype (Fig. 1b,

level 4). To that end, we computed the total number and the number of hyper- and

hypomethylation changes in every single patient within each B-cell tumor subtype as

compared to HPC (Fig. 4a). As each B-cell tumor entity is derived from a distinct

cellular origin, this approach has the advantage of fixing a reference point for all B-cell

tumors, and the changes observed can be subsequently dissected into those

modulated in normal B-cell maturation and those taking place exclusively in the

context of neoplastic transformation (i.e. B-cell independent changes). Overall, we

found large differences in the numbers of DNA methylation changes per patient (Fig.

4a and Supplementary Table 8). To analyze whether some B-cell neoplasms show an

intrinsically more variable epigenome, we studied the degree of DNA methylation

variability in each group, but no differences were observed (Extended Data Fig. 4a).

The total number of altered CpGs per case was classified into four categories,

depending whether DNA methylation was gained or lost and whether it was

modulated or not during normal B cell development [6] (Fig. 2b and Fig. 4a). Regardless

of the cellular origin of each B-cell tumor entity, this analysis uncovered striking

correlations between the degree of B-cell related and B-cell independent DNA

methylation changes, a finding that was maintained when hypermethylation and

hypomethylation were studied separately (Fig. 4b and Extended Data Fig. 4b). This

association suggests that the overall DNA methylation burden of the tumor in each

individual patient may be shaped by a similar underlying phenomenon, which is

manifested in all four CpG categories. This statement is further supported by a

thorough annotation of the CpGs sites. Patient-specific CpGs that undergo

hypomethylation in the B-cell related and B-cell independent fractions are consistently

located in low CpG-content (open sea) low-signal heterochromatin, and the associated

genes are constitutively silent both in normal and neoplastic B cells (Fig. 4c-e and

Extended Data Fig. 4c-f). In the case of patient-specific hypermethylation, CpGs in both

fractions are located in promoter regions and CpG islands (CGIs) with H3K27me3-

repressed and poised-promoter chromatin states, and affect genes that remain silent

across normal differentiation and neoplastic transformation of B cells (Fig. 4f-h,

Extended Data Fig. 4c, g-i). These findings indicate that most DNA methylation

changes in B-cell tumor patients occur in silent chromatin regions in the absence of

concurrent phenotypic changes, suggesting that a mechanism independent from gene

regulation may underlie their overall DNA methylation landscape.

Beyond the classical role of DNA methylation as gene regulator, an accumulated body

of published evidence supports the concept that DNA methylation changes accumulate

during cell division (Fig. 4i) [33–39]. Studies in fibroblasts and hematopoietic stem cells

have reported that DNA methylation loss in late-replication heterochromatic regions

and DNA methylation gain of polycomb targets increase as cells proliferate without an

apparent impact on gene expression [33–35]. Furthermore, DNA methylation loss in

heterochromatic, late-replicating domains has been recently associated with mitotic

cell division in cancer [39]. Additionally, early studies detected preferential methylation

of CGI marked by H3K27me3 in cancer [40–42], and DNA methylation changes in regions

marked by H3K27me3 in embryonic stem cells have been recently related to mitotic

cell division in cancer [38].

In order to explore whether DNA methylation changes in silent regions reflect

the proliferative history of primary human B cells, we used DNA methylation data of an

in vitro differentiation model of primary NBCs into plasma cells (Fig. 4j) [43]. At days 4

and 6, different B cells were separated based on their proliferation history measured

by CFSE dilution. We evaluated the DNA methylation profile in repressed regions at

these time points and we detected the presence of hypermethylation at repressed

H3K27me3-containing regions and hypomethyation of low signal/H3K9me3-containing

heterochomatic regions in cells that have proliferated (Fig. 4k). This finding was

particularly marked at day 6, in which the gradual accumulation of DNA methylation

changes in silent regions was directly associated with their proliferative history (from

less divided P3 to highly divided P1). The genes associated to the measured CpGs did

not show any change of expression levels regardless of their methylation status, and

thus were unlikely to be related to the phenotype of the cells (Fig. 4l).

Collectively, all these data indicate that a great fraction of the patient-specific DNA

methylation   changes   in   B-cell   tumors   accumulate   at   H3K27me3   and   low

signal/H3K9me3 regions during cell division without affecting gene expression levels, a

finding that is consistent with an epigenetic mitotic clock.

Development and validation of an epigenetic mitotic clock reflecting the proliferative

history of normal and neoplastic B cells

We performed a step-wise selection of CpGs whose methylation status would reflect

the cell mitotic activity regardless of its normal or neoplastic nature (Methods). Using

this strategy, we finally retained 184 CpGs in stable polycomb regions that tend to gain

DNA methylation and 1,164 CpGs in constitutive heterochromatin that tend to lose

DNA methylation during normal B-cell maturation and neoplastic transformation (Fig.

5a, Supplementary Table 9 and SI2). We next constructed two scores, one for

hypermethylation and one for hypomethylation, which we respectively named

epiCMIT-hyper and epiCMIT-hypo, ranging from 0 to 1 depending on low or high

proliferative histories, respectively (Methods). Remarkably, epiCMIT-hypo contained a

significant proportion of solo-WCGW CpGs, which have been recently associated with

human mitotic cell division (N=1e5 permutations, pval<0.0001) [39]. As normal B cell

differentiation entails cell division, we initially evaluated both scores in normal B cells

and observed an expected but strikingly high correlation (R=0.96, pval<0.001), with B-

cell subpopulations distributed according to their maturation state (and thus according

to their accumulated proliferative history during B-cell differentiation) (Fig. 5b, left

panel). This finding suggests that mitotic cell division in normal B cells leaves both

hyper- and hypomethylated imprints. This high correlation between the two scores

was also observed for MCL, CLL and DLBCL (Fig. 5b) but not for ALL and MM. The

epiCMIT-hyper was greater than the epiCMIT-hypo in ALL samples, and the opposite

scenario was observed in MM. As these two neoplasias originate from B cells at the

two extremes of the maturation spectrum, our data suggests that the impact of cell

division onto the DNA methylome may be different depending on the maturation state

of the cellular origin. Upon neoplastic transformation, dividing precursor B cells do not

seem to acquire broad hypomethylation in heterochromatin but rather

hypermethylation in polycomb-repressed regions. In contrast, neoplastic plasma cells

acquire widespread hypomethylation in heterochromatin and virtually lack

hypermethylation of polycomb-repressed regions. As previously shown in Extended

Data Fig. 2d, this phenomenon may be related to the differential expression of DNMTs

of the respective cells of origin of ALL and MM. We then took into consideration both

scores to derive a unique epiCMIT score (Fig. 5c). The epiCMIT reflects the relative

accumulation of mitotic cell divisions of a particular sample and is able to capture

different tendencies in gaining or losing DNA methylation during mitotic cell division,

as it respectively happens in ALL and MM patients (Fig. 5b). Finally, epiCMIT cannot be

affected by different distribution of cell cycle phases in tumor samples, since the DNA

methylome remains rather stable during the whole cell cycle [44].

The applicability of the epiCMIT as mitotic clock was validated through several

perspectives. First, we compared it with two previously reported hypermethylation-

based mitotic clocks called epiTOC and MiAge [38,45] (Supplementary Table 8). The

epiCMIT showed excellent correlations with these clocks in B-cell neoplasms that tend

to acquire polycomb-related hypermethylation (e.g. mostly ALL, but also DLBCL and

MCL); a moderate correlation in the case of CLL, which acquires more hypo- than

hypermethylation (Fig. 4a and Extended Data Fig. 4b), and a total lack of correlation in

the case of MM, which mostly loses DNA methylation (Fig. 5d and Extended Data Fig.

5a). Identical observations were obtained comparing the epiCMIT and the widely-

reported CpG island methylator phenotype (CIMP) in human cancer [46] calculated as

previously proposed [47] (Supplementary Table 8). Therefore, our analysis also reveals

that the classical CIMP score in reality may represent a measure of the mitotic history

of the cells. A potentially confusing aspect related to epiCMIT is the fact that DNA

methylation has mainly been used to measure the chronological age of an individual

[48,49]. In order to study the relationship between mitotic and aging DNA methylation

clocks, we first compared the epiCMIT in naïve B cells from infant, adults and elderly

donors and showed that epiCMIT was stable in all age ranges regardless of donor's age

(Extended Data Fig. 5b). We next applied the most popular chronological age clock

proposed by Horvath [50] and showed that the age of donors was predicted with high

accuracy. This result indicates that, although both Horvath and epiCMIT scores are

based on DNA methylation, they measure distinct biological phenomena (Extended

Data Fig. 5b). Furthermore, the epiCMIT was highly variable in pediatric ALL samples

with a minimum age range (Extended Data Fig. 5c). Collectively, these analyses

indicate that the epiCMIT is a more universal mitotic clock than hypermethylation-

based clocks since it captures the proliferative history of neoplasms regardless of their

tendency of gaining or losing DNA methylation during mitotic cell division.

Second, the epiCMIT was validated using the *in vitro* B-cell differentiation

model shown in Fig. 4j, in which we observed that epiCMIT increases with the

proliferative history of the cells without altering gene expression levels of the epiCMIT-

associated genes (Fig. 5e). Third, using WGS data from 138 CLL patients from our

cohort [10,17,51,52], we observed that the epiCMIT was highly correlated with the total

number of somatic mutations (Extended Data Fig. 5d). We next extracted mutational

signatures as recently described [53] (Extended Data Fig. 5e) and observed significant

correlations with mutational signatures SBS1 and SBS5 (Fig. 5f), that have been

recently described as mitotic-like mutational processes [54]. We also identified a

significant link between the epiCMIT and the non-canonical AID signature (SBS9) (Fig.

5f) [52,55] in IGHV mutated CLL, possibly reflecting rounds of cell divisions in germinal

center B cells before differentiation into memory B cells and malignant transformation.

Fourth, although the epiCMIT reflects the proliferative history of the cell rather than

the actual proliferative status of the samples (e.g, bmPC have high epiCMIT and do not

proliferate, Fig. 5c), a relationship between epiCMIT and cell proliferation is expected

(more proliferative history implies overall more proliferation, although also depends

on time). Accordingly, MCL cases showing higher Ki-67 (a proliferation marker) also

had higher epiCMIT than cases with moderate Ki-67 expression (Extended Data Fig 5f).

Furthermore, GSEA analyses in ALL and CLL cases with high and low epiCMIT revealed

that cases with high epiCMIT showed higher expression of genes related with cell

proliferation (Extended Data Fig. 5g, h). Thus, these data suggest that cases with higher

proliferative history also seem to have higher proliferation at the time of sampling.

Collectively, the four lines of evidence presented above support that the epiCMIT may

represent a *bona fide* measure of the relative number of cumulative mitotic cell

divisions that normal and neoplastic B cells have undergone since the uncommitted

hematopoietic cell stage.

The epiCMIT is a strong independent variable predicting clinical behavior in B-cell

tumors

In normal B-cell maturation, the epiCMIT gradually augments as B cells proliferate

during cell differentiation, an increase that is particularly marked in proliferative GC B

cells (Fig. 5c). In neoplastic B cells, however, the interpretation of the epiCMIT is less

trivial and must be divided into two components: the epiCMIT of the cell of origin and

the epiCMIT acquired in the course of the neoplastic transformation and progression

(Extended Data Fig. 6a). Therefore, the relative epiCMIT must be compared among

patients from entities arising from the same B-cell maturation stage and should be a

dynamic variable during cancer progression. Thus, we compared the epiCMIT in two

paradigmatic transitions between precursor conditions and overt cancer, i.e.

monoclonal gammopathy of undetermined significance (MGUS) and MM, as well as

monoclonal B cell lymphocytosis (MBL) and CLL categorized according to their cellular

origin. This analysis showed an overall lower epiCMIT in precursor lesions compared

with overt cancer (Fig. 6a, upper panels), as would be expected due to their increased

leukemia-specific proliferative history. In line with this finding, the epiCMIT increased

in paired CLLs at diagnosis and progression before treatment as well as in sequential

ALL samples at diagnosis, first relapse and second relapse (Fig. 6a, lower panels). These

results suggest that the epiCMIT evolves together with clinical progression.

Based on our previous observations, we next wondered whether the epiCMIT

could be useful to predict the clinical behavior of B-cell neoplasms. We analyzed the

relationship of the epiCMIT with clinical outcome in specific B-cell tumor subtypes

based on their shared cytogenetics (i.e. ALL) or cell of origin (i.e. MCL, CLL and DLBCL),

and thus having a similar ground state B-cell specific proliferative history (Extended

Data Fig. 6a). In ALL, high epiCMIT was consistently associated with longer relapse-free

and overall survival (Fig. 6b and Extended Data Fig. 6b) of the patients within each

cytogenetic subtype, and thus better clinical outcome. Next, we additionally showed

epiCMIT as significant variable in a multivariate regression Cox model with epiCMIT as

continuous variable together with cytogenetic subtypes (Fig. 6b). This observation is in

line with published evidence showing that a higher CIMP, which is highly correlated

with the epiCMIT in ALL (Fig. 5d), is a good prognosis factor in this disease [56,57]. In

contrast to ALL, the opposite clinical scenario was observed in mature B-cell

neoplasms. In each of the CLL subtypes, a high epiCMIT was strongly associated with a

worse prognosis using time to treatment (TTT) as end-point variable, both from

sampling time (Fig. 6c, left panel) and in cases whose sample was obtained close to

diagnosis (Extended Data Fig. 6c, left panel). Additionally, a multivariate Cox regression

model for TTT revealed that epiCMIT as continuous variable was a highly significant

variable conferring dismal prognosis together with age, number of driver genetic

alterations [17,58] and epigenetic subgroups [10,11,59] (Fig. 6c right panel and Extened Data

Fig. 6c). The epiCMIT maintained its independent prognostic value using overall

survival as end-point variable, although its effect was moderate (Extended Data Fig.

6d). These findings were widely confirmed in an additional series of 136 CLLs treated

with chemo-immunotherapy (Fig. 6d and Extended Data Fig. 6c, d right panels). In the

case of MCL, the epiCMIT showed an independent poor prognostic impact in the two

cell-of-origin subtypes (C1 and C2), an observation that was confirmed in an extended

series in the more aggressive and prevalent C1 group (Fig. 6e, f). Lastly, although the

sample size was limited and requires further studies, our data suggest that high

epiCMIT could also represent a poor prognostic variable within the two cell-of-origin

DLBCL subtypes (Extended Data Fig. 6e). Overall, these results suggest that an

increased proliferative history of the neoplastic clone in precursor B-cell neoplasms at

diagnosis predicts for a better disease-free survival, as previously published in studies

analyzing the CIMP score [56,57]. In sharp contrast, in mature B-cell neoplasms, which are

overall less proliferative than ALL, neoplastic clones with high proliferation history

seem to predict for future proliferative capacity and consistently show worse clinical

outcomes.

epiCMIT is associated with specific genetic driver alterations in CLL

We next sought to assess which CLL driver alterations may confer a proliferative

advantage to neoplastic cells, and subsequently a higher epiCMIT. To that end, we

exploited 477 CLL samples in which we had DNA methylation data and whole exome

sequencing (WES) (Fig. 7a). We initially depicted all driver genetic changes in each CLL

subtype divided in high and low epiCMIT (Extended Data Fig. 7a). Next, we

interrogated the levels of epiCMIT in patients with each driver genetic alteration (with

at least 4 mutated patients). We performed the analyses in all CLL patients (showed as

Global in Fig. 7b) and then within each epigenetic subgroup separately (Fig. 7b,

Extended Data Fig. 7b and Methods). We showed significant and positive associations

of epiCMIT with 24 genetic driver alterations affecting the main signaling pathways

altered in CLL (Fig. 7b,c) [17,58]. The majority of these genetic alterations have been

previously linked to an adverse clinical behavior of patients, such as *NOTCH1, TP53,*

SF3B1, ATM, BIRC3 or EGR2. Interestingly, epiCMIT showed a clear association with a

new and recently identified non-coding genetic driver in CLL, the U1 spliceosomal RNA,

a finding that may help in explaining its suggested poor prognostic impact [60].

Remarkably, the presence of some genetic alterations was associated with high

epiCMIT indistinctly in all patients, such as *TP53*, while others were particularly

associated with epiCMIT within CLL subgroups, such as *NOTCH1* in n-CLL and i-CLL,

SF3B1 in i-CLL, and del(13q) and tri12 in m-CLL.

Collectively, these results suggest that the well-established clinical impact of genetic

alterations in CLL may be explained by their association with a high proliferative

potential, being this association different for certain genetic alterations depending on

the maturation state of the cellular origin.

Discussion

In this study, we have followed a systematic approach to dissect the sources of DNA

methylation variability of B-cell neoplasms in the context of the normal B-cell

differentiation program. Overall, we found that the methylation levels of 88% of all

CpGs are modulated in normal and/or neoplastic B cells, suggesting that the human

DNA methylome is even more dynamic than previously appreciated [6,61,62]. The

extensive DNA methylation variability among different B-cell neoplasms is in part

related to imprints of normal B-cell development. This phenomenon has been

previously observed and has led to a more accurate classification not only of B-cell

neoplasms [9–11,59], but also of solid tumors [2,63,64]. In addition to this epigenetic link to

normal cell maturation, each B-cell neoplasm also shows disease-specific hyper- and

hypomethylation. Of particular interest are the disease-specific *de novo*

hypomethylation signatures in active regulatory regions, which were associated with

binding of TFs involved in the pathogenesis of each respective B-cell tumor. This

phenomenon was particularly marked in ALL, MCL and CLL, whose *de novo*

hypomethylation was enriched in regulatory elements [23–25]. Unexpectedly, although

DLBCL and MM pathogenesis has been linked to TFs, *de novo* hypomethylation was

depleted in regulatory elements containing TF binding sites. In these two malignancies,

we detected few binding sites of TFs with potential involvement in the diseases.

However, we did not detect classical TFs such as BCL6 in DLBCL or IRF4 in MM, possibly

because they are key players during B-cell differentiation and their binding sites may

already be hypomethylated in the normal B-cell counterparts of DLBCL and MM.

In spite of the importance of DNA methylation in regulatory regions, we

identified that the majority of DNA methylation changes in B-cell neoplasms, and

especially those related to inter-patient variability within each B-cell tumor subtype,

are located in inactive chromatin. The magnitude of this inter-patient variability

affecting normal B-cell related and -independent regions is highly correlated,

suggesting that the cause of these DNA methylation changes is a phenomenon that

takes place both in normal and neoplastic B cells. Compelling published evidence as

well as new data presented in our study support the notion that mitotic cell division

leaves transcriptionally-inert epigenetic imprints onto the DNA. These take place in the

form of hypomethylation of heterochromatin and hypermethylation of polycomb-

repressed regions. This knowledge has recently led to the concept of using DNA

methylation as a mitotic clock [38,45], and has been recently used by single cell DNA

methylation data to track the evolutionary trajectory of CLL [65]. Notably, methylation-

based mitotic clocks seem to capture different biological information from aging clocks

(Extended Data Fig. 5b) [48]. As far as we are aware, there are two published mitotic

clocks [38,45] and both are hypermethylation-based. Furthermore, our analyses suggest

that the classical CIMP phenomenon in cancer [46] may indeed represent another

hypermethylation-based mitotic clock. However, only using hypermethylation to

determine the mitotic clock is insufficient to capture the mitotic activity of the cells, as

some neoplasms may not acquire hypermethylation upon cell division but rather

hypomethylation. For instance, neoplastic plasma cells in MM do not tend to acquire

polycomb-related hypermethylation. Thus, using hypermethylation to determine the

mitotic history of MM cells would incongruently lead to the conclusion that they have

not proliferated beyond their cellular origin (Fig. 5d and Extended Data Fig. 5a).

Therefore, to circumvent possible misleading interpretations, we generated a more

broadly applicable mitotic clock (called epiCMIT) that takes into consideration both

hyper- and hypomethylation. Importantly, epiCMIT aims at capturing the entire mitotic

history of cells, including cell division associated both with normal development as well

as neoplastic transformation and progression (Extended Data Fig. 6a). Thus, the

epiCMIT must not be compared among tumors arising from different normal

counterparts but its relative magnitude must be studied in those arising from a

particular maturation stage. Within each of these subgroups, the relative epiCMIT has

a profound independent prognostic value from other well-established clinical

variables. Increased epiCMIT is associated with worse clinical outcome within CLL and

MCL subgroups, thus indicating that superior proliferative history before treatment

seems to determine future proliferative capacity and is thus associated with worse

clinical outcome. Strikingly, we consistently found the opposite pattern in precursor

ALL subgroups, a finding in line with recent reports showing that the presence of CIMP

(a hypermethylation-based mitotic clock) is associated with better clinical outcome

[56,57]. These results suggest that children having more proliferative ALL cells at diagnosis

(and thus a larger proliferative history) are more prone to be cured with high intensive

chemotherapy regimens, [66], which cannot be administrated in elderly patients.

Finally, in order to identify the genetic lesions associated with higher

proliferative history in CLL, we exploited our thorough genetic characterization of our

cases to study the relationship between driver genetic alterations and epiCMIT. We

identified 24 driver genetic alterations that may confer a higher proliferative capacity

and thus are associated with higher epiCMIT or methylation evolution [67]. These genetic

alterations were distributed throughout the main altered signaling pathways in CLL

and were manifested differently in distinct CLL subgroups based on their cellular origin

(Fig. 7b, c). This finding suggests that specific mutations may predispose to a higher

proliferative advantage depending on the maturation stage and (epi)genetic makeup

of the cellular origin.

In summary, our comprehensive epigenetic evaluation of normal and neoplastic

B cells at different maturation stages uncovers multiple new insights into the biological

roles of DNA methylation in cancer, an analytic approach that may also benefit our

understanding of other cancers. From a clinical perspective, DNA methylation may

provide a holistic diagnostic and prognostic approach to B-cell neoplasms. Particularly,

we defined an accurate and easy-to-implement pan-B-cell tumor diagnostic tool and

generated a mitotic clock reflecting the proliferative history of the neoplastic cells of

each patient to estimate their clinical risk, which shall represent a valuable asset in the

precision medicine era.

ACKNOWLEDGEMENTS

This research was funded by the European Union's Seventh Framework Programme through the Blueprint Consortium (grant agreement 282510), Generalitat de Catalunya Suport Grups de Recerca AGAUR 2017-SGR-1142 (to E.C.) and 2017-SGR-736 (to J.I.M.-S.), Ministerio de Ciencia, Innovación y Universidades of the Spanish Government (MCIU), Grants RTI2018-094274-B-I00 (to E.C.) and SAF2017-86126-R (to J.I.M.-S.) as well as Proyecto Medicina Personalizada PERMED (Grant PMP15/00007), which is part of Plan Nacional de I+D+I and is co-financed by the ISCIII-Sub-Directorate General for Evaluation and the European Regional Development Fund (FEDER-"Una manera de Hacer Europa"), CIBERONC (CB16/12/00225, CB16/12/00334, CB16/12/00236, and CB16/12/00489), the Accelerator award CRUK/AIRC/AECC joint funder-partnership, research funding from Fondo de Investigaciones Sanitarias, Instituto de Salud Carlos III PI17/01061 (SB), Ministerio de Ciencia, Innovación y Universidades (MCIU), RTI2018-094274-B-I00, SAF2015-64885-R (EC), the NIH grant number 1 P01CA229100 (EC), and the European Regional Development Fund "Una manera de fer Europa", CERCA Programme/Generalitat de Catalunya. FN is supported by a pre-doctoral fellowship of the Ministerio de Economía y Competitividad (MINECO, BES-2016-076372). E.C. is an Academia Researcher of the "Institució Catalana de Recerca i Estudis Avançats" (ICREA) of the Generalitat de Catalunya. This work was partially developed at the Centro Esther Koplowitz (CEK, Barcelona, Spain). We thank Francesco Maura for his help with the analysis of mutational signatures.

AUTHOR CONTRIBUTIONS

Investigator contributions were as follows: T.B., J.N., Y.N-Z., G.L., A.R-D., S.M., R.O.,

G.C., M.K., A.C-Q., L.S-T., J.W., J.L., E.G., S.B., P.J., X.A., F.P., C.L-O., X.S.P., C.C.O., T.Z.,

J.D., A.L-G. and E.C. contributed to sample biological and/or clinical annotation; M.D-

F., G.C., F.N. and R.B. performed methylome, ChIP-Seq, transcriptome, genetic and/or

statistical analyses; R.R., M.P. and D.T. provided computational support. M.D-F. and

J.I.M.-S. participated in the study design. M.D-F., F.N., R.B., T.B., J.D., A.L-G., E.C. and

J.I.M.-S. participated in data interpretation. J.I.M.-S. directed the research and wrote

the manuscript together with M.D-F.

COMPETING INTERESTS

The authors declare no competing interests.

References

1.　Roy, N. & Hebrok, M. Regulation of Cellular Identity in Cancer. *Dev. Cell* **35**, 674–
84 (2015).

2.　Hoadley, K. A. *et al.* Cell-of-Origin Patterns Dominate the Molecular
Classification of 10,000 Tumors from 33 Types of Cancer. *Cell* **173**, 291-304.e6
(2018).

3.　Swerdlow SH, Campo E, Harris NL, Jaffe ES, Pileri SA, Stein H, T. J. *WHO*
Classification of Tumours of Haematopoietic and Lymphoid Tissues.
(International Agency for Research on Cancer (IARC), 2017).

4.　Bird, A. DNA methylation patterns and epigenetic memory. *Genes Dev.* **16**, 6–21
(2002).

5.　Luo, C., Hajkova, P. & Ecker, J. R. Dynamic DNA methylation: In the right place at
the right time. *Science* **361**, 1336–1340 (2018).

6.　Kulis, M. *et al.* Whole-genome fingerprint of the DNA methylome during human
B cell differentiation. *Nat. Genet.* **47**, 746–756 (2015).

7.　Nordlund, J. *et al. Genome-wide signatures of differential DNA methylation in*
pediatric acute lymphoblastic leukemia. Genome biology **14**, (2013).

8.　Lee, S.-T. *et al.* Epigenetic remodeling in B-cell acute lymphoblastic leukemia
occurs in two tracks and employs embryonic stem cell-like signatures. *Nucleic*
Acids Res. **43**, 2590–602 (2015).

9.　Queirós, A. C. *et al.* Decoding the DNA Methylome of Mantle Cell Lymphoma in
the Light of the Entire B Cell Lineage. *Cancer Cell* **30**, 806–821 (2016).

10.　Kulis, M. *et al.* Epigenomic analysis detects widespread gene-body DNA
hypomethylation in chronic lymphocytic leukemia. *Nat. Genet.* **44**, 1236–1242
(2012).

11.　Oakes, C. C. *et al.* DNA methylation dynamics during B cell maturation underlie a
continuum of disease phenotypes in chronic lymphocytic leukemia. *Nat. Genet.*
(2016). doi:10.1038/ng.3488

12.　Kretzmer, H. *et al.* DNA methylome analysis in Burkitt and follicular lymphomas
identifies differentially methylated regions linked to somatic mutation and
transcriptional control. *Nat. Genet.* (2015). doi:10.1038/ng.3413

13.　Shaknovich, R. *et al.* DNA methylation signatures define molecular subtypes of
diffuse large B-cell lymphoma. *Blood* **116**, e81-9 (2010).

14.　Agirre, X. *et al.* Whole-epigenome analysis in multiple myeloma reveals DNA
hypermethylation of B cell-specific enhancers. *Genome Res.* **25**, 478–87 (2015).

15.　Kaiser, M. F. *et al.* Global methylation analysis identifies prognostically
important epigenetically inactivated tumor suppressor genes in multiple
myeloma. *Blood* **122**, 219–226 (2013).

16.　Oakes, C. C. & Martin-Subero, J. I. Insight into origins, mechanisms & utility

of DNA methylation in B cell malignances. *Blood* **132**, blood-2018-02-692970
(2018).

17.   Puente, X. S. *et al.* Non-coding recurrent mutations in chronic lymphocytic
leukaemia. *Nature* (2015). doi:10.1038/nature14666

18.   Karube, K. *et al.* Integrating genomic alterations in diffuse large B-cell lymphoma
identifies new relevant pathways and potential therapeutic targets. *Leukemia*
675–684 (2017). doi:10.1038/leu.2017.251

19.   Stunnenberg, H. G., International Human Epigenome Consortium & Hirst, M.
The International Human Epigenome Consortium: A Blueprint for Scientific
Collaboration and Discovery. *Cell* **167**, 1145–1149 (2016).

20.   Viré, E. *et al.* The Polycomb group protein EZH2 directly controls DNA
methylation. *Nature* **439**, 871–874 (2006).

21.   Stadler, M. B. *et al.* Corrigendum: DNA-binding factors shape the mouse
methylome at distal regulatory regions. *Nature* **484**, 550–550 (2012).

22.   Burger, L., Gaidatzis, D., Schübeler, D. & Stadler, M. B. Identification of active
regulatory regions from DNA methylation data. *Nucleic Acids Res.* **41**, (2013).

23.   Somasundaram, R., Prasad, M. A. J., Ungerbäck, J. & Sigvardsson, M.
Transcription factor networks in B-cell differentiation link development to acute
lymphoid leukemia. *Blood* **126**, 144–152 (2015).

24.   Sánchez-Tilló, E. *et al.* The EMT activator ZEB1 promotes tumor growth and
determines differential response to chemotherapy in mantle cell lymphoma. *Cell*
Death Differ. **21**, 247–257 (2014).

25.   Wolf, C. *et al.* NFATC1 activation by DNA hypomethylation in chronic
lymphocytic leukemia correlates with clinical staging and can be inhibited by
ibrutinib. *Int. J. Cancer* **142**, 322–333 (2018).

26.   Blonska, M. *et al.* Jun-regulated genes promote interaction of diffuse large B-cell
lymphoma with the microenvironment. *Blood* **125**, 981–991 (2015).

27.   Huerta-Yepez, S. *et al.* Overexpression of Yin Yang 1 in bone marrow-derived
human multiple myeloma and its clinical significance. *Int. J. Oncol.* **45**, 1184–
1192 (2014).

28.   Sprynski, A. C. *et al.* Insulin is a potent myeloma cell growth factor through
insulin/IGF-1 hybrid receptor activation. *Leukemia* **24**, 1940–1950 (2010).

29.   Riz, I. & Hawley, R. G. Increased expression of the tight junction protein
TJP1/ZO-1 is associated with upregulation of TAZ-TEAD activity and an adult
tissue stem cell signature in carfilzomib-resistant multiple myeloma cells and
high-risk multiple myeloma patients. *Oncoscience* **4**, 79–94 (2017).

30.   Herath, N. I., Rocques, N., Garancher, A., Eychène, A. & Pouponnot, C. GSK3-
mediated MAF phosphorylation in multiple myeloma as a potential therapeutic
target. *Blood Cancer J.* **4**, e175–e175 (2014).

31.   Navarro, A. *et al.* Improved classification of leukemic B-cell lymphoproliferative
disorders using a transcriptional and genetic classifier. *Haematologica* **102**, 360–

363 (2017).

32.   Alizadeh, A. A. *et al.* Distinct types of diffuse large B-cell lymphoma identified by
690   gene expression profiling. *Nature* **403**, 503–511 (2000).

691   33.   Aran, D., Toperoff, G., Rosenberg, M. & Hellman, A. Replication timing-related
692   and gene body-specific methylation of active human genes. *Hum. Mol. Genet.*
693   **20**, 670–680 (2011).

694   34.   Beerman, I. *et al.* Proliferation-dependent alterations of the DNA methylation
695   landscape underlie hematopoietic stem cell aging. *Cell Stem Cell* **12**, 413–25
696   (2013).

697   35.   Landan, G. *et al.* Epigenetic polymorphism and the stochastic formation of
698   differentially methylated regions in normal and cancerous tissues. *Nat. Genet.*
699   **44**, 1207–14 (2012).

700   36.   Siegmund, K. D., Marjoram, P., Woo, Y.-J., Tavaré, S. & Shibata, D. Inferring
701   clonal expansion and cancer stem cell dynamics from DNA methylation patterns
702   in colorectal cancers. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 4828–4833 (2009).

703   37.   Spencer, D. H. *et al.* CpG Island Hypermethylation Mediated by DNMT3A Is a
704   Consequence of AML Progression. *Cell* **168**, 801-816.e13 (2017).

705   38.   Yang, Z. *et al.* Correlation of an epigenetic mitotic clock with cancer risk.
706   *Genome Biol.* **17**, 205 (2016).

707   39.   Zhou, W. *et al.* DNA methylation loss in late-replicating domains is linked to
708   mitotic cell division. *Nat. Genet.* **50**, 591–602 (2018).

709   40.   Schlesinger, Y. *et al.* Polycomb-mediated methylation on Lys27 of histone H3
710   pre-marks genes for de novo methylation in cancer. *Nat. Genet.* **39**, 232–236
711   (2007).

712   41.   Ohm, J. E. *et al.* A stem cell–like chromatin pattern may predispose tumor
713   suppressor genes to DNA hypermethylation and heritable silencing. *Nat. Genet.*
714   **39**, 237–242 (2007).

715   42.   Widschwendter, M. *et al.* Epigenetic stem cell signature in cancer. *Nat. Genet.*
716   **39**, 157–158 (2007).

717   43.   Caron, G. *et al.* Cell-Cycle-Dependent Reconfiguration of the DNA Methylome
718   during Terminal Differentiation of Human B Cells into Plasma Cells. *Cell Rep.* **13**,
719   1059–71 (2015).

720   44.   Vandiver, A. R., Idrizi, A., Rizzardi, L., Feinberg, A. P. & Hansen, K. D. DNA
721   methylation is stable during replication and cell cycle arrest. *Sci. Rep.* **5**, 1–8
722   (2015).

723   45.   Youn, A. & Wang, S. The MiAge Calculator: a DNA methylation-based mitotic age
724   calculator of human tissue types. *Epigenetics* **13**, 192–206 (2018).

725   46.   Issa, J. CpG island methylator phenotype in cancer. *Nat. Rev. Cancer* **4**, 988–93
726   (2004).

727   47.   Sánchez-Vega, F., Gotea, V., Margolin, G. & Elnitski, L. Pan-cancer stratification
728   of solid human epithelial tumors and cancer cell lines reveals commonalities and

tissue-specific features of the CpG island methylator phenotype. *Epigenetics and Chromatin* **8**, 1–24 (2015).

48. Field, A. E. *et al.* DNA Methylation Clocks in Aging: Categories, Causes, and Consequences. *Mol. Cell* **71**, 882–895 (2018).

49. Horvath, S. & Raj, K. DNA methylation-based biomarkers and the epigenetic clock theory of ageing. *Nat. Rev. Genet.* (2018). doi:10.1038/s41576-018-0004-3

50. Horvath, S. DNA methylation age of human tissues and cell types. *Genome Biol* **14**, R115 (2013).

51. Ferreira, P. G. *et al.* Transcriptome characterization by RNA sequencing identifies a major molecular and clinical subdivision in chronic lymphocytic leukemia Gene Regulation Stem Cells and Cancer Programme, Centre for Genomic Regulation (CRG). 212–226 (2014). doi:10.1101/gr.152132.112

52. Puente, X. S. *et al.* Whole-genome sequencing identifies recurrent mutations in chronic lymphocytic leukaemia. *Nature* **475**, 101–105 (2011).

53. Maura, F. et al. A practical guide for mutational signature analysis in hematological malignancies. *Nat. Commun.* (2019). doi:10.1038/s41467-019-11037-8

54. Alexandrov, L. B. *et al.* Clock-like mutational processes in human somatic cells. *Nat. Genet.* **47**, 1402–7 (2015).

55. Alexandrov, L. *et al.* The Repertoire of Mutational Signatures in Human Cancer. *bioRxiv* (2018). doi:https://doi.org/10.1016/j.cell.2011.02.013

56. Borssén, M. *et al.* DNA methylation holds prognostic information in relapsed precursor B-cell acute lymphoblastic leukemia. *Clin. Epigenetics* **10**, 31 (2018).

57. Sandoval, J. *et al.* Genome-wide DNA methylation profiling predicts relapse in childhood B-cell acute lymphoblastic leukaemia. *Br. J. Haematol.* **160**, 406–9 (2013).

58. Landau, D. A. *et al.* Mutations driving CLL and their evolution in progression and relapse. *Nature* **526**, 525–30 (2015).

59. Queirós, a C. *et al.* A B-cell epigenetic signature defines three biological subgroups of chronic lymphocytic leukemia with clinical impact. *Leukemia* 598–605 (2015). doi:10.1038/leu.2014.252

60. Shuai, S. *et al.* The U1 spliceosomal RNA is recurrently mutated in multiple cancers. *Nature* **574**, 712–716 (2019).

61. Ziller, M. J. *et al.* Charting a dynamic DNA methylation landscape of the human genome. *Nature* **500**, 477–81 (2013).

62. Lister, R. *et al.* Human DNA methylomes at base resolution show widespread epigenomic differences. *Nature* **462**, 315–322 (2009).

63. Rodríguez-Paredes, M. *et al.* Methylation profiling identifies two subclasses of squamous cell carcinoma related to distinct cells of origin. *Nat. Commun.* **9**, (2018).

769   64.   Bormann, F. *et al.* Cell-of-Origin DNA Methylation Signatures Are Maintained
770   during Colorectal Carcinogenesis. *Cell Rep.* **23**, 3407–3418 (2018).

771   65.   Gaiti, F. *et al.* Epigenetic evolution and lineage histories of chronic lymphocytic
772   leukaemia. *Nature* (2019). doi:10.1038/s41586-019-1198-z

773   66.   Rhein, P. *et al.* Gene expression shift towards normal B cells, decreased
774   proliferative capacity and distinct surface receptors characterize leukemic blasts
775   persisting during induction therapy in childhood acute lymphoblastic leukemia.
776   *Leukemia* **21**, 897–905 (2007).

777   67.   Oakes, C. C. *et al.* Evolution of DNA Methylation Is Linked to Genetic Aberrations
778   in Chronic Lymphocytic Leukemia. *Cancer Discov.* **4**, 348–361 (2014).

Figure legends

Fig. 1: Shared DNA methylation dynamics in B-cell tumors.

a, Number of normal and neoplastic B-cells included in the study with tumor cell

content of at least 60% (additional samples for validations are used in Fig. 3 and Fig. 6,

which are later detailed). HPC, hematopoietic precursor cells; pre-B, precursor B-cell

and immature B cells; NBC, naïve B cells; GC, germinal center B cells; MBC, memory B

cells; tPC, tonsillar plasma cells; bmPC, bone-marrow plasma cells; ALL, acute

lymphoblastic leukemia; MCL, mantle cell lymphoma; CLL, chronic lymphocytic

leukemia; DLBCL, Diffuse large B cell lymphoma; MM, multiple myeloma; BM, bone

marrow; PB, peripheral blood; LN, lymph node.

b, Different levels of DNA methylation variability addressed in the study.

c, Heatmaps showing shared DNA hyper- (top) and hypomethylation (bootm) in B-cell

tumors.

d, Chromatin state enrichments of regions sharing DNA hyper- and hypomethylation

(all CpGs of 450k array were used as background). ActProm, Active promoter;

WkProm, Weak promoter; StrEnh1, Strong enhancer 1 (promoter-related); StrEnh2,

Strong enhancer 2; WkEnh, Weak enhancer; TxnTrans, Transcription transition;

TxnElong, Transcription elongation; WkTxn, Weak transcription; PoisProm, Poised

promoter; H3K27me3, Polycomb-repressed region; H3K9me3, H3K9me3

heterochromatin; Het;LowSign, Het;LowSign heterochtomatin.

e, Overlap of the target genes of stably methylated and unmethylated CpGs.

f, Gene expression percentile within each sample of genes showing stable hyper- and

hypomethylation.

Fig. 2: Disease-specific hypomethylation at H3K27ac regions is associated with

specific transcription factor bindings and differential gene expression.

a, Principal component analysis of normal and neoplastic B-cells, including ALL, MCL,

CLL, DLBCL and MM. ALL, acute lymphoblastic leukemia; MCL, mantle cell lymphoma;

CLL, chronic lymphocytic leukemia; DLBCL, Diffuse large B cell lymphoma; MM,

multiple myeloma.

b, Number of *de novo* DNA methylation changes in each B-cell tumor entity.

c, Heatmap showing entity-specific hypomethylation and the number of CpGs falling at

regulatory regions. Per each B-cell tumor, the same number of *de novo* CpGs was

randomly chosen from the 450K array and interrogated the percentage falling at

regulatory regions.

d, Transcription factor binding site predictions for *de novo* hypomethylated CpGs

falling at regulatory regions in c, for each B-cell tumor.

e, Differential gene expression percentiles among B-cell tumors for genes showing

specific-hypomethylation at regulatory regions.

Fig. 3: Accurate classification of 14 clinico-biological subtypes of B cell neoplasms

using epigenetic biomarkers.

a, Heatmaps for the CpGs used for the pan B-cell cancer classifier. The classifier

consists of two steps: in the first step (1) an unknown B-cell tumor can be predicted

into ALL, MCL, CLL, DLBCL or MM, and subsequently (second step, using one of the

predictors (2), (3), (4) or (5)) to any B-cell tumor subtype, namely HeH, 11q23/MLL,

t(1;19), t(9;22), dic(9;20) for ALL, C1 and C2 for MCL, n-CLL, i-CLL and m-CLL for CLL,

and finally GCB or ABC for DLBCLs. ALL, acute lymphoblastic leukemia; MCL, mantle cell

lymphoma; CLL, chronic lymphocytic leukemia; DLBCL, Diffuse large B cell lymphoma;

MM, multiple myeloma.

b, Accuracy for the pan-B-cell cancer diagnostic classifier (formed by the 5 predictors in

Fig. 3a) for training and validation series. Sensitivity is represented as black circles or

triangles for training or validation series, respectively (sensitivity in the training series

was evaluated using 10-fold stratified cross-validation). The total number of samples

used for both training and validation is shown.

Fig. 4: Patient-specific DNA methylation changes are associated with silent chromatin

without an impact on gene expression.

a, Circos plot representing the number of DNA methylation changes with respect to

heamatopoietic precursor cells (HPC) in individual patients for normal and neoplastic B

cells (each bar represents one patient). Total number of DNA methylation changes,

hypomethylation changes and hypermethylation changes are depicted at outer,

middle and inner tracks, respectively. Changes are further classified and color-coded as

B-cell related or B-cell independent if they occur or not during normal B-cell

differentiation, respectively.

b, Number of B-cell related changes against B-cell independent changes in normal and

neoplastic B-cells. Fitted regression lines are shown at bottom per each B-cell tumor

subtype.

c, B-cell related CpGs losing DNA methylation in B-cell tumors and the percentages in

each chromatin state in normal and neoplastic B-cells.

d, Example CpGs from **c** in normal and neoplastic B-cells.

e, Density of genes distributed along gene expression percentiles of genes associated

with B-cell related CpGs losing DNA methylation in B-cell tumors in each B-cell tumor

subtype. Expressed genes are displayed at right as control (with presence of

H3K36me3). Means within each B-cell subpopulation as well as B-cell tumors are

represented.

f, B-cell related CpGs gaining DNA methylation in B-cell tumors and the percentages in

each chromatin state in normal and neoplastic B-cells.

g, Example CpGs from **f** in normal and neoplastic B-cells.

h, Density of genes distributed along gene expression percentiles of genes associated

with B-cell related CpGs gaining DNA methylation in B-cell tumors in each B-cell tumor

subtype. Expressed genes are displayed at right as control (with presence of

H3K36me3). Means within each B-cell subpopulation as well as B-cell tumors are

represented.

i, Model for DNA methylation changes occurring at repressed regions during mitotic

cell division.

j, *In vitro* model for plasma blast differentiation from human primary naïve B cells.

Primary naïve B cells are labeled with CFSE at day 0. DNA methylation profiles are

obtained at day 0 (primary naïve B cells) day 4 (CFSE-high and CFSE-low) and day 6 (P1-

CFSE-low and CD38+, P2-CFSE-intermediate, CD38-, P3-CFSE-high CD38-).

k, DNA methylation changes accumulate at repressed regions during mitotic cell

division upon *in vitro* proliferation and differentiation of primary naïve B cells to

plasma blasts. Hypermethylation takes place at H3K27me3 regions, whereas

hypomethylation at heterochromatin and H3K9me3 regions. ChIP-seq data for primary

NBC and GC B cells was used. The mean of several biological replicates is represented

(numbers are depicted at bottom of the heatmaps).

l, Density of genes distributed in gene expression percentiles for genes with changes in

DNA methylation in (K).

Fig. 5: Development and validation of an epigenetic mitotic clock reflecting the

proliferative history of normal and neoplastic B cells.

a, Selection of CpGs gaining DNA methylation at H3K27me3 and losing DNA

methylation in normal and neoplastic B cells to construct the epiCMIT-hyper and

epiCMIT-hypo scores. epiCMIT, epigenetically-determined Cumulative MIToses.

b, Correlation of epiCMIT-hyper and epiCMIT-hypo in normal and neoplastic B cells.

c, epiCMIT in normal and neoplastic B cells.

d, Correlation of epiCMIT with previously reported hypermethylation-based mitotic

clocks, including epiTOC (epigenetic Timer for Cancer risk) and MiAge. Correlation with

CIMP (CpG hypermethylator phenotype) is also shown, although it has not been

formally presented as mitotic clock.

e, epiCMIT score during the *in vitro* B-cell differentiation as well as gene expression of

genes containing epiCMIT-CpGs. Expressed genes are also shown as control (presence

of H3K36me3).

f, Correlation of epiCMIT and mutational signatures related to proliferative history for

138 CLL samples with available DNA methylation data and WGS. SBS1, SBS5 have been

reported as clock-like mutational signatures. SBS9 is related to ncAID mutations.

Fig. 6: The epiCMIT is a strong independent variable predicting clinical behavior in B-

cell tumors.

a, epiCMIT is a dynamic variable that reflects disease progression. Precursor conditions

MGUS and MBL show significantly lower epiCMIT levels than their respective cancer

conditions CLL and MM, respectively. epiCMIT increases from diagnosis to progression

in paired CLL samples and in paired ALL at diagnosis, first relapse and second relapse.

b, Kaplan-Meier curves for three ALL cytogenetic groups shown as example divided in

low and high epiCMIT using maxstat rank statistics. Multivariate cox regression model

for relapse-free survival with epiCMIT against cytogenetic groups. Hazard ratio for

epiCMIT correspond to 0.1 increments.

c, Kaplan-Meier curves for CLL epigenetic groups based on different cellular origin

divided in low and high epiCMIT using maxstat rank statistics. Multivariate cox

regression model for time to first treatment with epiCMIT against age, number of

driver alterations and epigenetic groups based on different cellular origin. Validation

series is shown in **d**. Hazard ratio for epiCMIT correspond to 0.1 increments.

e, Kaplan-Meier curves for MCL epigenetic groups based on different cellular origin

divided in low and high epiCMIT using maxstat rank statistics. Multivariate cox

regression model for overall survival with epiCMIT against epigenetic groups based on

different cellular origin. Hazard ratio for epiCMIT correspond to 0.1 increments.

f, Validation series for C1 MCL. Hazard ratio for epiCMIT correspond to 0.1 increments.

Fig. 7: epiCMIT is associated with specific genetic driver alterations in CLL

a, Are there specific genetic alterations in CLL that may predispose a proliferative

advantage to neoplastic cells?

b, Genetic driver alterations in CLL associated with higher epiCMIT grouped by

signaling pathways reported. Analyses were done globally for all CLL samples (although

adjusted by epigenetic groups) as well as within each epigenetic subgroup. Positive

point estimates relate to positive associations with epiCMIT. 95% confidence intervals

are shown with colors coding for FDR corrections. Number of patients with each

genetic driver alteration is shown at right.

c, Oncoprint showing genetic driver alterations associated with higher epiCMIT in CLL

epigenetic groups separately. Other clinicobiological features including MBL or CLL,

IGHV status, Age, Binet stage, epiCMIT subgroups based on Fig. 5c, need for treatment

and patient status are shown. Cases are ordered within each CLL subgroup from lower

to higher epiCMIT values. Distinct genetic driver alterations are depicted with different

colors and shapes. The percentage of mutated patients as well as barplots showing the

number of mutated patients for each alteration is shown at right.

Supplementary figure legends

Extended Data Fig.1

a, Principal component analysis and hierarchical clustering of paired un/purified DNA

methylation profiles obtained with EPIC array from MCL and CLL patients. Colors

represent the same un/purified sample, with FACS-based purities highlighted in each

sample. MCL, mantle cell lymphoma. CLL, chronic lymphocytic leukemia.

b, Correlations and Passing Bablock regression fits of gold-standard methods for tumor

purity prediction (FACS and genetic-based) against DNA methylation-based tumor

purity prediction for MCL and CLL patients.

c, Pearson correlations and Passing Bablock regression fits for gold-standard methods

for tumor purity predictions (FACS and genetic-based) against DNA methylation-based

tumor purity predictions for MM and DLBCL patients. MM, multiple myeloma. DLBCL,

Diffuse large B cell lymphoma.

d, Pan-B cell DNA methylation signature used to deconvolute DNA methylation and

obtain B-cell tumor purities in DLBCL and MM. Bar plots representing DNA-methylation

based predictions as well as gold standard-based predictions are represented at top

the heatmaps.

e, Genomic distribution of stably un/methylated CpGs in B-cell tumors.

f, Example gene showing stably un/methylated CpGs.

g, Gene ontology analysis of genes showing both stable un/methylated CpGs.

Extended Data Fig.2

a, Principal component analysis for normal and neoplastic B cells, including ALL, MCL,

CLL, DLBCL and MM. Each component is resented separately (until the ninth

component). ALL, acute lymphoblastic leukemia; MCL, mantle cell lymphoma; CLL,

chronic lymphocytic leukemia; DLBCL, Diffuse large B cell lymphoma; MM, multiple

myeloma.

b, Heatmap showing B-cell tumor-specific hypermethylation and the number of CpGs

falling at regulatory regions. Per each B-cell tumor, the same number of *de novo* CpGs

was randomly chosen from the 450K array and interrogated the percentage falling at

regulatory regions.

c, Genomic distribution for *de novo* DNA methylation changes in B-cell tumors.

d, Gene expression of DNA methyltransferases *DNMT1*, *DNMT3A* and *DNMT3B*

throughout B cell differentiation.

e, Gene expression percentile of TFs showing the most significant p-values and

frequencies in *de novo* hypomethylation signatures in each B-cell tumor in Fig. 2d.

Extended Data Fig. 3

a, Sensitivity of the pan-B-cell diagnostic algorithm for the classification of an unknown

B-cell tumor into ALL, MCL, CLL, DLBCL or MM while incrementing the number of CpGs

used in the classifier algorithm (first step of Fig. 3A). The number of CpGs selected for

the predictor is selected by maximizing the highest balanced accuracy and is indicated

with a red circle. This strategy was applied also in the remaining 4 predictors to classify

B-cell tumor subtypes in panels b,, c,, d, and e, (second step of Fig. 3A). Each B-cell

tumor is represented with different shapes and colors. ALL, acute lymphoblastic

leukemia; MCL, mantle cell lymphoma; CLL, chronic lymphocytic leukemia; DLBCL,

Diffuse large B cell lymphoma; MM, multiple myeloma

b, Sensitivity of predictor 2 for the pan-B-cell diagnostic algorithm (predictor 2 of Fig.

3a) for the classification of ALL into the subtypes HeH, 11q23/MLL, t(12;21), t(1;19),

t(9;22) and dic(9;20) while incrementing the number of CpGs.

c, Sensitivity of predictor 3 of the pan-B-cell diagnostic algorithm (predictor 3 of Fig.

3a) for the classification of MCL into the subtypes C1 or C2 while incrementing the

number of CpGs.

d, Sensitivity of predictor 4 of the pan-B-cell diagnostic algorithm (predictor 4 of Fig.

3a) for the classification of CLL into the subtypes n-CLL, i-CLL or m-CLL while

incrementing the number of CpGs.

e, Sensitivity of predictor 5 of the pan-B-cell diagnostic algorithm (predictor 5 of Fig.

3a) for the classification of DLBCL into the subtypes ABC and GCB while incrementing

the number of CpGs.

1001

Extended Data Fig.4

a, Variability of DNA methylation changes (IQR) in normal neoplastic B cells against the

median number of DNA methylation changes.

b, Correlations of B-cell independent DNA methylation changes in B-cell tumors

against B-cell related changes for hypermethylation (top) and hypomethylation

(bottom).

c, Number of B-cell related or B-cell independent hyper- or hypomethylation in B-cell

tumors (Methods).

d, B-cell independent CpGs losing DNA methylation in B-cell tumors and the

percentages in each chromatin state in normal and neoplastic B-cells.

e, Example CpGs from **d,** in normal and neoplastic B-cells.

f, Density of genes distributed along gene expression percentiles of genes associated

with B-cell independent CpGs losing DNA methylation in B-cell tumors in each B-cell

tumor subtype. Expressed genes are displayed at right as control (with presence of

H3K36me3). Means within each B-cell subpopulation as well as B-cell tumors are

represented.

g, B-cell independent CpGs gaining DNA methylation in B-cell tumors and the

percentages in each chromatin state in normal and neoplastic B-cells.

h, Example CpGs from g, in normal and neoplastic B-cells.

i, Density of genes distributed along gene expression percentiles of genes associated

with B-cell independent CpGs gaining DNA methylation in B-cell tumors in each B-cell

tumor subtype. Expressed genes are displayed at right as control (with presence of

H3K36me3). Means within each B-cell subpopulation as well as B-cell tumors are

represented.

1026

Extended Data Fig.5.

a, Correlations between epiCMIT-hyper and epiCMIT-hypo with other

hypermethylation-based mitotic clocks including epiTOC (epigenetic Timer for Cancer

risk) and MiAge as well as the CIMP (CpG hypermethylator phenotype, not formally

conceived as mitotic clock). epiCMIT, epigenetically-determined Cumulative MIToses.

b, Poor effect of age in epiCMIT in normal B cells. epiCMIT correlation with age in naïve

B cells from healthy infants, young-adults and older adults. Horvath model correctly

predicted the real age. epiCMIT and Horvath are poorly correlated.

c, Poor effect of age in epiCMIT in B-ALL. Wide epiCMIT spectrum in children with B-

ALL.

d, Correlation between the number of mutations detected by WGS against epiCMIT in

CLL subtypes with different cellular origin (n-CLL, i-CLL and m-CLL).

e, Mutational signatures for 138 CLL samples with available WGS and DNA methylation

data divided into different subtypes of CLL with different cellular origin, namely n-CLL,

i-CLL and m-CLL.

f, Correlation of epiCMIT with Ki67 in nodal MCL patients.

g, Gene expression signatures resulting after GSEA analysis related to proliferation in

CLL samples with low and high epiCMIT.

h, Gene expression signatures resulting after GSEA analysis related to proliferation in

ALL samples with low and high epiCMIT.

1047

Extended Data Fig.6.

a, epiCMIT interpretation in normal and neoplastic B cells. Mitotic cell division occurs

in normal B cell development and DNA methylation changes accumulate at repressed

regions (blue component of the epiCMIT on the x axis). B-cell tumors arise from

different maturation stages, and thus they contain different baseline epiCMIT from the

cell of origin from which they originate (blue bar in B-cell tumors). When B-cell tumors

progress, they acquire additional DNA methylation changes at repressed regions

(shown as red component of the epiCMIT). Notably, these new DNA methylation

changes occur in both axes, i.e. changes occurring in normal B cell maturation as well

as new B-cell independent changes. This concept is shown with real data at Fig. 4b.

b, Multivariate cox regression model for overall survival in ALL for epiCMIT against

cytogenetic groups. Hazard ratio for epiCMIT correspond to 0.1 increments.

c, Multivariate cox regression model for time to first treatment in CLL samples near

diagnosis (30 months) for epiCMIT against age, number of driver alterations and

epigenetic groups based on different cellular origin. At right is shown the multivariate

cox regression model for validation series. Hazard ratio for epiCMIT correspond to 0.1

increments.

d, Multivariate cox regression model in CLL for overall survival for epiCMIT against age,

number of driver alterations and epigenetic groups for the initial and validation series

(left and right, respectively). Treated and untreated patients were included in the

validation series. Hazard ratio for epiCMIT correspond to 0.1 increments.

e, Kaplan-Meyer curves for DLBCL separated by ABC and GCB groups and low and high

epiCMIT groups based on maxstat rank statistics. Multivariate cox regression model

treating epiCMIT as continuous variable is shown at bottom. Hazard ratio for epiCMIT

correspond to 0.1 increments.

1073

Extended Data Fig.7.

a, Oncoprint showing all genetic driver alterations considered in CLL grouped by

epigenetic subgroups and ordered according to increasing levels of epiCMIT (from left

to right within each epigenetic subgroup). Other clinicobiological features including

MBL or CLL, IGHV status, Age, Binet stage, epiCMIT subgroups based on Fig. 5C, need

for treatment and patient status are shown. Distinct genetic driver alterations are

depicted with different colors and shapes. The percentage of mutated patients as well

as barplots showing the number of mutated patients for each alteration is shown at

right.

b, Driver genetic alterations without clear associations with epiCMIT. Analyses were

done globally for all CLL samples (although adjusted by epigenetic groups) as well as

within each epigenetic subgroup. Positive point estimates relate to positive

associations with epiCMIT. 95% confidence intervals are shown with colors coding for

FDR corrections. Number of patients with each genetic driver alteration is shown at

bottom.

1089

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Extended Data Fig.1

Extended Data Fig. 2

Extended Data Fig. 3

Extended Data Fig. 4

Extended Data Fig. 5

Extended Data Fig. 6

Extended Data Fig. 7