ContamLD: Estimation of Ancient Nuclear DNA Contamination Using Breakdown of Linkage Disequilibrium

Nathan Nakatsuka[1,2,3,*,†], Éadaoin Harney[1,3,4,* ,†], Swapan Mallick[1,3], Matthew Mah[1,3], Nick Patterson[3], and David Reich[1,3,5,6,†]

[1]Department of Genetics, Harvard Medical School, New Research Building, 77 Ave. Louis Pasteur, Boston, MA 02115, USA
[2]Harvard-MIT Division of Health Sciences and Technology, Harvard Medical School, Boston, MA 02115, USA
[3]Department of Human Evolutionary Biology, Harvard University, 16 Divinity Ave., Cambridge, MA 02138, USA
[4]Department of Organismic and Evolutionary Biology, Harvard University, 16 Divinity Ave., Cambridge, MA 02138, USA
[5]Broad Institute of Harvard and Massachusetts Institute of Technology, Cambridge, MA 02141, USA
[6]Howard Hughes Medical Institute, Harvard Medical School, Boston, MA 02115, USA

*co-first authors

† Corresponding authors: Nathan Nakatsuka (nathan_nakatsuka@hms.harvard.edu), Éadaoin Harney (harney@g.harvard.edu), and David Reich (reich@genetics.med.harvard.edu)

Abstract

We report a method, *ContamLD*, for estimating autosomal ancient DNA (aDNA) contamination

by measuring the breakdown of linkage disequilibrium in a sequenced individual due to the

introduction of contaminant DNA, leveraging the idea that contaminants should have haplotypes

uncorrelated to those of the studied individual. Using simulated data, we confirm that *ContamLD*

accurately infers contamination rates with low standard errors (e.g. less than 1.5% standard

error in cases with <10% contamination and data from at least 500,000 sequences covering

SNPs). This method is optimized for application to aDNA, leveraging characteristic aDNA

damage patterns to provide calibrated contamination estimates. Availability:

https://github.com/nathan-nakatsuka/ContamLD.

Keywords

Ancient DNA, linkage disequilibrium, contamination, autosomal DNA, nuclear DNA

Background

Ancient DNA (aDNA) data has emerged as a powerful tool for learning about ancient population

history, allowing direct study of the genomes of individuals who lived thousands of years in the

past [1-3]. Unfortunately, these inferences can be distorted by contamination during the

excavation and storage of skeletal material, as well as the intensive processing required to

extract the DNA and convert it into a form that can be sequenced.

Accurate measurement of the proportion of contamination in ancient DNA data is important,

because it can provide guidance about whether analysis should be restricted to sequences that

show the characteristic pattern of C-to-T mismatch to the reference genome of authentic aDNA

(if contamination is high) [4], or carried out at all. When analysis is restricted to focus only on

sequences showing evidence of characteristic ancient DNA damage, the substantial majority of

authentic sequences are usually removed from the analysis dataset, as only a fraction of

genuinely ancient sequences typically carry characteristic damage. In addition, if a sample is

contaminated by another individual with damaged DNA—which can arise for example as a

result of cross-contamination from other specimens handled in the same ancient DNA

laboratory—it is impossible to distinguish authentic sequences from contaminating ones based

on the presence or absence of characteristic ancient DNA damage.

Current methods for estimating contamination have significant limitations. Methods based on

testing for heterogeneity in mitochondrial DNA sequences (which are almost always

homogeneous in an uncontaminated individual) can be biased, because there are several

orders of magnitude of variation in the ratio of the mitochondrial to nuclear DNA copy number

across samples. Thus, samples that have evidence of mitochondrial contamination can be

nearly uncontaminated in their nuclear DNA, while samples that have no evidence of

mitochondrial contamination can have high nuclear contamination [5]. Another reliable set of

methods for estimating rates of contamination in ancient DNA leverage polymorphism on the X

chromosome in males, including the popular *ANGSD* method [6-9] and an improved

methodology that enhances power for low-coverage samples [10]. However, these methods do

not work in females.

Several methods for estimating contamination rates in nuclear DNA from modern genomes

have been published, including *ContEst* [11] and *ContaminationDetection* [12]. However, these

methods generally rely on access to uncontaminated genotype data from the individual of

interest or access to all possible contaminating individuals, neither of which is typically available

for aDNA. Another method estimated modern human autosomal contamination in aDNA from

archaic Denisovans [13] and Neanderthals [14] by producing maximum likelihood co-estimation

of sequence error, contamination, and parameters correlated with divergence and

heterozygosity. However, this method heavily relies on the significant divergence between

archaic and modern humans. A similar method, *DICE*, expanded on this method and jointly

estimates contamination rate and error rate along with demographic history based on allele

frequency correlation patterns [15]. However, this method requires both explicit demographic

modeling and high genome coverage. While this may be effective for estimation of

contamination in archaic genomes like Neanderthals and Denisovans that are highly genetically

diverged from likely contaminant individuals, it is not optimized for study of contamination

among closely related present-day human groups with complex demographic relationships

relative to each other, or contamination from individuals of the same population. In Racimo *et al.*

2016 [15], *DICE* required over 3x genome sequence coverage and solved the distinctive

problem of measuring contamination of present-day humans in a Neanderthal genome.

We report a method for estimating autosomal aDNA contamination using patterns of linkage

disequilibrium (LD) within a sample. This approach, implemented in our software *ContamLD*, is

based on the idea that when sequences from one or more contaminating individuals are present

in a sample, LD among sequences derived from that sample is expected to be diminished,

because the contaminant DNA derives from different haplotypes and therefore should have no

LD with the authentic DNA of the ancient individual of interest. Thus, the goal of the algorithm is

to determine the LD pattern the ancient individual would have had without contamination and

compare it to the LD pattern found in the sample. The LD patterns of ancient individuals are

determined using reference panels from 1000 Genomes Project populations to compute

approximate background haplotype frequencies, where haplotypes are defined as pairs of SNPs

with high correlation to each other. Contamination is then estimated by fitting a maximum

likelihood model of a mixture of haplotypes from an uncontaminated individual and a proportion

of contamination (to be estimated from the data) from an unrelated individual. *ContamLD*

corrects for mismatch of the ancestry of the ancient individual with the reference panels using

two different user-specified options. In the first option, mismatch is corrected using estimates

from damaged sequences (which, ideally, lack present-day contaminants). In the second option,

ContamLD performs an "external" correction by subtracting the sample's contamination estimate

from estimates for individuals of the same population believed to have negligible contamination

(the user could obtain this value from a *ContamLD* calculation on a male individual with a very

low estimate of contamination based on *ANGSD*). The second option has more power than the

first option and allows detection of cross-contamination by other ancient samples, but it could be

biased if a reliable estimate from an un-contaminated individual from the same population is not

available for the external correction.

We show that *ContamLD* accurately infers contamination in both ancient and present-day

individuals of widely divergent ancestries with simulated contamination coming from individuals

of different ancestries. The contamination estimates are highly correlated with estimates based

on X chromosome analysis in ancient samples that are male, as assessed using *ANGSD* [16].

ContamLD run with the first option has standard errors less than 1.5% in samples with at least

500,000 sequences covering SNPs (~0.5x coverage for data produced by in-solution

enrichment for ~1.2 million SNPs [2, 17], or ~0.1x coverage for data produced using whole-

genome shotgun sequences). With the second option, *ContamLD* has standard errors less than

0.5% in these situations, allowing users to detect samples with 5% or more contamination with

high confidence so they can be removed from subsequent analyses.

Results

Simulations of Contamination in Present-Day Individuals:

To test the performance of *ContamLD*, we simulated sequence level genetic data. For our first

simulations, each uncontaminated individual was simulated based on genotype calls from a

present-day individual from the 1000 Genomes Project dataset. To determine the sequence

coverage at each site, we used data from an ancient individual for which we had data at 1.02x

coverage and in each case generated the same number of simulated sequences at each site,

with the allele drawn from the present-day individual e.g. if the present day individual is

homozygous for the reference allele at a site, all simulated alleles are of the reference type,

while if the present day individual is heterozygous, simulated alleles are either of the reference

or alternative variant, with 50% probability of each). The damage status (i.e. whether it carries

the characteristic C-to-T damage often observed in ancient DNA sequences) of each sequence

was also determined based on the status of the ancient reference individual. Contaminating

sequences were then "spiked-in" at varying proportions (0 to 40%), using an additional present-

day individual from the 1000 Genomes Project to determine the contaminating allele type (see

Methods). All contaminating sequences were defined to be undamaged, as would be expected if

the contamination came from a non-ancient source.

For most of the analyses reported in this study, we simulate data for SNP sites targeted in the

1.24 million SNP capture reagent [2, 17] that intersect with 1000 Genomes sites, after removing

sites on the X and Y chromosomes (this leaves ~1.1 million SNPs). The *ContamLD* software

also allows users to make panels based on their own SNP sets, and in a later section we report

results from a larger panel (~5.6 million SNPs) provided with the software that we recommend

for shotgun sequenced samples, and which provides more power to measure contamination.

We first analyzed data generated using a reference individual from the 1000 Genomes CEU

population (Utah Residents (CEPH) with Northern and Western European Ancestry) and the

SNP coverage profile of a 1.02x coverage ancient individual of West Eurasian ancestry (Iberian

Bronze individual I3756 who lived 2014-1781 calBCE; see Methods). Supplementary Figure 1

illustrates the distribution of LOD (logarithm of the odds) scores generated when *ContamLD* is

run on samples with 0%, 7% and 15% simulated contamination. Supplementary Figure 2 shows

the contamination rate estimates generated for data with simulated contamination rates

between 0 to 40%. At very high contamination (above 15%) *ContamLD* often overestimates

contamination, but in practice samples with above 10% contamination are generally removed

from population genetic analyses, so inaccuracies in the estimates at these levels are not a

concern in our view (the importance of a contamination estimate in many cases is to flag

problematic samples, not to accurately estimate the contamination proportion). *ContamLD*

assumes that the individual making up the majority of the sequences is the base individual, so

we do not explore contamination rates greater than 50% in these simulation studies.

We observe a linear shift in the contamination estimates such that most estimates are biased to

be slightly higher than the actual value, with even greater overestimates occurring at higher

contamination rates (Supplementary Figure 2). This is likely due to the difference between the

haplotype distribution of the test individual and that of the haplotype panel, since the magnitude

of this shift increases as the test individual increases in genetic distance from the haplotype

panel. Even in cases where the test individual is of the same ancestry as the haplotype panel

(as in Supplementary Figure 2) there is expected to be a shift, because the test individual's

haplotypes are a particular sampling of the population's haplotypes, and the difference between

having only frequencies of the haplotype panel and a particular instantiation of those

frequencies in the test individual will lead to the artificial need for an external source

("contaminant") to fit the model properly.

In contrast to the upward bias in contamination estimates due to mismatch of the individual's

haplotypes with the reference panel haplotype frequencies, we observe negative shifts for

inbred individuals, as expected because *ContamLD* assumes the paternal and maternal copy of

a chromosome are unrelated. In contrast, if the two chromosomes are related, extra LD will be

induced and more contamination will be necessary to produce the expected LD pattern. In

principle, this inbreeding effect could be corrected explicitly by estimating the total amount of

ROH in each individual and applying this as a correction, although we do not provide such

functionality as part of our software. If a reliable methodology for quantifying the proportion of

the genome that is affected by inbreeding in ancient individuals becomes available, *ContamLD*

could be further improved by using this information as an input parameter.

A final type of bias could be expected to arise if the contamination comes from an individual

related to the target individual. In this case the true contamination rate is expected to be under-

estimated, because *ContamLD* only detects contamination where the contaminant sequence

differs from the target individual's sequence. If the contaminant carries the same haplotypes as

the target individual, in the most extreme case as expected for an identical twin, then the

existence of contamination will be missed altogether. In general, contamination from closely

related individuals is unlikely to be a concern for many population genetic analyses, as close

relatives usually (but with important exceptions) have very similar ancestry.

In our implementation, we correct for these systematic biases in two ways, implemented as

different options in *ContamLD*.

The first option leverages information from sequences that contain evidence of the C-to-T

damage that is characteristic of ancient sequences. This option assumes these sequences are

authentically ancient and not derived from a contaminating source (assumed to be from present-

day individuals), so the *ContamLD* estimate based on undamaged sequences is corrected by

estimates based on the damaged sequences (see Methods for more details). In the second

option, we allow the user to subtract the contamination estimate from the estimate of an

individual of the same ancestry assumed to be uncontaminated. An advantage of the second

option compared to the first is that it has smaller standard errors (Figure 1), reflecting the fact

that it does not rely on estimates from damaged sequences (reliance on damaged sequences

reduces power since it often reflects a very small subset of the data). A second advantage of

the second option is that it allows estimation of contamination in cases where the source of

contamination is also ancient in origin, as would be expected if the contamination occurred

anciently or due to cross contamination with other ancient samples (the first option would be

expected to produce an underestimate of contamination in such cases, since it assumes that

sequences that contain C-to-T damage are not contaminated). On the other hand, a drawback

of the second option is that it requires users to identify a relatively high coverage,

uncontaminated, ancestry-matched sampled for benchmarking purposes; the method is also

only expected to work if there is minimal inbreeding in either the sample of interest or the

matched sample. Identifying such benchmarking samples may be impossible when analyzing

samples from previously unsampled contexts (e.g. early modern humans), and indeed verifying

that a benchmarking sample is uncontaminated is very difficult if it is female (if it is male a

method like *ANGSD* can be used).

In what follows, we report results of analyses based on the first option, but *ContamLD* includes

both methods as options. The uncorrected score also forms the basis for warning output by the

software, namely high contamination or possible contamination with another ancient sample

leading to an inaccurate damage correction estimate.

A)   **B)**

Figure 1. *ContamLD* **estimates when the target individual, contaminant, and haplotype panel are**

from the CEU population. Contamination estimates when the simulated contamination rate is between

0.00-0.15. **A)** Estimates with damage restricted correction (option 1). **B)** Estimates with external

correction from an uncontaminated sample (option 2). The black dotted line is y=x, which would

correspond to a perfect estimate of contamination. Error bars are 1.96*standard error (95% confidence

interval determined via jackknife resampling across chromosomes).

Simulated Contamination of Ancient Samples with Present-Day Samples:

ContamLD is designed to work on ancient individuals, so we simulated contamination of real

ancient individuals with present-day individuals from the 1000 Genomes Project, a scenario that

would occur when skeletal material from ancient individuals is contaminated by present-day

individuals during excavation or at some point during the processing of the material. We used

data from male individuals selected due to very low X chromosome contamination estimates

(less than 1%) based on *ANGSD* [16] (developed first in Rasmussen *et al.* [9]; we used method

1 of that software). (We subtracted the *ANGSD* estimates from the *ContamLD* estimates to

correct for any residual contamination.) Figure 2A shows results from the Iberian Bronze Age

sample [18] (I3756) with 1.02x coverage at the targeted ~1.24 million SNP positions,

demonstrating that *ContamLD* produces highly accurate contamination estimates for this

simulation.

Effect of Different Haplotype Panels

There are many potential cases in which ancient individuals can come from populations with

very different genetic profiles compared to present-day 1000 Genomes populations, leading to

an ancestry mismatch to the haplotype reference panels. *ContamLD* provides panels from all

1000 Genomes populations as well as tools to identify the panel most closely matching to the

ancestry of their ancient individual (based on outgroup-f_3 statistics [19] to determine the most

shared genetic drift), which they can then select for the analysis. However, due to the potential

for ancestry mismatch to still occur, we tested the effect of choosing haplotype panels that are

genetically diverged from the individual of interest (Figure 2A). For the ancient Iberian sample,

the CEU and TSI (Toscani in Italia) panels—representing northern and southern European

ancestry, respectively—yielded contamination estimates that are close to the true contamination

rate, especially for rates below 5%. However, *ContamLD* underestimates contamination by ~2%

when the CHB (Han Chinese in Beijing, China) and YRI (Yoruba in Ibadan, Nigeria) panels were

used instead (though we view these as very pessimistic cases, because the user should usually

be able to choose a panel more closely related to their ancient individual than these scenarios).

We thus recommend that users take care to choose an appropriate panel that is within the same

continental ancestry as their ancient individual. Nevertheless, we note that we were able to

obtain reasonably accurate estimates for Upper Paleolithic European hunter-gatherers, such as

the Kostenki14 individual [20], who is ~37,470 years old, even when using present-day

European panels that have significantly different ancestry from the hunter-gatherers

(Supplementary Figure 3).

A)   **B)**

Figure 2. Genetic divergence between uncontaminated individual and contamination sources or

haplotype panels impacts *ContamLD* estimates. A) Ancient Iberian (I3756, 1.02x coverage)

contaminated with CEU with haplotype panels generated from CEU, TSI, CHB, and YRI populations. **B)**

Contamination estimates from the same ancient Iberian contaminated with TSI, CHB, or YRI and

analyzed with a CEU panel; from an ancient East Asian (DA362.SG, 1.10x coverage) contaminated with

CEU and analyzed with a CHB panel; and from an ancient South African (I9028.SG, 1.21x coverage)

contaminated with CEU and analyzed with a YRI panel. The black dotted line is y=x, corresponding to a

perfect estimate of contamination. All estimates use the damage restricted correction (option 1).

Effect of Mismatch Between the Ancestry of the True Sample and Contaminating Individual

Contamination can come from a wide variety of sources, including, but not limited to, members

of the archaeological excavation team, the aDNA laboratory, or residual human DNA on the

plastic and glassware or in laboratory reagents. Thus, we sought to understand the effect of

mismatch in the ancestry of the true sample and the contaminating individual in our

contamination estimates. We found that as the ancestry of the two diverged, *ContamLD* over-

estimated contamination (Figure 2B and Supplementary Figure 4). This occurred when we

tested an ancient European with different contaminant ancestries and when we tested ancient

East Asian [21] and ancient South African [22] samples contaminated with European DNA.

Nevertheless, the over-estimation was not severe at contamination levels below 5 percent, and

samples above this proportion would likely be flagged as problematic. We also explored

scenarios where the ancestry of the panel matches the contaminant rather than the true sample

(Supplementary Figure 4) and found a ~2% under-estimate at low levels of contamination and

an over-estimate at high levels of contamination; these are modest effects and are unlikely to

change our qualitative assessment. When we tested the effect of having multiple contaminant

individuals (Supplementary Figure 5), we found only a slight over-estimate at higher levels of

contamination, as expected given *ContamLD* normally assumes contamination from a single

individual where the haplotypes are re-formed if they are created from two contaminant reads

(which will happen at lower rates with more contaminant individuals).

Estimating Contamination in Admixed Individuals

ContamLD relies on measuring the difference between the LD pattern of the sample and that

expected from an uncontaminated individual. However, individuals from groups recently

admixed between two highly divergent ancestral groups have LD patterns that are similar in

some ways to that of an unadmixed individual with contamination from a group with ancestry

diverged from that of the individual of interest. To understand how this would impact *ContamLD*,

we ran the software on an ASW (Americans of African Ancestry in Southwest USA) individual

with different levels of added CEU contamination. When we ran *ContamLD* with a YRI panel

and no correction on an individual with no contamination, the individual was inferred to have a

contamination of ~20% (likely because the individual had ~15% European ancestry, and this

was interpreted by the software as contamination). Using an ASW panel did not perform any

better. However, the concerns were mostly addressed by the damage-restricted correction

(option 1) at low contamination levels (Supplementary Figure 6). The simulation with African-

Americans represents an extreme of difficulty, because the individual is from a group with very

recent admixture (~6 generations [23]) of ancestries highly divergent from each other with one

of the ancestries very genetically similar to the reference panel. It highlights how the damage-

restricted correction is still able to produce accurate estimates in these difficult cases.

Effect of Coverage:

We tested the power of our procedure at different coverages with simulations of ancient West

Eurasian ancestry individuals contaminated with CEU on the 1240K SNP set (Figure 3). We

found that while our estimates were not biased to produce estimates consistently above or

below the true value, the standard errors increased significantly at lower coverages, as

expected for the decreased power for accurate estimation in these scenarios. We provide a

much larger panel with ~5.6 million SNPs (vs. ~1.1 million for the 1240K panel) that usually

decreases standard errors for samples that are shotgun sequenced (Supplementary Figure 7).

This panel increases *ContamLD*'s compute time and memory requirements, so we recommend

that it only be used for individuals with lower than 0.5x coverage. As an additional feature, we

provide users tools to create their own panels to meet their specific needs.

Different Ancient Individuals contaminated
with CEU using CEU panel

Individual
- Iberian (0.46x coverage)
- Hungarian (0.27x coverage)
- Ukranian (0.015x coverage)

Figure 3. *ContamLD* estimates for ancient European samples of different coverages after damage restricted correction (option 1). An ancient Iberian of 0.46x coverage, an ancient Hungarian of 0.27x coverage, and an ancient Ukranian of 0.015x coverage (~16,000 snps) were contaminated with CEU and analyzed using a CEU panel with *ContamLD* option 1 (damage restricted correction). The black dotted line is y=x. Error shading is 1.96*standard error (95% confidence interval).

Simulations to Compare ContamLD to ANGSD X Chromosome Estimates

We performed simulations where we randomly added sequences at increasing levels from 0 to 15% from an ancient West Eurasian individual (I10895) into the BAM files of 65 ancient male individuals of variable ancestries and ages (we set the damaged sequences to be only from the non-contaminant individual; see Methods). We chose ancient male individuals that had average coverage over 0.5X and X chromosome contamination estimates under 2% (using method 1 of *ANGSD*) when no artificial contamination was added (and also corrected even for this baseline contamination by setting damaged reads to be a 5% down-sampling of the files that had no artificial contamination; see Methods). We then analyzed the individuals with *ContamLD* and

ANGSD and found that compared to *ANGSD*, *ContamLD* consistently had similar errors relative

to the true contamination level (Figure 4, Supplementary Online Table 2).

Figure 4. Contamination estimates with *ContamLD* and *ANGSD* for ancient individuals with

different levels of contamination added. 65 ancient individuals with average coverage over 0.5X had

increasing levels of artificial contamination added in (from I10895, an ~1200BP ancient West Eurasian

individual) and were then analyzed with *ContamLD* (with panels most genetically similar to the ancient

individual and using damage restricted correction, option 1) and *ANGSD*. Details of all estimates

(including standard errors) are provided in Supplementary Online Table 2. The black dotted line is y=x,

which would correspond to a perfect estimate of the contamination.

Comparing ContamLD, ANGSD, and Mitochondrial Estimates (ContamMix) in Ancient

Individuals without Added Contamination

We tested 439 ancient males with *ContamLD*, *ANGSD* (X chromosome contamination

estimates), and *ContamMix* (mitochondrial contamination estimates) without adding additional

contamination. For this analysis, we included published data generated with the ~1.24 million

SNP enrichment reagent, as well as data from libraries that failed quality control due to

evidence of contamination (Supplementary Online Table 3). Similar to prior studies [5], the

mitochondrial estimates often differed from the nuclear (*ANGSD* and *ContamLD*) estimates,

showing high contamination in some libraries with low nuclear contamination, and low

mitochondrial contamination in some libraries with high nuclear contamination (Figure 5A). In

contrast, *ANGSD* and *ContamLD* had better concordance. However, we observed that some of

the samples with high contamination estimates based on *ANGSD* had much lower *ContamLD*

estimates, reflecting over-correction from analyzing the damaged sequences, perhaps because

the contamination was actually cross-contamination from other ancient individuals, violating the

assumptions of our damage-correction (Figure 5B). This problem was mitigated in part,

however, because *ContamLD* produces a warning of "Very_High_Contamination" if the

uncorrected estimate is above 15% (even in cases where the corrected estimate is very low),

and all samples with X chromosome estimates over 5% were flagged with this warning and/or

had estimates of over 5% contamination with *ContamLD* (all samples with less than 5%

contamination in *ANGSD* had lower than 5% contamination with *ContamLD*). It is unfortunately

not possible to know the true contamination of the samples we tested in Figure 5, but the fact

that our software produced results with good correlation to X chromosome estimates shows that

it works well in real ancient data.

It is possible for there to be samples with moderately high contamination from another

ancient individual but both a low damage restricted correction estimate and no warning

generated, because these would have high uncorrected estimates, yet not high enough to reach

the threshold required for the warning. These samples would have to be identified with an

external correction. Lowering the threshold for the "Very_High_Contamination" warning would

produce too many false positives, because there are many cases with high uncorrected

estimates that have low corrected estimates that are likely not contaminated (e.g. due to

ancestry mismatches of the panel and the test individual). To understand these issues better,

we performed a simulation in which an ancient Iberian (I3756) was contaminated with another

ancient West Eurasian individual (I10895) and the damaged sequences were set to be a 5%

down-sampling of the set of contaminated sequences (thus simulating a case in which all of the

contamination is from another ancient individual who has the same damage proportion as the

ancient individual of interest). We found that, as expected, the contamination from the ancient

individual was not detected (the contamination estimates were always near 0%) by the damage

restricted correction version of *ContamLD* until the contamination reached 15% at which point

the "Very_High_Contamination" flag came up (Supplementary Figure 9). The contamination

would have been detected with the external correction version of *ContamLD* (since the damage

restricted correction continued to go up with increasing contamination; see Supplementary

Online Table 4), but without an uncontaminated ancient individual of the same group as the

target individual, this would be difficult to do without bias in the contamination estimate.

(A)   **(B)**

Figure 5. Contamination estimates from *ContamLD*, *ANGSD*, and *ContamMix* in 439 ancient

individuals of variable ancestry. *ANGSD* estimates (method 1) are plotted on the X-axis, and on the Y-

axis are either **(A)** *ContamMix* or **(B)** *ContamLD* estimates. In red are samples that were flagged in

ContamLD as "Very_High_Contamination" based on having uncorrected estimates over 15%. All

ContamLD estimates below 0 were set to 0.

Discussion and Conclusion

We have presented a tool, *ContamLD*, for estimating rates of autosomal DNA contamination in

aDNA samples. *ContamLD* is able to measure contamination accurately in both male and

female individuals, with standard errors less than 1.5% for individuals with coverage above 0.5X

on the 1240K SNP set (for contamination levels less than 10%) for the damage restricted

correction method (option 1). On the shotgun panel we provide, standard errors are less than

1.5% for coverages above 0.1x. *ContamLD* is best suited to scenarios in which the contaminant

and the ancient individual of interest are similar ancestry, which is useful, because *DICE* [15]

and many population genetic tools (e.g. PCA or ADMIXTURE [24]) are better suited for

detecting cases where the contaminant is of very different ancestry from the ancient individual

of interest. *ContamLD* works even for recently admixed individuals. Lastly, *ContamLD* can

detect cases of contamination from other ancient individuals, though this works best if it is large

amounts of contamination that can reach the threshold required for the

"Very_High_Contamination" flag.

We tested *ContamLD* in multiple simulation scenarios to determine when bias or less reliable

results could be expected. When applied to the situation with a test individual (ancient or

present-day), contaminant, and haplotype reference panel all from the same continental

ancestry, *ContamLD* provides an accurate, unbiased estimate of contamination. When the

contaminant comes from a population that is of a different continental ancestry from the

population used for the base and haplotype panel, the contamination appears to be slightly

overestimated, particularly for higher contamination. This should not be a large problem in

analyses of real (i.e. non-simulated) data, because the effect is small at the contamination levels

of interest (<5%). When we varied haplotype panels, we found that the estimator is robust when

applied to simulated datasets using haplotype panels that are moderately divergent from the

base sample (within-continent levels of variation). We provide users tools for automatically

determining the panel that shared the most genetic drift with the sample so that the user can

select the panel most closely related to the sample. In other simulations, we found that the

performance of the algorithm declines as the coverage of the sample decreases. The estimates

are not biased, the standard errors substantially increase when fewer than 300,000 sequences

are available. In these cases, if the individual was shotgun sequenced, we recommend that

users choose the shotgun panel, which will substantially increase power for the analyses.

We applied the algorithm to estimate contamination levels in dozens of ancient samples and

compared them to X chromosome-based contamination estimates. There was generally good

correlation with the X chromosome estimates, except that when the true contamination was very

high, the LD based estimates were sometimes estimated incorrectly, likely because the

contamination was due to cross-contamination from another ancient individual and there was

over-correction from the damage estimates. This problem is mitigated, however, because the

software indicates if the uncorrected estimate is very high so users can identify highly

contaminated samples and remove them from further analyses. A difficult case for the software

is if there is contamination in part from another ancient sample. This can cause an over-

correction and lead to an under-estimate of the contamination. The "Very_High_Contamination"

warning catches very high contamination from other ancient samples, but it will miss cases of

moderate levels of contamination from other ancient samples, because it will not reach the

threshold required for the warning. In theory, the user can determine the true contamination in

these cases using the external correction, but the external correction can be difficult if the user

does not have an adequate sample to correct the estimate of the sample of interest. The

damage correction of the software also does not work if the samples have undergone full UDG

treatment (no damaged sequences), and for this case, the external correction is the only option.

The software run-time is dependent on SNP coverage. If ~1,000,000 SNPs are covered (the

depth of the coverage on each SNP does not affect run-time), the analysis takes approximately

2 hours if 3 cores are available on CentOS 7.2.15 Linux machines (~25 GB of memory). The

software is designed for samples to be run in parallel, so the total time for analysis even for

large numbers of samples is often not much greater than the time for a single sample.

In summary, *ContamLD* is able to estimate autosomal nuclear contamination in ancient DNA

accurately with standard errors that depend on the coverage of the sample. This will be

particularly useful for female samples where X chromosome estimates are not possible. As a

general recommendation for users, we believe in most cases all samples with a contamination

estimate that is greater than 0.05 (5%) should be removed from further analyses, or the

contamination should be explicitly modeled in population genetic analyses.

Supplementary Data:

The Supplementary Data include 4 Excel spreadsheets detailing all ancient samples used and the contamination estimates for this algorithm. Also included are 10 supplementary figures.

Materials and Methods

Datasets:

Present-day samples:

Genome wide datasets from individuals that were part of the 1000 Genomes Project [25] were used as present-day reference samples. We restricted to autosomal sites included in the aDNA ~1.24 million SNP capture reagent [2, 17] and to SNPs at greater than 10% minor allele frequency in the pooled 1000 Genomes Project dataset [25]. However, the software allows users to make panels based on their own SNP set. In the analyses presented here, we filtered for SNPs that were present in the 1000 Genomes dataset and also removed all sex chromosome SNPs leading to 1,085,678 SNPs in the final 1240K dataset and 5,633,773 SNPs in the final shotgun dataset.

Ancient data set:

We analyzed mitochondrial and X chromosome contamination estimates [16, 26] from ancient individuals from previous studies generated by shotgun sequencing or targeted enrichment with 1.24 million SNP enrichment, including many samples that failed quality control due to contamination but were from the same archaeological sites [2, 22, 27-33]. Information about the ancient individual data are detailed in Supplementary Online Table 1 and below.

Obtaining sequence information:

For each ancient individual, we generated the sequence-depth data from the sample bam file,

counting the number of reference and alternative alleles at each SNP site in the analysis

dataset. Damage-restricted data was generated by restricting to sequences with PMD scores

greater than or equal to 3 [4]. Our software can accommodate both genotype call data as well

as sequence data (the sequence data adds additional power to the analyses), but all analyses

were performed using the sequence-based method. We provide users with tools to pull down

read count data from BAM files in the format required for *ContamLD*.

Haplotype Calculation

527
To create haplotype panels, we obtained all SNP pairs in high LD for each 1000 Genomes

population using PLINK version 1.9 [34] with r^2 cut-off of 0.2. (Users can increase power slightly

at the expense of increased computational time by creating their own haplotype panel with a

lower r^2 cutoff.) We then calculated the frequencies of each SNP in all of these pairs as well as

the haplotype frequencies at each of these pairs while holding out the present-day individuals

used for contamination simulation.

Algorithm to Estimate Contamination

536
Overview:

Our goal is to estimate α, the level of contamination, by examining the frequencies of allele pairs

that should be in LD (we term this two-allele pair a haplotype) and determining how much their

frequencies differ from what would be expected under no contamination. To estimate this, we

need both the distribution underlying the haplotypes (q) that an uncontaminated test sample

should have as well as the distribution of "unrelated haplotypes" ($\tilde{h}$) that would form by chance

543 from background allele frequencies. Here and below, "distribution" refers to the set of

544 frequencies of the different possible haplotypes (all possible combinations of ancestral and

545 derived alleles at the SNP pair) across all haplotypes in the genome. Supplementary Figure 10

546 is a schematic of the algorithm.

548 *Determining Haplotype Distributions Based on Reference Panels:*

549 To determine q we must account for the fact that the test individual's genotypes do not have

550 diploid calls and are not phased. Due to the low sequence depths at each SNP in many ancient

551 DNA datasets, it is difficult to make confident heterozygous calls, so instead we create pseudo-

552 haploid calls by randomly choosing a sequence to represent the genotype at that position (this

553 holds when we are using genotype calls or the sequence information directly, and when multiple

554 sequences cover the same SNP, we use all of them and treat them as independent). Thus, for

555 this analysis, when examining a pair of SNPs, it is equally likely for the SNP pair to have been

556 formed from the true haplotype (if the same parental chromosome is sampled from in both

557 SNPs of the haplotype) or the background distribution (if the opposite parental chromosome is

558 sampled from). We therefore can estimate q as:

559
$$q = h/2 + \widetilde{h}/2$$

560 where h is the distribution of true haplotypes and $\tilde{h}$ is the distribution of unrelated haplotypes

561 that would form by chance from background allele frequencies. For inbred samples, the weight

562 on h is more than 1/2, because the two parental chromosomes are more related, but this can

563 generally be corrected (see below).

565 $\tilde{h}$ can be determined by multiplying the SNP frequencies to obtain the haplotype frequencies

566 that would form after randomly pairing SNPs of unrelated individuals. h can be estimated from

567 an external reference panel using a maximum likelihood estimator (MLE) to obtain haplotypes

568   frequencies in the population from the counts (necessary because the panels are not phased).

569   The MLE set-up is:

$$log(L(h|c)) = \sum_{j=1}^{n} \sum_{i=1}^{4} c_{ij} log(P(i,j|h))$$

573   with:

$$P(i,j|h) = \sum_{a_1,a_2,b_1,b_2=0,1;a,b \rightarrow (i,j)} h_{(a_1,b_1)} * h_{(a_2,b_2)}$$

where $P(i, j|h)$ is the (unknown) diploid count distribution of the haplotypes of the population the

test individual is from (approximated by the external panel), n is the number of SNP pairs, c is

the vector of observed haplotypes in the diploid count panel (from 1000 Genomes), i sums over

all 4 haplotype possibilities, $h_{(a,b)}$ are the (also unknown) haplotype distributions of the parents of

the test individual (the haploid chromosomes they pass on to their child), and a, $b \rightarrow (i, j)$ implies

that $a_1 + a_2 = i$ and $b_1 + b_2 = j$, meaning that one adds up all cases where the haplotype

combination would lead to a particular diploid count (e.g. in the notation, for example, 01,11

means the first parent contributes a haplotype that has 0 alternative alleles at the first SNP and

1 alternative allele at the second SNP, and the second parent contributes a haplotype where

both SNPs have the alternative allele. The test individual with these parents would then have a

12 diploid count, which means at the first SNP the individual has 1 alternative allele and at the

second SNP the individual has 2 alternative alleles. Since our observed data are not phased,

both 01,11 and 11,01 would lead to a 12 diploid count). This assumes independence of SNP

pairs, which is not true, but because our standard errors are based on jackknife resampling

across chromosomes, correlation among SNP pairs is corrected for in our error estimates.

The MLE would be computationally intractable to solve due to our lack of knowledge of which

parent contributed to each count, so we instead used an EM algorithm to obtain h, where

knowledge of the parents' contribution is the unobserved latent variable. The algorithm involves

an expectation step of:

$$n_1 = \frac{C_{(i,j)} * \sum_{a,b \to (i,j)} h_{(a,b)} * h_{(a_2,b_2)}}{P(i,j|h)}$$

where n_1 is the expected number of times that the (a, b) configuration of the father's

chromosome contributed to a particular diploid count (this is the same value for the mother, n_2,

because they are assumed to be from the same haplotype distribution). In other words, given

the observed haplotype counts in the reference panel, how many times would it be expected

that a particular haplotype configuration (e.g. ancestral at SNP1, derived at SNP2) in one of the

parents contributed to those counts?

Once the counts (n_1 and n_2) of the haploid parents are obtained, they are added together to

produce the diploid individual (i.e. the expected number of all possible haplotype

configurations). Then the expected value of the haplotype distribution can be maximized by

averaging over the possible haplotype distributions. Thus, the maximization step is:

611
$$D_{(a,b)} = \sum_{(i,j)} C_{(i,j)} * [n_1 + n_2]$$

$$\hat{h_{(a,b)}} = \frac{D_{(a,b)}}{\sum_{a,b} D_{(a,b)}}$$

where $D(a,b)$ is the sum of the probabilities of a particular haplotype configuration over all

diploid count configurations.

We initially set all $h(a,b)$ to be 0.25 and then iterated through the algorithm until convergence

(using a squared distance summed over all SNPs and a threshold of 0.001). We then used this

estimate of h to get an estimate of q (based on the first equation above).

Estimating Contamination Based on Haplotype Distributions and Test Individual's Haplotypes:

To estimate α, we used the equation:

$$T = (1-2\alpha'+2\alpha'^2)q+2\alpha'(1-\alpha')\tilde{h}$$

Here T is the expected distribution underlying the observed haplotypes of the sample, which is a

mix of the test individual and contaminant. This means that assuming the test individual comes

from a population with a haplotype distribution (frequency of the different haplotype possibilities

at each SNP pair throughout the genome) that can be approximated by the chosen reference

panel (and estimated as above), T is the haplotype distribution expected for the sample given a

particular amount of contamination (α', where ' is used to indicate that this is an estimate of the

real α). q is the haplotype distribution for an uncontaminated sample. A fraction $(1 - \alpha')^2 + \alpha'^2$ of

the distribution should look like this, where $(1 - \alpha')^2$ is the probability that two uncontaminated

sequences form the SNP pair and α'^2 is the probability that two contaminated sequences form

the SNP pair, assuming the contaminating sequences are from a single individual, which would

"re-form" a SNP pair with LD (note: this also makes the simplifying assumption that the

contaminant and the test individual have the same background haplotype and SNP distribution).

$\tilde{h}$ is the distribution of unrelated "haplotypes" that would form by chance from background allele

frequencies in the population. Contamination would form these unrelated haplotypes by

breaking up LD, so a fraction $2\alpha'(1 - \alpha')$ of the distribution should look like this (the probability

that the SNP pair is formed from a contaminated sequence and an uncontaminated sequence).

This expression can be used to solve for α' by maximizing the LOD (log of the odds) scores

under the null hypothesis that $\alpha' = 0$ and the alternative hypotheses of different α'. A LOD score

is assigned to each estimate of the contamination rate (α) between -0.1 to 0.5 (negative scores

are included to allow correction for inbreeding). The grid of α' is scaled by intervals of 0.0001.

The α' with the highest LOD score is the best estimate of α, and is returned. When we have

multiple sequences on the same SNP we assume independence of the sequences, which

provides additional power. The assumption of independence does not bias the error estimation

for the same reason as explained above for independence of SNP pairs.

Correcting for Bias in Contamination Estimates:

In practice, the α' we obtain is not equal to the true α, because the reference panel does not

perfectly capture the SNP and haplotype frequencies of the test sample. We found that this

difference causes a linear shift in contamination estimate where the mismatch between the

sample individual and the reference panel leads to a positive shift while inbreeding leads to a

negative shift. These biases can be addressed in either of two ways.

First, for the "damage correction" approach, we performed an α' estimate only on alleles from

sequences with evidence of damage characteristic of ancient samples. Under the assumption

that these sequences are not affected by present-day contamination, the inferred α' would be an

estimate of the bias, which can be subtracted out from the estimate based on all sites. We

separately analyzed the following pairs of SNPs: UU (both SNPs at undamaged sequences),

DU (one site damaged and the other undamaged), and DD (both SNPs at damaged

sequences). For the UU pairs, the value we calculate would be $\alpha + k$, where k is the linear shift.

For DU pairs the value calculated would be $\alpha/2 + k$, and for DD pairs the value calculated would

be k. We added the likelihoods for these pairs and maximized the likelihood to solve for α and k.

After solving for α, we multiply by (1-damage rate) to obtain the contamination level across all

sequences, because α is the contamination rate at undamaged sequences.

Second, for the "external correction" approach, we took individuals from the test individual's

population that were high coverage and samples we believed had very low contamination

(based on X chromosome estimates with *ANGSD* using method 1 as developed first by

Rasmussen *et al*. [9]) and measured α'. We assumed a true contamination of 0 for these

samples and thus subtracted this α' from all other contamination estimates. We caution that this

method does not correct for uncertainty in the contamination estimate in the external sample

used for benchmarking.

Comparison to a Similar Method:

The approach of *ContamLD* is similar to that of Vohr *et al*. [35] except the two have the opposite

goals. Vohr *et al*. searches for LD in reads from two different samples in an attempt to

determine whether the two samples are from the same individual (or closely related individuals),

using a reference panel to determine LD patterns. In contrast, *ContamLD* searches for breaks in

LD in the sequences of a single sample to determine if sequences from other individuals are

present in the sample.

Data simulation:

To test the accuracy of the algorithm, we applied it to a variety of scenarios with both present-

day DNA as well as real aDNA samples that had simulated present-day DNA contamination. In

all our simulations with 1000 Genomes individuals, we removed the individual being used from

our haplotype panel before performing the analyses.

Simulating Contamination of Present-day Individuals:

We first simulated contamination of present-day individuals with other present-day individuals as

contaminants (this allowed us to be sure that there was no baseline contamination). In order to

best approximate the distribution of both the damaged and undamaged sequences that is

characteristic of aDNA data, we used sequence-depth information from an ancient individual as

a reference. At each SNP, the total number of simulated "damaged" and "undamaged"

sequences was determined based on the number of damaged and undamaged sequences at

the SNP in the reference ancient individual. The identity of each allele for the present-day

"base" sample was randomly chosen based on the genotype of the "base" present-day 1000

Genomes individual at each SNP, as described above for the contamination. The addition of

contaminant sequences to the dataset was performed using the method described above. In

order to reduce bias caused by the damage correction procedure, the damage restricted dataset

was generated only once for each simulation type (which included multiple simulations across

varying contamination rates) and combined with the undamaged dataset to produce the overall

dataset. This method was used to generate a simulated individual using present-day CEU

(NA06985) or ASW (NA19625) from the 1000 Genomes dataset as the "base" sample from the

sequence distributions of a 1.02x coverage ancient Iberian individual (I3756) (the "reference")

[18]. The CEU (NA06984) individual was used as "contaminant" in each case.

We generated simulated data with contamination from multiple sources by adjusting the

present-day contamination simulation method to randomly sample from two or more present-

day source contaminant genomes with equal probability. In each case, a 1000 Genomes Project

CEU individual (NA06985) was used as a "base" genome with the sequence distribution of

I3756 (the "reference"). In the case of 2 sources of contamination (Supplementary Figure 5), two

CEU individuals from the 1000 Genomes Project dataset (NA06984 and NA06986) were used

as contamination sources, and in the case of three contamination sources, an additional CEU

individual was used (NA06989). Data was generated for all combinations of undamaged

contamination rates, α, from 0-15%.

Simulated contamination of ancient individuals:

We performed two sets of simulations contaminating different ancient individuals. In both cases

we selected ancient male individuals with minimal contamination (as assessed by X

chromosome contamination levels from *ANGSD* [16]) to act as the "base" uncontaminated

genome. In the first simulation set, we tested *ContamLD*'s performance with different ancient

individuals and different present-day contaminant individuals from the 1000 Genomes dataset

[25] to assess the impact of contaminant ancestry and coverage of the ancient individual. In this

case we were only using *ContamLD* and thus we performed the simulated contamination on the

genotype level. In the second simulation set, we compared *ContamLD* to *ANGSD* and used a

~1200BP ancient West Eurasian individual (I10895) to contaminate the BAM files directly.

In the first simulation set, we assumed that sequences with C-to-T damage are highly unlikely to

be the product of contamination (this assumption would be falsified in the context of cross-

contamination by another ancient DNA sample). Thus, we exclusively added contamination to

the "undamaged" fraction of sequences. At each SNP site, we classified sequences present in

the damage-restricted dataset as "damaged" and added to the simulated data. We classified all

other sequences as "undamaged" and also added them to the simulated data, but for each

"undamaged sequence" we added a contaminant sequence to the simulated SNP data with

probability $\alpha/(1-\alpha)$, where α is equal to the contamination rate (since the added sequences

contribute to the total number of sequences, we needed to add a higher proportion than the

contamination rate to obtain our desired contamination rate). The identity of the added

contaminant allele was randomly chosen based on the genotype of the chosen "contaminant"

present-day genome at the site (i.e. if the contaminant individual was homozygous at the site,

the allele it possesses would be added to the simulated individual, while if it were heterozygous

at the site, either the reference or alternative allele would be selected randomly and added to

the simulated individual). This method maintains the underlying distribution of "uncontaminated"

reference and alternative alleles at each SNP site, while adding additional "contaminant" alleles

to each site, producing an overall contamination rate of α in the undamaged sequences.

For each simulation, we generated two output files: (1) a file reporting the total number of

sequences carrying reference and alternative alleles at each SNP and (2) a damage restricted

file reporting the total number of damaged sequences carrying reference and alternative alleles

at each SNP. We used a 1.02x coverage ancient Iberian individual (I3756) (Supplementary

Online Table 1) with contamination from either the 1000 Genomes CEU individual NA06984, the

TSI individual NA20502, the CHB individual NA18525, or the YRI individual NA18486. We also

used 5 other ancient individuals: I1845 (an ancient Iberian sample of 0.46x coverage) [18],

I2743 (an ancient Hungarian of 0.27x coverage) [30], I5891 (a Neolithic Ukrainian individual of

0.016x coverage) [36], DA362.SG (a Russian early Neolithic Shamanka East Asian individual of

1.10x coverage) [21], and I9028.SG (a South African individual of 1.21x coverage) [22]. In each

case, we simulated individuals with 0-15% contamination.

For the second simulation set, we analyzed 65 ancient individuals of average coverage over

0.5X and baseline *ANGSD* estimates under 2% (Supplementary Online Table 2). In these

cases, we added artificial contamination with sequences from a ~1200BP ancient West

Eurasian individual (I10895) into the BAM files at the amounts: (0.000, 0.005, 0.010, 0.020,

0.025, 0.030, 0.040, 0.050, 0.060, 0.070, 0.080, 0.090, 0.100, 0.150). We removed two base

pairs from the end of each sequence of partial UDG treated samples and ten nucleotides for

non-UDG treated samples and pulled down the genotypes by randomly selecting a single

sequence at each site covered by at least one sequence in each individual to represent the

individual's genotype at that position ("pseudo-haploid" genotyping). To ensure that the damage

sequences were only from the non-contaminant individual (so that we could use the damage

restricted correction mode, option 1, of *ContamLD* without bias), we created the "damaged"

sequence set as a randomly chosen 5% of the sequences from the non-contaminant individual.

We then analyzed the data with *ContamLD* (damage restricted correction version, option 1) and

ANGSD using default settings (Method 1). We also performed simulations with a 1.0x coverage

ancient West Eurasian ancestry individual (DA57.SG, an ancient Krgyzstanian individual) [37]

down-sampled to 0.5x coverage and contaminated with I10895. To simulate different damage

rates, we varied the damage rate to the proportions (0.005, 0.01, 0.02, 0.03, 0.04, 0.05, 0.075)

by setting the amount of "damaged" sequences to be those proportions.

As a last simulation, we examined the case of an ancient individual contaminating another

ancient individual where some of the damaged sequences would also come from the

contaminating individual. In this simulation, we analyzed a 1.02x coverage ancient Iberian

individual (I3756) and contaminated the BAM with sequences from a ~1200BP ancient West

Eurasian individual (I10895) in the proportions (0.000, 0.005, 0.010, 0.020, 0.025, 0.030, 0.040,

0.050, 0.060, 0.070, 0.080, 0.090, 0.100, 0.150, 0.200, 0.300). We then down-sampled the

BAM, taking a random 5% of the sequences of these contaminated BAM files to act as the

"damaged" sequences, because this would correct for any baseline contamination in the I3756

individual yet would simulate additional contamination of I3756 by an ancient individual with the

same damage rate as I3756 (i.e. if there is 5% contamination, then also 5% of the damaged

sequences would be from the contaminant individual in this simulation). We then performed the

standard processing of both the full contaminated BAMs and the 5% down-sampled BAMs

(simulated to be "damaged" sequences), removing two base pairs from the end of each

sequence and carrying out a "pseudo-haploid" genotype pulldown. We ran *ContamLD* on the

resulting data with damage restricted correction, option 1.

Direct Analyses of Contamination Levels in Ancient Individuals:

As our last set of analyses, we directly measured contamination levels in ancient individuals

without simulated contamination. We used *ContamLD* to examine shotgun sequenced

individuals analyzed at the1240K SNP set and the large 5.6 million SNP shotgun panel. The

ancient shotgun sequenced individuals were of 0.1-0.5x coverage from Allentoft *et al*., 2015

[31], Damgaard *et al.*, Nature 2018 [37], and Damgaard *et al*., Science 2018 [21]. In addition, we

analyzed 439 individuals from a variety of ancestries with *ContamLD* (damage corrected

version), *ANGSD* [16, 38] using default settings (we report the results from Method 1), and

contamMix [39] with the settings: down-sampling to 50X for samples above that coverage, --

trimBases X (2 bases for UDG-half samples and 10 bases for UDG-minus samples), 8 threads,

4 chains, and 2 copies, taking the first one that finishes. Supplementary Online Table 1 includes

all information from these individuals.

Declarations

Ethics Approval and Consent to Participate

Not applicable (all samples were from previously published studies).

Consent for publication

Not applicable.

Availability of Data and Materials:

All data analyzed in this article are available in [2, 21, 22, 27-33, 37]. The software is available

at: https://github.com/nathan-nakatsuka/ContamLD. It requires Python 3 and R (any version

should suffice). Archived version (1.0) used for analyses in this manuscript:

https://zenodo.org/record/3736774#.XoTbj257mgQ (DOI: 10.5281/zenodo.3736774)

Scripts for data simulations are available in the Github folder "Simulation_Scripts".

Competing interests:

The authors declare that they have no competing interests.

Funding:

Funding was provided by an NIGMS (GM007753) fellowship to NN and a MHAAM fellowship to

EH. DR is an Investigator of the Howard Hughes Medical Institute this work was supported by

grants HG006399 and GM100233 from the National Institutes of Health, by an Allen Discovery

Center grant from the Paul Allen Foundation, and by grant 61220 from the John Templeton

Foundation.

Authors' Contributions:

N.N., E.H., N.P., and D.R. conceived the study. N.N., E.H., and S.M. performed analysis. N.N.,

E.H., and D.R., wrote the manuscript with the help of all co-authors.

Acknowledgements:

We thank Iosif Lazaridis and Mark Lipson for helpful discussions.

References

844   1.   Dabney J, Knapp M, Glocke I, Gansauge MT, Weihmann A, Nickel B, Valdiosera C,
845   Garcia N, Paabo S, Arsuaga JL, Meyer M: **Complete mitochondrial genome**
846   **sequence of a Middle Pleistocene cave bear reconstructed from ultrashort DNA**
847   **fragments.** *Proc Natl Acad Sci U S A* 2013, **110:**15758-15763.

848   2.   Haak W, Lazaridis I, Patterson N, Rohland N, Mallick S, Llamas B, Brandt G, Nordenfelt
849   S, Harney E, Stewardson K, et al: **Massive migration from the steppe was a source**
850   **for Indo-European languages in Europe.** *Nature* 2015, **522:**207-211.

851   3.   Rohland N, Harney E, Mallick S, Nordenfelt S, Reich D: **Partial uracil-DNA-**
852   **glycosylase treatment for screening of ancient DNA.** *Philos Trans R Soc Lond B Biol*
853   *Sci* 2015, **370:**20130624.

854   4.   Skoglund P, Northoff BH, Shunkov MV, Derevianko AP, Paabo S, Krause J, Jakobsson
855   M: **Separating endogenous ancient DNA from modern day contamination in a**
856   **Siberian Neandertal.** *Proc Natl Acad Sci U S A* 2014, **111:**2229-2234.

857   5.   Sawyer S, Renaud G, Viola B, Hublin JJ, Gansauge MT, Shunkov MV, Derevianko AP,
858   Prufer K, Kelso J, Paabo S: **Nuclear and mitochondrial DNA sequences from two**
859   **Denisovan individuals.** *Proc Natl Acad Sci U S A* 2015, **112:**15696-15700.

860   6.   Fu Q, Meyer M, Gao X, Stenzel U, Burbano HA, Kelso J, Paabo S: **DNA analysis of an**
861   **early modern human from Tianyuan Cave, China.** *Proc Natl Acad Sci U S A* 2013,
862   **110:**2223-2227.

863   7.   Green RE, Malaspinas AS, Krause J, Briggs AW, Johnson PL, Uhler C, Meyer M, Good
864   JM, Maricic T, Stenzel U, et al: **A complete Neandertal mitochondrial genome**
865   **sequence determined by high-throughput sequencing.** *Cell* 2008, **134:**416-426.

866   8.   Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, Patterson N, Li H, Zhai
867   W, Fritz MH, et al: **A draft sequence of the Neandertal genome.** *Science* 2010,
868   **328:**710-722.

869   9.   Rasmussen M, Guo X, Wang Y, Lohmueller KE, Rasmussen S, Albrechtsen A, Skotte L,
870   Lindgreen S, Metspalu M, Jombart T: **An Aboriginal Australian genome reveals**
871   **separate human dispersals into Asia.** *Science* 2011, **334:**94-98.

872   10.   Moreno-Mayar JV, Korneliussen TS, Dalal J, Renaud G, Albrechtsen A, Nielsen R,
873   Malaspinas A-S: **A likelihood method for estimating present-day human**
874   **contamination in ancient male samples using low-depth X-chromosome data.**
875   *Bioinformatics* 2020, **36:**828-841.

876   11.   Cibulskis K, McKenna A, Fennell T, Banks E, DePristo M, Getz G: **ContEst: estimating**
877   **cross-contamination of human samples in next-generation sequencing data.**
878   *Bioinformatics* 2011, **27:**2601-2602.

879   12.   Jun G, Flickinger M, Hetrick KN, Romm JM, Doheny KF, Abecasis GR, Boehnke M,
880   Kang HM: **Detecting and estimating contamination of human DNA samples in**
881   **sequencing and array-based genotype data.** *Am J Hum Genet* 2012, **91:**839-848.

882   13.   Meyer M, Kircher M, Gansauge M-T, Li H, Racimo F, Mallick S, Schraiber JG, Jay F,
883   Prüfer K, De Filippo C: **A high-coverage genome sequence from an archaic**
884   **Denisovan individual.** *Science* 2012, **338:**222-226.

885   14.   Prüfer K, Racimo F, Patterson N, Jay F, Sankararaman S, Sawyer S, Heinze A, Renaud
886   G, Sudmant PH, De Filippo C: **The complete genome sequence of a Neanderthal**
887   **from the Altai Mountains.** *Nature* 2014, **505:**43-49.

888   15.   Racimo F, Renaud G, Slatkin M: **Joint estimation of contamination, error and**
889   **demography for nuclear DNA from ancient humans.** *PLoS genetics* 2016, **12**.

16.   Korneliussen TS, Albrechtsen A, Nielsen R: **ANGSD: Analysis of Next Generation**
Sequencing Data. *BMC Bioinformatics* 2014, **15:**356.
17.   Mathieson I, Lazaridis I, Rohland N, Mallick S, Patterson N, Roodenberg SA, Harney E,
Stewardson K, Fernandes D, Novak M, et al: **Genome-wide patterns of selection in**
230 ancient Eurasians. *Nature* 2015, **528:**499-503.
18.   Olalde I, Mallick S, Patterson N, Rohland N, Villalba-Mouco V, Silva M, Dulias K,
Edwards CJ, Gandini F, Pala M: **The genomic history of the Iberian Peninsula over**
the past 8000 years. *Science* 2019, **363:**1230-1234.
19.   Patterson N, Moorjani P, Luo Y, Mallick S, Rohland N, Zhan Y, Genschoreck T, Webster
T, Reich D: **Ancient admixture in human history.** *Genetics* 2012, **192:**1065-1093.
20.   Seguin-Orlando A, Korneliussen TS, Sikora M, Malaspinas A-S, Manica A, Moltke I,
Albrechtsen A, Ko A, Margaryan A, Moiseyev V: **Genomic structure in Europeans**
dating back at least 36,200 years. *Science* 2014, **346:**1113-1118.
21.   de Barros Damgaard P, Martiniano R, Kamm J, Moreno-Mayar JV, Kroonen G, Peyrot
M, Barjamovic G, Rasmussen S, Zacho C, Baimukhanov N: **The first horse herders**
and the impact of early Bronze Age steppe expansions into Asia. *Science* 2018,
360:eaar7711.
22.   Skoglund P, Thompson JC, Prendergast ME, Mittnik A, Sirak K, Hajdinjak M, Salie T,
Rohland N, Mallick S, Peltzer A, et al: **Reconstructing Prehistoric African Population**
Structure. *Cell* 2017, **171:**59-71 e21.
23.   Patterson N, Hattangadi N, Lane B, Lohmueller KE, Hafler DA, Oksenberg JR, Hauser
SL, Smith MW, O'Brien SJ, Altshuler D: **Methods for high-density admixture mapping**
of disease genes. *The American Journal of Human Genetics* 2004, **74:**979-1000.
24.   Alexander DH, Novembre J, Lange K: **Fast model-based estimation of ancestry in**
unrelated individuals. *Genome Res* 2009, **19:**1655-1664.
25.   Genomes Project C, Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, Korbel
JO, Marchini JL, McCarthy S, McVean GA, Abecasis GR: **A global reference for**
human genetic variation. *Nature* 2015, **526:**68-74.
26.   Renaud G, Slon V, Duggan AT, Kelso J: **Schmutzi: estimation of contamination and**
endogenous mitochondrial consensus calling for ancient DNA. *Genome Biol* 2015,
16:224.
27.   Lazaridis I, Patterson N, Mittnik A, Renaud G, Mallick S, Kirsanow K, Sudmant PH,
Schraiber JG, Castellano S, Lipson M, et al: **Ancient human genomes suggest three**
ancestral populations for present-day Europeans. *Nature* 2014, **513:**409-413.
28.   Lazaridis I, Nadel D, Rollefson G, Merrett DC, Rohland N, Mallick S, Fernandes D,
Novak M, Gamarra B, Sirak K, et al: **Genomic insights into the origin of farming in**
the ancient Near East. *Nature* 2016, **536:**419-424.
29.   Lazaridis I, Mittnik A, Patterson N, Mallick S, Rohland N, Pfrengle S, Furtwangler A,
Peltzer A, Posth C, Vasilakis A, et al: **Genetic origins of the Minoans and**
Mycenaeans. *Nature* 2017, **548:**214-218.
30.   Lipson M, Szecsenyi-Nagy A, Mallick S, Posa A, Stegmar B, Keerl V, Rohland N,
Stewardson K, Ferry M, Michel M, et al: **Parallel palaeogenomic transects reveal**
complex genetic history of early European farmers. *Nature* 2017, **551:**368-372.
31.   Allentoft ME, Sikora M, Sjogren KG, Rasmussen S, Rasmussen M, Stenderup J,
Damgaard PB, Schroeder H, Ahlstrom T, Vinner L, et al: **Population genomics of**
Bronze Age Eurasia. *Nature* 2015, **522:**167-172.
32.   Keller A, Graefen A, Ball M, Matzas M, Boisguerin V, Maixner F, Leidinger P, Backes C,
Khairat R, Forster M, et al: **New insights into the Tyrolean Iceman's origin and**
phenotype as inferred by whole-genome sequencing. *Nat Commun* 2012, **3:**698.

33.   Olalde I, Brace S, Allentoft ME, Armit I, Kristiansen K, Booth T, Rohland N, Mallick S,
940   Szécsényi-Nagy A, Mittnik A: **The Beaker phenomenon and the genomic**
941   **transformation of northwest Europe.** *Nature* 2018, **555:**190.

942   34.   Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P,
943   de Bakker PI, Daly MJ, Sham PC: **PLINK: a tool set for whole-genome association**
944   **and population-based linkage analyses.** *Am J Hum Genet* 2007, **81:**559-575.

945   35.   Vohr SH, Najar CFBA, Shapiro B, Green RE: **A method for positive forensic**
946   **identification of samples from extremely low-coverage sequence data.** *BMC*
947   *genomics* 2015, **16:**1034.

948   36.   Mathieson I, Alpaslan-Roodenberg S, Posth C, Szécsényi-Nagy A, Rohland N, Mallick
949   S, Olalde I, Broomandkhoshbacht N, Candilio F, Cheronet O: **The genomic history of**
950   **southeastern Europe.** *Nature* 2018, **555:**197.

951   37.   de Barros Damgaard P, Marchi N, Rasmussen S, Peyrot M, Renaud G, Korneliussen T,
952   Moreno-Mayar JV, Pedersen MW, Goldberg A, Usmanova E: **137 ancient human**
953   **genomes from across the Eurasian steppes.** *Nature* 2018, **557:**369.

954   38.   Durvasula A, Hoffman PJ, Kent TV, Liu C, Kono TJ, Morrell PL, Ross-Ibarra J: **angsd-**
955   **wrapper: utilities for analysing next-generation sequencing data.** *Mol Ecol Resour*
956   2016, **16:**1449-1454.

957   39.   Fu Q, Li H, Moorjani P, Jay F, Slepchenko SM, Bondarev AA, Johnson PL, Aximu-Petri
958   A, Prufer K, de Filippo C, et al: **Genome sequence of a 45,000-year-old modern**
959   **human from western Siberia.** *Nature* 2014, **514:**445-449.
960

Supplementary Figures

A)   **B)**

C)

Supplementary Figure 1. Distribution of LOD scores in simulated data. The distribution of LOD scores is depicted for samples with **A)** 0%, **B)** 7%, and **C)** 15% simulated contamination. These data were generated as part of tests using 1000 Genomes CEU individuals as the sample and contaminant DNA and for the haplotype panel.

Supplementary Figure 2. *ContamLD* **estimates when the individual, contaminant, and haplotype**

panel are all from CEU. *ContamLD* was run with no correction. The black dotted line is y=x, which would

correspond to a perfect estimation of the contamination. Error bars are 1.96*standard error (95%

confidence interval).

Supplementary Figure 3. *ContamLD* **estimates for Upper Paleolithic European individual after**

damage restricted correction (option 1). Kostenki14 (2.81x coverage) was contaminated with CEU and

analyzed using a CEU panel with *ContamLD* using damage correction and *ANGSD* [16] (Method 1). The

black dotted line is y=x, which would correspond to a perfect estimation of the contamination. Error

shading is 1.96*standard error (95% confidence interval).

Supplementary Figure 4. *ContamLD* estimates where the sample, contaminant, and haplotype

panels have varying ancestries. *ContamLD* was run with damage restricted correction (option 1). An

ancient Iberian of 1.02x coverage (I3756) was analyzed after contamination with **A)** CHB or **B)** YRI. An

ancient East Asian of 1.10x coverage (DA362.SG) is analyzed after contamination with **C)** CEU, **D)** CHB,

996   or **E)** YRI. An ancient South African of 1.21x coverage (I9028.SG) is analyzed after contamination with **F)**

997   CEU, **G)** CHB, or **H)** YRI. The black dotted line is y=x, which would correspond to a perfect estimation of

998   the contamination.

1000

1001

1002

1003

1004

1005   **A)**   **B)**

1006

Supplementary Figure 5. *ContamLD* estimates with CEU as the sample and multiple CEU

individuals as contaminants. *ContamLD* was run with CEU haplotype panels and damage restricted

correction (option 1). A CEU individual of 1.02x coverage (from the sequence distribution of the ancient

Iberian above) is contaminated with **A)** two CEU individuals or **B)** three CEU individuals. The black dotted

line is y=x, which would correspond to a perfect estimation of the contamination. Error bars are

1.96*standard error (95% confidence interval).

1013

1014

1015

Supplementary Figure 6. *ContamLD* **estimates with an ASW (African-American) individual and YRI**

panel. *ContamLD* was run with the 1240K SNP set and damage restricted correction (option 1). The

black dotted line is y=x, which would correspond to a perfect estimate of the contamination. Error bars are

1.96*standard error (95% confidence interval).

1020

Supplementary Figure 7. Contamination estimate standard errors of shotgun sequenced ancient

individuals comparing the 1240K and shotgun panels. Ancient shotgun sequenced individuals of 0.1-

0.5x coverage from Allentoft *et al*., 2015 [31], Damgaard *et al.*, Nature 2018 [37], and Damgaard *et al*.,

Science 2018 [21] were analyzed with *ContamLD* damage restricted correction (option 1) using the

1240K SNP set and a shotgun panel created using all variants above 10% frequency in the 1000

Genomes dataset. This test shows that analyses with the shotgun panel generally have smaller error bars

relative to those done with the 1240K panel, though it is unclear why there are two outliers with high

standard errors on the shotgun panel and low standard errors on the 1240K panel. All estimates are in

Supplementary Online Table 1.

1030

1031

1032

Supplementary Figure 8. *ContamLD* **standard errors from an ancient individual contaminated with**

another individual at different damage rates. The target individual was a 0.5x coverage ancient West

Eurasian-related individual (DA57.SG), and the contaminating individual was an ancient Iberian (I10895).

ContamLD was run with the 1240K SNP set with the CEU panel. The damaged sequences were

simulated as 0.005, 0.01, 0.02, 0.03, 0.04, 0.05, and 0.075.

1039

Supplementary Figure 9. *ContamLD* **estimates with an ancient individual contaminated with**

another ancient individual including its damaged sequences. The target individual was an ancient

Iberian (I3756) and the contaminating individual was another ancient Iberian (I10895). *ContamLD* was run

with the IBS panel and 1240K SNP set using damage restricted correction (option 1). The damaged

sequences were simulated as a 5% down-sampling of each respective contaminated BAM file. IBS are

1000 Genomes Project present-day Iberians from Spain. The black dotted line is y=x, which would

correspond to a perfect estimation of the contamination. Error bars are 1.96*standard error (95%

confidence interval). Points in red are those flagged with "Very_High_Contamination" by the software.

See Supplementary Online Table 4 for all values.

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1055 **Supplementary Figure 10.** Schematic of *ContamLD* algorithm (see text for additional details).