Prochlorococcus phage ferredoxin: structural characterization and interactions with cyanobacterial sulfite reductases

Ian J. Campbell[1,2], Jose L. Olmos, Jr. [1,2], Weijun Xu[2], Dimithree Kahanda[2], Joshua T. Atkinson[2], Othneil N. Sparks[2], Mitchell D. Miller[2], George N. Phillips,[2,3] George N. Bennett,[2,4] and Jonathan J. Silberg[2,4,5]*

From the [1]Biochemistry and Cell Biology Graduate Program, Rice University, 6100 Main Street, MS-140, Houston, Texas 77005; [2]Department of Biosciences, Rice University, 6100 Main Street, Houston, MS-140, TX, 77005; [3]Department of Chemistry, Rice University, 6100 Main Street, Houston, MS-140, TX, 77005; [4]Department of Chemical and Biomolecular Engineering, Rice University, 6100 Main Street, MS-362, Houston, TX, 77005; [5]Department of Bioengineering, Rice University, 6100 Main Street, MS-142, Houston, TX, 77005

Running title: *Phage Fd characterization and host SIR interactions*

*To whom correspondence should be addressed: Jonathan J. Silberg: Biosciences Department, Rice University, Houston TX 77005; joff@rice.edu; Tel. (713) 348-3849

Keywords: cyanobacteria, electron transfer, ferredoxin, marine, phage, sulfite reductase, virus

ABSTRACT

Marine cyanobacteria are infected by phage whose genomes encode ferredoxin (Fd) electron carriers. While these Fds are thought to redirect the energy harvested from light to phage-encoded oxidoreductases that enhance viral fitness, it is not clear how the biochemical and biophysical properties of phage Fds relate to those in photosynthetic organisms. Bioinformatic analysis using a sequence similarity network revealed that phage Fds display the highest similarity with cyanobacterial Fds that transfer electrons from photosystems to oxidoreductases involved in nutrient assimilation. Structural analysis of myovirus P-SSM2 Fd (pssm2-Fd), which infects *Prochlorrococcus marinus*, revealed high similarity to cyanobacterial Fds ($\leq$0.5Å RMSD). Additionally, pssm2-Fd exhibits a low midpoint reduction potential (-336 mV vs. SHE) similar to other photosynthetic Fds, albeit lower thermostability (T_m = 29°C) than many Fds. When expressed in an *Escherichia coli* strain with a sulfite assimilation defect, pssm2-Fd complemented growth when coexpressed with a *Prochlorrococcus marinus* sulfite reductase, revealing that pssm2-Fd can transfer electrons to a host sulfite reductase. The high structural similarity with cyanobacterial Fds and cross reactivity with sulfite reductase suggests that phage Fds evolved to transfer electrons to both phage- and cyanobacterial-encoded oxidoreductases.

Prochloroccus marinus is thought to be the most prevalent photosynthetic organism on Earth, with an estimated global abundance of 10^{27} cells (1), making it a key player in oceanic biogeochemical processes. This oligotrophic cyanobacterium inhabits the euphotic zone of oceans (40°N to 40°S) and is projected to expand both in density and range as global temperatures rise (1). Collectively fixing four gigatons of carbon annually (1), the diverse ecotypes of *P. marinus* thrive at many depths, where light, nutrient availability, and temperature vary (2–4). Up to 24% of the CO_2 fixed by *P. marinus* is released as dissolved organic carbon, providing critical feedstocks for heterotrophic organisms and supporting the larger oceanic ecosystem (5).

The core metabolic systems in *P. marinus* have diverged significantly from other cyanobacteria due in part to genome streamlining (6–10). Additionally, *P. marinus* lacks

phycobillisomes widely used by other cyanobacteria for light harvesting and instead uses a divinyl chlorophyll a_2/b_2 complex (11). Furthermore, this cyanobacterium contains only one set of core photosystem components while other cyanobacteria maintain several homologs (8, 12). Nitrogen assimilation pathways have been pruned as well, with species often lacking the ability to reduce either nitrate or nitrite (13, 14), and various ecotypes are incapable of heterotrophic growth, lacking several tricarboxylic acid cycle genes (8). Regardless of gene count, however, *P. marinus* ecotypes undergo a complex life cycle, facilitated in part by the role viruses play in their metabolism.

Cyanophages that infect *P. marinus* have evolved genomes with as many as 327 open reading frames (15). Viruses from three clades infect *P. marinus*, including T4-like myoviruses, T7-like podoviruses, and, less commonly, members of *Siphoviridae* (16). In some ecosystems, as many as 50% of cyanobacteria may be infected with phage (17). T4-like myoviruses and T7-like podoviruses exhibit infection cycles that are closely tied to day-night rhythms of cyanobacteria, although individual viruses vary in their specific responses (18). Once infection begins, expression of many host genes is repressed in favor of phage homologs (19), which can lead to a decrease in cyanobacterial carbon fixation, potentially keeping 5.39 gigatons of carbon in the atmosphere (20). While recent estimates suggest that cyanophages are limited by energy rather than nutrients (21), our understanding of their controls over *P. marinus* electron transfer (ET) remain limited.

Following infection, cyanophages express auxiliary metabolic genes to modulate photosynthesis (22). Phage express their own reaction core (D1 protein) of photosystem II during infection (17, 23, 24), with viral transcripts equaling or even exceeding bacterial transcripts in some environments. Certain cyanophages encode the entire photosystem I (PSI) that are thought to promiscuously reduce cytochromes, rather than plastocyanin like the host PSI, which may push the cells toward cyclic photosynthesis (25). Modified versions of viral plastocyanin, which possess altered charged surfaces compared with host plastocyanin, are also expressed in cyanophages and are hypothesized to transfer electrons directly to cytochrome aa₃ oxidase rather than PSI in the case of a heavily reduced quinone pool (26). Other phage genes implicated in oxygenic photosynthesis include *hli* (high-light inducible protein), *pebA* (15,16-dihydrobiliverdin:ferredoxin oxidoreductase), *psbD* (phosystem II D2 protein), *pcyA* (phycocyanobilin:ferredoxin oxidoreductase) and *petF* (ferredoxin) (15). Although the sequence similarities of these genes to their cyanobacterial homologs give clues to their function, we do not know how these proteins interact with host proteins to tune the host electron fluxome.

The *Prochlorococcus* P-SSM2 phage ferredoxin (pssm2-Fd) supports the reduction of phage phycoerythrobilin synthase (PebS), which catalyzes the four electron reduction of biliverdin IX_α to *3E/3Z* phycoerythrobilin (27). While previous studies have shown that some Fds can support ET to diverse partner oxidoreductases (28–31), it remains unclear how the biophysical properties of this and other phage Fds compare to cyanobacterial Fds, which function as central ET hubs that couple light harvesting to reductive metabolic reactions (30–32). In addition, it is not known how the specific constraints of viral evolution might have modulated the primary structure of phage Fd, which was likely acquired from its cyanobacterial host (22).

In this study, we used a sequence similarity network (SSN) to examine how cyanophage and cyanobacterial Fds relate to one another. In addition, we report the first structure of a cyanophage Fd, which has high structural similarity to cyanobacterial Fds that transfer electrons from photosystems to proteins involved in nutrient assimilation. We show that this Fd has a low midpoint potential characteristic of photosynthetic Fds, establish that this protein has an unusually low thermostability (T_m = 29°C), and demonstrate that it can support ET to a cyanobacterial oxidoreductase involved in sulfur assimilation. These results suggest that viral Fds support ET to both phage- and cyanobacterial-encoded oxidoreductases, and they extend our understanding about the ways viruses might alter the sulfur cycling mediated by cyanobacteria in the oceans.

Results
Cyanophage Fds share evolutionary links with cyanobacterial Fds

To better understand how cyanophage and

cyanobacterial Fd sequences relate to one another, we searched all 8392 viral genomes available in the Integrated Microbial Genomes and Microbiomes database for ORFs matching the Interpro signature for [2Fe-2S] Fds, IPR001041 (33). This genome mining yielded twenty six Fds from phages that have been shown to infect *Prochlorococcus* and *Synechococcus* (including pssm2-Fd), a Fd from an uncultured marine virus, and a Fd from a giant virus (*Pseudomonas* phage OBP) whose genome was recently reported (34). All viral ORFs encoded Fds with 95 to 97 amino acids, with the exception of the *Pseudomonas* phage ORF which was 83 residues. We compared the primary structures of the viral Fds with 807 cyanobacterial Fds obtained from a recent genome mining effort (35). *Zea mays* Fd (zm-Fd1) was included as a frame of reference because this protein has been intensively studied (36, 37). When the Fds were compiled into a SSN (38) and a stringent sequence similarity score (a cutoff of 30) was applied, the cyanobacterial Fds presented over a dozen distinct clusters, organized by average Fd length within each cluster (Figure 1A). Across the twelve largest clusters, the average Fd length varies by ~100 amino acids. These clusters also vary in the number of cyanobacterial species represented and the average size of the Fds, with the top five most species diverse clusters being VI (100 ±8 residues) > IX (115 ±10 residues) > VII (106 ±13 residues) > I (79 ±2 residues) > XI (163 ±8 residues).

Viral Fds are all localized to Cluster VI in close association with cyanobacterial Fds (Figure 1B), with the exception of *Pseudomonas* phage OBP Fd, which does not cluster with any other Fds under these conditions. Cluster VI includes numerous Fds with reported structures, including cyanobacterial Fds from *Synechocystis sp.* PCC 6803 (s6803-Fd1), *Nostoc sp.* PCC 7119 (nsp-Fd1), *Thermosynechococcus elongatus* (te-Fd1), and *Mastigocladus laminosus* (ml-Fd1). Additionally, this cluster contains zm-Fd1.

Among the Fds in Cluster VI, interactions with a range of partner oxidoreductases have been documented. s6803-Fd1 binds PSI, glutamate synthase (GS), and Fd-thioredoxin reductase (FTR) (39–42). nsp-Fd1 interacts with FNR and nitrate reductase (NIR) (43, 44). te-Fd1 interacts with FNR, PSI, heme oxygenase (HO), PcyA, and, 4-Hydroxy-3-methylbut-2-enyl diphosphate synthase (GcpE) (45–47). This latter oxidoreductase is in the non-mevalonate pathway for isoprenoid

biosynthesis (47). The tight association of cyanophage and cyanobacterial Fds having a wide range of protein-protein interactions suggest cyanophage Fds might be able to support ET to host oxidoreductases. In the case of pssm2-Fd, which is encoded by a myophage that infects *P. marinus*, it has the potential to interact with a total of eight host oxidoreductases (Figure 1C), in addition to the virally-encoded PebS (15, 48).

pssm2-Fd contains many sequence features of host cyanobacterial Fds

Among the phage Fds, pssm2-Fd has been the most intensively studied biochemically (27). To better understand how pssm2-Fd structure relates to cyanobacterial Fds, we analyzed the pairwise sequence identity of pssm2-Fd and cyanobacterial Fds in *P. marinus* hosts that are infected by cyanophage P-SSM2 (48). Fds from the host strains (ecotypes NATL1A, NATL2A, and MIT9211) were collected by protein BLAST (49) and designated pmN1-Fd (NATL1A), pmN2-Fd (NATL2A), and pm9211-Fd (MIT9211). Like pssm2-Fd, these Fds reside within Cluster VI. We also evaluated the pairwise identity of pssm2-Fd with five Fds whose structures have been reported, including cyanobacterial (s6803-Fd1, nsp-Fd1, te-Fd1, and ml-Fd1) and plant (zm-Fd1) Fds. This analysis revealed most (5/7) cyanobacterial Fds cluster tightly and exhibit ≥70% identity (Figure 2A), while pssm2-Fd exhibits a range of sequence identities (40-60%) with the cyanobacterial Fds. Interestingly, pssm2-Fd presents higher sequence identities with Fds from non-host cyanobacteria (52-59%) compared to host cyanobacteria (43-52%). The Fds from *P. marinus* ecotypes NATL1A and NATL2A (pmN1-Fd1 and pmN2-Fd1) exhibit the lowest sequence identity with pssm2-Fd of any Fd pairs tested (43%), and appear to be the two most divergent Fds in the set. pm9211-Fd1 exhibits higher sequence identity with pssm2-Fd (52%) and appears more closely related to other cyanobacterial Fds (s6803-Fd1, te-Fd1, nspFd1, and ml-Fd1) than the other *P. marinus* ecotypes. zm-Fd1 shows low identity with pssm2-Fd, pmN1-Fd1, and pmN2-Fd2 (48-50%), and is more similar to the cyanobacterial Fds including pm9211-Fd1 (65-72%).

Fd binding specificities are influenced by charged surface residues and residues that make direct contacts with partner proteins (36, 37). To establish whether pssm2-Fd sequence is conserved

at positions implicated in forming residue-residue contacts with partner proteins, a multiple sequence alignment was generated (Figure 2B). Residues implicated in mediating the interactions with Fd partners found in the host *P. marinus* are highlighted, which were identified using crystal structures of Fd-partner complexes (36, 37, 45). In the case of the Fd-FNR complex (PDB = 1GAQ), all of the Fd residues that make contacts with FNR are completely conserved (37), suggesting a ubiquitous binding strategy to FNR across cyanobacteria (37). For the Fd-PSI complex (PDB = 5ZF0) and Fd-SIR (PDB = 5H8V), greater variability is observed in the Fd residues that mediate partner contacts (36, 45). Eight of the eleven Fd residues that contact PSI are either absolutely conserved or have residues with similar physicochemical properties. Similarly, four of the six Fd residues that contact SIR are either absolutely conserved or similar. Among the *P. marinus* Fds, two (pmN1-Fd1 and pmN2-Fd1) have a pair of residues that have diverged from the Fds that mediate contacts in the Fd-SIR and Fd-PSI structures, while one (pm9211-Fd1) does not vary in either case. For both PSI- and SIR-contacting residues, pssm2-Fd varies from the partner-contacting residue at one contacting position observed in each crystal structure.

pssm2-Fd binds a low-potential iron sulfur cluster and is thermally unstable

The tight association of pssm2-Fd with cyanobacterial Fds in Cluster VI suggested that these proteins may exhibit similar biophysical properties. To obtain recombinant pssm2-Fd for *in vitro* characterization, the gene encoding pssm2-Fd was cloned into a pET expression vector, overexpressed in *Escherichia coli* at 37°C, and purified using a combination of anion exchange and size exclusion chromatography. This protocol yielded 0.6 mg/L of phage Fd, which was estimated to be >95% homogeneous by SDS-PAGE analysis. This yield is ~25-fold lower than that obtained with a thermophilic cyanobacterial Fd (15 mg/L) using an identical expression protocol (50).

To determine if pssm2-Fd contains a [2Fe-2S] cluster like photosynthetic Fds, we acquired pssm2-Fd absorbance and circular dichroism (CD) spectra (Figures 3A, 3B). The absorption spectrum contains peaks (465, 420, and 330 nm) that are characteristic of Fds coordinating a [2Fe-2S] cluster

(51, 52). Additionally, the CD spectrum presented ellipticity maxima (427 and 360 nm) and minima (550 and 505 nm) that are characteristic of holoFds (51, 52). The ratio of the ellipticity at 427 nm to the absorbance at 280 was comparable to a thermophilic cyanobacterial Fd that was recently reported (50), suggesting that a large fraction of pssm2-Fd contains a [2Fe-2S] cluster.

Because the yield of pssm2-Fd was lower than a thermophilic Fd that we recently purified (50), we postulated that pssm2-Fd might have a lower melting temperature. To test this idea, we evaluated how the ellipticity of pssm2-Fd at 427 nm changes as a function of temperature (Figure 3C). Assuming a two-state unfolding mechanism (53), the fraction folded of pssm2-Fd at each temperature was analyzed in a Van't Hoff plot of $\ln(K_{eq})$ vs $1/T$, yielding the ΔH and ΔS of the unfolding reaction as the slope and intercept of the linear regression, respectively (Supplementary Figure 1). The ratio of these two properties revealed a melting temperature (T_M) of 29 °C.

pssm2-Fd is thought to serve as an electron acceptor for PSI, but it is not clear how the midpoint potential of this protein relates to the photosynthetic Fds with a similar cellular role. To address this question, we characterized the electrochemical properties of pssm2-Fd using thin film, square wave voltammetry (54). With this analysis, pssm2-Fd presented a low midpoint potential (E_m = -336 mV) relative to a standard hydrogen electrode (Figure 3D).

The pssm2-Fd structure is highly similar to cyanobacterial homologs

To investigate structural features of pssm2-Fd that underlie its low midpoint potential and low thermostability, we determined the pssm2-Fd crystal structure at 1.6 Å resolution; the R_{cryst}/R_{free} are 0.1783/0.2134. The resulting model contains all 96 residues for each of the two monomers in the asymmetric unit, excluding the initiator methionines. Mass spectrometry data corroborates the absence of the initiator methionine. The two chains of the asymmetric crystal align well, with a root-mean-square deviation of all atomic coordinates (n_A= 1392, n_B=1431) of 0.17 Å. The main source of variation between the chains arises from subtle differences within the beta sheets. While chain A in the asymmetric unit is well ordered, chain B lacks density for a variety of

residues around the iron-sulfur cluster with a poor fit to the electron density resulting in a large real-space R-value. Therefore, NCS restraints refinements were carried out using chain A as a reference for chain B. The apo-ferredoxin is unstable and evidence in the literature indicates that zinc can interfere with and disrupt iron-sulfur clusters (55–58). In chain B, the iron-sulfur cluster site was disrupted. The density surrounding the cluster in chain B is consistent with a mixture of zinc bound and an intact 2Fe-2S cluster. To account for the poor density and the anomalous data, chain B has been modeled as a grouped occupancy, where the model is partially comprised of intact cysteine-coordinated iron-sulfur cluster, and a zinc ion with a hydration shell. An interpretation of these data is that as the cluster is disrupted, the surrounding residues become disordered. The crystallographic data collection and refinement statistics are provided in Supplementary Table 2.

The structure of the pssm2-Fd Fd is shown in Figure 4A. With five β-sheet strands and three α-helices, pssm2-Fd possesses a canonical β-grasp topology, which consists of a β-sheet that is four to five strands long with α-helices interspersed and packed against the sheet (59). This topology is observed in other Fds, such as ml-Fd1 and s6803-Fd1 (40, 60). To quantify how similar the pssm2-Fd structure is to other cyanobacterial Fds in Cluster VI (1OFF, 5AUI, 1CZP, 1RFK, 3B2F), we calculated the pairwise RMSD between these structures (40, 60–63). The pssm2-Fd structure presents low RMSD ($\leq$0.5 Å) in all atom positions with all of the cyanobacterial Fds (Supplementary Table 3), including two thermophilic Fds (45, 60) and a plant Fd (zm-Fd1).

We next mapped the sequence variation of structurally-characterized Fds within Cluster VI onto the pssm2-Fd structure (Figure 4B). To quantify variation, we calculated the conservation scores at each pssm2-Fd residue using *ConSurf* (64). This analysis revealed that the largest density of absolutely conserved residues occurs in the loop surrounding the [2Fe-2S] cluster, with the next most conserved regions being those adjacent to this loop. There are additional residues that are highly conserved across all of the Fds analyzed. However, these are dispersed across the structure and primarily localized to regions that are packed within the center of the pssm2-Fd structure or forming salt-bridges on the surface.

Analysis of the intramolecular interactions within pssm2-Fd and those observed within cyanobacterial Fds found in Cluster VI revealed additional differences. Among the proteins analyzed, pssm2-Fd presents a lower number of salt bridges (n = 3) compared with all other Fds analyzed (n = 4 to 6). However, pssm2-Fd contains a similar number of total hydrogen bonds (n = 133) as the other Fds (n = 105 to 154). Only one salt bridge was conserved in all of the Fds, which arises from an interaction between an arginine at position 41 or 42 and either a glutamate or aspartate situated between positions 27 and 30. This interaction may be conserved because it plays a role in stabilizing [2Fe-2S] cluster coordination, as the arginine is part of the cluster binding loop (Figure 2B).

pssm-Fd conducts in vivo electron transfer with host partners

The high structural similarity between pssm2-Fd and Fds that couple with a wide range of partner oxidoreductases suggested that pssm2-Fd might be able to transfer electrons from PSI to oxidoreductases encoded by cyanobacterial hosts. In a previous study, we examined whether pssm2-Fd could support ET between Zea mays FNR and SIR using a synthetic ET pathway in *E. coli* that is required for cellular growth (50). In this cellular assay, Fd-mediated ET from zm-FNR to zm-SIR is required for sulfur assimilation and growth on medium containing sulfate as the only sulfur source (65). Initial studies did not observe ET from zm-FNR to zm-SIR (50). However, these experiments were performed at a temperature (37°C) where only a small fraction of pssm2-Fd is expected to be folded.

To increase the fraction of holo-pssm2-Fd in the cellular assay, we built an aTc-regulated expression vector that initiates translation of pssm2-Fd using a stronger RBS than previously utilized. When the cellular assay was performed using this pssm2-Fd expression construct, aTc-dependent complementation was observed at 37°C (Figure 5). This finding indicates that pssm2-Fd supports ET from zm-FNR to zm-SIR, but requires increased protein expression to overcome the relative instability of pssm2-Fd.

To test whether pssm2-Fd can also support ET to host SIRs, we created vectors that constitutively express *P. marinus* NATL2A SIR (pmN2-SIR) and MIT9211 SIR (pm9211-SIR) in

parallel with zm-FNR. SIR from the NATL1A host was omitted because it shares high sequence identity (98%) with pmN2-SIR (Supplementary Figure 2). The charge conservation of five of the seven zmFd1-contacting residues in the *P. marinus* homologs suggested the cyanobacterial SIRs would function similarly to zm-SIR (Supplementary Figure 3) We also built aTc-inducible vector systems to express host Fds (pmN1-Fd1, pmN2-Fd1 and pm9211-Fd1) as a frame of reference. Using these vectors, we tested the ability of cyanophage and host Fds to support ET from zm-FNR to three different SIRs (zm-SIR, pmN2-SIR, zm-9211-SIR). Because ml-Fd1 supports ET from zm-FNR to zm-SIR (50), it was included in the analysis. Assays were performed across a range of temperatures (23 to 37°C) in the presence and absence of 100 ng/mL aTc.

At the highest temperature assayed (37°C), only cells expressing zm-SIR complement cellular growth when coexpressed with a Fd and FNR (Figure 6A). This growth was observed with three different Fds, including ml-Fd1, pssm2-Fd, and pm9211-Fd1. Cells expressing pmN2-Fd did not support ET from zm-FNR to zm-SIR. At 30°C, complementation was again observed with the same three-component ET pathways (Figure 6B). Additionally, cells expressing pm9211-SIR complemented growth when coexpressing the zm-FNR and either the cognate host Fd, pssm2-Fd, or ml-Fd1. At 23°C, similar trends were observed with zm-SIR and pm9211-SIR as at 30°C (Figure 6C). Additionally, significant complementation, albeit small, was observed when pmN2-SIR was coexpressed with the pssm2-Fd and ml-Fd1.

To investigate if differences in Fd-dependent growth arise because each Fd is expressed at different levels, we created expression vectors that produce these proteins as RFP fusions and performed whole cell fluorescence measurements with and without 100 ng/mL aTc, and compared them to an unfused ml-Fd1 (Figure 7). This analysis revealed that pmN2-Fd is expressed at higher levels than the other three Fds under identical expression conditions as the cellular assay. This finding suggests that the weaker complementation observed with pmN2-Fd does not arise because it is expressed at lower levels than the Fds that complement growth.

Discussion

Our bioinformatic analysis revealed 12 major clusters of cyanobacterial Fds, with many clusters having a narrow Fd length distribution. This analysis also revealed that some clusters represent Fds which are widespread across cyanobacteria, while others contain Fds from a small number of organisms, illustrating how Fd gene duplications have diversified controls over ET in specific cyanobacterial genera. The extent to which the SSN clusters represent Fds with distinct ET roles is not known. In the case of the cyanobacterial Fds found in Cluster VI, which includes all cyanophage Fds analyzed, several of these proteins are the Fd predominately expressed under phototrophic conditions (30, 66, 67). In some cases, this expression is likely due to their critical role as the electron acceptor for PSI (41, 45). Since Cluster VI contains cyanobacterial Fds that have evolved to support ET to a wide range of oxidoreductases (FNR, GcpE, SIR, NIR, GS, HO, and PycA), it seems probable that cyanophage Fds arose from the duplication of a cyanobacterial Fd that mediates ET to one or more of these partner oxidoreductases.

The cellular roles of the cyanobacterial Fds found in each SSN cluster are not well established, although some clues can be gathered from studies in the model cyanobacterium *Synechocystis* sp. PCC6803. This organism contains a Fd (s6803-Fd1), within Cluster VI and also has five more [2Fe-2S] Fd paralogs that map onto Clusters V, VI, IX and X (30). The dispersion of Fds from one organism into these different clusters with a wide range of primary structure lengths suggests that evolutionary pressures have diversified *Synechocystis* sp. PCC6803 Fds into multiple roles. The variable regulation of these Fds supports this notion (30, 66, 68). Aside from the s6803-Fd1, transcripts of the other five [2Fe-2S] Fds change markedly under various stresses, with differences observed in response to salt stress, high concentrations of metals, low light conditions, and blue light conditions. These results suggest that more remains to be learned about how cyanobacterial Fds control the proportion of electrons that flow between different partner oxidoreductases, particularly within pathways that remediate metals and oxidative species as well as pathways that modulate photosynthetic machinery in response to changes in light. Analyzing Fd orthologs from the same cluster may reveal larger

patterns in partner specificity.

Some additional unique features are observed in the SSN clusters. Clusters I and II consist of Fds that have a tight size distribution ($\mu = 79 \pm 2$, $\mu = 81 \pm 5$) and have primary structures that all terminate in tryptophan. This terminal residue is an unusual feature in comparison to canonical Fds (*e.g.*, Cluster VI) which generally end with a tyrosine. These "tryptophan-terminating" Fds are found in >40 species, although Cluster II consists wholly of the two related genera *Prochlorococcus* and *Synechococcus* (10). Cluster III is solely populated by *P. marinus* ecotypes. None of these ecotypes are known to be infected by P-SSM2 cyanophage. Cluster V consists completely of Fds from *Synechocystis* strains. Of the 87 species with Fds in Cluster IX, 96% have two Fds. The smaller set of Fds exhibit a tight size distribution ($\mu = 106 \pm 5$) as do the larger set ($\mu = 122 \pm 3$). Finally*,* Cluster X is populated primarily by Fds from *P. marinus* and *Synechococcus sp.* strains, although it also contains two species of *Cyanobium*. To better understand how sequence similarities relate to various roles Fds play within Cyanobacteria, functional studies of many cyanobacterial Fds with a suite of native partner oxidoreductases are required.

Focusing on Cluster VI, our analysis of the pairwise identities of well characterized Fd from this group shows that pssm2-Fd presents the lowest sequence identity with Fds found in host strains NATL1A and NATL2A. This is surprising, as previous phylogenetics has established that another P-SSM2 phage encoded gene, PSII reaction center D1, is more closely related to NATL1A and NATL2A homologs than viral homologs, suggesting direct transfer of D1 from host to phage (22). Taken together, these findings support a model in which pssm2-Fd is under more selective pressure than D1. Perhaps this disparity arises because the cyanophage P-SSM2 benefits from redistributing electrons in the infected host using mutated Fds. However, the photosynthetic machinery obtained from the host functions sufficiently for viral replication, so large mutations in D1 are not observed. The difference in sequence divergence may also be attributable to Fd being acquired virally at an earlier time and experiencing genetic drift for a longer period.

The cyanophage Fd characterized herein exhibits higher sequence identities with non-host cyanobacterial Fds than with host Fds. The underlying cause of this trend is not known. Biophysical studies revealed that pssm2-Fd exhibits a similar overall structure to non-host cyanobacterial Fds (40, 60–62), presenting ≤ 0.5Å RMSD with mesophilic (nsp-Fd1 and s6803-Fd1) and thermophilic (te-Fd1 and ml-Fd1) Fds. Additionally, pssm2-Fd exhibits a low midpoint potential that is identical to that of ml-Fd1 ($E° = -336$ mV) and higher than s6803-Fd1 ($E° = -412$ mV) and nsp-Fd1 ($E° = -384$ mV) (50, 69, 70). While structures and electrochemical properties of *P. marinus* Fds have not been established, they are expected to be similar to characterized orthologs in other cyanobacteria that function as electron acceptors for PSI.

One aspect of pssm2-Fd that is particularly noteworthy is the low melting temperature ($T_m = 29°C$). This T_m is lower than all of the characterized Fds in Cluster VI, which includes the thermophilic ml-Fd1 ($T_m = 76°C$) (71). This low thermostability may have been selected to support rapid turnover of cyanophage Fds or to regulate protein turnover in response to temperature changes, or perhaps in the low temperature waters (15-25°C) that P-SSM2 hosts colonize there is not strong selection for Fds to maintain thermostability (4). Currently, it is not known how Fds tune their relative stabilities without altering their low potentials. The native residues that control Fd stabilities ($T_m = 29$ to $76°C$) could be established in future studies by creating chimeras between pssm2-Fd and more thermostable Fds with similar midpoint potentials (*e.g.*, ml-Fd1) and then evaluating how inheritance of different blocks of sequence relate to the thermostability of the resulting chimeras. This approach has previously been used in parallel with structure-guided recombination to identify modules in proteins that can be exchanged to control thermostability (72, 73).

The growth selection used to monitor Fd-mediated ET illustrates how a cellular assay simplifies the analysis of Fd and partner oxidoreductase interactions (28). In previous studies, this cellular assay was used to evaluate the cross reactivity between Fds from a range of species and plant FNR and SIR (50, 65). In addition, this approach has been used to select for engineered Fds that support cellular ET from FNR to SIR (50, 74, 75). The measurements described herein illustrate how this cellular assay can be used to rapidly assess

whether phage-encoded Fds are capable of supporting ET to host-encoded oxidoreductases. The finding that pssm2-Fd supports ET to *P. marinus* SIRs indicates that this ET may occur during the cyanophage life cycle. However, the extent to which this reaction occurs in nature is not known, and it is unclear how phage Fds control the proportion of electrons transferred to host oxidoreductases like SIR vs phage-encoded oxidoreductases like PebS. One way to begin evaluating this competition in the future would be to examine how coexpression viral PebS and host SIR in the cellular assay described herein affects growth complementation.

How cyanophages influence both demand for and production of sulfur in their hosts will require further studies. Our finding that host SIRs can be reduced by virally-encoded Fds supports the idea that P-SSM2 has evolved to supply reducing power to host sulfide production machinery during infection. However, we still do not know how the temporal expression of phage Fds modulates the cyclical lifecycle of cyanobacteria. Recent studies have revealed that cyanophage gene expression and replication is diurnally linked to host replication, with both processes occurring during the afternoon-evening period when light intensity is high (18, 76, 77). Additionally, phage transcripts drop at night and when photosystem is compromised (18). Despite these advances, we don't know what feedback loops govern expression of phage genes in infected cells, especially photosyntheis-related genes derived from host homologs.

While the relationship of cyanophage infection cycles to the sulfur metabolism of widespread cyanobacteria like *Prochlorococcus* and *Synechococcus* is unclear, such interactions are likely critical to the health of oligotrophic oceanic gyres. Although sulfate is highly abundant in these ecosystems (up to 28 mM) (78), reduced sulfur species are more scarce, with concentrations of dissolved dimethylsulfoniopropionate (DMSP) at 0.5-10 nM (78–80) and methionine at 10-40 pM, during the day (80). Reduced sulfur in oceanic gyres appears to be in high demand, as DMSP is readily assimilated by gyre communities even when sulfate $>10^6$ -fold more abundant. (78). Further, DMSP uptake is repressed by nanomolar additions of more highly reduced compounds that are ostensibly easier to metabolize, such as methionine, cysteine, sulfide, and even dimethylsulfoxide (DMSO), indicating a hierarchy of preferred sulfur species among gyre plankton (78, 80). Additionally, some heterotrophs from the Sar11 clade, which constitutes 30% of sea surface bacteria (81), are unable to metabolize sulfate and rely on dissolved reduced sulfur sources (82), potentially those produced by phototrophs. Conversely, recent work has demonstrated that two ecotypes of *P. marinus* (MED4 and MIT9313) were unable to supply sufficient reduced sulfur to enable SAR11 growth in laboratory co-cultures (83). However, this study also demonstrated that only one of the same two strains were able to complement glycine auxotrophy in SAR11, suggesting that sulfide complementation may be observed in other *P. marinus* ecotypes or other species of cyanobacteria. The fact that environmental methionine levels peak during the day and drop below detection limits after sunset suggests that the diurnal cycle of cyanobacteria contributes to sulfur cycling in surface communities (80). Uncovering these relationships could be key to understanding nutrient availability across the Earth's oceans.

Experimental procedures
Materials

Tris Base was from Fisher Scientific, N-cyclohexyl-3-aminopropanesulfonic (CAPS) was from Acros Organics, 2-(N-Morpholino)ethanesulfonic acid (MES) and N-[Tris(hydroxymethyl)methyl]-3-aminopropanesulfonic acid (TAPS) were from Fluka Biochemika, and isopropyl β-D-1-thiogalactopyranoside (IPTG), dithiothreitol (DTT), kanamycin, chloramphenicol, and streptomycin were from Research Product International. 3-Morpholinopropane-1-sulfonic acid (MOPS), N-Cyclohexyl-2-aminoethanesulfonic acid (CHES), and all other chemicals were purchased from Sigma-Aldrich. *E. coli* EW11 was a gift from Pam Silver (Harvard University)(65), XL1-Blue *E. coli* competent cells were from Agilent, and Rosetta™(DE3) *E. coli* were from Novagen.

Vector Design

All plasmids are listed in Supplementary Table 1. Genes were synthesized by either Integrated DNA technologies or Twist Bioscience as G-blocks and Gene Fragments, respectively. Plasmids were constructed by ligating PCR

products amplified with Q5 polymerase (New England Biolabs) using Golden Gate DNA assembly (84). Ribosome binding sites were designed using the Salis Lab RBS calculator (85). All plasmids were sequence verified.

Bioinformatics

To harvest viral Fd open reading frames (ORFs), all available viral genomes from Integrated Microbial Genomes and Microbiomes (n=8392) were scanned for genes matching the [2Fe-2S] Fd Interpro sequence signature IPR001041 (33). Viral genes with this signature were compared with the 807 cyanobacterial [2Fe-2S] Fds recently characterized (35). All Fd sequences encoded by these ORFs were to analyzed using the Enzyme Function Initiative-Enzyme Similarity Tool to obtain pairwise sequence similarity scoring and generate a SSN (86). The resulting networks were analyzed and images were generated using Cytoscape with the prefuse force directed layout based on alignment score (87). Protein sequence alignments were generated using MUSCLE (88). Root-mean-squared-deviation calculations and structural images were performed using PyMOL (89). Conservation of Fd residues mapped onto the pssm2-Fd structure was generated using the ConSurf Server with the WAG evolution model, seeded with all Fd sequences from Cluster VI (64).

Protein purification

Rosetta *E. coli* cells transformed with a vector encoding pssm2-Fd under control of the lac promoter (pJTA022) were grown at 37˚C in lysogeny broth (LB) containing 50 µg/mL kanamycin to midlog phase, induced to express protein using 50 µM IPTG, and grown overnight at 37˚C while shaking at 250 rpm. Cells harvested by centrifugation (4,000 g) were resuspended in 25 mL lysis buffer (per L of culture), which contained 10 mM Tris pH 8, 5 mM DTT, 10 mg/L DNase I, and 0.5 mg/mL lysozyme. After freezing at −80˚C, cells were thawed and mixed with a cOmplete Mini, EDTA-Free protease inhibitor tablet (Sigma-Aldrich) at a ratio of one tablet per 100 mL lysate. Samples were kept on ice or at 4˚C for all subsequent purification steps. Clarified lysate harvested by centrifugation (47,000 g) was diluted three-fold with TED buffer (25 mM Tris pH 8, 1 mM EDTA, and 1 mM DTT) to lower the salt concentration. This mixture was loaded onto a DE52 anion exchange column (Whatman) that had been equilibrated with TED buffer, the column was washed with TED containing 200 mM NaCl, and the Fd was eluted using sequential isocratic washes with TED containing 250 and 300 mM NaCl. Fractions appearing brown were mixed and diluted with TED to bring NaCl below 100 mM. These pooled fractions were then loaded onto HiTrap Q XL column (GE Healthcare) equilibrated with TED using an AKTA Start FPLC system (GE Healthcare). The column was washed using 100 mL TED buffer. Protein was eluted using a linear gradient (0 to 375 mM NaCl in TED) followed by an isocratic wash (500 mM NaCl in TED). Brown fractions were pooled and then further purified using a HiLoad 16/600 Superdex 75 size exclusion column containing TED. SDS-PAGE was performed to analyze purity at each step using NuPage 12% Bis-Tris Gels (Invitrogen) and the PageRuler Unstained Broad Range Protein Ladder (Thermo Scientific). Samples appearing homogeneous were pooled and concentrated using an Amicon Ultra 10 K MWCO spin column and then flash frozen with liquid nitrogen.

Spectroscopy

Prior to all measurements, pssm2-Fd was dialyzed into TED. Absorbance spectra and ellipticity were acquired using a J-815 spectropolarimeter (Jasco, Inc) using quartz cuvettes with a 1 cm path length. Scans were conducted using a 1 nm bandwidth, a 0.5 nm data pitch, and a 200 nm/min scan rate at 20°C. To assess protein stability, a cuvette containing 50 µM Fd was heated from 5 to 50°C at a rate of 1°C/minute while monitoring ellipticity and absorbance.

Electrochemistry

Electrochemical measurements were performed anaerobically using a three-electrode system. A Ag/AgCl/1M KCl electrode (CH Instruments) was used as the reference electrode, and a platinum wire was used as the counter electrode. An edge-plane pyrolytic graphite electrode was used as the working electrode to perform protein film electrochemistry. Prior to adding Fd, this electrode was treated with 100 mM neomycin trisulfate (Sigma-Aldrich) to improve the electrochemical signal (90). An aliquot (3 µL) of Phage Fd (620 uM) was applied

directly to the electrode surface following neomycin treatment, and the protein was allowed to adhere to the surface for 1 min at 23°C. The electrode was then placed in a glass vial containing a pH 7 buffer solution (5 mM acetate, MES, MOPS, TAPS, CHES, CAPS) containing 100 mM NaCl at 23.5 °C. Square wave voltammograms were collected at 10 Hz frequency, and electrochemical signals were analyzed using QSoas open software. Similar results were obtained when experiments were performed using pssm2-Fd from different purifications. A CH Instruments potentiostat and CHI660E electrochemical analyzer were used for all measurements. All data is reported relative to Standard Hydrogen Electrode (SHE), taking into account the potential difference between SHE and Ag/AgCl/1M KCl, which is 0.222 V.

Crystallization

Purified Fd was concentrated to ~20 mg/mL for initial crystal screening using the mosquitoLCP pipetting robot (TTP Labtech, Boston, MA) and commercially available screens including Wizard Classic 1 and 2 and Wizard Classic 3 and 4 (Rigaku Reagents, Inc., Bainbridge Island, WA), MORPHEUS and MIDAS (Molecular Dimensions, Holland, OH), PEGRx and IndexHT (Hampton Research, Alisa Viejo, CA). Protein crystals were grown via the sitting drop method of vapor diffusion at a 1:1 (v:v) ratio of protein:precipitant 200:200 nL, at 20 ºC. Crystals for diffraction and data collection were optimized and grown with a precipitant solution containing 10% (w/v) PEG 3000, 200 mM zinc acetate, and 100 mM sodium acetate/acetic acid pH 4.5. The crystals were cryo-protected with 10 % glycerol, harvested using a MiTeGen micromesh loop (MiTeGen, Ithaca, NY), and flash-cooled in liquid nitrogen prior to data collection.

X-ray data collection, structure solution and refinement

Data were collected at the Advanced Photo Source GM/CA beamline 23ID-B using a 20 μm beam and 0.2 degree oscillations at a wavelength of 1.033 Å with an EIGER X 16M (DECTRIS Ltd, Philadelphia, PA) detector. The data were processed using the *autoPROC* toolbox (91), indexed and integrated using *XDS* (92), and scaled with *AIMLESS* (93). Initial phases were obtained by

molecular replacement with *Phaser-MR* using the *Zea mays* Fd PDB ID: 5H57 (94) as a search model. Model building and refinement were performed with *Coot* (95) and *phenix.refine* (96). A custom graphics system was used for collaborative visualization (97). NCS restraints were used in initial refinements, with chain A as a reference for chain B, and removed in the final rounds of refinement. For TLS refinement, groups were automatically determined for chain A while chain B was treated as one group. Anomalous difference maps were calculated using *ANODE* and zinc from the precipitant solution was modeled in the structure (98). Data processing and refinement software were compiled and supported by the SBGrid Consortium (99). The structural model is available in the worldwide Protein Data Bank (PDB ID: 6VJV) (100). Data collection and refinement statistics are provided in Supplementary Table 2.

Cellular assay

To assess Fd-mediated ET from Fd-NADP reductase (FNR) to sulfite reductase (SIR), *E. coli* EW11 was transformed with two plasmids using electroporation as previously described (50, 65, 74, 75)). One plasmid constitutively expressed the different FNR and SIR pairs, while the other expressed different Fds under control of an anhydrotetracycline (aTc)-inducible promoter. To maintain both plasmids, all growth steps included chloramphenicol (34 μg/mL) and streptomycin (100 μg/mL). Starter cultures inoculated using single colonies were grown in deep-well 96-well plates for 18 h at 37 °C in 1 mL of a nonselective modified M9 medium (M9c), which contained sodium phosphate heptahydrate, dibasic (6.8 g/L), potassium phosphate, monobasic (3 g/L), sodium chloride (0.5 g/L), 2% glucose, ammonium chloride (1 g/L), calcium chloride (0.1 mM), magnesium sulfate (10 mM), ferric citrate (24 mg/L), p-aminobenzoic acid (2 mg/L), inositol (20 mg/L), adenine (5 mg/L), uracil (20 mg/L), tryptophan (40 mg/L), tyrosine (1.2 mg/L), and the remaining 18 amino acids (80 mg/L each). Starter cultures grown to stationary phase were then diluted 1:100 into a selective modified M9 medium (M9sa), which is identical to M9c but lacks cysteine and methionine, containing varying amounts of aTc. Cultures were inoculated in sterile Nunc™ Edge 2.0 96-well plates with side reservoirs filled with 1 mL of water. The edge between plate tops and bottoms were

sealed with parafilm. Cells were then grown in a Spark plate reader (Tecan) at 37 °C while shaking (90 rpm) at an amplitude of 3 mm in double-orbital mode. Optical density (OD) was measured at 600 nm every 5 min for 48 h. For endpoint measurements, all conditions were the same except cells were grown in incubators at the indicated temperature while shaking at 250 rpm for 48 hours.

Fluorescence measurements

To evaluate the relative expression of Fds in the cellular assay, the genes encoding each Fd were placed under control of an aTc-inducible promoter and expressed as fusions to a monomeric red fluorescent protein (RFP) through a twelve amino acid linker (101). Each vector was transformed into *E. coli* EW11 cells, alongside a plasmid (pSAC01) constitutively expressing *Zea mays* SIR (zm-SIR) and *Zea mays* FNR (zm-FNR)(50). Transformed cells were grown under identical conditions to those described fror the cellular assay, with the exception of being grown in M9c at all steps. After 24 hours, endpoint OD and RFP fluorescence ($\lambda_{excitation}$ = 560 nm; $\lambda_{emission}$ = 650 nm) were measured.

Statistics

Error bars represent standard deviation calculated from three or more biological replicates. Independent two-tailed t-tests were applied to compare differences between all relevant samples with α=0.05.

Acknowledgements: This project was supported by DOE grant DE-SC0014462 (to J.J.S), NASA NAI grant number 80NSSC18M0093 (to J.J.S), and NSF grant 1231306 (to G.N.P). GM/CA@APS has been funded in whole or in part with Federal funds from the National Cancer Institute (ACB-12002) and the National Institute of General Medical Sciences (AGM-12006). The Eiger 16M detector at GM/CA-XSD was funded by NIH grant S10 OD012289 .This research used resources of the Advanced Photon Source, a U.S. Department of Energy (DOE) Office of Science User Facility operated for the DOE Office of Science by Argonne National Laboratory under Contract No. DE-AC02-06CH11357. Additionally, IJC and JTA were supported by a Lodieska Stockbridge Vaughn Fellowship.

Conflict of interest: The authors declare that they have no conflicts of interest with the contents of this article.

Author contributions: IJC, JTA, GNB, and JJS conceptualized the project. IJC, NS, and JTA purified recombinant protein. IJC conducted spectroscopy. JLO, WX, MDM, and GNP crystallized phage Fd and generated the structure. DK conducted electrochemistry. IJC constructed vectors and performed cellular assays. IJC and JJS wrote the manuscript, and all other authors refined the text.

References

1. Flombaum, P., Gallegos, J. L., Gordillo, R. A., Rincon, J., Zabala, L. L., Jiao, N., Karl, D. M., Li, W. K. W., Lomas, M. W., Veneziano, D., Vera, C. S., Vrugt, J. A., and Martiny, A. C. (2013) Present and future global distributions of the marine Cyanobacteria *Prochlorococcus* and *Synechococcus*. *Proc. Natl. Acad. Sci.* **110**, 9824–9829

2. Malmstrom, R. R., Coe, A., Kettler, G. C., Martiny, A. C., Frias-Lopez, J., Zinser, E. R., and Chisholm, S. W. (2010) Temporal dynamics of *Prochlorococcus* ecotypes in the Atlantic and Pacific oceans. *ISME J.* **4**, 1252–1264

3. Martiny, A. C., Tai, A. P. K., Veneziano, D., Primeau, F., and Chisholm, S. W. (2009) Taxonomic resolution, ecotypes and the biogeography of *Prochlorococcus*. *Environ. Microbiol.* **11**, 823–832

4. Zinser, E. R., Johnson, Z. I., Coe, A., Karaca, E., Veneziano, D., and Chisholm, S. W. (2007) Influence of light and temperature on *Prochlorococcus* ecotype distributions in the Atlantic Ocean. *Limnol. Oceanogr.* **52**, 2205–2220

5. Bertilsson, S., Berglund, O., Pullin, M., and Chisholm, S. (2005) Release of dissolved organic matter by *Prochlorococcus*. *Vie Milieu*. **55**, 225–231

6. Dufresne, A., Garczarek, L., and Partensky, F. (2005) Accelerated evolution associated with genome reduction in a free-living prokaryote. *Genome Biol.* **6**, R14

7. Kettler, G. C., Martiny, A. C., Huang, K., Zucker, J., Coleman, M. L., Rodrigue, S., Chen, F., Lapidus, A., Ferriera, S., Johnson, J., Steglich, C., Church, G. M., Richardson, P., and Chisholm, S. W. (2007) Patterns and Implications of Gene Gain and Loss in the Evolution of *Prochlorococcus*. *PLoS Genet.* **3**, e231

8. Rocap, G., Larimer, F. W., Lamerdin, J., Malfatti, S., Chain, P., Ahlgren, N. A., Arellano, A., Coleman, M., Hauser, L., Hess, W. R., Johnson, Z. I., Land, M., Lindell, D., Post, A. F., Regala, W., Shah, M., Shaw, S. L., Steglich, C., Sullivan, M. B., Ting, C. S., Tolonen, A., Webb, E. A., Zinser, E. R., and Chisholm, S. W. (2003) Genome divergence in two *Prochlorococcus* ecotypes reflects oceanic niche differentiation. *Nature*. **424**, 1042–1047

9. Sun, Z., and Blanchard, J. L. (2014) Strong Genome-Wide Selection Early in the Evolution of *Prochlorococcus* Resulted in a Reduced Genome through the Loss of a Large Number of Small Effect Genes. *PLoS ONE*. **9**, e88837

10. Urbach, E., Scanlan, D. J., Distel, D. L., Waterbury, J. B., and Chisholm, S. W. (1998) Rapid Diversification of Marine Picophytoplankton with Dissimilar Light-Harvesting Structures Inferred from Sequences of *Prochlorococcus* and *Synechococcus* (Cyanobacteria). *J. Mol. Evol.* **46**, 188–201

11. Chisholm, S. W., Frankel, S. L., Goericke, R., Olson, R. J., Palenik, B., Waterbury, J. B., West-Johnsrud, L., and Zettler, E. R. (1992) *Prochlorococcus marinus* nov. gen. nov. sp.: an oxyphototrophic marine prokaryote containing divinyl chlorophyll *a* and *b*. *Arch. Microbiol.* **157**, 297–300

12. Golden, S. S., Brusslan, J., and Haselkorn, R. (1986) Expression of a family of psbA genes encoding a photosystem II polypeptide in the cyanobacterium *Anacystis nidulans* R2. *EMBO J.* **5**, 2789–2798

13. Berube, P. M., Rasmussen, A., Braakman, R., Stepanauskas, R., and Chisholm, S. W. (2019) Emergence of trait variability through the lens of nitrogen assimilation in *Prochlorococcus*. *eLife*. **8**, e41043

14. Biller, S. J., Berube, P. M., Lindell, D., and Chisholm, S. W. (2015) *Prochlorococcus*: the structure and function of collective diversity. *Nat. Rev. Microbiol.* **13**, 13–27

15. Sullivan, M. B., Coleman, M. L., Weigele, P., Rohwer, F., and Chisholm, S. W. (2005) Three *Prochlorococcus* Cyanophage Genomes: Signature Features and Ecological Interpretations. *PLoS Biol.* **3**, e144

16. Labrie, S. J., Frois-Moniz, K., Osburne, M. S., Kelly, L., Roggensack, S. E., Sullivan, M. B., Gearin, G., Zeng, Q., Fitzgerald, M., Henn, M. R., and Chisholm, S. W. (2013) Genomes of marine cyanopodoviruses reveal multiple origins of diversity: Cyanopodovirus genomics. *Environ. Microbiol.* **15**, 1356–1376

17. Sieradzki, E. T., Ignacio-Espinoza, J. C., Needham, D. M., Fichot, E. B., and Fuhrman, J. A. (2019) Dynamic marine viral infections and major contribution to photosynthetic processes shown by spatiotemporal picoplankton metatranscriptomes. *Nat. Commun.* **10**, 1169

18. Liu, R., Liu, Y., Chen, Y., Zhan, Y., and Zeng, Q. (2019) Cyanobacterial viruses exhibit diurnal rhythms during infection. *Proc. Natl. Acad. Sci.* **116**, 14077–14082

19. Lindell, D., Jaffe, J. D., Coleman, M. L., Futschik, M. E., Axmann, I. M., Rector, T., Kettler, G., Sullivan, M. B., Steen, R., Hess, W. R., Church, G. M., and Chisholm, S. W. (2007) Genome-wide expression dynamics of a marine virus and host reveal features of co-evolution. *Nature.* **449**, 83–86

20. Puxty, R. J., Millard, A. D., Evans, D. J., and Scanlan, D. J. (2016) Viruses Inhibit CO_2 Fixation in the Most Abundant Phototrophs on Earth. *Curr. Biol.* **26**, 1585–1589

21. Puxty, R. J., Evans, D. J., Millard, A. D., and Scanlan, D. J. (2018) Energy limitation of cyanophage development: implications for marine carbon cycling. *ISME J.* **12**, 1273–1286

22. Lindell, D., Sullivan, M. B., Johnson, Z. I., Tolonen, A. C., Rohwer, F., and Chisholm, S. W. (2004) Transfer of photosynthesis genes to and from *Prochlorococcus* viruses. *Proc. Natl. Acad. Sci.* **101**, 11013–11018

23. Clokie, M. R. J., Shan, J., Bailey, S., Jia, Y., Krisch, H. M., West, S., and Mann, N. H. (2006) Transcription of a "photosynthetic" T4-type phage during infection of a marine cyanobacterium. *Environ. Microbiol.* **8**, 827–835

24. Sharon, I., Tzahor, S., Williamson, S., Shmoish, M., Man-Aharonovich, D., Rusch, D. B., Yooseph, S., Zeidner, G., Golden, S. S., Mackey, S. R., Adir, N., Weingart, U., Horn, D., Venter, J. C., Mandel-Gutfreund, Y., and Béjà, O. (2007) Viral photosynthetic reaction center genes and transcripts in the marine environment. *ISME J.* **1**, 492–501

25. Sharon, I., Alperovitch, A., Rohwer, F., Haynes, M., Glaser, F., Atamna-Ismaeel, N., Pinter, R. Y., Partensky, F., Koonin, E. V., Wolf, Y. I., Nelson, N., and Béjà, O. (2009) Photosystem I gene cassettes are present in marine virus genomes. *Nature.* **461**, 258–262

26. Puxty, R. J., Millard, A. D., Evans, D. J., and Scanlan, D. J. (2015) Shedding new light on viral photosynthesis. *Photosynth. Res.* **126**, 71–97

27. Dammeyer, T., Bagby, S. C., Sullivan, M. B., Chisholm, S. W., and Frankenberg-Dinkel, N. (2008) Efficient Phage-Mediated Pigment Biosynthesis in Oceanic Cyanobacteria. *Curr. Biol.* **18**, 442–448

28. Atkinson, J. T., Campbell, I., Bennett, G. N., and Silberg, J. J. (2016) Cellular Assays for Ferredoxins: A Strategy for Understanding Electron Flow through Protein Carriers That Link Metabolic Pathways. *Biochemistry.* **55**, 7047–7064

29. Burkhart, B. W., Febvre, H. P., and Santangelo, T. J. (2019) Distinct Physiological Roles of the Three Ferredoxins Encoded in the Hyperthermophilic Archaeon *Thermococcus kodakarensis*. *mBio.* **10**, e02807-18

30. Cassier-Chauvat, C., and Chauvat, F. (2014) Function and Regulation of Ferredoxins in the Cyanobacterium, *Synechocystis* PCC6803: Recent Advances. *Life.* **4**, 666–680

31. Terauchi, A. M., Lu, S., Zaffagnini, M., Tappa, S., Hirasawa, M., Tripathy, J. N., Knaff, D. B., Farmer, P. J., Lemaire, D., Hase, T., and Merchant, S. S. (2009) Pattern of Expression and Substrate Specificity of Chloroplast Ferredoxins from *Chlamydomonas reinhardtii*. **284**, 25867–25878

32. Onda, Y., Matsumura, T., Kimata-Ariga, Y., Sakakibara, H., Sugiyama, T., and Hase, T. (2000) Differential interaction of maize root ferredoxin:NADP(+) oxidoreductase with photosynthetic and non-photosynthetic ferredoxin isoproteins. *Plant Physiol.* **123**, 1037–1045

33. Finn, R. D., Attwood, T. K., Babbitt, P. C., Bateman, A., Bork, P., Bridge, A. J., Chang, H.-Y., Dosztányi, Z., El-Gebali, S., Fraser, M., Gough, J., Haft, D., Holliday, G. L., Huang, H., Huang, X., Letunic, I., Lopez, R., Lu, S., Marchler-Bauer, A., Mi, H., Mistry, J., Natale, D. A., Necci, M., Nuka, G., Orengo, C. A., Park, Y., Pesseat, S., Piovesan, D., Potter, S. C., Rawlings, N. D., Redaschi, N., Richardson, L., Rivoire, C., Sangrador-Vegas, A., Sigrist, C., Sillitoe, I., Smithers, B., Squizzato, S., Sutton, G., Thanki, N., Thomas, P. D., Tosatto, S. C. E., Wu, C. H., Xenarios, I., Yeh, L.-S., Young, S.-Y., and Mitchell, A. L. (2017) InterPro in 2017—beyond protein family and domain annotations. *Nucleic Acids Res.* **45**, D190–D199

34. Cornelissen, A., Hardies, S. C., Shaburova, O. V., Krylov, V. N., Mattheus, W., Kropinski, A. M., and Lavigne, R. (2012) Complete Genome Sequence of the Giant Virus OBP and Comparative Genome Analysis of the Diverse KZ-Related Phages. *J. Virol.* **86**, 1844–1852

35. Campbell, I. J., Bennett, G. N., and Silberg, J. J. (2019) Evolutionary Relationships Between Low Potential Ferredoxin and Flavodoxin Electron Carriers. *Front. Energy Res.* **7**, 79

36. Kim, J. Y., Nakayama, M., Toyota, H., Kurisu, G., and Hase, T. (2016) Structural and mutational studies of an electron transfer complex of maize sulfite reductase and ferredoxin. *J. Biochem. (Tokyo)*. **160**, 101–109

37. Kurisu, G., Kusunoki, M., Katoh, E., Yamazaki, T., Teshima, K., Onda, Y., Kimata-Ariga, Y., and Hase, T. (2001) Structure of the electron transfer complex between ferredoxin and ferredoxin-NADP(+) reductase. *Nat. Struct. Biol.* **8**, 117–21

38. Atkinson, H. J., Morris, J. H., Ferrin, T. E., and Babbitt, P. C. (2009) Using Sequence Similarity Networks for Visualization of Relationships Across Diverse Protein Superfamilies. *PLoS ONE*. **4**, e4345

39. Dai, S., Friemann, R., Glauser, D. A., Bourquin, F., Manieri, W., Schürmann, P., and Eklund, H. (2007) Structural snapshots along the reaction pathway of ferredoxin–thioredoxin reductase. *Nature*. **448**, 92–96

40. van den Heuvel, R. H. H., Svergun, D. I., Petoukhov, M. V., Coda, A., Curti, B., Ravasio, S., Vanoni, M. A., and Mattevi, A. (2003) The active conformation of glutamate synthase and its binding to ferredoxin. *J. Mol. Biol.* **330**, 113–128

41. Lelong, C., Boekema, E. J., Kruip, J., Bottin, H., Rögner, M., and Sétif, P. (1996) Characterization of a redox active cross-linked complex between cyanobacterial photosystem I and soluble ferredoxin. *EMBO J.* **15**, 2160–2168

42. Palma, P. N., Lagoutte, B., Krippahl, L., Moura, J. J. G., and Guerlesquin, F. (2005) *Synechocystis* ferredoxin/ferredoxin-NADP $^+$ -reductase/NADP $^+$ complex: Structural model obtained by NMR-restrained docking. *FEBS Lett.* **579**, 4585–4590

43. Hirasawa, M., Rubio, L. M., Griffin, J. L., Flores, E., Herrero, A., Li, J., Kim, S.-K., Hurley, J. K., Tollin, G., and Knaff, D. B. (2004) Complex formation between ferredoxin and *Synechococcus* ferredoxin:nitrate oxidoreductase. *Biochim. Biophys. Acta BBA - Bioenerg.* **1608**, 155–162

44. Morales, R., Kachalova, G., Vellieux, F., Charon, M.-H., and Frey, M. (2000) Crystallographic studies of the interaction between the ferredoxin-NADP $^+$ reductase and ferredoxin from the cyanobacterium *Anabaena*: looking for the elusive ferredoxin molecule. *Acta Crystallogr. D Biol. Crystallogr.* **56**, 1408–1412

45. Kubota-Kawai, H., Mutoh, R., Shinmura, K., Sétif, P., Nowaczyk, M. M., Rögner, M., Ikegami, T., Tanaka, H., and Kurisu, G. (2018) X-ray structure of an asymmetrical trimeric ferredoxin–photosystem I complex. *Nat. Plants*. **4**, 218–224

46. Okada, K. (2009) HO1 and PcyA proteins involved in phycobilin biosynthesis form a 1:2 complex with ferredoxin-1 required for photosynthesis. *FEBS Lett.* **583**, 1251–1256

47. Okada, K., and Hase, T. (2005) Cyanobacterial non-mevalonate pathway: (*E*)-4-hydroxy-3-methylbut-2-enyl diphosphate synthase interacts with ferredoxin in *Thermosynechococcus elongatus* BP-1. *J. Biol. Chem.* **280**, 20672–20679

48. Sullivan, M. B., Waterbury, J. B., and Chisholm, S. W. (2003) Cyanophages infecting the oceanic cyanobacterium *Prochlorococcus*. *Nature*. **424**, 1047–1051

49. Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990) Basic local alignment search tool. *J. Mol. Biol.* **215**, 403–410

50. Atkinson, J. T., Campbell, I. J., Thomas, E. E., Bonitatibus, S. C., Elliott, S. J., Bennett, G. N., and Silberg, J. J. (2019) Metalloprotein switches that display chemical-dependent electron transfer in cells. *Nat. Chem. Biol.* **15**, 189–195

51. Stephens, P. J., Thomson, A. J., Dunn, J. B. R., Keiderling, T. A., Rawlings, J., Rao, K. K., and Hall, D. O. (1978) Circular dichroism and magnetic circular dichroism of iron-sulfur proteins. *Biochemistry*. **17**, 4770–4778

52. Ta, D. T., and Vickery, L. E. (1992) Cloning, sequencing, and overexpression of a [2Fe-2S] ferredoxin gene from *Escherichia coli*. *J. Biol. Chem.* **267**, 11120–11125

53. Moczygemba, C., Guidry, J., Jones, K. L., Gomes, C. M., Teixeira, M., and Wittung-Stafshede, P. (2001) High stability of a ferredoxin from the hyperthermophilic archaeon *A. ambivalens*: Involvement of electrostatic interactions and cofactors. *Protein Sci.* **10**, 1539–1548

54. Osteryoung, J. G., and Osteryoung, R. A. (1985) Square wave voltammetry. *Anal. Chem.* **57**, 101–110

55. Jin, S., Kurtz, D. M., Liu, Z.-J., Rose, J., and Wang, B.-C. (2004) Displacement of iron by zinc at the diiron site of *Desulfovibrio vulgaris* rubrerythrin: X-ray crystal structure and anomalous scattering analysis. *J. Inorg. Biochem.* **98**, 786–796

56. Iametti, S., Uhlmann, H., Sala, N., Bernhardt, R., Ragg, E., and Bonomi, F. (1996) Reversible, Non-Denaturing Metal Substitution in Bovine Adrenodoxin and Spinach Ferredoxin and the Different Reactivities of [2Fe-2S]-Cluster-Containing Proteins. *Eur. J. Biochem.* **239**, 818–826

57. Kojoh, K., Matsuzawa, H., and Wakagi, T. (1999) Zinc and an N-terminal extra stretch of the ferredoxin from a thermoacidophilic archaeon stabilize the molecule at high temperature. *Eur. J. Biochem.* **264**, 85–91

58. Sazinsky, M. H., LeMoine, B., Orofino, M., Davydov, R., Bencze, K. Z., Stemmler, T. L., Hoffman, B. M., Argüello, J. M., and Rosenzweig, A. C. (2007) Characterization and Structure of a Zn^{2+} and [2Fe-2S]-containing Copper Chaperone from *Archaeoglobus fulgidus*. *J. Biol. Chem.* **282**, 25950–25959

59. Lo Conte, L. (2000) SCOP: a Structural Classification of Proteins database. *Nucleic Acids Res.* **28**, 257–259

60. Fish, A., Danieli, T., Ohad, I., Nechushtai, R., and Livnah, O. (2005) Structural Basis for the Thermostability of Ferredoxin from the Cyanobacterium *Mastigocladus laminosus*. *J. Mol. Biol.* **350**, 599–608

61. Morales, R., Charon, M. H., Hudry-Clergeon, G., Pétillot, Y., Norager, S., Medina, M., and Frey, M. (1999) Refined X-ray structures of the oxidized, at 1.3 A, and reduced, at 1.17 A, [2Fe-2S] ferredoxin from the cyanobacterium *Anabaena* PCC7119 show redox-linked conformational changes. *Biochemistry*. **38**, 15764–15773

62. Mutoh, R., Muraki, N., Shinmura, K., Kubota-Kawai, H., Lee, Y.-H., Nowaczyk, M. M., Rögner, M., Hase, T., Ikegami, T., and Kurisu, G. (2015) X-ray Structure and Nuclear Magnetic Resonance Analysis of the Interaction Sites of the Ga-Substituted Cyanobacterial Ferredoxin. *Biochemistry*. **54**, 6052–6061

63. Sakakibara, Y., Kimura, H., Iwamura, A., Saitoh, T., Ikegami, T., Kurisu, G., and Hase, T. (2012) A new structural insight into differential interaction of cyanobacterial and plant ferredoxins with nitrite reductase as revealed by NMR and X-ray crystallographic studies. *J. Biochem. (Tokyo)*. **151**, 483–492

64. Ashkenazy, H., Abadi, S., Martz, E., Chay, O., Mayrose, I., Pupko, T., and Ben-Tal, N. (2016) ConSurf 2016: an improved methodology to estimate and visualize evolutionary conservation in macromolecules. *Nucleic Acids Res.* **44**, W344-350

65. Barstow, B., Agapakis, C. M., Boyle, P. M., Grandl, G., Silver, P. A., and Wintermute, E. H. (2011) A synthetic system links FeFe-hydrogenases to essential *E. coli* sulfur metabolism. *J. Biol. Eng.* **5**, 7

66. Houot, L., Floutier, M., Marteyn, B., Michaut, M., Picciocchi, A., Legrain, P., Aude, J.-C., Cassier-Chauvat, C., and Chauvat, F. (2007) Cadmium triggers an integrated reprogramming of the metabolism of *Synechocystis* PCC6803, under the control of the Slr1738 regulator. *BMC Genomics*. **8**, 350

67. Kimata, Y., and Hase, T. (1989) Localization of Ferredoxin Isoproteins in Mesophyll and Bundle Sheath Cells in Maize Leaf. *Plant Physiol.* **89**, 1193–1197

68. Singh, A. K., Elvitigala, T., Cameron, J. C., Ghosh, B. K., Bhattacharyya-Pakrasi, M., and Pakrasi, H. B. (2010) Integrative analysis of large scale expression profiles reveals core transcriptional

response and coordination between multiple cellular processes in a cyanobacterium. *BMC Syst. Biol.* **4**, 105

69. Bottin, H., and Lagoutte, B. (1992) Ferrodoxin and flavodoxin from the cyanobacterium *Synechocystis* sp. PCC 6803. *Biochim. Biophys. Acta BBA - Bioenerg.* **1101**, 48–56

70. Hurley, J. K., Weber-Main, A. M., Hodges, A. E., Stankovich, M. T., Benning, M. M., Holden, H. M., Cheng, H., Xia, B., Markley, J. L., Genzor, C., Gomez-Moreno, C., Hafezi, R., and Tollin, G. (1997) Iron-sulfur cluster cysteine-to-serine mutants of *Anabaena* -2Fe-2S- ferredoxin exhibit unexpected redox properties and are competent in electron transfer to ferredoxin:NADP+ reductase. *Biochemistry*. **36**, 15109–15117

71. Nechushtai, R., Lammert, H., Michaeli, D., Eisenberg-Domovich, Y., Zuris, J. A., Luca, M. A., Capraro, D. T., Fish, A., Shimshon, O., Roy, M., Schug, A., Whitford, P. C., Livnah, O., Onuchic, J. N., and Jennings, P. A. (2011) Allostery in the ferredoxin protein motif does not involve a conformational switch. *Proc. Natl. Acad. Sci.* **108**, 2240–2245

72. Chang, C.-J., Lee, C.-C., Chan, Y.-T., Trudeau, D. L., Wu, M.-H., Tsai, C.-H., Yu, S.-M., Ho, T.-H. D., Wang, A. H.-J., Hsiao, C.-D., Arnold, F. H., and Chao, Y.-C. (2016) Exploring the Mechanism Responsible for Cellulase Thermostability by Structure-Guided Recombination. *PLOS ONE*. **11**, e0147485

73. Otey, C. R., Landwehr, M., Endelman, J. B., Hiraga, K., Bloom, J. D., and Arnold, F. H. (2006) Structure-Guided Recombination Creates an Artificial Family of Cytochromes P450. *PLoS Biol.* **4**, e112

74. Mutter, A. C., Tyryshkin, A. M., Campbell, I. J., Poudel, S., Bennett, G. N., Silberg, J. J., Nanda, V., and Falkowski, P. G. (2019) De novo design of symmetric ferredoxins that shuttle electrons *in vivo*. *Proc. Natl. Acad. Sci.* 10.1073/pnas.1905643116

75. Wu, B., Atkinson, J. T., Kahanda, D., Bennett, G. N., and Silberg, J. J. (2019) Combinatorial design of chemical-dependent protein switches for controlling intracellular electron transfer. *AIChE J.* 10.1002/aic.16796

76. Aylward, F. O., Boeuf, D., Mende, D. R., Wood-Charlson, E. M., Vislova, A., Eppley, J. M., Romano, A. E., and DeLong, E. F. (2017) Diel cycling and long-term persistence of viruses in the ocean's euphotic zone. *Proc. Natl. Acad. Sci.* **114**, 11446–11451

77. Yoshida, T., Nishimura, Y., Watai, H., Haruki, N., Morimoto, D., Kaneko, H., Honda, T., Yamamoto, K., Hingamp, P., Sako, Y., Goto, S., and Ogata, H. (2018) Locality and diel cycling of viral production revealed by a 24 h time course cross-omics analysis in a coastal region of Japan. *ISME J.* **12**, 1287–1295

78. Kiene, R. P., Linn, L. J., González, J., Moran, M. A., and Bruton, J. A. (1999) Dimethylsulfoniopropionate and methanethiol are important precursors of methionine and protein-sulfur in marine bacterioplankton. *Appl. Environ. Microbiol.* **65**, 4549–4558

79. Andreae, M. O. (1990) Ocean-atmosphere interactions in the global biogeochemical sulfur cycle. *Mar. Chem.* **30**, 1–29

80. del Valle, D., Martínez-García, S., Sañudo-Wilhelmy, S., Kiene, R., and Karl, D. (2015) Methionine and dimethylsulfoniopropionate as sources of sulfur to the microbial community of the North Pacific Subtropical Gyre. *Aquat. Microb. Ecol.* **75**, 103–116

81. Morris, R. M., Rappé, M. S., Connon, S. A., Vergin, K. L., Siebold, W. A., Carlson, C. A., and Giovannoni, S. J. (2002) SAR11 clade dominates ocean surface bacterioplankton communities. *Nature*. **420**, 806–810

82. Tripp, H. J., Kitner, J. B., Schwalbach, M. S., Dacey, J. W. H., Wilhelm, L. J., and Giovannoni, S. J. (2008) SAR11 marine bacteria require exogenous reduced sulphur for growth. *Nature*. **452**, 741–744

83. Becker, J. W., Hogle, S. L., Rosendo, K., and Chisholm, S. W. (2019) Co-culture and biogeography of *Prochlorococcus* and SAR11. *ISME J.* **13**, 1506–1519

84. Engler, C., Kandzia, R., and Marillonnet, S. (2008) A one pot, one step, precision cloning method with high throughput capability. *PloS One*. **3**, e3647

85. Salis, H. M. (2011) The ribosome binding site calculator. *Methods Enzymol.* **498**, 19–42

86. Gerlt, J. A. (2017) Genomic Enzymology: Web Tools for Leveraging Protein Family Sequence–Function Space and Genome Context to Discover Novel Functions. *Biochemistry.* **56**, 4293–4308

87. Shannon, P. (2003) Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks. *Genome Res.* **13**, 2498–2504

88. Edgar, R. C., Drive, R. M., and Valley, M. (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* **32**, 1792–1797

89. Schrödinger, LLC (2015) The PyMOL Molecular Graphics System, Version 1.8

90. Li, B., and Elliott, S. J. (2016) The Catalytic Bias of 2-Oxoacid:ferredoxin Oxidoreductase in CO_2: evolution and reduction through a ferredoxin-mediated electrocatalytic assay. *Electrochimica Acta.* **199**, 349–356

91. Vonrhein, C., Flensburg, C., Keller, P., Sharff, A., Smart, O., Paciorek, W., Womack, T., and Bricogne, G. (2011) Data processing and analysis with the *autoPROC* toolbox. *Acta Crystallogr. D Biol. Crystallogr.* **67**, 293–302

92. Kabsch, W. (2010) *XDS. Acta Crystallogr. D Biol. Crystallogr.* **66**, 125–132

93. Evans, P. R., and Murshudov, G. N. (2013) How good are my data and what is the resolution? *Acta Crystallogr. D Biol. Crystallogr.* **69**, 1204–1214

94. Shinohara, F., Kurisu, G., Hanke, G., Bowsher, C., Hase, T., and Kimata-Ariga, Y. (2017) Structural basis for the isotype-specific interactions of ferredoxin and ferredoxin: NADP+ oxidoreductase: an evolutionary switch between photosynthetic and heterotrophic assimilation. *Photosynth. Res.* **134**, 281–289

95. Emsley, P., and Cowtan, K. (2004) *Coot*: model-building tools for molecular graphics. *Acta Crystallogr. D Biol. Crystallogr.* **60**, 2126–2132

96. Afonine, P. V., Grosse-Kunstleve, R. W., Echols, N., Headd, J. J., Moriarty, N. W., Mustyakimov, M., Terwilliger, T. C., Urzhumtsev, A., Zwart, P. H., and Adams, P. D. (2012) Towards automated crystallographic structure refinement with *phenix.refine*. *Acta Crystallogr. D Biol. Crystallogr.* **68**, 352–367

97. Yennamalli, R., Arangarasan, R., Bryden, A., Gleicher, M., and Phillips, G. N. (2014) Using a commodity high-definition television for collaborative structural biology. *J. Appl. Crystallogr.* **47**, 1153–1157

98. Thorn, A., and Sheldrick, G. M. (2011) *ANODE*: anomalous and heavy-atom density calculation. *J. Appl. Crystallogr.* **44**, 1285–1287

99. Morin, A., Eisenbraun, B., Key, J., Sanschagrin, P. C., Timony, M. A., Ottaviano, M., and Sliz, P. (2013) Collaboration gets the most out of software. *eLife.* **2**, e01456

100. Berman, H., Henrick, K., and Nakamura, H. (2003) Announcing the worldwide Protein Data Bank. *Nat. Struct. Mol. Biol.* **10**, 980–980

101. Campbell, R. E., Tour, O., Palmer, A. E., Steinbach, P. A., Baird, G. S., Zacharias, D. A., and Tsien, R. Y. (2002) A monomeric red fluorescent protein. *Proc. Natl. Acad. Sci.* **99**, 7877–7882

Figure 1. Network analysis of cyanophage and cyanobacterial Fd sequence similarities. (**A**) A network illustrating the relationships of 807 cyanobacterial Fds and 28 cyanophage Fds. Edges are drawn between two Fds with a sequence alignment score ≥30. For the twelve largest clusters (Clusters I to XII), average gene lengths encoded by the ORFs are listed with standard deviation and clusters are colored on a viridis gradient by the number of species present in each cluster. (**B**) Cyanophage Fds, cyanobacterial Fds, and a plant Fd are highlighted in Cluster VI. (**C**). Partner oxidoreductases that have the potential to interact with cyanophage Fd during infection of *P. marinus*.

Figure 2. Cyanophage and host cyanobacterial Fd sequence comparisons. (**A**) A matrix showing the pairwise sequence identity of pssm2-Fd and Fds from host cyanobacteria, cyanobacteria having reported structures, and zm-Fd1, colored darker shades of blue to indicate greater identity. (**B**) A multiple sequence alignment illustrates the conservation of cysteines, charged residues, and residues implicated in binding FNR (triangles), SIR (asterisks), and PSI (circles) (36, 37, 45). Negatively charged residues are colored red, positively charged residues are colored blue, and cysteines are colored yellow.

Figure 3. Purified pssm2-Fd is a low potential protein with low thermostability. (**A**) The absorbance and (**B**) circular dichroism spectra of pssm2-Fd (50 µM) reveal features characteristic of [2Fe-2S] Fds that support photosynthesis. (**C**). Effect of temperature on the ellipticity of pssm2-Fd (50 µM) at 427 nm. The midpoint of the transition occurs at 29 °C. (**D**) Square wave voltammetry of pssm2-Fd (620 µM) reveals a midpoint reduction potential of -334 mV. Voltammetry prior to background subtraction is shown as an inset.

Figure 4. Comparison of the pssm2-Fd structure with cyanobacterial Fd structures. (**A**) Front- and back-facing ribbon structures of pssm2-Fd. Iron atoms are shown as brown spheres, while sulfur atoms are colored yellow. (**B**) Front- and back-facing ribbon structures of pssm2-Fd colored with ConSurf conservation scores generated using Cluster VI Fds (Figure 1). The most conserved residues are shown in magenta, while the least conserved residues are in cyan.

Figure 5. pssm2-Fd supports ET from plant FNR to SIR. (**A**) An *E. coli* auxotroph unable to produce sulfide endogenously can be rescued by a three-component pathway consisting of FNR, Fd, and SIR. Fd and SIR can be readily substituted for homologs. (**B**) The effect of aTc concentration on the growth complementation of *E. coli* EW11 transformed with vectors that express pssm2-Fd using an aTc-inducible promoter and zm-FNR/zm-SIR using constitutive promoters. aTc concentrations (0, 1.6, 3.1, 6.3, 12.5, 25, 50, 100 ng/mL) are color coded as gradients of dark purple (highest) to light purple (lowest). Error bars represent standard deviation from three biological replicates.

Figure 6. pssm2-Fd can transfer electrons to host SIRs. Growth complementation observed at (**A**) 37°C, (**B**) 30°C, and (**C**) 23°C with *E. coil* EW11 transformed with a vector that constitutively expresses zm-FNR with three different SIR (zm-SIR, pmN2-SIR, and pm9211-SIR) and a vector that uses an aTc-inducible promoter to express four different Fds (ml-Fd1, pssm2-Fd, pmN2-Fd1, and pm9211-Fd1). Experiments were performed in the absence (grey bars) and presence (orange, yellow, and blue bars) of aTc to evaluate the dependence of growth upon Fd-mediated ET from FNR to SIR. Error bars represent standard deviation from three biological replicates. For each strain, an independent two-tailed t-test was used to compare OD in the absence or presence of aTc ($\alpha=0.05$). Significant differences are noted with asterisks.

Figure 7. Relative expression of Fds evaluated using RFP fusion proteins. Fluorescence is aTc-inducible for all cells expressing cyanophage and cyanobacterial Fds fused to RFP under expression conditions that are identical to those in the cellular assay, with the host pmN2-Fd exhibiting the highest expression levels. For each strain, an independent two-tailed t-test was used to compare AU/OD in the absence or presence of aTc ($\alpha=0.05$). Significant differences are noted with asterisks.

ABBREVIATONS

Fd	ferredoxin
pssm2-Fd	myovirus P-SSM2 Fd
ET	electron transfer
PSI	photosystem I
PcyA	phycocyanobilin:ferredoxin oxidoreductase
PebS	phycoerythrobilin synthase
FNR	Fd-NADP reductase
SIR	sulfite reductase
GS	glutamate synthase
FTR	Fd-thioredoxin reductase
NIR	nitrate reductase
HO	heme oxygenase
GcpE	4-Hydroxy-3-methylbut-2-enyl diphosphate synthase
SSN	sequence similarity network
CAPS	N-cyclohexyl-3-aminopropanesulfonic
MES	2-(N-Morpholino)ethanesulfonic acid
TAPS	N-[Tris(hydroxymethyl)methyl]-3-aminopropanesulfonic acid
IPTG	isopropyl β-D-1-thiogalactopyranoside
DTT	dithiothreitol
MOPS	3-Morpholinopropane-1-sulfonic acid
CHES	N-Cyclohexyl-2-aminoethanesulfonic acid

ORFs	open reading frames
LB	lysogeny broth
SHE	Standard Hydrogen Electrode
aTc	anhydrotetracycline
M9c	nonselective modified M9 medium
M9sa	selective modified M9 medium
DMSP	dimethylsulfoniopropionate
DMSO	dimethylsulfoxide
OD	Optical density
RFP	red fluorescent protein
CD	circular dichroism
zm-Fd1	*Zea mays* Fd 1
s6803-Fd1	*Synechocystis sp.* PCC 6803 Fd 1
nsp-Fd1	*Nostoc sp.* PCC 7119 Fd 1
te-Fd1	*Thermosynechococcus elongatus* Fd 1
ml-Fd1	*Mastigocladus laminosus* Fd 1
pmN1-Fd1	*Prochlorococcus marinus* NATL1A Fd 1
pmN2-Fd1	*Prochlorococcus marinus* NATL2A Fd 1
pm9211-Fd1	*Prochlorococcus marinus* MIT9211 Fd 1
zm-SIR	*Zea mays* SIR
zm-FNR	*Zea mays* FNR
pmN2-SIR	*Prochlorococcus marinus* NATL2A SIR

pm9211-SIR *Prochlorococcus marinus*
MIT9211 SIR