A novel factor essential for unconventional secretion
of chitinase Cts1

Michèle Reindl[1,2¶], Janpeter Stock[1,2,#a¶] Kai P. Hussnaetter[1,2], Aycin Genc[1,#b], Andreas Brachmann[3], and Kerstin Schipper[1,2*]

[1] Institute for Microbiology, Heinrich Heine University Düsseldorf, Universitätsstraße 1, 40225 Düsseldorf, Germany

[2] Bioeconomy Science Center (BioSC), Forschungszentrum Jülich, Jülich, Germany

[3] Ludwig-Maximilians-Universität München, Faculty of Biology, Genetics, Großhaderner Str. 2-4, 82152, Planegg-Martinsried, Germany

[#a] Current Address: NUMAFERM GmbH, Merowingerplatz 1a, 40225 Düsseldorf, Germany

[#b] Current Address: VITROMED GmbH, Raiffeisenstr. 15A, 40764 Langenfeld, Germany

[*] Corresponding author:
E-mail: kerstin.schipper@uni-duesseldorf.de (KS)

[¶] These authors contributed equally to this work.

Running Header: A novel unconventional secretion factor

Abstract

Subcellular targeting of proteins is essential to orchestrate cytokinesis in eukaryotic cells.

During cell division of *Ustilago maydis*, for example, chitinases must be specifically targeted

to the fragmentation zone at the site of cell division to degrade remnant chitin and thus

separate mother and daughter cells. Chitinase Cts1 is exported to this location via an

unconventional secretion pathway. The underlying mechanism is largely unexplored. Here,

we applied a forward genetic screen based on UV mutagenesis to identify components

essential for Cts1 export. The screen revealed a novel factor termed Jps1 lacking known

protein domains. Deletion of the corresponding gene confirmed its essential role for Cts1

secretion. Localization studies demonstrated that Jps1 co-localizes with Cts1 in the

fragmentation zone of dividing yeast cells. While loss of Jps1 leads to exclusion of Cts1 from

the fragmentation zone and strongly reduced unconventional secretion, deletion of the

chitinase does not disturb Jps1 localization. In essence, we identified a novel component of

unconventional secretion that operates in the fragmentation zone to enable export of Cts1 via

a lock-type mechanism.

Author summary

The development of eukaryotic cells strongly depends on the exact localization of proteins in

space and time. Cell wall remodeling during fungal development is mediated by hydrolytic

enzymes like glucanases and chitinases. During cytokinesis of the fungus *Ustilago maydis*,

chitinases orchestrate cell division by degrading the connecting cell wall in the fragmentation

zone between mother and daughter cell. One of these chitinases, Cts1, is exported to the

fragmentation zone via a recently described lock-type unconventional secretion mechanism.

However, detailed molecular knowledge of the pathway is lacking. Here we describe the

establishment of a genetic screen to identify novel factors for Cts1 export. An unknown

protein termed Jps1 was identified. Absence of this protein leads to impaired Cts1 secretion

and its aberrant localization. We speculate that Jps1 might deal as an anchoring factor needed

for successful Cts1 targeting to the fragmentation zone. Since Cts1 is also exploited for

unconventional secretion of biotechnologically relevant proteins, these results also have a

strong impact for applied sciences.

Keywords

Forward genetic screen, β-galactosidase, β-glucuronidase, unconventional secretion, *Ustilago*

maydis, UV mutagenesis.

Abbreviations

aa, amino acid; Gus, β-glucuronidase; LacZ, β-galactosidase; MUC, 4-methylumbelliferyl β-

D-N,N′,N″-triacetylchitotrioside; MUG, 4-methylumbelliferyl-ß-D-glucuronide; ONPG, o-

nitrophenyl-β-D-galactopyranoside; SNP, single nucleotide polymorphism; X-Gal, 5-brome-

4-chlor-3-indoxyl-β-D-galactopyranoside; X-Gluc, 5-brome-4-chlor-3-indolyl-β-D-

glucuronic acid.

Introduction

Protein targeting is required to orchestrate essential cellular functions. Eukaryotic cells particularly rely on this process because of their compartmentalization and the necessity of equipping membrane-enclosed organelles with cognate protein subsets (1). Protein targeting is mediated mostly by signal sequences. This is exemplified by the N-terminal signal peptide for entry of the endoplasmic reticulum. The endomembrane system was thought to be the only export route for long time. However, recent years challenged this view by the finding that many proteins lacking a signal peptide are secreted by other mechanisms. The term unconventional secretion collectively describes protein export mechanisms that circumvent signal peptide-mediated passage through the canonical endoplasmic reticulum (ER) – Golgi pathway (2, 3). Unconventional secretion has been discovered in lower eukaryotes like the fungal model *Saccharomyces cerevisiae* or the amoeba *Dictyostelium discoideum*, but also plays important roles in higher eukaryotes. It is even involved in human disease like in infections with the human immunodeficiency (HIV) or Epstein Barr viruses (4-6).

Recently, a novel mechanism of unconventional secretion has been described for chitinase Cts1 in the model microorganism *Ustilago maydis* (7). In its yeast form the fungus grows by budding. In these cells, Cts1 acts in concert with a second chitinase, Cts2, and mediates cell separation during cytokinesis. Elimination of both enzymes results in a cytokinesis defect and the formation of cell aggregates (8). Cts2 has a predicted N-terminal signal peptide and is thus thought to be secreted via the conventional secretion route, pointing towards an intricate interplay between both pathways.

In line with its cellular function, Cts1 translocates into the fragmentation zone of budding yeast cells (8, 9). This unique small compartment arises between mother and daughter cell after consecutive formation of two septa at the cell boundary (7). Recently we

demonstrated that Cts1 release depends on cytokinesis and that the fragmentation zone is its

most likely site of secretion, suggesting a lock-type mechanism (9). The septation proteins

Don1 and Don3 are essential for Cts1 release (9). Don1 is a guanosine triphosphate exchange

factor (GEF) that is delivered into the fragmentation zone by motile early endosomes, and

Don3 is a Germinal Centre kinase (10, 11). Both proteins are required for secondary septum

formation and their absence results in a cytokinesis defect similar to the one observed for the

cts1/cts2 deletion strain (8). To obtain further insights into subcellular targeting and

unconventional secretion of Cts1, we here developed and applied a UV-mutagenesis screen to

identify components of the unconventional secretion pathway.

Results

A genetic screen for mutants impaired in Cts1 secretion identifies a novel component

To identify mutants impaired in Cts1 secretion a forward genetic screen based on UV mutagenesis was established (Fig. 1, Fig. S1; for experimental details see Materials and Methods section). To this end a screening strain derived from the haploid FB2 wild type strain (mating type *a2b2*) was developed. This allows for co-infections of the host plant maize with compatible strains of differing *a* and *b* alleles like FB1 (mating type *a1b1*) to obtain meiotic progeny (12).

Fig. 1. Rationale of the forward genetic screen. The three reporters LacZ-Cts1, Gus-Cts1 and endogenous Cts1 were used to identify mutants with diminished Cts1 secretion. After

UV-mutagenesis, a high-throughput screen for absence of LacZ activity of LacZ-Cts1 was
conducted on X-Gal containing plates. Next, colonies were patched on X-Gluc containing
plates to verify the result with the Gus-Cts1 reporter. Remaining candidates were assayed in
quantitative liquid assays for extra- and intracellular LacZ, Gus, and Cts1 activity. Candidates
with diminished extracellular activity of all three reporters but unimpaired intracellular
activity were collected. After genome sequencing, the mutation responsible for diminished
secretion was identified by PCR analysis of different meiotic progeny showing a similar
secretion phenotype on loci containing SNPs as identified in the genome comparison with the
progenitor strain. For details see Fig. S1.
131

To minimize false positive screening hits, three different reporters for Cts1 secretion

were employed: the bacterial reporter enzymes β-glucuronidase (Gus; published in 13, 14)

and β-galactosidase (LacZ; newly established for *U. maydis*) as fusion proteins with Cts1

(Gus-Cts1, LacZ-Cts1), and endogenous Cts1 (15). The genetic constructs for Gus-Cts1 and

LacZ-Cts1 were stably inserted in the genome at distinct loci resulting in screening strain

FB2^{CGL} (Table 1) (Fig. 2A). Strains harboring cytoplasmic Gus or LacZ were used as lysis

controls and to mimic defective Cts1 secretion with intracellular reporter accumulation

(control strains FB2 Gus$_{cyt}$ and FB2 LacZ$_{cyt}$). Plate assays with colorimetric substrates

demonstrated that the screening strain developed the expected blue color indicative for

extracellular Gus and LacZ activity while the controls showed no color. Artificial cell lysis

demonstrated the presence of intracellular reporter activity in the controls confirming the

absence of significant cell lysis (Fig. 2B). Furthermore, quantitative liquid assays detecting

extracellular LacZ, Gus and Cts1 activity revealed strongly enhanced extracellular activities

of all reporters in FB2^{CGL} compared to the respective control strains containing cytoplasmic

versions or lacking the reporters (Fig. 2C). Activity measurements in cell extracts again

confirmed that all reporters were functional (Fig. 2D). In addition, since we intended to use

LacZ activity for high-throughput screening a mixing experiment to assay its suitability was

conducted. Mixing of FB2^{CGL} with FB2 LacZ$_{cyt}$ in different ratios on indicator plates showed

that the ratio of mixing was reflected by the ratio of colonies with a blue halo versus colorless

colonies (Fig. 2E-G). This demonstrated that mutants with defective secretion can be

identified vis-a-vis colonies with intact secretion as an important requirement for the

screening procedure. Finally, plant infection experiments indicated that pathogenicity of

FB2^{CGL} was not impaired, since FB2^{CGL} (*a2b2*) crossed with the compatible mating partner

FB1 (*a1b1*) elicited the typical symptoms including tumor formation on maize seedlings (Fig.

S3). Thus, genetic back-crossing experiments are feasible with this strain. In summary, we

successfully designed the strain FB2^{CGL} for screening mutants defective in unconventional

secretion of Cts1.

Fig. 2. Establishing screening strain FB2^{CGL} harboring three reporters for
unconventional secretion. A, Scheme of the genetic constructs for the three reporters present
in distinct loci of *U. maydis* strain FB2^{CGL}. The reporter genes encoding endogenous Cts1 as

well as Gus-Cts1 and LacZ-Cts1 are located on three different chromosomes (chr.). While the

cts1 gene on chr. 12 has not been modified and is present in its natural setup controlled by its

native promoter active in yeast cells, both *lacZ:cts1* and *gus:cts1* gene fusions were inserted

artificially by homologous recombination using described protease loci (16). Both

translational fusion genes are hooked up to the strong synthetic promoter P_{oma} which is

constitutively active in yeast cells grown on CM-glc. Insertion of the two reporters results in

deletion of the genes *upp1* and *pep4* which both encode harmful extracellular proteases (16).

B, Plate assay to determine the suitability of the used reporters in FB2^{CGL} in comparison to

the lysis controls FB2 Gus$_{cyt}$ and FB2 LacZ$_{cyt}$ harboring cytoplasmic reporter enzymes

(middle, bottom). FB2 was used as negative control containing neither Gus nor LacZ (top).

X-Gluc and X-Gal were applied as colorimetric substrates for Gus and LacZ, respectively. To

visualize intracellular reporter activity, cells were lysed using liquid nitrogen. cyt, control

strains with intracellular reporter activity or absent reporter activity. lys, cells lysed by

treatment with liquid nitrogen. **C,** Liquid assays using the substrates MUG, ONPG and MUC

to determine the suitability of the three reporters Gus, LacZ and Cts1, respectively. Culture

supernatants (Gus-Cts1/LacZ-Cts1) or intact cells (Cts1) of indicated strains were tested to

determine extracellular reporter activities. Upper panel: Cts1 activity based on conversion of

MUC; middle panel: Gus activity based on conversion of MUG; lower panel: LacZ activity

based on conversion of ONPG. Error bars represent standard deviation of three biological

replicates. **D,** Liquid assays using the substrates MUG, ONPG and MUC to determine the

suitability of the three reporters Gus-Cts1, LacZ-Cts1 and Cts1, respectively. 10 µg cell

extracts of the indicated strains were used to assay intracellular reporter activities. Upper

panel: Cts1 activity based on conversion of MUC; middle panel: Gus activity based on

conversion of MUG; lower panel: LacZ activity based on conversion of ONPG. Error bars

represent standard deviation of three biological replicates. **E-G,** Mixed culture experiments to

verify applicability of the LacZ reporter for high-throughput screening on the colorimetric

substrate X-Gal. Screening strain FB2^{CGL} and control strain FB2 LacZ$_{cyt}$ were mixed in the

indicated ratios and plated onto indicator plates containing X-Gal to visualize extracellular

LacZ activity (blue color). Photographs were taken after incubation for 1 d. **E**, 100% FB2^{CGL};

F, 50% FB2^{CGL}/50% FB2 LacZ$_{cyt}$; **G**, 10% FB2^{CGL}/90% FB2 LacZ$_{cyt}$.

In order to screen for diminished Cts1 activity, strain FB2^{CGL} was subjected to UV

irradiation with approximately 1% survival rate (Fig. 1, Fig. S1). Mutagenized cells were

plated on X-Gal plates to detect extracellular LacZ activity based on the reporter LacZ-Cts1.

Approximately 185,000 colonies were screened with a focus on mutants that exhibited

normal growth (i.e. normal colony size) but impaired Cts1 secretion. 2,087 candidate mutants

showing strongly reduced or absent blue halos were patched on X-Gluc plates employing the

Gus marker (Fig. 1). Of those, 566 that stayed colorless again were retested in plate assays

using both the LacZ and the Gus marker to confirm these results. 112 remaining colorless

candidates were assayed for Cts1, Gus and LacZ activity in quantitative liquid assays. Again,

mutants showing reduced growth in liquid culture were sorted out to ensure that reduced

secretion is not connected to growth problems. Finally, three mutants were left, in which

Cts1, LacZ and Gus activity was present intracellularly, but strongly reduced extracellularly

(FB2^{CGL}mut1-3; Fig. 3A,B). Western blot analysis confirmed equal protein amounts for

LacZ-Cts1 and Gus-Cts1 in cell extracts of these mutants (Fig. S2A). All mutants showed

wildtype growth rates suggesting that reduced secretion is not resulting from growth defects

(Fig. S2B).

212
213

214
215
Fig. 3. The screen identifies the uncharacterized protein Jps1. A, Plate assays of indicated
strains for extracellular LacZ activity using X-Gal and extracellular Gus activity using X-
Gluc. The assays are based on secretion of the reporters LacZ-Cts1 and Gus-Cts1 and the
respective substrates are only converted, if the fusion protein is secreted. This leads to the
formation of blue colonies (and blue halos in case of Gus) resulting from extracellular
substrate conversion. Mutants FB2^{CGL}mut1, mut2 and mut3 were identified after UV
mutagenesis of strain FB2^{CGL} on screening plates (white colonies on both X-Gluc and X-Gal)
and show the expected reduced extracellular activities indicating deficiency in Cts1 secretion.
B, Quantitative liquid assays detecting intracellular (int, filled columns) and extracellular (ex,
open columns) Gus, LacZ and Cts1 reporter activity. Screening strain FB2^{CGL} and a *cts1*
deletion mutant (AB33 cts1Δ, 15) were used as positive and negative controls, respectively.

Activities obtained for FB2^{CGL} were set to 100% to allow for a direct comparison of all
strains. The assay was conducted thrice with similar results and a representative replicate is
shown. **C**, In all three identified mutants, base exchanges in gene *umag_03776*, now termed
jps1, were identified (left). At the aa level, mutations result in the introduction of premature
stop codons leading to the production of truncated proteins in all three mutants (right). Mut1
carries an additional aa exchange at position 335. Mutations identified in FB2^{CGL}mut1-3 are
indicated with red lines in the schematic representation of the protein. The size of native Jps1
is 609 aa.
235

To identify the responsible mutations whole genome sequencing comparing

FB2^{CGL}mut1 with its progenitor FB2^{CGL} was performed, using the published sequence of *U.*

maydis UM521 as a template for the assembly

(https://mycocosm.jgi.doe.gov/Ustma2_1/Ustma2_1.home.html; last accessed 2019/07/25)

(17). 32 base-pair substitutions were detected in the comparison of FB2^{CGL} and FB2^{CGL}mut1.

The majority of mutations were found to be 5´-C→T or 5´-CC→TT transitions which are

expected for UV-induced mutations (Fig. S3) (18). 9 single nucleotide polymorphisms

(SNPs) were located in non-coding regions and were thus unlikely to cause the observed

defect in unconventional secretion. 11 of 23 substitutions in coding regions were irrelevant

silent mutations. The remaining 12 SNPs led to 10 aa replacements in 9 encoded proteins and

hence constituted the top remaining candidates (Fig. S3). To locate which of these mutations

was responsible for the defective Cts1 secretion, genetic back-crossing experiments were

performed using plant infections. FB2^{CGL}mut1 was pathogenic in crosses with compatible

FB1^{CGL} which carries similar reporter genes but an opposite mating type (Fig. S4, Fig. S5; see

Materials and Methods). Meiotic progeny was obtained and assayed for extracellular Gus,

LacZ and Cts1 activity on indicator plates and by quantitative liquid assays. Based on these

results the progeny was grouped into mutants with defective and intact secretion (Fig. S6).

Sequencing of the 9 candidate genes obtained from comparative genome sequencing revealed

that all progeny with reduced extracellular reporter activity harbored mutations in gene

umag_03776 (Fig. 3, Fig. S6). Strikingly, also strains FB2^{CGL}mut2 and mut3 carried

detrimental mutations in *umag_03776*, leading to synthesis of C-terminally truncated proteins

(Fig. 3C). This strongly suggested that this gene is essential for Cts1 secretion. The

corresponding gene product was subsequently termed Jps1 (jammed in protein secretion

screen 1). While the protein Jps1 has a predicted length of 609 aa, the truncated versions

produced in mutants FB2^{CGL}mut1, mut2 and mut3 only contained 448, 338 and 446 aa,

respectively (Fig. 3C). In essence we identified a crucial factor for Cts1 secretion by genetic

screening.

Jps1 is essential for Cts1 localization and secretion

Jps1   is   annotated   as   hypothetical   protein   with   unknown   function

(https://mycocosm.jgi.doe.gov/Ustma2_1/Ustma2_1.home.html; last accessed 2019/07/25)

and does not contain yet known domains (SMART; http://smart.embl-heidelberg.de/). None

of its homologs in other Basidiomycetes has been characterized so far and Ascomycetes like

S. cerevisiae lack proteins with significant similarities. Thus, to obtain first insights into its

function we initially validated the screen by deleting the respective gene in the background of

laboratory strain AB33 (19) using homologous recombination (AB33jps1Δ). Microscopic

analysis revealed that budding cells of the deletion strain had a normal morphology

comparable to controls like AB33 or the chitinase deletion strain AB33cts1Δ (Fig. 4A) (8).

Chitinase assays showed strongly reduced extracellular activity, confirming the essential role

of Jps1 for Cts1 secretion (Fig. 4B). To analyze the localization of Cts1 in a *jpsΔ* background

strain AB33jps1Δ/CtsG expressing a functional Cts1-Gfp (Cts1G) fusion was generated (Fig.

4C) (15). Microscopic studies revealed that in contrast to its native localization in the

fragmentation zone (Fig. 4D) (8, 9) the Gfp signal for Cts1G now accumulated intracellularly

and at the septa (Fig. 4E,F). Interestingly, in about 56% of the cases, the signal was detected

at the primary septum only, while in the remaining 44% the signal was present at both septa

(Fig. 4G). Cts1G was never observed in the fragmentation zone of the *jps1* deletion strain

(Fig. 4H).

Fig. 4. Jps1 is crucial for unconventional Cts1 secretion. A, Micrographs of yeast-like
growing cells of indicated strains. The *jps1* deletion strain AB33jps1Δ (*jps1Δ*) does not show
any morphological abnormalities. The *cts1* deletion strain AB33cts1Δ (*cts1Δ*) and the
progenitor laboratory strain AB33 are shown for comparison. Scale bars, 10 μm. **B,** *jps1*
deletion in the AB33 verifies its essential function in Cts1 secretion. Extracellular Cts1
activity of AB33, AB33jps1Δ and the control AB33cts1Δ is depicted. The assay was

conducted in three biological replicates. Error bars indicate standard deviation. ***, p value
0.001 (two sample t-test). **C,** Extracellular Cts1 activity of Cts1G expressing strains are
comparable to the progenitor strain AB33, suggesting that the protein is functional. The assay
was conducted in five biological replicates. Error bars indicate standard deviation. ***, p
value 0.001; n.s., not significant (two sample t-test). **D-F,** Localization of Cts1-Gfp (Cts1G)
in AB33 (**D**) and AB33jps1Δ (**E,F**). While Cts1 accumulates in the fragmentation zone of
dividing cells with two septa in AB33 (**D**), it enriches in the cytoplasm and at the septa in the
jps1 deletion strain. Two different scenarios were observed: Either Cts1 was found only at the
primary septum at the mother cell side (**E**) or at both septa (**F**). White arrows depict septa
with Cts1 signal. Scale bars, 10 µm. **G,** Distribution of Cts1G signal at the primary septum
only (PS) and at both septa (PS+SS) of dividing AB33jps1Δ cells with completely assembled
fragmentation zones (AB33Cts1G: 900 cells analyzed; AB33jps1Δ/Cts1G: 1370 cells
analyzed; three biological replicates). **H,** Cts1G is restricted from the fragmentation zone.
The graph depicts the fraction of cells in exponentially growing cultures of indicated strains
with Cts1G accumulation in the fragmentation zone (similar cells analyzed as shown in G).
306
307

Next, a strain expressing Jps1 fused to Gfp (Jps1G) was generated to localize the

protein (AB33Jps1G). Chitinase assays verified the functionality of the fusion protein (Fig.

5A). Intriguingly, microscopic analysis revealed that Jps1G accumulated in the fragmentation

zone of budding cells, similar to Cts1 (Fig. 5B). To investigate if Jps1 localization depends

on chitinase function, Cts1 was deleted in the background of AB33JpsG (AB33cts1Δ/Jps1G).

Interestingly, this did not disturb Jps1 localization (Fig. 5C). This suggests that Cts1 is

dispensable for Jps1 function but not *vice versa*, indicating a unidirectional dependency of

Cts1 on Jps1. To further substantiate the apparent co-localization, both proteins were

differentially tagged in a single strain, expressing Jps1 fused to mCherry and Cts1 fused to

Gfp (AB33Cts1G/Jps1mC). Indeed, both signals completely overlapped in the fragmentation

zone in each observed case (Fig. 5D, E).

In essence, our genetic screen identified a novel essential protein for unconventional

Cts1 secretion. We hypothesize that it might act as an anchoring factor for Cts1 that supports

its local accumulation in the fragmentation zone.

322
323

Fig. 5. Jps1 co-localizes with Cts1 in the fragmentation zone. A, Extracellular Cts1 activity of indicated strains. AB33cts1Δ lacking Cts1 was used as negative control. Both Jps1G and Jps1mC fusions are functional. The assay was conducted in five biological replicates. Error bars indicate standard deviation. ***, p value 0.001; n.s., not significant (two sample t-test). **B,** Localization of Jps1G in AB33. The protein accumulates in the fragmentation zone of dividing cells. Scale bar, 10 µm. **C,** Localization of Jps1G in AB33cts1Δ. Localization of Jps1 in the fragmentation zone is not altered in the absence of Cts1. Scale bar, 10 µm. **D,** Micrographs of strain AB33Cts1G/Jps1mC indicating co-localization of Cts1G and Jps1mC in the fragmentation zone. **E**, Quantification of co-localizing signals (Co-Loc) and not co-localizing signals (No Co-Loc) of strain AB33Cts1G/Jps1mC in the fragmentation zones of 970 cells observed. The experiment was conducted in 3 biological replicates.

Discussion

In this study we identified Jps1, a novel factor essential for unconventional export of

chitinase Cts1 via the fragmentation zone of budding cells. Jps1 was found in a forward

genetic screen. Such genetic screens are powerful tools to identify important players in

unknown pathways (20). The prime example is the screen for components of the conventional

secretion pathway which has been performed in *S. cerevisiae*. Initially, this temperature-

sensitive screen was based on the fact that proteins accumulate in the endomembrane system

of cells in which secretion is disturbed. These dense cells can be separated from cells with

intact secretion by gradient centrifugation (21). The screen was continuously further

developed and finally provided a detailed view on the key components of the canonical

pathway including all stages of protein export (22).

Similarly, the here employed screen proved to be very efficient and powerful: in all

three obtained mutants mutations localized to the same gene (*jps1*). On the one hand this

underlines the quality of the screening procedure. On the other hand, this observation may

also limit the screen in that other factors may be hard to identify in this screening set-up.

Alternatively, the repeated identification of the same mutant could be due to the fact that

there is only one major factor involved in Cts1 secretion. However, we consider that unlikely.

Therefore, in the next step a second copy of Jps1 under its natural promoter will be inserted

into the genome. This will minimize the risk of identifying the gene again in further screening

attempts.

Furthermore, we will streamline identification of responsible mutations. Here, we

used genetic back-crosses via plant infection in combination with a PCR approach to identify

Jps1. For future studies, we will use batch-sequencing of mutants with or without secretion of

the reporters (Fig. S1 steps 10 & 11). Such pooled linkage analysis based on next-generation

sequencing is known to have a great statistical power and thus allows an efficient

identification of underlying mutations (23).

In the first screening round we concentrated on mutants that showed a normal growth

behavior (i.e. colony sizes similar to untreated cells after UV mutagenesis). Mutants that are

impaired in growth are much more complicated to analyze, as discrimination between true

secretory defects and reduced secretion due to poor fitness of the cells is very difficult.

Hence, it is well conceivable that we missed other important factors, especially because

unconventional Cts1 secretion is tightly connected to cytokinesis (9). This also explains why

we did not detect mutants defective in the septation factors Don1 or Don3 which we have

shown to be essential for unconventional Cts1 secretion (9). The cognate deletion mutants

have a cytokinesis defect and grow in tree-like structures (10).

As an alternative application, the screen will be applied to optimize our recently

established protein expression platform (24, 25). Here we use Cts1 as a carrier for valuable

heterologous proteins. Exploiting the unconventional secretion route brings the advantage

that *N*-glycosylation is circumvented and thus, sensitive proteins like bacterial enzymes can

be exported in an active state (16, 26, 27). The screen will be adapted to select for mutants

with enhanced marker secretion to eliminate existing bottlenecks and thus enhance yields

(28). Random mutagenesis screens for hypersecretors were for example key to establishing

industrial production strains like the cellulase producing filamentous fungus *Trichoderma*

reesei strain RutC-30 in which amongst further changes carbon catabolite repression was

eliminated (29). A restriction enzyme mediated insertion (REMI) screen based on a β-

galactosidase reporter has also led to the discovery of a set of mutants showing supersecretion

of the reporter in *Pichia pastoris* (30).

The identified component Jps1 and its orthologs have not been investigated to date

and our bioinformatic analyses did not reveal the presence of any annotated domains or

motives. Interestingly, Jps1 is not conserved in *S. cerevisiae* or other Ascomycetes. This fits

well with the observation that septation in *S. cerevisiae* does not involve formation of a

fragmentation zone, suggesting that molecular details of cell division differ between the two

organisms (7). Jps1 is essential for efficient Cts1 localization and secretion while in turn Cts1

is not needed for Jps1 accumulation in the fragmentation zone. This one-sided dependency

indicates that Jps1 is an important factor for Cts1 secretion but not *vice versa*. The underlying

molecular details and thus the exact role of Jps1 during Cts1 secretion remain to be addressed

by detailed biochemical and cell biological studies in the future. It is also not clear how the

two proteins reach the fragmentation zone. Moving early endosomes enrich in the

fragmentation zone prior to budding and where shown to carry Don1 (10, 31). They are thus

prime candidates for transporting proteins into the small compartment. However, we neither

observed Cts1 nor Jps1 on these motile organelles (data not shown). Based on our results, it is

conceivable that Jps1 functions as an anchoring factor for Cts1, thus supporting its local

accumulation in the fragmentation zone where it degrades remnant chitin together with

conventionally secreted Cts2 to support cytokinesis (Fig. 6).

Fig. 6. Current model for subcellular targeting and unconventional secretion of Cts1 via
anchoring factor Jps1. Cts1 is targeted to the fragmentation zone via an unknown
mechanism. We hypothesize that the newly identified factor Jps1 functions in anchoring Cts1
in the small compartment. Cts1 acts in degrading remnant chitin for detaching mother and
daughter cell in concert with Cts2. Motile early endosomes shuttle bidirectionally through the
cells and transport the septation factor Don1 which is essential for secondary septum
formation. Together with Don3 they localize to the fragmentation zone formed between
mother and daughter cell during cytokinesis.
413
414

In essence, our genetic screen has already proven very efficient and an improved

version will deal as a basis to identify further key components of unconventional secretion

and optimize the connected protein expression platform in the future.

Material and methods

Molecular biology methods

All plasmids (pUMa vectors, see below and Table 1) generated in this study were obtained

using standard molecular biology methods established for *U. maydis* including Golden Gate

cloning (32-35). Oligonucleotides applied for sequencing and cloning are listed in Table 2.

Genomic DNA of strain UM521 (17) was used as template for PCR reactions. All plasmids

were verified by restriction analysis and sequencing. Detailed cloning strategies and vector

maps will be provided upon request.

Table 1. *U. maydis* strains used in this study.

Strains	Relevant genotype/ Resistance	Strain collection no. (UMa)	Plasmids transformed / Resistance	Manipulated locus	Pro-genitor (UMa)	Reference
FB1[a]	*a1b1* (wild type)	51	/	/	Cross of wild type strains UM518 x UM521	(12)
FB2[a]	*a2b2* (wild type)	52	/	/	Cross of wild type strains UM518 x UM521	(12)
AB33	*a2 P$_{nar}$bW2bE1 PhleoR*	133	pAB33 (19)	*B*	UMa52	(19)
FB6a[a]	*a2b1*	55	/	/	Cross of wild type strains UM518 x UM521	(12)

FB6b [a]	*a1b2*	56	/	/	Cross of wild type strains UM518 x UM521	(12)
FB2 lacZ-cts1 (screening progenitor)	*a2b2* *pep4::[P_oma:lacZ:shh:cts1]* NatR	1501	pUMa2373 / lacZ-Cts1_NatR	*umag_04926* (*pep4*) (16)	UMa52	This study.
FB2^{CGL} (screening strain)	*a2b2* *upp1:: [P_omagus:shh:cts1]* HygR *pep4::[P_oma:lacZ:shh:cts1]* NatR	1502	pUMa2374 / *gus-cts1*_HygR	*umag_02178* (*upp1*) (16)	UMa1501 (Screening progenitor)	This study.
FB1^{CGL} (strain used for back-crossings)	*a1b1* *upp1:: [P_omagus:shh:cts1]* HygR *pep4::[P_oma:lacZ:shh:cts1]* NatR	1547	/	/	Derivative of crossing between FB1 and FB2^{CGL}	This study.
FB2 Gus$_{cyt}$	*a2b2* *ip^S* *[P_omagus:shh]ip^R* cbxR	1507	pUMa2335 / *gus*_cbxR	*cbx*	UMa52	This study.
FB2 LacZ$_{cyt}$	*a2b2* *ip^S* *[P_omalacZ:shh]ip^R* cbxR	1508	pUMa2336 / *lacZ_cbxR*	*cbx*	UMa52	This study.
FB2^{CGL} mut1	*a2b2* *upp1:: [P_omagus:shh:cts1]* HygR *pep4::[P_oma:lacZ:shh:cts1]* NatR + UV-induced mutations #24-8	1795	/	UV mutagenized (Fig. 3; Fig. S3).	UMa1502 (screening strain)	This study.
FB2^{CGL} mut2	*a2b2* *upp1:: [P_omagus:shh:cts1]* HygR	1831	/	UV mutagenized (Fig. 3).	UMa1502 (screening strain)	This study.

	pep4::[P_{oma}:lacZ:shh:cts1] NatR + UV-induced mutations #4-13					
FB2^{CGL} mut3	*a2b2 upp1:: [P_{oma}gus:shh:cts1]* HygR *pep4::[P_{oma}:lacZ:shh:cts1]* NatR + UV-induced mutations #3-10	1830	/	UV mutagenized (Fig. 3).	UMa1502 (screening strain)	This study.
AB33 jps1G	*a2 P_{nar}bW2bE1* PhleoR *ip^{S}[P_{jps1}::umag_03776:gfp]ip^{R}* CbxR	2299	pUMa3293 / CbxR	*cbx*	UMa133	This study.
AB33 cts1G	*a2 P_{nar}bW2bE1* PhleoR *umag_10419:gfp* NatR	388	pUMa828 (pCts1G-NatR) (15)	*umag_10419* (*cts1*)	UMa133	(15)
AB33 jps1mC/ Cts1G	*a2 P_{nar}bW2bE1* PhleoR *umag_10419:gfp* NatR *umag_03776:mcherry* HygR	2048	pUMa3034 / HygR	*umag_ 03776* (*jps1*)	UMa388	This study.
AB33 jps1Δ	*a2 P_{nar}bW2bE1* PhleoR	2092	pUMa2775 / HygR	*umag_ 03776* (*jps1*)	UMa133	This study.
AB33cts 1Δ	*a2 P_{nar}bW2bE1 PhleoR umag_10419Δ* HygR	387	pUMa780 (pCts1Δ- HygR) (15)	*umag_10419* (*cts1*)	133	(15)

429　[a] FB1, FB2, FB6a and FB6b are used as tester strains for mating and are derived from the

430　same spore obtained after crossing of the wild type strains UM518 and UM521 (12).

432　**Table 2.** DNA oligonucleotides used in this study.

Designation	Nucleotide sequence (5´- 3´)
oDD691	TGACCGTCAACGCATGGC
oDD692	TCGAGAATCGTGGTACCG

oDD693	TTGCAGCCTACAGGCAGG
oDD694	ACGGTGTGGCTGGAGTGG
oDD695	CATCGGTATGCTTGGCTC
oDD696	AGCGTCGAATTGACCGCC
oDD697	GCTGTCAGAGGCGTTTCA
oDD698	CCCAGAACAGCGCGTTCA
oDD699	CGCGCAAACAAGCCAAGA
oDD729	CGTCGTGCAATGCTGCCG
oDD730	TGTCGAGCCTGCCGGTGG
oDD731	AGACTCGGCTGCAGCAGC
oDD732	AAGCTGGACAGGAGTGGG
oDD733	AGATCGTAGCCGCCTTCG
oDD734	TTGCTCCATCGTTGCCCG
oDD735	CGTGAACGTCGCCCCGTA
oDD736	ATGACCAAATCGCCGCCC
oDD737	TGACGCTCCCTGCTCTCC
oDD815	ATAGCTCTTCCGTGCAATATTGTGCTGTGAAGAGTCTCG
oDD816	ATAGCTCTTCCGGCCGATTTGCAAGTCGTGGGC
oDD819	ATAGCTCTTCCCCTCCGCTCCGCATCCCTCGACC
oDD820	ATAGCTCTTCCGACAATATTCATCTACGACGAGATTGGAGG
oDD824	ATAGCTCTTCCGTGCAATATTTGACAACCTCGTCGGG
oDD825	ATAGCTCTTCCCGAGGATTCCGCATCGATTGGG
oMB190	GATTACAGGATCCATGCCAGGCATCTCC
oMB372	TTAGGCGCGCCATGCCAGGCATCTCC
oMB373	TTAGGGCCCTTAGGATTCCGCATCGATTGGGG
oMB520	CATGAATTCGGATTCCGCATCGATTGGGG
oMB521	TCAGAATTCATGGTGAGCAAGGGCGAGG
oMB522	CATGCGGCCGCCTTACTTGTACAGCTCGTCC
oMF502	ACGACGTTGTAAAACGACGGCCAG
oMF503	TTCACACAGGAAACAGCTATGACC
oRL272	GACCATGGAGACAACTTCGGTCATCTCCGCG
oRL273	GCACTAGTATTGATCGTTCCAGAGCACG
oRL1124	AGAGTTTGATCMTGGCTCAG
oRL1125	GACGGGCRGTGWGTRCA
oRL1982	GGTCTCGCCTGCATTTAAATAGGAACGCCGCGTCGGC
oRL1983	GGTCTCCAGGCCTGTCTTGAAGTGAATGTCGG
oRL1984	GGTCTCCGGCCTGTTGTTCAGTAGCAATGTG
oRL1985	GGTCTCGCTGCATTTAAATCACCCATTCGTGATTCACCAC
oUP65	GGAATTCCATATGGCGAGCCTTGAGGCTGCGTTCC
oUP66	CGGGATCCGATTTGCAAGTCGTGGGCCTTCG

pUMa2373 (pDest-pep4D_Poma-LacZ-SHH-Cts1_NatR) was obtained in a Golden

Gate cloning reaction using two flanking regions obtained by PCR using oligonucleotide

combinations oRL1982 x oRL1983 (upstream flank *pep4* locus) and oRL1984 x oRL1985

(downstream flank *pep4* locus), the destination vector pUMa1476 (32) and pUMa2372.

Storage vector pUMa2372 contained a P_{oma} controlled non-optimized version of the *lacZ*

gene (beta-D-galactosidase, accession NP_414878.1) from *Escherichia coli* Rosetta 2 in

translational fusion to the *cts1* gene (*umag_10419*) via an SHH linker (16) and a

nourseothricin-resistance marker cassette (NatR) flanked by BsaI sites. To generate

pUMa2335 (pRabX1-Poma_Gus-SHH_cbx), pUMa2113 was hydrolyzed with NcoI and NotI

and a Gus-SHH encoding fragment was inserted, replacing the previous insert coding for

Gus-SHH-Cts1. Similary, in pUMa2336 (pRabX1-Poma_LacZ-SHH_cbx) the insert in

pUMa2113 was replaced via NcoI/NotI restriction/ligation by a fusion gene encoding LacZ-

SHH. Both plasmids were integrated in the *ip* locus under the control of the strong,

constitutively active *oma* promoter. pUMa2605 was obtained by hydrolysis of p*cts2*Δ_hyg

(8) with SfiI and replacing the HygR cassette with the G418-resistance cassette (G418R)

from pUMa1057 (pMF1g) (36). For assembly of pUMa3012 (pRabX1-Poma_Gus-SHH-

Jps1) the *jps1* gene was amplified by PCR using oMB372 x oMB373 yielding a 1844 bp

product flanked by AscI and ApaI restriction sites. After hydrolysis with these enzymes the

jps1 gene replaced the *cts1* gene in pUMa2113 (16). For assembly of pUMa3034 flanking

regions were amplified with oDD824 x oDD825 (upstream flank *umag_03776* gene) and

oDD819 x oDD820 (3´region of *umag_03776*) and used for SapI mediated Golden Gate

cloning including destination vector pUMa2074 and the storage vector pUMa3035.

pUMa3035 contained the mCherry gene for translational fusions as well as a HygR cassette.

pUMa3111 (pDest-jps1D_G418R) was generated by replacing the HygR cassette in

pUMa2775 by a G418R cassette from pUMa1057 using flanking SfiI sites. The progenitor

vector pUMa2775 (pDest-jps1D_HygR) was synthetized by SapI mediated Golden Gate

cloning with flanking regions obtained by PCR with oDD815 x oDD816 (upstream flank for

umag_03776) and oDD819 x oDD820 (downstream flank for *umag_03776*), destination

vector pUMa2074 and storage vector pUMa2242 harboring a HygR cassette (9). For

generation of pUMa3293 (pRabX1-Pjps1_Jps1_eGfp_CbxR) a 1952 bp PCR product

obtained with oUP65 x oUP66 (*umag_03776* promoter region) was hydrolyzed with NdeI

and BamHI and inserted into a pRabX1 derivative (pUMa3095) upstream of a *jps1*:*gfp* fusion

gene (13). pUMa3095 was assembled in a three-fragment ligation of a 1849 bp PCR product

of oMB190 x oMB120 (*umag_03776* gene) hydrolyzed with BamHI and EcoRI, a 741 bp

PCR product of oMB521 x oMB522 (*gfp* gene) hydrolyzed with EcoRI and NotI and a

6018 bp fragment of vector pUMa2113 (16) hydrolyzed with BamHI and NotI.

Strains and cultivation conditions

U. maydis strains used in this study were obtained by homologous recombination yielding

genetically stable strains (Table 1). For genome insertion at the *ip* locus, integrative plasmids

were used (13). These plasmids contain an ip^R allele that mediates carboxin resistance (37).

Integrative plasmids were linearized within the ip^R and subsequently used to transform *U.*

maydis protoplasts. Mutants harboring a single copy of the plasmid were obtained via

homologous recombination (33, 34). Gene deletion and translational fusions *in locus* were

performed with plasmids obtained by either the SfiI- or Golden Gate cloning strategy using

plasmids   deposited   in   the   Institutes   plasmid   collection   (32,   34)

http://www.mikrobiologie.hhu.de/ustilago-community.html). Gene insertion at the *pep4*

(*umag_04926*) and *upp1* (*umag_02178*) locus resulted in the deletion of the respective

protease-encoding genes and hence, as a positive side-effect for the screen diminished

proteolytic activity of the strains (16). The corresponding plasmids were obtained by Golden

Gate cloning (32). All strains generated were verified by Southern blot analysis using

digoxygenin labeled probes (Roche). For *ip* insertions, the probe was obtained with the

primer combination oMF502/oMF503 and the template pUMa260 (38). For insertions at the

pep4 or *upp1* locus (16) and other in-locus manipulations, the two flanking regions (upstream

and downstream flanks) were amplified as probes.

U. maydis strains were grown at 28 °C in complete medium (CM) supplemented with

1% (w/v) glucose (CM-glc) (39) or YepsLight (modified from 40). Solid media were

supplemented with 2% (w/v) agar agar. CM-glc plates containing 1% (w/v) charcoal (Sigma

C-9157) were used for mating assays (41).

Generation of a compatible strain by genetic crossings

To enable genetic back-crosses with FB2^{CGL} the reporters were also introduced into the

compatible strain FB1 by genetic crosses. Therefore, wild type strain FB1 was crossed with

screening strain FB2^{CGL} (mating type *a2b2*) using plant infections (see below) to obtain

meiotic progeny. These were tested for their ability to mate with FB2. Compatible mating

was screened on CM-glc plates containing 1% (w/v) charcoal (Fig. S4) on which strains

harboring different alleles of both mating type loci form fuzzy colonies while strains which

do not mate grow in smooth colonies. First, progeny was screened for the presence of the two

artificial reporters Gus-Cts1 and LacZ-Cts1. Positive candidates were then tested in mating

experiments with tester strains for induction of the fuzzy phenotype in FB2 crosses.

Mixing experiments to distinguish intra- and extracellular reporter activities

To distinguish intra- and extracellular LacZ activity defined cell amounts were mixed on

CM-glc/X-Gal plates containing 1% (w/v) glucose and X-Gal (5-bromo-4-chloro-3-indolyl-

β-D-galactopyranoside; 20 mg/ml in DMSO, f.c. 60 mg/L in CM-glc). Screening strain

FB2^{CGL} and the AB33LacZ$_{cyt}$ control were grown in liquid CM-glc medium until logarithmic

phase. Cells were harvested, washed in PBS (1x, pH 7.2) and adjusted to an OD$_{600}$ of 1.0. A

10^{-4} serial dilution was prepared in PBS. The diluted suspensions of both strains were mixed

in defined ratios and plated in a total volume of 150 µl on CM-glc/X-Gal plates. After

incubation for 4 days at 28 °C protected from light, growth and conversion of substrate was

photographed. For illumination a Ledgo CN-B150 LED On-Camera Light was used.

Gus/LacZ activity plate and membrane assays

Gus and LacZ activity were tested by indicator plate assays using CM plates containing 1%

(w/v) glucose (CM-glc) and the respective chromogenic substrate X-Gluc (5-bromo-4-chloro-

3-indolyl-beta-d-glucuronic acid; 0.5 mg/ml in DMSO) or X-Gal (20 mg/ml in DMSO),

respectively. Tested strains were grown in CM-glc for 16 hours. After adjusting the cultures

to an OD_{600} of 1.0 in sterile PBS, 10 µl suspension were spotted on CM-glc plates and

incubated at 28 °C for 2 days. Intracellular Gus and LacZ activity was visualized by placing a

nitrocellulose membrane (Amersham ^{TM} Protran TM 0.45 µM NC, GE Healthcare Life

Sciences) on top for 24 h at 28 °C. The membrane was then removed and treated with liquid

nitrogen for 3 min for cell lysis. Subsequently, it was soaked in X-Gluc buffer (25 µg X-

Gluc/ml, solved in DMSO, 5 mM sodium phosphate buffer pH 7.0, 14 µM β-

mercaptoethanol, 0.4 mM EDTA, 0.0021% (v/v) lauroyl-sarcosin, 0.002% (v/v) Triton X-

100, 0.1 mg/ml (w/v) BSA) or X-Gal Buffer (1 mg X-Gal/ml, solved in DMSO, 15 mM

Sodium phosphate buffer pH 7, 5 mM KCl, 0.5 mM $MgSO_4$, 34 mM β-mercaptoethanol) for

Gus and LacZ activity, respectively, and incubated at 37 °C for 18 h.

UV mutagenesis

For UV mutagenesis a 20 ml YepsLight pre-culture of screening strain $FB2^{CGL}$ was

inoculated from a fresh plate and incubated overnight (200 rpm, 28 °C). In the morning, the

culture was diluted to an OD_{600} of 0.1 in 20 ml (200 rpm, 28 °C). The culture was incubated

until it reached an OD_{600} of 0.5. Subsequently it was diluted stepwise to an OD_{600} of 0.00125

(1: 400) in 20 ml YepsLight. 150 µl of the 1:400 dilutions were spread evenly onto CM-glc

screening plates containing 10 µg/ml X-Gal. The dried plates were exposed to UV irradiation

(30 mJ/cm^2) using a Stratalinker device (Stratagene). Plate lids were removed during

exposition. Subsequently, plates were incubated for 2 to 3 days at 28 °C until single colonies

were grown.

Screening for diminished reporter secretion

Clones that showed reduced or absent LacZ activity (i.e., colorless appearing colonies) after

UV mutagenesis on CM-glc screening plates containing X-Gal (see above) were patched on

plates containing X-Gluc to additionally assay for extracellular Gus activity. Plates were

prepared by spreading 100 µl X-Gluc solution (100 mg/ml stock in DMSO) on CM-glc

plates. Plates were incubated for 2 to 3 days on 28 °C. Colorless colonies were patched again

on X-Gluc and X-Gal plates simultaneously, this time streaking out larger areas of about 0.5

x 0.5 cm. During the procedure, control strains producing intracellular LacZ or Gus (FB2

$LacZ_{cyt}$ and FB2 Gus_{cyt}, respectively), the non-mutagenized screening strain (FB2^{CGL}) and the

precursor strain lacking any reporters (FB2) were handled in parallel to verify the results

(Table 1).

Generation of cell extracts and supernatants

Strains were inoculated in 20 ml CM-glc and incubated at 28 °C overnight. Next morning, the

culture was used to inoculate a new culture of 70 ml CM-glc with a starting OD_{600} of 0.05. To

detect potential growth defects, growth of the culture was followed by determining the OD_{600}

every hour for at least 8 to 10 hours. 2 ml aliquots of supernatants for Gus/LacZ assays and

whole cells for Cts1 assays were harvested at OD_{600} of 0.3. Once the culture reached an

OD_{600} of 0.7 50 ml were harvested (5 min, 3000 rpm, Falcon tubes, 4 °C). The supernatant

was transferred to a new tube and stored on 4 °C. The cell pellet was used to prepare native

cell extracts used for the Cts1, Gus and LacZ assay (modified from 14). To this end the cell

pellet was resuspended in 2 ml ice-cold native extraction buffer (1 mM

phenylmethylsulfonylfluorid (PMSF), 2.5 mM benzamidine hydrochloride hydrate, 1 µM

pepstatin; 100 µL protease inhibitor cocktail 50×; dissolve in PBS pH 7.4). The suspension

was then frozen in pre-chilled metal pots (25 ml, Retsch) using liquid nitrogen. Cells were

ruptured at 4 °C using the Retsch mill (10 min, 30 Hz) and then the metal pots were thawed

at 4 °C for 1 hour. The cell extracts were transferred to a reaction tube and centrifuge for 30

min (4 °C, 13000 rpm, benchtop centrifuge). Bradford assays were conducted to determine

the protein concentrations in the samples (42).

574   Quantitative determination of Gus and LacZ activity

Importantly, quantitative Gus, LacZ and Cts1 assays were conducted from a single culture

(see below for Cts1 assay). Quantitative Gus and LacZ liquid assays were based on the

chromogenic and fluorescent substrates ONPG for the liquid LacZ activity assay (o-

nitrophenyl-β-D-galactopyranoside) and MUG for the liquid Gus activity assay (4-

methylumbelliferyl-ß-D-glucuronide trihydrate; BioWorld, 30350000-2 (714331),

respectively (13, 14). For both the ONPG and the MUG liquid assays (modified from 13, 14),

activity was determined in native cell extracts and in the cell-free culture supernatant of

candidate mutants in comparison to control strains (8, 13, 15).

The Gus and LacZ assays with cell extracts and cell-free supernatants were conducted

according to slightly modified published protocols (13, 43). To this end, native cell extracts

were adjusted to a total protein concentration of 100 µg/ml using PBS buffer. 10 µl of native

cell extracts were then mixed in a black 96-well plate (96 Well, PS, F-Bottom, µCLEAR,

black, CELLSTAR) with 90 µl of Gus- or Z-buffer and 100 µl of the respective substrate

solution (Gus: 2 mM MUG, 1/50 vol. bovine serum albumin fraction V (BSA) in 1x Gus

buffer; LacZ: 1 mg/ml ONPG in 2x Z-buffer). For supernatant measurements, 100 µl cell-free

supernatants were mixed with 100 µl of the respective substrate solution. 2x Gus buffer (14)

was used for the Gus assay and 2x Z-buffer (80 mM Na_2HPO_4, 120 mM $NaH_2PO_4*H_2O$, 20

mM KCl, 2 mM $Mg_SO_4*7H_2O$; adjust to pH 7; add 100 mM β-mercaptoethanol freshly) was

used for the LacZ assay. The assays were conducted in the Tecan device (Tecan Group Ltd.,

Männedorf, Switzerland) for 1 h at 37 °C with measurements every 10 min

(excitation/emission wavelengths: 365/465 nm for Gus activity; OD_{420} for LacZ activity). A

fixed gain of 150 was used for cell extract measurements and fixed values of 60 and 100 for

Gus and Cts1 activity assays of culture supernatants, respectively.

For data evaluation the slope of the connected data points was determined. Values for

the screening strain FB2^{CGL} were set to 100% to judge the activities in the mutants.

Quantitative determination of Cts1 activity

The fluorescent substrate MUC was applied for the Cts1 liquid assay (4-methylumbelliferyl

β-D-N,N′,N″-triacetylchitotrioside hydrate; M5639 Sigma-Aldrich) (13, 15). Strain AB33

cts1Δ (UMa387) (15) carrying a *cts1* deletion dealt as negative control for the Cts1 assays.

For the MUC assay whole cells were subjected to the assay after washing to detect Cts1

activity at the cell surface. For intracellular activities, cell extracts were used (see above).

The Cts1 activity liquid assay with whole cells or 10 µg of native cell extracts was

conducted according to published protocols with minor changes (13, 15). Once the culture

had reached an OD_{600} of 0.3 a 2 ml sample was taken and cells were harvested by

centrifugation (3 min, 8000 rpm, bench-top centrifuge). Cells were resuspended in 1 ml KHM

buffer (110 mM potassium acetate, 20 mM HEPES, 2 mM $MgCl_2$) (13, 15), the OD_{600} was

documented and the suspension was subjected to the MUC assay. A MUC working solution

was prepared form a stock solution (2 mg/ml MUC in DMSO) by diluting it 1:10 with KHM

buffer (protect from light, store at 4 °C). Black 96-well plates (96 Well, PS, F-Bottom,

μCLEAR, black, CELLSTAR) were used for the assay. 70 µl of working solution were

mixed with 30 µl of the cell suspension in one well. Activities for each strain were

determined in triplicates. The plates were sealed with parafilm and incubated in the dark for 1

h at 37 °C. The reaction was then stopped by adding 200 µl 1 M Na_2CO_3 and relative

fluorescence units were determined in a plate reader at excitation and emission wave length

of 360/450 nm, respectively, at 37 °C with a fixed gain of 100 (Tecan Reader).

Plant infection and genetic back-crosses

Compatible *U. maydis* strains were subjected to genetic crosses on corn plants. Strain FB1[CGL]

was obtained by genetic crossing of FB1 (mating type *a1b1*) with FB2[CGL] (mating type

a2b2). Furthermore, mutagenized strains (derived from the FB2[CGL] strain background; mating

type *a2b2*) which showed strongly reduced Cts1 secretion assayed by all three reporters were

subjected to back-crosses with the compatible wild type strain FB1[CGL] (mating type *a1b1*).

For infection of the host plant *Z. mays* (Early Golden Bantam), strains were grown to an

OD_{600} of 0.8 in CM-glc, washed three times with H_2O, and resuspended to an OD_{600} of 1 in

H_2O. For infection, compatible strains were then mixed in a 1:1 ratio. The cell suspension

was injected into seven day-old maize seedlings. Virulence of strain crosses was quantified

using established pathogenicity assays (17). Therefore, 7 days post infection plants were

scored for symptom formation according to the following categories: (1) no symptoms, (2)

chlorosis, (3) anthocyanin accumulation, (4) small tumors (<1 mm), (5) medium tumors (>1

mm), and heavy tumors associated with bending of stem. For spore collection, mature tumor

material was harvested two to three weeks after infection and dried at 37 °C for about 7 days.

Spore germination and analysis of progeny

Spore germination and analysis was conducted according to published protocols (44). To

germinate spores for progeny analysis after genetic back-crossing of FB1 x FB2^{CGL} or

FB1^{CGL} x FB2^{CGL}mut1 dried tumor material was homogenized in a mortar, treated with 2 ml

of a solution of 3.0% (w/v) copper sulfate for 15 min and washed twice with 1 ml sterile H_2O.

The spores were then resuspended in 500 µl sterile water. Prior to plating, the spore solutions

were supplemented with ampicillin and tetracyclin to avoid bacterial contaminations (final

concentrations: 600 µg/ml ampicillin; 150 µg/ml tetracyclin). Then, 200 µl of 1:1, 1:10 and

1:100 dilutions were spread on CM-glc plates, and incubated for 2 d at 28 °C. Resulting

colonies were singled out again on CM-glc plates to guarantee that each colony results from

one clone. To identify FB1^{CGL} after crossing of FB1 x FB2^{CGL} progeny was assayed on

charcoal plates for their mating types (see below) and on X-Gal and X-Gluc plates for the

presence of the two reporters Gus and LacZ. To analyze FB1^{CGL} and FB2^{CGL}mut1 again

indicator plates containing X-Gal and X-Gluc were used to pre-sort the cells into secretion

competent and deficient clones. Candidates were further tested using liquid assays for all

three reporters.

Mating assay

For mating assays, cells were grown in CM medium to an OD_{600} of 1.0 and washed once with

sterile H_2O. Washed cells were adjusted to an OD_{600} of 3 in sterile H_2O. Indicated strains

were pre-mixed in equal amounts and then co-spotted on CM-glc plates containing 1% (w/v)

charcoal. Plates were incubated at 28 °C for 24 h. Tester strains were used as controls (Table

2).

Genome sequencing and assembly

Genomic DNA extraction was performed according to published protocols (35), two separate

preparations were combined, and the DNA concentration was adjusted to 100 ng/µl using

TE-RNase in approximately 500 µl final volume. gDNA quality was verified by PCR

reactions using primers specific for the bacterial 16sRNA gene (oRL1124 x oRL1125) and

the intrinsic gene *uml2* (*umag_01422*; oRL272 x oRL273). For genome sequencing, DNA

libraries were generated using the Nextera XT Kit (Illumina) according to manufacturer's

instructions. Sequencing (v3 chemistry) was performed with a MiSeq sequencer (Illumina) at

the Genomics Service Unit (LMU Biocenter). Obtained reads were quality trimmed and

filtered with trimmomatic version 0.30 and fasta toolkit version 0.13.2. Sequence assemblies

with passed single and paired reads were performed using CLC Genomics Workbench 8.0

(QIAGEN). The genome sequence of UM521 (*Ustilago maydis* 521;

http://www.ncbi.nlm.nih.gov/nuccore/AACP00000000.2; accessed on 07/27/2016) was used

as template for read assembly. Genomic DNA of screening strain FB2^{CGL} was sequenced as

reference.

Identification of genomic mutations

For identification of the underlying mutation in progeny of FB1^{CGL} and FB2^{CGL}mut1, PCR

products were generated from gDNA of progeny clones #3 and #5 which showed no Cts1

activity and no blue halos on X-Gal and X-Gluc indicator plates. To this end, the following

primer combinations were used to amplify the indicated genes (Table 2): *umag_00493*:

oDD691 x oDD729; *umag_06269*: oDD692 x oDD730; *umag_02631*: oDD693 x oDD731;

umag_03776: oDD694 x oDD732; *umag_04298*: oDD695 x oDD733; *umag_05386*: oDD696

x oDD734; *umag_04385*: oDD697 x oDD735; *umag_04494*: oDD698 x oDD736 and

umag_11876: oDD699 x oDD737. gDNA was obtained like described (35). Additional

sequencing was conducted for *umag_06269* and *umag_03776* using progeny #9, #26, #31 and

#39 (not shown). PCR products were sequenced and analyzed for the presence or absence of

the mutations identified in genome sequence alignments of FB2^{CGL} and FB2^{CGL}mut1.

Protein precipitation from culture supernatants

Secreted proteins were enriched from supernatant samples using trichloric acid (TCA)

precipitation. Therefore, culture supernatants were supplemented with 10% (w/v) TCA and

incubated overnight on 4 °C. After washing twice in -20 °C acetone the protein pellets were

resuspended in minimal amounts of 3x Laemmli-Buffer (45) and the pH was eventually

neutralized with 1 M NaOH. For SDS-Page analysis the samples were first boiled for 10 min

and then centrifuged (22,000 x g, 5 min, room temperature).

SDS-Page and Western blot analysis

Boiled protein samples were separated by SDS-Page using 10% (w/v) acrylamide gels.

Subsequently, proteins were blotted to methanol-activated PVDF membranes. The analyzed

proteins contained an SHH-tag (16) and were detected using anti-HA (Sigma-Aldrich, USA)

antibodies and anti-mouse IgG-HRP (Promega, USA) conjugates as primary and secondary

antibodies, respectively. HRP activity was detected using AceGlow Western blotting

detection reagent (PeqLab, Germany) and a LAS4000 chemiluminescence imager (GE

LifeScience, Germany).

Microscopy, image processing and staining procedures

Microscopic analysis was performed with a wide-field microscope from Visitron Systems

(Munich, Germany), Zeiss (Oberkochen, Germany) Axio Imager M1 equipped with a Spot

Pursuit CCD camera (Diagnostic Instruments, Sterling Heights, MI) and objective lenses Plan

Neofluar (40x, NA 1.3) and Plan Neofluar (63x, NA 1.25). Fluorescence proteins were

excited with an HXP metal halide lamp (LEj, Jena, Germany) in combination with filter sets

for Gfp (ET470/40BP, ET495LP, ET525/50BP), mCherry (ET560/40BP, ET585LP,

ET630/75BP, Chroma, Bellow Falls, VT), and DAPI (HC387/11BP, BS409LP, HC

447/60BP; AHF Analysentechnik, Tübingen, Germany). The system was operated with the

software MetaMorph (Molecular Devices, version 7, Sunnyvale, CA). Image processing

including adjustments of brightness and contrast was also conducted with this software. To

visualize fungal cell walls and septa 1 ml of cell culture was stained with calcofluor white

(1 µg/ml) before microscopy.

Author contributions

J.S. designed, conducted and evaluated the genetic screen with support of M.R., K.H. and A.G., M.R. characterized Jps1 and prepared micrographs, quantifications and enzyme assays. A.B. performed genome sequencing. K.S. prepared the manuscript with input of all co-authors, directed the project and acquired funding.

Acknowledgements

We acknowledge Dr. M. Feldbrügge for continuous support and valuable discussion. We thank B. Axler and U. Meyer for excellent technical support of the project. T.E. Hyland and L. Mielke contributed to the project in the framework of an internship and M. Tulinski in his Master project.

Funding

This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Projektnummer 267205415 – SFB 1208 (M.F., K.S., M.R.). The scientific activities of the Bioeconomy Science Center were financially supported by the Ministry of Culture and Science within the framework of the NRW Strategieprojekt BioSC (No. 313/323-400-002 13).

References

1. Sommer MS, Schleiff E. Protein targeting and transport as a necessary consequence of increased cellular complexity. Cold Spring Harb Perspect Biol. 2014;6(8).

2. Malhotra V. Unconventional protein secretion: an evolving mechanism. EMBO J. 2013;32(12): 1660-1664.

3. Rabouille C. Pathways of Unconventional Protein Secretion. Trends Cell Biol. 2017;27(3): 230-240.

4. Debaisieux S, Rayne F, Yezid H, Beaumelle B. The ins and outs of HIV-1 Tat. Traffic. 2012;13(3): 355-363.

5. Rayne F, Debaisieux S, Bonhoure A, Beaumelle B. HIV-1 Tat is unconventionally secreted through the plasma membrane. Cell Biol Int. 2010;34(4): 409-413.

6. Nowag H, Munz C. Diverting autophagic membranes for exocytosis. Autophagy. 2015;11(2): 425-427.

7. Reindl M, Hänsch S, Weidtkamp-Peters S, Schipper K. A Potential Lock-Type Mechanism for Unconventional Secretion in Fungi. Int J Mol Sci. 2019;20(3).

8. Langner T, Öztürk M, Hartmann S, Cord-Landwehr S, Moerschbacher B, Walton JD, et al. Chitinases Are Essential for Cell Separation in *Ustilago maydis*. Eukaryot Cell. 2015;14(9): 846-857.

9. Aschenbroich J, Hussnaetter KP, Stoffels P, Langner T, Zander S, Sandrock B, et al. The germinal centre kinase Don3 is crucial for unconventional secretion of chitinase Cts1 in *Ustilago maydis*. Biochim Biophys Acta Proteins Proteom. 2019;1867(12): 140154.

780   10.   Weinzierl G, Leveleki L, Hassel A, Kost G, Wanner G, Bölker M. Regulation of cell

781   separation in the dimorphic fungus *Ustilago maydis*. Mol Microbiol. 2002;45(1): 219-

782   231.

783   11.   Böhmer C, Ripp C, Bölker M. The germinal centre kinase Don3 triggers the dynamic

784   rearrangement of higher-order septin structures during cytokinesis in *Ustilago maydis*.

785   Molecular microbiology. 2009;74(6): 1484-1496.

786   12.   Banuett F, Herskowitz I. Different *a* alleles of *Ustilago maydis* are necessary for

787   maintenance of filamentous growth but not for meiosis. PNAS. 1989;86(15): 5878-

788   5882.

789   13.   Stock J, Sarkari P, Kreibich S, Brefort T, Feldbrügge M, Schipper K. Applying

790   unconventional secretion of the endochitinase Cts1 to export heterologous proteins in

791   *Ustilago maydis*. J Biotechnol. 2012;161(2): 80-91.

792   14.   Stock J, Terfrüchte M, Schipper K. A Reporter System to Study Unconventional

793   Secretion of Proteins Avoiding *N*-Glycosylation in *Ustilago maydis*. Unconventional

794   Protein Secretion: Methods and Protocols Springer Protocols. 2016;1459: 149-160.

795   15.   Koepke J, Kaffarnik F, Haag C, Zarnack K, Luscombe NM, Konig J, et al. The RNA-

796   binding protein Rrm4 is essential for efficient secretion of endochitinase Cts1. Mol Cell

797   Proteomics. 2011;10(12): M111 011213.

798   16.   Sarkari P, Reindl M, Stock J, Müller O, Kahmann R, Feldbrügge M, et al. Improved

799   expression of single-chain antibodies in *Ustilago maydis*. J Biotechnol. 2014;191: 165-

800   175.

801   17.   Kämper J, Kahmann R, Bölker M, Ma LJ, Brefort T, Saville BJ, et al. Insights from the

802   genome of the biotrophic fungal plant pathogen *Ustilago maydis*. Nature.

803   2006;444(7115): 97-101.

804   18.   Pfeifer GP, You YH, Besaratinia A. Mutations induced by ultraviolet light. Mutation

805   Res. 2005;571(1-2): 19-31.

806   19.   Brachmann A, Weinzierl G, Kämper J, Kahmann R. Identification of genes in the

807   bW/bE regulatory cascade in *Ustilago maydis*. Mol Microbiol. 2001;42(4): 1047-1063.

808   20.   Forsburg SL. The art and design of genetic screens: yeast. Nat Rev Genet. 2001;2(9):

809   659-68.

810   21.   Novick P, Field C, Schekman R. Identification of 23 complementation groups required

811   for post-translational events in the yeast secretory pathway. Cell. 1980;21(1): 205-215.

812   22.   Mellman I, Emr SD. A Nobel Prize for membrane traffic: vesicles find their journey's

813   end. J Cell Biol. 2013;203(4): 559-561.

814   23.   Birkeland SR, Jin N, Ozdemir AC, Lyons RH, Jr., Weisman LS, Wilson TE. Discovery

815   of mutations in *Saccharomyces cerevisiae* by pooled linkage analysis and whole-

816   genome sequencing. Genetics. 2010;186(4): 1127-1137.

817   24.   Feldbrügge M, Kellner R, Schipper K. The biotechnological use and potential of plant

818   pathogenic smut fungi. Appl Microbiol Biotechnol. 2013;97(8): 3253-3265.

819   25.   Sarkari P, Feldbrügge M, Schipper K. The corn smut fungus *Ustilago maydis* as an

820   alternative expression system for biopharmaceuticals. In: Fungal Biology Gene

821   expression systems in fungi: advancements and applications. Edited by M Schmoll and

822   C Dattenböck 2016;Springer Book Series: 183-200.

823   26.   Terfrüchte M, Reindl M, Jankowski S, Sarkari P, Feldbrügge M, Schipper K. Applying

824   Unconventional Secretion in *Ustilago maydis* for the Export of Functional Nanobodies.

825   Int J Mol Sci. 2017;18(5).

826   27.   Stoffels P, Müller MJ, Stachurski S, Terfrüchte M, Schröder S, Ihling N, et al.

827   Complementing the intrinsic repertoire of *Ustilago maydis* for degradation of the pectin

828   backbone polygalacturonic acid. J Biotechnol. 2019;307: 148-163.

28.   Terfrüchte M, Wewetzer S, Sarkari P, Stollewerk D, Franz-Wachtel M, Macek B, et al.

Tackling destructive proteolysis of unconventionally secreted heterologous proteins in

Ustilago maydis. J. Biotechnol. 2018;284: 37-51.

29.   Peterson R, Nevalainen H. *Trichoderma reesei* RUT-C30 - thirty years of strain

improvement. Microbiology. 2012;158(Pt 1): 58-68.

30.   Larsen S, Weaver J, de Sa Campos K, Bulahan R, Nguyen J, Grove H, et al. Mutant

strains of *Pichia pastoris* with enhanced secretion of recombinant proteins.

Biotechnology letters. 2013;35(11): 1925-1935.

31.   Schink KO, Bölker M. Coordination of cytokinesis and cell separation by endosomal

targeting of a Cdc42-specific guanine nucleotide exchange factor in *Ustilago maydis*.

Mol Biol Cell. 2009;20(3): 1081-1088.

32.   Terfrüchte M, Joehnk B, Fajardo-Somera R, Braus GH, Riquelme M, Schipper K, et al.

Establishing a versatile Golden Gate cloning system for genetic engineering in fungi.

Fungal Gen Biol. 2014;62: 1-10.

33.   Kämper J. A PCR-based system for highly efficient generation of gene replacement

mutants in *Ustilago maydis*. Mol Gen Genomics. 2004;271(1): 103-110.

34.   Brachmann A, König J, Julius C, Feldbrügge M. A reverse genetic approach for

generating gene replacement mutants in *Ustilago maydis*. Mol Gen Genomics.

2004;272(2): 216-126.

35.   Bösch K, Frantzeskakis L, Vranes M, Kämper J, Schipper K, Göhre V. Genetic

Manipulation of the Plant Pathogen *Ustilago maydis* to Study Fungal Biology and Plant

Microbe Interactions. J Vis Exp. 2016(115).

36.   Baumann S, Pohlmann T, Jungbluth M, Brachmann A, Feldbrügge M. Kinesin-3 and

dynein mediate microtubule-dependent co-transport of mRNPs and endosomes. J Cell

Sci. 2012;125(Pt 11): 2740-2752.

37.   Keon JP, White GA, Hargreaves JA. Isolation, characterization and sequence of a gene

855   conferring resistance to the systemic fungicide carboxin from the maize smut pathogen,

856   *Ustilago maydis*. Curr Gen. 1991;19(6): 475-481.

857   38.   Loubradou G, Brachmann A, Feldbrügge M, Kahmann R. A homologue of the

858   transcriptional repressor Ssn6p antagonizes cAMP signalling in *Ustilago maydis*. Mol

859   Microbiol. 2001;40(3): 719-730.

860   39.   Holliday R. *Ustilago maydis*. Handbook of Genetics. 1. New York, NY: Plenum Press;

861   1974. p. 575-95.

862   40.   Tsukuda T, Carleton S, Fotheringham S, Holloman WK. Isolation and characterization

863   of an autonomously replicating sequence from *Ustilago maydis*. Mol Cell Biol.

864   1988;8(9): 3703-3709.

865   41.   Hartmann HA, Kahmann R, Bölker M. The pheromone response factor coordinates

866   filamentous growth and pathogenicity in *Ustilago maydis.* EMBO. 1996;15(7): 1632-

867   1641.

868   42.   Bradford MM. A rapid and sensitive method for the quantitation of microgram

869   quantities of protein utilizing the principle of protein-dye binding. Anal Biochm.

870   1976;72: 248-254.

871   43.   Miller GL. Dinitrosalicylic Acid Reagent for Determination of Reducing Sugar. Anal

872   Chem. 1959;31(3): 426-428.

873   44.   Eichhorn H, Lessing F, Winterberg B, Schirawski J, Kämper J, Müller P, et al. A

874   ferroxidation/permeation iron uptake system is required for virulence in *Ustilago*

875   *maydis*. Plant Cell. 2006;18(11): 3332-3345.

876   45.   Laemmli UK. Cleavage of structural proteins during the assembly of the head of

877   bacteriophage T4. Nature. 1970;227(5259): 680-685.

878 **Supplementary**

880 The Supplementary contains Figures S1-S7.

Fig. S1. Detailed step-by-step description of the forward genetic screen. All steps
performed to identify factors for unconventional secretion of chitinase Cts1 are indicated in
detail. First of all, cells were mutagenized by UV light (1). Next, plate assays were performed
(2-4). Initial screening of the mutagenized cells was conducted on X-Gal containing plates
(2). White colonies lacking extracellular LacZ activity were then subjected to further plate
assays on both X-Gal and X-Gluc (3, 4). Colonies that stayed white on both substrates were
analyzed in detail using quantitative liquid assays with substrates for all three unconventional
secretion reporters (Cts1, LacZ-Cts1, Gus-Cts1) (5,6). Both intra- (5) and extracellular
activities (6) were determined. Genome sequencing and alignment with the progenitor strain
FB2^{CGL} revealed multiple mutations (candidate loci) likely resulting from UV treatment (7).
Genetic back-crosses with a non-mutagenized FB1 derivative harboring all three reporters

(FB1^{CGL}) were performed and led to progeny showing defective unconventional secretion
while harboring cytoplasmic activity for all three reporters (8). Amplification and sequencing
of the candidate loci identified in the alignment identify the responsible mutation (9). In
future studies, progeny of the back-crosses could also be pooled according to the absence or
presence of unconventional secretion. Genome sequencing of the two pools should also
reveal the responsible mutation (10,11).

Fig. S2. Mutants impaired in Cts1 secretion grow normal and produce similar levels of Gus-Cts1 and LacZ-Cts1 as the screening strain FB2^{CGL}**. A**, Growth curves of FB2^{CGL} and the indicated UV mutants. Growth was followed by determining the optical density at 600 nm for eight hours. All strains duplicate with comparable rates. **B**, Western blot detecting Gus-Cts1 and LacZ-Cts1 levels in FB2^{CGL} and the indicated UV mutants. Antibodies directed against the HA-tag in the SHH linker of the fusion proteins were used for detection. Asterisks and rhombs depict the expected sizes of the fusion proteins LacZ-Cts1 and Gus-Cts1, respectively. A strain lacking the reporters was used as negative control (AB33 *cts1Δ*).

Fig. S3. SNPs identified in genome sequence comparisons between screening strain FB2^CGL and mutant candidate FB2^CGLmut1. The schematic representation shows all 23 chromosomes. Bars depict 32 base changes detected in FB2^CGLmut1 compared to its progenitor FB2^CGL. Red bars indicate 12 mutations which go along with amino acid replacements in 9 ORFs and are thus candidates likely responsible for the observed defective secretion. Unexpected base transitions are indicated in grey letters while black letters depict transitions typically observed after UV treatment. Figure dimensions are not to scale.

Fig. S4. Mating assay to identify FB1$^{\text{CGL}}$ with *a1b1* mating type harboring the reporter LacZ-Cts1 and Gus-Cts1 using charcoal plates. Mating of compatible strains was observed on CM-charcoal plates. Fuzzy colonies indicate mating while smooth colonies show incompatibility. FB1 (mating type *a1b1*), FB2 (mating type *a2b2*), FB6a (mating type *a2b1*) and FB6b (mating type *a1b2*) are tester strains with known genotypes. Progeny carrying the two reporters Gus-Cts1, LacZ-Cts1 was obtained by crossing the screening strain FB2$^{\text{CGL}}$ with wild type strain FB1 and tested for mating to identify an *a1b1* clone. One clone of the two displayed clones has an *a1b1* mating type because it shows a fuzzy colony with FB2 but not with FB6a (strain termed FB1$^{\text{CGL}}$). This strain was used for genetic back-crosses with mutagenized FB2$^{\text{CGL}}$ during the screen.

981 **Fig. S5. Plant infections with compatible FB1 and FB2 derivatives.** Maize seedlings were
982 inoculated with the indicated compatible strains and disease symptoms scored 7 days post
983 infection. The infection was performed once with 40 plants each. All tested strain
984 combinations were pathogenic and showed symptoms comparable to the wild type cross FB1
985 (*a1b1*) x FB2 (*a2b2*).

998 **A**

1000 **B**

1001
1002
1003 **C**

	Progeny FB1CGL x FB2CGL (# artificially numbered)
Progeny with intact Cts1 secretion	#1, #2, #4, #6, #7, #8, #10, #11, #12, #13, #15, #16, #17, #18, #20, #22, #24
Progeny with defective Cts1 secretion	#3, #5, #9, #14, #19, #21

D

Gene	#3	#5	#9
UMAG_00493			
UMAG_06269			
UMAG_02631			
UMAG_03776			
UMAG_04298			
UMAG_05386			
UMAG_04385			
UMAG_04494			
UMAG_11876			

1004

1005
Fig. S6. Extracellular Cts1 activity of progeny from the genetic cross of FB1CGL and
FB2CGLmut1. A, Back-crossing strategy to allow for identification of responsible mutations
in strain FB2CGLmut1. Spores obtained after crossing FB1CGL and FB2CGLmut1 were
germinated and singled out. Single colonies were sorted into two groups (extracellular
reporter activity: yes/no) by assaying extracellular reporter activities on X-Gal and X-Gluc
plates and by liquid Cts1 assays with whole cells. **B,** Liquid Gus, LacZ and Cts1 assays to
sort progeny obtained from FB1CGL and FB2CGLmut1 crosses. Both extracellular (ex, filled
columns) and intracellular (int, open columns) activities were determined. Numbers (#)
indicate the different clones obtained from germinated spores. The progenitor strains FB2CGL,

FB2^CGL mut1 and FB1^CGL were used as references for normal and strongly diminished
unconventional secretion. Clones with strongly reduced extracellular activities for all three
reporters are listed in C and were used for sequencing. **C**, Progeny groups obtained after
genetic back-crossing based on the results shown in B. **D**, Sequencing of diagnostic PCR
products was conducted on progeny which did not show extracellular reporter activity using
primers that encompass the 12 potential mutations leading to aa exchanges in 9 ORFs.
Sequencing revealed mutagenized *umag_03776* to be responsible for the reduction in
unconventional secretion. Red: candidate locus mutagenized; green: candidate locus
wildtypic.

Figure S7. Aa alignment of Jps1 wild type protein and three mutant versions obtained in the UV mutagenesis for candidates with diminished Cts1 secretion. Red background color indicates aa changes in the mutants in comparison to the native protein (upper line). Asterisks depict stop codons. Mutations result in the production of C-terminally truncated proteins (compare Fig. 3). Numbers show base pair counts.