Mechanisms underlying the cooperation between loss of epithelial polarity and

Notch signaling during neoplastic growth in *Drosophila*

Rémi Logeay[1], Charles Géminard[1], Patrice Lassus[2], Diala Kantar[1], Lisa Héron-

Milhavet[1], Bettina Fischer[3], Sarah J. Bray[4], Jacques Colinge[1], and Alexandre

Djiane[1]*

[1] IRCM, Inserm, Univ Montpellier, ICM, Montpellier, France

[2] IRCM, Inserm, Univ Montpellier, ICM, CNRS, Montpellier, France

[3] Dept. of Genetics, University of Cambridge, Cambridge, United Kingdom

[4] Dept. of Physiology Development and Neuroscience, University of Cambridge,

Cambridge, United Kingdom

*Author for correspondence: IRCM, Inserm U1194

16   208, rue des Apothicaires

17   34298 Montpellier, Cedex, France

18   tel: +33 (0) 467 612 441

19   email: alexandre.djiane@inserm.fr

KEY WORDS:

Notch signaling, *Drosophila*, epithelial polarity, neoplasia, cell competition

SUMMARY

Aggressive neoplastic growth can be initiated by a limited number of genetic

alterations, such as the well-established cooperation between loss of cell architecture

and hyperactive signaling pathways. However, our understanding of how these

different alterations interact and influence each other remains very incomplete. Using

Drosophila paradigms of imaginal wing disc epithelial growth, we have monitored

the changes in Notch pathway activity according to the polarity status of cells and

show that epithelial polarity changes directly impact the transcriptional output of the

Notch pathway. Importantly, we show that this Notch pathway redirection is not

mediated by a redeployment of Su(H), the Notch dedicated transcription factor, but

relies on the cooperation with a combination of oncogenic transcription factors. Our

work highlights in particular the role of the stress response CEBPG homologue

CG6272/Irbp18 and of its partner Xrp1 suggesting that parts of the cellular

competition program might promote neoplastic growth.

INTRODUCTION

In multicellular organisms, cells constantly integrate many stimuli, to adapt their behaviors and responses. How cells achieve this integration is a fundamental question in biology and its disruption underlies many pathologies. Cancer genomes harbor many mutations and understanding how different mutations interact within a given cell is a major goal in cancer research.

Epithelial cells represent the basic unit of many organs. Their apico-basal (A/B) polarity is controlled by the asymmetric segregation of highly conserved protein complexes such as the Scrib/Dlg/Lgl complex (Bilder et al., 2003; Coopman and Djiane, 2016; St Johnston and Ahringer, 2010). The far-reaching effects of A/B polarity is epitomized by the observation that many tumors of epithelial origin exhibit impaired polarity, and that several viral oncoproteins target polarity complexes (Banks et al., 2012; Huang and Muthuswamy, 2010).

Studies in human cell lines and in animal models have also suggested a contributing role of polarity alterations to tumor formation. For instance, mutations in the baso-lateral determinant *SCRIB1* have been shown to control proliferation and invasion in MCF-10A human mammary cells (Cordenonsi et al., 2011). Similarly in *Drosophila,* *scrib*, *dlg* or *lgl* mutations, result in multilayered overgrowth of larval epithelial imaginal discs (Bilder et al., 2003; Bunker et al., 2015). However, this uncontrolled growth is at least partly achieved because larvae exhibiting *scrib* mutations fail to undergo proper metamorphosis and imaginal discs grow for an extended period. Indeed, *scrib* mutant cells actually grow slower than wild-type cells and are eliminated by wild-type neighbors (Cordero et al., 2010; Igaki et al., 2009, 2006; Ohsawa et al., 2011). Interestingly, this is reversed when additional mutations are

introduced, such as overexpression of the BTB/POZ chromatin remodelers Abrupt or

Chinmo (Doggett et al., 2015; Turkel et al., 2013) or the constitutive activation of

signaling pathways (e.g. Ras or Notch), converting *scrib* mutant cells into aggressive,

invasive and hyperproliferative cells (Brumby and Richardson, 2003; Pagliarini and

Xu, 2003). Similar observations have been reported in mouse, where Notch or Ras

activation and *Par3* depletion cooperate to generate aggressive neoplasms in mouse

mammary glands (McCaffrey et al., 2012; Xue et al., 2013).

The Notch pathway is a highly conserved cell-signaling pathway mis-regulated in

several cancers (Ntziachristos et al., 2014; Ranganathan et al., 2011). Upon activation,

Notch receptors undergo two proteolytic cleavages to release their intra-cellular

domain or NICD, which enters the nucleus, binds to the Notch pathway specific

transcription factor CSL (Rbpj in mammals; Suppressor of Hairless, Su(H) in

Drosophila), and converts it from a repressor to an activator to turn on the

transcription of specific target genes (Bray, 2016). These Notch direct target genes

differ depending on cell type and account for the variety of outcomes triggered by

Notch activity. Increased Notch activity has been associated with several epithelial

cancers such as non-small-cell lung carcinomas (Maraver et al., 2012; Ntziachristos et

al., 2014), but in animal models, the sole increase in Notch activity either promotes

differentiation or only results in benign over-proliferation (hyperplasia) (Brumby and

Richardson, 2003; Djiane et al., 2013; Fre et al., 2005; Ho et al., 2015; McCaffrey et

al., 2012). However, as mentioned previously, Notch pathway activation cooperates

with loss of polarity to generate invasive neoplasms (Brumby and Richardson, 2003;

Ho et al., 2015; McCaffrey et al., 2012; Pagliarini and Xu, 2003).

So, while the cooperation between loss of cell architecture and hyperactive signaling

pathways is well established, the underlying mechanisms remain poorly understood. It

could merely reflect an additive effect where the consequences of both events

combine (the additive model). Alternatively, it could indicate a more profound

integration within epithelial cells where these two events impact on each other to

generate unique new behaviors (the redirective model). Using *Drosophila* paradigms

of imaginal wing disc epithelial growth, we have monitored the changes in Notch

pathway activity according to the polarity status of cells and show that epithelial

polarity changes directly impact the transcriptional output of the Notch pathway. We

further provide evidence that this Notch redirection is not mediated by new genomic

binding regions for Su(H), but relies on the cooperation with Su(H) of a combination

of transcription factors, such as basic leucine zipper (bZIP), whose activity is

triggered in response to JNK signaling during polarity loss, extending earlier reports

on the cooperation between oncogenic Ras and polarity loss (Atkins et al., 2016;

Davie et al., 2015; Külshammer et al., 2015; Uhlirova and Bohmann, 2006). Our work

highlights in particular the role of the stress response CEBPG homologue

CG6272/Irbp18 and of its partner Xrp1, key factors in mediating the loser fate during

cell competition (Baillon et al., 2018; Blanco et al., 2020; Ji et al., 2019; Lee et al.,

2018), suggesting that cellular competition, or parts of the cellular competition

program, are co-opted during neoplastic growth.

RESULTS

Notch activation and scrib mutation cooperate to promote neoplastic growth

In order to gain insights into the mechanisms underlying neoplastic growth, we first characterized the effects of Notch activation and *scrib* mutation mediated epithelial polarity impairment on wing disc growth. Using precisely controlled *Drosophila* larvae culture conditions (crowding and timing), we compared the phenotypes of wild-type (WT), Nicd overexpressing (N), *scrib* mutant (S), and Nicd overexpressing and *scrib* mutant (NS) 3rd instar wing imaginal discs at 6 days after egg laying at 25C. These different paradigms are shown in Figure 1A-D. For clarity, in all figures N will be shown in green, S in red, and NS in blue. Reproducing our previous observations (Djiane et al., 2013), N discs overgrew compared to WT, but remained as monolayered epithelia with properly localized E-Cadherin-based adherens junctions, and represent a paradigm of hyperplastic-like growth (Fig. 1A&B). S discs were smaller than WT, but grew as unstratified mass of cells with weak uniform E-Cadherin (E-Cad). These discs however showed an extensive expression of the JNK signaling target Mmp1 (Fig. 1C), a metallo-protease implicated in the digestion of the extracellular matrix, indicative that *scrib-* cells activate JNK signaling and are prone to invasiveness (Igaki et al., 2006; Uhlirova and Bohmann, 2006). It is noteworthy that S larvae did not pupariate and if left to grow for longer, the S discs ultimately developed as massive overgrowths with very disrupted epithelial polarity, that invaded and fused with neighboring tissues such as other discs (Bilder et al., 2003). Strikingly, NS discs combined aspects of N and S discs. They were overgrown like N discs, but also expressed low levels of E-Cad and high levels of Mmp1 like S discs (Fig. 1D). These discs grew as multilayered tissues and were able to invade the surrounding tissues such as haltere discs, and represent therefore a paradigm for neoplastic-like growth.

Neoplastic and hyperplastic discs have different transcriptomes

The context of the scribble mutation converts the Notch-based *Drosophila* wing disc into a paradigm of neoplastic growth. Until now, the cooperation between activated Ras and *scrib* mutations has been studied extensively, mainly in the imaginal eye disc (Atkins et al., 2016; Cordero et al., 2010; Davie et al., 2015; Igaki et al., 2009; Katheder et al., 2017; Pagliarini and Xu, 2003; Toggweiler et al., 2016; Wu et al.,

2010), but less attention has been given to the way that other activated pathways, such

as Notch or Hedgehog, also cooperate with polarity loss beyond the initial studies

(Brumby and Richardson, 2003; Pagliarini and Xu, 2003). Exploring how these other

pathways cooperate is important to evaluate the extent that are generally important for

neoplasia or specific to Ras signaling induced transformations.

First, we compared the transcriptomes of the different genetic conditions, to identify

genes whose expression was significantly different. Differential expression analyses

of genome-wide RNA-seq profiles (using DESeq with adjusted p-value for multiple

testing <0.05; (Anders and Huber, 2010) identified the cohorts of genes that were

significantly up-regulated or down-regulated in each condition compared to WT

controls (Fig. 1E&F and Supplemental Tables 1-3). The numbers were broadly

similar in each condition N (503 up; 663 dw), S (757 up; 1029 dw), and NS (1003 up;

991 dw). Using semi-quantitative qRT-PCR we then validated a subset of the

transcriptional changes and could confirm the specific differences. The N only

affected genes *E2f1*, *Sdr*, and *mxc*, were activated only in N discs and not in S or NS

while the S only gene *p38a* and the NS only genes *Act87E* and *Wnt10*, were activated

only in their respective conditions. In addition, *Ets21C*, *ftz-f1*, and *Atf3* were up-

regulated in all three conditions, as detected by the RNA-Seq profiles (Fig. 1G)

Comparing these data with previously published transcriptome analyses on similar or

related genetic backgrounds revealed significant overlap, validating our experimental

approaches. For instance, 174 of the 503 up-regulated genes in N, and 285 of the 663

down-regulated genes were also detected in our previous analysis using dual color

differential expression arrays (significant overlap p=1.11e-273, hypergeometric test,

(Djiane et al., 2013). Similarly, 676 of the 757 up-regulated genes in S were identified

in a previous analysis of *scrib* depleted discs (significant overlap p=4.90e-193;

(Bunker et al., 2015).

To gain insights into the nature of the mis-regulated genes we performed a GO term

analysis on the genes affected in N, S, and NS (p-value < 0.05). As expected from

their genetic composition, the N and NS tissues were over-represented for genes in

the Notch signaling pathway (GO:0007219) and the NS and S had changes in A/B

polarity (GO:0045197; GO0019991) (Fig. 2A and Supplemental Tables 4&5).

Focusing on the other most robust GO terms (q-value < 0.05) revealed common

alterations in all three growth paradigms which included cell adhesive properties

(GO:0030198 "extracellular matrix organization"; GO:0007155 "cell adhesion"),

ribosomal biology (GO:0042274/GO:0042273 "ribosomal small/large subunit

biogenesis") suggesting that protein synthesis might be affected, mis-regulated

metabolic processes, in particular with respect to glucose and glycolysis

(GO:0006002), or oxidative stress (GO:0006979 "response to oxidative stress"; Fig.

2A and Supplemental Tables S4&5).

Despite these similarities, there were also differences in the profiles. For example,

markers of increased proliferation were found in both NS and N, but not S, consistent

with the overgrowth phenotypes (e.g. "mitotic cytokinesis" and "mitotic spindle

organization"; GO:0000281; GO:0007052). The overgrowing behavior of NS tissues

may therefore be driven by the Notch activation (Fig. 2A and Supplemental Tables

S4&5) although it should be noted that N was enriched in additional mitosis related

categories such as "mitotic metaphase plate congression" or "centriole replication"

(GO:0007080; GO:0007099), suggesting a more robust proliferation signature in N

tissues. Similarly, both NS and S (but not N) showed characteristics of cell migration

(e.g. "border follicle cell migration"; GO:0007298) and cellular stress such as

"response to starvation" (GO:0042594), or "response to endoplasmic reticulum

stress" (GO:0034976) suggesting that these features in NS were likely contributed by

the *scrib* mutation (Fig. 2A and Supplemental Tables S4&5).

Notably there were several GO categories specific to NS including "positive

regulation of apoptotic signaling pathway" (GO:2001235) "negative regulation of

SAPK signaling cascade" (GO:0070303) and "mitotic G1/G2 DNA damage

checkpoint" (GO:0031571/GO:0007095). These results argue that the combined

Notch activation and polarity loss promoted the emergence of new cell behaviors and

responses, in particular related to DNA damage responses (Fig. 2A&B and

Supplemental Tables S4&5).

Distinguishing between the additive and redirective models of cooperation

The transcriptomic changes revealed that NS encompassed not only additive N and S

features but also the emergence of new behaviors. This raises the question of how the

defects in *Notch* and in *scrib* cooperate to produce these transcriptional consequences.

Two main models could be proposed for genetic cooperation: (1) an additive model,
in which the effect of each genetic change is independent and summed to produce the
phenotype, and (2) a redirective model, in which the two mutations influence each
other to change their effects. Since Notch signaling exerts a direct effect on
transcription, without intermediates (unlike Ras and RTK signaling for instance;
(Bray, 2016), our Notch-based paradigms may allow us to distinguish between the
two models. Indeed, in the N and NS paradigms overexpressing the transcriptional
activator Nicd, the genes directly activated by Notch (Notch Direct Targets, NDTs),
should behave differently depending on whether the cooperation is additive or
redirective. In the additive model, genes directly activated by Notch should be
insensitive to the polarity status of the cells and should remain the same in the
hyperplastic (Nicd only) and neoplastic (Nicd & *scrib-*). Alternatively, if the loss of
polarity affects Notch directly, the directly activated genes should be, at least in part,
different.
224
Genes that are directly regulated by Notch (NDTs) should have the transcription
complex, containing Su(H) bound at their regulatory regions (Djiane et al., 2013). To
identify potential NDTs in N and NS, we thus monitored the genomic regions
occupied by the Su(H) transcription factor by genome-wide Chromatin Immuno-
Precipitation (ChIP) (Supplemental Table S6). These overlapped significantly with
our previous analysis of Notch induced overgrowth, suggesting that we have captured
all the robust regions of Su(H) enrichment.
232
Strikingly, there was a strong overlap in the Su(H) bound regions in both conditions:
almost all (416 out of a total of 464) NS Su(H) peaks overlap with peaks present in N
discs. This implies that the vast majority of Su(H) binding remained the same in N
and NS (Fig. 3C & S1C). The overlap was also important between N and S peaks
(447/554 S peaks overlapping with N peaks; Fig. 3C & S1C). These results suggests
that in NS neoplastic discs (and to a lesser extent in S discs), a minority of Su(H)
peaks represent new binding regions compared to hyperplastic N discs, and that the
new NS behaviors are not the consequence of general redistribution of the Notch
specific transcription factor Su(H).
We note that the majority of binding in all conditions is associated with intronic
regions (39.70% and 42.89% in N and NS respectively; Fig. S1A), which are

frequently the site of enhancers in *Drosophila* patterning and growth genes. There

was a subtle shift in the distribution in other regions with a decrease in Su(H) peaks

that were associated with promoter regions in NS discs (from 31.55% in N to 23.06%

in NS; Fig. S1A) complemented by an increase in Su(H) binding associated with

intergenic regions (from 15.51% in N to 20.47% in NS; Fig. S1A), but the

significance of this shift is unclear.

In order to estimate the programs specifically activated by Notch in N, S, and NS, we

then intersected the transcriptomic data with the Su(H) ChIP data, considering that

upregulated genes located within 20kb of a Su(H) peak were likely NDTs. Using this

approach, we identified similar numbers of NDTs in N (177) and NS (171) (Fig.

3A&B, Supplemental Table S7). Again, there was substantial overlap with previous

data, with 64/177 NDTs in N conditions being identified in our previous study

(significant overlap p=5.89e-96, hypergeometric test (Djiane et al., 2013). When the

N and NS scenarios were compared, 69 genes were common to both and thus

represent core NDTs in the wing disc overgrowth. However, a significant proportion

of NDTs appeared specific for each condition: 108 for N, 102 for NS (Fig. 3B) which

supports the redirective model. Amongst the 102 NS-specific NDTs, only 24 were

also NDTs in S, arguing that the difference cannot be explained by specific

contribution of the S condition. Taking all of these comparisons into account, there

appear to be 78 genes that are NS-specific NDTs. Only a minority were associated

with new Su(H) binding regions: around the *87E* locus (*yellow-e3, yellow-e, Ir87a,*

and *Act87E*), and the *94A* locus (*CG18596, CG7059, CG13857,* and *CG13850*).

Amongst these 78 NS specific NDTs are p53, His2Av, and the Tip60 complex

component Act87E (Kusch et al., 2004), which have all been linked to DNA damage

response and could therefore account for the NS emergent behaviors identified in the

GO term analysis (Fig. 2A and Supplemental Tables S4&5). The Venn-diagrams

representing the overlaps in N, S, and NS, highlight the extent of additivity (the

yellow circles encompassing common NDTs) and the scale that there is redirection of

the Notch pathway output (red circles). Both losses and gains occur so that in the

conditions where there is neoplastic growth (NS), 78 new NDTs are acquired while

108 NDTs are no longer present compared to N (and 70 compared to S). The

observation that the NDTs differ substantially between NS and N is in strong support

of the redirective model.

Taken together, our results indicate that Notch activation and *scrib* mediated loss of

polarity cooperate both in an additive and redirective manner to promote neoplastic

growth, and that the redirection of the Notch pathway transcriptional output is not a

consequence of a general redeployment of Su(H) to new active sites. They argue that

alterations in the cell architecture can directly influence the outcome of the Notch

signaling pathway and support a model in which different genetic injuries within

tumor cells could thus influence each other to sustain neoplastic growth.

Identification of the transcriptional networks in our different growth paradigms

Using the NDT datasets, we sought to identify the factors that are required for the

transition from N hyperplastic to NS neoplastic growth. As the NS transcriptome

showed signatures consistent with "response to oxidative stress" (GO:0006979),

"cellular response to gamma radiation" (GO:0071480), and "DNA damage

checkpoints" (GO:0031571/0007095), we first asked whether interfering with such

pathways could block the growth and invasiveness of NS tissues.

To perform these genetic tests, we generated a stable fly line which overexpressed

Nicd and *scribRNAi* together with a *GFP* marker under the *BxGal4* driver (driving

expression in the pouch of the larval wing discs; *Bx>NS*), and monitored the size of

the overgrowth (GFP positive tissue), and its invasiveness potential (Mmp1

expressing cells; Fig. 5A). Blocking the oxidative stress response by overexpressing

the Reactive Oxygen Species (ROS) sponge CAT and SOD, or knocking down by

RNAi the expression of *ATM* or *ATR* (*tefu* or *mei-P26* in the fly), the two master

kinases mediating the early response to DNA damage (double or single strand breaks

respectively) did not have any significant effect on the NS overgrowth or the

expression of Mmp1 (Fig. 5I). Similarly, expression of RNAi or dominant negative

forms of the acute stress response and severe DNA damage major effector and NS

specific NDT *p53* could not modify the NS overgrowth phenotype (Fig. 5I). While we

cannot exclude that the tools used here were not strong enough, these results suggest

that even though activated in NS tissues, oxidative stress and DNA damage responses

were either not required to sustain NS growth, or that they could compensate for each

other converging ultimately on an as yet unidentified core response promoting NS

growth.

Next we sought to identify transcriptional factors that could account for the

cooperation between Notch and polarity loss. We used the iRegulon software to

identify the transcriptional network involved. When utilized to analyze genes

involved in sustaining the tumorous growth of *RasV12/scrib-* 3rd instar larval eye,

wing and leg discs, it highlighted the role of the Hippo pathway terminal effectors

Yki/Sd and their targets Myc, Crp, and Ftz-F1, the JNK pathway regulated AP-1

factors (in particular Atf3, Kay, and CEBPG), and the Jak/Stat pathway (Atkins et al.,

2016; Davie et al., 2015; Külshammer et al., 2015).

Implementing iRegulon on our step-wise Notch-based paradigms allowed us (i) to

identify transcriptional modules unique or shared between polarity loss only (S),

proliferation only (N) and proliferation plus invasiveness (NS), and (ii) to assess the

conservation of the "oncogenic" modules identified previously with Ras in a Notch-

driven neoplastic paradigm. We performed these analyses feeding iRegulon either

with the lists of up-regulated genes in N, S, and NS (Fig. 4A & Supplemental Tables

S7-11) or with the lists of NDTs (Fig. S2 & Supplemental Table S12-15). We decided

to focus our analyses on the up-regulated genes here since: i) the overexpressed Nicd

triggers transcriptional activation (Bray, 2016) and down-regulated genes might thus

represent very indirect effects of the cooperation, and ii) the Su(H) module should be

found enriched in the regulatory regions of the up-regulated genes and represent thus

an internal control to evaluate the bio-informatic predictions. These analyses

identified "modules of transcription factors" likely co-regulating the genes identified

in N, S, or NS. Modules identified are presented as Venn diagram (Fig. 4A & S4) to

highlight common and specific programs in the different conditions. Feeding either

up-regulated genes (Fig. 4A) or NDTs (Fig. S2) identified similar modules indicating

that the Notch pathway redirection is mediated, at least in part, by transcription

factors that broadly affect the whole transcriptome. This is likely also true for other

neoplastic paradigms such as Ras, even though this remains to be established.

Importantly, iRegulon identified the Notch pathway dedicated transcription factor

Su(H) in the N and NS transcriptomes.

First, focusing on NS, which most resembles the *RasV12/scrib-* paradigms, our

analysis identified the same major nodes and oncogenic module as described

previously:

- AP-1 basic Leucine Zipper factors related to stress kinase signaling; *Mmp1* is a

canonical JNK targets and is highly expressed in NS growing discs (Fig. 1D)

- Stat92E of the Jak/Stat pathway; *upd3* coding for the upstream ligand of the Jak/Stat

pathway is indeed highly expressed in NS discs

- Ftz-F1 nuclear receptor

- basic Helix-Loop-Helix factors of the Myc family.

In NS transcriptome, we also identified a contribution of the E(spl) bHLH

transcriptional repressors. E(spl)-HLH genes are canonical Notch targets, and they are

robustly up-regulated in N and NS, in particular E(spl)mγ-HLH. In NS there were

however two noteworthy differences when focusing only on NDTs, rather than all

differentially transcribed genes. (1) The absence of the E(spl) factors, suggesting that

when considering the whole NS transcriptomes a significant of genes could be

associated with E(spl)-mediated regulation while this is no longer the case when

considering the more restricted number of NS NDTs. (2) The identification of Ewg,

the fly NRF1 homologue. In mammals, NRF1 is implicated in the regulation of key

metabolic and respiratory genes (Herzig et al., 2000).

The iRegulon analyses also suggested that the AP-1 bZIP, and the Stat92E signatures

in NS are contributed by S, since they are also detected in the S transcriptomes, while

the Su(H) signature is contributed by N. Interestingly, the bHLH Myc signature is

found in N and S as well as NS. Finally, iRegulon identified a signature for the

Polycomb chromatin silencers specifically in N. Such factors include Pho, a zinc

finger protein which binds to Polycomb Responsive Elements (PREs) and recruits

Polycomb complexes, and the three Polycomb Repressor Complex 1 components Psc,

Su(z)2, and l(3)73Ah. Recently, PRC1 has been associated with specific and

unexpected transcriptional activation at larval stages, raising the possibility that in N,

such genes, normally repressed at embryonic stages, become active (Loubiere et al.,

2020).

A polarity-loss oncogenic module

In order to validate the functional relevance of the transcription factors identified in

the Notch-driven neoplastic growth, we then asked whether their depletion by RNAi

could alter the growth and invasiveness of the NS tissue, using the *Bx>NS* fly line

described previously. We first focused our analysis on the the oncogenic module.

We confirmed earlier reports that blocking JNK activity by the overexpression of a

JNK dominant negative construct, strongly abolishes NS driven growth (GFP positive

tissue size; Fig. 5B&J) and invasiveness (Mmp1 expression; Fig. 5B&J'). As shown

in the *RasV12/scrib-* paradigms, RNAi mediated knock-down of the Jak/Stat pathway

terminal transcription factor *stat92E*, the JNK pathway component *Ets21C*, and to a

lesser extent *ftz-f1* (Atkins et al., 2016; Davie et al., 2015; Külshammer et al., 2015;

Toggweiler et al., 2016) strongly suppressed both growth (GFP) and invasiveness

(Mmp1; Fig. 5J&J'). Unlike the RasV12 models, we did not identify any particular

enrichment for Sd/TEAD, the transcriptional factor mediating the effect of Yki and of

the Hippo pathway mediated growth in wing discs. However, impairing Yki activity

(through RNAi-mediated knock-down) strongly suppressed NS neoplastic behaviors

(Fig. 5C&J).

Taken together these results suggest that independently of the oncogenic driver, Ras

or Notch, relatively similar tumorous transcriptional networks (AP-1/Yki/Stat/Ftz-f1)

are put in place during their cooperation with polarity loss (Atkins et al., 2016; Davie

et al., 2015; Külshammer et al., 2015). Given that these nodes were also identified in

the S transcriptome, we suggest that they might represent a polarity loss module

cooperating with oncogenic signaling pathways Ras or Notch (and likely other

pathways such as Hh as was initially reported; (Brumby and Richardson, 2003).

The CG6272/Xrp1 module is required for neoplastic growth

A striking feature of the iRegulon analyses was that the JNK module, which

contained classic basic leucine zipper (bZIP) transcription factors such as Jun or Fos,

was found in NS and S. As the bZIP factors identified belonged to sub-families

involved in different cellular responses including oxidative stress (Maf-S, Cnc,

(Sykiotis and Bohmann, 2008), polarity loss (Atf3, (Donohoe et al., 2018), ER stress

(Atf6, Crc, (Harding et al., 2003; Ye et al., 2000), DNA damage… they were potential

candidates to mediate the transcriptional switch. First, we analyzed whether these

different bZIP factors were specifically up-regulated in NS (Fig. 4B). Most factors,

with the exception were upregulated in S, suggesting that polarity impairment

promotes the expression of many bZIP factors which could alter the response of the

transcriptional machinery. A further up-regulation in NS, even compared to S, could

be detected for kay, crc, maf-S, CG7786, and CrebB (Fig. 4B). Strikingly, cnc and

Xrp1were upregulated only in NS (Fig. 4B). It should be noted however, that gene up-

regulations do not always indicate functional relevance. Then, in order to identify

which bZIP factor/family is most relevant for NS, we therefore systematically

depleted by RNAi the different genes highlighted by iRegulon in this bZIP node,

irrespective of their potential up-regulations in NS, using the *Bx>NS* fly line.

Amongst the different factors tested, we observed a significant suppression of the

invasiveness (Mmp1; Fig. 5K') but not of the growth (GFP; Fig. 5K) of NS tissues

after impairing the AP-1 factors *kay* and *Jra* (fly homologues of *FOS* and *JUN*

respectively), and *CrebB-17A*. Conversely, knocking down the DNA damage

response factor *maf-S* led to a dramatic reduction of the GFP outgrowth but without

affecting the invasiveness in which the tissue was expressing high levels of Mmp1

(Fig. 5G&K). It should be noted here that these *maf-S* depleted discs appeared sick

with a severely irregular and misshapen GFP domain (Fig. 5G&K, star), suggesting

that maf-S might be an essential gene, and that the size suppression might merely

reflect enhanced cell death. In contrast, we did not observe any significant effect

when *cnc*, the fly homologue of *NFE2L1* implicated in oxidative stress response and

excision repair (Han et al., 2012; Sykiotis and Bohmann, 2008), *Atf2*, *gt*, *CG7786*, *vri*

(Fig. 5K), were knocked-down. Neither were any consequences on the phenotype

with knock-down of *Atf6* and *crc* (the *ATF4* fly homologue), which have been both

linked to the unfolded protein response and ER stress (Fig. 5D&K; (Harding et al.,

2003; Ye et al., 2000).

The most complete suppression of the NS behaviors (both GFP overgrowth and

Mmp1 invasiveness) was observed after depletion of *Atf3*, *Pdp1* or CG6272 (Fig.

5E,H,K). Atf3 has recently been shown to control the expression of genes involved in

the maintenance of epithelial polarity, and to be specifically activated in polarity

deficient cells and required in the RasV12/scrib- overgrowth models (Atkins et al.,

2016; Donohoe et al., 2018). Pdp1 (the homologue of Hepatic Leukemia Factor -

HLF) has previously been linked to mitotic cell cycle and growth (Reddy et al., 2006),

and shown in the *RasV12/scrib-* paradigms to have modest effects on invasion, but

none on growth. The role of *CG6272* (a.k.a. *Irbp18*, the homologue of *CEBPG*) was

also highlighted in the *RasV12/scrib-* paradigms where it was shown to control

growth but not invasion (Atkins et al., 2016). In the context of Notch (*Nicd/scrib-*),

the role of *Pdp1* and *CG6272* appeared more essential.

bZIP factors have been shown to act as homo or heterodimers. Knocking down the

function of *crc/Atf4*, the classic bZIP partner of CG6272/CEBPG, did not have any

effect on NS tissue behaviors (Fig. 5D,E,K) whereas the knock-down of *Xrp1* very

efficiently suppressed the growth and Mmp1 expression in NS tissues. This suggests

that .CG6272/CEBPG could be acting in combination with its bZIP partner Xrp1 in

this context (Francis et al., 2016; Reinke et al., 2013). Although the cellular processes

controlled by CG6272/Xrp1 remain to be determined, the fact that Xrp1 is an early

p53 target gene (Akdemir et al., 2007; Link et al., 2013), raises the possibility that its

functions could be related to DNA/genome integrity maintenance. This would align

with the NS specific GO categories "cellular response to gamma radiation"

(GO:0071480), and "DNA damage checkpoints" (GO:0031571/0007095). Indeed,

Xrp1 has recently been linked to cell competition and shown to control the expression

of a subset of genes activated in response to Ribosomal protein haplo-insufficiency

including *Ets21C*, *upd3*, or *ilp8* in developing wing discs (Baillon et al., 2018; Boulan

et al., 2019; Ji et al., 2019; Lee et al., 2018). Strikingly, we observed a very strong

overlap between the Xrp1 transcriptional program (Supplemental Table S16) and the

genes activated in NS (60 out of the 171 Xrp1 positively regulated genes are also

upregulated in NS; hypergeometric test, p=4.71 10e-21), while a much more modest

overlap could be observed with N or S (overlap of 17 and 33 with p values of 1.12

10e-3 and 1.56 10e-8 respectively). These results support the proposal that the

CG6272/Xrp1 module is functionally important in the acquisition of neoplastic

growth.

DISCUSSION

In this study, using Notch-driven paradigms of epithelial overgrowth in *Drosophila* wing discs, we describe the molecular mechanisms underlying the cooperation between Notch and polarity loss during neoplasia. We show that epithelial polarity alterations redirect the transcriptional outcome of the Notch signaling pathway. We further show that this redirection occurs mainly on pre-existing Su(H) bound regions rather than new ones. Finally, we show that similarly to what was previously described for Ras signaling (Atkins et al., 2016; Davie et al., 2015), the cooperation between Notch signaling and polarity loss is controlled by a "tumor transcriptional network" centered around the AP-1/Stat/Yki transcription factors, and including the critical bZIP factors Pdp1 and CEPBG (CG6272/Irbp18). But our analysis uncovered a previously unreported role for the cell competition regulator, and Irbp18 binding partner Xrp1 (Baillon et al., 2018; Blanco et al., 2020; Boulan et al., 2019; Ji et al., 2019; Lee et al., 2018), raising the interesting prospect that neoplastic growth could be mediated at least in part, by co-opting cell competition.

While cancer genomes exhibit multiple mutations in cancer cells, their functional interactions remain difficult to monitor and model. This analysis in the genetically controlled *Drosophila* wing disc, supports a model in which the effects of different mutations do not just simply add up, but interact with each other leading to the emergence of new cell behaviors (redirective model). Indeed, even though the combination showed characteristics that could be attributed to either Notch (proliferation and mitosis), or to polarity loss (invasion/cell migration), it also exerted signs of emerging behaviors such as DNA damage response. Neoplastic tissues appear thus to experience many cellular stresses: DNA damage responses, but also ER and unfolded protein response, starvation, or oxidative stresses. However, even though present, these different stresses and in particular oxidative stress and DNA damage are not individualy necessary in the context of polarity loss as blocking them or the cellular response they promote (CAT/SOD overexpression, or inhibition of ATM/ATR) could not significantly suppress the NS tumorous behaviors. These observations suggest that the different stress pathways activated during polarity loss might all converge leading ultimately to the activation of a common core response.

While *Drosophila* and mouse models have demonstrated that overactive signaling

pathways cooperate with epithelial polarity impairment to generate neoplastic growth

(Brumby and Richardson, 2003; McCaffrey et al., 2012; Pagliarini and Xu, 2003; Xue

et al., 2013), the vast majority of studies seeking to understand the underlying

mechanisms, have focused primarily on the cooperation between activated RasV12

and *scrib* mutants, especially in *Drosophila* (Atkins et al., 2016; Cordero et al., 2010;

Davie et al., 2015; Igaki et al., 2009; Katheder et al., 2017; Pagliarini and Xu, 2003;

Toggweiler et al., 2016; Wu et al., 2010). Importantly, the current study, investigating

the cooperation between Notch and polarity, shows that many observations made for

Ras can be extended to Notch, suggesting that the paradigms used are not a Ras

specificity but might represent a more general tumor growth paradigm.

But even though we could highlight the involvement of a core "oncogenic module"

(Atkins et al., 2016; Davie et al., 2015; Külshammer et al., 2015), there are specifics

that are likely oncogene specific. In the case of Ras, it was shown that Yki activity

could reprogram Ras by promoting the expression of the Ras pathway specific

regulators Capicua and Pointed to promote aggressive growth (Pascual et al., 2017).

Both genes were either unaffected (*capicua*) or downregulated (*pointed*) in NS Notch

driven neoplastic paradigm, suggesting that, even though Yki is clearly active (Fig.

5C), changes in the expression of capicua and pointed are unlikely mediators here.

Furthermore, in NS transcriptome, we identified a contribution of the E(spl) bHLH

transcriptional repressors, canonical Notch targets (Bray, 2016), which represents thus

a Notch specificity. However, the fact that motifs for E(spl)-HLH repressors are

found in the up-regulated transcriptome of NS and not N could suggest that in NS, the

repressive ability of E(spl)-HLH factors is antagonized (even though their expression

is not affected). It would be interesting to explore further the link between NS and

E(spl)-HLH-mediated repression, but due to the high redundancy between the seven

E(spl)-HLH factors (δ, γ, β, 3, 5, 7, 8) and Dpn, the requirement of E(spl)-HLH-

mediated repression in the Notch-driven neoplasia could not be formally tested.

Amongst the most dramatic suppressors of the neoplastic growth are the two basic

leucine zipper transcription factors CG6272/Irbp18 (the fly homologue of CEBPG),

and Xrp1. These two genes, and in particular *Xrp1*, were up-regulated in NS

compared to WT or N. Importantly, impairing with their function in wild-type tissues

did not have any effect, confirming observations made by other groups for CG6272

(Atkins et al., 2016), supporting a model in which these genes are dispensable in

healthy wild-type cells, but become indispensable for tumor cells, hence representing

a possible attack strategy specifically targeting tumor cells, while sparing healthy

tissues. In mammals, CEBPG represents a major regulator of stress responses. It is

recruited through its interaction with the bZIP factor ATF4 at the level of cis-

regulatory C/EBP:ATF response elements to activate the expression of stress

mitigation genes such as glutathione biosynthesis pathway genes in the case of

oxidative stress (Huggins et al., 2015). Interestingly, we did not observe any effect

knocking down *Crc*, the fly *Atf4* homologue, suggesting that during NS overgrowth in

Drosophila, CG6272/Irbp18 could be acting in combination with another partner.

CG6272/Irbp18 also interacts with Xrp1, and the heterodimer has been implicated in

DNA repair (Akdemir et al., 2007; Francis et al., 2016). We thus propose that the

Irbp18/Xrp1 dimer is activated in NS and is required to facilitate DNA repair and

ensure genomic stability to prevent catastrophic genotoxic effect upon the combined

cellular stresses of S and replication stress of N. Recently, *Xrp1* together with its

binding partner CG6272/Irbp18 (but not Atf4) has been shown to mediate a loser

status in ribosomal genes deficient cells, through the implementation of a specific

transcriptional program (Baillon et al., 2018; Blanco et al., 2020; Ji et al., 2019; Lee et

al., 2018). This "loser state" promoter role of Xrp1/Irbp18 is however in conflict with

our observation that *Xrp1* is strongly upregulated in NS overgrowing discs, and that

Xrp1 and *Irbp18* are required for the overgrowth and invasive capacities of NS

neoplastic cells. This could be reconciled by proposing that in NS neoplastic discs, the

role of the Xrp1/Irbp18 is modified by an as yet unknown factor. Alternatively, and

more attractively, it is also possible that in NS cells, cell death is prevented as was

shown for RasV12 expressing cells (Pinal et al., 2018), for instance by upregulating

DIAP1 (a Notch direct target, (Djiane et al., 2013) leading to a perverted cell

competition. There an incomplete "loser" program would be initiated, but not fully

implemented (no cell death), leading to the secretion of growth factors (e.g. the

Jak/Stat ligands upd…) that would act in an autocrine manner to further promote their

growth. These factors do not actually need to act cell autonomously, since it was

shown that at least in the *RasV12 / scrib-* paradigm, delaminating cells cooperate with

non-delaminating proliferating cells to sustain tumor growth (Muzzopappa et al.,

2017; Uhlirova et al., 2005; Wu et al., 2010). In NS, preventing the

losing/delaminating cells by impairing the Xrp1/Irbp18 nexus, would thus prevent

neoplastic growth. This intriguing possibility highlights that more studies are needed

to better understand the role of the Xrp1/Irbp18 module, its links to cell competition,

and to the growth of neoplastic tissues.

MATERIALS AND METHODS

Drosophila genetics

The different overgrowth paradigms were obtained by generating random clones in 3rd instar wing discs at high frequency as previously published in (Djiane et al., 2013). In brief, the *abxUbxFLPase; Act>y>Gal4, UAS GFP; FRT82B tubGal80* flies were crossed either to *FRT82B* (to generate Ctrl discs), or to *UAS-Nicd; FRT82B* (to generate hyperplastic N discs), or to *UAS-Nicd; FRT82B scrib1* (to generate neoplastic NS discs). *scrib1* represents a loss of function allele for the *scribble* gene. Because *scrib1* clones are eliminated in growing discs, the dysplasic S discs were obtained from *FRT82B scrib1 / Df(3R)BSC752* 3rd instar larvae. All crosses were performed at 25°C and carefully staged (time after egg laying and tube crowding). For functional studies, neoplastic growth was obtained by driving *UAS-Nicd* and the *scrib* RNAi *P{TRiP.HMS01490}attP2* by the *Bx-Gal4* (pouch of larval wing discs). Modifications of the overgrowth phenotype and of the expression of the Mmp1 invasive marker were performed by crossing in F1 *Bx-Gal4, UAS GFP;; UAS Nicd, UAS scribHMS01490* to the desired *UAS RNAi* or control lines (*UAS white RNAi* or *UAS GFP*), to ensure similar UAS load. List of lines tested in Supplemental Materials.

Information on gene models and functions, and on *Drosophila* lines available were obtained from FlyBase (flybase.org – (Thurmond et al., 2019).

RNA extraction and RNA-Seq

RNA from 60 or 80 dissected third instar larva wing discs of WT, N, NS and S discs was extracted using TriZOL. Genomic DNA was eliminated using Ambion's DNA-free kit (#AM1906). cDNA bank preparation were then performed from 1μg of RNA and sequencing on a Illumina HisSeq 2000 by the Biocampus genomic facility MGX of Montpellier. After sequencing, reads obtained were filtered based on their quality (circa 40 millions reads were kept per conditions). The reads were then align on *Drosophila* dm6 genome by the ABIC facility in Montpellier producing a matrix of reads per gene and per condition. This matrix was then normalized and pair-wise differential expression was performed using DESeq (Anders and Huber, 2010). Other differential expression tools were tested such as DESeq2 and edgeR with default parameters but appeared either less stringent, or inadequate.

qPCR

qPCR was performed on biological triplicates on a Roche LightCycler 480, and fold

change was estimated by the δδCT approach. List of primers used in Supplemental

Materials.

Su(H) Chromatin Immuno Precipitation

After dissection in PBS 1X, Protein/DNA complexes from 60 wing discs (80 for S

condition) were cross-linked with 1% formaldehyde for 10 minutes. The reaction was

then quenched by 0.125 M Glycine and washed 3x in PBS. Wing disc cells were

resuspended in 50μL Nuclear Lysis Buffer (Tris-HCl pH 8.1 20mM, EDTA 10mM,

SDS 1%). Lysates were sonicated on a Bioruptor (Diagenode), and diluted 10x in

Immunoprecipitation Dilution Buffer (Tris-HCl pH 8.1 20mM, EDTA 2mM, SDS

0.01%, NaCl 150mM, Triton X-100 1%) and precleared with rabbit IgG (Sigma) and

protein G Agarose (Santa Cruz Biotechnology). ChIP reactions were performed by

incubating lysates overnight at 4°C with 1ng of Goat anti-Su(H) (Santa Cruz

Biotechnology, sc15813), and immunocomplexes were then isolated with Protein G

Agarose for 2h, washed 2x with Wash Buffer 1 (Tris-HCl pH 8.1 20mM, EDTA

2mM, SDS 0.1%, NaCl 50mM, Triton X-100 1%) and 2x with Wash Buffer 2 (Tris-

HCl pH 8.1 10mM, EDTA 1mM, LiCl 250mM, NP-40 1%, Deoxycholic acid 0.4%),

before a decross-linking step at 65°C in 0.25M NaCl. Samples were then treated with

0.2 mg/mL proteinase K and 50mg/mL RNase A. The DNA was then purified on

columns (Qiagen, 28106). ChIP efficiency was checked by qPCR normalized on input

chromatin with the following primer couples, corresponding to known strong binding

sites of Su(H). List of primers used in Supplemental Materials.

For whole-genome analysis, 1 μg double-stranded ChIP or input DNA (corresponding

to 180 discs for each replicate) was labelled with either Cy3- or Cy5-random primers

using the Nimblegen Dual Colour kit. Both ChIP and input were co-hybridised to

NimbleGen D. melanogaster ChIP-chip 2.1 M whole-genome tiling arrays in the

NimbleGen hybridisation station at 42°C for 16 h and then washed according to the

NimbleGen Wash Buffer kit instructions. The data obtained were normalized using

quantile normalization across the replicate arrays in R. Window smoothing and peak

calling were performed using the Bioconductor package Ringo (Toedling et al., 2007)

with a winHalfSize of 300 bp and min.probes = 5. Probe levels were then assigned P-

values based on the normalNull method, corrected for multiple testing using the

Hochberg–Benjamini algorithm and then condensed into regions using distCutOff of

200 bp.

In order to determine the Notch Direct Targets (NDTs), ChIP and RNA-Seq results

were compared: NDTs are defined as up-regulated genes with Su(H) enrichment

within 20kb. As such one Su(H) peak could be assigned to several upregulated genes

consistent with its role in enhancer regions. The 20kb window was chosen as it

allowed the recovery of more than 85% of NDTs in our previous study that was based

on closest gene assignment irrespective of distance (Djiane et al., 2013).

GO Term analyses

The lists of significantly regulated genes in the various comparisons were submitted

to gene ontology (GO) term enrichment analysis. We used the GO biological process

(GOBP) ontology and applied hypergeometric tests (p-values) followed by

Benjamini-Hochberg multiple hypothesis correction (q-values).

iRegulon analyses

In order to determine the likely transcriptional modules in our transcriptomic and

NDTs datasets, we used the online tool iRegulon (http://iregulon.aertslab.org/), with

the standard settings using the 6K Motif collection (6383 PWMs) and a Putative

regulatory region of "10kb upstream, full transcript and 10kb downstream".

Importantly, these settings allowed the recovery of the "positive control" Su(H)

module.

Immunocytochemistry

Antibody staining of wing imaginal discs were performed using standard protocols.

Briefly, larval heads containing the imaginal discs (LH) were dissected in cold PBS

and fixed for 20min in 4% Formaldehyde in PBS at room temperature (RT), before

being rinsed 3x 10min in PBS 0.2% TritonX100 (PBT), and blocked in PBT + 0.5%

BSA (PBTB) for 30min at RT. LH were then incubated overnight at 4°C with primary

antibodies in PBTB. LH were then rinsed 3x 10min in PBT at RT and before being

incubated with secondary antibody in PBTB for 90min at RT. LH were then rinsed 3x

20min in PBT at RT, before being equilibrated overnight in Citifluor mounting media

(Agar). Discs were then further dissected and mounted. Images were acquired on a

Zeiss Apotome2 microscope and processed and quantified using Zen or ImageJ.
Primary antibodies used were rat anti-DE-Cadherin (DCAD2, Developmental Studies
Hybridoma Bank – DHSB, 1:25), rabbit anti-GFP (A6455, Molecular Probes, 1:200),
and mouse anti-Mmp1 (3A6B4, DHSB, 1:25). Secondary antibodies used conjugated
to Alexa-350, Alexa-488, or Cy3 were from Jackson Labs Immuno Research (1:200).
686

Quantification methods
Genotypes were tested in batches with controls and 13-17 images corresponding to
13-17 different discs were all acquired on the same microscope with the same
exposure settings.
Growth was estimated by the size of the GFP positive area and normalized to that of
controls. A disc was considered overgrown if its size was over the average size of
controls plus 1.5 the stdev of controls. A disc was undergrown, if its size was under
the average of controls minus 1.5 the stdev of controls. Using this "conservative" set
up identified only circa 5% of controls as undergrown.
Mmp1 intensities were ranked as High, Low, Null by independent observer with
genotypes masked and processed in random order.
698

ACKNOWLEDGEMENTS

We thank G. Alvès, D. Andrew, D. Eberl, P. Léopold, R. Levayer, C. Maurange, A.

Teleman, J. Terman, and T.T. Su for sharing flies. We acknowledge the Bloomington

Stock Center, the Vienna Stock Center, the DGRC Kyoto Stock Center, the

Developmental Studies Hybridoma Bank, the *Drosophila* facility, the MGX

sequencing facility (BioCampus Montpellier, CNRS, INSERM, Université de

Montpellier), and FlyBase for their support to our research.

The lab of SJB is supported by the MRC (MRC programme grant MR/L007177/1).

RL was supported by fellowships from "Ligue Nationale Contre le Cancer - LNCC"

and from "Fondation ARC pour la recherche sur le cancer". The lab of AD is

supported by the "Fondation ARC pour la recherche sur le cancer", "Marie Curie

CIG", and "Agence Nationale de la Recherche - ANR".

AUTHOR CONTRIBUTIONS

RL, CG, PL, DK, and AD performed experiments; RL, CG, LHM, JC, and AD

analyzed data; JC provided expertise for RNA-Seq; BF and SJB provided technical

support and expertise for ChIP; RL, SJB, and AD wrote the manuscript; AD

conceived and supervised the project.

DECLARATION OF INTEREST

The authors declare no competing interests.

REFERENCES

Akdemir F, Christich A, Sogame N, Chapo J, Abrams JM. 2007. p53 directs focused
genomic responses in Drosophila. *Oncogene* **26**:5184–5193.
doi:10.1038/sj.onc.1210328
Anders S, Huber W. 2010. Differential expression analysis for sequence count data.
Genome Biol **11**:R106. doi:10.1186/gb-2010-11-10-r106
Atkins M, Potier D, Romanelli L, Jacobs J, Mach J, Hamaratoglu F, Aerts S, Halder G. 2016.
An Ectopic Network of Transcription Factors Regulated by Hippo Signaling
Drives Growth and Invasion of a Malignant Tumor Model. *Curr Biol* **26**:2101–
2113. doi:10.1016/j.cub.2016.06.035
Baillon L, Germani F, Rockel C, Hilchenbach J, Basler K. 2018. Xrp1 is a transcription
factor required for cell competition-driven elimination of loser cells. *Sci Rep*
8:17712. doi:10.1038/s41598-018-36277-4
Banks L, Pim D, Thomas M. 2012. Human tumour viruses and the deregulation of cell
polarity in cancer. *Nat Rev Cancer* **12**:877–886. doi:10.1038/nrc3400
Bilder D, Schober M, Perrimon N. 2003. Integrated activity of PDZ protein complexes
regulates epithelial polarity. *Nat Cell Biol* **5**:53–58. doi:10.1038/ncb897
Blanco J, Cooper JC, Baker NE. 2020. Roles of C/EBP class bZip proteins in the growth
and cell competition of Rp ('Minute') mutants in Drosophila. *Elife* **9**.
doi:10.7554/eLife.50535
Boulan L, Andersen D, Colombani J, Boone E, Léopold P. 2019. Inter-Organ Growth
Coordination Is Mediated by the Xrp1-Dilp8 Axis in Drosophila. *Dev Cell* **49**:811-
818.e4. doi:10.1016/j.devcel.2019.03.016
Bray SJ. 2016. Notch signalling in context. *Nat Rev Mol Cell Biol* **17**:722–735.
doi:10.1038/nrm.2016.94
Brumby AM, Richardson HE. 2003. scribble mutants cooperate with oncogenic Ras or
Notch to cause neoplastic overgrowth in Drosophila. *EMBO J* **22**:5769–5779.
doi:10.1093/emboj/cdg548
Bunker BD, Nellimoottil TT, Boileau RM, Classen AK, Bilder D. 2015. The transcriptional
response to tumorigenic polarity loss in Drosophila. *Elife* **4**.
doi:10.7554/eLife.03189
Coopman P, Djiane A. 2016. Adherens Junction and E-Cadherin complex regulation by
epithelial polarity. *Cell Mol Life Sci* **73**:3535–3553. doi:10.1007/s00018-016-
2260-8
Cordenonsi M, Zanconato F, Azzolin L, Forcato M, Rosato A, Frasson C, Inui M,
Montagner M, Parenti AR, Poletti A, Daidone MG, Dupont S, Basso G, Bicciato S,
Piccolo S. 2011. The Hippo transducer TAZ confers cancer stem cell-related
traits on breast cancer cells. *Cell* **147**:759–772. doi:10.1016/j.cell.2011.09.048
Cordero JB, Macagno JP, Stefanatos RK, Strathdee KE, Cagan RL, Vidal M. 2010.
Oncogenic Ras diverts a host TNF tumor suppressor activity into tumor
promoter. *Dev Cell* **18**:999–1011. doi:10.1016/j.devcel.2010.05.014
Davie K, Jacobs J, Atkins M, Potier D, Christiaens V, Halder G, Aerts S. 2015. Discovery of
transcription factors and regulatory regions driving in vivo tumor development
by ATAC-seq and FAIRE-seq open chromatin profiling. *PLoS Genet* **11**:e1004994.
doi:10.1371/journal.pgen.1004994
Djiane A, Krejci A, Bernard F, Fexova S, Millen K, Bray SJ. 2013. Dissecting the
mechanisms of Notch induced hyperplasia. *EMBO J* **32**:60–71.
doi:10.1038/emboj.2012.326
Doggett K, Turkel N, Willoughby LF, Ellul J, Murray MJ, Richardson HE, Brumby AM.
2015. BTB-Zinc Finger Oncogenes Are Required for Ras and Notch-Driven
Tumorigenesis in Drosophila. *PLoS ONE* **10**:e0132987.
doi:10.1371/journal.pone.0132987

Donohoe CD, Csordás G, Correia A, Jindra M, Klein C, Habermann B, Uhlirova M. 2018. Atf3 links loss of epithelial polarity to defects in cell differentiation and cytoarchitecture. *PLoS Genet* **14**:e1007241. doi:10.1371/journal.pgen.1007241

Francis MJ, Roche S, Cho MJ, Beall E, Min B, Panganiban RP, Rio DC. 2016. Drosophila IRBP bZIP heterodimer binds P-element DNA and affects hybrid dysgenesis. *Proc Natl Acad Sci USA* **113**:13003–13008. doi:10.1073/pnas.1613508113

Fre S, Huyghe M, Mourikis P, Robine S, Louvard D, Artavanis-Tsakonas S. 2005. Notch signals control the fate of immature progenitor cells in the intestine. *Nature* **435**:964–968. doi:10.1038/nature03589

Han W, Ming M, Zhao R, Pi J, Wu C, He Y-Y. 2012. Nrf1 CNC-bZIP protein promotes cell survival and nucleotide excision repair through maintaining glutathione homeostasis. *J Biol Chem* **287**:18788–18795. doi:10.1074/jbc.M112.363614

Harding HP, Zhang Y, Zeng H, Novoa I, Lu PD, Calfon M, Sadri N, Yun C, Popko B, Paules R, Stojdl DF, Bell JC, Hettmann T, Leiden JM, Ron D. 2003. An integrated stress response regulates amino acid metabolism and resistance to oxidative stress. *Mol Cell* **11**:619–633. doi:10.1016/s1097-2765(03)00105-9

Herzig RP, Andersson U, Scarpulla RC. 2000. Dynein light chain interacts with NRF-1 and EWG, structurally and functionally related transcription factors from humans and drosophila. *J Cell Sci* **113 Pt 23**:4263–4273.

Ho DM, Pallavi SK, Artavanis-Tsakonas S. 2015. The Notch-mediated hyperplasia circuitry in Drosophila reveals a Src-JNK signaling axis. *Elife* **4**:e05996. doi:10.7554/eLife.05996

Huang L, Muthuswamy SK. 2010. Polarity protein alterations in carcinoma: a focus on emerging roles for polarity regulators. *Curr Opin Genet Dev* **20**:41–50. doi:10.1016/j.gde.2009.12.001

Huggins CJ, Mayekar MK, Martin N, Saylor KL, Gonit M, Jailwala P, Kasoji M, Haines DC, Quiñones OA, Johnson PF. 2015. C/EBPγ Is a Critical Regulator of Cellular Stress Response Networks through Heterodimerization with ATF4. *Mol Cell Biol* **36**:693–713. doi:10.1128/MCB.00911-15

Igaki T, Pagliarini RA, Xu T. 2006. Loss of cell polarity drives tumor growth and invasion through JNK activation in Drosophila. *Curr Biol* **16**:1139–1146. doi:10.1016/j.cub.2006.04.042

Igaki T, Pastor-Pareja JC, Aonuma H, Miura M, Xu T. 2009. Intrinsic tumor suppression and epithelial maintenance by endocytic activation of Eiger/TNF signaling in Drosophila. *Dev Cell* **16**:458–465. doi:10.1016/j.devcel.2009.01.002

Ji Z, Kiparaki M, Folgado V, Kumar A, Blanco J, Rimesso G, Chuen J, Liu Y, Zheng D, Baker NE. 2019. Drosophila RpS12 controls translation, growth, and cell competition through Xrp1. *PLoS Genet* **15**:e1008513. doi:10.1371/journal.pgen.1008513

Katheder NS, Khezri R, O'Farrell F, Schultz SW, Jain A, Rahman MM, Schink KO, Theodossiou TA, Johansen T, Juhász G, Bilder D, Brech A, Stenmark H, Rusten TE. 2017. Microenvironmental autophagy promotes tumour growth. *Nature* **541**:417–420. doi:10.1038/nature20815

Külshammer E, Mundorf J, Kilinc M, Frommolt P, Wagle P, Uhlirova M. 2015. Interplay among Drosophila transcription factors Ets21c, Fos and Ftz-F1 drives JNK-mediated tumor malignancy. *Dis Model Mech* **8**:1279–1293. doi:10.1242/dmm.020719

Kusch T, Florens L, Macdonald WH, Swanson SK, Glaser RL, Yates JR, Abmayr SM, Washburn MP, Workman JL. 2004. Acetylation by Tip60 is required for selective histone variant exchange at DNA lesions. *Science* **306**:2084–2087. doi:10.1126/science.1103455

Lee C-H, Kiparaki M, Blanco J, Folgado V, Ji Z, Kumar A, Rimesso G, Baker NE. 2018. A Regulatory Response to Ribosomal Protein Mutations Controls Translation, Growth, and Cell Competition. *Dev Cell* **46**:456-469.e4. doi:10.1016/j.devcel.2018.07.003

Link N, Kurtz P, O'Neal M, Garcia-Hughes G, Abrams JM. 2013. A p53 enhancer region
regulates target genes through chromatin conformations in cis and in trans.
Genes Dev **27**:2433–2438. doi:10.1101/gad.225565.113
Loubiere V, Papadopoulos GL, Szabo Q, Martinez A-M, Cavalli G. 2020. Widespread
activation of developmental gene expression characterized by PRC1-dependent
chromatin looping. *Sci Adv* **6**:eaax4001. doi:10.1126/sciadv.aax4001
Maraver A, Fernández-Marcos PJ, Herranz D, Muñoz-Martin M, Gomez-Lopez G,
Cañamero M, Mulero F, Megías D, Sanchez-Carbayo M, Shen J, Sanchez-Cespedes
M, Palomero T, Ferrando A, Serrano M. 2012. Therapeutic effect of γ-secretase
inhibition in KrasG12V-driven non-small cell lung carcinoma by derepression of
DUSP1 and inhibition of ERK. *Cancer Cell* **22**:222–234.
doi:10.1016/j.ccr.2012.06.014
McCaffrey LM, Montalbano J, Mihai C, Macara IG. 2012. Loss of the Par3 polarity protein
promotes breast tumorigenesis and metastasis. *Cancer Cell* **22**:601–614.
doi:10.1016/j.ccr.2012.10.003
Muzzopappa M, Murcia L, Milán M. 2017. Feedback amplification loop drives malignant
growth in epithelial tissues. *Proc Natl Acad Sci USA* **114**:E7291–E7300.
doi:10.1073/pnas.1701791114
Ntziachristos P, Lim JS, Sage J, Aifantis I. 2014. From fly wings to targeted cancer
therapies: a centennial for notch signaling. *Cancer Cell* **25**:318–334.
doi:10.1016/j.ccr.2014.02.018
Ohsawa S, Sugimura K, Takino K, Xu T, Miyawaki A, Igaki T. 2011. Elimination of
oncogenic neighbors by JNK-mediated engulfment in Drosophila. *Dev Cell*
20:315–328. doi:10.1016/j.devcel.2011.02.007
Pagliarini RA, Xu T. 2003. A genetic screen in Drosophila for metastatic behavior. *Science*
302:1227–1231. doi:10.1126/science.1088474
Pascual J, Jacobs J, Sansores-Garcia L, Natarajan M, Zeitlinger J, Aerts S, Halder G,
Hamaratoglu F. 2017. Hippo Reprograms the Transcriptional Response to Ras
Signaling. *Dev Cell* **42**:667-680.e4. doi:10.1016/j.devcel.2017.08.013
Pinal N, Martín M, Medina I, Morata G. 2018. Short-term activation of the Jun N-terminal
kinase pathway in apoptosis-deficient cells of Drosophila induces tumorigenesis.
Nat Commun **9**:1541. doi:10.1038/s41467-018-04000-6
Ranganathan P, Weaver KL, Capobianco AJ. 2011. Notch signalling in solid tumours: a
little bit of everything but not all the time. *Nat Rev Cancer* **11**:338–351.
doi:10.1038/nrc3035
Reddy KL, Rovani MK, Wohlwill A, Katzen A, Storti RV. 2006. The Drosophila Par domain
protein I gene, Pdp1, is a regulator of larval growth, mitosis and endoreplication.
Dev Biol **289**:100–114. doi:10.1016/j.ydbio.2005.10.042
Reinke AW, Baek J, Ashenberg O, Keating AE. 2013. Networks of bZIP protein-protein
interactions diversified over a billion years of evolution. *Science* **340**:730–734.
doi:10.1126/science.1233465
St Johnston D, Ahringer J. 2010. Cell polarity in eggs and epithelia: parallels and
diversity. *Cell* **141**:757–774. doi:10.1016/j.cell.2010.05.011
Sykiotis GP, Bohmann D. 2008. Keap1/Nrf2 signaling regulates oxidative stress
tolerance and lifespan in Drosophila. *Dev Cell* **14**:76–85.
doi:10.1016/j.devcel.2007.12.002
Thurmond J, Goodman JL, Strelets VB, Attrill H, Gramates LS, Marygold SJ, Matthews BB,
Millburn G, Antonazzo G, Trovisco V, Kaufman TC, Calvi BR, FlyBase Consortium.
2019. FlyBase 2.0: the next generation. *Nucleic Acids Res* **47**:D759–D765.
doi:10.1093/nar/gky1003
Toedling J, Skylar O, Sklyar O, Krueger T, Fischer JJ, Sperling S, Huber W. 2007. Ringo--an
R/Bioconductor package for analyzing ChIP-chip readouts. *BMC Bioinformatics*
8:221. doi:10.1186/1471-2105-8-221

Toggweiler J, Willecke M, Basler K. 2016. The transcription factor Ets21C drives tumor
growth by cooperating with AP-1. *Sci Rep* **6**:34725. doi:10.1038/srep34725
Turkel N, Sahota VK, Bolden JE, Goulding KR, Doggett K, Willoughby LF, Blanco E,
Martin-Blanco E, Corominas M, Ellul J, Aigaki T, Richardson HE, Brumby AM.
2013. The BTB-zinc finger transcription factor abrupt acts as an epithelial
oncogene in Drosophila melanogaster through maintaining a progenitor-like cell
state. *PLoS Genet* **9**:e1003627. doi:10.1371/journal.pgen.1003627
Uhlirova M, Bohmann D. 2006. JNK- and Fos-regulated Mmp1 expression cooperates
with Ras to induce invasive tumors in Drosophila. *EMBO J* **25**:5294–5304.
doi:10.1038/sj.emboj.7601401
Uhlirova M, Jasper H, Bohmann D. 2005. Non-cell-autonomous induction of tissue
overgrowth by JNK/Ras cooperation in a Drosophila tumor model. *Proc Natl*
Acad Sci USA **102**:13123–13128. doi:10.1073/pnas.0504170102
Wu M, Pastor-Pareja JC, Xu T. 2010. Interaction between Ras(V12) and scribbled clones
induces tumour growth and invasion. *Nature* **463**:545–548.
doi:10.1038/nature08702
Xue B, Krishnamurthy K, Allred DC, Muthuswamy SK. 2013. Loss of Par3 promotes
breast cancer metastasis by compromising cell-cell cohesion. *Nat Cell Biol*
15:189–200. doi:10.1038/ncb2663
Ye J, Rawson RB, Komuro R, Chen X, Davé UP, Prywes R, Brown MS, Goldstein JL. 2000.
ER stress induces cleavage of membrane-bound ATF6 by the same proteases
that process SREBPs. *Mol Cell* **6**:1355–1364. doi:10.1016/s1097-
2765(00)00133-7
905
906

 FIGURE LEGENDS

Figure 1. Notch-based neoplastic growth paradigms in *Drosophila* wing discs

A-D. 3rd instar wing imaginal discs at precisely 5 days after egg-laying either wild-

type (WT; A), overexpressing activated Notch (N; B), mutant for *scrib* (S; C), or

combining overexpressed Notch and *scrib* mutation (NS; D) and marked for E-Cad

(blue) and Mmp1 (red). **A,B&D.** MARCM clones (positively marked by GFP; green)

of the indicated genotypes: expressing only GFP (A; WT), expressing Nicd & GFP

(B; N), and expressing Nicd & GFP and mutant for *scrib* (C; NS). **C.** Discs fully

mutant for *scrib*.

E-F. Differentially expressed genes as compared to WT in the different growth

paradigms N (green), S (red), and NS (blue) identified by RNA-Seq. This color code,

green for N, red for S, and blue for NS is used in all figures. **E.** Heatmap of gene

expressions after unsupervised clustering. **F.** Venn diagram of up-regulated and

down-regulated genes in N, S, and NS.

G. Semi-quantitative RT-PCR of the indicated genes represented as fold change

compared to WT (grey) in the different N (green), S (red), and NS (blue) growth

paradigms and normalized to *Atc5C* expression. Biological triplicates, standard error

to the mean (s.e.m.) is shown.

Figure 2. GO term analysis of the N, S, and NS transcriptomes

A. Enrichment diagram as measured by adjusted p-value for selected GO terms (full

list as supplemental material) and represented as bars for N (green), S (red), and NS

(blue). GO terms color reflect whether they are shared or specific: shared by all

(black), common N&S (orange: mix of green and red), common N&NS (dark green:

mix of green and blue), common S&NS (purple: mix of red and blue).

B. Venn diagram showing the domains of overlap of GO terms identified

(significantly enriched) in N, S, and NS.

Figure 3. Polarity loss redirects the transcriptional output of Notch during

neoplastic growth

A. Experimental set-up to identify the Notch Direct Targets genes (NDTs): Genes up-

regulated in N, S, or NS (Transcriptomic), and located within 20kb of a Su(H) binding

site (ChIP).

B. Venn diagram showing the NDTs overlap in N, S, and NS, showing core Notch

responses, but also significant condition specific NDTs.

C. Overlap of the Su(H) binding sites identified by ChIP in N, S, and NS, showing

that almost all S and NS Su(H) peaks are also found in N. The overlap is shown in

white. Numbers are slightly different because in the Su(H) peaks calling protocol, in

some rare cases, some peaks can be split between conditions where one peak in one

condition would overlap with two peaks in the other.

D. Genome Viewer snapshots of several NDTs (shown in purple) such as the NS

specific *Act87E* and *Wnt10* (but also *yellow-e* and *Ir87a*), the common *Ets21C*, and

the N/NS NDT *upd3*. For each condition, the Su(H) ChIP enrichment is shown in the

upper lane, and the Su(H) peaks identified are represented by the blocks underneath.

Figure 4. Identification of potential transcriptional modules mediating N, S, and

NS growth

A. Venn diagram of significant transcription factors (TFs) identified by iRegulon as

potential key mediators for the expression of the N (in the green circle), S (red circle),

and NS (blue circle) up-regulated genes. Fed with lists of co-regulated genes, and

analyzing the genomic features in the vicinity of the transcription start sites of these

genes, iRegulon identifies potential groups of TFs and DNA-binding factors, that are

enriched in the dataset of regulatory sequences, and could thus represent potential

mediators of the N, S, and NS transcriptomes. TFs were color-coded according to

their molecular class and/or belonging to the similar regulon (see supplemental tables

S8-15). Numbers represent the number of TFs identified. See also Figure S2 for the

iRegulon analyses of the NDTs, and the detailed lists of both iRegulon analyses in

Supplemental tables S8 to S15.

B. Heat map for the expression of some of the transcription factors identified by

iRegulon and grouped as "oncogenic module", "basic Leucine Zippers", "DNA

damage and genome integrity".

Figure 5. Neoplastic growth is mediated by the "oncogenic module" and by a

diverse network of bZIP transcription factors including Xrp1

A-H. 3rd instar wing imaginal discs expressing *GFP*, an activated form of Notch

(*Nicd*) and an RNAi for *scrib* under the control of the *Bx-Gal4* driver (dorsal wing

pouch) and stained for GFP (green, white A'-H') to assess tissue overgrowth and for

Mmp1 (red, white A''-H'') to assess tissue invasiveness. Discs also expressed under

Bx-Gal4 control the either *UAS Bsk DN* (B) or the indicated RNAi constructs:

w[HMS00045] (A; Ctrl), *yki[KK109756]* (C), *crc[JF02007]* (D),

CG6272[HMS00057] (E), *Xrp1[HMJ21189]* (F), *maf-S[HMS02020]* (G),

pdp1[HMS02030] (H). Representative discs are shown.

I-K. Quantification of the overgrowth of the GFP territory in the indicated genotypes.

Results are shown as percentage of discs. In blue are shown the discs smaller than the

average of controls (size < Average size of controls – 1.5x StDev of controls). In

green are shown discs that are the same size as the controls average or bigger. Results

for the "DNA damage module" in **I**, for the "oncogenic module" in **J**, and for the

"AP-1 and bZIP module" in **K**. Green arrowheads indicate modification. Star indicate

a genotype with misshapen and "sick" discs. Sample size 13 – 17 discs.

For more details of RNAi lines either from the TRiP collection (labelled with trip

superscript) or from the Vienna collection (labelled with KK superscript) are provided

in supplemental information.

I'-K'. Quantification of the Mmp1 intensity in the indicated genotypes. Results are

shown as percentage where discs were classified to fall in three categories: High

Mmp1 staining intensity (red, similar to that shown in A''), Low Mmp1 staining

intensity (blue, similar to that shown in C or H), No Mmp1 staining (light blue,

similar to that shown in B''). Results for the "DNA damage module" in **I'**, for the

"oncogenic module" in **J'**, and for the "AP-1 and bZIP module" in **K'**. Red

arrowheads indicate modification. Star indicate a genotype with misshapen and "sick"

discs. Sample size 13 – 17 discs.

For more details of RNAi lines either from the TRiP collection (labelled with trip

superscript) or from the Vienna collection (labelled with KK superscript) are provided

in supplemental information.

SUPPLEMENTAL INFORMATION

1001

SUPPLEMENTAL MATERIAL AND METHODS

Drosophila genetics

Overexpression lines tested in the *Bx-Gal4, UAS GFP;; UAS Nicd, UAS*

scribHMS01490 screen were UAS GFP, UAS bskK53R [20.1a] BL#9311, and UAS

SOD CAT (gift from P. Leopold). RNAi lines used are listed in the table below with

an indication of the labels used in Fig. 5I-K&I'-K' and whether pictures are shown in

Fig. 5A-H. TRiP collection lines have a stock BL#, and Vienna collection lines have a

stock v#.

1010

Gene	RNAi ID	Stock #	Label in Fig. 5 I-K and I'-K'	Discs shown in Fig. 5 A-H
Atf-2	HMC05118	BL#60124	Atf2 trip	
Atf3	JF02303	BL#26741	Atf3 trip	
Atf6	JF02109	BL#26211	Atf6 trip	
ATM/tefu	KK100008	v#108074	ATM KK	
ATR/mei-P26	HMC04662	BL#57268	ATR trip	
CG6272	KK110056	v#101871	CG6272 KK	
CG6272	HMS00057	BL#33652	CG6272 trip	Fig. 5E
CG7786	HMC05169	BL#62162	CG7786 trip	
cnc	HMS00650	BL#32863	cnc trip	
crc	JF02007	BL#25985	crc trip	Fig. 5D
CrebB-17A	HMJ30249	BL#63681	CrebB-17A trip	
Ets21C	HMS01989	BL#39069	Ets21C trip	
ftz-f1	KK108995	v#104463	ftz-f1 KK	
gt	HMS01105	BL#34631	gt trip	
Jra	JF01184	BL#31595	Jra trip	
kay	HMS00254	BL#33379	kay trip	
maf-S	HMS02020	BL#40853	maf-S trip	Fig. 5G
p53	HMS02286	BL#41720	p53 trip	
pdp1	HMS02030	BL#40863	pdp1 trip	Fig. 5H

Stat92E	HMS00035	BL#33637	Stat92E trip	
vri	HMS02029	BL#40862	vri trip	
w	HMS00045	BL#33644	Ctrl	Fig. 5A
Xrp1	HMS00053	BL#34521	Xrp1 trip#1	
Xrp1	HMJ21189	BL#51054	Xrp1 trip#2	Fig. 5F
Xrp1	HMJ22533	BL#60356	Xrp1 trip#3	
yki	KK109756	v#111001	yki KK	Fig. 5C

1011

1012

Primers

qPCR primers:

Act5C_F: GAGCGCGGTTACTCTTTCAC

Act5C_R: ACTTCTCCAACGAGGAGCTG

Act87E_F: GTCCACCGCAAGTGCTTCTA

Act87E_R: TTTCTTTGGATGGCAGGGCA

Atf3_F: CAGCATGGCAACATTGGGAC

Atf3_R: ATGAAGGCAGTGGCTGAGTC

Diap1_F: CAGCCACACGCATCTTCAAC

Diap1_R: ACTTTGTCACAGAGGAGGCG

E2f1_F: ACAGAATCCTCGCCTCCAAC

E2f1_R: GACTGCTGCCGTAGCCTATT

Ets21C_F: CTGCTCGCTGATTCGTCCAA

Ets21C_R: TAGGCATACCGCTTTCCGTG

ftz-f1_F: ATTCCTGGTCGGACATGCTT

ftz-f1_R: TTCATGCAGACATAGTCGCCC

mxc_F: ACTAGAGGAGGAGCAGCGAA

mxc_R: CTAGTGGACAGCGGCGTATT

p38a_F: TACGGACAGGTGTCAAAGGC

p38a_R: CAGCGATCCATTAGCGGGAT

p53_F: TGCGTGTGTTCCTTTGCTTC

p53_R: GTTCAGGGGGACTACAACGG

puc_F: ATTGACCTCGCCGCCAATTA

puc_R: ATTCCGCTTGAACAGAGCCA

1037	sd_F:	AGGGTCCACAGAATGCGTTT
1038	sd_R:	TCGCTTTCCACCTTCTCCAC
1039	Sdr_F:	CGCTCCCTCAATCCCAAAGT
1040	Sdr_R:	ACAACGTCCATCAGCCAGTT
1041	Ser_F:	GCACGAATCTCTGGTGTGGA
1042	Ser_R:	TAGATTTGGCTGGCAGTCGG
1043	wg_F:	GCAGTCTGGTCTGGTCTACG
1044	wg_R:	ATTGTGCGGGTTCAGTTGGA
1045	Wnt10_F:	AATGGCATCGGTGGAACTGT
1046	Wnt10_R:	CAGCGTCTTGCGATTGATGG
1047		
1048	qChIP primers:	
1049	E(spl)mβ_F:	AAGTCGGAGCTTTGAATGAG
1050	E(spl)mβ_R:	CAAGTCATTTTATTGCCCTCAC
1051	E(spl)m5_F:	GTTTCCGCAGGTCCAGTTAC
1052	E(spl)m5_R:	GTTTGATGTTCACGCTGCTG
1053	white_F:	CGAAGGACGTTGACACATTG
1054	white_R:	GAATTGCCGCTTTTTCTCAC
1055	DDC_F:	AAGTGGGATTTGCCAGTGAC
1056	DDC_R:	TGCTGGTGAACTTTGACTGC
1057	CG42808_F:	CTCGTTAAGAGCAACTGCGA
1058	CG42808_R:	GTGAGAACTCCGAATCGAGG
1059	CG6191_F:	CGAAAAATGCGGACGATTCC
1060	CG6191_R:	CCCACCAATCTAGGGTTTCA
1061	Ilp8_F:	TCATCTCCGGTGTCTGACTT
1062	Ilp8_R:	AAAGAATTGGCTGCGGAAGA
1063		
1064		

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. Features of the Notch Direct Targets (NDTs) in N, S, and NS (relates

to Fig. 3)

A. Genome-wide localization of the Su(H) ChIP enrichment peaks with respect to

genome features, showing that the major changes were found in NS where Su(H)

"peaks" were more prevalent in introns but less prevalent at promoter regions than in

N and S.

B. Heatmaps for the expression of the different NDTs in WT, N, S, and NS. From left

to right are presented the N, S, NS, and finally All NDTs, highlighting that NDTs

could be transcriptionally up-regulated in more than in one condition.

C. Genome browser view of the whole left arm of the 2nd chromosome, and showing

the Su(H) ChIP enrichment (upper rows)) and the intervals called as Su(H) peaks

(lower rows) in N (green), NS (blue), and S (red). Note the higher number of peaks in

N, and the rarity of NS, or S peaks not found in N.

1079

Figure S2. Identification of potential transcriptional modules mediating N, S,

and NS growth (relates to Fig. 4)

A. Venn diagram of significant transcription factors identified by iRegulon in N, S,

and NS Notch Direct Targets, and color-coded according to their molecular class.

Numbers represent the number of transcription factors identified in each group. See

also the detailed lists of all iRegulon analyses in Supplemental tables S8 to S15.

1086

1087

SUPPLEMENTAL TABLES

Table S1-3. Differentially expressed genes in N, S, and NS identified by DESeq

(related to Fig. 1). Columns are:

FBgn_ID: Unique FlyBase gene ID

Symbol: Current FlyBase gene symbol

qval: adjusted p-value for multiple testing

logFC: log2 of the Fold Change "Condition N, S, or NS" / "Control WT"

1095

Table S4. Gene Ontology (GO) enrichment in N, S, and NS (related to Fig. 2).

Columns are:

experiment: type of comparison

GO_term: GO term number

description: GO term description

intersect.size: number of genes in query within the GO term

n.pw: total number of genes within the given GO term

pval: p value after hypergeometric test

qval: adjusted pvalue for multiple testing

genes: genes in query falling in the specific GO term

1106

Table S5. Curated Gene Ontology (GO) enrichment in N, S, and NS (related to Fig.

2). Columns are:

experiment: type of comparison

GO_term: GO term number

description: GO term description

intersect.size: number of genes in query within the GO term

n.pw: total number of genes within the given GO term

pval: p value after hypergeometric test

qval: adjusted pvalue for multiple testing

genes: genes in query falling in the specific GO term

1117

Table S6. Su(H) ChIP enrichment peaks coordinates in N, S, and NS (related to Fig.

3). Columns are:

Exp: N, S, or NS

Chr: Chromosome arm

MIN: smallest peak coordinate

MAX: biggest peak coordinate

1124

1125

Table S7. All Notch Direct Targets (NDTs) ordered by genomic position. This table

includes an indication whether the genes are transcriptionally upregulated or have a

Su(H) peak in the vicinity in each N, S, and NS condition. Columns are:

N/NS/S: NDT in the corresponding condition

Type: NDT in different conditions.

FBgn_ID: Unique FlyBase gene ID

SYMBOL: Current FlyBase gene symbol

K_ARM: Chromosome arm location of the gene

MIN (gene pos): smallest gene coordinate

MAX (gene pos): biggest gene coordinate

STRAND: +1 or -1

N Fold: Log2 Fold Change in gene expression N/WT (n.s. not significant)

N ChIP: Su(H) ChIP enrichment peak within 20kb in N (green yes, red no)

NS Fold: Log2 Fold Change in gene expression NS/WT (n.s. not significant)

NS ChIP: Su(H) ChIP enrichment peak within 20kb in NS (green yes, red no)

S Fold: Log2 Fold Change in gene expression S/WT (n.s. not significant)

S ChIP: Su(H) ChIP enrichment peak within 20kb in S (green yes, red no)

1143

1144 **Table S8-10.** iRegulon analyses of the significantly upregulated genes in N, S, and

1145 NS (related to Fig. 4). Analyses were performed using the 6K-PWM and 10kb

1146 upstream and downstream set-ups.

1147

1148 **Table S11.** Curated iRegulon analyses corresponding to Tables S8-10 (related to Fig.

1149 4)

1150

1151 **Table S12-14.** iRegulon analyses of the Notch Direct Targets in N, S, and NS (related

1152 to Fig. S2). Analyses were performed using the 6K-PWM and 10kb upstream and

1153 downstream set-ups.

1154

1155 **Table S15.** Curated iRegulon analyses corresponding to Tables S12-14 (related to

1156 Fig. S2)

1157

1158 **Table S16.** Xrp1 target genes. Xrp1 regulated genes identified in Lee et al. (2018)

1159 (positively and negatively) and in Baillon et al. (2018). In bold are highlighted genes

1160 found in both studies. Columns are:

1161 FBgn_ID: Unique FlyBase gene ID

1162 Symbol: Current FlyBase gene symbol

Logeay et al. Figure 2

A Localisation of the Su(H) binding on the genome in each condition (in %)

Legend:
- Promoter
- 5' UTR
- 3' UTR
- Exon
- Intron
- Downstream
- Intergenic

B Heat-maps of the expression of N, S, or NS NDTs (left to right) in each condition

N NDTs   S NDTs   NS NDTs   All NDTs

C Genome browser view of chromosome 2L with Su(H) peak calling in N, S, and NS

N ChIP
peak calling
NS ChIP
peak calling
S ChIP
peak calling
Gene (+)
Coordinates
Gene (−)

A