Loss of epithelial polarity redirects Notch signaling and triggers a Xrp1 response

during neoplastic growth in *Drosophila*

Rémi Logeay[1], Charles Géminard[1], Patrice Lassus[2], Diala Kantar[1], Lisa Héron-

Milhavet[1], Bettina Fischer[3], Sarah J. Bray[4], Jacques Colinge[1], and Alexandre

Djiane[1]*

[1] IRCM, Inserm, Univ Montpellier, ICM, Montpellier, France

[2] IRCM, Inserm, Univ Montpellier, ICM, CNRS, Montpellier, France

[3] Dept. of Genetics, University of Cambridge, Cambridge, United Kingdom

[4] Dept. of Physiology Development and Neuroscience, University of Cambridge,

Cambridge, United Kingdom

*Author for correspondence: IRCM, Inserm U1194

15   208, rue des Apothicaires

16   34298 Montpellier, Cedex, France

17   tel: +33 (0) 467 612 441

18   email: alexandre.djiane@inserm.fr

KEY WORDS:

Notch signaling, *Drosophila*, epithelial polarity, neoplasia, cell competition

SUMMARY

Aggressive neoplastic growth can be initiated by a limited number of genetic

alterations, such as the well-established cooperation between loss of cell architecture

and hyperactive signaling pathways. However, our understanding of how these

different alterations interact and influence each other remains very incomplete. Using

Drosophila paradigms of imaginal wing disc epithelial growth, we have monitored

the changes in Notch pathway activity according to the polarity status of cells and

show that epithelial polarity changes directly impact the transcriptional output of the

Notch pathway. We further provide evidence that this Notch redirection is not

mediated by new genomic binding regions for Su(H), but relies on the cooperation

with Su(H) of a combination of oncogenic transcription factors. Our work highlights

in particular the role of the stress response CEBPG homologue CG6272/Irbp18 and of

its partner Xrp1 suggesting that cellular competition, or parts of the cellular

competition program, are co-opted during neoplastic growth.

INTRODUCTION

In multicellular organisms, cells constantly integrate many stimuli, to adapt their behaviors and responses. How cells achieve this integration is a fundamental question in biology and its disruption underlies many pathologies. Cancer genomes harbor many mutations and understanding how different mutations interact within a given cell is a major goal in cancer research.

Epithelial cells represent the basic unit of many organs. Their apico-basal (A/B) polarity is controlled by the asymmetric segregation of highly conserved protein complexes such as the Scrib/Dlg/Lgl complex (Bilder et al., 2003; Coopman and Djiane, 2016; St Johnston and Ahringer, 2010). The far-reaching effects of A/B polarity is epitomized by the observation that many tumors of epithelial origin exhibit impaired polarity, and that several viral oncoproteins target polarity complexes (Banks et al., 2012; Huang and Muthuswamy, 2010).

Studies in human cell lines and in animal models have also suggested a contributing role of polarity alterations to tumor formation. For instance, mutations in the baso-lateral determinant *SCRIB1* have been shown to control proliferation and invasion in MCF-10A human mammary cells (Cordenonsi et al., 2011). Similarly in *Drosophila, scrib*, *dlg* or *lgl* mutations, result in multilayered overgrowth of larval epithelial imaginal discs (Bilder et al., 2003; Bunker et al., 2015). However, this uncontrolled growth is at least partly achieved because larvae exhibiting *scrib* mutations fail to undergo proper metamorphosis and imaginal discs grow for an extended period. Indeed, *scrib* mutant cells actually grow slower than wild-type cells and are eliminated by wild-type neighbors (Cordero et al., 2010; Igaki et al., 2009, 2006; Ohsawa et al., 2011). Interestingly, this is reversed when additional mutations are

introduced, such as constitutive activation of signaling pathways (e.g. Ras or Notch),

converting *scrib* mutant cells into aggressive, invasive and hyperproliferative cells

(Brumby and Richardson, 2003; Pagliarini and Xu, 2003). Similar observations have

been reported in mouse, where Notch or Ras activation and *Par3* depletion cooperate

to generate aggressive neoplasms in mouse mammary glands (McCaffrey et al., 2012;

Xue et al., 2013).

The Notch pathway is a highly conserved cell-signaling pathway mis-regulated in

several cancers (Ntziachristos et al., 2014; Ranganathan et al., 2011). Upon activation,

Notch receptors undergo two proteolytic cleavages to release their intra-cellular

domain or NICD, which enters the nucleus, binds to the Notch pathway specific

transcription factor CSL (Rbpj in mammals; Suppressor of Hairless, Su(H) in

Drosophila), and converts it from a repressor to an activator to turn on the

transcription of specific target genes (Bray, 2016). These Notch direct target genes

differ depending on cell type and account for the variety of outcomes triggered by

Notch activity. Increased Notch activity has been associated with several epithelial

cancers such as non-small-cell lung carcinomas (Maraver et al., 2012; Ntziachristos et

al., 2014), but in animal models, the sole increase in Notch activity either promotes

differentiation or only results in benign over-proliferation (hyperplasia) (Brumby and

Richardson, 2003; Djiane et al., 2013; Fre et al., 2005; Ho et al., 2015; McCaffrey et

al., 2012). However, as mentioned previously, Notch pathway activation cooperates

with loss of polarity to generate invasive neoplasms (Brumby and Richardson, 2003;

Ho et al., 2015; McCaffrey et al., 2012; Pagliarini and Xu, 2003).

So, while the cooperation between loss of cell architecture and hyperactive signaling

pathways is well established, the underlying mechanisms remain poorly understood. It

could merely reflect an additive effect where the consequences of both events

combine (the additive model). Alternatively, it could indicate a more profound

integration within epithelial cells where these two events impact on each other to

generate unique new behaviors (the redirective model). Using *Drosophila* paradigms

of imaginal wing disc epithelial growth, we have monitored the changes in Notch

pathway activity according to the polarity status of cells and show that epithelial

polarity changes directly impact the transcriptional output of the Notch pathway. We

further provide evidence that this Notch redirection is not mediated by new genomic

binding regions for Su(H), but relies on the cooperation with Su(H) of a combination

of transcription factors, such as basic leucine zipper (bZIP), whose activity is

triggered in response to JNK signaling during polarity loss, extending earlier reports

on the cooperation between oncogenic Ras and polarity loss (Atkins et al., 2016;

Külshammer et al., 2015; Uhlirova and Bohmann, 2006). Our work highlights in

particular the role of the stress response CEBPG homologue CG6272/Irbp18 and of

its partner Xrp1, key factors in mediating the loser fate during cell competition

(Baillon et al., 2018; Blanco et al., 2020; Ji et al., 2019), suggesting that cellular

competition, or parts of the cellular competition program, are co-opted during

neoplastic growth.

RESULTS

Notch activation and scrib mutation cooperate to promote neoplastic growth

In order to gain insights into the mechanisms underlying neoplastic growth, we first characterized the effects of Notch activation and *scrib* mutation mediated epithelial polarity impairment on wing disc growth. Using precisely controlled *Drosophila* larvae culture conditions (crowding and timing), we compared the phenotypes of wild-type (WT), Nicd overexpressing (N), *scrib* mutant (S), and Nicd overexpressing and *scrib* mutant (NS) 3rd instar wing imaginal discs at 6 days after egg laying at 25C. Reproducing our previous observations (Djiane et al., 2013), N discs overgrew compared to WT, but remained as monolayered epithelia with properly localized E-Cadherin-based adherens junctions, and represent a paradigm of hyperplastic-like growth (Fig. 1A&B). S discs were smaller than WT, but grew as unstratified mass of cells with weak uniform E-Cadherin (E-Cad). These discs however showed an extensive expression of the JNK signaling target Mmp1 (Fig. 1C), a metallo-protease implicated in the digestion of the extracellular matrix, indicative that *scrib-* cells activate JNK signaling and are prone to invasiveness (Igaki et al., 2006; Uhlirova and Bohmann, 2006). It is noteworthy that S larvae did not pupariate and if left to grow for longer, the S discs ultimately developed as massive overgrowths with very disrupted epithelial polarity, that invaded and fused with neighboring tissues such as other discs (Bilder et al., 2003). Strikingly, NS discs combined aspects of N and S discs. They were overgrown like N discs, but also expressed low levels of E-Cad and high levels of Mmp1 like S discs (Fig. 1D). These discs grew as multilayered tissues and were able to invade the surrounding tissues such as haltere discs, and represent therefore a paradigm for neoplastic-like growth.

Neoplastic and hyperplastic discs have different transcriptomes

The context of the scribble mutation converts the Notch-based *Drosophila* wing disc into a paradigm of neoplastic growth. Until now, the cooperation between activated Ras and *scrib* mutations has been studied extensively, mainly in the imaginal eye disc (Atkins et al., 2016; Cordero et al., 2010; Igaki et al., 2009; Katheder et al., 2017; Pagliarini and Xu, 2003; Toggweiler et al., 2016; Wu et al., 2010), but less attention has been given to the way that other activated pathways, such as Notch or Hedgehog, also cooperate with polarity loss beyond the initial studies (Brumby and Richardson,

2003; Pagliarini and Xu, 2003). Exploring how these other pathways cooperate is

important to evaluate the extent that are generally important for neoplasia or specific

to Ras signaling induced transformations.

First, we compared the transcriptomes of the different genetic conditions, to identify

genes whose expression was significantly different. Differential expression analysis

of genome-wide RNA-seq profiles (using DESeq with adjusted p-value for multiple

testing <0.05; (Anders and Huber, 2010) identified the cohorts of genes that were

significantly up-regulated or down-regulated in each condition compared to WT

controls (Fig. 1E&F and Supplemental Tables 1-3). The numbers were broadly

similar in each condition N (503 up; 663 dw), S (757 up; 1029 dw), and NS (1003 up;

991 dw). Using semi-quantitative qRT-PCR we then validated a subset of the

transcriptional changes and could confirm the specific differences. The N only

affected genes *E2f1*, *Sdr*, and *mxc*, were activated only in N discs and not in S or NS

while the S only gene *p38a* and the NS only genes *Act87E* and *Wnt10*, were activated

only in their respective conditions. In addition, *Ets21C*, *ftz-f1*, and *Atf3* were up-

regulated in all three conditions, as detected by the RNA-Seq profiles (Fig. 1G)

Comparing these data with previously published transcriptome analyses on similar or

related genetic backgrounds revealed significant overlap, validating our experimental

approaches. For instance, 174 of the 503 up-regulated genes in N, and 285 of the 663

down-regulated genes were also detected in our previous analysis using dual color

differential expression arrays (significant overlap p=1.11e-273, hypergeometric test,

(Djiane et al., 2013). Similarly, 676 of the 757 up-regulated genes in S were identified

in a previous analysis of *scrib* depleted discs (significant overlap p=4.90e-193;

(Bunker et al., 2015).

To gain insights into the nature of the mis-regulated genes we performed a GO term

analysis on the genes affected in N, S, and NS (p-value < 0.05). As expected from

their genetic composition, the N and NS tissues were over-represented for genes in

the Notch signaling pathway (GO:0007219) and the NS and S had changes in A/B

polarity (GO:0045197; GO0019991) (Fig. 2A and Supplemental Tables 4&5).

Focusing on the other most robust GO terms (q-value < 0.05) revealed common

alterations in all three growth paradigms which included cell adhesive properties

(GO:0030198 "extracellular matrix organization"; GO:0007155 "cell adhesion"),

ribosomal biology (GO:0042274/GO:0042273 "ribosomal small/large subunit

biogenesis") suggesting that protein synthesis might be affected, mis-regulated

metabolic processes, in particular with respect to glucose and glycolysis

(GO:0006002), or oxidative stress (GO:0006979 "response to oxidative stress"; Fig.

2A and Supplemental Tables S4&5).

Despite these similarities, there were also differences in the profiles. For example,

markers of increased proliferation were found in both NS and N, but not S, consistent

with the overgrowth phenotypes (e.g. "mitotic cytokinesis" and "mitotic spindle

organization"; GO:0000281; GO:0007052). The overgrowing behavior of NS tissues

may therefore be driven by the Notch activation (Fig. 2A and Supplemental Tables

S4&5) although it should be noted that N was enriched in additional mitosis related

categories such as "mitotic metaphase plate congression" or "centriole replication"

(GO:0007080; GO:0007099), suggesting a more robust proliferation signature in N

tissues. Similarly, both NS and S (but not N) showed characteristics of cell migration

(e.g. "border follicle cell migration"; GO:0007298) and cellular stress such as

"response to starvation" (GO:0042594), or "response to endoplasmic reticulum

stress" (GO:0034976) suggesting that these features in NS were likely contributed by

the *scrib* mutation (Fig. 2A and Supplemental Tables S4&5).

Notably there were several GO categories specific to NS including "positive

regulation of apoptotic signaling pathway" (GO:2001235) "negative regulation of

SAPK signaling cascade" (GO:0070303) and "mitotic G1/G2 DNA damage

checkpoint" (GO:0031571/GO:0007095). These results argue that the combined

Notch activation and polarity loss promoted the emergence of new cell behaviors and

responses, in particular related to DNA damage responses (Fig. 2A&B and

Supplemental Tables S4&5).

Distinguishing between the additive and redirective models of cooperation

The transcriptomic changes revealed that NS encompassed not only additive N and S

features but also the emergence of new behaviors. This raises the question of how the

defects in *Notch* and in *scrib* cooperate to produce these transcriptional consequences.

Two main models could be proposed for genetic cooperation: (1) an additive model,

in which the effect of each genetic change is independent and summed to produce the

phenotype, and (2) a redirective model, in which the two mutations influence each

other to change their effects. Since Notch signaling exerts a direct effect on

transcription, without intermediates (unlike Ras and RTK signaling for instance;

(Bray, 2016), our Notch-based paradigms may allow us to distinguish between the

two models. Indeed, the genes directly activated by Notch (Notch Direct Targets,

NDTs), should behave differently depending on whether the cooperation is additive or

redirective. In the additive model, genes directly regulated by Notch should be

insensitive to the polarity status of the cells and should remain the same in the

hyperplastic (Nicd only) and neoplastic (Nicd & *scrib-*). Alternatively, if the loss of

polarity affects Notch directly, the regulated targets should be, at least in part,

different.

Genes that are directly regulated by Notch (NDTs) should have the transcription

complex, containing Su(H) bound at their regulatory regions (Djiane et al., 2013). To

identify potential NDTs in N and NS, we thus monitored the genomic regions

occupied by the Su(H) transcription factor by genome-wide Chromatin Immuno-

Precipitation (ChIP) (Supplemental Table S6). These overlapped significantly with

our previous analysis of Notch induced overgrowth, suggesting that we have captured

all the robust regions of Su(H) enrichment.

Strikingly, there was a strong overlap in the Su(H) bound regions in both conditions:

almost all (416 out of a total of 464) NS Su(H) peaks overlap with peaks present in N

discs. This implies that the vast majority of Su(H) binding remained the same in N

and NS (Fig. 3C & S1C). The overlap was also important between N and S peaks

(447/554 S peaks overlapping with N peaks; Fig. 3C & S1C). These results suggests

that in NS neoplastic discs (and to a lesser extent in S discs), a minority of Su(H)

peaks represent new binding regions compared to hyperplastic N discs, and that the

new NS behaviors are not the consequence of general redistribution of the Notch

specific transcription factor Su(H).

We note that the majority of binding in all conditions is associated with intronic

regions (39.70% and 42.89% in N and NS respectively; Fig. S1A), which are

frequently the site of enhancers in *Drosophila* patterning and growth genes. There

was a subtle shift in the distribution in other regions with a decrease in Su(H) peaks

that were associated with promoter regions in NS discs (from 31.55% in N to 23.06%

in NS; Fig. S1A) complemented by an increase in Su(H) binding associated with

intergenic regions (from 15.51% in N to 20.47% in NS; Fig. S1A), but the

significance of this shift is unclear.

In order to estimate the Notch specific programs in N, S, and NS, we then intersected

the transcriptomic data with the Su(H) ChIP data, considering that upregulated genes

located within 20kb of a Su(H) peak were likely NDTs. Using this approach, we

identified similar numbers of NDTs in N (177) and NS (171) (Fig. 3A&B,

Supplemental Table S7). Again, there was substantial overlap with previous data,

with 64/177 NDTs in N conditions being identified in our previous study (significant

overlap p=5.89e-96, hypergeometric test (Djiane et al., 2013). When the N and NS

scenarios were compared, 69 genes were common to both and thus represent core

NDTs in the wing disc overgrowth. However, a significant proportion of NDTs

appeared specific for each condition: 108 for N, 102 for NS (Fig. 3B) which supports

the redirective model. Amongst the 102 NS-specific NDTs, only 24 were also NDTs

in S, arguing that the difference cannot be explained by specific contribution of the S

condition. Taking all of these comparisons into account, there appear to be 78 genes

that are NS-specific NDTs. Only a minority were associated with new Su(H) binding

regions: around the *87E* locus (*yellow-e3, yellow-e, Ir87a,* and *Act87E*), and the *94A*

locus (*CG18596, CG7059, CG13857,* and *CG13850*). Amongst these 78 NS specific

NDTs are p53, His2Av, and the Tip60 complex component Act87E (Kusch et al.,

2004), which have all been linked to DNA damage response and could therefore

account for the NS emergent behaviors identified in the GO term analysis (Fig. 2A

and Supplemental Tables S4&5). The Venn-diagrams representing the overlaps in N,

S, and NS, highlight the extent of additivity (the yellow circles encompassing

common NDTs) and the scale that there is redirection of the Notch pathway output

(red circles). Both losses and gains occur so that in the conditions where there is

neoplastic growth (NS), 78 new NDTs are acquired while 108 NDTs are no longer

present compared to N (and 70 compared to S). The observation that the NDTs differ

substantially between NS and N is in strong support of the redirective model.

Taken together, our results indicate that Notch activation and *scrib* mediated loss of

polarity cooperate both in an additive and redirective manner to promote neoplastic

growth, and that the redirection of the Notch pathway transcriptional output is not a

consequence of a general redeployment of Su(H) to new active sites. They argue that

alterations in the cell architecture can directly influence the outcome of the Notch

signaling pathway and support a model in which different genetic injuries within

tumor cells could thus influence each other to sustain neoplastic growth.

Identification of the transcriptional networks in our different growth paradigms

Using the NDT datasets, we sought to identify the factors that are required for the

transition from N hyperplastic to NS neoplastic growth. As the NS transcriptome

showed signatures consistent with "response to oxidative stress" (GO:0006979),

"cellular response to gamma radiation" (GO:0071480), and "DNA damage

checkpoints" (GO:0031571/0007095), we first asked whether interfering with such

pathways could block the growth and invasiveness of NS tissues.

To perform these genetic tests, we generated a stable fly line which overexpressed

Nicd and *scribRNAi* together with a *GFP* marker under the *BxGal4* driver (driving

expression in the pouch of the larval wing discs; *Bx>NS*), and monitored the size of

the overgrowth (GFP positive tissue), and its invasiveness potential (Mmp1

expressing cells; Fig. 5A). Blocking the oxidative stress response by overexpressing

the Reactive Oxygen Species (ROS) sponge CAT and SOD, or knocking down by

RNAi the expression of *ATM* or *ATR* (*tefu* or *mei-P26* in the fly), the two master

kinases mediating the early response to DNA damage (double or single strand breaks

respectively) did not have any significant effect on the NS overgrowth or the

expression of Mmp1 (Fig. 5I). Similarly, expression of RNAi or dominant negative

forms of the acute stress response and severe DNA damage major effector and NS

specific NDT *p53* could not modify the NS overgrowth phenotype (Fig. 5I). While we

cannot exclude that the tools used here were not strong enough, these results suggest

that even though activated in NS tissues, oxidative stress and DNA damage responses

were either not required to sustain NS growth, or that they could compensate for each

other converging ultimately on an as yet unidentified core response promoting NS

growth.

Next we sought to identify transcriptional factors that could account for the

cooperation between Notch and polarity loss. We used the iRegulon software to

identify the transcriptional network involved. When utilized to analyze genes

involved in sustaining the tumorous growth of *RasV12/scrib-* 3rd instar larval eye,

wing and leg discs, it highlighted the role of the Hippo pathway terminal effectors

Yki/Sd and their targets Myc, Crp, and Ftz-F1, the JNK pathway regulated AP-1

factors (in particular Atf3, Kay, and CEBPG), and the Jak/Stat pathway (Atkins et al.,

2016; Külshammer et al., 2015).

Implementing iRegulon on our step-wise Notch-based paradigms allowed us (i) to

identify transcriptional modules unique or shared between polarity loss only (S),

proliferation only (N) and proliferation plus invasiveness (NS), and (ii) to assess the

conservation of the "oncogenic" modules identified previously with Ras in a Notch-

driven neoplastic paradigm. We performed these analyses feeding iRegulon either

with the lists of up-regulated genes in N, S, and NS (Fig. 4A & Supplemental Tables

S7-11) or with the lists of NDTs (Fig. S2 & Supplemental Table S12-15). These

analyses identified "modules of transcription factors" likely co-regulating the genes

identified in N, S, or NS. Modules identified are presented as Venn diagram (Fig. 4A

& S4) to highlight common and specific programs in the different conditions. Feeding

either up-regulated genes (Fig. 4A) or NDTs (Fig. S2) identified similar modules

indicating that the Notch pathway redirection is mediated, at least in part, by

transcription factors that broadly affect the whole transcriptome. This is likely also

true for other neoplastic paradigms such as Ras, even though this remains to be

established. Importantly, iRegulon identified the Notch pathway dedicated

transcription factor Su(H) in the N and NS transcriptomes.

First, focusing on NS, which most resembles the *RasV12/scrib-* paradigms, our

analysis identified the same major nodes and oncogenic module as described

previously:

- AP-1 basic Leucine Zipper factors related to stress kinase signaling; *Mmp1* is a

canonical JNK targets and is highly expressed in NS growing discs (Fig. 1D)

- Stat92E of the Jak/Stat pathway; *upd3* coding for the upstream ligand of the Jak/Stat

pathway is indeed highly expressed in NS discs

- Ftz-F1 nuclear receptor

- basic Helix-Loop-Helix factors of the Myc family.

In NS transcriptome, we also identified a contribution of the E(spl) bHLH

transcriptional repressors. E(spl)-HLH genes are canonical Notch targets, and they are

robustly up-regulated in N and NS, in particular E(spl)mγ-HLH. In NS there were

however two noteworthy differences when focusing only on NDTs, rather than all

differentially transcribed genes. (1) The absence of the E(spl) factors, suggesting that

when considering the whole NS transcriptomes a significant of genes could be

associated with E(spl)-mediated regulation while this is no longer the case when

considering the more restricted number of NS NDTs. (2) The identification of Ewg,

the fly NRF1 homologue. In mammals, NRF1 is implicated in the regulation of key

metabolic and respiratory genes (Herzig et al., 2000).

The iRegulon analyses also suggested that the AP-1 bZIP, and the Stat92E signatures

in NS are contributed by S, since they are also detected in the S transcriptomes, while

the Su(H) signature is contributed by N. Interestingly, the bHLH Myc signature is

found in N and S as well as NS. Finally, iRegulon identified a signature for the

Polycomb chromatin silencers specifically in N. Such factors include Pho, a zinc

finger protein which binds to Polycomb Responsive Elements (PREs) and recruits

Polycomb complexes, and the three Polycomb Repressor Complex 1 components Psc,

Su(z)2, and l(3)73Ah. Recently, PRC1 has been associated with specific and

unexpected transcriptional activation at larval stages, raising the possibility that in N,

such genes, normally repressed at embryonic stages, become active (Loubiere et al.,

2020).

A polarity-loss oncogenic module

In order to validate the functional relevance of the transcription factors identified in

the Notch-driven neoplastic growth, we then asked whether their depletion by RNAi

could alter the growth and invasiveness of the NS tissue, using the *Bx>NS* fly line

described previously. We first focused our analysis on the the oncogenic module.

We confirmed earlier reports that blocking JNK activity by the overexpression of a

JNK dominant negative construct, strongly abolishes NS driven growth (GFP positive

tissue size; Fig. 5B&J) and invasiveness (Mmp1 expression; Fig. 5B&J'). As shown

in the *RasV12/scrib-* paradigms, RNAi mediated knock-down of the Jak/Stat pathway

terminal transcription factor *stat92E*, the JNK pathway component *Ets21C*, and to a

lesser extent *ftz-f1* (Atkins et al., 2016; Külshammer et al., 2015; Toggweiler et al.,

2016) strongly suppressed both growth (GFP) and invasiveness (Mmp1; Fig. 5J&J').

Unlike the RasV12 models, we did not identify any particular enrichment for

Sd/TEAD, the transcriptional factor mediating the effect of Yki and of the Hippo

pathway mediated growth in wing discs. However, impairing Yki activity (through

RNAi-mediated knock-down) strongly suppressed NS neoplastic behaviors (Fig.

5C&J).

Taken together these results suggest that independently of the oncogenic driver, Ras

or Notch, relatively similar tumorous transcriptional networks (AP-1/Yki/Stat/Ftz-f1)

are put in place during their cooperation with polarity loss (Atkins et al., 2016;

Külshammer et al., 2015). Given that these nodes were also identified in the S

transcriptome, we suggest that they might represent a polarity loss module

cooperating with oncogenic signaling pathways Ras or Notch (and likely other

pathways such as Hh as was initially reported; (Brumby and Richardson, 2003).

The CG6272/Xrp1 module is required for neoplastic growth

A striking feature of the iRegulon analyses was that the JNK module, which

contained classic basic leucine zipper (bZIP) transcription factors such as Jun or Fos,

was found in NS and S. As the bZIP factors identified belonged to sub-families

involved in different cellular responses including oxidative stress (Maf-S, Cnc,

(Sykiotis and Bohmann, 2008), polarity loss (Atf3, (Donohoe et al., 2018), ER stress

(Atf6, Crc, (Harding et al., 2003; Ye et al., 2000), DNA damage… they were potential

candidates to mediate the transcriptional switch. In order to identify which bZIP

factor/family is most relevant for NS, we therefore systematically depleted by RNAi

the different genes highlighted by iRegulon in this bZIP node, using the *Bx>NS* fly

line.

Amongst the different factors tested, we observed a significant suppression of the

invasiveness (Mmp1; Fig. 5K') but not of the growth (GFP; Fig. 5K) of NS tissues

after impairing the AP-1 factors *kay* and *Jra* (fly homologues of *FOS* and *JUN*

respectively), and *CrebB-17A*. Conversely, knocking down the DNA damage

response factor *maf-S* led to a dramatic reduction of the GFP outgrowth but without

affecting the invasiveness in which the tissue was expressing high levels of Mmp1

(Fig. 5G&K). It should be noted here that these *maf-S* depleted discs appeared sick

with a severely irregular and misshapen GFP domain (Fig. 5G&K, star), suggesting

that maf-S might be an essential gene, and that the size suppression might merely

reflect enhanced cell death. In contrast, we did not observe any significant effect

when *cnc*, the fly homologue of *NFE2L1* implicated in oxidative stress response and

excision repair (Han et al., 2012; Sykiotis and Bohmann, 2008), *Atf2*, *gt*, *CG7786*, *vri*

(Fig. 5K), were knocked-down. Neither were any consequences on the phenotype

with knock-down of *Atf6* and *crc* (the *ATF4* fly homologue), which have been both

linked to the unfolded protein response and ER stress (Fig. 5D&K; (Harding et al.,

2003; Ye et al., 2000).

The most complete suppression of the NS behaviors (both GFP overgrowth and

Mmp1 invasiveness) was observed after depletion of *Atf3*, *Pdp1* or CG6272 (Fig.

5E,H,K). Atf3 has recently been shown to control the expression of genes involved in

the maintenance of epithelial polarity, and to be specifically activated in polarity

deficient cells and required in the RasV12/scrib- overgrowth models (Atkins et al.,

2016; Donohoe et al., 2018). Pdp1 (the homologue of Hepatic Leukemia Factor -

HLF) has previously been linked to mitotic cell cycle and growth (Reddy et al., 2006),

and shown in the *RasV12/scrib-* paradigms to have modest effects on invasion, but

none on growth. The role of *CG6272* (a.k.a. *Irbp18*, the homologue of *CEBPG*) was

also highlighted in the *RasV12/scrib-* paradigms where it was shown to control

growth but not invasion (Atkins et al., 2016). In the context of Notch (*Nicd/scrib-*),

the role of *Pdp1* and *CG6272* appeared more essential.

bZIP factors have been shown to act as homo or heterodimers. Knocking down the

function of *crc/Atf4*, the classic bZIP partner of CG6272/CEBPG, did not have any

effect on NS tissue behaviors (Fig. 5D,E,K) whereas the knock-down of *Xrp1* very

efficiently suppressed the growth and Mmp1 expression in NS tissues. This suggests

that .CG6272/CEBPG could be acting in combination with its bZIP partner Xrp1 in

this context (Francis et al., 2016; Reinke et al., 2013). Although the cellular processes

controlled by CG6272/Xrp1 remain to be determined, the fact that Xrp1 is an early

p53 target gene (Akdemir et al., 2007; Link et al., 2013), raises the possibility that its

functions could be related to DNA/genome integrity maintenance. This would align

with the NS specific GO categories "cellular response to gamma radiation"

(GO:0071480), and "DNA damage checkpoints" (GO:0031571/0007095). Indeed,

Xrp1 has recently been linked to cell competition and shown to control the expression

445 of a subset of genes activated in response to Ribosomal protein haplo-insufficiency

446 including *Ets21C*, *upd3*, or *ilp8* in developing wing discs (Baillon et al., 2018; Boulan

447 et al., 2019; Ji et al., 2019). Strikingly, we observed a very strong overlap between the

448 Xrp1 transcriptional program (Supplemental Table S16) and the genes activated in NS

449 (60 out of the 171 Xrp1 positively regulated genes are also upregulated in NS;

450 hypergeometric test, p=4.71 10e-21), while a much more modest overlap could be

451 observed with N or S (overlap of 17 and 33 with p values of 1.12 10e-3 and 1.56 10e-

452 8 respectively). These results support the proposal that the CG6272/Xrp1 module is

453 functionally important in the acquisition of neoplastic growth.

DISCUSSION

In this study, using Notch-driven paradigms of epithelial overgrowth in *Drosophila* wing discs, we describe the molecular mechanisms underlying the cooperation between Notch and polarity loss during neoplasia. We show that epithelial polarity alterations redirect the transcriptional outcome of the Notch signaling pathway. We further show that this redirection occurs mainly on pre-existing Su(H) bound regions rather than new ones. Finally, we show that similarly to what was previously described for Ras signaling (Atkins et al., 2016), the cooperation between Notch signaling and polarity loss is controlled by a "tumor transcriptional network" centered around the AP-1/Stat/Yki transcription factors, and including the critical bZIP factors Pdp1 and CEPBG (CG6272/Irbp18). But our analysis uncovered a previously unreported role for the cell competition regulator, and Irbp18 binding partner Xrp1 (Baillon et al., 2018; Blanco et al., 2020; Boulan et al., 2019; Ji et al., 2019), raising the interesting prospect that neoplastic growth could be mediated at least in part, by co-opting cell competition.

While cancer genomes exhibit multiple mutations in cancer cells, their functional interactions remain difficult to monitor and model. This analysis in the genetically controlled *Drosophila* wing disc, supports a model in which the effects of different mutations do not just simply add up, but interact with each other leading to the emergence of new cell behaviors (redirective model). Indeed, even though the combination showed characteristics that could be attributed to either Notch (proliferation and mitosis), or to polarity loss (invasion/cell migration), it also exerted signs of emerging behaviors such as DNA damage response. Neoplastic tissues appear thus to experience many cellular stresses: DNA damage responses, but also ER and unfolded protein response, starvation, or oxidative stresses. However, even though present, these different stresses and in particular oxidative stress and DNA damage are not individualy necessary in the context of polarity loss as blocking them or the cellular response they promote (CAT/SOD overexpression, or inhibition of ATM/ATR) could not significantly suppress the NS tumorous behaviors. These observations suggest that the different stress pathways activated during polarity loss might all converge leading ultimately to the activation of a common core response.

While *Drosophila* and mouse models have demonstrated that overactive signaling

pathways cooperate with epithelial polarity impairment to generate neoplastic growth

(Brumby and Richardson, 2003; McCaffrey et al., 2012; Pagliarini and Xu, 2003; Xue

et al., 2013), the vast majority of studies seeking to understand the underlying

mechanisms, have focused primarily on the cooperation between activated RasV12

and *scrib* mutants, especially in *Drosophila* (Atkins et al., 2016; Cordero et al., 2010;

Igaki et al., 2009; Katheder et al., 2017; Pagliarini and Xu, 2003; Toggweiler et al.,

2016; Wu et al., 2010). Importantly, the current study, investigating the cooperation

between Notch and polarity, shows that many observations made for Ras can be

extended to Notch, suggesting that the paradigms used are not a Ras specificity but

might represent a more general tumor growth paradigm.

But even though we could highlight the involvement of a core "oncogenic module"

(Atkins 2016, Külshammer), there are specifics that are likely oncogene specific. In

the case of Ras, it was shown that Yki activity could reprogram Ras by promoting the

expression of the Ras pathway specific regulators Capicua and Pointed to promote

aggressive growth (Pascual et al., 2017). Both genes were either unaffected (*capicua*)

or downregulated (*pointed*) in NS Notch driven neoplastic paradigm, suggesting that,

even though Yki is clearly active (Fig. 5C), capicua and pointed regulations are

unlikely mediators here. Furthermore, in NS transcriptome, we identified a

contribution of the E(spl) bHLH transcriptional repressors, canonical Notch targets

(Bray, 2016), which represents thus a Notch specificity. However, the fact that motifs

for E(spl)-HLH repressors are found in the up-regulated transcriptome of NS and not

N could suggest that in NS, the repressive ability of E(spl)-HLH factors is

antagonized (even though their expression is not affected). It would be interesting to

explore further the link between NS and E(spl)-HLH-mediated repression, but due to

the high redundancy between the seven E(spl)-HLH factors (δ, γ, β, 3, 5, 7, 8) and

Dpn, the requirement of E(spl)-HLH-mediated repression in the Notch-driven

neoplasia could not be formally tested.

Amongst the most dramatic suppressors of the neoplastic growth are the two basic

leucine zipper transcription factors CG6272/Irbp18 (the fly homologue of CEBPG),

and Xrp1. Importantly, impairing with their function in wild-type tissues did not have

any effect, confirming observations made by other groups for CG6272 (Atkins et al.,
2016), supporting a model in which these genes are dispensable in healthy wild-type
cells, but become indispensable for tumor cells, hence representing a possible attack
strategy specifically targeting tumor cells, while sparing healthy tissues. In mammals,
CEBPG represents a major regulator of stress responses. It is recruited through its
interaction with the bZIP factor ATF4 at the level of cis-regulatory C/EBP:ATF
response elements to activate the expression of stress mitigation genes such as
glutathione biosynthesis pathway genes in the case of oxidative stress (Huggins et al.,
2015). Interestingly, we did not observe any effect knocking down *Crc*, the fly *Atf4*
homologue, suggesting that during NS overgrowth in *Drosophila*, CG6272/Irbp18
could be acting in combination with another partner. CG6272/Irbp18 also interacts
with Xrp1, and the heterodimer has been implicated in DNA repair (Akdemir et al.,
2007; Francis et al., 2016). We thus propose that the Irbp18/Xrp1 dimer is activated in
NS and is required to facilitate DNA repair and ensure genomic stability to prevent
catastrophic genotoxic effect upon the combined cellular stresses of S and replication
stress of N. Recently, *Xrp1* together with its binding partner CG6272/Irbp18 (but not
Atf4) has been shown to mediate a loser status in ribosomal genes deficient cells,
through the implementation of a specific transcriptional program (Baillon et al., 2018;
Blanco et al., 2020; Ji et al., 2019). This "loser state" promoter role of Xrp1/Irbp18 is
however in conflict with our observation that *Xrp1* is strongly upregulated in NS
overgrowing discs, and that *Xrp1* and *Irbp18* are required for the overgrowth and
invasive capacities of NS neoplastic cells. This could be reconciled by proposing that
in NS neoplastic discs, the role of the Xrp1/Irbp18 is modified by an as yet unknown
factor. Alternatively, and more attractively, it is also possible that in NS cells, cell
death is prevented as was shown for RasV12 expressing cells (Pinal et al., 2018), for
instance by upregulating DIAP1 (a Notch direct target, (Djiane et al., 2013) leading to
a perverted cell competition. There an incomplete "loser" program would be initiated,
but not fully implemented (no cell death), leading to the secretion of growth factors
(e.g. the Jak/Stat ligands upd…) that would act in an autocrine manner to further
promote their growth. These factors do not actually need to act cell autonomously,
since it was shown that at least in the *RasV12 / scrib-* paradigm, delaminating cells
cooperate with non-delaminating proliferating cells to sustain tumor growth
(Muzzopappa et al., 2017; Uhlirova et al., 2005; Wu et al., 2010). In NS, preventing
the losing/delaminating cells by impairing the Xrp1/Irbp18 nexus, would thus prevent

neoplastic growth. This intriguing possibility highlights that more studies are needed

to better understand the role of the Xrp1/Irbp18 module, its links to cell competition,

and to the growth of neoplastic tissues.

MATERIALS AND METHODS

Drosophila genetics

The different overgrowth paradigms were obtained by generating random clones in 3rd instar wing discs at high frequency as previously published in (Djiane et al., 2013). In brief, the *abxUbxFLPase; Act>y>Gal4, UAS GFP; FRT82B tubGal80* flies were crossed either to *FRT82B* (to generate Ctrl discs), or to *UAS-Nicd; FRT82B* (to generate hyperplastic N discs), or to *UAS-Nicd; FRT82B scrib1* (to generate neoplastic NS discs). Because *scrib1* clones are eliminated in growing discs, the dysplasic S discs were obtained from *FRT82B scrib1 / Df(3R)BSC752* 3rd instar larvae. All crosses were performed at 25°C and carefully staged (time after egg laying and tube crowding).

For functional studies, neoplastic growth was obtained by driving *UAS-Nicd* and the *scrib* RNAi *P{TRiP.HMS01490}attP2* by the *Bx-Gal4* (pouch of larval wing discs). Modifications of the overgrowth phenotype and of the expression of the Mmp1 invasive marker were performed by crossing in F1 *Bx-Gal4, UAS GFP;; UAS Nicd, UAS scribHMS01490* to the desired *UAS RNAi* or control lines (*UAS white RNAi* or *UAS GFP*), to ensure similar UAS load. List of lines tested in Supplemental Materials.

Information on gene models and functions, and on *Drosophila* lines available were obtained from FlyBase (flybase.org – (Thurmond et al., 2019).

RNA extraction and RNA-Seq

RNA from 60 or 80 dissected third instar larva wing discs of WT, N, NS and S discs was extracted using TriZOL. Genomic DNA was eliminated using Ambion's DNA-free kit (#AM1906). cDNA bank preparation were then performed from 1µg of RNA and sequencing on a Illumina HisSeq 2000 by the Biocampus genomic facility MGX of Montpellier. After sequencing, reads obtained were filtered based on their quality (circa 40 millions reads were kept per conditions). The reads were then align on *Drosophila* dm6 genome by the ABIC facility in Montpellier producing a matrix of reads per gene and per condition. This matrix was then normalized and pair-wise differential expression was performed using DESeq (Anders and Huber, 2010). Other differential expression tools were tested such as DESeq2 and edgeR with default parameters but appeared either less stringent, or inadequate.

qPCR

qPCR was performed on biological triplicates on a Roche LightCycler 480, and fold

change was estimated by the δδCT approach. List of primers used in Supplemental

Materials.

Su(H) Chromatin Immuno Precipitation

After dissection in PBS 1X, Protein/DNA complexes from 60 wing discs (80 for S

condition) were cross-linked with 1% formaldehyde for 10 minutes. The reaction was

then quenched by 0.125 M Glycine and washed 3x in PBS. Wing disc cells were

resuspended in 50μL Nuclear Lysis Buffer (Tris-HCl pH 8.1 20mM, EDTA 10mM,

SDS 1%). Lysates were sonicated on a Bioruptor (Diagenode), and diluted 10x in

Immunoprecipitation Dilution Buffer (Tris-HCl pH 8.1 20mM, EDTA 2mM, SDS

0.01%, NaCl 150mM, Triton X-100 1%) and precleared with rabbit IgG (Sigma) and

protein G Agarose (Santa Cruz Biotechnology). ChIP reactions were performed by

incubating lysates overnight at 4°C with 1ng of Goat anti-Su(H) (Santa Cruz

Biotechnology, sc15813), and immunocomplexes were then isolated with Protein G

Agarose for 2h, washed 2x with Wash Buffer 1 (Tris-HCl pH 8.1 20mM, EDTA

2mM, SDS 0.1%, NaCl 50mM, Triton X-100 1%) and 2x with Wash Buffer 2 (Tris-

HCl pH 8.1 10mM, EDTA 1mM, LiCl 250mM, NP-40 1%, Deoxycholic acid 0.4%),

before a decross-linking step at 65°C in 0.25M NaCl. Samples were then treated with

0.2 mg/mL proteinase K and 50mg/mL RNase A. The DNA was then purified on

columns (Qiagen, 28106). ChIP efficiency was checked by qPCR normalized on input

chromatin with the following primer couples, corresponding to known strong binding

sites of Su(H). List of primers used in Supplemental Materials.

For whole-genome analysis, 1 μg double-stranded ChIP or input DNA (corresponding

to 180 discs for each replicate) was labelled with either Cy3- or Cy5-random primers

using the Nimblegen Dual Colour kit. Both ChIP and input were co-hybridised to

NimbleGen D. melanogaster ChIP-chip 2.1 M whole-genome tiling arrays in the

NimbleGen hybridisation station at 42°C for 16 h and then washed according to the

NimbleGen Wash Buffer kit instructions. The data obtained were normalized using

quantile normalization across the replicate arrays in R. Window smoothing and peak

calling were performed using the Bioconductor package Ringo (Toedling et al., 2007)

with a winHalfSize of 300 bp and min.probes = 5. Probe levels were then assigned P-

values based on the normalNull method, corrected for multiple testing using the

Hochberg–Benjamini algorithm and then condensed into regions using distCutOff of

200 bp.

In order to determine the Notch Direct Targets (NDTs), ChIP and RNA-Seq results

were compared: NDTs are defined as up-regulated genes with Su(H) enrichment

within 20kb. As such one Su(H) peak could be assigned to several upregulated genes

consistent with its role in enhancer regions. The 20kb window was chosen as it

allowed the recovery of more than 85% of NDTs in our previous study that was based

on closest gene assignment irrespective of distance (Djiane et al., 2013).

GO Term analyses

The lists of significantly regulated genes in the various comparisons were submitted

to gene ontology (GO) term enrichment analysis. We used the GO biological process

(GOBP) ontology and applied hypergeometric tests (p-values) followed by

Benjamini-Hochberg multiple hypothesis correction (q-values).

iRegulon analyses

In order to determine the likely transcriptional modules in our transcriptomic and

NDTs datasets, we used the online tool iRegulon (http://iregulon.aertslab.org/), with

the standard settings using the 6K Motif collection (6383 PWMs) and a Putative

regulatory region of "10kb upstream, full transcript and 10kb downstream".

Importantly, these settings allowed the recovery of the "positive control" Su(H)

module.

Immunocytochemistry

Antibody staining of wing imaginal discs were performed using standard protocols.

Images were acquired on a Zeiss Apotome2 microscope and processed and quantified

using Zen or ImageJ. Primary antibodies used were rat anti-DE-Cadherin (DCAD2,

Developmental Studies Hybridoma Bank – DHSB, 1:25), rabbit anti-GFP (A6455,

Molecular Probes, 1:200), and mouse anti-Mmp1 (3A6B4, DHSB, 1:25). Secondary

antibodies used conjugated to Alexa-350, Alexa-488, or Cy3 were from Jackson Labs

Immuno Research (1:200).

Quantification methods

Genotypes were tested in batches with controls and 13-17 images corresponding to

13-17 different discs were all acquired on the same microscope with the same

exposure settings.

Growth was estimated by the size of the GFP positive area and normalized to that of

controls. A disc was considered overgrown if its size was over the average size of

controls plus 1.5 the stdev of controls. A disc was undergrown, if its size was under

the average of controls minus 1.5 the stdev of controls. Using this "conservative" set

up identified only circa 5% of controls as undergrown.

Mmp1 intensities were ranked as High, Low, Null by independent observer with

genotypes masked and processed in random order.

ACKNOWLEDGEMENTS

We thank G. Alvès, D. Andrew, D. Eberl, P. Léopold, R. Levayer, C. Maurange, A. Teleman, J. Terman, and T.T. Su for sharing flies. We acknowledge the Bloomington Stock Center, the Vienna Stock Center, the DGRC Kyoto Stock Center, the Developmental Studies Hybridoma Bank, the *Drosophila* facility, the MGX sequencing facility (BioCampus Montpellier, CNRS, INSERM, Université de Montpellier), and FlyBase for their support to our research.

The lab of SJB is supported by the MRC (MRC programme grant MR/L007177/1). RL was supported by fellowships from "Ligue Nationale Contre le Cancer - LNCC" and from "Fondation ARC pour la recherche sur le cancer". The lab of AD is supported by the "Fondation ARC pour la recherche sur le cancer", "Marie Curie CIG", and "Agence Nationale de la Recherche - ANR".

AUTHOR CONTRIBUTIONS

RL, CG, PL, DK, and AD performed experiments; RL, CG, LHM, JC, and AD analyzed data; JC provided expertise for RNA-Seq; BF and SJB provided technical support and expertise for ChIP; RL, SJB, and AD wrote the manuscript; AD conceived and supervised the project.

DECLARATION OF INTEREST

The authors declare no competing interests.

REFERENCES

Akdemir F, Christich A, Sogame N, Chapo J, Abrams JM. 2007. p53 directs focused genomic responses in Drosophila. *Oncogene* **26**:5184–5193. doi:10.1038/sj.onc.1210328

Anders S, Huber W. 2010. Differential expression analysis for sequence count data. *Genome Biol* **11**:R106. doi:10.1186/gb-2010-11-10-r106

Atkins M, Potier D, Romanelli L, Jacobs J, Mach J, Hamaratoglu F, Aerts S, Halder G. 2016. An Ectopic Network of Transcription Factors Regulated by Hippo Signaling Drives Growth and Invasion of a Malignant Tumor Model. *Curr Biol* **26**:2101–2113. doi:10.1016/j.cub.2016.06.035

Baillon L, Germani F, Rockel C, Hilchenbach J, Basler K. 2018. Xrp1 is a transcription factor required for cell competition-driven elimination of loser cells. *Sci Rep* **8**:17712. doi:10.1038/s41598-018-36277-4

Banks L, Pim D, Thomas M. 2012. Human tumour viruses and the deregulation of cell polarity in cancer. *Nat Rev Cancer* **12**:877–886. doi:10.1038/nrc3400

Bilder D, Schober M, Perrimon N. 2003. Integrated activity of PDZ protein complexes regulates epithelial polarity. *Nat Cell Biol* **5**:53–58. doi:10.1038/ncb897

Blanco J, Cooper JC, Baker NE. 2020. Roles of C/EBP class bZip proteins in the growth and cell competition of Rp ('Minute') mutants in Drosophila. *Elife* **9**. doi:10.7554/eLife.50535

Boulan L, Andersen D, Colombani J, Boone E, Léopold P. 2019. Inter-Organ Growth Coordination Is Mediated by the Xrp1-Dilp8 Axis in Drosophila. *Dev Cell* **49**:811-818.e4. doi:10.1016/j.devcel.2019.03.016

Bray SJ. 2016. Notch signalling in context. *Nat Rev Mol Cell Biol* **17**:722–735. doi:10.1038/nrm.2016.94

Brumby AM, Richardson HE. 2003. scribble mutants cooperate with oncogenic Ras or Notch to cause neoplastic overgrowth in Drosophila. *EMBO J* **22**:5769–5779. doi:10.1093/emboj/cdg548

Bunker BD, Nellimoottil TT, Boileau RM, Classen AK, Bilder D. 2015. The transcriptional response to tumorigenic polarity loss in Drosophila. *Elife* **4**. doi:10.7554/eLife.03189

Coopman P, Djiane A. 2016. Adherens Junction and E-Cadherin complex regulation by epithelial polarity. *Cell Mol Life Sci* **73**:3535–3553. doi:10.1007/s00018-016-2260-8

Cordenonsi M, Zanconato F, Azzolin L, Forcato M, Rosato A, Frasson C, Inui M, Montagner M, Parenti AR, Poletti A, Daidone MG, Dupont S, Basso G, Bicciato S, Piccolo S. 2011. The Hippo transducer TAZ confers cancer stem cell-related traits on breast cancer cells. *Cell* **147**:759–772. doi:10.1016/j.cell.2011.09.048

Cordero JB, Macagno JP, Stefanatos RK, Strathdee KE, Cagan RL, Vidal M. 2010. Oncogenic Ras diverts a host TNF tumor suppressor activity into tumor promoter. *Dev Cell* **18**:999–1011. doi:10.1016/j.devcel.2010.05.014

Djiane A, Krejci A, Bernard F, Fexova S, Millen K, Bray SJ. 2013. Dissecting the mechanisms of Notch induced hyperplasia. *EMBO J* **32**:60–71. doi:10.1038/emboj.2012.326

Donohoe CD, Csordás G, Correia A, Jindra M, Klein C, Habermann B, Uhlirova M. 2018. Atf3 links loss of epithelial polarity to defects in cell differentiation and cytoarchitecture. *PLoS Genet* **14**:e1007241. doi:10.1371/journal.pgen.1007241

Francis MJ, Roche S, Cho MJ, Beall E, Min B, Panganiban RP, Rio DC. 2016. Drosophila IRBP bZIP heterodimer binds P-element DNA and affects hybrid dysgenesis. *Proc Natl Acad Sci USA* **113**:13003–13008. doi:10.1073/pnas.1613508113

Fre S, Huyghe M, Mourikis P, Robine S, Louvard D, Artavanis-Tsakonas S. 2005. Notch signals control the fate of immature progenitor cells in the intestine. *Nature* **435**:964–968. doi:10.1038/nature03589

Han W, Ming M, Zhao R, Pi J, Wu C, He Y-Y. 2012. Nrf1 CNC-bZIP protein promotes cell
survival and nucleotide excision repair through maintaining glutathione
homeostasis. *J Biol Chem* **287**:18788–18795. doi:10.1074/jbc.M112.363614
Harding HP, Zhang Y, Zeng H, Novoa I, Lu PD, Calfon M, Sadri N, Yun C, Popko B, Paules R,
Stojdl DF, Bell JC, Hettmann T, Leiden JM, Ron D. 2003. An integrated stress
response regulates amino acid metabolism and resistance to oxidative stress.
Mol Cell **11**:619–633. doi:10.1016/s1097-2765(03)00105-9
Herzig RP, Andersson U, Scarpulla RC. 2000. Dynein light chain interacts with NRF-1 and
EWG, structurally and functionally related transcription factors from humans
and drosophila. *J Cell Sci* **113 Pt 23**:4263–4273.
Ho DM, Pallavi SK, Artavanis-Tsakonas S. 2015. The Notch-mediated hyperplasia
circuitry in Drosophila reveals a Src-JNK signaling axis. *Elife* **4**:e05996.
doi:10.7554/eLife.05996
Huang L, Muthuswamy SK. 2010. Polarity protein alterations in carcinoma: a focus on
emerging roles for polarity regulators. *Curr Opin Genet Dev* **20**:41–50.
doi:10.1016/j.gde.2009.12.001
Huggins CJ, Mayekar MK, Martin N, Saylor KL, Gonit M, Jailwala P, Kasoji M, Haines DC,
Quiñones OA, Johnson PF. 2015. C/EBPγ Is a Critical Regulator of Cellular Stress
Response Networks through Heterodimerization with ATF4. *Mol Cell Biol*
36:693–713. doi:10.1128/MCB.00911-15
Igaki T, Pagliarini RA, Xu T. 2006. Loss of cell polarity drives tumor growth and invasion
through JNK activation in Drosophila. *Curr Biol* **16**:1139–1146.
doi:10.1016/j.cub.2006.04.042
Igaki T, Pastor-Pareja JC, Aonuma H, Miura M, Xu T. 2009. Intrinsic tumor suppression
and epithelial maintenance by endocytic activation of Eiger/TNF signaling in
Drosophila. *Dev Cell* **16**:458–465. doi:10.1016/j.devcel.2009.01.002
Ji Z, Kiparaki M, Folgado V, Kumar A, Blanco J, Rimesso G, Chuen J, Liu Y, Zheng D, Baker
NE. 2019. Drosophila RpS12 controls translation, growth, and cell competition
through Xrp1. *PLoS Genet* **15**:e1008513. doi:10.1371/journal.pgen.1008513
Katheder NS, Khezri R, O'Farrell F, Schultz SW, Jain A, Rahman MM, Schink KO,
Theodossiou TA, Johansen T, Juhász G, Bilder D, Brech A, Stenmark H, Rusten TE.
2017. Microenvironmental autophagy promotes tumour growth. *Nature*
541:417–420. doi:10.1038/nature20815
Külshammer E, Mundorf J, Kilinc M, Frommolt P, Wagle P, Uhlirova M. 2015. Interplay
among Drosophila transcription factors Ets21c, Fos and Ftz-F1 drives JNK-
mediated tumor malignancy. *Dis Model Mech* **8**:1279–1293.
doi:10.1242/dmm.020719
Kusch T, Florens L, Macdonald WH, Swanson SK, Glaser RL, Yates JR, Abmayr SM,
Washburn MP, Workman JL. 2004. Acetylation by Tip60 is required for selective
histone variant exchange at DNA lesions. *Science* **306**:2084–2087.
doi:10.1126/science.1103455
Link N, Kurtz P, O'Neal M, Garcia-Hughes G, Abrams JM. 2013. A p53 enhancer region
regulates target genes through chromatin conformations in cis and in trans.
Genes Dev **27**:2433–2438. doi:10.1101/gad.225565.113
Loubiere V, Papadopoulos GL, Szabo Q, Martinez A-M, Cavalli G. 2020. Widespread
activation of developmental gene expression characterized by PRC1-dependent
chromatin looping. *Sci Adv* **6**:eaax4001. doi:10.1126/sciadv.aax4001
Maraver A, Fernández-Marcos PJ, Herranz D, Muñoz-Martin M, Gomez-Lopez G,
Cañamero M, Mulero F, Megías D, Sanchez-Carbayo M, Shen J, Sanchez-Cespedes
M, Palomero T, Ferrando A, Serrano M. 2012. Therapeutic effect of γ-secretase
inhibition in KrasG12V-driven non-small cell lung carcinoma by derepression of
DUSP1 and inhibition of ERK. *Cancer Cell* **22**:222–234.
doi:10.1016/j.ccr.2012.06.014

McCaffrey LM, Montalbano J, Mihai C, Macara IG. 2012. Loss of the Par3 polarity protein
promotes breast tumorigenesis and metastasis. *Cancer Cell* **22**:601–614.
doi:10.1016/j.ccr.2012.10.003
Muzzopappa M, Murcia L, Milán M. 2017. Feedback amplification loop drives malignant
growth in epithelial tissues. *Proc Natl Acad Sci USA* **114**:E7291–E7300.
doi:10.1073/pnas.1701791114
Ntziachristos P, Lim JS, Sage J, Aifantis I. 2014. From fly wings to targeted cancer
therapies: a centennial for notch signaling. *Cancer Cell* **25**:318–334.
doi:10.1016/j.ccr.2014.02.018
Ohsawa S, Sugimura K, Takino K, Xu T, Miyawaki A, Igaki T. 2011. Elimination of
oncogenic neighbors by JNK-mediated engulfment in Drosophila. *Dev Cell*
20:315–328. doi:10.1016/j.devcel.2011.02.007
Pagliarini RA, Xu T. 2003. A genetic screen in Drosophila for metastatic behavior. *Science*
302:1227–1231. doi:10.1126/science.1088474
Pascual J, Jacobs J, Sansores-Garcia L, Natarajan M, Zeitlinger J, Aerts S, Halder G,
Hamaratoglu F. 2017. Hippo Reprograms the Transcriptional Response to Ras
Signaling. *Dev Cell* **42**:667-680.e4. doi:10.1016/j.devcel.2017.08.013
Pinal N, Martín M, Medina I, Morata G. 2018. Short-term activation of the Jun N-terminal
kinase pathway in apoptosis-deficient cells of Drosophila induces tumorigenesis.
Nat Commun **9**:1541. doi:10.1038/s41467-018-04000-6
Ranganathan P, Weaver KL, Capobianco AJ. 2011. Notch signalling in solid tumours: a
little bit of everything but not all the time. *Nat Rev Cancer* **11**:338–351.
doi:10.1038/nrc3035
Reddy KL, Rovani MK, Wohlwill A, Katzen A, Storti RV. 2006. The Drosophila Par domain
protein I gene, Pdp1, is a regulator of larval growth, mitosis and endoreplication.
Dev Biol **289**:100–114. doi:10.1016/j.ydbio.2005.10.042
Reinke AW, Baek J, Ashenberg O, Keating AE. 2013. Networks of bZIP protein-protein
interactions diversified over a billion years of evolution. *Science* **340**:730–734.
doi:10.1126/science.1233465
St Johnston D, Ahringer J. 2010. Cell polarity in eggs and epithelia: parallels and
diversity. *Cell* **141**:757–774. doi:10.1016/j.cell.2010.05.011
Sykiotis GP, Bohmann D. 2008. Keap1/Nrf2 signaling regulates oxidative stress
tolerance and lifespan in Drosophila. *Dev Cell* **14**:76–85.
doi:10.1016/j.devcel.2007.12.002
Thurmond J, Goodman JL, Strelets VB, Attrill H, Gramates LS, Marygold SJ, Matthews BB,
Millburn G, Antonazzo G, Trovisco V, Kaufman TC, Calvi BR, FlyBase Consortium.
2019. FlyBase 2.0: the next generation. *Nucleic Acids Res* **47**:D759–D765.
doi:10.1093/nar/gky1003
Toedling J, Skylar O, Sklyar O, Krueger T, Fischer JJ, Sperling S, Huber W. 2007. Ringo--an
R/Bioconductor package for analyzing ChIP-chip readouts. *BMC Bioinformatics*
8:221. doi:10.1186/1471-2105-8-221
Toggweiler J, Willecke M, Basler K. 2016. The transcription factor Ets21C drives tumor
growth by cooperating with AP-1. *Sci Rep* **6**:34725. doi:10.1038/srep34725
Uhlirova M, Bohmann D. 2006. JNK- and Fos-regulated Mmp1 expression cooperates
with Ras to induce invasive tumors in Drosophila. *EMBO J* **25**:5294–5304.
doi:10.1038/sj.emboj.7601401
Uhlirova M, Jasper H, Bohmann D. 2005. Non-cell-autonomous induction of tissue
overgrowth by JNK/Ras cooperation in a Drosophila tumor model. *Proc Natl*
Acad Sci USA **102**:13123–13128. doi:10.1073/pnas.0504170102
Wu M, Pastor-Pareja JC, Xu T. 2010. Interaction between Ras(V12) and scribbled clones
induces tumour growth and invasion. *Nature* **463**:545–548.
doi:10.1038/nature08702

Xue B, Krishnamurthy K, Allred DC, Muthuswamy SK. 2013. Loss of Par3 promotes
breast cancer metastasis by compromising cell-cell cohesion. *Nat Cell Biol*
15:189–200. doi:10.1038/ncb2663
Ye J, Rawson RB, Komuro R, Chen X, Davé UP, Prywes R, Brown MS, Goldstein JL. 2000.
ER stress induces cleavage of membrane-bound ATF6 by the same proteases
that process SREBPs. *Mol Cell* **6**:1355–1364. doi:10.1016/s1097-
2765(00)00133-7
860
861

 FIGURE LEGENDS

Figure 1. Notch-based neoplastic growth paradigms in *Drosophila* wing discs

A-D. 3rd instar wing imaginal discs at precisely 5 days after egg-laying either wild-

type (WT; A), overexpressing activated Notch (N; B), mutant for *scrib* (S; C), or

combining overexpressed Notch and *scrib* mutation (NS; D) and marked for E-Cad

(blue) and Mmp1 (red). **A,B&D.** MARCM clones (positively marked by GFP; green)

of the indicated genotypes: expressing only GFP (A; WT), expressing Nicd & GFP

(B; N), and expressing Nicd & GFP and mutant for *scrib* (C; NS). **C.** Discs fully

mutant for *scrib*.

E-F. Differentially expressed genes as compared to WT in the different growth

paradigms N (green), S (red), and NS (blue) identified by RNA-Seq. **E.** Heatmap of

gene expressions after unsupervised clustering. **F.** Venn diagram of up-regulated and

down-regulated genes in N, S, and NS.

G. Semi-quantitative RT-PCR of the indicated genes represented as fold change

compared to WT (grey) in the different N (green), S (red), and NS (blue) growth

paradigms and normalized to *Atc5C* expression. Biological triplicates, standard error

to the mean (s.e.m.) is shown.

Figure 2. GO term analysis of the N, S, and NS transcriptomes

A. Enrichment diagram as measured by adjusted p-value for selected GO terms (full

list as supplemental material) and represented as bars for N (green), S (red), and NS

(blue). GO terms color reflect whether they are shared or specific: shared by all

(black), common N&S (orange), common N&NS (dark green), common S&NS

(purple).

B. Venn diagram showing the domains of overlap of GO terms identified

(significantly enriched) in N, S, and NS.

Figure 3. Polarity loss redirects the transcriptional output of Notch during

neoplastic growth

A. Experimental set-up to identify the Notch Direct Targets genes (NDTs): Genes up-

regulated in N, S, or NS (Transcriptomic), and located within 20kb of a Su(H) binding

site (ChIP).

B. Venn diagram showing the NDTs overlap in N, S, and NS, showing core Notch

responses, but also significant condition specific NDTs.

C. Overlap of the Su(H) binding sites identified by ChIP in N, S, and NS, showing

that almost all S and NS Su(H) peaks are also found in N. The overlap is shown in

white. Numbers are slightly different because in the Su(H) peaks calling protocol, in

some rare cases, some peaks can be split between conditions where one peak in one

condition would overlap with two peaks in the other.

D. Genome Viewer snapshots of several NDTs (shown in purple) such as the NS

specific *Act87E* and *Wnt10* (but also *yellow-e* and *Ir87a*), the common *Ets21C*, and

the N/NS NDT *upd3*. For each condition, the Su(H) ChIP enrichment is shown in the

upper lane, and the Su(H) peaks identified are represented by the blocks underneath.

Figure 4. Identification of potential transcriptional modules mediating N, S, and

NS growth

A. Venn diagram of significant transcription factors (TFs) identified by iRegulon as

potential key mediators for the expression of the N (in the green circle), S (red circle),

and NS (blue circle) up-regulated genes. TFs were color-coded according to their

molecular class and/or belonging to the similar regulon. Numbers represent the

number of TFs identified. See also Figure S2 for the iRegulon analyses of the NDTs,

and the detailed lists of both iRegulon analyses in Supplemental tables S8 to S15.

B. Heat map for the expression of some of the transcription factors identified by

iRegulon and grouped as "oncogenic module", "basic Leucine Zippers", "DNA

damage and genome integrity".

Figure 5. Neoplastic growth is mediated by the "oncogenic module" and by a

diverse network of bZIP transcription factors including Xrp1

A-H. 3rd instar wing imaginal discs expressing *GFP*, an activated form of Notch

(*Nicd*) and an RNAi for *scrib* under the control of the *Bx-Gal4* driver (dorsal wing

pouch) and stained for GFP (green, white A'-H') to assess tissue overgrowth and for

Mmp1 (red, white A''-H'') to assess tissue invasiveness. Discs also expressed under

Bx-Gal4 control the indicated constructs (*UAS-RNAi* or *UAS Bsk DN*).

I-K. Quantification of the overgrowth of the GFP territory in the indicated genotypes.

Results are shown as percentage of discs. In blue are shown the discs smaller than the

average of controls (size < Average size of controls – 1.5x StDev of controls). In

green are shown discs that are the same size as the controls average or bigger. Results

for the "DNA damage module" in **I**, for the "oncogenic module" in **J**, and for the

"AP-1 and bZIP module" in **K**. Green arrowheads indicate modification. Star indicate

a genotype with misshapen and "sick" discs.

I'-K'. Quantification of the Mmp1 intensity in the indicated genotypes. Results are

shown as percentage where discs were classified to fall in three categories: High

Mmp1 staining intensity (red, similar to that shown in A''), Low Mmp1 staining

intensity (blue, similar to that shown in C or H), No Mmp1 staining (light blue,

similar to that shown in B''). Results for the "DNA damage module" in **I'**, for the

"oncogenic module" in **J'**, and for the "AP-1 and bZIP module" in **K'**. Red

arrowheads indicate modification. Star indicate a genotype with misshapen and "sick"

discs.

SUPPLEMENTAL INFORMATION

SUPPLEMENTAL MATERIAL AND METHODS

Drosophila genetics

Overexpression lines tested in the *Bx-Gal4, UAS GFP;; UAS Nicd, UAS*

scribHMS01490 screen were UAS GFP, UAS bskK53R [20.1a] BL#9311, and UAS

SOD CAT (gift from P. Leopold). RNAi lines used are listed below:

Atf-2   HMC05118   BL#60124

Atf-2   JF02108   BL#26210

Atf3   KK113077   v#105036

Atf3   JF02303   BL#26741

Atf6   JF02109   BL#26211

ATM/tefu   KK100008   v#108074

ATM/tefu   HMS02790   BL#44073

ATR/mei-P26 HMC04662   BL#57268

CG6272   HMS00057   BL#33652

CG6272   KK110056   v#101871

CG7786   HMC05169   BL#62162

cnc   HMS00650   BL#32863

crc   JF02007   BL#25985

crc   KK111018   v#109014

CrebB HMJ30249   BL#63681

gt   HMS01105   BL#34631

Jra   JF01184   BL#31595

kay   HMS00254   BL#33379

mafS   HMS02020   BL#40853

p53   HMS02286   BL#41720

pdp1   HMS02030   BL#40863

vri   HMS02029   BL#40862

w   HMS00045   BL#33644

Xrp1   HMS00053   BL#34521

Xrp1   HMJ21189   BL#51054

Xrp1   HMJ22533   BL#60356

Primers

qPCR primers:

976	Act5C_F:	GAGCGCGGTTACTCTTTCAC
977	Act5C_R:	ACTTCTCCAACGAGGAGCTG
978	Act87E_F:	GTCCACCGCAAGTGCTTCTA
979	Act87E_R:	TTTCTTTGGATGGCAGGGCA
980	Atf3_F:	CAGCATGGCAACATTGGGAC
981	Atf3_R:	ATGAAGGCAGTGGCTGAGTC
982	Diap1_F:	CAGCCACACGCATCTTCAAC
983	Diap1_R:	ACTTTGTCACAGAGGAGGCG
984	E2f1_F:	ACAGAATCCTCGCCTCCAAC
985	E2f1_R:	GACTGCTGCCGTAGCCTATT
986	Ets21C_F:	CTGCTCGCTGATTCGTCCAA
987	Ets21C_R:	TAGGCATACCGCTTTCCGTG
988	ftz-f1_F:	ATTCCTGGTCGGACATGCTT
989	ftz-f1_R:	TTCATGCAGACATAGTCGCCC
990	mxc_F:	ACTAGAGGAGGAGCAGCGAA
991	mxc_R:	CTAGTGGACAGCGGCGTATT
992	p38a_F:	TACGGACAGGTGTCAAAGGC
993	p38a_R:	CAGCGATCCATTAGCGGGAT
994	p53_F:	TGCGTGTGTTCCTTTGCTTC
995	p53_R:	GTTCAGGGGGACTACAACGG
996	puc_F:	ATTGACCTCGCCGCCAATTA
997	puc_R:	ATTCCGCTTGAACAGAGCCA
998	sd_F:	AGGGTCCACAGAATGCGTTT
999	sd_R:	TCGCTTTCCACCTTCTCCAC
1000	Sdr_F:	CGCTCCCTCAATCCCAAAGT
1001	Sdr_R:	ACAACGTCCATCAGCCAGTT
1002	Ser_F:	GCACGAATCTCTGGTGTGGA
1003	Ser_R:	TAGATTTGGCTGGCAGTCGG
1004	wg_F:	GCAGTCTGGTCTGGTCTACG
1005	wg_R:	ATTGTGCGGGTTCAGTTGGA
1006	Wnt10_F:	AATGGCATCGGTGGAACTGT
1007	Wnt10_R:	CAGCGTCTTGCGATTGATGG

1008

qChIP primers:

E(spl)mβ_F: AAGTCGGAGCTTTGAATGAG

E(spl)mβ_R: CAAGTCATTTTATTGCCCTCAC

E(spl)m5_F: GTTTCCGCAGGTCCAGTTAC

E(spl)m5_R: GTTTGATGTTCACGCTGCTG

white_F: CGAAGGACGTTGACACATTG

white_R: GAATTGCCGCTTTTCTCAC

DDC_F: AAGTGGGATTTGCCAGTGAC

DDC_R: TGCTGGTGAACTTTGACTGC

CG42808_F: CTCGTTAAGAGCAACTGCGA

CG42808_R: GTGAGAACTCCGAATCGAGG

CG6191_F: CGAAAAATGCGGACGATTCC

CG6191_R: CCCACCAATCTAGGGTTTCA

Ilp8_F: TCATCTCCGGTGTCTGACTT

Ilp8_R: AAAGAATTGGCTGCGGAAGA

1024

1025

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. Features of the Notch Direct Targets (NDTs) in N, S, and NS (relates

to Fig. 3)

A. Genome-wide localization of the Su(H) ChIP enrichment peaks with respect to

genome features, showing that the major changes were found in NS where Su(H)

"peaks" were more prevalent in introns but less prevalent at promoter regions than in

N and S.

B. Heatmaps for the expression of the different NDTs in WT, N, S, and NS. From left

to right are presented the N, S, NS, and finally All NDTs, highlighting that NDTs

could be transcriptionally up-regulated in more than in one condition.

C. Genome browser view of the whole left arm of the 2nd chromosome, and showing

the Su(H) ChIP enrichment (upper rows)) and the intervals called as Su(H) peaks

(lower rows) in N (green), NS (blue), and S (red). Note the higher number of peaks in

N, and the rarity of NS, or S peaks not found in N.

1040

Figure S2. Identification of potential transcriptional modules mediating N, S,

and NS growth (relates to Fig. 4)

A. Venn diagram of significant transcription factors identified by iRegulon in N, S,

and NS Notch Direct Targets, and color-coded according to their molecular class.

Numbers represent the number of transcription factors identified in each group. See

also the detailed lists of all iRegulon analyses in Supplemental tables S8 to S15.

1047

1048

SUPPLEMENTAL TABLES

Table S1-3. Differentially expressed genes in N, S, and NS identified by DESeq

(related to Fig. 1). Columns are:

FBgn_ID: Unique FlyBase gene ID

Symbol: Current FlyBase gene symbol

qval: adjusted p-value for multiple testing

logFC: log2 of the Fold Change "Condition N, S, or NS" / "Control WT"

1056

Table S4. Gene Ontology (GO) enrichment in N, S, and NS (related to Fig. 2).

Columns are:

experiment: type of comparison

GO_term: GO term number

description: GO term description

intersect.size: number of genes in query within the GO term

n.pw: total number of genes within the given GO term

pval: p value after hypergeometric test

qval: adjusted pvalue for multiple testing

genes: genes in query falling in the specific GO term

1067

Table S5. Curated Gene Ontology (GO) enrichment in N, S, and NS (related to Fig.

2). Columns are:

experiment: type of comparison

GO_term: GO term number

description: GO term description

intersect.size: number of genes in query within the GO term

n.pw: total number of genes within the given GO term

pval: p value after hypergeometric test

qval: adjusted pvalue for multiple testing

genes: genes in query falling in the specific GO term

1078

Table S6. Su(H) ChIP enrichment peaks coordinates in N, S, and NS (related to Fig.

3). Columns are:

Exp: N, S, or NS

Chr: Chromosome arm

MIN: smallest peak coordinate

MAX: biggest peak coordinate

1085

1086

Table S7. All Notch Direct Targets (NDTs) ordered by genomic position. This table

includes an indication whether the genes are transcriptionally upregulated or have a

Su(H) peak in the vicinity in each N, S, and NS condition. Columns are:

N/NS/S: NDT in the corresponding condition

Type: NDT in different conditions.

FBgn_ID: Unique FlyBase gene ID

SYMBOL: Current FlyBase gene symbol

K_ARM: Chromosome arm location of the gene

MIN (gene pos): smallest gene coordinate

MAX (gene pos): biggest gene coordinate

STRAND: +1 or -1

N Fold: Log2 Fold Change in gene expression N/WT (n.s. not significant)

N ChIP: Su(H) ChIP enrichment peak within 20kb in N (green yes, red no)

NS Fold: Log2 Fold Change in gene expression NS/WT (n.s. not significant)

NS ChIP: Su(H) ChIP enrichment peak within 20kb in NS (green yes, red no)

S Fold: Log2 Fold Change in gene expression S/WT (n.s. not significant)

S ChIP: Su(H) ChIP enrichment peak within 20kb in S (green yes, red no)

1104

1105   **Table S8-10.** iRegulon analyses of the significantly upregulated genes in N, S, and

1106   NS (related to Fig. 4). Analyses were performed using the 6K-PWM and 10kb

1107   upstream and downstream set-ups.

1108

1109   **Table S11.** Curated iRegulon analyses corresponding to Tables S8-10 (related to Fig.

1110   4)

1111

1112   **Table S12-14.** iRegulon analyses of the Notch Direct Targets in N, S, and NS (related

1113   to Fig. S2). Analyses were performed using the 6K-PWM and 10kb upstream and

1114   downstream set-ups.

1115

1116   **Table S15.** Curated iRegulon analyses corresponding to Tables S12-14 (related to

1117   Fig. S2)

1118

1119   **Table S16.** Xrp1 target genes. Xrp1 regulated genes identified in Ji et al. (2019)

1120   (positively and negatively) and in Baillon et al. (2018). In bold are highlighted genes

1121   found in both studies. Columns are:

1122   FBgn_ID: Unique FlyBase gene ID

1123   Symbol: Current FlyBase gene symbol

Logeay et al. Figure 1

Logeay et al. Figure 2

A Localisation of the Su(H) binding on the genome in each condition (in %)

Legend:
- Promoter
- 5' UTR
- 3' UTR
- Exon
- Intron
- Downstream
- Intergenic

B Heat-maps of the expression of N, S, or NS NDTs (left to right) in each condition

N NDTs S NDTs NS NDTs All NDTs

C Genome browser view of chromosome 2L with Su(H) peak calling in N, S, and NS

N ChIP
peak calling
NS ChIP
peak calling
S ChIP
peak calling
Gene (+)
Coordinates
Gene (−)

A