The Pathogenicity of SARS-CoV-2 in hACE2 Transgenic Mice

Linlin Bao[†,1], Wei Deng[†,1], Baoying Huang[†,2], Hong Gao[†,1], Jiangning Liu[†,1], Lili Ren[3],

Qiang Wei[1], Pin Yu[1], Yanfeng Xu[1], Feifei Qi[1], Yajin Qu[1], Fengdi Li[1], Qi Lv[1], Wenling

Wang[2], Jing Xue[1], Shuran Gong[1], Mingya Liu[1], Guanpeng Wang[1], Shunyi Wang[1], Zhiqi

Song[1], Linna Zhao[1], Peipei Liu[2], Li Zhao[2], Fei Ye[2], Huijuan Wang[2], Weimin Zhou[2], Na

Zhu[2], Wei Zhen[2], Haisheng Yu[1], Xiaojuan Zhang[1], Li Guo[3], Lan Chen[3], Conghui Wang[3],

Ying Wang[3], Xinming Wang[3], Yan Xiao[3], Qiangming Sun[4], Hongqi Liu[4], Fanli Zhu[4],

Chunxia Ma[4], Lingmei Yan[4], Mengli Yang[4], Jun Han[2], Wenbo Xu[2], Wenjie Tan[2],

Xiaozhong Peng[4], Qi Jin[3], Guizhen Wu[*,2], Chuan Qin[*,1]

[1] Key Laboratory of Human Disease Comparative Medicine, Chinese Ministry of Health,

Beijing Key Laboratory for Animal Models of Emerging and Remerging Infectious

Diseases, Institute of Laboratory Animal Science, Chinese Academy of Medical Sciences

and Comparative Medicine Center, Peking Union Medical College, Beijing, China.

[2] MHC Key Laboratory of Biosafety, National Institute for Viral Disease Control and

Prevention, China CDC, Beijing, China.

[3] Institute of Pathogen Biology, Chinese Academy of Medical Sciences, Beijing, China.

[4] Institute of Medical Biology, Chinese Academy of Medical Sciences, Beijing, China.

[†]These authors contributed equally to this work.

[*]Correspondence should be addressed to Chuan Qin, Email: qinchuan@pumc.edu.cn, or

Guizhen Wu, Email: wugz@ivdc.chinacdc.cn.

Abstract

Severe acute respiratory syndrome CoV-2 (SARS-CoV-2) caused the Corona Virus Disease 2019 (COVID-19) cases in China has become a public health emergency of international concern (PHEIC). Based on angiotensin converting enzyme 2 (ACE2) as cell entry receptor of SARS-CoV, we used the hACE2 transgenic mice infected with SARS-CoV-2 to study the pathogenicity of the virus. Weight loss and virus replication in lung were observed in hACE2 mice infected with SARS-CoV-2. The typical histopathology was interstitial pneumonia with infiltration of significant lymphocytes and monocytes in alveolar interstitium, and accumulation of macrophages in alveolar cavities. Viral antigens were observed in the bronchial epithelial cells, alveolar macrophages and alveolar epithelia. The phenomenon was not found in wild type mice with SARS-CoV-2 infection. The pathogenicity of SARS-CoV-2 in hACE2 mice was clarified and the Koch's postulates were fulfilled as well, and the mouse model may facilitate the development of therapeutics and vaccines against SARS-CoV-2.

Key words: SARS-CoV-2, pathogenicity, pneumonia, mouse model, hACE2

In late December of 2019, the coronavirus disease 2019 (COVID-19) caused by severe

acute respiratory syndrome CoV-2 (SARS-CoV-2), linked to a seafood market in which

exotic animals were also sold and consumed, were identified and reported from Wuhan

City, Hubei Province, China[1,2]. The number of confirmed cases has since soared, with

almost 78,000 cases reported and over 2,700 deaths as of February 25, 2020 in China[3],

and imported cases from travelers of mainland China in several other countries. It is

critical to find the pathogenicity and biology of the virus for prevention and treatment of

the disease.

Because SARS-CoV-2 was highly homologous with severe acute respiratory

syndrome coronavirus (SARS-CoV), human angiotensin-converting enzyme 2 (hACE2),

which was the entry receptor of SARS-CoV, was also considered to have a high binding

ability with the SARS-CoV-2 by molecular biological analysis[4,5]. Therefore, we used the

hACE2 transgenic and wild type (WT) mice infected with SARS-CoV-2 to study the

pathogenicity of the virus.

Specific pathogen-free, 6-11-month-old, male and female WT mice (WT-HB-01,

n=15) and hACE2 mice (ACE2-HB-01, n=19) were inoculated intranasally with SARS-

CoV-2 stock virus (HB-01) at a dosage of 10^5 TCID$_{50}$/50 μl inoculum volume per mouse

after the mice were anesthetized by 2.5% avertin, and the mock-treated hACE2 mice

(ACE2-Mock, n=15) were used as control. Clinical manifestations were recorded from

thirteen mice (WT-HB-01, n=3; ACE2-Mock, n=3; ACE2-HB-01, n=7). Compared to

WT-HB-01 mice and ACE2-Mock mice, slight bristles and weight loss were only

observed in ACE2-HB-01 mice during the 14 days observation, and other clinical

symptoms such as arched back and decreased projection of external stimuli were not

found. Notably, the weight loss of ACE2-HB-01 mice was up to 8% at 5 days post

infection (dpi) (Figure 1a).

Next, viral replication and pathological changes were examined from three animals

per group at each time point, and the primary organs were collected periodically,

including heart, liver, spleen, lung, kidney, brain, intestine and testis. As shown in Figure

1b, viral loads were detected by qRT-PCR at 1 dpi, 3 dpi, 5 dpi and 7 dpi in the lungs of

ACE2-HB-01 mice but not in that of WT-HB-01mice (data not shown), and the viral

RNA copies reached a peak at 3 dpi ($10^{6.77}$ copies/ml). Interestingly, the viral RNA was

also detected at 1 dpi in the intestine of ACE2-HB-01 mice, which was failed to be

detected in other tissues along the timeline (Figure 1b). Consistent to the results of viral

loads, infectious virus was respectively isolated from the lungs of ACE2-HB-01 mice at 1

dpi, 3 dpi and 5 dpi, and the highest virus titers were detected at 3 dpi ($10^{2.44}$ TCID$_{50}$/100

μl) (Figure 1c). Meanwhile, the infectious virus was isolated by Vero E6 cells culture

from lung, and the SARS-CoV-2 particles were observed by electron microscopy (Figure

1d). However, the virus was failed to be isolated from the lungs in WT-HB-01 mice and

ACE2-Mock mice along the detecting timeline (Figure 1c), which suggested that hACE2

was essential for SARS-CoV-2 infection and replication in mice. Moreover, specific IgG

antibodies against S protein of SARS-CoV-2 were positively measured in the sera of

ACE2-HB-01 mice at 21 dpi (Figure 1e).

There were no obviously gross and histopathological changes at 1 dpi in all the

animals of each group. Compared to WT-HB-01 mice or ACE2-mock mice with

homogeneously pink and slightly deflated, all the ACE2-HB-01 mice at 3 dpi displayed

comparable gross lesions with focal to multifocal dark red discoloration in certain lung

lobes. The lesions progressed into multifocal to coalescent scattered-dark reddish purple

areas and focal palpable nodules throughout the lung lobes at 5 dpi (Figure 2a). The

damaged lungs became swollen and enlarged in size. Microscopically, consistently, lung

tissues from ACE2-HB-01 mice at 3 dpi showed moderate, multifocal, interstitial

pneumonia. Inflammatory cells including lymphocytes and monocytes accumulated in the

alveolar interstitium and caused thickening of the alveolar walls (Figure 2a). Increased

collagen fiber in the thickened alveolar interstitium in the ACE2-HB-01 mice was

confirmed by Modified Masson's Trichrome stain with blue colors (Supplementary

Figure 1a). The bronchiolar epithelial cells showed swelling, degeneration, some of

which dissolved (Figure 2a). Affected bronchioles were filled with occasionally small

amounts of periodic acid-Schiff (PAS) positive-exudation or denatured and detached

bronchiolar epithelium (Supplementary Figure 1b). The alveolar cavities were dispersed

mainly by swollen and degenerative alveolar macrophages and lymphocytes (Figure 2a).

To investigate the infiltration of specific inflammatory cells, immunohistochemical (IHC)

was carried out to identify MAC2$^+$ macrophages (Supplementary Figure 2a), CD3$^+$ T

lymphocytes, and CD19$^+$ B lymphocytes (Supplementary Figure 2b). Compared to the

WT-HB-01 mice, more macrophages, T lymphocytes, and B lymphocytes were observed

in the lungs of ACE2-HB-01 mice along with lasting prolonged infection time. The

MAC2$^+$ macrophages were diffusely infiltrated in the alveolar cavities (at 3 dpi) or

focally aggregated together in the thickened alveolar septum (at 5 dpi, Supplementary

Figure 2a). CD3$^+$ T lymphocytes and CD19$^+$ B lymphocytes were scattered dispersed or

occasionally aggregated in the alveolar interstitium in the ACE2-HB-01 mice, by contrast,

they were rarely observed in the WT-HB-01 mice (Supplementary Figure 2b).

Perivascular infiltrated inflammatory cells, including lymphocytes and monocytes, were

observed multifocally within and adjacent to affected areas of the lungs. At 5 dpi, the

lung progressed into coalescing interstitial pneumonia with diffuse lesions. Thickened

alveolar septa were filled with lymphocytes and monocytes (Figure 2a). IHC staining of

sequential section revealed that viral antigens were found in the alveolar macrophages,

alveolar epithelia, and the degenerative and being desquamated bronchial epithelial cells

(Figure 2b). There were no significant histopathological lesions (Supplementary Figure 3)

or viral antigens of SARS-CoV-2 (Supplementary Figure 4) in the other organs, including

myocardium, liver, spleen, kidney, cerebrum, intestine, and testis. At 7 dpi, the

pneumonia became mild with focal lesion areas (data not shown).

In addition, the co-localization of SARS-CoV-2 S protein (Fifure 3f) and hACE2

receptor (Figure 3g) was demonstrated in alveolar epithelial cells of ACE2-HB-01 mice

by immunofluorescence (Figures 3h). The phenomenon was not found in the ACE2-

Mock mice (Figures 3a, b, c and d) or WT-HB-01 mice (data not shown), indicating that

the SARS-CoV-2, as same as SARS-CoV, also utilizes the hACE2 as a receptor for

entry[4].

The speed of geographical spread of COVID-19 caused by SARS-CoV-2 has been

declared as public health emergency of international concern (PHEIC), with cases

reported on multiple continents only weeks after the disease was first reported[6]. Although

it has been determined by bioinformatics that the pathogen of this epidemic is a novel

coronavirus, it is necessary to be confirmed by animal experiments following Koch's

postulates.

In the present study, after the experimental infection of hACE2 transgenic mice with

one of the earliest known isolates of SARS-CoV-2, the mice lost weight and showed

interstitial pneumonia, which are comparable with initial clinical reports of pneumonia

caused by SARS-CoV-2[7]. Meanwhile, virus replication was detected in the lung of

ACE2-HB-01 mice and elicited the specific IgG against SARS-CoV-2. In clinical

autopsy, histological examination showed bilateral diffuse alveolar damage with cellular

fibromyxoid exudates[8]. Correspondingly, ACE2-HB-01 mice showed coalescing

interstitial pneumonia in this study. Furthermore, specifically histopathology changes and

progress were observed in the mice, e.g. affected bronchioles were filled with

occasionally small amounts of periodic acid schiff (PAS) positive-exudation or denatured

and detached brochiolar epithelium.

The case fatality rate of currently reported cases is about 2%, which implies that so

far, this novel coronavirus does not seem to cause the high fatality rates as SARS-CoV

(9-11%)[9], which suggested the difference in pathogenicity between the two viruses. The

pathogenicity of SARS-CoV-2 seems mild compared to SARS-CoV in mice, the latter

caused extrapulmonary organ damage, includes brain, kidney, intestine, heart and liver,

furthermore, the neurons are susceptible for SARS-CoV infection, and cerebral casculitis

and hemorrhage were observed in hACE2 transgenic mice[10,11]. However, only interstitial

pneumonia was observed in SARS-CoV-2-infected hACE2 mice, implying the disparity

in pathogenicity of the coronavirus.

Therefore, our results clarified the pathogenicity of SARS-CoV-2 in mice, together

with the previous clinical studies[7], completely fulfills the Koch's postulates[12] and

confirmed SARS-CoV-2 was the pathogen of COVID-19. The mouse model may

facilitate the development of therapeutics and vaccines against SARS-CoV-2.

Materials and methods

Ethics statement

Murine studies were performed in an animal biosafety level 3 (ABSL3) facility using

HEPA-filtered isolators. All procedures in this study involving animals were reviewed

and approved by the Institutional Animal Care and Use Committee of the Institute of

Laboratory Animal Science, Peking Union Medical College (BLL20001).

Viruses and cells

The SARS-CoV-2 (strain HB-01) was kindly provided by Professor Wenjie Tan[1], from

the China Centers for Disease Control and Prevention (China CDC). The complete

genome for this SARS-CoV-2 was submitted to GISAID (BetaCoV/Wuhan/IVDC-HB-

01/2020|EPI_ISL_402119), and deposited in the China National Microbiological Data

Center (accession number NMDC10013001 and genome accession numbers

MDC60013002-01). Seed SARS-CoV-2 stocks and virus isolation studies were

performed in Vero cells, which are maintained in Dulbecco's modified Eagle's medium

(DMEM, Invitrogen, Carlsbad, USA) supplemented with 10% fetal bovine serum (FBS),

100 IU/ml penicillin, and 100 µg/ml streptomycin, and incubated at 37°C, 5% CO_2. For

infected mice, the lung homogenates were used for virus titration tests using endpoint

titration in Vero E6 cells. Virus titer of the supernatant were determined using a standard

50% tissue culture infection dose (TCID50) assay.

Animal experiments

For the animal experiments, specific pathogen-free, 6-11-month-old, male and female transgenic hACE2 mice were obtained from the Institute of Laboratory Animal Science, Peking Union Medical College, China. Transgenic mice were generated by microinjection of the mice ACE2 promoter driving the human ACE2 coding sequence into the pronuclei of fertilized ova from ICR mice, and then human ACE2 integrated was identified by PCR as previous described[10], the hACE2 mainly expressed in lung, heart, kidney, and intestine of transgenic mice. The hACE2 mice or WT (ICR) mice were respectively inoculated intranasally with SARS-CoV-2 stock virus at a dosage of 10^5 $TCID_{50}$ and hACE2 mice intranasally inoculated with equal volume of PBS was used as mock-infection control. The infected animals were continuously observed daily to record body weights, clinical symptoms, responsiveness to external stimuli and death. And the mice were dissected at 1 dpi 3 dpi, 5 dpi and 7 dpi respectively to collect different tissues to screen virus replication and histopathological changes.

Preparation of Homogenate Supernatant

Tissues homogenates were prepared by homogenizing perfused whole tissue using an electric homogenizer for 2 min 30 s in 1 ml of DMEM. The homogenates were centrifuged at 3,000 rpm for 10 min at 4°C. The supernatant was collected and stored at −80°C for viral isolation and viral load detection.

RNA extraction and qRT-PCR

Total RNA was extracted from organs using the RNeasy Mini Kit (Qiagen, Hilden,

Germany), and reverse transcription was performed using the PrimerScript RT Reagent

Kit (TaKaRa, Japan) following manufacturer instructions. A quantitative real-time

reverse transcription-PCR (qRT-PCR) reactions were performed using the PowerUp

SYBG Green Master Mix Kit (Applied Biosystems, USA), in which samples were

processed in duplicate using the following cycling protocol: 50°C for 2 min, 95°C for 2

min, followed by 40 cycles at 95°C for 15 s and 60°C for 30 s, and then 95°C for 15 s,

60°C for 1 min, 95°C for 45 s. The primer sequences used for RT-PCR are targeted

against the envelope (E) gene of SARS-CoV-2 and are as follows: Forward: 5'-

TCGTTTCGGAAGAGACAGGT-3', Reverse: 5'-GCGCAGTAAGGATGGCTAGT-3'.

The PCR products were verified by sequencing using the dideoxy method on an ABI

3730 DNA sequencer (Applied Biosystems, CA, USA). During the sequencing process,

amplification was performed using specific primers. The sequences for this process are

available upon request. The sequencing reads obtained were linked using DNAMAN, and

the results were compared using the Megalign module in the DNAStar software package.

ELISA method

The specific IgG against SARS-CoV from ACE2-HB-01 mice and WT-HB-01 mice were

determined by enzyme-linked immunosorbent assay (ELISA). 96-well plates were coated

with Spike 1 protein of SARS-CoV-2 (0.1μg/100 μL, Sino Biological, 40591-V08H), the

tested sera were diluted at 1:100 and added to each well, and 3 multiple wells were set for

each sample, and then incubated at 37℃ for 30 minutes, followed by the goat anti-mouse

secondary antibodies conjugated with HRP (ZB-2305, zhongshan,1:10,000 dilution), and

incubated at room temperature for 30 minutes. The reaction was developed by TMB

substrate and the optical densities at 450 nm were determined (Metertech960 enzyme

marker with 450 nm wavelength).

Laboratory preparation of the antibody of SARS-CoV-2 Spike-1 (S1) protein

Mice were immunized with purified SARS-CoV-2 S1 protein (Sino biological) and

splenocytes of hyper immunized mice were fused with myeloma cells. Positive clones

were selected by ELISA using SARS-CoV-2 S1 protein (Supplementary Figure 1). The

cell supernatant of 7D2 clone, binding to SARS-CoV-2 S1 protein, was collected for

immunofluorescence analysis.

Pathological Examination

Autopsies were performed in the animal biosafety level 3 (ABSL3) laboratory. Major

organs were grossly examined and then fixed in 10% buffered formalin solution, and

paraffin sections (3-4 μm in thickness) were prepared routinely. Hematoxylin and Eosin

(H&E) stain, periodic acid-Schiff (PAS) stain and modified Masson's Trichrome stain

were used to identify histopathological changes in all the organs. The histopathology of

the lung tissue was observed by light microscopy.

Immunohistochemistry (IHC)

The organs were fixed in 10% buffered formalin solution, and paraffin sections (3-4 μm

in thickness) were prepared routinely. Sections were treated with an antigen retrieval kit

(Boster, AR0022) for 1 min at 37℃ and quenched for endogenous peroxidases in 3%

H_2O_2 in methanol for 10 min. After blocking in 1% normal goat serum, the sections were

incubated with 7D2 monoclonal antibody (laboratory preparation) at 4℃ overnight,

followed by HRP-labeled goat anti-mouse IgG secondary antibody (HRP) (Beijing ZSGB

Biotechnology, ZDR-5307). Alternatively, the sections were stained with MAC2

antibody (Cedarlane Laboratories, CL8942AP), CD3 antibody (Dako, A0452) or CD19

antibody (Cell Signaling Technology, 3574) at 4℃ overnight. Subsequently, the sections

were goat anti-rat IgG secondary antibody (HRP) (Beijing ZSGB Biotechnology,

PV9004), goat anti-rabbit IgG secondary antibody (HRP) (Beijing ZSGB Biotechnology,

PV9001) for 60 min, and visualized by incubation with 3,30-diaminobenzidine

tetrahydrochloride (DAB). The slices were counterstained with hematoxylin, dehydrated

and mounted on a slide and viewed under an Olympus microscope.

Confocal microscopy

For viruses and hACE2 receptor co-localization analysis, the lung tissue sections were

washed twice with PBS, fixed by Immunol Staining Fix Solution (P0098), blocked 1 hour

at room temperature by Immunol Staining Blocking Buffer (P0102) and then incubated

overnight at 4°C with the appropriate primary and secondary antibodies. The nuclei were

stained with DAPI. Anti-S protein antibody (mouse monoclonal 7D2, laboratory

preparation, 1:200) and anti-hACE2 antibody (rabbit polyclonal, ab15348, Abcam1:200)

were used as the primary antibody. The sections were washed with PBS and incubated

with secondary antibodies conjugated with FITC (goat anti-mouse, ZF-0312, Beijing

ZSGB Biotechnology, 1:200) and TRITC (goat anti rabbit, ZF-0317, Beijing ZSGB

Biotechnology, 1:200), dried at room temperature and observed via fluorescence

microscopy. For the expression of hACE2, the sections from WT mice stained with anti-

ACE2 antibody were used as the negative control, and the stable cell line expressing

hACE2 was used as the positive control. For the viral antigen, the sections from ACE2-

Mock mice incubated with anti-S protein antibody were used as the negative control.

Transmission Electron Microscopy

Supernatant from Vero E6 cell cultures that showed cytopathic effects was collected,

inactivated with 2% paraformaldehyde for at least 2 hours, and ultracentrifuged to

sediment virus particles. The enriched supernatant was negatively stained on film-coated

grids for examination. The negative stained grids were observed under transmission

electron microscopy.

Statistical analysis

All data were analyzed with GraphPad Prism 8.0 software. Statistically significant

differences between two groups were determined using unpaired Student's *t*-tests. The

statistical significance among three groups was assessed by one-way ANOVA. A two-

sided p value<0.05 was considered statistically significant.

Acknowledgement

We thank Dr George F Gao for his advice and coordination on this work. We thank

Hongkui Deng, Xiuhong Yang and Lianfeng Zhang for providing the hACE2 mice as a

gift. We also thank Gary Wong for helping us proofread the language. This work was

supported by National Research and Development Project of China (Grant No.

2020YFC0841100), Fundamental Research Funds for CAMS of China (2020HY320001),

National Key Research and Development Project of China (Grant No.

2016YFD0500304), CAMS initiative for Innovative Medicine of China (Grant No. 2016-

I2M-2-006), and National Mega projects of China for Major Infectious Diseases

(2017ZX10304402).

References

1. Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med*, 10.1056/NEJMoa2001017, doi:10.1056/NEJMoa2001017 (2020).

2. Ren, L. L. *et al.* Identification of a novel coronavirus causing severe pneumonia in human: a descriptive study. *Chinese medical journal*, doi:10.1097/cm9.0000000000000722 (2020).

3. Commission, C. N. H. *Update on the novel coronavirus pneumonia outbreak (Jan 24, 2020). Beijing: China National Health Commission, 2020,* < http://www.nhc.gov.cn/yjb/s7860/202002/84faf71e096446fdb1ae44939ba5c528.shtml> (2020).

4. Xu, X. *et al.* Evolution of the novel coronavirus from the ongoing Wuhan outbreak and modeling of its spike protein for risk of human transmission. *Science China. Life sciences*, doi:10.1007/s11427-020-1637-5 (2020).

5. Kuba, K. *et al.* A crucial role of angiotensin converting enzyme 2 (ACE2) in SARS coronavirus–induced lung injury. *Nature Medicine* **11**, 875-879, doi:10.1038/nm1267 (2005).

6. Chan, J. F. *et al.* A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet* **395**, 514-523, doi:10.1016/s0140-6736(20)30154-9 (2020).

7. Huang, C. *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* **395**, 497-506, doi:10.1016/s0140-6736(20)30183-5 (2020).

8. Xu, Z. *et al.* Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *The Lancet. Respiratory medicine*, doi:10.1016/s2213-2600(20)30076-x (2020).

9. de Wit, E., van Doremalen, N., Falzarano, D. & Munster, V. J. SARS and MERS: recent insights into emerging coronaviruses. *Nature reviews. Microbiology* **14**, 523-534, doi:10.1038/nrmicro.2016.81 (2016).

10. Yang, X. H. *et al.* Mice transgenic for human angiotensin-converting enzyme 2 provide a model for SARS coronavirus infection. *Comparative medicine* **57**, 450-459 (2007).

11. Netland, J., Meyerholz, D. K., Moore, S., Cassell, M. & Perlman, S. Severe acute respiratory syndrome coronavirus infection causes neuronal death in the absence of encephalitis in mice transgenic for human ACE2. *J Virol* **82**, 7264-7275, doi:10.1128/jvi.00737-08 (2008).

12. Rivers, T. M. Viruses and Koch's Postulates. *Journal of bacteriology* **33**, 1-12 (1937).

Figure 1. Weight loss, virus replication and specific IgG production in hACE2 mice post infection with SARS-CoV-2. For weigh loss record, hACE2 mice (n=7) and WT mice (n=3) were experimentally challenged intranasally with SARS-CoV-2, and the ACE2-Mock mice (n=3) were used as control. And then the weight loss was recorded over 14 days (a, ANOVA, ***$p<0.001$). To screen virus replication, 12 mice were infected in each group, and 3 mice per group were sacrificed and their major organs harvested for viral load and virus titer at 1 dpi, 3 dpi, 5 dpi and 7 dpi respectively. The distribution of SARS-CoV-2 in the primary organs of ACE2-HB-01 mice was detected by qRT-PCR (b). Virus titers of lungs were determined on Vero E6 cells (c, unpaired t-test, ***$p<0.001$), and the virus isolated from lungs of ACE2-HB-01 mice at 3 dpi was observed by electron microscope (d). The specific IgG against SARS-CoV-2 was detected at 21 dpi by ELISA (e, unpaired t-test, ***$p<0.001$).

Figure 2. Gross pathology, histopathology, and immunohistochemistry of the lungs in SARS-CoV-2-infected hACE2 mice. a. Gross pathology and histopathology of lungs from WT-HB-01 mice (3 dpi), ACE2-Mock mice (3 dpi) and ACE2-HB-01 mice (3 dpi and 5 dpi). Postmortem examinations showed focal dark red lesions (red arrow) throughout the dorsal of the right middle lobe of the lung at 3dpi. The lesions progressed into multifocally scattered-dark reddish purple areas and palpable nodules (red arrow) throughout the right lobe of the lung at 5 dpi. Histopathological observation indicated that

moderate interstitial pneumonia with thickened alveolar septa (black arrows) and infiltration of lymphocytes (red arrows). The swollen and degenerative alveolar macrophages (green arrows) are scattered within the alveolar cavities at 3dpi and 5 dpi. b. Immunohistochemical examination of lungs of each group. The sequential sections were stained by HE and IHC, respectively. The viral antigens were observed in the alveolar macrophage (green arrows), the alveolar epithelia (blue arrows), and the degenerative and being desquamated bronchial epithelial cells (black arrows). Black bar = 100 µm, red bar = 50 µm.

Figure 3. Immunofluorescence analysis of viral antigens in lungs of SARS-CoV-2-

infected hACE2 mice. Co-localization of SARS-CoV-2 S protein and hACE2 receptor in

hACE2 mouse lungs, the sections were incubated with anti-SARS-CoV-2 S protein

antibody, anti-human ACE2 antibody, and DAPI. The lung sections of ACE2-Mock mice

(a-d). The lung sections of ACE2-HB-01 mice (e-h). The white arrows showed the viral S

protein (f) and hACE2 (g), respectively, the yellow arrow showed the merge of viral S

protein and hACE2 (h). White bar=25 µm.

Supplementary Figure 1. Special stains of the lungs in the WT-HB-01 and ACE2-HB-01 mice at 3dpi and 5 dpi. a. Modified Masson's Trichrome of the lung. Compared to the WT-HB-01 group, the increased collagen fiber (blue-stained fibers) in the thickened alveolar interstitium were observed in both the ACE2-HB-01 mice at 3dpi and 5 dpi. b. Periodic acid schiff (PAS) staining of the respiratory epithelium in bronchioles. A small amount of mucus was accumulated on the surface of bronchial epithelial cells. Black bar=40 μm.

Supplementary Figure 2. IHC was carried out to identify macrophages with MAC2, T lymphocytes with CD3, and B lymphocytes with CD19. a. Diffuse infiltration of macrophages (red arrow) in the expanded alveolar septum in the ACE2-HB-01 mice at 3 dpi and 5 dpi. b. Many T lymphocytes (red arrow) infiltrated the thickened alveolar septa in the first row of b at 5 dpi in the ACE2-HB-01 mice. A few B lymphocytes (red arrow) were observed in the ACE2-HB-01 mice. Black bar=40 µm.

Supplementary Figure 3. The histopathological observation of the organs in the WT-HB-01 and ACE2-HB-01 mice. There were no significant histopathological lesions in the organs, including myocardium, liver, spleen, kidney, cerebrum, intestine and testis in the ACE2-HB-01 mice compared with the ACE2-Mock mice. Black bar = 100 µm.

Supplementary Figure 4. The immunohistochemical (IHC) observation of the organs in the ACE2-HB-01 mice. There were no SARS-CoV-2 antigens in the organs, including myocardium, liver, spleen, kidney, cerebrum, intestine and testis. Red bar = 50 µm.

Supplementary Figure 5. Identification of 7D2 antibody against SARS-CoV-2 S1

protein. The plate coated by 0.2 ug SARS-CoV-2 S1 protein was incubated with 7D2

antibody as primary antibody (1:200) and detected using HRP-conjugated goat anti-

mouse secondary antibody. The titer of antibody was determined using enzyme-linked

immunosorbent assay (ELISA) assay. Significant differences are indicated with an

asterisk (unpaired t-test, $*p<0.05$).

412