Distinct Hierarchical Alterations of Intrinsic Neural Timescales Account for Different Manifestations of Psychosis

Kenneth Wengler, PhD[a,b*], Andrew T. Goldberg[b], George Chahine, MD[c], and Guillermo Horga, MD, PhD[a,b*]

a. Department of Psychiatry, Columbia University, New York, NY, USA

b. New York State Psychiatric Institute, New York, NY, USA

c. Department of Psychiatry, Yale University, New Haven, CT, USA

Corresponding Authors

Kenneth Wengler, PhD

Postdoctoral Research Fellow, Columbia University

Division of Translational Imaging, New York State Psychiatric Institute

Kenneth.Wengler@nyspi.columbia.edu

646-774-5571

Guillermo Horga, MD, PhD

Florence Irving Assistant Professor of Psychiatry, Columbia University

Division of Translational Imaging, New York State Psychiatric Institute

Guillermo.Horga@nyspi.columbia.edu

646-774-5810

Keywords

Psychosis, Intrinsic Neural Timescale, Schizophrenia, Excitation-Inhibition Balance, Hierarchical Perceptual Inference

Abstract

Hierarchical perceptual-inference models of psychosis may provide a holistic framework for understanding psychosis in schizophrenia, including heterogeneity in its clinical presentations. Particularly, hypothesized alterations at distinct levels of the perceptual-inference hierarchy may explain why positive symptoms such as hallucinations and delusions tend to cluster together yet sometimes manifest in isolation. To test this, we used a recently developed resting-state fMRI measure of intrinsic neural timescale (INT), which reflects the time window of neural integration and has been shown to capture hierarchical brain gradients. In analyses examining extended sensory hierarchies that we first validated in an independent sample, we found distinct hierarchical INT alterations for hallucinations versus delusions in the auditory and somatosensory systems, thus providing support for hierarchical perceptual-inference models of psychosis. Simulations using a large-scale biophysical model further suggested local elevations of the excitation-inhibition ratio at different hierarchical levels as a potential underlying mechanism. More generally, our work highlights the robustness and the utility of INT for studying hierarchical processes relevant to basic and clinical neuroscience.

Introduction

Hallucinations and delusions are burdensome symptoms that typically manifest together as the psychotic syndrome of schizophrenia. Perceptual-inference models of psychosis suggest that these symptoms result from alterations in the updating of internal models of the environment that are used to make inferences about external sensory events and their causes (Adams et al., 2013; Horga and Abi-Dargham, 2019; Sterzer et al., 2018). These models are receiving increasing empirical support (Adams et al., 2018; Baker et al., 2019; Cassidy et al., 2018; Davies et al., 2017; Powers et al., 2017; Teufel et al., 2015), yet current theories do not provide a satisfactory explanation for how hallucinations and delusions tend to co-occur but sometimes manifest in isolation. This suggests that these psychotic symptoms may share a common neurobiological mechanism but simultaneously depend on symptom-specific pathways.

We and others have proposed that this apparent tension may be resolved in the context of *hierarchical* perceptual-inference models (Adams et al., 2013; Baker et al., 2019; Corlett et al., 2009a; Corlett et al., 2018; Fletcher and Frith, 2009; Powers III et al., 2016; Sterzer et al., 2018). One possibility is that alterations at higher levels—supporting inferences on abstract hidden states like someone's intentions—may drive delusions, and alterations at lower levels—supporting inferences about lower-level features of stimuli such as stimulus presence or absence—may drive hallucinations (Baker et al., 2019; Corlett et al., 2009b; Davies et al., 2017; Horga and Abi-Dargham, 2019; Powers et al., 2017; Schmack et al., 2013). In addition to these symptom-specific pathways, alterations at any level may naturally propagate throughout the interdependent levels of the hierarchy (Chaudhuri et al., 2015), potentially explaining symptom co-occurrence. Importantly, neural systems supporting inference are thought to feature a hierarchical architecture of timescales that mirrors the hierarchical temporal dynamics of natural environments, where rapidly changing events are typically nested within slower changing contexts (Kiebel et al., 2008, 2009). Thus, inferences pertaining to slower changing contexts require neural systems with the ability to integrate information over longer periods of time; an ability consistent with the persistent neuronal activity that characterizes higher-level regions (Major and Tank, 2004; Mazurek et al., 2003).

A hierarchy of neural timescales has been observed in both single-neuron recordings in non-human primates (Murray et al., 2014) and functional magnetic resonance imaging (fMRI) recordings in humans (Hasson et al., 2015; Hasson et al., 2008; Honey et al., 2012; Lerner et al., 2011; Stephens et al., 2013), and was recapitulated by a large-scale biophysical model (Chaudhuri et al., 2015). Furthermore, a newly developed method based on resting-state fMRI, which was validated against EEG, was shown to capture a hierarchy of intrinsic neural timescales (INT) as well as alterations in psychopathology (Watanabe et al., 2019). Here, we specifically applied this fMRI measure to test whether hallucinations and delusions are associated with

distinct changes in INT at low and high levels of neural hierarchies, respectively. We

hypothesized that INT at these respective levels would increase with more severe symptoms,

reflecting increased neural integration of prior information, in line with behavioral findings in

hallucinations and delusions (Baker et al., 2019; Cassidy et al., 2018; Glaze et al., 2015; Mante et

al., 2013; Mazurek et al., 2003). If present, these INT changes should manifest as symptom-

specific differences in hierarchical gradients.

8 Results

INT maps were estimated as previously described (Watanabe et al., 2019) (**Materials and**

Methods). Briefly, the autocorrelation function of the fMRI signal at each voxel (or vertex) was

estimated and the sum of the autocorrelation coefficients during the initial positive period was

calculated. This initial positive period included all timepoints from the current timepoint (zero lag)

until the timepoint immediately preceding the first lagged timepoint with a non-positive

autocorrelation coefficient. To adjust for differences in temporal resolution, the sum was multiplied

by the repetition time (TR) of the fMRI data. This product was used as an index for INT (note that

values are similar to those from an exponential fit (Murray et al., 2014); **Figure 1—figure**

supplement 1). INT maps were parcellated using the HCP-multimodal parcellation (Glasser et al.,

2016) to facilitate further analysis. Additionally, T1w/T2w (myelin) and cortical-thickness maps were

obtained from high-resolution structural scans from the HCP database. Both of these measures

have previously been shown to capture the structural hierarchy (Burt et al., 2018; Fischl et al., 2008;

Wagstyl et al., 2015), consistent with the classic use of myeloarchitecture and cytoarchitecture for

cortical parcellation (Brodmann, 1909; Sarkissov et al., 1955; Vogt, 1911; Von Economo, 1929).

24 Selection and Multimodal Validation of Neural Hierarchies

Our hypothesis of symptom-specific INT differences in hierarchical gradients was agnostic with

respect to the specific neural hierarchies involved in psychosis, but the involvement of most sensory

modalities has been reported (Lewandowski et al., 2009; Postmes et al., 2014). Consistent with

prior empirical and theoretical work (Chaudhuri et al., 2015; Vázquez-Rodríguez et al., 2019), in a

subset of 100 unrelated young and healthy subjects from the HCP dataset we did not observe a

systematic relationship across the whole brain between INT (**Figure 1A**), an index of functional

hierarchy (Chaudhuri et al., 2015; Murray et al., 2014), and the two indices of structural hierarchy,

T1w/T2w and cortical thickness (Burt et al., 2018; Fischl et al., 2008; Wagstyl et al., 2015) (**Figure**

1—figure supplement 2). We thus decided to focus on specific, well-studied hierarchies of the

auditory, visual, and somatosensory systems that have been parcellated in humans, where the

functional and structural indexes of hierarchy appeared better aligned.

Figure 1. Model comparison to determine the hierarchical orderings of auditory, visual, and somatosensory systems. **A)**
Group-averaged intrinsic neural timescale (INT) map from the Human Connectome Project (HCP) dataset (N = 100; top),
parcellated group-averaged INT map (middle), and flattened cortex showing the parcels in the auditory, visual, and
somatosensory systems (bottom). Color coding of parcels indicates their anatomical location. **B)** Goodness-of-fit (R^2) of
linear mixed-effects models predicting different hierarchical orderings for the auditory system (left), visual system (middle),
and somatosensory system (right) from T1w/T2w and cortical thickness values in the HCP dataset (see **Materials and**
Methods). First, the winning ordering (i.e., the model with the best goodness of fit) for each system was determined for the
7 sensory cortex regions (bottom 4 models); then, winning orderings were determined for extended models with 2
downstream prefrontal cortex regions added to the respective winning models for sensory cortex (top 2 models). Note that,
for each of the four considered orderings within the sensory cortex for each system, only the 4 regions whose order is varied
(out of 7 regions) are shown to denote the models. For the auditory cortex, A1 was always the lowest order region while A4
and A5 were always the two highest order regions. For the visual cortex, V1, V2, and V3 were always the three lowest order
regions. For the somatosensory cortex, 5m, 7b, and 7a were always the three highest order regions. Null distributions were
generated by randomly permuting the hierarchical ordering (0^{th} – 95^{th} percentiles shown). **C)** Scatterplots showing INT
values plotted as a function of hierarchical level for the PFC-extended winning models in **B** (red outline) for the HCP dataset
(top) and the healthy control group in the schizophrenia combined dataset (N = 158; bottom).

Despite ample anatomical investigation in primates, ambiguities in the definition of anatomical hierarchies of these sensory systems still remain (Hilgetag et al., 1996; Kaas and Hackett, 2000) that have not been fully addressed in human MRI work. To address this issue, we used an anatomically informed, data-driven approach to determine the most suitable hierarchical orderings for each system. First, we determined the hierarchical orderings of the sensory cortex parcels (auditory, visual, or somatosensory) by selecting the ordering that was best predicted by T1w/T2w and cortical thickness parcel-wise values for each system (i.e., the ordering for which these values explained the most variance) in the HCP sample. We specifically constrained this comparison to the four most plausible hierarchical orderings for each system based on previous anatomical studies (Felleman and Van, 1991; Galaburda and Pandya, 1983; Hyvärinen and Poranen, 1978; Morel et al., 1993). The winning orderings were A1 → LBelt → MBelt → PBelt → RI → A4 → A5 for the auditory cortex, V1 → V2 → V3 → MT → V4 → V6 → V7 for the visual cortex, and 3b → 3a → 1 → 2 → 5m → 7b → 7a for the somatosensory cortex (**Figure 1B**). Using the same approach to build upon these winning orderings and capture the broadest possible range of the hierarchies, we then determined the hierarchical position of two additional prefrontal cortex (PFC) regions known to be downstream projections of the auditory, visual, and somatosensory cortices: area 8a and area 46 (Felleman and Van, 1991; Kaas and Hackett, 2000). For all sensory systems, area 46 was selected as the highest hierarchical level (**Figure 1B**). Notably, each of the PFC-extended winning models explained more variance than chance based on a null distribution of 10,000 random orderings (auditory system: $P_{permutation}$ < 10^{-4}; visual system: $P_{permutation}$ = 0.003; somatosensory system: $P_{permutation}$ = 0.001).

We then evaluated whether these winning hierarchies, selected solely based on structural measures of hierarchy, could also capture functional variability in the INT measure such that higher levels exhibited higher INT. Within the HCP sample, hierarchical position significantly correlated with INT for the auditory system (r_s = 0.87, P = 0.005; **Figure 1C**) and this correlation was above chance level based on a null distribution of 10,000 random orderings ($P_{permutation}$ = 0.003). The hierarchical ordering was further validated in an out-of-sample group of 158 healthy controls from the schizophrenia combined dataset (see **Materials and Methods**), where hierarchy similarly correlated with INT for the auditory system (r_s = 0.80, P = 0.01; **Figure 1C**). Although not significant, similar positive effects were observed for the visual system (in-sample: r_s = 0.27, P = 0.49, $P_{permutation}$ = 0.47; out-of-sample: r_s = 0.22, P = 0.58; **Figure 1C**). Trend-level results were observed in the somatosensory system (in-sample: r_s = 0.58, P = 0.108, $P_{permutation}$ = 0.097; out-of-sample: r_s = 0.80, P = 0.014; **Figure 1C**). Thus, we empirically validated extended sensory hierarchies that captured variability in structural and functional hierarchical indices across two independent samples, although this was surprisingly less evident for the visual system.

Assessment of Robustness and Reliability in the HCP Dataset

Next, we set out to determine the robustness and reliability of INT. We focused on head motion, a

common source of artifacts in fMRI data (Power et al., 2012). Head motion during data acquisition

was associated with generally decreased INT values (181 out of 188 parcels, $P_{permutation}$ = 0.01;

Figure 1—figure supplement 3). Yet these effects were comparable across hierarchical levels

(auditory system: r_s = -0.23, $P_{permutation}$ = 0.805; visual system: r_s = -0.38, $P_{permutation}$ = 0.649;

somatosensory system: r_s = -0.88, $P_{permutation}$ = 0.054; **Figure 1—figure supplement 3**). No effects

were observed for gender or age (all $P_{permutation}$ > 0.174). Finally, INT maps showed excellent

reliability between the first and last 5 minutes of the fMRI acquisition [median ICC(2,1) $\pm$

interquartile range across voxels: 0.94 $\pm$ 0.03; **Figure 1—figure supplement 4]**.

Table 1. Participant Characteristics

Variable	Healthy Controls					Schizophrenia Subjects				
	BGS	COBRE	NMCH	UCLA	All	BGS	COBRE	NMCH	UCLA	All
N	24	42	25	67	158	40	31	26	30	127
Age, mean (SD), y	36.0 (13.1)	33.9 (10.4)	29.8 (7.2)	32.3 (8.5)	32.9 (9.7)	31.1 (12.6)	30.5 (11.9)	30.5 (6.2)	35.0 (8.9)	31.8 (10.6)
Male sex, No. (%)	22 (92)	34 (81)	16 (64)	54 (81)	126 (80)	38 (95)	26 (84)	19 (73)	22 (73)	105 (83)
Framewise Displacement, mean (SD), mm	0.15 (0.04)	0.14 (0.05)	0.14 (0.11)	0.10 (0.04)	0.13 (0.06)	0.16 (0.06)	0.17 (0.06)	0.12 (0.06)	0.13 (0.04)	0.15 (0.06)
Delusion Score, mean (SD)	NA	NA	NA	NA	NA	2.3 (1.6)	1.7 (1.5)	3.2 (1.9)	2.5 (1.4)	2.4 (1.7)
Hallucination Score, mean (SD)	NA	NA	NA	NA	NA	2.1 (1.6)	1.7 (1.4)	2.9 (2.0)	2.2 (1.6)	2.2 (1.7)
Conceptual Disorganization Score, mean (SD)	NA	NA	NA	NA	NA	0.9 (1.3)	0.6 (1.0)	2.0 (1.6)	1.4 (1.4)	1.2 (1.4)
Emotional Withdrawal Score, mean (SD)	NA	NA	NA	NA	NA	1.8 (1.2)	1.2 (1.3)	3.4 (1.7)	2.3 (1.5)	2.1 (1.6)
Social Withdrawal Score, mean (SD)	NA	NA	NA	NA	NA	1.8 (1.4)	1.3 (1.4)	3.2 (1.7)	2.7 (1.6)	2.2 (1.6)
Blunted Affect Score, mean (SD)	NA	NA	NA	NA	NA	1.8 (1.6)	1.6 (1.5)	3.3 (1.6)	1.1 (1.1)	1.9 (1.6)
Alogia Score, mean (SD)	NA	NA	NA	NA	NA	1.3 (1.6)	1.2 (1.4)	2.1 (1.7)	0.9 (1.6)	1.3 (1.6)

BGS, BrainGluSchi; NMCH, NMorphCH; SD, standard deviation

Exploratory Analysis of Intrinsic Neural Timescales in Schizophrenia versus Health

Although our primary hypothesis dealt with hierarchical differences between hallucinations and

delusions, we first present exploratory analyses of diagnosis effects on INT. Participant

characteristics are listed in **Table 1**. Relative to controls (N = 158), patients (N = 127) exhibited a

small-to-moderate, but widespread, reduction of INT (98 out of 188 parcels, $P_{permutation}$ = 0.013;

Figure 2E). A similar result was observed in a voxel-wise analysis (**Figure 2—figure supplement**

1). The INT reductions in patients were however comparable across hierarchical levels (all

$P_{permutation}$ > 0.40; **Figure 2F**). *In silico* simulations using a large-scale biophysical model (Chaudhuri

et al., 2015) suggested that the global INT reduction in patients could be neuronally implemented

by globally reduced excitation-inhibition (E/I) ratio (**Figure 2—figure supplement 2**).

Figure 2. Exploratory analyses show that patients with schizophrenia exhibit widespread reductions of intrinsic neural

timescales compared to healthy controls. **A)** Parcellated group-averaged intrinsic neural timescale (INT) map for healthy

controls (N = 158). **B)** Parcellated group-averaged INT map for patients with schizophrenia (N = 127). **C)** t-statistic

(Cohen's d) map showing the contrast of patients greater than controls. Across most parcels, INT is shorter in patients

than controls in a regression model ($M1_{exploratory}$) controlling for age, gender, framewise displacement, and data-collection

site. Only the left hemisphere is shown because statistical analyses were performed after averaging the values in each

parcel across left- and right-hemispheres. **D)** To illustrate the effect of reduced INT in patients with schizophrenia, the

group-averaged, whole-brain-averaged autocorrelation functions were estimated from subjects with fMRI data acquired

with the same repetition time (top; controls: N = 132; patients: N = 101). The group-averaged autocorrelation function for

patients crosses the zero point on the y axis (i.e., autocorrelation coefficient = 0) sooner than in controls, demonstrating

the global reduction of INT in patients. The flattened cortex shows the parcels in the auditory, visual, and somatosensory

systems for reference (bottom). **E)** Scatterplot showing t-statistic values from the regression model ($M1_{exploratory}$), plotted as

a function of the INT rank (determined from the group-averaged INT map from HCP subjects). Each datapoint represents

one parcel. **F)** Scatterplots showing t-statistic values from parcels within the auditory (left), visual (middle), and

somatosensory (right) hierarchies plotted as a function of hierarchical level. No hierarchical effects for schizophrenia

diagnosis were observed.

Hierarchical Differences of Intrinsic Neural Timescales Between Hallucinations and

Delusions

Our *a priori* hypothesis was that hallucinations and delusions were associated with alterations of

INT at different hierarchical levels, leading to distinct changes in hierarchical gradients. To test this,

we determined the unique variance associated with the effect of interindividual variability in
hallucination and delusion severity on INT. As expected, severity of hallucinations and delusions in
our sample were correlated (r_s = 0.62, P < 0.01) but exhibited sufficient unique variance [$(1 - R^2)$ =
0.62] to evaluate their independent contributions. Severity of these symptoms was uncorrelated
with antipsychotic dose among the 109 patients with available data (chlorpromazine equivalents:
both P > 0.86), making it an unlikely confound (**Figure 3—figure supplement 1**).

The main model (M2; **Materials and Methods**) we used as a primary test of main effects and
interactions of symptoms on hierarchical INT gradients—which also included interaction terms for
each sensory system—was significant (omnibus $F_{11,41}$ = 5.52, P < 10^{-4}). Critically, within this model
we found hierarchical-gradient effects that differed significantly between hallucinations and
delusions in the expected directions for 2/3 systems (auditory system, symptom-by-hierarchical-
level interaction: t_{42} = 4.59 [95% bootstrap confidence interval: 3.39, 9.08], Cohen's f^2 = 1.00,
$P_{permutation}$ = 0.001; visual: t_{42} = -2.06 [-6.19, 0.16], f^2 = 0.11, $P_{permutation}$ = 0.083; and somatosensory:
t_{42} = 3.50 [2.19, 7.35], f^2 = 0.41, $P_{permutation}$ = 0.011; **Figure 3**). For the auditory system, this
interaction was driven by significant hierarchical-gradient effects in opposite directions for
hallucinations (hierarchical-level effect: t_{42} = -3.50 [-8.42, -2.24], f^2 = 0.41, $P_{permutation}$ = 0.010) and
delusions (hierarchical-level effect: t_{42} = 2.99 [1.18, 6.37], f^2 = 0.27, $P_{permutation}$ = 0.025). For the
somatosensory system, this effect was driven by a trend-level negative hierarchical-gradient effect
for hallucinations (hierarchical-level effect: t_{42} = -2.35 [-5.91, -1.00], f^2 = 0.15, $P_{permutation}$ = 0.056)
and a significant positive hierarchical-gradient effect for delusions (hierarchical-level effect: t_{42} =
2.60 [1.43, 5.57], f^2 = 0.19, $P_{permutation}$ = 0.042; **Figure 3**). For the visual system, hierarchical-gradient
effects were not significant for either symptom (hallucination hierarchical-level effect: t_{42} = 0.27 [-
1.06, 3.62], f^2 = 0.00, $P_{permutation}$ = 0.901; delusion hierarchical-level effect: t_{42} = -2.01 [-6.01, 0.53],
f^2 = 0.11, $P_{permutation}$ = 0.087). We also found significant three-way interactions with sensory system
(see statistics in **Figure 3B**) but these were not *a priori* tests (see also **Discussion** for issues of
interpretability). Examining the significant symptom effects further, we observed that, within the
auditory system, patients with high-severity hallucinations exhibited a numeric increase of INT at
lower levels of the hierarchy relative to those with low-severity hallucinations, leading to a
compression of the INT hierarchical gradient (**Figure 3C**); in contrast, in both the auditory and
somatosensory systems, patients with high-severity delusions exhibited a numeric increase of INT
at higher levels of the hierarchy relative to those with low-severity delusions, leading to a more
pronounced INT hierarchical gradient.

To correct for multiple comparisons, we carried out a family-wise permutation test determining
the probability of spuriously obtaining the set of significant *a priori* effects we observed in support
of our original hypothesis. Based on the chance level of jointly observing negative hierarchical-
gradient effects for hallucination severity in at least 1/3 systems, *and* positive hierarchical-gradient

effects for delusion severity in at least 2/3 systems, *and* interaction effects of hierarchy-by-symptom

in the expected direction in at least 2/3 systems; this analysis suggested that the observed set of

results was statistically above chance (set-level $P_{permutation}$ = 0.014). Furthermore, based on the

chance level of observing a significant negative hierarchical-gradient effect for hallucinations, *and*

a significant positive hierarchical-gradient effect for delusions, *and* a significant symptom-by-

hierarchical-level interaction (i.e., all 3 effects in one system); this analysis suggested that the

observed set of results in the auditory system was also statistically above chance (set-level

$P_{permutation}$ = 0.043).

Figure 3. *A priori* analyses show that hallucinations and delusions exhibit distinct hierarchical-gradient effects on intrinsic

neural timescales in the auditory and somatosensory systems. **A)** Scatterplots showing *t*-statistic values—from a

regression model (M1$_{primary}$) including all 7 symptoms and controlling for age, gender, framewise displacement, and data-

collection site—for hallucination-severity (top) and delusion-severity (bottom) effects from parcels within the auditory (left),

visual (middle), or somatosensory (right) hierarchies plotted as a function of hierarchical level (using PFC-extended

winning hierarchies; **Figure 1**). **B)** Summary of results from a model (M2) including symptom-severity effect

(hallucinations or delusions), hierarchical level, sensory system (auditory, visual, or somatosensory), and their

interactions. These results demonstrate: (1) in the auditory system, a significant difference in the relationship between hallucination severity and hierarchical level versus that for delusion severity and hierarchical level (b); (2) significant hierarchical effects of hallucination severity (a) and delusion severity (c) in the auditory system; (3) in the somatosensory system, a significant difference in the relationship between hallucination severity and hierarchical level versus that for delusion severity and hierarchical level (e); and (4) a significant hierarchical effect of delusion severity in the somatosensory system (d). Note that different symptoms and systems are used as reference (implicit variable) across three plots to show each of the relevant effects, which were tested within a single model (M2). **C)** To illustrate the effects, the group-averaged autocorrelation functions were estimated from subjects with fMRI data acquired with the same repetition time (N = 10 for each group). High severity patients were the 10 subjects with the highest residual symptom scores after regressing out all other symptoms; low severity patients were the 10 subjects with the lowest residual symptom scores. The group-averaged autocorrelation functions are shown for high-severity (solid lines) and low-severity (dashed lines) hallucination patients from low and high levels of the auditory hierarchy (A1 and area 46, top). The group-averaged autocorrelation functions are also shown for high-severity and low-severity delusion patients from low and high levels of the auditory hierarchy (middle). The group-averaged autocorrelation functions are finally shown for high-severity and low-severity delusion patients from low and high levels of the somatosensory hierarchy (area 3b and area 46, bottom). These plots depict high-severity hallucination patients exhibiting a compression of the auditory hierarchical gradient and high-severity delusion patients instead exhibiting an expansion of this gradient.

Post-Hoc Analysis on the Specificity of INT Hierarchical-Gradient Effects

In a post-hoc analysis, we then investigated the specificity of these hierarchical-gradient effects to the positive psychotic symptoms under investigation. To this end, we determined hierarchical-gradient effects individually for each symptom in the auditory and somatosensory systems using a model including symptom-severity effect (only one symptom), hierarchical level, sensory system (auditory, visual, and somatosensory), and their interactions. In the auditory system, conceptual disorganization was the only symptom—other than hallucinations and delusions—that showed a significant effect (hierarchical-level effect: t_{21} = -2.80 [-5.31, -1.10], f^2 = 0.60, $P_{permutation}$ = 0.036; **Figure 3—figure supplement 2**), although this effect was weaker than that for hallucinations (hierarchical-level effect: t_{21} = -4.38 [-10.31, -3.28], f^2 = 10.57, $P_{permutation}$ = 0.005; **Figure 3—figure supplement 2**). These results thus suggest some specificity to positive symptoms, which conceptual disorganization is classically defined as (Association, 2013; VandenBos, 2007) (but see (van der Gaag et al., 2006)), consistent with a stronger correlation of conceptual disorganization with positive symptoms (average r_s = 0.48) versus negative symptoms (average r_s = 0.23) in our sample. Indeed, a permutation test comparing the average strength of hierarchical-gradient effects (i.e., mean absolute t-statistics) for positive versus negative symptoms (i.e., blunted affect, social withdrawal, emotional withdrawal, and alogia) showed the effects of positive symptoms to be significantly larger than the effects of negative symptoms ($P_{permutation}$ = 0.043).

In the somatosensory system, no symptoms other than delusions showed a significant hierarchical-gradient effect. Hallucinations however showed the strongest negative effect (hierarchical-level effect: t_{21} = -2.23 [-6.79, -1.75], f^2 = 0.31, $P_{permutation}$ = 0.079; **Figure 3—figure supplement 2**).

1   Thus, although the hierarchical-gradient effects were not unique to the two symptoms under
2   investigation—which is not to be required under perceptual-inference models of psychosis and
3   which could suggest model extensions to account for additional phenomena—these effects were
4   strongest for, and relatively specific to, positive symptoms.
5
Altered E/I Ratio as a Potential Biological Mechanism
To explore candidate biological mechanisms for the effects we observed *in vivo*, we leveraged a
large-scale biophysical model previously shown to capture intrinsic timescale hierarchies
(Chaudhuri et al., 2015). This model includes 29 recurrently connected cortical nodes, with
connection strengths based on tract-tracing macaque studies. Given growing evidence for E/I
imbalance in schizophrenia (see (Foss-Feig et al., 2017; Jardri et al., 2016) for review) and the
hypothesized local increases of INT, we fit the biophysical model to our data to explore whether
our results could be driven by local increases in E/I ratio. To do this, for each symptom
(hallucinations and delusions) we ft free parameters controlling the hierarchical level of the peak
increase in local E/I ratio, the magnitude of the E/I ratio increase (percent change in local recurrent
connection strength) at this level, and the E/I ratio change at the minimum (the most distant level
from the peak), to the observed difference of INTs across three different symptomatic profiles
(based on the M1$_{primary}$ fitted INT values; **Figure 4**) in 6 nodes that directly correspond to levels of
our visual hierarchy (V1, V2, V4, MT, 8l, and 46d). The *in silico* simulations using the best-fitting
parameters were able to recapitulate the INT effects of hallucinations and delusions (compression
or expansion of the timescale gradient, respectively) via local increases in E/I ratio at low or high
levels of the hierarchy, respectively; specifically, the best-fitting levels of the peak increase in local
E/I ratio were levels 1 and 8, respectively (**Figure 4**). Interestingly, given the relatively greater
strength of both recurrent and long-range connections at higher levels that is built into the model,
the required E/I ratio increase to achieve the observed changes of INT was considerably smaller
for the delusion-related alteration in the higher level (ΔE/I = 4.02%) compared to the hallucination-
related alteration in the lower level (ΔE/I = 21.61%). Importantly, an additive combination of the
low-level (hallucinations) and the high-level (delusions) changes closely approximated the
combined effect of hallucinations and delusions, which consisted of a general increase of INT with
no clear change in the hierarchical gradient. This suggests that additivity of these alterations could
explain symptom co-occurrence.

Figure 4. Hallucination-severity and delusion-severity effects are recapitulated by elevated excitatory-inhibitory (E/I) ratios at different hierarchical levels. Simulation results are from a large-scale biophysical model of primate cortex (**Materials and Methods**). **A)** The model consists of 29 nodes with local excitatory (red triangles) and inhibitory (blue circles) pools of neurons; only two nodes—high (top) and low (bottom) hierarchical levels—are shown for illustrative purposes. These nodes have both local (recurrent) and long-range (across nodes) connections. The hierarchy within the visual system (V1, V2, V4, MT, 8l, and 46d) was used for these simulations given the lack of tract-tracing data for the auditory cortex, but the qualitative pattern of results applies to hierarchical effects in any given sensory system. Theoretical perturbations to the model are marked by lightning bolts. Thicker or thinner lines with respect to the reference scenario reflect increased or decreased connection strengths, respectively. **B)** Scatterplots show intrinsic neural timescale (INT) values estimated for a simulated reference (unaltered model; left column) scenario, which demonstrates the natural hierarchy of timescales (note that similar results are observed when using an altered model with global decrease in E/I ratio as the reference scenario). Relative to the reference scenario, increased excitatory-to-excitatory recurrent connection strengths or decreased excitatory-to-inhibitory recurrent connection strengths (elevated E/I ratio) at low hierarchical levels lead to increased INT at these levels (middle-left), emulating the compressed hierarchical gradient related to hallucinations. Relative to the

reference scenario, increased excitatory-to-excitatory recurrent connection strengths or decreased excitatory-to-inhibitory recurrent connection strengths (elevated E/I ratio) at high hierarchical levels lead to increased INT at these levels (middle-right), emulating the expanded hierarchical gradient related to delusions. Relative to the reference scenario, increased excitatory-to-excitatory recurrent connection strengths or decreased excitatory-to-inhibitory recurrent connection strengths (elevated E/I ratio) at both low and high hierarchical levels lead to increased INT throughout the hierarchy (right), with an unaltered hierarchical gradient similar to that for patients with both hallucinations and delusions. Insets for **A** show predicted INT values in the auditory system for the 4 scenarios based on *in vivo* results (fitted data for M1$_{primary}$ for each scenario, with the presence of a symptom corresponding to fitted values for a score of 5 for that symptom); the x-axis scale is identical to the main scatter plots. **C)** Scatterplots showing the difference in simulated INT values between each of the 3 pathological scenarios and the reference scenario. By allowing the E/I ratios to vary, the fitted biophysical model can recapitulate the *in vivo* INT changes with a negative hierarchical-gradient effect for hallucinations, a positive hierarchical-gradient effect for delusions, and an overall INT increase (without a manifest hierarchical-gradient effect) for both hallucinations and delusions. Red arrowheads depict the location and strength of the maximum E/I ratio increase. Insets for **B** show the INT changes in the auditory hierarchy for the 3 pathological scenarios based on *in vivo* results (fitted data for M1$_{primary}$); both the x-axis and y-axis scales are identical to the main scatterplots. **D)** Scatterplots showing the fitted changes to E/I ratios for the 3 pathological scenarios: the *in vivo* INT changes associated with hallucinations can be recapitulated by elevated E/I ratio at the lowest hierarchical level and those associated with delusions by elevated E/I ratio (of smaller magnitude) at the highest hierarchical level, with the addition of these two alterations capturing the changes in patients with both hallucinations and delusions.

In a follow-up analysis, we further explored our *in vivo* data for evidence of this additive effect, focusing on the auditory system. We first compared the average INT across all auditory system parcels between patients with both high hallucination and delusion scores (i.e., subjects with a score of 5 for both symptoms; $N = 11$) and patients with neither hallucinations nor delusions (i.e., subjects with a score of 0 for both symptoms; $N = 18$). Here, we observed significantly higher average INT in patients with high-severity hallucinations and delusions ($t_{27} = 1.84$, $P = 0.038$; one-tailed two-sample t-test). Second, we fit a linear model predicting auditory parcel INT as a function of hierarchical level, allowing separate intercepts for each of the two groups, and an interaction between hierarchical level and group. Here, we found that the intercept was indeed higher for patients with high-severity hallucinations and delusions compared to patients with neither symptom ($t_{257} = 2.04$, $P = 0.043$). Furthermore, we found no difference in the hierarchical gradients between these groups, with a non-significant hierarchical-level-by-group interaction ($t_{257} = 0.65$, $P = 0.519$). Although preliminary, these results provide some support for the notion of additive hierarchical alterations underlying hallucinations and delusions.

Control Analyses Examining Alternative Definitions of Auditory Hierarchies

Given that the hierarchical-gradient effects supporting our initial hypotheses were clearest in the auditory system—a system thought to comprise dual processing streams—we considered the impact of alternative definitions of the auditory hierarchy on our results.

Diverging auditory streams with downstream projections to dorsal (areas 8a and 46) versus

ventral (areas 10 and 12vl) PFC have been described (Kaas and Hackett, 2000). We thus conducted an ordering-selection analysis for the ventral stream similar to that presented above for the dorsal stream. Within the ventral stream, area 12vl was better predicted as the highest hierarchical level, and the winning ordering explained more variance than chance based on a null distribution of random orderings (ventral auditory system: $P_{permutation} < 10^{-4}$; **Figure 1—figure supplement 5A**). Furthermore, similar to the dorsal stream, hierarchical level in the ventral stream correlated with INT (in-sample: $r_s = 0.87$, $P = 0.005$, $P_{permutation} = 0.003$; out-of-sample: $r_s = 0.80$, $P = 0.014$; **Figure 1—figure supplement 5B**). Given these dual auditory streams and the corresponding validated hierarchies, we explored potential differences in hierarchical-gradient effects for hallucination and delusion severity for the dorsal versus ventral streams. The model explaining symptom effects and their differences by hierarchical level and their interaction by symptoms and auditory stream was significant (omnibus $F_{7,27} = 8.60$, $P < 10^{-4}$). We further found a significant difference in the symptom-by-hierarchical-level effects between the dorsal and ventral auditory streams (symptom-by-hierarchical-level-by-processing-stream interaction: $t_{28} = 1.75$ [1.02, 3.90], $f^2 = 0.12$, $P_{permutation} = 0.005$; **Figure 3—figure supplement 3**). In the ventral stream, we did not find a significant difference in the hierarchical-gradient effects between hallucinations and delusions (symptom-by-hierarchical-level interaction: $t_{28} = 2.01$ [-0.37, 6.43], $f^2 = 0.17$, $P_{permutation} = 0.159$; **Figure 3—figure supplement 3**); we found a trend-level hierarchical-gradient effect of hallucination severity (hierarchical-level effect: $t_{28} = -2.56$ [-7.04, -0.73], $f^2 = 0.31$, $P_{permutation} = 0.098$; **Figure 3—figure supplement 3**) and no effect of delusion severity (hierarchical-level effect: $t_{28} = 0.28$ [-2.42, 3.36], $f^2 = 0.00$, $P_{permutation} = 0.445$; **Figure 3—figure supplement 3**). Interestingly, comparing the dorsal and ventral auditory streams, we observed a significant difference in the hierarchical-gradient effect of delusion severity (hierarchical-level-by-processing-stream interaction: $t_{28} = 1.89$ [1.39 – 3.49], $f^2 = 0.15$, $P_{permutation} = 0.003$; **Figure 3—figure supplement 3**). These results thus support an involvement of dorsolateral PFC in delusions, consistent with prior work (Corlett et al., 2007).

As an additional control for the uncertainty in defining the auditory hierarchy, we also adopted an anatomically agnostic, data-driven approach. First, the symptom effects (M1$_{primary}$) were estimated for each voxel within the 9 auditory system parcels (600 voxels total). Second, each voxel was ranked based on its INT value from the average INT map in the HCP dataset. Third, 10 equally spaced bins along the INT ranking (60 voxels per bin) were created, which comprised the levels of the data-driven hierarchy, and the voxelwise t-statistics (from M1$_{primary}$) were averaged per bin. Similar to the main analysis (**Figure 3**), a model that included main effects and interactions of symptoms on the hierarchical INT gradient was significant (omnibus $F_{4,15} = 10.20$, $P = 0.001$). Within this model, we found hierarchical-gradient effects that differed significantly between hallucinations and delusions (symptom-by-hierarchical-level interaction: $t_{16} = 5.19$ [0.48 23.26], f^2

= 10.12, $P_{permutation}$ = 0.003; **Figure 3—figure supplement 4**). This interaction was driven by significant hierarchical-gradient effects in opposite directions for hallucinations (hierarchical-level effect: t_{16} = -2.79 [-16.72, 2.40], f^2 = 0.25, $P_{permutation}$ = 0.049; **Figure 3—figure supplement 4**) and delusions (hierarchical-level effect: t_{16} = 4.55, [1.04, 19.76], f^2 = 4.04, $P_{permutation}$ = 0.008; **Figure 3—figure supplement 4**).

Discussion

Using a recently developed method for measuring neural timescales from resting-state fMRI data, we set out to test the hypothesis that hallucinations and delusions are associated with dysfunctions at different levels of neural hierarchies. Using established structural indices of hierarchy (myelin and cortical thickness) and INT (a functional index of hierarchy) in independent samples, we first validated extended sensory hierarchies for the auditory, visual, and somatosensory systems that captured substantial variability in the hierarchical MRI indices. After further showing excellent reliability of the INT measure, in exploratory analyses we showed for the first time that patients with schizophrenia have globally reduced INT. Most importantly, our primary analyses comparing INT effects for hallucinations versus delusions in the validated hierarchies demonstrated that these symptoms are associated with distinct changes along the hierarchical gradients in the auditory and somatosensory systems, an effect we failed to observe in the visual system.

Hierarchical models of perceptual inference posit that perceptions are shaped by prior beliefs (Dayan et al., 1995; Friston and Kiebel, 2009; Kiebel et al., 2009; Lee and Mumford, 2003; Rao and Ballard, 1999) through reciprocal message-passing across different levels of sensory-processing hierarchies; an architecture that mirrors the known anatomy of sensory systems (Felleman and Van, 1991; Glasser et al., 2016; Kaas and Hackett, 2000; Markov et al., 2012; Van Essen et al., 1992; Young, 1993). In this scheme, higher levels of the neural hierarchy are thought to represent increasingly abstract belief states that evolve at slower timescales (Kiebel et al., 2009). For instance, during speech perception, the hierarchical structure of linguistic units can be parsed such that lower levels of auditory processing encode syllable information at faster timescales while higher levels encode sentence information at slower timescales (Ding et al., 2016). An emerging body of work in psychosis has linked hallucinations to preferential biases towards prior beliefs in low-level inferences during detection or estimation of stimulus features (Cassidy et al., 2018; Davies et al., 2017; Powers et al., 2017; Teufel et al., 2015) and delusions to preferential biases towards prior beliefs in higher-level inferences about more abstract hidden-states (Baker et al., 2019). The observed biases towards prior beliefs in past behavioral work can be framed as primacy biases (Baker et al., 2019), where past information is weighted more heavily during the inferential process, or equivalently, where information is integrated over longer

timescales (Glaze et al., 2015). Temporal integration is at the core of the neural implementation

of perceptual inference (Mazurek et al., 2003) and is thought to depend crucially on recurrent

network activity (Chaudhuri et al., 2015; Mante et al., 2013). Thus, a plausible neuronal

implementation of primacy biases at a given level of the hierarchy would be through increases in

the strength of recurrent excitation or decreases in the strength of recurrent inhibition (i.e.,

elevated E/I ratio) leading to relative increases in neural timescales.

Here, we observed changes in neural timescales across anatomical levels of neural

hierarchies, that differed between hallucinations and delusions; an effect that was most evident in

the auditory hierarchy. Patients with more severe hallucinations exhibited a less pronounced

hierarchical gradient, consistent with increased timescales at lower levels compared to those with

less severe hallucinations; and those with more severe delusions instead exhibited a more

pronounced hierarchical gradient, consistent with increased timescales at higher levels compared

to those with less severe delusions (**Figure 3C**). We further recapitulated these findings by

respectively elevating E/I ratios at low or high hierarchical levels of a large-scale biophysical

model (Chaudhuri et al., 2015). These E/I ratio elevations could, in principle, result from

alterations in NMDA or dopamine activation at these levels and are thus plausible under widely

supported glutamatergic and dopaminergic theories of psychosis (Brunel and Wang, 2001; Corlett

et al., 2009a; Corlett et al., 2011; Durstewitz and Seamans, 2002; Jardri et al., 2016; Javitt et al.,

2012; Weinstein et al., 2017). These results thus demonstrate distinct hierarchical alterations for

hallucinations and delusions that are generally consistent with our hypothesized hierarchical

framework, where distinct hierarchical alterations provide symptom-specific pathways that

together may explain symptom co-occurrence and present a candidate biological mechanism for

the psychotic syndrome.

Such hierarchical alterations may also fit well with the phenomenological timescale of these

symptoms. Clinical observation indicates that hallucinations—like rapidly changing sensory

events—change transiently and intermittently over seconds or minutes, while delusions—like

slowly changing 'conceptual' beliefs—evolve more slowly over days or months, but their average

severity over a given period of time typically evolves in parallel. All of these clinical features are

consistent with a hierarchical structure of nested timescales (Kiebel et al., 2008). While our

findings generally support this notion, computational work explicitly laying out the proposed model

in the context of inferential alterations in psychosis and empirical confirmations are warranted.

One outstanding question is how the alterations in neural timescales we observed here in relation

to delusions, which may predominate in high levels of the hierarchy yet manifest as changes on

the order of seconds, may drive delusions evolving over much longer timescales. One possible

explanation is that, while delusion maintenance may involve longer-term memory processes, the

underlying mechanism originating delusions transpires more rapidly and disrupts inferences at

timescales on the order of seconds, consistent with prior work (Baker et al., 2019). Since encoded memories likely reflect inferences summarizing information at a given time point (Shadlen and Shohamy, 2016), high-level inferential biases at shorter timescales may be sufficient to shape long-term conceptual memories in a way that further propagates biases over long time-periods, particularly under primacy biases that decrease the relative influence of newer information. Although less critical, it is also worth noting that INT reflects differences in resting circuit dynamics, the timescale of which is likely to be substantially magnified when these circuits are engaged (Chaudhuri et al., 2015; Hasson et al., 2008).

Our opposing findings for diagnosis (globally reduced INT) and symptom severity (focally increased INT) may be reconciled within pathophysiological models of psychosis which posit a key role for compensatory processes in schizophrenia. Hallucinations and delusions have been proposed to represent a temporary state of the illness that results from a failed attempt to compensate for a trait-like, baseline deficit (Adams et al., 2013; Moutoussis et al., 2011). Relatedly, long-standing circuit-level theories have suggested that psychosis-related increases in striatal dopamine transmission is secondary to a primary cortical deficit (Weinberger, 1987). In particular, previous frameworks suggest that psychotic states are associated with excessive prior biases in inferential processes arising as an overcompensation for a baseline trait consisting of the opposite bias (Adams et al., 2013; Horga and Abi-Dargham, 2019). From a biophysical-modeling standpoint, the trait-like baseline deficit in schizophrenia could consist of reduced E/I ratio (for instance, arising from NMDA-receptor hypofunction of excitatory neurons (Cavanagh et al., 2019)), which behaviorally would translate into general recency biases. In contrast, a failed compensatory mechanism could result in local increases in E/I ratio at different levels leading to distinct primacy biases and psychotic symptoms (Lam et al., 2017). While speculative, the compensatory changes could arise from dopaminergic alterations that effectively increase E/I ratio by preferentially boosting NMDA-receptor function of excitatory neurons (or other changes dampening NMDA-receptor function of inhibitory neurons) (Brunel and Wang, 2001).

Our finding of a preferential involvement of the auditory system for hallucinations is not surprising, given that in schizophrenia this symptom tends to predominate in the auditory modality despite also presenting in other modalities (Lim et al., 2016; Waters and Fernyhough, 2017); auditory-cortex abnormalities in schizophrenia are also well established (Javitt and Sweet, 2015). Our finding of somatosensory system involvement for delusions is also consistent with previous work on delusions of passivity (Brüne et al., 2008; Spence et al., 1997) and deficits in sensory attenuation via motor predictions in schizophrenia (Shergill et al., 2005; Shergill et al., 2014). However, despite our failure to detect differential alterations in the visual system, substantial evidence also suggests visual-cortex abnormalities in schizophrenia (Butler et al., 2008; Çavuş et al., 2012; Dorph-Petersen et al., 2007). And evidence from subclinical populations suggests

symptom-specific hierarchical alterations in visual tasks (Davies et al., 2017). Furthermore, the

general differences of INT values between sensory systems (**Figure 1**), while potentially relevant

to psychosis in and of themselves, could imply differential sensitivity in our analyses across

sensory domains. The negative findings in the visual system could similarly result from the poorer

correspondence between levels of visual hierarchy and hierarchical MRI indices (not only for INT,

but also surprisingly for the structural indices) compared to the other systems (**Figure 1**).

Previous empirical work using structural (Bassett et al., 2008) and functional measures

(Dondé et al., 2019; Leitman et al., 2010; Yang et al., 2016), suggests hierarchical alterations in

schizophrenia. This work however did not evaluate hierarchical differences between symptoms

and used measures that differ fundamentally from INT. In exploratory analyses testing diagnostic

effects, we found global INT reductions in schizophrenia but no clear shifts in the hierarchical INT

gradients (see **Figure 2—figure supplement 2** for initial evidence of an exponential effect). We

used the same approach as a previous study measuring INT in individuals with autism, which

reported decreased INT in visual cortex (and increased INT in the caudate) (Watanabe et al.,

2019). Consistent with our interpretation, this INT phenotype was linked to other data in autism

supporting excessive weighting of sensory evidence (Gollo, 2019; Lawson et al., 2017)—akin to a

decreased primacy bias, or a recency bias.

Some limitations are worth discussing. Because 93% of the patients (with available

medication data) were taking antipsychotics, we cannot definitively rule out medication

confounds, particularly on diagnosis effects. However, we observed similar effects when

controlling for dose, no correlations between dose and symptoms, and did not expect differential

neural modulations for hallucinations versus delusions (**Figure 3—figure supplement 1**). Future

studies should elucidate medication effects on INT. Furthermore, our study was limited to

investigating the effects of global severity of hallucinations and delusions and could not resolve

effects of symptom subtype or content, since detailed assessments were only available in a small

subset of our patients. Larger studies with more detailed assessments are needed to tease out

these potential effects.

In conclusion, we have presented evidence for distinct hierarchical alterations in neural

timescales as a function of hallucination and delusion severity, lending initial neural support for

hierarchical views of psychosis. Additionally, our work suggests that INT (Watanabe et al., 2019)

provides a reliable and interpretable measure of neural function with potential to elucidate

hierarchical alterations and dysfunctions in circuit dynamics in schizophrenia and other

neuropsychiatric disorders.

Materials and Methods

Human Connectome Project Dataset

T1w/T2w maps and resting-state fMRI data were obtained for a subset of 100 unrelated young

and healthy subjects from the Human Connectome Project (HCP) WU-Minn Consortium (Van

Essen et al., 2013). The first fMRI run (single-shot EPI with left-to-right phase encoding direction)

from the first fMRI session were collected for each subject during an eyes-open-on-fixation

session with the following scanning parameters: repetition time (TR) = 720 ms; spatial resolution

= 2×2×2 mm; time points = 1200. High-resolution (0.7-mm isotropic voxels) T1w and T2w

anatomical images were also acquired. Details regarding subject recruitment and MRI data

acquisition have been previously reported (Smith et al., 2013; Van Essen et al., 2012).

Preprocessing of the HCP data was performed using the HCP minimal preprocessing pipeline

(Glasser et al., 2013). The preprocessed fMRI data were then used for estimation of INT maps in

32k Conte69 mesh surface space and in MNI152_ICBM2009a_nlin volume space with native

spatial resolution. The T1w/T2w (myelin) maps (Glasser and Van Essen, 2011) in 32k Conte69

mesh surface space were used to compare functional (INT) and structural (T1w/T2w) measures

of hierarchy.

Schizophrenia Combined Dataset

T1w images and resting-state fMRI data were obtained for 331 healthy control subjects and 254

patients diagnosed with either schizophrenia (N = 241) or schizoaffective disorder (N = 13) from

four publicly available datasets. Three of these datasets were from the SchizConnect repository

(BrainGluSchi (Bustillo et al., 2016), COBRE (Aine et al., 2017; Çetin et al., 2014), and

NMorphCH (Alpert et al., 2016)) and one was from the OpenfMRI repository (UCLA (Poldrack et

al., 2016)). Data that survived a quality-control check (~95%) and motion-censoring check (~64%)

included 140 patients and 225 controls. The quality control check consisted of visual inspection of

the spatially normalized images. The motion censoring check consisted of determining if there

were sufficient degrees of freedom after motion censoring to perform nuisance variable

regression. A subset of 158 controls was then selected that matched patients with regards to

gender and age. To minimize scanner- and site-related differences we excluded subjects if the

signal-to-noise ratio (SNR) was less than 100 for any of the standard regions-of-interest (Power

et al., 2011). The final sample after quality-control checks consisted of 127 patients and 152 age-

and gender-matched controls (**Table 1**).

The fMRI data were collected for each subject during an eyes-open-on-fixation session with

the following scanning parameters: TR = 2000 ms (except for NMorphCH, where TR = 2200 ms);

time points (BrainGluSchi/COBRE/NMorphCH/UCLA) = 160/145/318/147; spatial resolution (mm)

= 3.5×3.5×3.5/3.5×3.5×3.5/4×4×4/3×3×3. Data were preprocessed using the AFNI afni_proc.py

function. The following steps were performed: (1) removal of the first five volumes with the 3dTcat

function; (2) slice-timing correction; (3) motion correction; (4) 12-parameter affine registration of

the fMRI images to the T1w image; (5) spatial normalization of fMRI images to

MNI152_ICBM2009a_nlin volume space using nonlinear warping via the T1w image; (6) single-

interpolation resampling of fMRI images combining motion correction and spatial normalization.

Before estimating the voxelwise INT values, preprocessed fMRI data were further processed

with the following steps: (1) regression of white-matter signal, cerebrospinal-fluid signal, global-

brain signal, and the 6 motion parameters along with their first derivatives; (2) bandpass filtering

in the 0.01–0.1 Hz range; (3) motion censoring to remove volumes with framewise displacement

(FD)(Power et al., 2012) greater than 0.3 mm along with the volumes directly preceding and

following that volume; (4) spatial smoothing with a 4 mm full-width-at-half-maximum Gaussian

kernel. After INT estimation, the INT maps for subjects from the BrainGluSchi, COBRE, and

NMorphCH samples were resampled to a spatial resolution of $3\times3\times3$ mm to match the UCLA

sample.

Symptom severity in patients was assessed with the Positive and Negative Syndrome Scale

(PANSS) (Kay et al., 1987) in the COBRE and BrainGluSchi samples, or with the Scale for the

Assessment of Positive Symptoms (SAPS) (Andreasen, 1984) and the Scale for the Assessment

of Negative Symptoms (SANS) (Andreasen, 1983) in the UCLA and NMorphCH samples. To

appropriately combine the scores across all four samples, we chose the subset of 7 items that

constituted unequivocal matches between the PANSS and SAPS/SANS (in parentheses):

delusions (global rating of delusions), conceptual disorganization (global rating of positive formal

thought disorder), hallucinatory behavior (global rating of hallucinations), blunted affect (global

rating of affective flattening), emotional withdrawal (global rating of anhedonia/asociality),

passive/apathetic social withdrawal (global rating of avolition/apathy), lack of spontaneity and flow

of conversation (global rating of alogia). PANSS scores were decreased by 1 point for all levels of

severity and the severe and moderately severe levels were combined into a single level so that

scoring conformed to the SAPS/SANS scale (from 0 to 5 with increasing severity).

HCP Dataset Analysis

Based on previous work showing that lower T1w/T2w map values co-localize with higher

hierarchical levels (Burt et al., 2018), as do neural timescales (Chaudhuri et al., 2015; Murray et

al., 2014), we examined the spatial relationship between T1w/T2w, cortical thickness, and INT

values. We restricted this examination to the HCP dataset since its high-resolution and good-

quality structural MRI data allows for precise estimation of myelin maps. Group-averaged

T1w/T2w, cortical thickness, and INT maps in surface space were parcellated using the HCP-

multimodal parcellation (HCP-MMP1.0) (Glasser et al., 2016). The parcels were then separated

into either 6 parcel groups [the 22 sections described by Glasser et al. are divided into 6 parcel

groups: (1) visual (sections 1 – 5); (2) sensorimotor (sections 6 – 9); (3) auditory (sections 10 –

12); (4) remaining temporal cortex (sections 13 – 14); (5) remaining posterior cortex (sections 15

– 18); (6) remaining anterior cortex (sections 19 – 22) (Glasser et al., 2016)] or 12 networks (Ji et

al., 2019). We tested the parcel-wise spatial relationship between T1w/T2w or cortical thickness

and INT values using linear regression.

Findings from the parcel-wise analysis did not support a brain-wide, systematic alignment of

structural and functional hierarchies. This motivated us to determine anatomically informed

hierarchies within the sensory systems (auditory, visual, and somatosensory). Linear mixed-

effects models were used to determine the best-fitting hierarchical ordering for each system.

Models predicted hierarchical level from fixed- and random-effects (per subject) of T1w/T2w and

cortical thickness values for parcels ordered accordingly. Hierarchical orderings were first

determined for the sensory cortices. The four most likely orderings for each sensory system were

determined based on the primate anatomy literature (Felleman and Van, 1991; Galaburda and

Pandya, 1983; Hyvärinen and Poranen, 1978; Kaas and Hackett, 2000; Morel et al., 1993). The

auditory cortex regions were A1, lateral belt (LBelt), medial belt (MBelt), parabelt (PBelt),

retroinsular cortex (RI), A4, and A5. The positions of LBelt and MBelt were allowed to take either

level 2 or 3 of the hierarchy; PBelt and RI were allowed to take either level 4 or 5; A1 was level 1,

A4 was level 6, and A5 was level 7 in all cases. The visual regions were V1, V2, V3, V4, MT, V6,

and V7. The positions of V4 and MT were allowed to take either level 4 or 5 of the hierarchy; V6

and V7 were allowed to take either level 6 or 7; V1 was level 1, V2 was level 2, and V3 was level

3 in all cases. The somatosensory cortex regions were areas 3b, 3a, 1, 2, 5m, 7b, and 7a. Areas

3b and 3a were allowed to take either level 1 or 2 of the hierarchy; areas 1 and 2 were allowed to

take either level 3 or 4; area 5m was level 5, area 7b was level 6, and area 7a was level 7 in all

cases. Since all compared models had the same number of variables, the winning models for

each system were simply determined based on the orderings that explained the most variance

(R^2). Following determination of the hierarchies in the sensory cortices, 2 downstream prefrontal

cortex regions—areas 8a and 46—were added as either level 8 or 9 of the hierarchy based on a

second model comparison. The PFC-extended winning hierarchies were then validated by

determining the relationship of hierarchical levels with INT values using non-parametric

Spearman correlations (r_s) both in the HCP (in-sample) dataset and in the control group from the

schizophrenia combined (out-of-sample) dataset.

HCP Robustness Analyses

Because the schizophrenia combined dataset was analyzed in volume space, HCP dataset

group-averaged INT maps in volume space were created and parcellated into 180 cortical parcels

using HCP-MMP1.0 in volume space (https://identifiers.org/neurovault.collection:1549) and 8

FreeSurfer (Fischl, 2012) subcortical parcels relevant to schizophrenia (thalamus, caudate,

putamen, pallidum, hippocampus, amygdala, nucleus accumbens, and ventral diencephalon).

The reliability of INT maps was assessed at the voxel level in volume space using the two-way

random, single score intraclass correlation coefficient [ICC(2,1)] (Shrout and Fleiss, 1979). INT

maps for each of the 100 subjects estimated using the first 5 minutes of data acquisition (similar

to the amount of data available for the schizophrenia datasets) were compared to those

estimated using the last 5 minutes of data acquisition of a single 14-minute run. To evaluate

potential confounds of the INT values, we examined their relationship with age, gender, and head

motion—based on mean framewise displacement (Power et al., 2012) (FD)—using linear

regression

Schizophrenia Combined Dataset Analysis

INT maps were parcellated into 180 cortical parcels using HCP-MMP1.0 in volumetric space

(https://identifiers.org/neurovault.collection:1549) and 8 FreeSurfer (Fischl, 2012) subcortical

parcels (thalamus, caudate, putamen, pallidum, hippocampus, amygdala, nucleus accumbens,

and ventral diencephalon). In an exploratory analysis, differences in INT map values between

patients with schizophrenia and healthy controls were investigated using a linear-regression

model (M1$_{exploratory}$) predicting INT values as a function of diagnosis while controlling for age,

gender, mean FD, and sample-acquisition site (BrainGluSchi, COBRE, NMorphCH, and UCLA).

To test our hypothesis of hallucination- and delusion-specific alterations of INT, we evaluated the

relationships between symptom severity and INT values using a linear-regression model

(M1$_{primary}$) predicting INT with each of the 7 symptoms (hallucinations, delusions, conceptual

disorganization, emotional withdrawal, social withdrawal, blunted affect, and alogia) as regressors

while controlling for age, gender, mean FD, and sample-acquisition site. We did not use

voxelwise statistical parametric-mapping approaches because our main focus was on effects

along hierarchical gradients not necessarily dependent on anatomical proximity.

Permutation Testing

To assess statistical significance while controlling for multiple comparisons, we used permutation

tests, which provide adequate protection against false positives in fMRI analyses (Eklund et al.,

2016). Our main test focused on differences between hallucinations and delusions in INT gradient

effects within anatomically informed hierarchies of the auditory, visual, and somatosensory

systems—reflecting symptom-specific INT alterations at different hierarchical levels. We

specifically tested our primary hypothesis using a linear-regression model (M2) featuring

interactions of symptom-by-hierarchical-level within anatomical gradients in sensory systems. We

included a symptom-by-hierarchical-level-by-sensory-system interaction to allow for differences

between sensory systems. A post-hoc power analysis for M2 showed our analyses had between

88% and 99% power to detect effect sizes (Cohen's f^2) between 0.19 and 0.36 (α = 0.05).

Permutation tests compared observed effects (parcel-wise t-statistics of individual regression

coefficients from M2 [or M1$_{exploratory}$]) to those in a null distribution obtained from 10,000 surrogate

datasets in which the values of the predictor variables of interest in M1$_{primary}$ (or M1$_{exploratory}$) were

randomly shuffled. Corrected P-values at 0.05 ('$P_{permutation}$'), two-sided, are reported. Permutation

tests were also used to determine null distributions of the hierarchy model comparison for

determining the hierarchical orderings. There, null distributions were obtained from 10,000

surrogate datasets in which the assigned hierarchical level of each region was randomly

assigned.

Bootstrap Confidence Intervals

Bootstrap confidence intervals were determined for the results from M2 using the accelerated

bias-corrected (BC$_a$) percentile method (Efron, 1987). 10,000 bootstraps were performed at the

level of M1$_{primary}$ and two-sided 95% confidence intervals were determined.

Large-Scale Biophysical Model of Cortical Neural Timescales

We implemented the model introduced by Chaudhuri et al. (Chaudhuri et al., 2015), where "A

Large-Scale Circuit Mechanism for Hierarchical Dynamical Processing in the Primate Cortex" was

introduced. We chose this model because it was constructed using gold-standard tract-tracing

experiments to determine the directed- and weighted-connectivity strengths between nodes

(unlike similar models of the human cortex). Additionally, this model was previously shown to

recapitulate the observed hierarchy of intrinsic neural timescales. The model contains 29 nodes,

each consisting of an excitatory and inhibitory population. The populations are described by:

$$\tau_E \frac{d}{dt} v_E = -v_E + \beta_E [I_E]_+$$
$$\tau_I \frac{d}{dt} v_I = -v_I + \beta_I [I_I]_+$$

(1)

v_E is the firing rate of the excitatory population, with intrinsic time constant τ_E and input current I_E,

and for which the f-I curve has slope β_E. $[I_E]_+ = \max(I_E, 0)$. The inhibitory population has

corresponding parameters v_I, τ_I, I_I and β_I. Values for τ_E, τ_I, β_E, and β_I are given below and were

taken from Binzegger et al. (Binzegger et al., 2009).

At each node, the input currents have a component originating within the area (i.e. local

input) and another originating from other areas (i.e. long-range input):

$$I_E^i = (1 + \eta h_i)(w_{EE} v_E^i + I_{lr,E}^i) - w_{EI} v_I^i + I_{ext,E}^i$$

(2)

$$I_I^i = (1 + \eta h_i)(w_{IE} v_E^i + I_{lr,I}^i) - w_{II} v_I^i + I_{ext,I}^i$$

The super- and sub-script, i, denotes the node $(1 - 29)$, w_{EE} and w_{EI} are couplings to the

excitatory population from the local excitatory and inhibitory population respectively, $I_{lr,E}^i$ is the

long-range input to the excitatory population, and $I_{ext,E}^i$ is external input (both stimulus input and

any noise added to the system). w_{IE}, w_{II}, and $I_{lr,I}^i$ are the corresponding parameters for the

inhibitory population.

The excitatory inputs to an area, both local and long-range, are scaled by its position in the

hierarchy, h_i (see below for details). h_i is normalized between 0 and 1, and η is a scaling

parameter that controls the effect of hierarchy. By setting $\eta = 0$, the intrinsic differences between

areas are removed. Note that both local and long-range projections were scaled by hierarchy,

rather than just local projections, in accordance with the observations of Markov et al. (Markov et

al., 2010), who found that the proportion of local to long-range connections is approximately

conserved across areas.

Long-range input is modeled as excitatory current to both excitatory and inhibitory cells:

$$I_{lr,E}^i = \mu_{EE} \sum_{j=1}^{29} FLN_{ij} v_E^j$$

(3)

$$I_{lr,I}^i = \mu_{IE} \sum_{j=1}^{29} FLN_{ij} v_E^j$$

Here, j ranges over all areas. $I_{lr,E}^i$ and $I_{lr,I}^i$ are the inputs to the excitatory and inhibitory

populations, v_E^j is the firing rate of the excitatory population in area j and FLN_{ij} is the fraction of

labeled neurons (FLN; see below for details) from area j to area i. μ_{EE} and μ_{IE} are scaling

parameters that control the strengths of long-range input to the excitatory and inhibitory

populations, respectively, and do not vary between connections; all the specificity comes from the

FLN. Long-range connectivity is thus determined by three parameters: μ_{EE} and μ_{IE} control the

connection strengths of long-range projections, and η maps the hierarchy into excitatory

connection strengths. The excitatory to inhibitory ratio of an input current, $\gamma = I_{inp,E}/I_{inp,I}$, was

chosen such that the steady-state firing rate of the excitatory population does not change when

the current is present. Given input of $I_{inp,E}$ to the excitatory population, an input of $\gamma I_{inp,E}$ to the

inhibitory population increases the inhibitory firing rate sufficiently to cancel out the additional

input to the excitatory population. μ_{EE} and μ_{IE} were chosen with a ratio slightly above this value

so that projections are weakly excitatory.

The parameter values were: $\tau_E = 20$ ms, $\tau_I = 10$ ms, $\beta_E = 0.066$ Hz/pA, $\beta_I = 0.351$ Hz/pA,

$w_{EE} = 24.3$ pA/Hz, $w_{IE} = 12.2$ pA/Hz, $w_{EI} = 19.7$ pA/Hz, $w_{II} = 12.5$ pA/Hz, $\mu_{EE} = 33.7$ pA/Hz, $\mu_{IE} =$

25.3 pA/Hz and $\eta = 0.68$. Background input for each area was chosen so that the excitatory and

inhibitory populations had rates of 10 and 35 Hz, respectively. As in Chaudhuri et al., we added

an external input of white-noise to all areas with a mean of 0 Hz and a SD of 10^{-5} Hz.

Connectivity data are from an ongoing project that is quantitatively measuring all

connections between cortical areas in the macaque cortex (Markov et al., 2013; Markov et al.,

2012). The connection strengths between areas are measured by counting the number of

neurons labeled by retrograde tracer injections. To control for injection size, these counts are

normalized by the total number of neurons labeled in the injection, giving a fraction of labeled

neurons (FLN):

$$FLN_{j \to i} = \frac{\text{number of neurons projecting to area } i \text{ from area } j}{\text{total number of neurons projecting to area } i \text{ from all areas}} \quad (4)$$

9   These data were also used to estimate the fraction of neurons in a projection originating in the

10   supragranular layers (SLN):

$$SLN_{j \to i} = \frac{\text{number of supragranular neurons projecting to area } i \text{ from area } j}{\text{number of neurons projecting to area } i \text{ from area } j} \quad (5)$$

11   The hierarchy was constructed following a similar framework to Markov et al. (Markov et al.,

12   2014), using a generalized linear model. Hierarchical values were assigned to each area such

13   that the difference in values predicts SLN (Barone et al., 2000):

$$SLN_{j \to i} \approx g^{-1}\left(h_i - h_j\right) \quad (6)$$

where g^{-1} is a logistic function (logistic regression) and h_i is the hierarchy value of area i. In the

fit, the contribution of each projection is weighted by the log of its FLN to preferentially match

stronger and less noisy projections. These hierarchy values were previously shown to correlate

with spine counts (Chaudhuri et al., 2015), providing support for the choice of introducing a

gradient of excitatory input strength across the cortex in the model. All connectivity data can be

downloaded from www.core-nets.org.

The simulated neuronal activity was converted to blood-oxygen-level-dependent (BOLD)

fMRI signal using the Balloon-Windkessel hemodynamic model (Stephan et al., 2007); a

dynamical model that describes the transduction of neuronal activity (v_E) to changes in a

vasodilatory signal (s) that is subject to autoregulatory feedback. This vasodilatory signal is

coupled to changes in cerebral blood flow (f) that result in changes to the normalized total

deoxyhemoglobin content (q) and normalized venous blood volume (v). For each area (i), these

biophysical variables are defined by the following equations:

$$\frac{ds_i}{dt} = v_E^i - \kappa s_i - \gamma(f-1) \quad (7)$$

$$\frac{df_i}{dt} = s_i \quad (8)$$

$$\tau_{MTT}\frac{dv_i}{dt} = f_i - v_i^{1/\alpha} \quad (9)$$

$$\tau_{MTT} \frac{dq_i}{dt} = f_i \frac{1 - (1 - \rho)^{1/f_i}}{\rho} - v_i^{1/\alpha} \frac{q_i}{v_i} \tag{10}$$

1   where τ_{MTT} is the mean transit time of blood, ρ is the resting oxygen extraction fraction, and α

2   represents the resistance of the veins (i.e., stiffness). For each area (i), the BOLD signal (B), is a

3   static nonlinear function of deoxyhemoglobin content (q) and venous blood volume (v), that

4   comprises a volume weighted sum of extravascular and intravascular signals:

$$B_i = V_0 \left[k_1(1 - q_i) + k_2 \left(1 - \frac{q_i}{v_i} \right) + k_3(1 - v_i) \right]$$

$$k_1 = 4.3\vartheta_0 \rho TE$$

$$k_2 = \varepsilon r_0 \rho TE$$

$$k_3 = 1 - \varepsilon \tag{11}$$

where V_0 is the resting venous blood volume fraction, ϑ_0 is the frequency offset at the outer

surface of the magnetized vessel for fully deoxygenated blood, ε is the ratio of intra- and extra-

vascular signals, r_0 is the slope of the relation between the intravascular relaxation rate $R_{2I}*$ and

oxygen saturation, and TE is the echo time of the fMRI acquisition. All parameters for the Balloon-

Windkessel model were chosen according to Stephan et al. for a 3T fMRI experiment (Stephan et

al., 2007). Simulated BOLD signals were down sampled to a temporal resolution of 2 seconds

(i.e., TR = 2 seconds) to match the *in vivo* data and INTs were estimated in the same was as for

the *in vivo* data.

To simulate the observed effects of hallucination and delusion severity on INTs, we

perturbed the strength of the couplings to the excitatory population from the local excitatory

population (w_{EE}) or to the inhibitory population from the local excitatory population (w_{IE}) for

specific nodes. Note that in the definition of the model, all nodes had the same values for w_{EE} and

w_{IE} but here we are assigning node-specific values. We investigated alterations in excitation-

inhibition (E/I) ratios by allowing the strength of recurrent connections to vary in 5 of the 6 nodes

that correspond to levels of our hierarchy (V1, V2, V4, MT, 8I, and 46d, with the latter being fixed)

to recapitulate our *in vivo* observations. Recurrent connection strength was fixed for 46d to avoid

model instability upon small parameter changes (E/I ratio changes of ~1%) associated with the

strong connectivity at this level. The biophysical model parameters were fitted by minimizing the

sum of squared differences between the simulated ΔINT (differences of INT with respect to the

reference scenario, i.e., unaltered model) and the ΔINT based on the fitted data from M1primary

(differences of INT with respect to a scenario of no hallucinations or delusions, where the scores

for hallucination and delusion scores were set to 0 and all other variables to the group average) in

the 6 auditory parcels that correspond to the hierarchical levels of the biopsychical model nodes

(A1, LBelt, PBelt, RI, 8a, and 46). Three pathological scenarios were jointly fitted: (1)

hallucinations only (hallucination score set to 5, delusion score set to 0, all other variables set to

the group average); (2) delusions only (hallucination score set to 0, delusion score set to 5, all other variables set to the group average); and (3) hallucinations and delusions (hallucination and delusion scores set to 5, all other variables set to the group average).

Using a constrained fitting procedure, we fit the following 3 free parameters for each symptom (6 parameters total) to the 3 pathological scenarios simultaneously (6 data points corresponding to each node for a total of 18 data points). (1) Hierarchical level of the peak increase in E/I ratio, which was allowed to take any integer between 1 and 8. Given the stationary nature of these parameters, they were held constant such that fitting was performed for each combination of peak E/I ratio increase (1–8 for hallucinations and 1–8 for delusions). (2) Magnitude of the increase in peak local E/I ratio, expressed as percent change to the local recurrent connection strength, which was allowed to vary between 0% and 40%. (3) E/I ratio change at the minimum (most distant hierarchical level from the peak), which was allowed to vary between -30% and 40%. A linear gradient in E/I ratio change was assumed between the peak and the minimum that was symmetrical around the peak. Results are shown for reductions to w_{IE}, but similar effects can be observed when increasing w_{EE} since both effectively increase the E/I ratio.

Acknowledgments

The authors would like to thank Drs. Rishidev Chaudhuri and Xiao-Jing Wang for their guidance in implementing the large-scale biophysical model. This work was supported by the National Institute of Mental Health under awards R01MH117323 and R01MH114965. BrainGluSchi: data were downloaded from the COllaborative Informatics and Neuroimaging Suite Data Exchange tool (COINS;http://coins.mrn.org/dx) and data collection was funded by NIMH R01MH084898-01A1. COBRE: Data was downloaded from the COllaborative Informatics and Neuroimaging Suite Data Exchange tool (COINS; http://coins.mrn.org/dx), data collection was performed at the Mind Research Network, and funded by a Center of Biomedical Research Excellence (COBRE) grant 5P20RR021938/P20GM103472 from the NIH to Dr. Vince Calhoun. NMorphCH: data were obtained from the Neuromorphometry by Computer Algorithm Chicago (NMorphCH) dataset (http://nunda.northwestern.edu/nunda/data/projects/NMorphCH); the investigators within NMorphCH contributed to the design and implementation of NMorphCH and/or provided data but did not participate in analysis or writing of this report; data collection and sharing for this project was funded by NIMH grant R01MH056584. UCLA: data was obtained from the OpenfMRI database (its accession number is ds000030) and data collection was funded by the Consortium for Neuropsychiatric Phenomics (NIH Roadmap for Medical Research grants UL1-DE019580, RL1MH083268, RL1MH083269, RL1DA024853, RL1MH083270, RL1LM009833, PL1MH083271, and PL1NS062410). HCP: Data were provided by the Human Connectome Project, WU-Minn

1 Consortium (Principal Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657)

2 funded by the 16 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience

3 Research; and by the McDonnell Center for Systems Neuroscience at Washington University.

Figure Supplements

Figure 1—figure supplement 1. Comparison of Different Methods for INT Estimation: Sum of Initial Positive Period vs Exponential Fit. Relationship between neural timescales estimated using the sum of initial positive period (INT, x axis) and an exponential fit (τ, y axis) to the autocorrelation function. Results for vertices (left) and parcels (right). Lines of best-fit are shown for comparison. The INT method of Watanabe et al. (Watanabe et al., 2019) is similar to estimating the decay rate of an exponential fit to the autocorrelation function (Murray et al., 2014). The former method was used in our study to maintain consistency with the previous study, which validated the technique against INT measured by electroencephalography (Watanabe et al., 2019). Additionally, this method avoids the need for nonlinear fitting—which is known to be challenging (Transtrum et al., 2010) and computationally expensive,—and may be more robust. The difference between timescales estimated with the two methods were assessed in the 100 HCP subjects by comparing the group averaged timescale maps using Spearman correlation. The correlation between the two methods was almost perfect (vertices: r_s = 0.9986, P << 0.01; parcels: r_s = 0.9985, P << 0.01) but there is an upwards bias in values estimated using the method of Watanabe et al. (Watanabe et al., 2019).

Figure 1—figure supplement 2. Relationships Between Intrinsic Neural Timescales and either T1w/T2w (Myelin) or Cortical Thickness. As reported in the **Main Text**, we did not observe a clear parcel-wide relationship between structural markers of hierarchy and INT. These relationships were investigated using two parcel-groupings; those described by Glasser et al. (left) and the Cole-Anticevic networks described by Jing et al. (right). To better characterize the relationship between myelin (top) or cortical thickness (bottom) and INT, intercepts for each parcel group (visual, auditory, sensorimotor, posterior, anterior, and temporal) or network (primary visual, secondary visual, somatomotor, cingulo-opercular, dorsal attention, language, frontoparietal, auditory, default mode, posterior multimodal, ventral multimodal, and orbito-affective) were introduced to allow for overall differences in INT-by-parcel-group (or INT-by-network) given the apparent differences observed in the data. Parcel-group-by-T1w/T2w-value (or network-by-T1w/T2w-value) interactions and parcel-group-by-cortical-thickness (or network-by-cortical-thickness) were also introduced to allow for differences in the relationship between

T1w/T2w (or cortical thickness) and INT values across parcel groups (or networks).

We found that an extended model with parcel-group intercepts and interactions of parcel-

group-by-T1w/T2w explained significantly more variance in INT than the reduced model (no

parcel-group intercepts or interactions): parcel groups differed in INT (model with parcel-group

intercepts versus reduced model: $F_{10,167}$ = 12.03, P < 10^{-14}) and their relationship between

T1w/T2w and INT (model with parcel-group intercepts and interactions of parcel-group-by-

T1w/T2w versus model with parcel-group intercepts but no interaction terms: $F_{5,167}$ = 5.55, P < 10^{-}

^{4}). Similar results were observed for the Cole-Anticevic Networks, where we found that an

extended model with network intercepts and interactions of network-by-T1w/T2w explained

significantly more variance in INT than the reduced model (no network intercepts or interactions):

networks differed in INT (model with network intercepts versus reduced model: $F_{22,155}$ = 4.62, P <

10^{-8}) and their relationship between T1w/T2w and INT (model with network intercepts and

interactions of network-by-T1w/T2w versus model with network intercepts but no interactions:

$F_{11,155}$ = 2.15, P = 0.020).

For cortical thickness, we found that an extended model with parcel-group intercepts and

interactions of parcel-group-by-cortical-thickness explained significantly more variance in INT

than the reduced model (no parcel-group intercepts or interactions): parcel groups differed in INT

(model with parcel-group intercepts versus reduced model: $F_{10,167}$ = 8.59, P < 10^{-10}) and their

relationship between cortical-thickness and INT (model with parcel-group intercepts and

interactions of parcel-group-by-cortical-thickness versus model with parcel-group intercepts but

no interaction terms: $F_{5,167}$ = 5.05, P < 10^{-3}). Similar results were observed for the Cole-Anticevic

Networks, where we found that an extended model with network intercepts and interactions of

network-by-cortical-thickness explained significantly more variance in INT than the reduced

model (no network intercepts or interactions): networks differed in INT (model with network

intercepts versus reduced model: $F_{22,155}$ = 3.58, P < 10^{-5}) and their relationship between cortical-

thickness and INT (model with network intercepts and interactions of network-by-cortical-

thickness versus model with network intercepts but no interactions: $F_{11,155}$ = 1.96, P = 0.036).

These results suggest that the relationship between structural hierarchies (T1w/T2w and

cortical thickness) and functional hierarchy (INT) is not constant across the whole brain but rather

changes for different parcel groups and networks.

Figure 1—figure supplement 3. Effect of framewise displacement (FD) on estimated

intrinsic neural timescales (INT). A) t-statistic values from a regression model predicting INT

from the average FD during fMRI data acquisition plotted as a function of the INT rank for 100

HCP subjects. Each datapoint represents one parcel. 181 out of 188 parcels showed significantly

lower INT with greater FD ($P_{permutation}$ = 0.01). Although there was a significant effect of FD on INT

throughout the whole brain, this effect was not modulated by order within the INT hierarchy (i.e.

the correlation in **A** is not significant against the null distribution). **B)** t-statistic values from parcels

within the well-defined auditory hierarchy (left) and visual hierarchy (right) plotted as a function of

hierarchical level. The effect of FD on estimated INT was not modulated by position within well-

defined sensory hierarchies across sensory systems (i.e. no significant difference between the

correlations in **B** tested against the null distribution). To determine the effect of head motion

during resting-state fMRI data acquisition on estimated INT values, we used a linear-regression

model to predict INT as a function of average FD while controlling for age and gender. Because

average FD represents a summary measure of head motion during the entire acquisition, this

regression does not provide information regarding differential effects of sustained small-levels of

head motion versus infrequent but large-levels of head motion.

Figure 1—figure supplement 4. Reliability of Intrinsic Neural Timescale Estimation. INT

maps for 100 HCP subjects were estimated in volume space and the values at each voxel falling

within the 188 volume parcels were compared when INT was estimated from the first 5 minutes or

last 5 minutes of fMRI acquisition. The INT maps showed excellent reliability based on the

ICC(2,1) between the first and last 5 minutes of the fMRI acquisition (median ICC $\pm$ interquartile

range across voxels: 0.94 $\pm$ 0.03).

Figure 1—figure supplement 5. Model comparison to determine the hierarchical ordering of

the ventral auditory pathway. **A)** Goodness-of-fit (R^2) of linear mixed-effects models predicting

different hierarchical orderings for the ventral auditory system from T1w/T2w and cortical thickness

values in the HCP dataset. First, the winning ordering (i.e., the model with the best goodness of fit)

for each system was determined for the 7 sensory cortex regions (bottom 4 models); then, winning

orderings were determined for extended models with 2 downstream prefrontal cortex regions added

to the respective winning models for sensory cortex (top 2 models). Note that, for each of the four

considered orderings within the sensory cortex for each system, only the 4 regions whose order is

varied (out of 7 regions) are shown to denote the models. For the auditory cortex, A1 was always

the lowest order region while A4 and A5 were always the two highest order regions. Null

distributions were generated by randomly permuting the hierarchical ordering (0^{th} – 95^{th} percentiles

shown). **B)** Scatterplots showing INT values plotted as a function of hierarchical level for the PFC-

extended winning model in **B** (red outline) for the HCP dataset (top) and the healthy control group

in the schizophrenia combined dataset (*N* = 158; bottom).

Figure 2—figure supplement 1. Voxel-Wise Analysis of Intrinsic Neural Timescales in

Schizophrenia versus Health. Patients with schizophrenia show widespread voxelwise

reductions of intrinsic neural timescales compared to healthy controls. t-statistic values from a

regression model predicting INT controlling for age, gender, framewise displacement and data

collection site, plotted as a function of the INT rank. Each datapoint represents one voxel. We

repeated the exploratory analysis of INT in patients with schizophrenia versus healthy controls at

the level of individual voxels. Only voxels that fell within the 188 parcels were included in this

analysis. Similar results were observed as reported in the **Main Text (Figure 3E)** where we

investigated effects at the level of parcels. Relative to controls, patients with schizophrenia

exhibited a small-to-moderate but widespread reduction of INT (overall effect of diagnosis: 5,511

out of 27,884, $P_{permutation}$ = 0.02;). A non-significant number of voxels also showed significantly

higher INT in patients with schizophrenia compared to healthy controls (93 out of 27,884 voxels,

$P_{permutation}$ = 0.49).

Figure 2—figure supplement 2. Biophysical Model Simulation of Reduced Intrinsic Neural

Timescales in Schizophrenia versus Health. An *in silico* simulation using a large-scale

biophysical model suggested that the globally lower INT observed in patients with schizophrenia

could be neuronally implemented by globally reduced E/I ratio. A scatterplot shows the difference

of intrinsic neural timescale (INT) values estimated for the reference (healthy) simulated scenario

and the globally reduced excitatory-to-excitatory recurrent connection strength (reduced E/I ratio);

each data point denotes 1 of the 29 nodes (top). Relative to the reference scenario, reduced

excitatory-to-excitatory recurrent connection strength leads to a global reduction of INT values

with the effect becoming more pronounced for regions with longer timescales in the healthy

condition. The data are well described by an exponential model (R^2 = 0.94). A scatterplot shows

the t-statistic values from a regression model predicting INT group differences controlling for age,

gender, framewise displacement and data collection site, plotted as a function of the INT value in

healthy subjects (determined from group-averaged INT map from HCP subjects to reduce

circularity); each data point denotes the average of 6 parcels along the INT hierarchy for a similar

number of data points as the biophysical model results. Similar to the biophysical model

simulation, the data are better fit by an exponential model (Matlab *fitnlm:* $F_{2,28}$ = 10.3, P = 0.003;

BIC = 4.8), than a linear model (Matlab *fitnlm:* $F_{1,29}$ = 1.4, P = 0.25; BIC = 9.3). Similar results are

observed when no binning of the data is performed (exponential model: $F_{2,185}$ = 8.3, P = 0.005;

BIC = 366.4; linear model: $F_{1,186}$ = 0.8, P = 0.38; BIC = 368.6). The findings are similar to a recent

study that recapitulated behavior of macaque monkeys on an evidence-varying decision-making

task following intramuscular administration of a subanesthetic dose of ketamine by reducing the

strength of the couplings to the excitatory population from the local excitatory population (w_{EE}) by

1.75% to achieve a global reduction of E/I ratio. This is consistent with the NMDA-receptor

hypofunction hypothesis of schizophrenia (Corlett et al., 2011; Kehrer et al., 2008; Krystal et al.,

2003; Lisman et al., 2008; Olney and Farber, 1995), and the use of ketamine as a model of

schizophrenia (Becker et al., 2003; Corlett et al., 2011; Frohlich and Van Horn, 2014; Krystal et

al., 1994). Furthermore, our biophysical model results support recent evidence of how global

changes can preferential effect higher order brain regions (Yang et al., 2016); and our in vivo

results lend additional—although preliminary—evidence.

Figure 3—figure supplement 1. Controlling for medication dose does not change the

distinct hierarchical-gradient effects of hallucination and delusion severity in the auditory

and somatosensory systems. A) Scatterplots showing *t*-statistic values for hallucination-

severity (top) or delusion-severity (bottom) effects from parcels within the auditory (left), visual

(middle), or somatosensory (right) hierarchies plotted as a function of hierarchical level. **B)**

Summary of results from a model including symptom-severity effect (hallucination or delusion),

hierarchical level, sensory system (auditory, visual, or somatosensory), and their interactions. We

repeated the main analysis comparing the effects of hallucination and delusion severity when

including medication dose as a covariate. Only 109 of the 127 patients had available data for

chlorpromazine equivalents. All results were similar to those reported in the **Main Text** where the

effect of medication dose was not controlled for. The model explaining symptom effects and their

differences by hierarchical-level and their interaction by symptoms and sensory system was

significant (omnibus $F_{11,41}$ = 22.4, $P < 10^{-13}$). Critically, we found significant hierarchical-gradient

effects that differed between hallucinations and delusions (auditory system symptom-by-

hierarchical-level interaction: t_{42} = 4.48, $P_{permutation}$ = 0.002; visual: t_{42} = -0.88, $P_{permutation}$ = 0.560;

and somatosensory: t_{42} = 3.92, $P_{permutation}$ = 0.001). These interactions were in the expected

direction for 2/3 of the sensory systems auditory and somatosensory; negative hierarchical

gradient for hallucinations and positive hierarchical gradient for delusions). For the auditory

system, this effect was driven by significant hierarchical-gradient effects in opposite directions for

hallucinations (hierarchical-level effect: t_{42} = -3.40, $P_{permutation}$ = 0.013) and delusions (hierarchical-

level effect: t_{42} = 2.94, $P_{permutation}$ = 0.028). For the somatosensory system, this effect was driven

by a trend-level negative hierarchical-gradient effect for hallucinations (hierarchical-level effect: t_{42}

= -2.25, $P_{permutation}$ = 0.064) and a significant negative hierarchical-gradient effect for delusions

(hierarchical-level effect: t_{42} = 3.30, $P_{permutation}$ = 0.017).

Figure 3—figure supplement 2. Only positive symptoms show hierarchical-gradient effects in the auditory system and only delusions show a hierarchical-gradient effect in the somatosensory system. *t*-statistic values for the hierarchical-gradient effect of each symptom in the auditory and somatosensory systems. In the auditory system, all three positive symptoms (including conceptual disorganization) show significant effects while none of the negative symptoms show significant effects. In the somatosensory system, only delusions show a significant effect, although hallucinations show the strongest negative effect. These results reflect some level of selectivity that should be examined in more detail in future studies. It is important to note that the perceptual inference model of psychosis does not require these effects to be specific to hallucinations and delusions. Furthermore, an effect of conceptual disorganization—a positive symptom that unlike negative symptoms tends to correlate with hallucinations and delusions—may suggest extensions of the model to account for additional phenomena.

Figure 3—figure supplement 3. Comparison of hierarchical effects in the dorsal and ventral auditory systems. A) Scatterplots showing *t*-statistic values for hallucination-severity (top) or delusion-severity (bottom) effects from parcels within the dorsal auditory (left) and ventral auditory (right) hierarchies plotted as a function of hierarchical level. **B)** Summary of results from a model including symptom-severity effect (hallucination or delusion), hierarchical level, sensory system (dorsal or ventral auditory), and their interactions. These results demonstrate: (1) in both the dorsal and ventral auditory systems, there is a significant hierarchical effect of hallucination severity and these effects are not significantly different from another (a); (2) there is no significant difference in the relationship between hallucination severity and hierarchical level versus that for delusion severity and hierarchical level in the ventral auditory system (b); (3) there is no significant hierarchical effect of delusion severity in the ventral auditory system. These findings add to previous evidence suggesting that the dorsolateral prefrontal cortex (area 46; include in dorsal but not ventral auditory system) may play a key role in delusions.

Figure 3—figure supplement 4. Distinct hierarchical-gradient effects of hallucination and delusion severity in the auditory system are observed when using an anatomically agnostic definition of the hierarchy. A) Scatterplots showing *t*-statistic values for hallucination-severity (left) or delusion-severity (right) effects from bins consisting of the average from 60 voxels within the 9 auditory system parcels (600 voxels total) as a function of hierarchical level. Voxels where first ranked using their corresponding intrinsic neural timescale (INT) value in the HCP dataset group-averaged INT map and then averaged within 10 equally spaced bins. Similar

to our main analysis results, we observe a significant symptom-by-hierarchical-level interaction, a

significant negative hierarchical-gradient effect for hallucinations and a significant positive

hierarchical-gradient effect for delusions.

References

Adams, R.A., Napier, G., Roiser, J.P., Mathys, C., Gilleen, J., 2018. Attractor-like dynamics in
belief updating in schizophrenia. Journal of neuroscience 38, 9471-9485.

Adams, R.A., Stephan, K.E., Brown, H.R., Frith, C.D., Friston, K.J., 2013. The computational
anatomy of psychosis. Frontiers in psychiatry 4, 47.

Aine, C., Bockholt, H.J., Bustillo, J.R., Cañive, J.M., Caprihan, A., Gasparovic, C., Hanlon, F.M.,
Houck, J.M., Jung, R.E., Lauriello, J., 2017. Multimodal neuroimaging in schizophrenia:
description and dissemination. Neuroinformatics 15, 343-364.

Alpert, K., Kogan, A., Parrish, T., Marcus, D., Wang, L., 2016. The northwestern university
neuroimaging data archive (NUNDA). NeuroImage 124, 1131-1136.

Andreasen, N., 1983. The scale for the assessment of negative symptoms (SANS) Iowa City. IA:
University of Iowa.

Andreasen, N.C., 1984. Scale for the Assessment of Positive Symptons:(SAPS). University of
Iowa.

Association, A.P., 2013. Diagnostic and statistical manual of mental disorders (DSM-5®).
American Psychiatric Pub.

Baker, S.C., Konova, A.B., Daw, N.D., Horga, G., 2019. A distinct inferential mechanism for
delusions in schizophrenia. Brain 142, 1797-1812.

Barone, P., Batardiere, A., Knoblauch, K., Kennedy, H., 2000. Laminar distribution of neurons in
extrastriate areas projecting to visual areas V1 and V4 correlates with the hierarchical rank and
indicates the operation of a distance rule. Journal of neuroscience 20, 3263-3281.

Bassett, D.S., Bullmore, E., Verchinski, B.A., Mattay, V.S., Weinberger, D.R., Meyer-Lindenberg,
A., 2008. Hierarchical organization of human cortical networks in health and schizophrenia.
Journal of neuroscience 28, 9239-9248.

Becker, A., Peters, B., Schroeder, H., Mann, T., Huether, G., Grecksch, G., 2003. Ketamine-
induced changes in rat behaviour: a possible animal model of schizophrenia. Progress in Neuro-
Psychopharmacology and Biological Psychiatry 27, 687-700.

Binzegger, T., Douglas, R.J., Martin, K.A., 2009. Topology and dynamics of the canonical circuit
of cat V1. Neural Networks 22, 1071-1078.

Brodmann, K., 1909. Vergleichende Lokalisationslehre der Grosshirnrinde in ihren Prinzipien
dargestellt auf Grund des Zellenbaues. Barth.

Brüne, M., Lissek, S., Fuchs, N., Witthaus, H., Peters, S., Nicolas, V., Juckel, G., Tegenthoff, M.,
2008. An fMRI study of theory of mind in schizophrenic patients with "passivity" symptoms.
Neuropsychologia 46, 1992-2001.

Brunel, N., Wang, X.-J., 2001. Effects of neuromodulation in a cortical network model of object
working memory dominated by recurrent inhibition. Journal of computational neuroscience 11, 63-
85.

Burt, J.B., Demirtaş, M., Eckner, W.J., Navejar, N.M., Ji, J.L., Martin, W.J., Bernacchia, A.,
Anticevic, A., Murray, J.D., 2018. Hierarchy of transcriptomic specialization across human cortex
captured by structural neuroimaging topography. Nature neuroscience 21, 1251.

Bustillo, J.R., Jones, T., Chen, H., Lemke, N., Abbott, C., Qualls, C., Stromberg, S., Canive, J.,
Gasparovic, C., 2016. Glutamatergic and neuronal dysfunction in gray and white matter: a
spectroscopic imaging study in a large schizophrenia sample. Schizophrenia Bulletin 43, 611-
619.

Butler, P.D., Silverstein, S.M., Dakin, S.C., 2008. Visual perception and its impairment in
schizophrenia. Biological psychiatry 64, 40-47.

Cassidy, C.M., Balsam, P.D., Weinstein, J.J., Rosengard, R.J., Slifstein, M., Daw, N.D., Abi-
Dargham, A., Horga, G., 2018. A perceptual inference mechanism for hallucinations linked to
striatal dopamine. Current Biology 28, 503-514. e504.

Cavanagh, S.E., Lam, N.H., Murray, J.D., Hunt, L.T., Kennerley, S.W., 2019. A circuit mechanism
for irrationalities in decision-making and NMDA receptor hypofunction: behaviour, computational
modelling, and pharmacology. bioRxiv, 826214.

Çavuş, I., Reinhart, R.M., Roach, B.J., Gueorguieva, R., Teyler, T.J., Clapp, W.C., Ford, J.M.,
Krystal, J.H., Mathalon, D.H., 2012. Impaired visual cortical plasticity in schizophrenia. Biological
psychiatry 71, 512-520.

Çetin, M.S., Christensen, F., Abbott, C.C., Stephen, J.M., Mayer, A.R., Cañive, J.M., Bustillo,
J.R., Pearlson, G.D., Calhoun, V.D., 2014. Thalamus and posterior temporal lobe show greater
inter-network connectivity at rest and across sensory paradigms in schizophrenia. NeuroImage
97, 117-126.

Chaudhuri, R., Knoblauch, K., Gariel, M.-A., Kennedy, H., Wang, X.-J., 2015. A large-scale circuit
mechanism for hierarchical dynamical processing in the primate cortex. Neuron 88, 419-431.

Corlett, P.R., Frith, C.D., Fletcher, P.C., 2009a. From drugs to deprivation: a Bayesian framework
for understanding models of psychosis. Psychopharmacology 206, 515-530.

Corlett, P.R., Honey, G.D., Krystal, J.H., Fletcher, P.C., 2011. Glutamatergic model psychoses:
prediction error, learning, and inference. Neuropsychopharmacology 36, 294.

Corlett, P.R., Horga, G., Fletcher, P.C., Alderson-Day, B., Schmack, K., Powers III, A.R., 2018.
Hallucinations and strong priors. Trends in cognitive sciences.

Corlett, P.R., Krystal, J.H., Taylor, J.R., Fletcher, P.C., 2009b. Why do delusions persist?
Frontiers in human neuroscience 3, 12.

Corlett, P.R., Murray, G.K., Honey, G.D., Aitken, M.R., Shanks, D.R., Robbins, T.W., Bullmore, E.T., Dickinson, A., Fletcher, P.C., 2007. Disrupted prediction-error signal in psychosis: evidence for an associative account of delusions. Brain 130, 2387-2400.

Davies, D.J., Teufel, C., Fletcher, P.C., 2017. Anomalous perceptions and beliefs are associated with shifts toward different types of prior knowledge in perceptual inference. Schizophrenia Bulletin 44, 1245-1253.

Dayan, P., Hinton, G.E., Neal, R.M., Zemel, R.S., 1995. The helmholtz machine. Neural computation 7, 889-904.

Ding, N., Melloni, L., Zhang, H., Tian, X., Poeppel, D., 2016. Cortical tracking of hierarchical linguistic structures in connected speech. Nature neuroscience 19, 158.

Dondé, C., Silipo, G., Dias, E.C., Javitt, D.C., 2019. Hierarchical deficits in auditory information processing in schizophrenia. Schizophrenia research 206, 135-141.

Dorph-Petersen, K.A., Pierri, J.N., Wu, Q., Sampson, A.R., Lewis, D.A., 2007. Primary visual cortex volume and total neuron number are reduced in schizophrenia. Journal of Comparative Neurology 501, 290-301.

Durstewitz, D., Seamans, J.K., 2002. The computational role of dopamine D1 receptors in working memory. Neural Networks 15, 561-572.

Efron, B., 1987. Better bootstrap confidence intervals. Journal of the American statistical Association 82, 171-185.

Eklund, A., Nichols, T.E., Knutsson, H., 2016. Cluster failure: Why fMRI inferences for spatial extent have inflated false-positive rates. Proceedings of the National Academy of Sciences 113, 7900-7905.

Felleman, D.J., Van, D.E., 1991. Distributed hierarchical processing in the primate cerebral cortex. Cerebral cortex (New York, NY: 1991) 1, 1-47.

Fischl, B., 2012. FreeSurfer. NeuroImage 62, 774-781.

Fischl, B., Rajendran, N., Busa, E., Augustinack, J., Hinds, O., Yeo, B.T., Mohlberg, H., Amunts, K., Zilles, K., 2008. Cortical folding patterns and predicting cytoarchitecture. Cerebral cortex 18, 1973-1980.

Fletcher, P.C., Frith, C.D., 2009. Perceiving is believing: a Bayesian approach to explaining the positive symptoms of schizophrenia. Nature Reviews Neuroscience 10, 48.

Foss-Feig, J.H., Adkinson, B.D., Ji, J.L., Yang, G., Srihari, V.H., McPartland, J.C., Krystal, J.H., Murray, J.D., Anticevic, A., 2017. Searching for cross-diagnostic convergence: neural mechanisms governing excitation and inhibition balance in schizophrenia and autism spectrum disorders. Biological psychiatry 81, 848-861.

Friston, K., Kiebel, S., 2009. Predictive coding under the free-energy principle. Philosophical Transactions of the Royal Society B: Biological Sciences 364, 1211-1221.

Frohlich, J., Van Horn, J.D., 2014. Reviewing the ketamine model for schizophrenia. Journal of psychopharmacology 28, 287-302.

Galaburda, A.M., Pandya, D.N., 1983. The intrinsic architectonic and connectional organization of the superior temporal region of the rhesus monkey. Journal of Comparative Neurology 221, 169-184.

Glasser, M.F., Coalson, T.S., Robinson, E.C., Hacker, C.D., Harwell, J., Yacoub, E., Ugurbil, K., Andersson, J., Beckmann, C.F., Jenkinson, M., 2016. A multi-modal parcellation of human cerebral cortex. Nature 536, 171.

Glasser, M.F., Sotiropoulos, S.N., Wilson, J.A., Coalson, T.S., Fischl, B., Andersson, J.L., Xu, J., Jbabdi, S., Webster, M., Polimeni, J.R., 2013. The minimal preprocessing pipelines for the Human Connectome Project. NeuroImage 80, 105-124.

Glasser, M.F., Van Essen, D.C., 2011. Mapping human cortical areas in vivo based on myelin content as revealed by T1-and T2-weighted MRI. Journal of neuroscience 31, 11597-11616.

Glaze, C.M., Kable, J.W., Gold, J.I., 2015. Normative evidence accumulation in unpredictable environments. Elife 4, e08825.

Gollo, L.L., 2019. Computational Psychiatry: Exploring atypical timescales in the brain. Elife 8, e45089.

Hasson, U., Chen, J., Honey, C.J., 2015. Hierarchical process memory: memory as an integral component of information processing. Trends in cognitive sciences 19, 304-313.

Hasson, U., Yang, E., Vallines, I., Heeger, D.J., Rubin, N., 2008. A hierarchy of temporal receptive windows in human cortex. Journal of neuroscience 28, 2539-2550.

Hilgetag, C.-C., Van Essen, D.C., O'Neill, M.A., Young, M.P., Felleman, D.J., 1996. Indeterminate organization of the visual systemon hierarchies: Response to hilgetag et al. rejoinder: Further commentary: Determinate or indeterminate organization. Science 271, 776-776.

Honey, C.J., Thesen, T., Donner, T.H., Silbert, L.J., Carlson, C.E., Devinsky, O., Doyle, W.K., Rubin, N., Heeger, D.J., Hasson, U., 2012. Slow cortical dynamics and the accumulation of information over long timescales. Neuron 76, 423-434.

Horga, G., Abi-Dargham, A., 2019. An integrative framework for perceptual disturbances in psychosis. Nature Reviews Neuroscience, 1-16.

Hyvärinen, J., Poranen, A., 1978. Receptive field integration and submodality convergence in the hand area of the post-central gyrus of the alert monkey. The Journal of physiology 283, 539-556.

Jardri, R., Hugdahl, K., Hughes, M., Brunelin, J., Waters, F., Alderson-Day, B., Smailes, D., Sterzer, P., Corlett, P.R., Leptourgos, P., 2016. Are hallucinations due to an imbalance between excitatory and inhibitory influences on the brain? Schizophrenia Bulletin 42, 1124-1134.

Javitt, D.C., Sweet, R.A., 2015. Auditory dysfunction in schizophrenia: integrating clinical and basic features. Nature Reviews Neuroscience 16, 535-550.

Javitt, D.C., Zukin, S.R., Heresco-Levy, U., Umbricht, D., 2012. Has an angel shown the way? Etiological and therapeutic implications of the PCP/NMDA model of schizophrenia. Schizophrenia Bulletin 38, 958-966.

Ji, J.L., Spronk, M., Kulkarni, K., Repovš, G., Anticevic, A., Cole, M.W., 2019. Mapping the
human brain's cortical-subcortical functional network organization. NeuroImage 185, 35-57.

Kaas, J.H., Hackett, T.A., 2000. Subdivisions of auditory cortex and processing streams in
primates. Proceedings of the National Academy of Sciences 97, 11793-11799.

Kay, S.R., Fiszbein, A., Opler, L.A., 1987. The positive and negative syndrome scale (PANSS) for
schizophrenia. Schizophrenia Bulletin 13, 261-276.

Kehrer, C., Maziashvili, N., Dugladze, T., Gloveli, T., 2008. Altered excitatory-inhibitory balance in
the NMDA-hypofunction model of schizophrenia. Frontiers in molecular neuroscience 1, 6.

Kiebel, S.J., Daunizeau, J., Friston, K.J., 2008. A hierarchy of time-scales and the brain. PLoS
computational biology 4, e1000209.

Kiebel, S.J., Daunizeau, J., Friston, K.J., 2009. Perception and hierarchical dynamics. Frontiers in
neuroinformatics 3, 20.

Krystal, J.H., D'Souza, D.C., Mathalon, D., Perry, E., Belger, A., Hoffman, R., 2003. NMDA
receptor antagonist effects, cortical glutamatergic function, and schizophrenia: toward a paradigm
shift in medication development. Psychopharmacology 169, 215-233.

Krystal, J.H., Karper, L.P., Seibyl, J.P., Freeman, G.K., Delaney, R., Bremner, J.D., Heninger,
G.R., Bowers, M.B., Charney, D.S., 1994. Subanesthetic effects of the noncompetitive NMDA
antagonist, ketamine, in humans: psychotomimetic, perceptual, cognitive, and neuroendocrine
responses. Archives of general psychiatry 51, 199-214.

Lam, N.H., Borduqui, T., Hallak, J., Roque, A.C., Anticevic, A., Krystal, J.H., Wang, X.-J., Murray,
J.D., 2017. Effects of altered excitation-inhibition balance on decision making in a cortical circuit
model. bioRxiv, 100347.

Lawson, R.P., Mathys, C., Rees, G., 2017. Adults with autism overestimate the volatility of the
sensory environment. Nature neuroscience 20, 1293.

Lee, T.S., Mumford, D., 2003. Hierarchical Bayesian inference in the visual cortex. JOSA A 20,
1434-1448.

Leitman, D.I., Sehatpour, P., Higgins, B.A., Foxe, J.J., Silipo, G., Javitt, D.C., 2010. Sensory
deficits and distributed hierarchical dysfunction in schizophrenia. American Journal of Psychiatry
167, 818-827.

Lerner, Y., Honey, C.J., Silbert, L.J., Hasson, U., 2011. Topographic mapping of a hierarchy of
temporal receptive windows using a narrated story. Journal of neuroscience 31, 2906-2915.

Lewandowski, K.E., DePaola, J., Camsari, G.B., Cohen, B.M., Ongur, D., 2009. Tactile, olfactory,
and gustatory hallucinations in psychotic disorders: a descriptive study. Ann Acad Med Singapore
38, 383-385.

Lim, A., Hoek, H.W., Deen, M.L., Blom, J.D., Bruggeman, R., Cahn, W., de Haan, L., Kahn, R.S.,
Meijer, C.J., Myin-Germeys, I., 2016. Prevalence and classification of hallucinations in multiple
sensory modalities in schizophrenia spectrum disorders. Schizophrenia research 176, 493-499.

Lisman, J.E., Coyle, J.T., Green, R.W., Javitt, D.C., Benes, F.M., Heckers, S., Grace, A.A., 2008.
Circuit-based framework for understanding neurotransmitter and risk gene interactions in
schizophrenia. Trends in neurosciences 31, 234-242.

Major, G., Tank, D., 2004. Persistent neural activity: prevalence and mechanisms. Current
opinion in neurobiology 14, 675-684.

Mante, V., Sussillo, D., Shenoy, K.V., Newsome, W.T., 2013. Context-dependent computation by
recurrent dynamics in prefrontal cortex. Nature 503, 78.

Markov, N., Misery, P., Falchier, A., Lamy, C., Vezoli, J., Quilodran, R., Gariel, M., Giroud, P.,
Ercsey-Ravasz, M., Pilaz, L., 2010. Weight consistency specifies regularities of macaque cortical
networks. Cerebral cortex 21, 1254-1272.

Markov, N.T., Ercsey-Ravasz, M., Lamy, C., Gomes, A.R.R., Magrou, L., Misery, P., Giroud, P.,
Barone, P., Dehay, C., Toroczkai, Z., 2013. The role of long-range connections on the specificity
of the macaque interareal cortical network. Proceedings of the National Academy of Sciences
110, 5187-5192.

Markov, N.T., Ercsey-Ravasz, M., Ribeiro Gomes, A., Lamy, C., Magrou, L., Vezoli, J., Misery, P.,
Falchier, A., Quilodran, R., Gariel, M., 2012. A weighted and directed interareal connectivity
matrix for macaque cerebral cortex. Cerebral cortex 24, 17-36.

Markov, N.T., Vezoli, J., Chameau, P., Falchier, A., Quilodran, R., Huissoud, C., Lamy, C.,
Misery, P., Giroud, P., Ullman, S., 2014. Anatomy of hierarchy: feedforward and feedback
pathways in macaque visual cortex. Journal of Comparative Neurology 522, 225-259.

Mazurek, M.E., Roitman, J.D., Ditterich, J., Shadlen, M.N., 2003. A role for neural integrators in
perceptual decision making. Cerebral cortex 13, 1257-1269.

Morel, A., Garraghty, P., Kaas, J., 1993. Tonotopic organization, architectonic fields, and
connections of auditory cortex in macaque monkeys. Journal of Comparative Neurology 335,
437-459.

Moutoussis, M., Bentall, R.P., El-Deredy, W., Dayan, P., 2011. Bayesian modelling of Jumping-
to-Conclusions bias in delusional patients. Cognitive neuropsychiatry 16, 422-447.

Murray, J.D., Bernacchia, A., Freedman, D.J., Romo, R., Wallis, J.D., Cai, X., Padoa-Schioppa,
C., Pasternak, T., Seo, H., Lee, D., 2014. A hierarchy of intrinsic timescales across primate
cortex. Nature neuroscience 17, 1661.

Olney, J.W., Farber, N.B., 1995. Glutamate receptor dysfunction and schizophrenia. Archives of
general psychiatry 52, 998-1007.

Poldrack, R.A., Congdon, E., Triplett, W., Gorgolewski, K., Karlsgodt, K., Mumford, J., Sabb, F.,
Freimer, N., London, E., Cannon, T., 2016. A phenome-wide examination of neural and cognitive
function. Scientific data 3, 160110.

Postmes, L., Sno, H., Goedhart, S., Van Der Stel, J., Heering, H., De Haan, L., 2014.
Schizophrenia as a self-disorder due to perceptual incoherence. Schizophrenia research 152, 41-
50.

Power, J.D., Barnes, K.A., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., 2012. Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. NeuroImage 59, 2142-2154.

Power, J.D., Cohen, A.L., Nelson, S.M., Wig, G.S., Barnes, K.A., Church, J.A., Vogel, A.C., Laumann, T.O., Miezin, F.M., Schlaggar, B.L., 2011. Functional network organization of the human brain. Neuron 72, 665-678.

Powers, A.R., Mathys, C., Corlett, P., 2017. Pavlovian conditioning–induced hallucinations result from overweighting of perceptual priors. Science 357, 596-600.

Powers III, A.R., Kelley, M., Corlett, P.R., 2016. Hallucinations as top-down effects on perception. Biological Psychiatry: Cognitive Neuroscience and Neuroimaging 1, 393-400.

Rao, R.P., Ballard, D.H., 1999. Predictive coding in the visual cortex: a functional interpretation of some extra-classical receptive-field effects. Nature neuroscience 2, 79.

Sarkissov, S., Filimonoff, I., Kononowa, E., Preobraschenskaja, I., Kukuew, L., 1955. Atlas of the cytoarchitectonics of the human cerebral cortex. Moscow: Medgiz 20.

Schmack, K., de Castro, A.G.-C., Rothkirch, M., Sekutowicz, M., Rössler, H., Haynes, J.-D., Heinz, A., Petrovic, P., Sterzer, P., 2013. Delusions and the role of beliefs in perceptual inference. Journal of neuroscience 33, 13701-13712.

Shadlen, M.N., Shohamy, D., 2016. Decision making and sequential sampling from memory. Neuron 90, 927-939.

Shergill, S.S., Samson, G., Bays, P.M., Frith, C.D., Wolpert, D.M., 2005. Evidence for sensory prediction deficits in schizophrenia. American Journal of Psychiatry 162, 2384-2386.

Shergill, S.S., White, T.P., Joyce, D.W., Bays, P.M., Wolpert, D.M., Frith, C.D., 2014. Functional magnetic resonance imaging of impaired sensory prediction in schizophrenia. JAMA psychiatry 71, 28-35.

Shrout, P.E., Fleiss, J.L., 1979. Intraclass correlations: uses in assessing rater reliability. Psychological bulletin 86, 420.

Smith, S.M., Beckmann, C.F., Andersson, J., Auerbach, E.J., Bijsterbosch, J., Douaud, G., Duff, E., Feinberg, D.A., Griffanti, L., Harms, M.P., 2013. Resting-state fMRI in the human connectome project. NeuroImage 80, 144-168.

Spence, S.A., Brooks, D.J., Hirsch, S.R., Liddle, P.F., Meehan, J., Grasby, P.M., 1997. A PET study of voluntary movement in schizophrenic patients experiencing passivity phenomena (delusions of alien control). Brain: a journal of neurology 120, 1997-2011.

Stephan, K.E., Weiskopf, N., Drysdale, P.M., Robinson, P.A., Friston, K.J., 2007. Comparing hemodynamic models with DCM. NeuroImage 38, 387-401.

Stephens, G.J., Honey, C.J., Hasson, U., 2013. A place for time: the spatiotemporal structure of neural dynamics during natural audition. Journal of neurophysiology 110, 2019-2026.

Sterzer, P., Adams, R.A., Fletcher, P., Frith, C., Lawrie, S.M., Muckli, L., Petrovic, P., Uhlhaas, P., Voss, M., Corlett, P.R., 2018. The predictive coding account of psychosis. Biological psychiatry 84, 634-643.

Teufel, C., Subramaniam, N., Dobler, V., Perez, J., Finnemann, J., Mehta, P.R., Goodyer, I.M., Fletcher, P.C., 2015. Shift toward prior knowledge confers a perceptual advantage in early psychosis and psychosis-prone healthy individuals. Proceedings of the National Academy of Sciences 112, 13401-13406.

Transtrum, M.K., Machta, B.B., Sethna, J.P., 2010. Why are nonlinear fits to data so challenging? Physical review letters 104, 060201.

van der Gaag, M., Hoffman, T., Remijsen, M., Hijman, R., de Haan, L., van Meijel, B., van Harten, P.N., Valmaggia, L., De Hert, M., Cuijpers, A., 2006. The five-factor model of the Positive and Negative Syndrome Scale II: a ten-fold cross-validation of a revised model. Schizophrenia research 85, 280-287.

Van Essen, D.C., Anderson, C.H., Felleman, D.J., 1992. Information processing in the primate visual system: an integrated systems perspective. Science 255, 419-423.

Van Essen, D.C., Smith, S.M., Barch, D.M., Behrens, T.E., Yacoub, E., Ugurbil, K., Consortium, W.-M.H., 2013. The WU-Minn human connectome project: an overview. NeuroImage 80, 62-79.

Van Essen, D.C., Ugurbil, K., Auerbach, E., Barch, D., Behrens, T., Bucholz, R., Chang, A., Chen, L., Corbetta, M., Curtiss, S.W., 2012. The Human Connectome Project: a data acquisition perspective. NeuroImage 62, 2222-2231.

VandenBos, G.R., 2007. APA dictionary of psychology. American Psychological Association.

Vázquez-Rodríguez, B., Suárez, L.E., Markello, R.D., Shafiei, G., Paquola, C., Hagmann, P., Van Den Heuvel, M.P., Bernhardt, B.C., Spreng, R.N., Misic, B., 2019. Gradients of structure–function tethering across neocortex. Proceedings of the National Academy of Sciences 116, 21219-21227.

Vogt, O., 1911. Die myeloarchitektonik des isocortex parietalis. J Psychol Neurol 18, 107-118.

Von Economo, C., 1929. The cytoarchitectonics of the human cerebral cortex. H. Milford Oxford University Press.

Wagstyl, K., Ronan, L., Goodyer, I.M., Fletcher, P.C., 2015. Cortical thickness gradients in structural hierarchies. NeuroImage 111, 241-250.

Watanabe, T., Rees, G., Masuda, N., 2019. Atypical intrinsic neural timescale in autism. Elife 8, e42256.

Waters, F., Fernyhough, C., 2017. Hallucinations: a systematic review of points of similarity and difference across diagnostic classes. Schizophrenia Bulletin 43, 32-43.

Weinberger, D.R., 1987. Implications of normal brain development for the pathogenesis of schizophrenia. Archives of general psychiatry 44, 660-669.

Weinstein, J.J., Chohan, M.O., Slifstein, M., Kegeles, L.S., Moore, H., Abi-Dargham, A., 2017. Pathway-specific dopamine abnormalities in schizophrenia. Biological psychiatry 81, 31-42.

1 Yang, G.J., Murray, J.D., Wang, X.-J., Glahn, D.C., Pearlson, G.D., Repovs, G., Krystal, J.H.,
2 Anticevic, A., 2016. Functional hierarchy underlies preferential connectivity disturbances in
3 schizophrenia. Proceedings of the National Academy of Sciences 113, E219-E228.

4 Young, M.P., 1993. The organization of neural systems in the primate cerebral cortex.
5 Proceedings of the Royal Society of London. Series B: Biological Sciences 252, 13-18.