The insert sequence in SARS-CoV-2 enhances spike protein cleavage by

TMPRSS

Tong Meng[1,2,10,11], Hao Cao[3,4,11], Hao Zhang[5,10,11], Zijian Kang[6,10], Da Xu[7,10], Haiyi Gong[5,10], Jing Wang[8], Zifu Li[8], Xingang Cui[7], Huji Xu[4,6], Haifeng Wei[5], Xiuwu Pan[7], Rongrong Zhu[9], Jianru Xiao[5]*, Wang Zhou[4,10]*, Liming Cheng[1]*, Jianmin Liu[8]*.

Division of Spine, Department of Orthopedics, Tongji Hospital affiliated to Tongji University School of Medicine, 200065 Shanghai, China

Tongji University Cancer Center, School of Medicine, Tongji University, 200092 Shanghai, China

School of Life Science and Biopharmaceutics, Shenyang Pharmaceutical University, 103 Wenhua Road, 110016 Shenyang, China

Peking-Tsinghua Center for Life Sciences, TsinghuaUniversity, 100084 Beijing, China

Department of Orthopaedic Oncology, Changzheng Hospital, Second Military Medical University, 200003 Shanghai, China

Department of Rheumatology and Immunology, Changzheng Hospital, Second Military Medical University, 200003 Shanghai, China

Depanrtment of Urology, The Third Affiliated Hospital of Second Military Medical University, 201805 Shanghai, China

Department of Neurosurgery, Changhai hospital, Second Military Medical University, 200003 Shanghai, China

9 Key Laboratory of Spine and Spinal Cord Injury Repair and Regeneration of

Ministry of Education, Orthopaedic Department of Tongji Hospital, School of Life

Science and Technology, Tongji University, 200092 Shanghai, China

10 Qiu-Jiang Bioinformatics Institute, 200003 Shanghai, China

11 These authors contributed equally to this work: Tong Meng, Hao Cao, Hao Zhang

*e-mail: chstroke@163.com; limingcheng@tongji.edu.cn; brilliant212@163.com;

jianruxiao83@163.com

Abstract

At the end of 2019, the SARS-CoV-2 induces an ongoing outbreak of pneumonia in

China[1], even more spread than SARS-CoV infection[2]. The entry of SARS-CoV into

host cells mainly depends on the cell receptor (ACE2) recognition and spike protein

cleavage-induced cell membrane fusion[3,4]. The spike protein of SARS-CoV-2 also

binds to ACE2 with a similar affinity, whereas its spike protein cleavage remains

unclear[5,6]. Here we show that an insertion sequence in the spike protein of

SARS-CoV-2 enhances the cleavage efficiency, and besides pulmonary alveoli,

intestinal and esophagus epithelium were also the target tissues of SARS-CoV-2.

Compared with SARS-CoV, we found a SPRR insertion in the S1/S2 protease

cleavage sites of SARS-CoV-2 spike protein increasing the cleavage efficiency by the

protein sequence aligment and furin score calculation. Additionally, the insertion

sequence facilitates the formation of an extended loop which was more suitable for

protease recognition by the homology modeling and molicular docking. Furthermore,

the single-cell transcriptomes identified that ACE2 and TMPRSSs are highly

coexpressed in AT2 cells of lung, along with esophageal upper epithelial cells and

absorptive enterocytes. Our results provide the bioinformatics evidence for the

increased spike protein cleavage of SARS-CoV-2 and indicate its potential target

cells.

Introduction

At the end of 2019, a rising number of pneumonia patients with unknown pathogen

emerged from Wuhan to nearly the entire China[7]. A novel coronavirus was isolated

and based on its phylogeny, taxonomy and established practice, the Coronavirus

Study Group (CSG) recognized it as a sister to severe acute respiratory syndrome

coronaviruses (SARS-CoVs) and labeled it as severe acute respiratory syndrome

coronavirus 2 (SARS-CoV-2) [1,8]. Although SARS-CoV-2 is generally less pathogenic

than SARS-CoV and Middle East respiratory syndrome coronavirus (MERS-CoV), it

has a relatively high transmissibility [9].

With regard to human coronavirus, the transmissibility and infectivity is largely

controlled by the spike (S) surface envelope protein [10]. Its surface unit (S1) mediates

the entry into host cells by binding to cell receptor and the transmembrane unit (S2)

subunit regulates the fusion of viral and cellular membranes [3]. Prior to membrane

fusion, the S protein should be cleaved and activated to allow for the fusion peptide

releasing onto host cell membranes (Fig. 1a) [11]. SARS-CoV-2 uses the same cell

receptor (angiotensin converting enzyme II, ACE2) as SARS-CoV, with a similar

binding affinity, whereas their transmissibility and infectivity are different [5,6,12,13].

Thus, the different virus transmission and infectivity may be associated with the

differentiated protease-induced S protein cleavage between SARS-CoV-2 and

SARS-CoV.

The transmembrane serine proteases (TMPRSSs) were the main host cell proteases on

the cell membrane[14]. The substrate specificity of TMPRSSs are almost similar and

revealing a strong preference for arginine or lysine residues in the P1 position.

Nowadays, their hydrolytic effects of TMPRSSs have been widely reported in

SARS-CoV and MERS-CoV pneumonia[15]. In the SARS-CoV-infected alveolar cells,

TMPRSSs, especially the TMPRSS2 and TMPRSS11D, cleave the SARS-CoV S

protein (SARS-S) at residue R667 (the S1/S2 cleavage site) and residue R797 (the S2'

cleavage site) (Fig. 1a)[15,16]. Besides cleaving S protein, they can also promote viral

spread in the host by cleaving ACE2 (Fig. 1b)[14,17]. Although SARS-CoV-2 and

SARS-CoV share the same host cell receptor with a similar affinity, however, the

SARS-CoV-2 S protein cleavage induced by TMPRSS remains unclear which may be

associated with the viral infectivity [4,5].

Results

The comparison of the S1/S2 and S2' cleavage sites between SARS-CoV-2 and

SARS-CoV

Generally, compared with SARS-CoV, the major differences in SARS-CoV-2 are the

three short insertions in the N-terminal domain and four out of five key residues

changes in the receptor-binding motif [5]. Here we used the alignment, furin score and

homology modeling to compare the sequence of the S1/S2 and S2' cleavage sites

between SARS-CoV-2 and SARS-CoV (Fig. 1c). The amino acid sequence of the

S1/S2 and S2' cleavage sites among ten beta-coronavirus were then analyzed and we

found that compared with SARS, there was an insertion sequence (SPRR) in the

S1/S2 cleavage sites of SARS-CoV-2 (Fig. 2a). The furin score was next used to

identify the cleavage efficiency of the insertion sequence in SARS-CoV-2. Its furin

score was 0.688, which was obviously higher than that of the corresponding sequence

in SARS-CoV (0.139), indicating that the insertion sequence may increase the

cleavage efficiency by proteases (Fig. 2b).

The structures of SARS-S and SARS-CoV-2 S protein were presented in Extended

Data Fig. 1a and 1b, along with their structural superimposition (Extended Data Fig.

1c). The structural comparison of homology modeling SARS-CoV-2 S protein with

SARS-S protein (PDB: 5x5b) showed that a exposed loop was formed by the insertion

which comprised R682 and R683 (S1/S2 site) on the surface of SARS-CoV-2 S

protein, and no significant difference of them in S2' site (Fig. 2c, d).

The insertion sequence of SARS-CoV-2 facilitating the TMPRSS recognition and

S protein cleavage

Structurally, TMPRSSs include extracellular domain, transmembrane domain and

intracellular domain in which extracellular domain is the main catalytic domain. They

show similar substrate-specificity and catalytic mechanism. Take TMPRSS2 as an

example. The catalytic triad consisted of H296, D345 and S441 and the substrate

binding residue D435, a conserved aspartate residue, was located in the bottom of

pocket[18,19]. The substrate binding pocket is deeper than most of serine proteinase

(Extended Data Fig. 2a, b). The bottom of the catalytic pocket has a negatively

charged aspartic acid residue which can facilitate the binding and stabilization of

arginine or lysine residues in the P1 position[18,19].

Polypeptide substrate analogue KQLR included arginine, glutamine, leucine and

lysine (Extended Data Fig. 2c). The substrate analogue could bind to the catalytic

pocket of TMPRSS2 (Extended Data Fig. 2d, e). The conformation of the insertion

sequence in SARS-CoV-2 S protein and TMPRSS2 was next simulated by molecular

docking. We found the insertion sequence formed a loop which was easily recognized

by the catalytic pocket of TMPRSS2 (Extended Data Fig. 2f, g). Thus, both the furin

score and molecular docking revealed that the insertion sequence of SARS-CoV-2

facilitates the TMPRSS2 recognition and S protein cleavage.

The potential target tissues of COVID-19

The entry of SARS-CoV-2 into host cells depends on the cell receptor recognition and

cell proteases cleaving. Thus, the target cells should coexpress both the cell receptor

ACE2 and cell proteases TMPRSSs. In order to identify the coexpressing cell

composition and proportion, we utilized 3 datasets including 32 samples and built the

largest single-cell transcriptome atlas of normal lung, the commonest infected organ

of SARS-CoV-2.

After initial quality controls, a total of 113,045 cells and 29 sub-clusters were

identified in the lung (Fig. 3a). The marker genes and dataset proportions of each

sub-cluster were presented in Extended Data Fig. 3-4.

We detected the expression of ACE2 and TMPRSSs in 29 cell groups, in which the

expression of the whole 17 TMPRSS genes is in the form of total signature value.

Pseudodyeing analysis was performed and we found that ACE2 was mainly expressed

in AT2 cells and marked with red (Fig. 3b, c). The total 17 TMPRSS genes was found

in AT1, AT2, airway secretory and ciliated cells colored with blue (Fig. 3b, d,

Extended Data Fig. 5a). Thus, we found an obvious coexpression between TMPRSSs

and ACE2 in AT2. Among the whole TMPRSS genes, TMPRSS1 and TMPRSS2

were highly expressed in AT2 and AT1 cells, which were co-expressed with ACE2 in

lung (Fig. 3b, Extended Data Fig. 5b). Due to the entry of virus into host cell is

related to endocytosis, we also detected the endocytosis-related genes among different

cells. We found that these genes had consistent distribution and highly expressed in

AT1, AT2, airway secretory, ciliated cells and M2 macrophage (Extended Data Fig.

5c).

Due to the RNA of SARS-CoV-2 was also found in the stool specimen of the

SARS-CoV-2-infected patient[20], the digestive system may also be the potential route

of COVID-19. Thus, in addition to lung, 4 datasets with the single-cell transcriptomes

of the esophagus, gastric, small intestine and colon were analyzed to identify the

expression of ACE2 and TMPRSSs in the digestive system. The co-expression of

ACE2 and TMPRSS was analyzed in esophagus, stomach, small intestine and colon

by 87947, 29678, 11218 and 47442 high-quality single cells, respectively (Extended

Data Fig. 6a). The coexpression of ACE2 and total TMPRSS genes were found in the

upper epithelial cells of esophagus, the absorptive enterocytes of ileum epithelia and

the enterocytes of colon epithelia (Extended Data Fig. 6b-e, 7a-d).

As both ACE2 and TMPRSSs are expressed in the lung and digestive system, we next

compared their relative expression values in the ACE2-expressing cells. A similar

distribution was found between ACE2 and TMPRSSs in all the 9 clusters with high

expressions in the esophageal upper epithelial cells, the ileal absorptive enterocytes

and the colonic enterocytes (Fig. 4a). In addition, their expression of AT2 was

relatively lower than that of epithelial cells in the digestive system. Among all the

TMPRSSs, TMPRSS1 and TMPRSS2 were relatively highly expressed in AT2, and

most TMPRSSs were highly found in the esophageal upper epithelial cells (Extended

Data Fig. 8a). The endocytosis- and exocytosis-associated genes which are related to

the entry of virus into host cells and virus infection were also detected in all the 9

clusters. The endocytosis signature was more expressed in AT1 and AT2 cells,

whereas the exocytosis signature was highly gathered in esophageal upper epithelial

cells. It can explain that the commonest infected tissue in COVID-19 is pulmonary

alveoli and SARS-CoV-2 can also be detected in the esophageal erosion (Fig. 4b)[21].

The RNA-seq data of lung, esophagus, stomach, small intestine, colon-transverse and

colon-sigmoid were obtained from GTEx database. The expressions of ACE2 and

TMPRSS2 also had a similar tendency and were highly expressed in small intestine

and colon, while the TMPRSS11D was mainly found in the esophagus (Extended

Data Fig. 8b).

Discussion

The coronaviruses is the common infection source of respiratory, enteric and central

nervous system in humans and other mammals[22]. At the beginning of the twenty-first

century, two betacoronaviruses, SARS-CoV and MERS-CoV, result in persistent

public panics and became the most significant public health events[23]. In December

2019, a novel identified coronavirus (SARS-CoV-2) induced an ongoing outbreak of

pneumonia in Wuhan, Hubei, China [7]. The rapidly increasing number of

SARS-CoV-2-infected cases suggests that SARS-CoV-2 may be transmitted

effectively among humans and give rise to a high pandemic potential [7,8,24].

Previous studies identified that SARS-CoV mutated between 2002 and 2004 to better

bind to its cell receptor, replicate in human cells and enhance the virulence [9]. Thus, it

is important to explore whether SARS-CoV-2 behaves like SARS-CoV to adapt to the

host cell. Notably, SARS-CoV and SARS-CoV-2 share the same receptor protein

ACE2[5,25]. Besides, the receptor-binding domain (RBD) in S protein of SARS-CoV-2

binds to ACE2 with the similar affinity as SARS-CoV RBD does[6]. Thus, during the

process of viral and host cellular membrane fusion, whether the specific structure of

SARS-CoV-2 S protein seems better suited to be activated by host cell proteases may

be related to the different virus infectivities and transmissibilities between

SARS-CoV and SARS-CoV-2[6].

In this study, we found the furin score of the S1/S2 cleavage sites in SARS-CoV-2

was higher than that of SARS, implying a more degree of hydrolysis. Through the

comparison of the two structures, R682, R683 and relative S680, P681 extended the

original exposed loop combined with R685 of SARS-CoV-2, which was more

suitable for hydrolysis by TMPRSSs. The substrate specificity of TMPRSSs are

almost similar, revealing a strong preference for arginine or lysine residues in the P1

position represented by R. More R (R682, R683 and R685) in the S1/S2 cleavage

sites of SARS-CoV-2 can enhance the cleavage of S1 with S2, which means that the

structurally constrains of S1 on S2 is removed, and the fusion peptides in S2 are

exposed and insert into the target host cell membrane, finally it increases the

efficiency of fuse membranes[18,19].

By the way, some researchers previously supposed the SARS-CoV-2 was artificial

due to four inserts in the S protein of SARS-CoV-2 from HIV sequence. However, the

results of protein sequence alignment revealed that the similar sequence of the

reported fourth insertion site (680-SPRR-683) in SARS-CoV-2 was commonly found

in many beta-coronavirus. Therefore, we supposed that based on the current evidence,

it is not scientific to consider the insertion sequence in SARS-CoV-2 S protein being

artificial.

With the help of single cell sequencing, we found a strong co-expression between

ACE2 and TMPRSSs, in especial TMPRSS1and TMPRSS2, in lung AT2 cells, which

was also the main infected cell type in SARS-CoV pneumonia[26]. Moreover, we also

found the endocytosis-associated genes was highly expressed in AT2 cells, implying

that endocytosis may also facilitate the entry of SARS-CoV-2 into host cells. As the

alveolar stem-like cells, AT2 cells are in charge of surfactant biosynthesis,

self-renewal and immunoregulation[27]. Thus, SARS-CoV-2 not only damages the AT2

cells leading to the direct injury to alveoli, but also raises alveolar surface tension to

induce dyspnea[28]. Additionally, the injuryed AT2 also damages the immunologic

balance in alveoli and results in inflammatory cascade[29]. In addition, they are also

highly co-expressed in absorptive enterocytes and upper epithelial cells of esophagus,

implying that intestinal epithelium and esophagus epithelium may also be the

potential target tissues. This can explain the cases whose SARS-CoV-2 was detected

in the esophageal erosion or stool specimen, implying that the digestive system is a

potential route of COVID-19 [7,20,21].

Due to the critical role of TMPRSSs in influenza virus and coronavirus infections,

serine protease inhibitors, such as camostat, nafamostat and leupeptin, have been used

in the antiviral treatment targeting TMPRSSs with high antiviral activities [14,30,31].

Nowadays, Remdesivir (GS-5734) has been used in the treatment of SARS-CoV-2,

however, the therapeutic effects are still unclear. Based on our results, we also

supposed that TMPRSSs may also serve as candidate antiviral targets for

1. SARS-CoV-2 infection and the clinical trials of serine protease inhibitors should also

2. be performed for COVID-19.

3. **Methods**

4. **Structure modelling**

5. The structures of SARS-CoV-2 S protein and TMPRSS2 were generated by

6. SWISS-MODEL online server[32]. The structures were marked, superimposed and

7. visualized by Chimera[33]. To further explore the possible catalytic mechanism of the

8. SARS-CoV-2 S protein cleaved by TMPRSS2, ZDOCK program was used to predict

9. their interaction [34]. A total of 5000 models were generated and were set to 50 clusters,

10. then the best scoring models from the 5 largest clusters were selected for further

11. analysis.

12. **Furin score**

13. The fragmentation maps, scoring and residue coverage analysis were conducted using

14. arginine and lysine propeptide cleavage sites prediction algorithms ProP 1.0 server[35].

15. **Single cell transcriptome data sources**

16. Single cell transcriptome data were obtained from Single Cell Portal

17. (https://singlecell.broadinstitute.org/single_cell), Human Cell Atlas Data Protal

18. (https://data.humancellatlas.org) and Gene Expression Omnibus (GEO;

19. https://www.ncbi.nlm.nih.gov/). Esophageal and lung data were obtained from the

20. research of E Madissoon *et al* containing 21 esophageal and 19 lung tissue samples[36].

21. Two lung datasets were further obtained from GSE122960[38] and GSE128169[39],

including eight and five lung tissues respectively. GSE134520 included 6 gastric

mucosal samples from 3 non-atrophic gastritis and 2 chronic atrophic gastritis

patients[40]. GSE134809 comprises 11 noninflammatory ileal samples from Crohn's

disease patients[41]. The data from Christopher S *et al* consisted of 12 normal colon

samples[42].

Quality control

Cells would be identified as poor-quality once (1) the number of expressed genes

fewer than 200 or greater than 5000, or (2) more than 20% of UMIs being mapped to

mitochondrial or ribosomal genes.

Data Integration, Dimension Reduction and Cell Clustering

Different methods were performed to process the downloaded data:

1. Esophagus dataset. Rdata were obtained and dimension reduction and clustering

had already been implemented by the authors [36].

2. Lung, stomach and ileum datasets. We utilized functions in the Seurat package to

normalize and scale the single-cell gene expression data[43]. Unique

molecularidentifier (UMI) counts were normalized by the total number of UMIs

per cell, multiplied by 10000 for normalization and log-transformed using the

NormalizeData" function. Then, multiple sample data within each dataset were

merged using the "FindIntegrationAnchors" and "Integratedata" functions. After

identifying highly variable genes (HVGs) using the "FindVariableGenes" function

a principal component analysis (PCA) was performed on the single-cell

expression matrix using the "RunPCA" function. The "FindClusters" function in

the Seurat package was next utilized to conduct the cell clustering analysis into a

graph structure in PCA space after constructing a K-nearest-neighbor graph based

on the Euclidean distance in PCA space. Uniform Manifold Approximation and

Projection (UMAP) visualization was performed for obtaining the clusters of

cells.

3.   Colon Dataset. The single cell data was processed with the R packages LIGER[44]

and Seurat[43]. The gene expression matrix was first normalized to remove

differences in sequencing depth and capture efficiency among cells. Variable

genes in each dataset were identified using the "selectGenes" function. Then we

used the "optimizeALS" function in LIGER to perform the integrative

nonnegative matrix factorization and selecte a k of 15 and lambda of 5.0 to obtain

a plot of expected alignment. The "quantileAlignSNF" function was then

performed to builds a shared factor neighborhood graph to jointly cluster cells,

then quantile normalizes corresponding clusters. Next nonlinear dimensionality

reduction was calculated using the "RunUMAP" function and the results were

visualized with UMAP.

Identification of cell types and Gene expression analysis

Clusters were annotated on the expression of known cell markers and the clustering

information provided in the articles. Then, we utilized the "RunALRA" function to

impute lost values in the gene expression matrix. The imputed gene expression was

shown in Feature plots and violin plots. We used "Quantile normalization" in the R

package preprocessCore (R package version 1.46.0.

https://github.com/bmbolstad/preprocessCore) to remove unwanted technical

variability across different datasets. The data were further denoised to compare the

gene expression levels of gene signature.

Endocytosis or exocytosis associated genes were obtained from Harmonizome dataset

[45].Mean expressions of the genesets were calculated to compare the ability of

endocytosis or exocytosis among clusters.

To minimize bias, external databases of Genotype-Tissue Expression (GTEx)[46] was

used to detect gene expression of ACE2, TMPRSS1 and TMPRSS2 at the tissue

levels including normal lung and digestive system, such as esophagus, stomach, small

intestine and colon.

Acknowledgements This study was jointly supported by the National Natural

Science Foundation of China (Grants 81702659 and 81572746) and National Key

R&D Program of China (Grants 2016YFA0100800).

Author contributions J.L., L.C., W.Z. and J.X. conceived the idea and directed the

team. T.M., H.C., H.Z. and W.Z. designed and coordinated the analysis and

characterization. H.Z., Z.K., D.X., H.G. performed single-cell sequencing and

characterization under the guidance of X.C., H.X., and H.W.. Data collection and

generation were performed by J.W., Z.L., R.Z. and X.P.. Data interpretation was

performed by J.L., L.C., W.Z. and J.X.. The alignment and structure comparison was

performed by H.C. under the guidance of W.Z. The manuscript was written by T.M.,

H.C., Z.K. and W.Z. All authors contributed to the analysis and discussion of the

results leading to the manuscript.

Competing interests The authors declare no competing interests.

1 Gorbalenya AE, B. S., Baric RS, de Groot RJ, Drosten C, Gulyaeva AA,

Haagmans BL, Lauber C, Leontovich AM, Neuman BW, Penzar D,

Perlman S, Poon LL, Samborskiy D, Sidorov IA, Sola I, Ziebuhr J.

Severe acute respiratory syndrome-related coronavirus: The species

and its viruses – a statement of the Coronavirus Study Group. *bioRxiv*

doi:doi: https://doi.org/10.1101/2020.02.07.937862 (2020).

2 Zhong, N. S. *et al.* Epidemiology and cause of severe acute respiratory

syndrome (SARS) in Guangdong, People's Republic of China, in

February, 2003. *Lancet* **362**, 1353-1358,

doi:10.1016/s0140-6736(03)14630-2 (2003).

3 Hofmann, H. & Pohlmann, S. Cellular entry of the SARS coronavirus.

Trends in microbiology **12**, 466-472, doi:10.1016/j.tim.2004.08.008

(2004).

4 Chan, J. F. *et al.* A familial cluster of pneumonia associated with the

novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet*, doi:10.1016/s0140-6736(20)30154-9 (2020).

P Zhou, X. Y., XG Wang, B Hu, L Zhang, W Zhang, HR Si, Y Zhu, B Li, CL Huang, HD Chen, J Chen, Y Luo, H Guo, RD Jiang, MQ Liu, Y Chen, XR Shen, X Wang, XS Zheng, K Zhao, QJ Chen, F Deng, LL Liu, B Yan, FX Zhan, YY Wang, GF Xiao, ZL Shi. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*, doi:https:// doi.org/10.1038/s41586-020-2012-7 (2020).

Tian XL, L. C., Huang A, Xia S, Lu SC, Shi ZL, Lu L, Jiang SB, Yang ZL, Wu YL, Ying TL. Potent binding of 2019 novel coronavirus spike protein by a SARS coronavirus-specific human monoclonal antibody. *bioRxiv* doi:doi: https://doi.org/10.1101/2020.01.28.923011 (2020).

Huang, C. *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*, doi:10.1016/s0140-6736(20)30183-5 (2020).

Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *The New England journal of medicine*, doi:10.1056/NEJMoa2001017 (2020).

Chen, J. Pathogenicity and Transmissibility of 2019-nCoV-A Quick Overview and Comparison with Other Emerging Viruses. *Microbes and*

infection, doi:10.1016/j.micinf.2020.01.004 (2020).

Walls, A. C. *et al.* Unexpected Receptor Functional Mimicry Elucidates Activation of Coronavirus Fusion. *Cell* **176**, 1026-1039.e1015, doi:10.1016/j.cell.2018.12.028 (2019).

Gallagher, T. M. & Buchmeier, M. J. Coronavirus spike proteins in viral entry and pathogenesis. *Virology* **279**, 371-374, doi:10.1006/viro.2000.0757 (2001).

Gui, M. *et al.* Cryo-electron microscopy structures of the SARS-CoV spike glycoprotein reveal a prerequisite conformational state for receptor binding. *Cell research* **27**, 119-129, doi:10.1038/cr.2016.152 (2017).

Zhao S, R. J., MUSA SS, Yang G, Lou Y, Gao D, Yang L, He D. . Preliminary estimation of the basic reproduction number of novel coronavirus (2019-nCoV) in China, from 2019 to 2020: A data-driven analysis in the early phase of the outbreak. . *bioRxiv*, 916395, doi:doi: https://doi.org/10.1101/2020.01.23.916395 (2020).

Zhou, Y. *et al.* Protease inhibitors targeting coronavirus and filovirus entry. *Antiviral research* **116**, 76-84, doi:10.1016/j.antiviral.2015.01.011 (2015).

Millet, J. K. & Whittaker, G. R. Host cell proteases: Critical determinants of coronavirus tropism and pathogenesis. *Virus research* **202**, 120-134,

doi:10.1016/j.virusres.2014.11.021 (2015).

Shirato, K., Kawase, M. & Matsuyama, S. Wild-type human coronaviruses prefer cell-surface TMPRSS2 to endosomal cathepsins for cell entry. *Virology* **517**, 9-15, doi:10.1016/j.virol.2017.11.012 (2018).

Heurich, A. *et al.* TMPRSS2 and ADAM17 cleave ACE2 differentially and only proteolysis by TMPRSS2 augments entry driven by the severe acute respiratory syndrome coronavirus spike protein. *Journal of virology* **88**, 1293-1307, doi:10.1128/jvi.02202-13 (2014).

Herter, S. *et al.* Hepatocyte growth factor is a preferred in vitro substrate for human hepsin, a membrane-anchored serine protease implicated in prostate and ovarian cancers. *The Biochemical journal* **390**, 125-136, doi:10.1042/bj20041955 (2005).

Limburg, H. *et al.* TMPRSS2 Is the Major Activating Protease of Influenza A Virus in Primary Human Airway Cells and Influenza B Virus in Human Type II Pneumocytes. *Journal of virology* **93**, doi:10.1128/jvi.00649-19 (2019).

Holshue, M. L. *et al.* First Case of 2019 Novel Coronavirus in the United States. *The New England journal of medicine*, doi:10.1056/NEJMoa2001191 (2020).

Guan WJ, N. Z., Hu Y, Liang WH, Ou CQ, He JX, Liu L, Shan H, Lei CL,

Hui David S.C., Du B   Clinical characteristics of 2019 novel coronavirus infection in China. *medRxiv*, doi:doi: http://dx.doi.org/10.1101/2020.02.06.20020974 (2020).

Perlman, S. & Netland, J. Coronaviruses post-SARS: update on replication and pathogenesis. *Nature reviews. Microbiology* 7, 439-450, doi:10.1038/nrmicro2147 (2009).

de Wit, E., van Doremalen, N., Falzarano, D. & Munster, V. J. SARS and MERS: recent insights into emerging coronaviruses. *Nature reviews. Microbiology* **14**, 523-534, doi:10.1038/nrmicro.2016.81 (2016).

Lee, P. I. & Hsueh, P. R. Emerging threats from zoonotic coronaviruses-from SARS and MERS to 2019-nCoV. *Journal of microbiology, immunology, and infection = Wei mian yu gan ran za zhi*, doi:10.1016/j.jmii.2020.02.001 (2020).

Li, W. *et al.* Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. *Nature* **426**, 450-454, doi:10.1038/nature02145 (2003).

Kuiken, T. *et al.* Newly discovered coronavirus as the primary cause of severe acute respiratory syndrome. *Lancet* **362**, 263-270, doi:10.1016/s0140-6736(03)13967-0 (2003).

Nabhan, A. N., Brownfield, D. G., Harbury, P. B., Krasnow, M. A. &

Desai, T. J. Single-cell Wnt signaling niches maintain stemness of alveolar type 2 cells. *Science (New York, N.Y.)* **359**, 1118-1123, doi:10.1126/science.aam6603 (2018).

Barkauskas, C. E. *et al.* Type 2 alveolar cells are stem cells in adult lung. *The Journal of clinical investigation* **123**, 3025-3036, doi:10.1172/jci68782 (2013).

Kroetz, D. N. *et al.* Type I Interferon Induced Epigenetic Regulation of Macrophages Suppresses Innate and Adaptive Immunity in Acute Respiratory Viral Infection. *PLoS pathogens* **11**, e1005338, doi:10.1371/journal.ppat.1005338 (2015).

Shen, L. W., Mao, H. J., Wu, Y. L., Tanaka, Y. & Zhang, W. TMPRSS2: A potential target for treatment of influenza virus and coronavirus infections. *Biochimie* **142**, 1-10, doi:10.1016/j.biochi.2017.07.016 (2017).

Shin, W. J. & Seong, B. L. Type II transmembrane serine proteases as potential target for anti-influenza drug discovery. *Expert opinion on drug discovery* **12**, 1139-1152, doi:10.1080/17460441.2017.1372417 (2017).

Biasini, M. *et al.* SWISS-MODEL: modelling protein tertiary and quaternary structure using evolutionary information. *Nucleic acids research* **42**, W252-258, doi:10.1093/nar/gku340 (2014).

Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for

exploratory research and analysis. *Journal of computational chemistry* **25**, 1605-1612, doi:10.1002/jcc.20084 (2004).

Wiehe, K. *et al.* ZDOCK and RDOCK performance in CAPRI rounds 3, 4, and 5. *Proteins* **60**, 207-213, doi:10.1002/prot.20559 (2005).

Duckert, P., Brunak, S. & Blom, N. Prediction of proprotein convertase cleavage sites. *Protein engineering, design & selection : PEDS* **17**, 107-112, doi:10.1093/protein/gzh013 (2004).

Madissoon, E. *et al.* scRNA-seq assessment of the human lung, spleen, and esophagus tissue stability after cold preservation. *Genome biology* **21**, 1, doi:10.1186/s13059-019-1906-x (2019).

Vieira Braga, F. A. *et al.* A cellular census of human lungs identifies novel cell states in health and in asthma. *Nature medicine* **25**, 1153-1163, doi:10.1038/s41591-019-0468-5 (2019).

Reyfman, P. A. *et al.* Single-Cell Transcriptomic Analysis of Human Lung Provides Insights into the Pathobiology of Pulmonary Fibrosis. *American journal of respiratory and critical care medicine* **199**, 1517-1536, doi:10.1164/rccm.201712-2410OC (2019).

Valenzi, E. *et al.* Single-cell analysis reveals fibroblast heterogeneity and myofibroblasts in systemic sclerosis-associated interstitial lung disease. *Annals of the rheumatic diseases* **78**, 1379-1387, doi:10.1136/annrheumdis-2018-214865 (2019).

Zhang, P. *et al.* Dissecting the Single-Cell Transcriptome Network Underlying Gastric Premalignant Lesions and Early Gastric Cancer. *Cell reports* 27, 1934-1947.e1935, doi:10.1016/j.celrep.2019.04.052 (2019).

Martin, J. C. *et al.* Single-Cell Analysis of Crohn's Disease Lesions Identifies a Pathogenic Cellular Module Associated with Resistance to Anti-TNF Therapy. *Cell* 178, 1493-1508.e1420, doi:10.1016/j.cell.2019.08.008 (2019).

Smillie, C. S. *et al.* Intra- and Inter-cellular Rewiring of the Human Colon during Ulcerative Colitis. *Cell* 178, 714-730.e722, doi:10.1016/j.cell.2019.06.029 (2019).

Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* 177, 1888-1902.e1821, doi:10.1016/j.cell.2019.05.031 (2019).

Welch, J. D. *et al.* Single-Cell Multi-omic Integration Compares and Contrasts Features of Brain Cell Identity. *Cell* 177, 1873-1887.e1817, doi:10.1016/j.cell.2019.05.006 (2019).

Rouillard, A. D. *et al.* The harmonizome: a collection of processed datasets gathered to serve and mine knowledge about genes and proteins. *Database : the journal of biological databases and curation* 2016, doi:10.1093/database/baw100 (2016).

Human genomics. The Genotype-Tissue Expression (GTEx) pilot

analysis: multitissue gene regulation in humans. *Science (New York,*

N.Y.) **348**, 648-660, doi:10.1126/science.1262110 (2015).

47   Uhlen, M. *et al.* Proteomics. Tissue-based map of the human proteome.

Science (New York, N.Y.) **347**, 1260419, doi:10.1126/science.1260419

(2015).

7
8   **Fig. 1 The schematic diagram of the project.**

9   a.   The entry of SARS-CoV into host cells: The spike protein of SARS-CoV binds to

ACE2 through its S1 subunit for viral recognition. Then it is cleaved by TMPRSS2 at the S1/S2 boundary or within S2 subunit, which removes the structural constraint of S1 on S2, and releases the internal fusion peptide combined with the Spike TM domain for the fusion of viral and cellular membranes. Finally, the viral genomes enter into the host cells.

b. ACE2 cleaving by TMPRSSs: TMPRSS2 can also cleave ACE2 amino acids 697 to 716, resulting in the shedding of 13kD ACE2 fragment in culture supernatants and augmented viral infectivity.

c. The difference between SARS-CoV-2 and SARS-CoV in the Spike protein cleavage: The Spike protein of SARS involves two cleavage sites recognized by TMPRSSs, one at arginine 667 and the other at arginine 797 (right). Compared with SARS-CoV, the Spike protein of SARS-CoV-2 (left) has an insertion sequence 680-SPRR-683 at the S1/S2 cleavage site. We speculated that R682, R683 and R685 (red box) could be used as the most suitable substrates for TMPRSSs, which can increase the Spike protein cleavage efficiency of TMPRSSs, promote its activation and enhance SARS-CoV-2 infection.

Fig. 2 The two potential Spike protein cleavage sites of SARS-CoV and SARS-CoV-2 by TMPRSS2.

a. Phylogenetic tree based on the protein sequences of Spike protein in SARS-CoV-2, SARS-CoV and other eight beta-coronaviruses are presented, along with the amino acid sequence alignment of two potential cleavage sites by TMPRSS2.

b. The putative furin scores of the two potential cleavage sites of the ten coronaviruses.

c. Structure comparison of the detailed Spike protein of the SARS-CoV and SARS-CoV-2. The insert 675-690 of SARS-CoV-2 Spike protein (yellow) and the corresponding loci to SARS-CoV Spike protein 661-672 (green). Three important

1 residues, R682, R683, R685, are specially marked.

2 d. The detail of c. The similarly SARS-CoV R797 with SARS-CoV-2 R815 are

3 marked with forest green and orange, respectively.

6 **Fig. 3 Single-cell analysis of the normal lung tissue.**

a. The UMAP plots of the landscape of lung cells. Thirteen clusters are colored,

distinctively labeled. T, B and myeloid cell subsets are further divided into finer

cell subsets according to the heterogeneity within the cell population.

b. The feature plots of the 17 TMPRSS genes, ACE2, TMPRSS1 and TMPRSS2.

c. The expression of ACE2 across clusters in the violin plot. The expression is

measured as the log2 (TP10K+1) value.

d. The mean expression of TMPRSS family genes across clusters in the boxplot. The

expression is measured as the mean log2 (TP10K+1) value.

Fig. 4 Expression levels of ACE2, TMPRSS restriction signature and functional

gene sets in lung and digestive tracts.

a. The expression levels of ACE2 and TMPRSS restriction signature in 2 lung

clusters and 7 digestive tract clusters. The expression is measured as the log2

(TP10K+1) value.

b. The expression levels of endocytosis and exocytosis-associated genes in 2 lung

clusters and 7 digestive tract clusters. The expression is measured as the log2

(TP10K+1) value.

2 **Extended Data Fig. 1 The overall structure of the Spike protein in SARS-CoV**

3 **and SARS-CoV-2 homo-trimers**

4 a. The structure of the SARS-CoV Spike protein (from PDB: 5X5B). The insert

5   aa675-690 to SARS-CoV Spike protein aa661-672 with the structural missed

6   residues are marked with green.

7 b. The structure of the SARS-CoV-2 Spike protein (Modelled by SWISS-MODEL).

8   The insert aa675-690 of 2019-nCoV Spike protein that corresponds to the insert

9   region of SARS-V Spike protein is marked with yellow.

10 c. The structural superimpose of Spike protein in the SARS-CoV (yellow) and

11   SARS-CoV-2 (blue).

Extended Data Fig. 2 The structure and catalytic mechanism of TMPRSS2

a-b. The overall structure and surface of TMPRSS2 (Modelled by SWISS-MODEL).

The TMPRSS2, catalytic triad comprised of H296, D345 and S441 are marked with

cyan, blue, cyan and green, respectively. The substrate binding residue D435 located

in the bottom of pocket is marked with red.

c. The polypeptide substrate analogue KQLR. The cleavage site Arg is marked with

orange. Gln and Leu are marked with yellow. Lys is marked with pink.

d. The state of substrate analogue binding in the catalytic pocket. The state of

substrate analogue binding in the catalytic pocket.

e. The detail of d. Arg of substrate analogue is strongly interacted with D435

f. The predicted state of SARS-CoV-2 Spike protein binding to the catalytic pocket of

TMPRSS2.

g. The detail of f. SARS-CoV-2 Spike protein and D345 of TMPRSS2 are marked

with wheat and medium blue, respectively.

Extended Data Fig. 3 Subset-specific markers.

a. The heatmap of marker genes (rows) across cell subsets (columns). The bubble

diagram of marker genes in thirteen clusters (b) and the sub-clusters of T cells (c),

B cells (d) and Myeloid cells (e).

2 **Extended Data Fig. 4 All cell subset distributions across samples.**

3 The fractions of cells (y axis) in each cell subset (bars) that are derived from each

4 sample in 3 databases (red, green and blue). The numbers of cells in each cluster are

5 labeled above.

Extended Data Fig. 5 The expression levels of ACE2, TMPRSS genes and

exocytosis-associated genes in lung subsets.

a. The Violin plots of ACE2 and TMPRSS family genes across clusters. The

expression is measured as the log2 (TP10K+1) value.

b. The boxplot of exocytosis-associated across clusters. The expression is measured

as the mean log2 (TP10K+1) value.

Extended Data Fig. 6 The single-cell analysis of esophageal cells, gastric mucosal

cells, ileal epithelial cells and colonic epithelial cells.

a. The UMAP plots of esophageal cells, gastric mucosal cells, ileal epithelial cells

and colonic epithelial cells. The Feature plots show the expression of ACE2 (red)

and TMPSS family genes (green). The plots were merged to reveal the

co-expression of these genes (brown).

c. The expression levels ACE2 and TMPRSS restriction signature across clusters in

esophagus (b), stomach(c), ileum(d) and colon(d). The expression is measured as

the mean log2 (TP10K+1) value.

2 **Extended Data Fig. 7 The expression levels of ACE2 and TMPRSS family genes**

3 **in lung and digestive tracts.**

4 The violin plots of ACE2 and TMPRSS family genes across clusters in esophagus (a),

5 stomach (b), ileum (c) and colon (d). The expression is measured as the mean log2

6 (TP10K+1) value.

Extended Data Fig. 8 The expression levels of ASE2 and TMPRSS family genes

in lung and digestive tracts

a. The violin plots of TMPRSS family genes in lung and digestive tracts. The
    expression is measured as the mean log2 (TP10K+1) value.

b. The expression levels of ACE2, TMPRSS1 and TMPRSS2 verified by RNA-seq
    data from the GTEx database.