Identification of a Pangolin Niche for a 2019-nCoV-like Coronavirus via an
Extensive Meta-metagenomic Search

Lamia Wahba[1]*, Nimit Jain[1,3]*, Andrew Z. Fire[1,2]*, Massa J. Shoura[1]*, Karen L.
Artiles[1]*, Matthew J. McCoy[1]*, Dae-Eun Jeong[1]*

[1]Department of Pathology, Stanford University School of Medicine, Stanford, CA 94305, USA

[2]Department of Genetics, Stanford University School of Medicine, Stanford, CA 94305, USA

[3]Department of Bioengineering, Stanford University, Stanford, CA 94305, USA

*Co-corresponding and equally contributing authors. Author order was chosen randomly.

Author for publication correspondence: *Andrew Fire (afire@stanford.edu)*

Abstract:

In numerous instances, tracking the biological significance of a nucleic acid sequence
can be augmented through the identification of environmental niches in which the sequence
of interest is present. Many metagenomic datasets are now available, with deep sequencing of
samples from diverse biological niches. While any individual metagenomic dataset can be
readily queried using web-based tools, meta-searches through all such datasets are less
accessible. In this brief communication, we demonstrate such a meta-meta-genomic
approach, examining close matches to the Wuhan coronavirus 2019-nCoV in all high-
throughput sequencing datasets in the NCBI Sequence Read Archive accessible with the
keyword "virome". In addition to the homology to bat coronaviruses observed in descriptions
of the 2019-nCoV sequence (F. Wu et al. 2020, Nature, doi.org/10.1038/s41586-020-2008-3;
P. Zhou et al. 2020, Nature, doi.org/10.1038/s41586-020-2012-7), we note a strong
homology to numerous sequence reads in a metavirome dataset generated from the lungs of
deceased Pangolins reported by Liu et al. (Viruses 11:11, 2019,
http://doi.org/10.3390/v11110979). Our observations are relevant to discussions of the
derivation of 2019-nCoV and illustrate the utility and limitations of meta-metagenomic
search tools in effective and rapid characterization of potentially significant nucleic acid
sequences.

Importance:

Meta-metagenomic searches allow for high-speed, low-cost identification of potentially significant biological niches for sequences of interest.

Introduction:

In the early years of nucleic acids sequencing, aggregation of the majority of published DNA and RNA sequences into public sequence databases greatly aided biological hypothesis generation and discovery. Search tools capable of interrogating the ever-expanding databases were facilitated by creative algorithm development and software engineering, and by the ever-increasing capabilities of computer hardware and the internet. As of the early 2000s, sequencing methodologies and computational technologies advanced in tandem, enabling quick homology results from a novel sequence without substantial cost.

With the development of larger-scale sequencing methodologies, the time and resources to search all extant sequence data became untenable for most studies. However, creative approaches involving curated databases and feature searches ensured that many key features of novel sequences remained readily accessible. At the same time, the nascent field of metagenomics began, with numerous studies highlighting the power of survey sequencing of DNA and RNA from samples as diverse as the human gut and Antarctic soil (1, 2). As the diversity and sizes of such datasets expand, the utility of searching them with a novel sequence increases. Meta-metagenomic searches are currently underutilized. In principle such searches would involve direct access to sequence data from a large set of metagenomic experiments on a terabyte scale, along with software able to search for similarity to a query sequence. We find that neither of these aspects of meta-metagenomic searches is infeasible with current data transfer and processing speeds. In this communication, we report the results of searching the recently-described 2019-nCoV coronavirus sequence through a set of metagenomic datasets with the tag "virome".

Materials and Methods:

Computing Hardware: A Linux workstation used for the bulk analysis of metagenomic datasets employs an 8-core i7 Intel Microprocessor, 128G of Random Access Memory, 12TB of conventional disk storage, and 1TB of SSD storage. Additional analyses of individual alignments were carried out with standard consumer-grade computers.

Sequence data: All sequence data for this analysis were downloaded from the National Center for Biotechnology Information (NCBI) website, with individual sequences downloaded through a web interface and metagenomic datasets downloaded from the NCBI Sequence Read Archive (SRA) using the SRA-tools package (version 2.9.1). The latter sequence data were downloaded as .sra files using the prefetch tool, with extraction to readable format (.fasta.gz) using the NCBI fastq-dump tool. Each of these manipulations can fail some fraction of the time. Obtaining the sequences can fail due to network issues, while extraction in readable format occasionally fails for unknown reasons. Thus the workflow continually requests .sra files with ncbi-prefetch until at least some type of file is obtained, followed by attempts to unpack into .fasta.gz format until one such file is obtained from each .sra file. Metagenomic datasets for analysis were chosen through a keyword search of the SRA descriptions for "virome" and downloaded between January 27 and January 31, 2020. We note that the "virome" keyword search will certainly not capture every metagenomic dataset

with viral sequences, and likewise not capture every virus in the short sequence read archive.
With up to 16 threads running simultaneously, total download time (prefetch) was
approximately 2 days. Similar time was required for conversion to gzipped fasta files. A total
of 9014 sequence datasets were downloaded and converted to fasta.gz files. Most files
contained large numbers of reads with a small fraction of files containing very little data (only
a few reads or reads of at most a few base pairs). The total dataset consists of 2.5TB of
compressed sequence data corresponding to approximately 10^{13} bases.

Search Software: For rapid identification of close matches among large numbers of
metagenomic reads, we used a simple dictionary based on the 2019-nCoV sequence (NCBI
MN908947.3Wuhan-Hu-1) and its reverse complement, querying every 8th k-mer along the
individual reads for matches to the sequence. As a reference, and to benchmark the workflow
further, we included several additional sequences in the query (Vaccinia virus, an arbitrary
segment of a flu isolate, the full sequence of bacteriophage P4, and a number of putative
polinton sequences from *Caenorhabditis briggsae*). The relatively small group of k-mers
being queried ($<10^6$) allows a rapid search for homologs. This was implemented in a Python
script run using the PyPy accelerated interpreter. We stress that this is by no means the most
comprehensive or fastest search for large datasets. However, it is more than sufficient to
rapidly find any closely matching sequence (with the downloading and conversion of the data,
rather than the search, being rate limiting).

Alignment of reads to 2019-nCoV: Reads from the positive pangolin datasets were
adapter-trimmed with cutadapt (version 1.18) (3), and mapped to the 2019-nCoV genome
with BWA-MEM (version 0.7.12) (4) using default settings for paired-end mode. Alignments
were visualized with the Integrated Genomics Viewer IGV tool (version 2.4.10) (5).

Assessment of nucleotide similarity between 2019-nCoV, pangolin metavirome reads,
and closely related bat coronaviruses: All pangolin metavirome reads that aligned to the
2019-nCoV genome with BWA-MEM after adapter trimming with cutadapt were used for
calculation. The bat coronavirus genomes were aligned to the 2019-nCoV genome in a
multiple sequence alignment using the web-interface for Clustal Omega
(https://www.ebi.ac.uk/Tools/msa/clustalo/) (6) with default settings. We note that
sequence insertions with respect to the 2019-nCoV genome in either the pangolin metavirome
reads or the bat coronavirus genomes are not accounted for in the similarity traces shown in
Figure 1b.

Regional assessment of synonymous and nonsynonymous mutations (Figure S1):
Although the incomplete nature of coverage in the Pangolin metavirome data somewhat
limits the application of measures such as normalized dN/dS values, it remains possible to
identify regions with the strongest matches of this inferred viral sequence with the human
and bat homologs and to determine the distribution of synonymous and nonsynonymous
variants in these regions. Details of this analysis are presented in Figure S1.

Accessibility of Software: Scripts used for the observations described in this
communication are available at https://github.com/firelabsoftware/Metasearch2020.

Results:

To identify biological niches that might harbor viruses closely related to 2019-nCoV,
we searched through publicly available metaviromic datasets. We were most interested in
viruses with highly similar sequences, as these would likely be most useful in forming
hypotheses about the origin and pathology of the recent human virus. We thus set a
threshold requiring matching of a perfect 32-nucleotide segment with a granularity of 8
nucleotides in the search (i.e., interrogating the complete database of k-mers from the virus
with k-mers starting at nucleotide 1, 9, 17, 25, 33 of each read from the metagenomic data for
a perfect match). This would catch any perfect match of 39 nucleotides or greater, with some
homologies as short as 32 nucleotides captured depending on the precise phasing of the read.

All metagenomic datasets with the keyword "virome" in NCBI SRA as of January 2020
were selected for analysis in a process that required approximately 2 days each for
downloading and conversion to readable file formats and one day for searching by k-mer
match on a desktop workstation computer (i7 8-core). Together the datasets included
information from 9014 NCBI Short Read Archive entries with (in total) $6.2*10^{10}$ individual
reads and $8.4*10^{12}$ base pairs.   Despite the relatively large mass of data, the 32-nucleotide k-
mer match remains a stringent measure, with spurious matches to the ~30 kb 2019-nCoV
genome expected at only 1 in $3*10^{14}$. Positive matches among the metagenomic datasets
analyzed were relatively rare, with the vast majority of datasets (8994/9014 or 99.8%)
showing no matched 32-mers to 2019-nCoV. Of the datasets with matched k-mers, one was
from an apparent synthetic mixture of viral sequences, while the remaining were all from
vertebrate animal sources. The matches were from five studies: two bat-focused studies (7,
8), one bird-focused study (9), one small-animal-and-rodent focused study (10), and a study
of pangolins (11) [Table 1].

The abundance and homology of viruses within a metagenomic sample are of
considerable interest in interpreting possible characteristics of infection and relevance to the
query virus. From the quick k-mer search, an initial indicator could be inferred from the
number of matching reads and k-mer match counts for those reads [Table 1, Supplementary
Table 1]. For the 2019-nCoV matches amongst the available metagenomic datasets, the
strongest and most abundant matches in these analyses came from the pangolin lung
metaviromes. The matches were observed throughout the 2019-nCoV query sequence and
many of the matching reads showed numerous matching 32-mer sequences. The vast
majority of matches were in two lung samples with small numbers of matches in two
additional lung datasets (11). No matches were detected for five additional lung datasets, and
no matches were seen in eight spleen samples and a lymph node sample (11). Further
analysis of coverage and homology through alignment of the entire metagenomic datasets
revealed an extensive, if incomplete, coverage of the 2019-nCoV genome [Figure 1a]. Percent
nucleotide similarity can be calculated for pangolin metavirome reads aligning to 2019-nCoV
[Figure 1b], and these segmental homologies consistently showed strong matches,
approaching (but still overall weaker than) the similarity of the closest known bat coronavirus
(RaTG13).

Discussion:

Meta-meta-genomic searching can provide unique opportunities to understand the
distribution of nucleic acid sequences in diverse environmental niches. As metagenomic
datasets proliferate and as both the need and capability to identify pathogenic agents through
sequencing increase, meta-metagenomic searching may prove extremely useful in tracing the
origins and spreading of causative agents. In the example we present in this paper, such a
search identifies a number of niches with sequences matching the genome of the recent 2019-
nCoV virus. These analyses raise a number of relevant points for the origin of 2019-nCoV.
Before describing the details of these points, however, it is important to stress that while
environmental, clinical, and animal-based sequencing is valuable in understanding how
viruses traverse the animal ecosphere, static sequence distributions cannot be used to
construct a virus' full transmission history amongst different biological niches. So even were
the closest relative of a virus causing disease in species X to be found in species Y, we cannot
define the source of the outbreak, or the direction(s) of transmission. As some viruses may
move more than once between hosts, the sequence of a genome at any time may reflect a
history of selection and drift in several different host species. This point is also accentuated
in the microcosm of our searches for this work. When we originally obtained the 2019-nCoV
sequence from the posted work of Wu et al., we recapitulated their result that bat-SL-
CoVZC45 was the closest related sequence in NCBI's non-redundant (nr/nt) database. In our
screen of metavirome datasets, we observed several pangolin metavirome sequences—which
are not currently in the (nr/nt) database—and which are more closely related to 2019-nCoV
than bat-SL-CoVZC45. An assumption that the closest relative of a sequence identifies the
origin would at that point have transferred the extant model to zoonosis from pangolin
instead of bat. To complicate such a model, an additional study from Zhou et al. (12)
described a previously unpublished Coronavirus sequence, designated RaTG13 with much
stronger homology to 2019-nCoV than either bat-SL-CoVZC45 or the pangolin reads from Liu
et al (11). While this observation certainly shifts the discussion (legitimately) toward a
possible bat-borne intermediate in the chain leading to 2019-nCoV, it remains difficult to
determine if any of these are true intermediates in the chain of infectivity.

The match of 2019-nCoV to the pangolin coronavirus sequences also enables a link to
substantial context on the pangolin samples from Liu et al. (11), with information on the
source of the rescued animals (from smuggling activity), the nature of their deaths despite
rescue efforts, the potential presence of both other coronaviruses and other non-corona
viruses in the same cells, and the accompanying pathology. That work describes analyses
indicating several coronaviruses present in two of the pangolin lungs as well as other viral
species in those lungs. The pangolins appear to have died from lung-related illness, which
may have involved the 2019-nCoV closely-related virus. Notably, however, two of the
deceased pangolin lungs had much lower coronavirus signals, while five showed no signal,
with sequencing depths in the various lungs roughly comparable. Although it remains
possible that the 2019-nCoV-like coronavirus was the primary cause of death for these
animals, it is also possible (as noted by Liu et al. (11)) that the virus was simply present in the
tissue, with mortality due to another virus, a combination of infectious agents, or other
exposures.

During the course of this work, the homology between 2019-nCoV and pangolin
coronavirus sequences in a particular genomic subregion was also noted and discussed in an
online forum ("Virological.org") with some extremely valuable analyses and insights.

Matthew Wong and colleagues bring up the homology to the pangolin metagenomic dataset in this thread and appear to have encountered it through a more targeted search than ours (this study has since been posted online in bioRxiv) (13). As noted by Wong et al. (13), the Spike region includes a segment of ~200 bases where the inferred divergence between RaTG13 and 2019-nCoV dramatically increases. This region is of interest as the spike protein is a determinant of viral host range and under heavy selection (14). The observed Spike region divergence indeed includes a substantial set of nonsynonymous differences (Supplemental Figure 1). Notably, while reads from the pangolin lung dataset mapped to this region do not show a similar increase in variation relative to the human 2019-nCoV, we also did not observe a significant drop in variance between human and pangolin in this region. While Wong et al. concluded that recombination likely occurred in the spike region in the derivation of 2019-nCoV, definitive conclusions regarding the origins of 2019-nCoV are difficult given the limited sequencing data available and without consideration to altered evolutionary rates in different lineages (15). Thus alternative models for the observed sequence variation seem most parsimonious, including that of selection acting on the RaTG13 sequences in bat or another intermediate host resulting in a rapid variation of this highly critical virus-receptor interface.

The availability of numerous paths (both targeted and agnostic) toward identification of natural niches for pathogenic sequences will remain useful to the scientific community and to public health, as will vigorous sharing of ideas, data, and discussion of potential origins and modes of spread for epidemic pathogens.

Acknowledgments:
Competing interests: The authors declare that they have no competing interests;
Funding: This study was supported by the following programs, grants and fellowships: Human Frontier Science Program (HFSP) to DEJ, Arnold O. Beckman Award to MJS, Stanford Genomics Training Program (5T32HG000044-22; PI: M. Snyder) to MJM, and R35GM130366 to AZF. **Authors' contributions:** All authors contributed equally, and author order was selected randomly.

References:

1.   Bry L, Falk PG, Midtvedt T, Gordon JI. 1996. A model of host-microbial interactions in an open mammalian ecosystem. Science 273:1380-3.

2.   Bowman JS. 2018. Identification of Microbial Dark Matter in Antarctic Environments. Front Microbiol 9:3165.

3.   Marcel M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnetjournal 17:10-12.

4.   Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler transform. Bioinformatics 26:589-95.

5.   Thorvaldsdottir H, Robinson JT, Mesirov JP. 2013. Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. Brief Bioinform 14:178-92.

245   6.   Madeira F, Park YM, Lee J, Buso N, Gur T, Madhusoodanan N, Basutkar P, Tivey ARN,
246   Potter SC, Finn RD, Lopez R. 2019. The EMBL-EBI search and sequence analysis tools
247   APIs in 2019. Nucleic Acids Res 47:W636-W641.

248   7.   Hu D, Zhu C, Wang Y, Ai L, Yang L, Ye F, Ding C, Chen J, He B, Zhu J, Qian H, Xu W,
249   Feng Y, Tan W, Wang C. 2017. Virome analysis for identification of novel mammalian
250   viruses in bats from Southeast China. Sci Rep 7:10917.

251   8.   Yinda CK, Zeller M, Conceicao-Neto N, Maes P, Deboutte W, Beller L, Heylen E,
252   Ghogomu SM, Van Ranst M, Matthijnssens J. 2016. Novel highly divergent reassortant
253   bat rotaviruses in Cameroon, without evidence of zoonosis. Sci Rep 6:34209.

254   9.   Wille M, Eden JS, Shi M, Klaassen M, Hurt AC, Holmes EC. 2018. Virus-virus
255   interactions and host ecology are associated with RNA virome structure in wild birds.
256   Mol Ecol 27:5263-5278.

257   10.   Wu Z, Lu L, Du J, Yang L, Ren X, Liu B, Jiang J, Yang J, Dong J, Sun L, Zhu Y, Li Y,
258   Zheng D, Zhang C, Su H, Zheng Y, Zhou H, Zhu G, Li H, Chmura A, Yang F, Daszak P,
259   Wang J, Liu Q, Jin Q. 2018. Comparative analysis of rodent and small mammal
260   viromes to better understand the wildlife origin of emerging infectious diseases.
261   Microbiome 6:178.

262   11.   Liu P, Chen W, Chen JP. 2019. Viral Metagenomics Revealed Sendai Virus and
263   Coronavirus Infection of Malayan Pangolins (Manis javanica). Viruses 11:11.

264   12.   Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, Si HR, Zhu Y, Li B, Huang CL,
265   Chen HD, Chen J, Luo Y, Guo H, Jiang RD, Liu MQ, Chen Y, Shen XR, Wang X, Zheng
266   XS, Zhao K, Chen QJ, Deng F, Liu LL, Yan B, Zhan FX, Wang YY, Xiao GF, Shi ZL.
267   2020. A pneumonia outbreak associated with a new coronavirus of probable bat origin.
268   Nature doi:10.1038/s41586-020-2012-7.

269   13.   Wong MC, Javornik Cregeen SJ, Ajami NJ, Petrosino JF. 2020. Evidence of
270   recombination in coronaviruses implicating pangolin origins of nCoV-2019. bioRxiv
271   doi:10.1101/2020.02.07.939207:2020.02.07.939207.

272   14.   Li F. 2016. Structure, Function, and Evolution of Coronavirus Spike Proteins. Annu
273   Rev Virol 3:237-261.

274   15.   Paraskevis D, Kostaki EG, Magiorkinis G, Panayiotakopoulos G, Sourvinos G, S. T.
275   2020. Full-genome evolutionary analysis of the novel corona virus (2019-nCoV) rejects
276   the hypothesis of emergence as a result of a recent recombination event. Infection,
277   Genetics and Evolution 79:104212.

Description	File	TotalReads	TotalBases	HitReads	HitKmers
Pangolin Lung	SRR10168376_1	18067615	2710142250	5	9
Pangolin Lung	SRR10168376_2	18067615	2710142250	6	21
Pangolin Lung	SRR10168377_1	16414925	2462238750	308	955
Pangolin Lung	SRR10168377_2	16414925	2462238750	285	904
Pangolin Lung	SRR10168378_1	19045923	2856888450	91	352
Pangolin Lung	SRR10168378_2	19045923	2856888450	96	337
Pangolin Lung	SRR10168392_1	39738679	5960801850	2	4
Pangolin Lung	SRR10168392_2	39738679	5960801850	4	6
"Mock virome"	SRR3458564_2	10957589	1654595939	1	1
Bat Feces	SRR5040897_1	4020145	607041895	5	5
Bat Feces	SRR5040897_2	4020145	607041895	4	4
Bat Feces	SRR5040918_1	4804340	725455340	4	4
Bat Feces	SRR5040918_2	4804340	725455340	3	3
Virome analysis of Rodents a	SRR5343975_1	19775296	1582023680	1	1
Virome analysis of Rodents a	SRR5343977_1	20227246	1618179680	1	2
Bat	SRR5351751_2	363186	32644600	1	1
Bat	SRR5351752_1	305206	27328640	24	57
Bat	SRR5351752_2	305206	27608440	20	55
Bat	SRR5351758_2	301889	27514580	1	3
Bat	SRR5351760_1	788212	70634120	407	962
Bat	SRR5351760_2	788212	71244040	378	1006
Virome analysis of Rodents a	SRR5365807_1	21646498	1731719840	2	5
Virome analysis of Rodents a	SRR5365809_1	22799703	1823976240	1	1
Virome analysis of Rodents a	SRR5431767_1	10213803	1031594103	2	2
Virome analysis of Rodents a	SRR5447167_1	13849913	1398841213	2	2
Virome analysis of Rodents a	SRR5447174_1	14063346	1420397946	1	1
Virome analysis of Rodents a	SRR5447175_1	7510256	758535856	1	1
Avocet (RNA Virome in Wild E	SRR7239364_1	22336985	2233698500	2	2

Metagenomic datasets with k=32-mer matches to MN908947.3 2019-nCoV
Details of search and are described in legend to Table S1

279
280 **Table 1. Metagenomic datasets with k=32-mer matches to MN908947.3 (2019-nCoV).**
281 Details of search are described in legend to Table S1.

Figure 1. (a) Integrated Genomics Viewer (IGV) snapshot of alignment. Reads from the pangolin lung virome samples (SRR10168377, SRR10168378 and SRR10168376) were mapped to a 2019-nCoV reference sequence (Genbank accession: MN908947.3). The total number of aligned reads from the three samples was 1107, 313 and 32 reads respectively. **(b) Quantification of nucleotide-level similarity between the 2019-nCoV genome and pangolin lung metavirome reads aligning to the 2019-nCoV genome.** Average similarity was calculated in 101-nucleotide windows along the 2019-nCoV genome, and is only shown for those windows where each nucleotide in the window had coverage ≥ 2. Average nucleotide similarity calculated (in 101-nucleotide windows) between the 2019-nCoV genome and reference genomes of three relevant bat coronaviruses (bat-SL-CoVZC45: accession MG772933.1, bat-SL-CoVZXC21: accession MG772934.1, RaTG13: accession MN996532.1) is also shown. Note that the Pangolin metavirome similarity trace is not directly comparable to the bat coronavirus similarity traces, because the former uses read data for calculation whereas the latter use reference genomes.

**Figure S1
(Wahba et al., 2000)**

Supplemental Figure 1: Detailed plot of inferred substitutions. The plot shows incidence of inferred synonymous, nonsynonymous, and noncoding substitution from comparison of RaTG13 assembly [2] and a pangolin coronavirus scaffold, to the 2019-nCoV isolate from Wu et al. [1]. The pangolin scaffold was generated from a Blast alignment of 2019-nCoV mapping reads in SRR10168377. This plot addresses several challenges associated with the limited sequencing data available by attempting to provide the most favorable alignment of that sequence possible. To maximize sensitivity in detecting potential recombination, ambiguities in which two or more reads apparently disagreed (which were rare; approximately 1.2% of assigned bases) were resolved in favor of "no substitution" at any position if one read matches the 2019-nCoV genome. This will provide a lower bound of variation, although regions covered by a single read are still subject to amplification and sequencing error. Near-perfect overlaps between reads from SRR10168377 argue that such error is relatively low as agreement in those regions is 99.6%. The cumulative count of synonymous, nonsynonymous, and noncoding variants are per base pair and are taken over a 75 bp window (approximately 1/2 the read length) and then averaged over each 75 bp window with a weighting inversely proportional to distance. This results in the observed inverted spike for each individual variant.

TableS1 (Wahba et al. 2020)
(Graphic below shows an abbreviated segment of Table S1, Full Table at GitHub.com/FireLabSoftware/MetaSearch)

File	TotalReads	TotalBases	HitReads	HitKmers	MatchList
DRR023333_1	6243181	1667084289	1	17	X51522.1_phageP4:p8531a_m17
DRR027642_1	133932	74769093	0	0	
DRR027643_1	175545	89568995	0	0	
DRR053207_1	389643	46137596	0	0	
DRR053207_2	389643	46084317	0	0	
DRR053208_1	596864	124512374	0	0	
DRR053208_2	596864	126069168	0	0	
DRR053209_1	1476975	165712432	0	0	
-------------	-------------	-------------	-------------	-------------	-------------
<17374 Rows Not Displayed>					
-------------	-------------	-------------	-------------	-------------	-------------
SRR9843092_1	12545909	1267136809	0	0	
SRR9843092_2	12545909	1267136809	0	0	
SRR9843093_1	11515623	1163077923	0	0	
SRR9843093_2	11515623	1163077923	0	0	
SRR9843094_1	11313284	1142641684	4	11	X51522.1_phageP4:p7952s_m2 p8127a_m5 p8873a_m2(2)
SRR9843094_2	11313284	1142641684	6	35	X51522.1_phageP4:p7952s_m2 p8204a_m9 p8241a_m9 p8282a_m1 p8472s_m9
SRR9892957_1	10302064	1030206400	1	4	X51522.1_phageP4:p5128a_m4
SRR9892957_2	10302064	1030206400	1	6	X51522.1_phageP4:p5012s_m6
SRR9892958_1	10170129	1017012900	0	0	
SRR9892958_2	10170129	1017012900	0	0	
SRR9892959_1	10042853	1004285300	0	0	
SRR9892959_2	10042853	1004285300	0	0	
--Metagenome Set: All NCBI SRA sequences as of January 27-31 2020 with keyword virome.					
--Query Set:					
>MN908947.3W	Initial Deposit of 2019-nCoV				
>cb1_chrUn:103	Control 1: A polinton transposon from C. briggsae				
>cb1_chrUn:164	Control 2: A polinton transposon from C. briggsae				
>cb1_chrUn:564	Control 3: A polinton transposon from C. briggsae				
>X51522.1_phag	Control 4: A bacteriophage genome				
>AF250364.2_InfluenzaH1N1_neuraminidase Control 5: An arbitrary segment from an H1N1 Flu isolate					
>AY603355.1_Vaccinia Control 6: An full vaccinia genome (note that inspection of the vaccinia hits seems to indicate that some or all hits are fr					
--Output Format					
Column 1: Dataset_ReadNumber (e.g. SRRXXX_2 is the read 2 file from SRXXX)					
Column 2: Number of Reads extracted and analyzed					
Column 3: Number of bases analzyed					
Column 4: Number of hits from the viral sampler query					
Column 5: Total number of hit k-mers					
Column 6: Summary of hits. Each element has a query name (e.g. X51522.1_phageP4), a position (e.g. p5012 starts at position 5012); a strand ('					
--Query Code: Jazz18Heap.py version AG					
Parameters of Jazz18Heap call were as follows					
Include=<Virus_Sampler_File> (fasta file of the above seven sequences)					
Data=*.fasta.gz					

ReportGranularity=0 (provides somewhat limited output reporting)					
MutiTask=16 (uses 16 threads-- this was used to provide multitasking on a system with 16 threads [8 cores=16threads]					
Klen=32					
SearchGranularity=8					
## Jazz18Heap.py version AG					
## This program is a fast metasearch that will look for instances of sequence reads matching					
## a reference in at least one k-mer.					
## Jazz18Heap is intended to look for evidence of matches to a relatively short reference sequence					
## e.g., less than 100KB, but probably workable up to several MB in a large number of high throughput sequencing experiments					
## Jazz18Heap is intended for finding relatively rare sequence (not common ones)					
## Jazz18Heap doesn't substitute for many tools to align and track coverage. It's main goal is rapid identification of potentially homologous se					
## Inputs are as follows (command line, Key=Value syntax)					
## ReferenceFile = <FastA file with list of sequences to match k-mers from>					
## ExcludeFile = <FastA file with sequences that will be excluded from matches>					
## DataFiles = <List of .fastq files, .fasta files, or NCBA-SRA accessions [e.g, SRR####]					
## Lists are comma delimited with no spaces in list					
## .fasta or .fastq files can be gzip compressed, although this somewhat slows the program down					
## .fasta and .fasta.gz files must have exactly one sequence per line (no multiline sequences)					
## For .fasta files to be downloaded from NCBI-SRA archive this entails the command line parameter --fasta 0					
## * Wildcards are allowed here as well, or list of files in a file with the extension .files					
## Providing a directory here will search all files in this directory or subdirectory for fasta and fastq data files					
## Optional Parameters (willdefault to reasonable values if not set)					
## OutFileBase = <Character String to Label Output Files with>					
## ReportGranularity = <How many reads to process before reporting hit numbers (default is 1Million)>					
## SearchGranularity = <How much distance between k-mers to be examined in each data read					
## setting SearchGranularity=1 makes Jass18Heap look at every k-mer in every read					
## setting SearchGranularity=8 makes Jass18Heap look at every 8th k-mer in every read					
## higher numbers may miss a few hits but can greatly improve speed.					
## setting to a large number (999999) ensures only one k-mer will be looked up per read					
## SearchOffset = <Where to start jumping through each read for potential k-mers (zero means start at first base)					
## fastqdump = <Where to look for the fasterq-dump binary [program will look for this but if it is not found, the full path to fasterq-dump bina					
## klen = <How long are the k-mers used>. Default klen = 32					
## snpAllow = <Set to True to allow a single mismatch in each k-mer (default is False)					
## Circular = <Set to True to force every Reference sequence to be treated as a circle>					
## Default : Uses the FastA name line-- if this line contains "Circular", the sequence is treated as a circle					
## Multithreading: Jazz18Heap has the very primative multitasking ability to spawn a number					
## of derivative processes for a large number of data files to be scanned. To use 16 Threads					
## Set MultiThreading=16 in the command line.					
## Output:					
## Output consists of a log file with information on the run and a FastA file.					
## FastA files have information about each read in the ID line					
## ID Structure:					
## DataFileName;LineNumberInDataFile;ReferenceFileName;ReferenceSequenceName;NumberOfMatchedKmers					
## Running the program:					
## Jazz18Heap only runs at full speed with a variant of Python (PyPY) that included a just-in-time compiler					
## Syntax Jazz18Heap RefFile=<MyRefFile> DataFiles=MyFastA1.fasta,MyFastA2.fasta.gz,MyFastQ*.fastq <Other_Parameters>					