Short title: ONAC127 and ONAC129 regulate rice grain filling

A Heat Stress Responsive NAC Transcription Factor Heterodimer Plays Key

Roles in Rice Caryopsis Filling

Ye Ren, Zhouquan Huang, Hao Jiang, Zhuo Wang, Fengsheng Wu, Yufei Xiong,

Jialing Yao [*]

College of Life Science and Technology, Huazhong Agricultural University, Wuhan

430070, China.

Corresponding author: Jialing Yao, Ph.D. and Professor, College of Life Science and

Technology, Huazhong Agricultural University, Wuhan 430070, China.

[*] Address correspondence to yaojlmy@mail.hzau.edu.cn

Telephone: 86-27-87282866

Fax: 86-27-87282866

ABSTRACT

High temperature usually leads to the failure of grain filling during caryopsis development, resulting in the loss of yield, however the mechanism is not yet well elucidated. Here, we report that two rice caryopsis-specific NAM/ATAF/CUC domain transcription factors, *ONAC127* and *ONAC129*, respond to heat stress and are involved in the caryopsis filling process. *ONAC127* and *ONAC129* are dominantly expressed in pericarp during grain filling and can form a heterodimer. To investigate the functions of these two ONACs, we obtained CRISPR/cas9 induced mutants and overexpression lines of them. Interestingly, we found that both knock-out and overexpression plants showed incompletely filling and shrunken phenotype of caryopses, which became more severe under heat stress. The shrunken caryopses of these transgenic lines are usually with ectopic accumulation of starch in the pericarp. Transcriptome analyses revealed that *ONAC127* and *ONAC129* mainly regulate stimulus response, cell wall construction and nutrient transport etc. ChIP-seq analyses identified the direct targets of ONAC127 and ONAC129 in developing caryopses, including monosaccharide transporter *OsMST6*, sugar transporter *OsSWEET4*, calmodulin-like protein *OsMSR2* and Ethylene-Response AP2/ERF Factor *OsEATB*. The result suggested that *ONAC127* and *ONAC129* might regulate the caryopsis filling through sugar transportation and abiotic stress responses. Overall, this study demonstrates the transcriptional regulatory networks involving ONAC127 and ONAC129, which coordinates multiple pathways to modulate caryopsis development and heat stress response at rice filling stage.

INTRODUCTION

Rice caryopsis development starts from the fertilized ovary and finishes with a dehydrated, hard and transparent grain. This process can be divided into three stages by landmark events: cell division, organogenesis and maturation. (Agarwal et al., 2011). A rice caryopsis is mainly composed of endosperm filled with starch grains, and the biosynthesis of starch is closely related to carbohydrates transportation. Carbohydrates are synthesized in leaves and delivered to caryopses via phloem (Patrick, 1997), then unloaded to the pericarp as the energy and materials for starch biosynthesis (Zhang et al., 2007). The dorsal vascular bundle passes through the pericarp is the main nutrient transport tissue in caryopsis (Oparka and Gates, 1981). While the vascular bundles are not contiguous with endosperm tissue (Hoshikawa, 1984), the apoplasmic pathway is the only way for nutrient to reach the starchy endosperm. (Matsuda et al., 1979).

Carbohydrates may enter the nucellar epidermis directly through plasmodesmata, then be transported to the apoplasmic space with sugar transporter protein OsSWEETs (sugar will eventually be exported transporters) and partially hydrolyzed into monosaccharide by cell wall invertase OsCINs. The monosaccharide is transported into the aleurone layer mainly by monosaccharide transporter OsMSTs while the rest un-hydrolyzed sucrose is directly transported into the aleurone layer via sucrose transporter OsSUTs (Yang et al., 2018). The nutrient transportation is also regulated by a range of transcription factors such as *OsNF-YB1* and *OsNF-YC12*, which activate the expression of OsSUTs to capture the leaked sucrose in apoplasmic space. Knockout of *NF-YB1* or *OsNF-YC12* leads to defective grains with chalky endosperms and similar phenotype has been observed in mutants of *OsSUT1* (Bai et al., 2016; Xiong et al., 2019).

The nutrient transport processes in caryopsis are susceptible to changing environmental conditions, extreme external stimuli (especially the extreme temperature) during grain-filling stage will lead to a reduction of nearly 50% in rice (Hu and Xiong, 2014). Plants respond to the unexpected high temperature through some stress-specific signaling pathways. During heat stress signal transduction, the heat shock transcription

factors (HSFs) will be activated to regulate the expression of downstream heat shock

proteins (HSPs) and other stress-related genes. For example, AP2/EREBP transcription

factor *DREB2A* can activate a heat shock transcription factor *hsfA3* under heat stress to

induce the expression of specific HSPs (Schramm et al., 2008). Similarly, the

expression of *OsbZIP60*, which play important roles in moisture retention and heat-

damage resistance, is induced under heat stress and endoplasmic reticulum stress (Oono

et al., 2010). Although several plant heat stress response related proteins have been

found and elucidated, the mechanisms of the response are still poorly understood.

NAC (NAM, ATAF1/2, CUC2) transcription factors are one of the largest family

of plant specific transcription factors with 151 members in rice (Nuruzzaman et al.,

2010). The typical structure of NAC transcription factors is a conserved NAC domain

with about 150 amino acids in N-terminus, followed by a variable transcriptional

regulation region (TRR) in C-terminus (Christianson et al., 2010). The NAC domain

can be further subdivided into five subdomains [A-E], subdomains C and D are highly

conserved and may mainly act in DNA-binding (Ooka et al., 2003). In addition, some

subdomain D contains a hydrophobic NAC repression domain (NARD) which can

suppress the transcriptional activity of NAC transcription factors by suppressing their

DNA-binding ability or nuclear localization ability. The specific functional motif in

NARD, LVFY can suppress not only the activity of NAC transcription factors, but also

the other transcription factors by its hydrophobicity (Hao et al., 2010).

NAC transcription factors are involved in many biological processes in plants, such

as organ development, secondary wall synthesis, stress response and so on. For instance,

NAC29 and *NAC31* regulate downstream cellulose synthase (CESA) by activating the

downstream transcription factor *MYB61* to control the synthesis of secondary wall

(Huang et al., 2015). For stress response, especially in abiotic ways, the NAC

transcription factors are master regulators in this process. *RD26* gene is the first NAC

transcription factor identified as a regulator in mediating abscisic acid (ABA) and

jasmonate (JA) signaling during stress responses in Arabidopsis (Fujita et al., 2004).

SNAC1 is a stress-responsive NAC transcription factor confers rice drought tolerance by closing stomata (Hu et al., 2006) and *SNAC3* confers heat and drought tolerance through regulation of reactive oxygen species (Fang et al., 2015). *OsNAC2* is also reported involving in drought stress response mediated by ABA (Shen et al., 2017). Meanwhile, it plays divergent roles in different biological processes, such as regulating shoot branching (Mao et al., 2007), affecting plant height and flowering time through mediating the gibberellic acid (GA) pathway (Chen et al., 2015). *VNI2* in Arabidopsis is a bifunctional NAC transcription factor reported as a transcriptional repressor functioning in xylem vessel formation (Yamaguchi et al., 2010), but the transcriptional activator activity of *VNI2* is induced by high salinity stress to regulate the leaf longevity (Yang et al., 2011).

Nine caryopsis-specific NAC transcription factors were found through transcriptome analyses in rice (Mathew et al., 2016).We selected four of these genes, *ONAC025*, *ONAC127*, *ONAC128* and *ONAC129* for further analyses, which located in the linked regions of chromosome 11 and were identified as a gene cluster (Fang et al., 2008). As recent study implied two NAC transcription factors specifically expressed in maize seeds, *ZmNAC128* and *ZmNAC130*, playing critical roles in starch and protein accumulation in seeds filling (Zhang et al., 2019). We inferred that the four NAC transcriptional factors might also be involved in some important processes in rice caryopsis development. In this study, we selected *ONAC127* and *ONAC129* from the four genes, which could form a heterodimer and participate in apoplasmic transportation as well as heat stress response to regulate rice caryopsis filling. These findings broadened the understanding of the regulation network of stress response and grain filling in rice.

RESULTS

***ONAC127* and *ONAC129* are specifically expressed in caryopsis**

To understand these four genes basically, we firstly checked the expression data of

ONAC025, *ONAC127*, *ONAC128* and *ONAC129* in the microarray database CREP

(http://crep.ncpgr.cn) (Wang et al., 2010). The data showed that the expression level of

ONAC025 in caryopsis was significantly higher than that of the rest genes, while that

of *ONAC128* was the lowest among the four genes. For *ONAC127* and *ONAC129*, they

showed the highest expression level at 7 d after pollination (DAP) and then decreased

gradually, such similar expression pattern attracted our further attention.

For validating the spatial and temporal expression patterns and further exploring

the specific expression distribution of *ONAC127* and *ONAC129* in caryopsis, we

mechanically isolated rice caryopses at 5-14 DAP to the mixture of pericarp and

aleurone layer, starch endosperm and embryo for qRT-PCR analyses by the method

described by Bai (Bai et al., 2016). The aleurone layer specifically expressed gene

oleosin and the starch endosperm specifically expressed gene *SDBE* were used as

markers (Ishimaru et al., 2015) (Supplemental Fig. S1). The result showed that

ONAC127 and *ONAC129* were mainly expressed in pericarp or aleurone layer while

they were weakly expressed in starch endosperm, and the expression went to the

summit at 5 DAP during the whole development process (Fig. 1A).

The mRNA *in situ* hybridization analyses using the unripe caryopses of ZH11

turned out that *ONAC127* and *ONAC129* were dominantly expressed in pericarp, when

the expression level in aleurone layer and starch endosperm were quite weak (Fig. 1B).

Histochemical GUS activity was also detected using the transgenic plants expressing

pONAC127::GUS and pONAC129::GUS, the result was almost identical to that of *in*

situ hybridization analyses, that *ONAC127* and *ONAC129* were preferentially

expressed in pericarp of caryopsis (Fig. 1C). Considering that 5 DAP is a key time point

of caryopsis development for the organogenesis is finished and maturation stage is just

initiated (Agarwal et al., 2011). *ONAC127* and *ONAC129* might be involved in some

substance exchange processes playing vital roles in caryopsis filling and maturation.

We then carried out the assay through transforming green fluorescent protein (GFP)

fused ONAC127 or ONAC129 protein into rice protoplast transiently to investigate the

subcellular localization pattern of the two genes. 35S::GFP was used as a positive

control and 35S::Ghd7-CFP was used as a nuclear marker (Xue et al., 2008). The

fluorescence generated by ONAC127-GFP and ONAC129-GFP was distributed in the

nucleus and cytoplasm just like the way positive control did (Fig. 1D). For further

confirming the subcellular localization, transgenic plants expressing pUbi::ONAC127-

GFP and pUbi::ONAC129-GFP were generated in Zhonghua11 (ZH11) background.

The subcellular localization pattern of ONAC127 and ONAC129 were validated by

detecting the fluorescence from the roots of two-week-old rice seedlings of the plants,

ONAC127 and ONAC129 proteins were indeed localized in the nucleus and cytoplasm

(Supplemental Fig. S2).

Fig. 1 Spatial and temporal expression patterns of *ONAC127* and *ONAC129*.

A. qRT-PCR analyses of *ONAC127* and *ONAC129* in different tissue. AL, aleurone layer; Pe, pericarp;

SE, starchy endosperm; Em, embryo; 1-21D, caryopses collected during 1-21 DAP. The values in each

column are the mean of three technical replicates and the error bars indicate the ±SD. Ubiquitin is used

as the reference gene. B. *In situ* hybridization of sectioned caryopses collected at 1, 3, 5, 7 and 10 DAP

using anti-sense and sense probes. Nu, nucellar; Pe, pericarp; SE, starchy endosperm; Em, embryo; AL,

aleurone layer. Scale bars are 50 μm. C. Histochemical GUS activity detection using pONAC127::GUS

and pONAC129::GUS. Pe, pericarp; SE, starchy endosperm; Em, embryo. Scale bar is 1 mm. D.

Subcellular localization of the *ONAC127* and *ONAC129* in rice protoplasts. 35S::Ghd7-CFP is used as a

nuclear marker. Scale bars are 5 μm.

ONAC127 and ONAC129 can form a heterodimer

It was noticed that the NAC transcription factors usually function as a dimer (Olsen

et al., 2005), thus we investigated if there was any interaction between ONAC127 and

ONAC129. The result indicated that ONAC127 could interact with ONAC129 in yeast

(Fig. 2A). To confirm the interaction between ONAC127 and ONAC129, the *in vitro*

GST pull-down assay was carried out. His-tagged ONAC127 and GST-tagged

ONAC129 were expressed respectively and incubated together. The protein mixture of

GST and ONAC127-His was used as a negative control. After purification by

Glutathione Agarose, ONAC127-His was detected in the sample containing

ONAC129-GST instead of the control with His-tag antibody (Fig. 2B). The *in vivo*

bimolecular fluorescence complementation (BiFC) assay was also performed with rice

protoplasts. Nuclear homodimer OsbZIP63 was chosen as positive control (Walter et

al., 2004), and 35S::Ghd7-CFP was used as a nuclear marker. Yellow fluorescence

generated from the interaction between ONAC127-YFP^N and ONAC129-YFP^C was

detected, which confirmed the heterodimer ONAC127 and ONAC129 formed in

nucleus (Fig. 2C). In view of previous studies, some of NAC transcription factors were

located in the cytoplasm or plasma membrane first, and then were imported to nuclei

under some specific conditions (Fang et al., 2014). Considering the subcellular

localization pattern of ONAC127 and ONAC129 (Fig. 1D), we speculated that these

two proteins might also exert their transcriptional regulation functions in this way.

Fig. 2 Interaction between ONAC127 & ONAC129.

A. The yeast two-hybrid assay. The full-length cDNA of ONAC127 and ONAC129 are both cloned into

pGBKT7 (BD) and pGADT7 (AD). The transformants are grown on [SD/–Trp /–Leu] and [SD/–Trp/–

Leu/–His/–Ade] plates with 10^0, 10^{-1} and 10^{-2} fold dilution. B. The pull-down assays showing that there

is a direct interaction between ONAC127-His and ONAC129-GST and *in vitro*. C. The BiFC assays of

ONAC127 and ONAC129. ONAC127-YFP^N and ONAC129-YFP^C interact to form a functional YFP in

rice protoplasts. 35S::Ghd7-CFP is used as a nuclear marker. Scale bars are 5 μm.

ONAC127 and *ONAC129* are key players in starch accumulation during rice

caryopsis filling

We obtained the specifically knockout *ONAC127* or *ONAC129* mutants (*onac127*

and *onac129*) in ZH11 background by CRISPR/Cas9 genome editing system (Ma et al.,

2015). Considering that these two proteins could form a heterodimer, double knockout

mutants (*onac127;129*) were also obtained. The sgRNA target sites were designed at

the exons of *ONAC127* and *ONAC129* using the web-based tool CRISPR-P (Liu et al.,

2017). There are two target sites of each gene, which were expected to generate

different mutations in the coding region of the genes (Fig. 3A). The grains of three T_0

homozygous plants of each mutant were selected for generating independent T_1

transgenic lines. The sequencing data of the transgenic lines was decoded by the method

described by Liu (Liu et al., 2015), and the genotypes of the homozygous lines were

showed in Supplemental Fig. S3. On the other hand, we also generated the

overexpression lines pUbi::ONAC127-FLAG and pUbi::ONAC129-FLAG (OX127

and OX129) (Fig. 3B). Relative expression level of the overexpression lines was

analyzed by qRT-PCR (Supplemental Fig. S4).

During the rice reproductive stage, we found the development of caryopses in

ONAC127 and *ONAC129* transgenic lines arrested during 5-7 DAP (Supplemental Fig.

S5), resulting in shrunken and incompletely filled grains after ripening (Fig. 3C, D). As

we mentioned before, the expression level of *ONAC127* and *ONAC129* was very high

during 5-7 DAP (Fig. 1A), the functions of *ONAC127* and *ONAC129* seem to be closely

related to grain filling, especially at the early stage of maturation. Since the phenotype

is illogical, we detected the expression level of the two ONACs in the transgenic lines

immediately. It turned out that the expression of *ONAC127* and *ONAC129* was up-

regulated in the overexpression lines and down-regulated in the mutants indeed

(Supplemental Fig. S6). We speculated that ONAC127 and ONAC129 may regulate

several different pathways contributing to caryopsis filling at the same time. The

expression of the two ONACs may need to be maintained at a steady level, any

fluctuations in their expression would interfere the normal running of different

pathways and have the chance to bring caryopsis filling defection.

To investigate the reason of the shrunken phenotype further, histological analyses

through resin embedded sections were prepared with 7 DAP incompletely filling

caryopses. In ZH11, starch was stored transiently in the pericarp during the early stage,

the degradation of starch in the pericarp correlated with starch accumulation in the

endosperm (Wu et al., 2016). As we expected, the starch in the endosperm of ZH11

caryopses was arrayed neatly, densely and hardly found in pericarp (Fig. 3E). In

contrast, many starch grains accumulated in the pericarp of the mutants and the

overexpression lines of *ONAC127* and *ONAC129*, especially in the double mutant

onac127;129, in which substantial starch grains were hardly seen in endosperm (Fig.

3E). The finding above suggested that *ONAC127* and *ONAC129* might play critical

roles in starch translocation and mobilization towards the developing endosperm.

Fig. 3 The construction of the CRISPR mutants and overexpression plants and their phenotypes.

A. Schematic diagram of the sgRNA target site in genomic region of *ONAC127* and *ONAC129*. The

target sequences of *onac127*, *onac129* and *onac127;129* are listed under the diagram and marked as a

colorful triangle. The Cas9 splice sites are marked as red triangle and the protospacer-adjacent motifs

are shown in orange. B. Schematic diagram of the construction of OX127 & OX129. C. Phenotypes of

grains of the overexpression lines and mutants of *ONAC127* and *ONAC129*. Scale bars are 2 mm. D.

Phenotypes of panicles at maturation stage. The panicles of transgenic plants are almost upright while

the panicles of ZH11 are bent due to the grain filling. Scale bars are 20 cm. E. Accumulation of starch

in the pericarp of the caryopsis of overexpression lines and mutants relative to ZH11. Cross sections of

caryopses stained with toluidine blue at 7DAP. Pe, pericarp; SE, starchy endosperm. Scale bar is 40

μm.

ONAC127 and ONAC129 are involved in heat stress response

With the measurement of agronomic traits, the transgenic lines of *ONAC127* and

ONAC129 showed significant reduction in both seed set rate (percentage of fully filled

seeds after harvest) and 1000-grains weight (Fig. 4A, B). The percentage of shrunken

grains showed an obvious increase while proportion of blighted grains (glumes with no

seed in it) did not change significantly in these transgenic lines (Fig. 4C). It proved that

the reduction of seed set rate and yield was mainly caused by the increase of shrunken

grains.

Notably, the transgenic plants during filling stages were suffered from severe heat

stress in the summer of 2018 (Fig. 4D, E). Previous studies have shown that the ambient

temperature higher than 35℃ would have a serious impact on the yield at flowering

and grain filling stages (Hakata et al., 2017). Therefore, we set 35℃ as the heat damage

temperature of rice (T_B), and calculated the heat damage accumulated temperature per

hour (TH_i), heat damage accumulated temperature during filling stage (T_S) and heat

damage hours during filling stage (H_S) as described by Chen (Chen et al., 2019). Given

that *ONAC127* and *ONAC129* function during early filling stage, we also calculated the

heat damage accumulated temperature during 0-7 DAP (T_{S7}) and heat damage hours

during 0-7 DAP (H_{S7}). Noting that we cultivated two batches of rice in the summer of

2018, the first batch of plants were flowering on about July 20 when the second batch

goes on about August 20. It's obvious that T_S and H_S of the first batch were much higher

than that of the second batch, especially at the early development stage of caryopsis

(Fig. 4D, E).Therefore, the caryopses of the plants cultivated in the first batch were

defined as suffering from heat stress while those in the second batch were defined as

planted under normal conditions with little or no heat stress. We found that heat stress

caused sharp increase of shrunken grains compared to normal conditions, and such

increase in *onac127;129* was more dramatic than the other transgenic lines (Fig. 4C),

which led to significant decrease of seed set rate and 1000-grains weight (Fig. 4A, B).

Meanwhile, the expression level of *ONAC127* and *ONAC129* was highly increased

under heat stress (Supplemental Fig. S6), which suggested that *ONAC127* and

ONAC129 might be involved in heat stress response during caryopsis filling stage.

Fig. 4 Agronomic traits analyses and ambient temperature.

A. Seed set rate. The ratio of full grains to all spikelets on a panicle. B. 1,000 grains weight. Only the

fully filled grains were counted and the data was calculated based on 100 grains. C. The composition of

the seed set of spikelets. Data in (A), (B) and (C) are meant (±SD) from three independent lines, each

of which are meant from three biological replicates. Significant differences between the ZH11 and the

transgenic lines were determined using Student's t-test (* P<0.05, ** P<0.01). D and E. The ambient

temperature of growing areas during rice caryopsis filling stage. 35°C is marked as yellow dotted lines

and 40°C is marked as red dotted lines. T_S, heat damage accumulated temperature during whole filling

stage; H_S, heat damage hours during whole filling stage; T_{S7}, heat damage accumulated temperature

during 0-7 DAP; H_{S7}, heat damage hours during 0-7 DAP.

ONAC129 negatively regulates ONAC127 transcriptional activity

For deeper exploring the relationship of *ONAC127* and *ONAC129*, we performed

a dual luciferase assay in rice protoplasts as the transcriptional activity of the two genes

was not detected in yeast assays (Fig. 2A). *ONAC127* and *ONAC129* were fused with

the yeast GAL4 binding domain (GAL4BD) as effectors to be co-transformed into rice

protoplasts with a firefly luciferase reporter gene driven by a CaMV35S promotor (35S-

GAL4-fLUC; Fig. 5A). A significant increase in the transcriptional activation activity

of ONAC127 and ONAC129 was detected, while the transcriptional activity of

ONAC127 was significantly restrained when ONAC129 was co-transformed into rice

protoplasts (Fig. 5B). These results hinted that ONAC129 might regulate the

transcriptional activity of ONAC127 negatively and we wonder if ONAC129 could

regulate the expression of *ONAC127*. However, we didn't find significant fluctuation

of the expression of *ONAC127* in the transgenic lines of *ONAC129* and vice versa.

(Supplemental Fig. S6).

As we know, several NAC transcription factors contain both transcriptional

activation domain and NAC repression domain (NARD), which means that the

combinatorial effects of both transcriptional regulation activities determine

downstream events (Hao et al., 2010). Hence, we decided to BLAST and analyze the

protein sequences of ONAC127 and ONAC129 through the method described by Hao,

and we found that there were NARD-like sequences in NAC subdomain D indeed in

both genes (Fig. 5C), which suggested that *ONAC127* and *ONAC129* might be

bifunctional transcription factors that could activate or repress downstream genes in

different situation.

Fig. 5 ONAC129 negatively regulates ONAC127 transcriptional activity.

A. Scheme of the constructs used in the rice protoplast cotransfection assay. B. Data of Dual-LUC

assay. The reporter and internal control are co-transformed with each experimental group. The

fLUC/rLUC ratio represents the relative activity of CaMV35S promoter. The values in each column are

the mean (±SD) of three replicates. Significant differences are determined using Student's t-test (**

P<0.01). C. Schematic diagram of the domains of ONAC127 & ONAC129. The color legends

indicating the domains are shown at the bottom of the figure.

ONAC127 and ONAC129 regulate the transcription of genes related to sugar

transportation and abiotic stimulus

RNA-Seq analyses worked in identifying differentially expressed genes. The 7

DAP caryopses of the overexpression lines and mutants of *ONAC127* and *ONAC129*

under heat stress and normal conditions were used to generate RNA-seq libraries. There

are 7056, 6485, 5765, and 1608 differentially expressed genes (DEGHs) identified in

onac127 (*onac127*H), *onac129* (*onac129*H), OX127 (OX127H), OX129 (OX129H)

respectively under heat stress, compared with ZH11 (ZH11H), and 15334, 13284, 1776,

2041 differentially expressed genes (DEGNs) which were identified in *onac127*

(*onac127*N), *onac129* (*onac129*N), OX127 (OX127N), OX129 (OX129N) respectively

under normal conditions compared with ZH11 (ZH11N) (Fig. 6A, B; Supplemental

Dataset S1). Gene Ontology (GO) enrichment analyses was performed to examine the

biological roles of DEGs in caryopsis development. The significantly enriched GO

terms were mainly associated with stimulus response, transcriptional activity regulation,

signal transduction, cell wall construction and substance transportation (Fig. 6C).

Subsequently, the genes directly bound by ONAC127 and ONAC129 in caryopsis

were identified using Chromatin Immunoprecipition Sequencing assays (ChIP-seq).

The ChIP assays were performed using the anti-FLAG antibody with 7 DAP caryopses

of OX127 (OX127H), OX129 (OX129H) under heat stress, and OX127 (OX127N),

OX129 (OX129N) under normal condition. The expression of the ONAC127-FLAG

and ONAC129-FLAG fusion proteins was verified by western blot analyses to validate

the effectiveness of the FLAG tags (Supplemental Fig. S7). After sequencing, 185, 6125,

220, 455 peaks were finally obtained respectively in OX127H, OX127N, OX129H and

OX129N (Supplemental Dataset S2), and 54, 5092, 82, 212 putative ONAC127 and

ONAC129 bound genes were identified in OX127H, OX127N, OX129H and OX129N

respectively among the peaks (Fig. 6D). Then, we identified the binding motifs of the

ONACs using MEME (Machanick and Bailey, 2011), and found that the significantly

enriched motif was 'CT(C)TTCT(C)TT' (Fig. 6E, F), in line with 'TT(A/C/G)CTT',

the specific motif of transmembrane NAC transcription factors *NTL6* and *NTL8*

(Lindemose et al., 2014). Both the two genes were associated with stimulus response,

implying that *ONAC127* and *ONAC129* were probably involved in stress response.

Notably, the motif 'CT(C)TTCT(C)TT' was only found significantly in OX127H and

OX129H. There was no significantly enriched motif in OX127N and OX129N,

implying that the target genes of ONAC127 and ONAC129 might be more specific

under heat stress.

For further exploring the target genes regulated by ONAC127 and ONAC129, we

defined the genes which were either consistently bound by ONAC127/129 or

simultaneously bound by ONAC127 and ONAC129 under heat stress or normal

conditions as ONAC127/129 preferable-bound genes (Fig. 6D, red trapezoid). Based

on the data from RNA-Seq analyses and phenotype of transgenic lines of ONAC127

and ONAC129, 8 genes with respect to sugar transportation or abiotic stress response

were selected from 136 preferable-bound genes as the potential target genes (PTGs) of

ONAC127 and ONAC129 for further investigation.

Fig. 6 Data analyses of RNA-Seq and ChIP-Seq.

A and B. Venn diagram showing the number of genes regulated by *ONAC127* and *ONAC129* based on

the RNA-seq analyses. Differentially expressed genes are defined as |Fold Change|≥2 and *p*-

value<0.05. *p*-values were calculated using the Fisher's exact test. C. A selection of enriched gene

ontology (GO) terms of the differentially expressed genes. GO terms with FDR <0.05 were kept. The

length of the bars represents the negative logarithm (base 10) of FDR. D. Venn diagram showing the

number of ONAC127 and ONAC129 direct target genes. The red trapezoid indicates the ONAC127

and ONAC129 preferable bound genes. E and F. Motif analyses of ONAC127 and ONAC129 binding

381   peaks. The E-value is the enrichment P-value multiplied by the number of candidate motifs tested.

ONAC127 and ONAC129 play a pivotal role in caryopsis filling through

regulating key factors directly in substance accumulation and stress response

To validate the interactions between ONAC127/129 and promotors of their

potential target genes (PTGs), the yeast one-hybrid assays were performed. It suggested

that ONAC127 and ONAC129 could bind to the promoter sequences of *OsEATB*,

OsHCI1, *OsMSR2*, *OsSWEET4*, *bHLH144* and *OsMST6* directly in yeast (Fig. 7A). The

Dual-LUC assays in rice protoplasts were performed to figure out whether ONAC127

and ONAC129 affect the transcription of these genes directly. ONAC127 and

ONAC129 were co-transformed into rice protoplasts with a firefly luciferase reporter

gene driven by target gene promotors (Fig. 7B). It turned out that ONAC127 activated

OsMSR2 and *OsMST6* promotors significantly, and both ONAC127 and ONAC129

repressed the *OsEATB* and *OsSWEET4* promotors strongly *in vivo*. It is noteworthy that

ONAC129 could suppress the transcriptional activation activity of ONAC127 to

OsMSR2 and *OsMST6* promotors, while ONAC129 barely regulated the transcription

of *OsMSR2* and *OsMST6* alone (Fig. 7C).

For testing whether the endogenous ONAC127 and ONAC129 bound to the target

genes specifically, ChIP-qPCR was performed using the same rice materials of ChIP-

Seq. The result indicated that ONAC127 and ONAC129 bound to the promoters of

OsEATB, *OsMSR2*, *OsSWEET4* and *OsMST6* respectively under heat stress and normal

conditions (Fig. 7D, E, F, G). The qRT-PCR for these genes was also performed in the

7 DAP caryopses of transgenic lines of *ONAC127* and *ONAC129*, showing that the

expression of *OsEATB* and *OsSWEET4* was significantly up-regulated in the mutants

and generally down-regulated in the overexpression lines. The expression of *OsMSR2*

and *OsMST6* was both up-regulated in *onac129* and down-regulated in *onac127*

(Supplemental Fig. S8). The results above indicated that *OsEATB*, *OsMSR2*,

OsSWEET4 and *OsMST6* were the direct targets of ONAC127 and ONAC129 in rice

during caryopsis filling.

Fig. 7 Identification and validation of ONAC127 and ONAC129 direct target genes in rice.

A. The interaction between ONAC127/129 and the promoters of PTGs as determined by yeast one-

hybrid analyses. The transformants were grown on [SD/–Leu /–His] plates with 3-AT which

concentrations were 50mM, 100mM and 150mM respectively. B. Scheme of the constructs used in the

rice protoplast cotransfection assay. C. Data of Dual-LUC assay. The fLUC/rLUC ratio represents the

relative activity of target gene promoters. D-G. ChIP-qPCR verification of ONAC127 and ONAC129

binding regions. The values in each column are the mean (±SD) of three replicates. Significant

differences were determined using Student's t-test (* P<0.05; ** P<0.01).

DISCUSSION

In this study, two rice caryopsis-specific NAC transcription factors *ONAC127* and

ONAC129 were identified, which can form a heterodimer and are predominantly

expressed during the early and middle stage of rice caryopsis development (Fig. 1, 2).

The caryopses of the transgenic plants were obviously unfilled (Fig. 3C, D, E;

Supplemental Fig. S5), and the proportion of shrunken grains was higher under natural

heat stress (Fig. 4C). By protein−DNA binding assays, we found that ONAC127 and

ONAC129 may regulate the expression of the sugar transporters, *OsMST6* and

OsSWEET4 directly. Both the two genes are involved in the transportation of

photosynthate from dorsal vascular bundles to endosperm during caryopsis filling

(Wang et al., 2008; Sosso et al., 2015). ONAC127 and ONAC129 also bind to the

promotors of *OsMSR2* and *OsEATB* directly (Fig. 7), which participate in abiotic stress

responses by responding to calcium ion (Ca^{2+}) or plant hormones (Qi et al., 2011; Xu

et al., 2011). These findings suggest that *ONAC127* and *ONAC129* may be involved in

multiple pathways, therefore interfere the abiotic stress response and caryopsis filling

process during rice reproductive stage.

ONAC127 and ONAC129 are involved in the apoplasmic transportation of

photosynthates as a balancer

At the early stage of caryopsis development, some of the photosynthate from the

dorsal vascular bundle was transported to endosperm through the apoplasmic space

formed by degenerative nucellar cell wall, the others synthesized the starch grains in

mesocarp cells (Hoshikawa, 1984). The starch accumulation in pericarp will reach a

maximum at 5 DAP. After that, the starch in mesocarp cells will be disintegrated and

the nutrient will be transported to the endosperm due to the continuous process of

endosperm cells filling. The cell contents of pericarp cells will disappear, and the cells

will eventually dehydrate and die, leaving only the cuticularized cell wall remnants (Wu

et al., 2016).

Since the abnormal expression of *ONAC127* and *ONAC129*, endosperm cells in

abnormal caryopses may not obtain enough photosynthate for filling due to the

functional defect of apoplasmic transport pathways. The photosynthates accumulates in

pericarp as starch grains without been disintegrated, thus stop the endosperm cells from

proliferation or filling by starch, eventually leading to incompletely filled and shrunken

caryopsis (Fig. 3C, E).

It is noteworthy that the target transporters *OsSWEET4* and *OsMST6* are

predominantly expressed at early stages of caryopsis development, and may act in

downstream of the cell wall invertase OsCIN2 to import hexose into aleurone layer

through apoplasmic space (Wang et al., 2008; Sosso et al., 2015; Yang et al., 2018). As

previously mentioned, ONAC127 and ONAC129 suppress the expression of

OsSWEET4 directly (Fig. 7C; Supplemental Fig. S8), and *ossweet4-1*, the mutant of

OsSWEET4, has a distinct phenotype with shrunken grains similar with the abnormal

caryopses in transgenic lines of *ONAC127* and *ONAC129* (Sosso et al., 2015). The fact

confirmed our hypothesis that *ONAC127* and *ONAC129* are involved in the apoplasmic

transport pathways directly by regulating the expression of transporters indeed.

Moreover, the shrunken caryopses we observed in mutants and overexpression

lines of *ONAC127* and *ONAC129* are also similar with the incompletely filled

caryopses of the double mutants *ossweet11;15*, in which two apoplasmic transporters

OsSWEET11 and OsSWEET15 were dysfunction (Yang et al., 2018). For that

OsSWEET15 has been excluded from the target genes by the yeast one-hybrid assays

(Fig. 7A), we detected the expression of *OsSWEET11* and *OsSWEET15* in transgenic

lines of *ONAC127* and *ONAC129* by qRT-PCR and found that both genes might be

activated by ONAC127 and ONAC129 indirectly (Supplemental Fig. S9). The fact

suggested that ONAC127 and ONAC129 might activate the expression of

OsSWEET11/15 and suppress the expression of *OsSWEET4* simultaneously. This may

explain the fact that the phenotypes of mutants and overexpression lines of *ONAC127*

and *ONAC129* are almost the same in incompletely filled caryopses (Fig. 3C, D, E;

Supplemental Fig. S5). Hence, we speculate that *ONAC127* and *ONAC129* may act as

a balancer in apoplasmic transportation by activating or suppressing the expression of

transporters dynamically (Fig. 8), therefore keeping the amount of several different

apoplasmic transporters at different stages of caryopsis development.

Fig. 8 Schematic diagram of the working model of ONAC127 and ONAC129 in rice caryopsis.

ONAC127 and ONAC129 regulate the expression of *OsMST6* and *OsSWEET4* directly to participate in

the apoplasmic transportation, and OsSWEET11 and OsSWEET15 may be also involved in this

process as the indirect downstream transporters of them. When heat stress occurs, ONAC127 and

ONAC129 may be involved in the response to heat stress during the filling stage by regulating the

expression of *OsMSR2* and *OsEATB* directly, while *OsHCI1* and *OsHSP101* are regulated indirectly.

ONAC127 and ONAC129 may act as the balancer of apoplasmic transportation and the core

transcription factors in the stress response during the early and middle stages of rice caryopsis

development.

ONAC129 suppresses the transcriptional activity of ONAC127 in some pathways,

meanwhile, ONAC127 and ONAC129 may regulate downstream genes differently at

dimer state and monomer state (Fig. 5B, 7C). We conjecture that the proportion of

ONAC127/129 dimers may be vital in adjusting the balance of the amount of

apoplasmic transporters at different stages of caryopsis development. Whether to

knockout or to over-express *ONAC127* or *ONAC129*, the proportion of the monomers

and dimers of ONAC127/129 will be changed inevitably, which induces various

changes in expression of the apoplasmic transporters, thus the apoplasmic

transportation defects, leading to the incomplete filling caryopsis phenotype (Fig. 3C).

In macroscopic view, the balance regulation of apoplasmic transportation may be a

"rate-limiting step" of rice reproductive growth to ensure that the photosynthate is

enough for both the filling of grains and the energy for plants to avoid premature aging,

thus may be vital for plants to keep the balance of vegetative growth and reproductive

growth.

ONAC127 and ONAC129 respond to heat stress during caryopsis filling

Heat stress has profound effects on plant growth, particularly in reproductive

development. When plants encounter heat stress, the rise of cytosolic Ca^{2+}

concentration is the most rapid response (Liu et al., 2003). With the increase of Ca^{2+}

concentration, the expression of calmodulins (CaMs) and Calmodulin-like genes

(CMLs) will be activated and modulate the phosphorylation state of HSFs to regulate

their DNA binding ability, thereby regulating the expression of heat stress related genes

including HSPs and some plant hormone responsive genes (Li et al., 2018).

The proportion of incompletely filled caryopses under heat stress is much higher

than that of normal condition (Fig. 4C), and the expression of *ONAC127* and *ONAC129*

was induced by heat stress (Supplemental Fig. S6), suggesting that heat stress might

interfere the dimerization of ONAC127 and ONAC129 to affect the balance of

apoplastic transportation. As previously mentioned, ONAC127 and ONAC129 regulate

the expression of *OsMSR2* and *OsEATB* directly (Fig. 6C; Supplemental Fig. S8).

Previous study showed that *OsMSR2*, which can be strongly induced by heat, cold and

drought stress, is a calmodulin-like protein working by binding Ca^{2+} and speculated

having the function as a Ca^{2+} sensor in plant cells (Xu et al., 2011). *OsEATB* is a rice

AP2/ERF gene which can suppress gibberellic acid (GA) synthesis while the GA

signaling pathway involved in the regulation of cell elongation and morphogenesis

under heat stress (Koini et al., 2009; Qi et al., 2011). We hereby speculated that

ONAC127 and *ONAC129* might participate in heat stress response by regulating the

Ca^{2+} and GA signaling pathway.

The expression level of *OsHSP101* and *OsHCI1* was changed significantly in the

transgenic lines of *ONAC127* and *ONAC129* (Fig. 7A, C; Supplemental Fig. S9). We

believe that they are also the potential target genes of ONAC127 and ONAC129, though

they may not be regulated directly. RING E3 ligase gene *OsHCI1* is induced by heat

stress and mediates nuclear–cytoplasmic trafficking of nuclear substrate proteins via

mono-ubiquitination to improve the heat tolerance under heat shock (Lim et al., 2013).

Meanwhile, OsHSP101 function as one of the important molecular chaperones that

interact with OsHSA32 and OsHsfA2c, the latter of which is one of the central

regulators of heat stress response (Singh et al., 2012; Lin et al., 2014). Accordingly, we

believe that *ONAC127* and *ONAC129* are involved in some core reactions in the heat

stress response during rice caryopsis development stage.

Ca^{2+} is one of the most important second messengers in heat stress response,
cytosolic Ca^{2+} concentration would change rapidly in a short time when heat stress
occurs (Li et al., 2018). The stress signal would be then transmitted by CaMs/CMLs
like OsMSR2 to influence downstream gene expression consequently. When the
downstream HSFs are activated, HSPs including OsHSP101 would be induced to
express and function, and some nuclear-cytoplasmic trafficking reactions involving
OsHCI1 would occur (Singh et al., 2012; Lim et al., 2013). Besides, a variety of plant
hormones including GA would act in heat stress response to maintain plant growth, and
OsEATB plays an important role in GA-mediated plant cell elongation (Qi et al., 2011;
Li et al., 2018). Considering that all these direct or indirect target genes of *ONAC127*
and *ONAC129* play vital roles in the key steps of the heat stress response regulatory
network, we speculate that ONAC127 and ONAC129 may be the core transcription
factors in heat stress response during rice caryopsis filling stage, and the stress and
hormone response may be involved in the complex regulatory network of apoplasmic
transportation.

MATERIALS AND METHODS

Generation of transgenic plants and growth conditions

For generating CRISPR mutants, we used the web-based tool CRISPR-P v2.0
(http://cbi.hzau.edu.cn/CRISPR2/) designed by Liu (Liu et al., 2017) to get the specific
gRNA cassettes targeting *ONAC127* and *ONAC129* and then cloned into a binary vector
pYLCRISPR/Cas9-MH (Ma et al., 2015). For overexpression plants, we amplified the
stop-code-less cDNA fragments of *ONAC127* and *ONAC129*, fused 3×Flag or EGFP
coding sequence at the 3' end. The fused sequences were cloned into the binary vector
pCAMBIA1301U (driven by a maize ubiquitin promotor). For tissue specific
expression analyses, a 2000bp fragment of the 5' upstream region of *ONAC127*, and a
2043bp fragment of the 5' upstream region of *ONAC129* were cloned into the binary
vector pDX2181G (with GUS, β-glucuronidase).These recombinant constructs were

introduced into rice Zhonghua11 (ZH11; *Oryza sativa* ssp. *japonica)* by *Agrobacterium tumefaciens* (EH105)-mediated transformation (Lin and Zhang, 2005). The plant materials were cultivated in paddy fields in Huazhong Agricultural University, Wuhan, China, under natural long-day conditions (approximately 12-14 h light/10-12 h dark) during May to October 2018. The temperature of growing areas during rice reproductive stage was showed in (Fig. 4D, E). We set 35℃ as the heat damage temperature of rice (T_B), and calculated the heat damage accumulated temperature per hour (TH_i), $TH_i = \begin{cases} T_i - T_B & T_i \geq T_B \\ 0 & T_i < T_B \end{cases}$ (T_i is the ambient temperature at *i* hour); heat damage hours during filling stage (H_S) , $H_s = \sum_{i=0}^{n} H_i$ $(H_i = \begin{cases} 1 & T_i \geq T_B \\ 0 & T_i < T_B \end{cases})$ and heat damage accumulated temperature during filling stage (T_S), $T_S = \sum_{i=0}^{n} TH_i$ using the method described by Chen (Chen et al., 2019). The heat damage accumulated temperature during 0-7 DAP (T_{S7}) and heat damage hours during 0-7 DAP (H_{S7}) was also calculated as the method above. The phenotypes were detected in homozygous T_1 generation of transgenic plants. The primer sequences used in this study were listed in Supplemental Table S1 and would not be repeated in the following article.

Histochemical GUS staining

Unripe caryopses at 5 and 7 DAP from pONAC127::GUS and pONAC129::GUS transgenic plants were collected for the GUS staining assays following the methods described by Yang (Yang et al., 2018).

Microscopy analyses

For semithin section microscopy analyses, the unripe caryopses at 7 DAP were collected in 2.5% glutaraldehyde for fixation, vacuum infiltrated on ice for 30 min and incubated at 4℃ for 24h. The plastic embedding and sectioning were performed as described by Wang (Wang et al., 2008). Slides were stained with toluidine blue and detected by a BX53 microscope (Olympus).

RNA *in situ* hybridization

Unripe caryopses (1-10 DAP) from rice ZH11 were collected for paraffin embedding. Paraffin section was performed according to the previous method (Xiong et al., 2019). Gene-specific fragments of *ONAC127* and *ONAC129* were amplified by the primers used in RT-PCR and cloned into the pGM-T vector. The probes were synthesized using the DIG RNA labeling kit (SP6/T7) (Roche) according to the manufacturer's recommendations. RNA hybridization and immunologic detection of the hybridized probes were performed on sections as described by Kouchi (Kouchi and Hata, 1993). Slides were observed using a BX53 microscope (Olympus).

Transient expression assays

To investigate the subcellular localization of ONAC127 and ONAC129, the CDS of these genes were cloned into the pM999-35S vector, the nuclear located gene *Ghd7* was used as a nuclear localization marker (Xue et al., 2008).For the BiFC assays, the CDS of *ONAC127* and *ONAC129* were cloned into the vectors pVYNE and pVYCE (Waadt et al., 2008) respectively. Plasmids were extracted and purified using the Plasmid Midi Kit (QIAGEN) and were then transformed into rice protoplasts according to the procedure described by Shen (Shen et al., 2017). The fluorescent signals were detected with a confocal laser scanning microscope (TCS SP8, Leica).

RNA isolation and transcript analyses

Total RNA was isolated using the TRIzol method (Invitrogen), and the first strand cDNA was synthesized using HiScript II Reverse Transcriptase (Vazyme). The real-time PCR was performed on QuantStudio 7 Flex Real-Time PCR System (Applied Biosystems), with the $2^{-\Delta\Delta Ct}$ method for relative quantification (Livak and Schmittgen, 2001). The significance of differences was estimated using Student's t-test. Relevant primers were designed according to qPrimerDB (https://biodb.swu.edu.cn/qprimerdb) (Lu et al., 2018).

Yeast assays

For yeast two-hybrid assay, the CDS of *ONAC127* and *ONAC129* were cloned into

the pGBKT7 and pGADT7 vector (Clontech). The fusion plasmids were transformed

into yeast strain AH109 or Y187. The pGBKT7-53 was co-transformed with pGADT7-

T as a positive control. For yeast one-hybrid assay, DNA fragments about 1000bp

corresponding to the promoters of target genes were independently inserted into the

pHisi-1 plasmid (Clontech). *ONAC127* and *ONAC129* were fused to GAL4

transcriptional activation domain (pGAD424). These constructs were transformed into

the yeast strain YM4271. Yeast two-hybrid assay and one-hybrid assay were performed

following the manufacturer's instructions (Clontech).

Chromatin immunoprecipitation (ChIP)

Overexpression lines of *ONAC127* and *ONAC129* (fused with 3×FLAG) were used

for ChIP-Seq analyses. We verified the expression of target proteins by western blot

using ANTI-FLAG M2 Monoclonal Antibody (Sigma-Aldrich). ChIP assay was

performed with ANTI-FLAG M2 Magnetic Beads (Sigma-Aldrich) as the method

described previously (Xiong et al., 2019). For each library, three independent replicated

samples were prepared. The immunoprecipitated DNA and input DNA were subjected

to being sequenced on the HiSeq 2000 platform (Illumina), which was processed by the

Novogene Corporation.

ChIP-Seq raw sequencing data was mapped to the rice reference genome (RGAP

ver. 7.0, http://rice.plantbiology.msu.edu) (Kawahara et al., 2013) using BWA (Li and

Durbin, 2009). MACS2 (Zhang et al., 2008) was used for peak calling and the peaks

were identified as significantly enriched (corrected P-value <0.05) in the IP libraries

compared with input DNA. Visual analyses were performed using IGV (Intergative

Genomics Viewer, v2.3.26) (Robinson et al., 2011). Motif enrichment analyses were

performed by MEME (Machanick and Bailey, 2011) with default parameters.

To validate the specific targets genes, the immunoprecipitated DNA and input DNA

was applied for ChIP-qPCR analyses. The enrichment value was normalized to the

input sample. The significance of differences was estimated using Student's t-test.

RNA-Seq

Total RNA was extracted from unripe caryopses at 7 DAP. For each library, three

independent replicated RNA samples were prepared. The RNA samples were then

sequenced on the HiSeq 2000 platform (Illumina), which was processed by the

Novogene Corporation.

The raw reads were filtered for adaptors and low-quality reads. Clean reads were

mapped to the reference genome of rice (RGAP v. 7.0) using HISAT2 (v.2.0.5) (Kim et

al., 2015). The gene expression level was calculated by FPKM method (Trapnell et al.,

2010), and the differentially expressed genes (|Fold Change| $\geq$ 2 and p-value<0.05)

were selected by DESeq2 software (Love et al., 2014). GOseq (Young et al., 2010) was

used for GO enrichment analyses, and p-value was converted to -$\log_{10}(p$-value) for

display.

In vitro **GST pull-down assay**

The CDS of *ONAC127* and *ONAC129* were cloned into pET28a and pGEX-4-1

vectors respectively for His-tagged ONAC127 and GST-tagged ONAC129 protein

expression *in vitro*. The fusion plasmids were transformed into *Escherichia coli* BL21

strain. The GST pull-down assays were performed according to the previous methods

(Xiong et al., 2019). The protein was separated on a 10% SDS-PAGE gel and further

analyzed by immunoblotting using anti-His and anti-GST antibody (Sigma-Aldrich).

Dual Luciferase Transcriptional Activity Assay

Full-length cDNAs of *ONAC127* and *ONAC129* were cloned to yeast GAL4

binding domain vectors (GAL4BD) and "None" as effectors. The 35S-GAL4-fLUC and

190-fLUC were used as reporters, and AtUbi::rLUC was used as an internal control.

The constructed plasmids were purified and transformed into rice protoplasts with the

procedure mentioned previously. Dual Luciferase Reporter Assay System (Promega)

was used to measure the luciferase activity by Tecan Infinite M200 (Tecan).

Agronomic traits analyses

Harvested ripe rice grains were air dried and stored at room temperature for at least

2 months before measuring. Only full grains were used for measuring 1000-grain

weight which was calculated based on 100 grains. The seed set rate/composition was

calculated based on the grains from 3-5 panicles (include the full, shrunken and blighted

grains). All measurements of the positive transgenic plants were undertaken using three

independent lines.

Accession numbers

The sequence data from this paper can be found in the RGAP database

(http://rice.plantbiology.msu.edu) under the following accession numbers: *ONAC025*

(LOC_Os11g31330), *ONAC127* (LOC_Os11g31340), *ONAC128* (LOC_Os11g31360),

ONAC129 (LOC_Os11g31380), *OsMST6* (LOC_Os07g37320), *OsSWEET4*

(LOC_Os02g19820), *OsEATB* (LOC_Os09g28440), *OsMSR2* (LOC_Os01g72530),

bHLH144 (LOC_Os04g35010), *OsHCI1* (LOC_Os10g30850), *HSP101*

(LOC_Os05g44340), *OsSWEET11* (LOC_Os08g42350), *OsSWEET15*

(LOC_Os02g30910), *OsbZIP63* (LOC_Os07g48820). The RNA-seq and ChIP-seq

data are deposited in the NCBI Gene Expression Omnibus (Edgar et al., 2002) with

accession number GSE140167.

Supplemental Material

Supplemental Fig. S1 Expression level of mechanical isolation marker in isolated

tissue.

Supplemental Fig. S2 Subcellular localization of the ONAC127 and ONAC129 in the

roots of two-week-old rice seedlings of pUbi::ONAC127-GFP and pUbi::ONAC129-
GFP transgenic plants.

Supplemental Fig. S3 Mutation sites in *onac127*, *onac129* and *onac127;129* lines, as
compared with wild-type (ZH11) sequences.

Supplemental Fig. S4 The relative expression level of the overexpression lines of
ONAC127 and *ONAC129*.

Supplemental Fig. S5 Different stages of caryopses development.

Supplemental Fig. S6 Expression level of *ONAC127* and *ONAC129* in 7 DAP
caryopses of transgenic lines compared with that of ZH11.

Supplemental Fig. S7 Detection of FLAG fusion proteins in the ZH11 and
overexpression lines.

Supplemental Fig. S8 Expression level of the target genes of ONAC127 and
ONAC129 in 7 DAP caryopses of transgenic lines compared with that of ZH11.

Supplemental Fig. S9 Expression level of the indirect target genes of ONAC127 and
ONAC129 in 7 DAP caryopses of transgenic lines compared with that of ZH11.

Supplemental Table S1   Primers used in this study.

Supplemental Dataset S1   Differentially expressed genes.

Supplemental Dataset S2   Binding sites identified by ChIP-seq.

ACKNOWLEDGEMENTS

We thank Prof. Min Chen, Prof. Honghong Hu, Prof. Yidan Ouyang and Ms.
Wenjing Guo for helping revise the manuscript. This research was supported by grants
from the National Natural Science Foundation of China (no. 31570321). The funders
had no role in the study design, data collection and analysis, the decision to publish, or
in the preparation of the manuscript.

Author contributions

Y.R. and J.Y. conceived the original research plans; Y.R. carried out most of the experiments; Z.H., Z.W., F.W. and Y.X. provided technical assistance to Y.R.; Y.R. and H.J. analyzed the data; Y.R. conceived the project and wrote the article with contributions of all the authors; J.Y. supervised and complemented the writing.

LITERATURE CITED

Agarwal P, Kapoor S, Tyagi AK (2011) Transcription factors regulating the progression of monocot and dicot seed development. Bioessays **33:** 189-202

Bai AN, Lu XD, Li DQ, Liu JX, Liu CM (2016) NF-YB1-regulated expression of sucrose transporters in aleurone facilitates sugar loading to rice endosperm. Cell Res **26:** 384-388

Chen S, Zhang T, Huang Q, Chen M, Zhao L, Zhang Y (2019) Effects of High Temperature Injury on Seed Setting Rate from Heading to Milky Stage of Early Season Rice in Hainan. China Tropical Agriculture: 65-69 (in Chinese with English abstract)

Chen X, Lu S, Wang Y, Zhang X, Lv B, Luo L, Xi D, Shen J, Ma H, Ming F (2015) OsNAC2 encoding a NAC transcription factor that affects plant height through mediating the gibberellic acid pathway in rice. Plant J **82:** 302-314

Christianson JA, Dennis ES, Llewellyn DJ, Wilson IW (2010) ATAF NAC transcription factors: regulators of plant stress signaling. Plant Signal Behav **5:** 428-432

Edgar R, Domrachev M, Lash AE (2002) Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. Nucleic Acids Res **30:** 207-210

Fang Y, Liao K, Du H, Xu Y, Song H, Li X, Xiong L (2015) A stress-responsive NAC transcription factor SNAC3 confers heat and drought tolerance through modulation of reactive oxygen species in rice. J Exp Bot **66:** 6803-6817

Fang Y, Xie K, Xiong L (2014) Conserved miR164-targeted NAC genes negatively regulate drought resistance in rice. J Exp Bot **65:** 2119-2135

Fang Y, You J, Xie K, Xie W, Xiong L (2008) Systematic sequence analysis and identification of tissue-specific or stress-responsive genes of NAC transcription factor family in rice. Mol Genet Genomics **280:** 547-563

Fujita M, Fujita Y, Maruyama K, Seki M, Hiratsu K, Ohme-Takagi M, Tran LS, Yamaguchi-Shinozaki K, Shinozaki K (2004) A dehydration-induced NAC protein, RD26, is involved in a novel ABA-dependent stress-signaling pathway. Plant J **39:** 863-876

Hakata M, Wada H, Masumoto-Kubo C, Tanaka R, Sato H, Morita S (2017) Development of a new heat tolerance assay system for rice spikelet sterility. Plant Methods **13:** 34

Hao YJ, Song QX, Chen HW, Zou HF, Wei W, Kang XS, Ma B, Zhang WK, Zhang JS, Chen SY (2010) Plant NAC-type transcription factor proteins contain a NARD domain for repression of transcriptional activation. Planta **232:** 1033-1043

Hoshikawa K (1984) Development of Endosperm Tissue with Special Reference to the Translocation of Reserve Substances in Cereals : III. Translocation pathways in rice

endosperm. Japanese journal of crop science **53:** 153-162

Hu H, Dai M, Yao J, Xiao B, Li X, Zhang Q, Xiong L (2006) Overexpressing a NAM, ATAF, and CUC (NAC) transcription factor enhances drought resistance and salt tolerance in rice. Proc Natl Acad Sci U S A **103:** 12987-12992

Hu H, Xiong L (2014) Genetic engineering and breeding of drought-resistant crops. Annu Rev Plant Biol **65:** 715-741

Huang D, Wang S, Zhang B, Shang-Guan K, Shi Y, Zhang D, Liu X, Wu K, Xu Z, Fu X, Zhou Y (2015) A Gibberellin-Mediated DELLA-NAC Signaling Cascade Regulates Cellulose Synthesis in Rice. Plant Cell **27:** 1681-1696

Ishimaru T, Ida M, Hirose S, Shimamura S, Masumura T, Nishizawa NK, Nakazono M, Kondo M (2015) Laser microdissection-based gene expression analysis in the aleurone layer and starchy endosperm of developing rice caryopses in the early storage phase. Rice (N Y) **8:** 57

Kawahara Y, de la Bastide M, Hamilton JP, Kanamori H, McCombie WR, Ouyang S, Schwartz DC, Tanaka T, Wu J, Zhou S, Childs KL, Davidson RM, Lin H, Quesada-Ocampo L, Vaillancourt B, Sakai H, Lee SS, Kim J, Numa H, Itoh T, Buell CR, Matsumoto T (2013) Improvement of the Oryza sativa Nipponbare reference genome using next generation sequence and optical map data. Rice (N Y) **6:** 4

Kim D, Langmead B, Salzberg SL (2015) HISAT: a fast spliced aligner with low memory requirements. Nat Methods **12:** 357-360

Koini MA, Alvey L, Allen T, Tilley CA, Harberd NP, Whitelam GC, Franklin KA (2009) High temperature-mediated adaptations in plant architecture require the bHLH transcription factor PIF4. Curr Biol **19:** 408-413

Kouchi H, Hata S (1993) Isolation and characterization of novel nodulin cDNAs representing genes expressed at early stages of soybean nodule development. Mol Gen Genet **238:** 106-119

Li B, Gao K, Ren H, Tang W (2018) Molecular mechanisms governing plant responses to high temperatures. J Integr Plant Biol **60:** 757-779

Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics **25:** 1754-1760

Lim SD, Cho HY, Park YC, Ham DJ, Lee JK, Jang CS (2013) The rice RING finger E3 ligase, OsHCI1, drives nuclear export of multiple substrate proteins and its heterogeneous overexpression enhances acquired thermotolerance. J Exp Bot **64:** 2899-2914

Lin MY, Chai KH, Ko SS, Kuang LY, Lur HS, Charng YY (2014) A positive feedback loop between HEAT SHOCK PROTEIN101 and HEAT STRESS-ASSOCIATED 32-KD PROTEIN modulates long-term acquired thermotolerance illustrating diverse heat stress responses in rice varieties. Plant Physiol **164:** 2045-2053

Lin YJ, Zhang Q (2005) Optimising the tissue culture conditions for high efficiency transformation of indica rice. Plant Cell Rep **23:** 540-547

Lindemose S, Jensen MK, Van de Velde J, O'Shea C, Heyndrickx KS, Workman CT, Vandepoele K, Skriver K, De Masi F (2014) A DNA-binding-site landscape and regulatory network analysis for NAC transcription factors in Arabidopsis thaliana. Nucleic Acids Res **42:** 7681-7693

Liu H, Ding Y, Zhou Y, Jin W, Xie K, Chen LL (2017) CRISPR-P 2.0: An Improved CRISPR-Cas9
Tool for Genome Editing in Plants. Mol Plant **10:** 530-532

Liu HT, Li B, Shang ZL, Li XZ, Mu RL, Sun DY, Zhou RG (2003) Calmodulin is involved in heat
shock signal transduction in wheat. Plant Physiol **132:** 1186-1195

Liu W, Xie X, Ma X, Li J, Chen J, Liu YG (2015) DSDecode: A Web-Based Tool for Decoding of
Sequencing Chromatograms for Genotyping of Targeted Mutations. Mol Plant **8:** 1431-
1433

Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods **25:** 402-408

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biology **15**

**Lu K, Li T, He J, Chang W, Zhang R, Liu M, Yu M, Fan Y, Ma J, Sun W, Qu C, Liu L, Li N, Liang
Y, Wang R, Qian W, Tang Z, Xu X, Lei B, Zhang K, Li J** (2018) qPrimerDB: a
thermodynamics-based gene-specific qPCR primer database for 147 organisms. Nucleic
Acids Res **46:** D1229-D1236

**Ma X, Zhang Q, Zhu Q, Liu W, Chen Y, Qiu R, Wang B, Yang Z, Li H, Lin Y, Xie Y, Shen R, Chen
S, Wang Z, Chen Y, Guo J, Chen L, Zhao X, Dong Z, Liu YG** (2015) A Robust CRISPR/Cas9
System for Convenient, High-Efficiency Multiplex Genome Editing in Monocot and Dicot
Plants. Mol Plant **8:** 1274-1284

Machanick P, Bailey TL (2011) MEME-ChIP: motif analysis of large DNA datasets. Bioinformatics
27: 1696-1697

Mao C, Ding W, Wu Y, Yu J, He X, Shou H, Wu P (2007) Overexpression of a NAC-domain
protein promotes shoot branching in rice. New Phytol **176:** 288-298

Mathew IE, Das S, Mahto A, Agarwal P (2016) Three Rice NAC Transcription Factors
Heteromerize and Are Associated with Seed Size. Front Plant Sci **7:** 1638

Matsuda T, Kawahara H, Chonan N (1979) Histo-Cytological Researches on Translocation and
Ripening in Rice Ovary : I. Histological changes and transfer pathways in the developing
ovary. Japanese journal of crop science **48:** 155-162

Nuruzzaman M, Manimekalai R, Sharoni AM, Satoh K, Kondoh H, Ooka H, Kikuchi S (2010)
Genome-wide analysis of NAC transcription factor family in rice. Gene **465:** 30-44

Olsen AN, Ernst HA, Leggio LL, Skriver K (2005) DNA-binding specificity and molecular functions
of NAC transcription factors. Plant Science **169:** 785-797

**Ooka H, Satoh K, Doi K, Nagata T, Otomo Y, Murakami K, Matsubara K, Osato N, Kawai J,
Carninci P, Hayashizaki Y, Suzuki K, Kojima K, Takahara Y, Yamamoto K, Kikuchi S**
(2003) Comprehensive analysis of NAC family genes in Oryza sativa and Arabidopsis
thaliana. DNA Res **10:** 239-247

Oono Y, Wakasa Y, Hirose S, Yang L, Sakuta C, Takaiwa F (2010) Analysis of ER stress in
developing rice endosperm accumulating beta-amyloid peptide. Plant Biotechnol J **8:**
691-718

Oparka KJ, Gates P (1981) Transport of assimilates in the developing caryopsis of rice (Oryza
sativa L.) : The pathways of water and assimilated carbon. Planta **152:** 388-396

Patrick JW (1997) PHLOEM UNLOADING: Sieve Element Unloading and Post-Sieve Element

Transport. Annu Rev Plant Physiol Plant Mol Biol **48**: 191-222

Qi W, Sun F, Wang Q, Chen M, Huang Y, Feng YQ, Luo X, Yang J (2011) Rice ethylene-response

AP2/ERF factor OsEATB restricts internode elongation by down-regulating a gibberellin

biosynthetic gene. Plant Physiol **157**: 216-228

Robinson JT, Thorvaldsdottir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP (2011)

Integrative genomics viewer. Nat Biotechnol **29**: 24-26

Schramm F, Larkindale J, Kiehlmann E, Ganguli A, Englich G, Vierling E, von Koskull-Doring P

(2008) A cascade of transcription factor DREB2A and heat stress transcription factor HsfA3

regulates the heat stress response of Arabidopsis. Plant J **53**: 264-274

Shen J, Liu J, Xie K, Xing F, Xiong F, Xiao J, Li X, Xiong L (2017) Translational repression by a

miniature inverted-repeat transposable element in the 3' untranslated region. Nat

Commun **8**: 14651

Shen J, Lv B, Luo L, He J, Mao C, Xi D, Ming F (2017) The NAC-type transcription factor OsNAC2

regulates ABA-dependent genes and abiotic stress tolerance in rice. Sci Rep **7**: 40641

Singh A, Mittal D, Lavania D, Agarwal M, Mishra RC, Grover A (2012) OsHsfA2c and OsHsfB4b

are involved in the transcriptional regulation of cytoplasmic OsClpB (Hsp100) gene in rice

(Oryza sativa L.). Cell Stress Chaperones **17**: 243-254

Sosso D, Luo D, Li QB, Sasse J, Yang J, Gendrot G, Suzuki M, Koch KE, McCarty DR, Chourey

PS, Rogowsky PM, Ross-Ibarra J, Yang B, Frommer WB (2015) Seed filling in

domesticated maize and rice depends on SWEET-mediated hexose transport. Nat Genet

47: 1489-1493

Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL, Wold BJ,

Pachter L (2010) Transcript assembly and quantification by RNA-Seq reveals unannotated

transcripts and isoform switching during cell differentiation. Nat Biotechnol **28**: 511-515

Waadt R, Schmidt LK, Lohse M, Hashimoto K, Bock R, Kudla J (2008) Multicolor bimolecular

fluorescence complementation reveals simultaneous formation of alternative CBL/CIPK

complexes in planta. Plant J **56**: 505-516

Walter M, Chaban C, Schutze K, Batistic O, Weckermann K, Nake C, Blazevic D, Grefen C,

Schumacher K, Oecking C, Harter K, Kudla J (2004) Visualization of protein interactions

in living plant cells using bimolecular fluorescence complementation. Plant J **40**: 428-438

Wang L, Xie W, Chen Y, Tang W, Yang J, Ye R, Liu L, Lin Y, Xu C, Xiao J, Zhang Q (2010) A

dynamic gene expression atlas covering the entire life cycle of rice. Plant J **61**: 752-766

Wang L, Zhou Z, Song X, Li J, Deng X, Mei F (2008) Evidence of ceased programmed cell death

in metaphloem sieve elements in the developing caryopsis of Triticum aestivum L.

Protoplasma **234**: 87-96

Wang Y, Xiao Y, Zhang Y, Chai C, Wei G, Wei X, Xu H, Wang M, Ouwerkerk PB, Zhu Z (2008)

Molecular cloning, functional characterization and expression analysis of a novel

monosaccharide transporter gene OsMST6 from rice (Oryza sativa L.). Planta **228**: 525-

535

Wu X, Liu J, Li D, Liu CM (2016) Rice caryopsis development II: Dynamic changes in the

endosperm. J Integr Plant Biol **58**: 786-798

Xiong Y, Ren Y, Li W, Wu F, Yang W, Huang X, Yao J (2019) NF-YC12 is a key multi-functional

regulator of accumulation of seed storage substances in rice. J Exp Bot **70:** 3765-3780

Xu GY, Rocha PS, Wang ML, Xu ML, Cui YC, Li LY, Zhu YX, Xia X (2011) A novel rice calmodulin-

like gene, OsMSR2, enhances drought and salt tolerance and increases ABA sensitivity in

Arabidopsis. Planta **234:** 47-59

Xue W, Xing Y, Weng X, Zhao Y, Tang W, Wang L, Zhou H, Yu S, Xu C, Li X, Zhang Q (2008)

Natural variation in Ghd7 is an important regulator of heading date and yield potential in

rice. Nat Genet **40:** 761-767

Yamaguchi M, Ohtani M, Mitsuda N, Kubo M, Ohme-Takagi M, Fukuda H, Demura T (2010)

VND-INTERACTING2, a NAC domain transcription factor, negatively regulates xylem

vessel formation in Arabidopsis. Plant Cell **22:** 1249-1263

Yang J, Luo D, Yang B, Frommer WB, Eom JS (2018) SWEET11 and 15 as key players in seed

filling in rice. New Phytol

Yang SD, Seo PJ, Yoon HK, Park CM (2011) The Arabidopsis NAC transcription factor VNI2

integrates abscisic acid signals into leaf senescence via the COR/RD genes. Plant Cell **23:**

2155-2168

Young MD, Wakefield MJ, Smyth GK, Oshlack A (2010) Gene ontology analysis for RNA-seq:

accounting for selection bias. Genome Biol **11:** R14

Zhang WH, Zhou YC, Dibley KE, Tyerman SD, Furbank RT, Patrick JW (2007) Nutrient loading

of developing seeds. Functional Plant Biology **34:** 314-331

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM,

Brown M, Li W, Liu XS (2008) Model-based analysis of ChIP-Seq (MACS). Genome Biol

9: R137

Zhang Z, Dong J, Ji C, Wu Y, Messing J (2019) NAC-type transcription factors regulate

accumulation of starch and protein in maize seeds. Proc Natl Acad Sci U S A **116:** 11223-

11228