Transcriptome-wide analysis of expressed resistance gene analogs (RGAs) in mango

Darlon V. Lantican[1]*, Cris Q. Cortaga[2], Anand Noel C. Manohar[1], Fe M. dela Cueva[3], and

Maria Luz J. Sison[2]

[1]Genetics Laboratory, [2]Entomology Laboratory and [3]Plant Pathology Laboratory, Institute of

Plant Breeding, College of Agriculture and Food Science, University of the Philippines Los

Baños, College, Laguna, Philippines 4031

*corresponding author: dvlantican@up.edu.ph

Abstract

Mango is an economically important fruit crop largely cultivated in the (sub)tropics and

thus, is constantly challenged by a myriad of insect pests and diseases. Here, we identified

and characterized the resistance gene analogs (RGAs) of mango from *de novo* assembly of

transcriptomic sequences. A core RGA database of mango with 747 protein models was

established and classified based on conserved domains and motifs: 53 nucleotide binding site

proteins (NBS); 27 nucleotide binding site-leucine rich repeat proteins (NBS-LRR); 17 coiled-

coil NBS-LRR (CNL); 2 toll/interleukin-1 receptor NBS-LRR (TNL); 29 coiled-coil NBS (CN); 4

toll/interleukin-1 receptor NBS (TN); 17 toll/interleukin-1 receptor with unknown domain (TX);

158 receptor-like proteins (RLP); 362 receptor-like kinases (RLK); 72 transmembrane coiled-

coil domain protein (TM-CC), and 6 NBS-encoding proteins with other domains. The various

molecular functions, biological processes, and cellular localizations of these RGAs were

functionally well-annotated through gene ontology (GO) analysis, and their expression profiles

across different mango varieties were also determined. Phylogenetic analysis broadly

clustered the core RGAs into 6 major clades based on their domain classification, while TM-

CC proteins formed subclades all across the tree. The phylogenetic results suggest highly

divergent functions of the RGAs which also provide insights into the mango-pest co-

evolutionary arms race. From the mango RGA transcripts, 134 unique EST-SSR loci were

identified, and primers were designed targeting these potential markers. To date, this is the

most comprehensive analysis of mango RGAs which offer a trove of markers for utilization in

resistance breeding of mango.

Keywords: mango, resistance gene analogs (RGAs), transcriptome, bioinformatics

Funding

The study is supported by the Department of Science and Technology-Philippine Council for

Agriculture, Aquatic and Natural Resources Research and Development (DOST-PCAARRD)

and co-funded by the University of the Philippines Los core funds.

Introduction

Mango (*Mangifera indica* L.) is a globally popular fruit crop with economical and agricultural significance, especially in the (sub)tropics where it is largely cultivated. One of the constraints in mango production worldwide include a myriad of insect pests (e.g., oriental fruit fly, mango hopper, cecid fly, etc.) and diseases, (e.g., anthracnose, stem-end rot, scab, etc.) which can affect mango at different life stages thus significantly reducing fruit quality and yield (Bally, 2006). These are major barriers to the export market and can impede international trade as many pests and diseases in mango are important quarantine considerations.

Along the course of evolution, plants have generally developed mechanisms to recognize pathogen and insect attacks and to activate defense response against them (Dangl and Jones., 2001; Acevedo et al., 2015). The molecules derived from pathogens, insects, and plant cell damage (e.g., due to pathogen and insect attack) that trigger defense responses are generally referred to as microbe-/pathogen-/herbivore-/damage-associated molecular patterns (MAMPs/ PAMPs/ HAMPs/ DAMPs) (Acevedo et al., 2015; Choi and Klessig, 2016). These molecular patterns are being recognized by the pattern recognition receptors (PRRs) in the plant cell and trigger defense responses as these are perceived by the plant as "non-self" and indicative of pest attack (Pitzschke, 2013; Zipfel et al., 2006, Acevedo et al., 2015; Choi and Klessig, 2016), Hence, these molecules are generally considered as elicitors of plant defense responses and initiate the first line of defense, known as PAMP or DAMP-triggered immunity (P/DTI) (Jones and Dangl, 2006; Hu et al., 2018). The first layer of defense, however, is often overcome by plant pests through effector proteins, leading to effector-triggered susceptibility (ETS). To counteract this, plants can have a second line of defense known as effector-triggered immunity (ETI) (Boller and He, 2009; Pieterse et al., 2012). In ETI, due to host-pest co-evolutionary "arms race", plants acquire a family of polymorphic and diverse resistance genes (*R* genes) which encode R proteins that recognize the attacker-specific effectors (Cui et al., 2015; Jones and Dangl, 2006) to correspond for a gene-for-gene

interaction (Thompson and Burdon, 1992). This recognition results in intracellular signaling

and switching 'on' of plant defense genes leading to plant resistance against pest attack.

Collectively, the PRRs and *R* genes are called resistance gene analogs (RGAs) which

share conserved domains and motifs (Li et al., 2016). PRRs are predominantly categorized

as receptor-like kinases (RLKs) and receptor like proteins (RLPs). RLKs have an extracellular

sensing domain [either leucine-rich repeat (LRR) type or lysin motif (LysM) type], a

transmembrane (TM) domain, and an intracellular kinase domain; while RLP possess similar

domain structure except that it lacks an intracellular kinase domain (Tör et al., 2009). The R

proteins are intracellular immune receptors (effector-recognition receptors) and majority of

which belongs to nucleotide binding-site LRR (NBS-LRR) class (Gururani et al., 2012;

Neupane et al., 2018). The R proteins contain certain domains or motifs such as

serine/threonine kinases (STKs), LRRs, NBS, TMs, leucine-zipper (LZ), coiled-coil (CC) and

toll/interleukin-1 receptor (TIR) (Li et al., 2016). Depending on the domain/motif architecture

combinations, the subgroups of NBS-encoding proteins were designated as NBS, CC-NBS-

LRR (CNL), TIR-NBS-LRR (TNL), CC-NBS (CN), TIR-NBS (TN), NBS-LRR (NL), TIR with

unknown domain (TX), and other NBS-protein that may have chimeric domain/motif

architecture. Since RGAs share conserved structural features, it is possible to predict them

extensively to obtain a deeper insight on the underlying molecular defenses of the plant. RGAs

have been widely studied and efficiently used for breeding of resistant crops against insect

pests and diseases (Karthika et al., 2019). One way to predict RGAs in plants is through

bioinformatics analyses of next-generation sequencing (NGS) genomic and transcriptomic

data (Lantican et al., 2019; Neupane et al., 2018; Zhang et al., 2016; Karthika et al., 2019).

With the advent of this technology, it is possible to unravel gene networks and develop

molecular markers tagging economically important traits, reveal other molecular information

such as intron-exon boundaries and presence of transposable elements, and discover novel

biological processes (Goodwin et al., 2016).

The RGAs were first comprehensively studied on a genome-wide scale using the

model plant *Arabidopsis thaliana* and since then, thousands of RGAs from numerous plant

genomes have been identified (Meyers et al., 2003; Sekhwal et al., 2015). Despite the

economical and agricultural importance of mango, especially in the tropics, its whole genome

sequence (WGS) is yet to be available. Mango is believed to be an allotetraploid (4n=40) with

genome size of around 450 Mb according to the on-going WGS effort (Chagné, 2015).

Therefore, no detailed genome-wide analysis of mango RGAs has been performed, which is

vital to facilitate breeding of resistant mango varieties. With the absence of such valuable

resource, this study aimed to comprehensively identify and characterize the expressed RGAs

of mango through *de novo* assembly of mango transcriptomes currently available in biological

repositories. To our knowledge, this paper provides the most comprehensive identification and

characterization of mango RGAs to date, which offer a trove of markers for direct utilization in

resistance breeding of mango against insect pests and pathogens.

Materials and Methods

De novo transcriptome assembly

To obtain the expressed RGAs in mango, raw transcriptome reads were accessed and

retrieved from public repository (NCBI; Table 1) and one directly from Hong et al. (2016). The

raw read sequences were pre-processed by removing the adapter sequences and low-quality

base score nucleotide sequences using Trimmomatic v0.36 (Bolger et al., 2014) with the

following parameters: SLIDINGWINDOW: 5:30; LEADING:5; TRAILING:5; MINLEN:85. Each

trimmed transcriptome FASTQ files was independently assembled *de novo* using Trinity

pipeline (Grabherr et al., 2011) at default parameters. Upon assembly of the contigs,

SuperTranscripts were then generated to provide an accurate representation of each

expressed genes in the transcriptome assembly (Davidson et al., 2017). TransDecoder (Haas

et al., 2013) was subsequently used to construct a polypeptide sequence database from the

coding region of the SuperTranscripts based on nucleotide composition and open reading

frame (ORF) length with default parameters.

RGA identification and classification

Mango transcriptome-wide resistance gene analogs (RGA) candidates belonging to

Nucleotide Binding Site (NBS) and transmembrane-coiled-coil (TM-CC) containing proteins,

and membrane associated receptor-like kinase (RLK) and receptor-like proteins (RLP)

families were identified in the generated gene models from genome annotation using

RGAugury (Li et al., 2016), an automated RGA prediction pipeline. The input protein

sequences were initially filtered using BLASTp search against RGAdb database integrated in

the pipeline using an *e*-value cut-off of $1e^{-5}$. The domain and motif of the initial set of candidate

RGAs were detected using nCoils (Lupas et al., 1991), phobius (Kall et al., 2004), pfam_scan

(Finn et al., 2010) and InterProScan 5 (Zdobnov and Apweiler, 2001) implemented within the

RGAugury pipeline. A core set of RGAs were generated by concatenation and clustering (at

90% identity) of the identified RGAs from each polypeptide database using CD-HIT (Fu et al.,

2012), and de-duplication based on BLAST description. The NBS-encoding proteins

containing chimeric domain/motif architecture and designated by RGAugury as OTHER were

manually checked for their domain/motif composition using Multiple Expectation Maximization

for Motif Elicitation (MEME) (Bailey and Elkan, 1994) and MOTIF Search

(https://www.genome.jp/tools/motif/).

RGA characterization and annotation

Gene ontology (GO) annotations such as molecular function (MF), biological

processes (BP), and cellular component (CC) were assigned to each protein represented in

the constructed mango RGA core database using BLAST2GO package (Conesa et al., 2005).

The homology of the predicted polypeptide sequences of each RGAs to existing entries in

UniProtKB/SwissProt protein database were determined using BLASTp (with e-value of $1e^{-5}$).

The mapped BLAST hits were then merged to InterProScan (Zdobnov and Apweiler, 2001)

search output to produce the GO annotations.

Profiling of mango RGA expression

RNA-Seq by Expectation Maximization (RSEM) (Li and Dewey, 2011) was used to

provide insights on the expression profile of each RGA obtained from different tissues and

diverse mango varieties. The pre-processed RNA-seq reads were independently mapped to

the constructed core mango RGA reference transcriptome using Bowtie2 (Langmead and

Salzberg, 2012). Transcripts per million (TPM) count values were then generated by RSEM

from each paired-end reads. A comma-separated value (csv) file containing the RGAs and

TPM values for each transcriptome data set was prepared and uploaded to Heatmapper

(Babicki et al., 2016) for heat-map visualization of the RGA expression profiles.

Evolutionary analysis

The FASTA amino acid sequences of the 747 core mango RGAs were used to

construct a phylogenetic tree. Multiple Sequence Alignment (MSA) of the RGA sequences

was performed using the CLUSTALW program (Thompson et al., 1994) with the following

parameters: Gap Opening Penalty: 10; Gap Extension Penalty: 0.2. The phylogeny of these

aligned sequences was reconstructed using Maximum Likelihood statistical method using IQ-

TREE (Nguyen et al., 2015) with best-fit substitution model selected through ModelFinder

(Kalyaanamoorthy et al., 2017). Based on the Bayesian Information Criterion (BIC) of the

models, General 'Variable Time' (VT) matrix (Mueller and Vingron, 2004) with empirical amino

acid frequencies (+F) and FreeRate (+R5) rate heterogeneity across sites (Yang 1995;

Soubrier et al., 2012) was used to generate the tree. The resulting phylogenetic tree was

validated with 1000 replicates of ultrafast bootstrapping (Hoang et al., 2017) and visualized

using FigTree (v1.4.4) (Rambaut, 2018).

Identification and design of EST-SSRs

The FASTA file corresponding to the RGA transcripts selected for analysis was

uploaded into GMATA (Genome-wide Microsatellite Analyzing Toward Application) Software

Package (Wang & Wang, 2016). SSR loci within the data set were identified using default

parameters [Min-length (nt) = 2; Max-length (nt) = 6; Min. repeat-times = 5]. Consequently,

oligonucleotide primer pairs were designed at regions flanking the identified SSRs using the

Primer3 (Untergasser et al., 2012; Koressaar and Remm, 2007) integrated within the package.

Default parameters were also used in primer design [Min. amplicon size (bp) = 120; Max.

amplicon size (bp) = 400; Optimal annealing T_m (°C) = 60; Flanking sequence length = 400;

Max. template length = 2000].

Results and Discussion

Identification of mango RGAs

Raw transcriptome sequences from diverse mango cultivars were retrieved,

assembled, and the RGAs were identified using RGAugury, an integrative NGS bioinformatics

pipeline that efficiently predicts RGAs in plants (Li et al., 2016). In general, this pipeline works

by identifying first the domains and motifs related to RGAs such as NBS, LRR,

serine/threonine and tyrosine kinase (STTK), CC, LysM, TM, and TIR. Based on the presence

of combinations of these domains and motifs, the RGA candidates are then identified and

categorized into four major families: NBS-encoding, TM-CC, and membrane associated RLP

and RLK. The number of RGAs predicted from the sequence reads archive (SRA) files ranged

from 56 to 454 (Table 1). In total, 7,391 RGAs were identified from the assembled

transcriptome sequences (Table 1; Supplemental File 1). These RGAs were further clustered

at 90% homology creating 2,892 clusters, followed by automated selection of a representative

protein sequence per cluster. The representative protein sequences were described using

BLAST and BLAST2GO, and proteins with duplicated BLAST description were removed

retaining only unique/non-redundant protein type per RGA domain to generate 747 core RGAs

with unique functions (Table 2, Supplemental Table 1 and Supplemental File 2). The core

RGA proteins were classified as follows: 53 nucleotide binding site protein (NBS); 27

nucleotide binding site-leucine rich repeat proteins (NBS-LRR); 17 coiled-coil NBS-LRR

(CNL); 2 toll/interleukin-1 receptor NBS-LRR (TNL); 29 coiled-coil NBS (CN); 4 toll/interleukin-

1 receptor NBS (TN); 17 toll/interleukin-1 receptor with unknown domain (TX); 158 receptor-

like proteins (RLP); 362 receptor-like kinases (RLK); 72 transmembrane coiled-coil domain

protein (TM-CC); and 6 NBS-encoding proteins with other domains (Table 2).

Upon comparison to other important agricultural crops, as reported by Li et al. (2016),

it appears that mango harbors more RGAs (747) than papaya (467) but has lower compared

to cacao (1,281), tomato (1,003), corn (1,062), rice (1,652) and banana (892) (Table 2).

Eventually, the RGA content and characteristics of a plant has been frequently associated

with crop resistance (Sekhwal et al., 2015). Like most crops, RLKs constitute the largest group

of RGAs in mango comprising almost 50% of the RGAs characterized, followed by RLPs, then

the NBS-encoding proteins (Table 2, Fig. 1). In dicots such as mango, all NBS-encoding

proteins are present (i.e., NBS, CNL, TNL, CN, TN, NL, TX, and other NBS-encoding proteins)

(Table 2) but in monocots such as corn, rice, banana, the TNL protein is usually absent (Table

2) (Tarr and Alexander, 2009). It is hypothesized that after the divergence of dicots and

monocots, the TNL genes might have been lost from the monocot lineage (Zhang et al., 2016).

Six mango RGA models showed chimeric domain/motif architecture and were classified as

'other' NBS-encoding proteins (Table 2). These were manually checked using the MEME

platform and appear to be NBS-encoding proteins containing TIR sequences and other

uncharacterized domain (Supplemental Figure 1).

Mango resistance (R)/defense proteins and expression profiles

BLAST analysis of the core mango RGA database revealed homology to a number of

well-known resistance (R) /defense proteins against pathogens and insects (Table 3). Among

these R proteins include RRS proteins (putative WRKY transcription factors), which confer

resistance against *Colletotrichum higginsianum* and *Ralstonia solanacearum* (Narusaka et al.,

2009; Saucet et al., 2015); RPP proteins (including At4g19530 and At4g19520 proteins) which

confer resistance against downy mildew (*Peronospora parasitica*) (Parker et al., 1997;

Sinapidou et al., 2004; Bittner-Eddy et al., 2000; Wan et al., 2019); the LRK10 proteins (all of

which are of RLK domains) which provide resistance against leaf rust (*Puccinia triticina*) (Shiu

and Bleecker, 2003; Feuillet et al., 1998); the RGA-blb and R1A-10 proteins which provide

resistance against the devastating late blight disease (*Phytophtora infestans*) (Lokossou et

al., 2010; Kuang et al., 2005); the RLM proteins which confer resistance against the fungal

pathogen *Leptosphaeria maculans* (Staal et al., 2008; 2006); the Pik-2 and RGA5 proteins

which confer resistance against blast disease (*Magnaporthe oryzae*) (Zhai et al., 2011;

Okuyama et al., 2011); and many homologues of TMV resistance protein N which are

responsible for tobacco mosaic virus (TMV) resistance (Dinesh-Kumar et al., 2000). The core

mango RGA also contained numerous sequences with homology to various RPS proteins,

including RPM1 and TAO1 proteins, which are important R proteins against the bacterium

Pseudomonas syringae, a widely studied biotrophic pathogen (Eitas et al., 2008; Mackey et

al., 2002; Warren et al., 1998; Kim et al., 2009).

Two RGA gene models showed homology to ERECTA protein which confer

quantitative resistance against the necrotrophic fungus *Plectosphaerella*, and bacterial wilt

caused by *Ralstonia solanacearum* (Sánchez-Rodríguez et al., 2009). This R protein is also

responsible in the regulation of efficient transpiration in plant (Masle et al., 2005). Homology

to the R protein EDR1 (ENHANCED DISEASE RESISTANCE 1), a MAPKKK

serine/threonine-protein kinase, was also found which confer non-host resistance against

Colletotrichum gloeosporioides in *Arabidopsis thaliana* through the induction of plant

defensins expression (Hiruma et al., 2011). One mango RGA showed homology to the R

protein ADR2 (ACTIVATED DISEASE RESISTANCE 2), an NBS-encoding protein that

confers broad-spectrum resistance against many pathogens such as strains *of Pseudomonas*

syringae and *Hyaloperonospora arabidopsis*, and full immunity to several races of the

pathogen *Albugo candida*, the causal agent of white rust disease (Borhan et al., 2008; Aboul-

Soud et al., 2009). This R protein is also reported to play a role in the response to UV stress

(Piofczyk et al., 2015). Predicted mango RGAs homologous to the R-like proteins DSC 1 and

2 (DOMINANT SUPPRESOR OF CAMTA3) were also found which act as guard of CAMTA3,

a negative regulator of immunity, during infection of pathogens (Lolle et al., 2017). Three NBS-

encoding RGAs in mango showed homology to SUMM2 (SUPPRESSOR OF *mkk1 mkk2* 2)

which acts as an R protein that becomes activated when the MEKK1-MKK1/MKK2-MPK4

cascade in the basal defense response is disrupted by the pathogen effector HopAI1 (Zhang

et al., 2012; Kong et al., 2012).

Other putative resistance proteins abundantly found in mango showed homology to

various DRL-coded (Uniprot/SwissProt ID) potential R proteins from *Arabidopsis thaliana*

which have not been fully studied (Supplemental Table 1). Homologues to CHS and CHL

proteins were also found in mango RGAs (Supplemental Table 1) which are important R

proteins against chilling/low temperature and also confer resistance against bacterial infection

such as *Pseudomonas syringae* (Zbierzak et al., 2013). Aside from resistance to pathogens,

R/defense proteins against insects were also identified by filtering the gene ontology results

with GO term "defense response to insect" (GO:0002213). The RGAs found include the

At4g11170 putative resistance protein, RPP5, DSC1 and 2, TRANSPORT INHIBITOR

RESPONSE 1 (TIR1), and CORONATINE INSENSITVE-1 (COI-1). TIR1 is an F-box protein

and a major receptor of the plant hormone auxin and mediate in the auxin/indole acetic acid

(IAA)-regulated signaling pathways involved in plant growth and development, as well as

defense response against pathogens and insect pests (Dharmasiri et al., 2005). COI-1 is a

jasmonic acid-isoleucine (JA-Ile) coreceptor and also an F-box protein involved in the JA

signaling pathway which regulate plant defense against insects and pathogens, wound

healing, and other vegetative and reproductive developmental functions (Xie et al., 1998). The

plant hormone JA is conjugated to isoleucine (Ile) as a consequence of insect/pathogen

attack, and JA-Ile binding with COI-1 stimulates degradation of JAZ coreceptor repressor

proteins, thereby promoting the expression of JA-responsive genes (Pieterse et al., 2012).

Further filtering of RGAs containing the GO term "response to insect" (GO:0009625) revealed

17 RGAs that putatively play vital roles in the defense pathways against insects (Table 4), all

of which were characterized to possess RLK/RLP domains. The complete and detailed

BLASTp analysis and description of the RGAs are provided in Supplemental Table 1.

Of all the mango varieties analyzed in this study, the three varieties that showed the

lowest expression of RGAs were SRR8449851 (CHOKANAN), SRR3359450 (DASHEHARI),

and SRR1562177 (KENT) (Fig. 2; Supplemental Table 2). On the other hand, the following

showed the highest expressions of the RGAs characterized: SRR2162953 (pooled RNA from

different varieties), SRR2162919 (THAI EVERBEARING), SRR2162878 (BURMA),

SRR2162889 (M. CASTURI "PURPLE"), SRR2159471 (TOMMY ATKINS), SRR2162929

(TURPENTINE), SRR2162836 (AMIN ABRAHIMPUR), SRR2162908 (NEELUM) (Fig. 2;

Supplemental Table 2). The relatively high expression of RGAs in these RNA-seq reads could

be explained by the fact that they are pooled RNA samples from different varieties or from

different plant tissues.

Gene ontology and functional annotation of mango RGAs

Gene ontology (GO) analysis and functional annotation of the core RGA database

provided a broader knowledge on the major roles of these proteins and their cellular

localizations. The molecular functions of the core RGAs are primarily associated with protein

and nucleotide binding, and kinase activity (Fig. 3) as they are known extra- and intracellular

binding receptors and relays defense signaling in the cells through a cascade of kinase

activities (Rodriguez et al., 2010). The absence of an intracellular kinase in RLPs suggests

that it relies/interact with other domains such as RLK-type receptors to relay the signalling

from extracellular matrix to the intracellular component (Tör et al., 2009). Meanwhile, the

biological processes of the RGAs are highly diverse, distributing the RGA sequences to a wide

array of GO IDs/terms. The RGAs are mainly involved in protein (auto)phosphorylation during

signal transductions in the cell, defense/immune responses to different biotic (insect pests and

diseases) and abiotic (water deprivation, ozone, UV stress, etc.) stresses, and in various

developmental processes (i.e., from embryonic to floral/pollen development) (Fig. 4).

Numerous RGAs were also described to be involved in hormone-mediated signaling pathways

and systemic acquired resistance which include the plant hormones brassinosteroids, abscisic

acid (ABA), salicylic acid (SA), auxin, jasmonic acid (JA), ethylene, and gibberellic acid (GA).

These plant hormones create a crosstalk to modulate defense signaling pathways and activate

systemic resistance against a broad spectrum of pathogens and insect pests. As an example

of this crosstalk, SA antagonizes JA to activate immune response against biotrophic

pathogens (Williams, 2011). Likewise, JA antagonizes SA to activate two branches of immune

response which also antagonizes each other: the ERF branch against necrotrophic pathogens

(synergistic interaction with ethylene) and the MYC branch against insects (synergistic

interaction with ABA) (Pieterse et al., 2012). In the cell, majority of the mango RGAs are

localized in the plasma membrane, plasmodesma, cytosol, and nucleus which are primary

sites for recognition of pathogen/insect invasion and effector proteins. In these cellular

components, the RGAs functions to convert extracellular stimuli into intracellular responses to

activate defense cascades and counteract the attack. In general, the results of GO analysis

and functional annotation show that plant immune response is a consequence of the activity

of a wide range of plant hormones, resistance/defense-related proteins, biochemical and

developmental processes, etc. that aim to suppress pest attack thereby improving plant

defense. The complete and detailed GO analysis and annotation of each RGAs are provided

in Supplemental Table 1 and Supplemental Table 3.

Evolutionary relationships among mango RGAs

Using the evolutionary best-fit model selected according to Bayesian Information

Criterion (BIC) (Supplemental File 3), a maximum likelihood phylogenetic tree was constructed

using the 747 core mango RGA proteins to investigate their evolutionary relationships and

diversity (Fig. 6; Supplemental File 4). It showed that clustering of RGAs based on domain

classification is very apparent but mingled with subclades from other RGA domains. As shown

in the tree, the mango RGAs have become highly diversified and are broadly clustered into 6

major clades with many subclades (Fig. 6). RGAs having TM-CC domain didn't emerge as a

separate cluster but formed subclades all across the tree suggesting that it shares similarity

to all other RGA classes. The clade 1 is composed mainly of NBS-encoding proteins and other

domains such as TM-CC and RLP. Notably, the clade 1 appeared to be the most distantly

related RGA group among other clades. The clade 2 is predominantly composed of RLP

proteins with subclades from all other RGA domains. This clade appears to be a sister clade

to clade 3, which contains some RLP, TM-CC, and most especially, numerous RLK proteins

which formed a major nested subclade indicating their close relationship with RLPs. The

clades 4 and 5 appeared to be sister clades and are composed mainly of NBS-encoding

proteins with subclades from all other RGA domains and most especially, from TM-CC

domains. Rody et al. (2019) also observed that TM-CC sequences in sugarcane form

widespread subclades across the phylogenetic tree and showed close relationship with NBS-

encoding proteins. The clade 6 is composed predominantly of RLKs and shared close

relationship to some other RGAs. In this clade, NBS-encoding proteins that shared close

similarity related to RLKs were observed to form a major nested subclade. The NBS and RLKs

are among the largest and diversified group of RGAs in plants (Hulbert et al., 2001; Meyers et

al., 2003; Tör et al., 2009, Li et al., 2016), which also holds true in mango based on this study.

The distribution of RGAs in the phylogenetic tree may indicate that their functions are highly

divergent and form clusters that may be related to functions but not necessarily in protein

sequence (Chang et al., 2002; Zhang et al., 2016). This could be the case in mango RGAs,

especially the NBS-encoding proteins which are broadly distributed into 3 major clades (clades

1, 4 and 5) and 1 major subclade (nested in clade 6), as well as in RLKs which formed a

major clade (clade 6) and a major subclade (nested in clade 3).

Many factors have been attributed to the evolutionary pattern and diversity of RGAs in

plants. One of which is the well-known co-evolutionary "arms race" model between the host

plant and associated pests and diseases which drives the selective forces to overcome each

other (Anderson et al., 2010; Edger et al., 2015; Boller and He, 2009). Other selection

pressures such as climatic conditions (e.g. rainfall, temperatures, humidity, etc.) that favors

pest and disease development have been correlated also to the diversity of RGAs (Sela et al.,

2009). Whole-genome duplications (WGDs) and genomic reorganizations that occurred

during ancient times, especially in angiosperms, have been associated to the expansion of

RGA families and in creating new gene functions (Perazzolli et al., 2014, Chagné, 2015;

Michelmore and Meyers, 1998). It is likely that in the evolutionary history of mango, it had

undergone at least one round of polyploidization i.e., allopolyploidy which is a type of WGD

via hybridization followed by genome doubling (Glover et al., 2016; Soltis and Soltis, 2009).

These are some factors that could have influenced the complex phylogenetic structure of

RGAs of mango, given also that the plant's tropical growing condition provides a thriving

environment to a plethora of insect pests and diseases.

RGA-derived EST-SSRs of mango

Using the RGA transcripts (Supplemental File 5), a total of 151 EST-SSR (Expressed

Sequence Tags – Simple Sequence Repeat) loci were identified. Among these, 134 loci

yielded unique potential markers. Mostly di-repeat motifs (2-mer) were observed, accounting

for 64.24% of the markers, followed by tri-repeats (3-mer) at 30.46%. The remaining 5.3% is

composed of tetra-, hexa-, and penta-repeat motifs. AT repeat is the most represented motif

among the di-repeats at 12.5%, followed by TC and TA at 11.26% each (Fig. 7). Meanwhile,

AG/CT is the most abundant paired repeat motif among the SSR loci analyzed at 16.56% (Fig.

8). Data on the distribution of *R* gene motif-linked SSRs (Fig. 9) revealed that RLK has the

most number of EST-SSRs designed with 67 markers, followed by RLP and TM-CC motifs at

34 and 18 markers, respectively. This distribution corresponds directly to the amount of *R*

genes per category identified in the whole mango RGA transcriptome.

Similar *in silico* approaches in EST-SSR marker development were performed by other

studies focusing on other plants, such as bamboo (Cai et al., 2019), mint (Kumar et al., 2015),

castor (Thatikunta et al., 2016), Ethiopian potato (Gadissa et al., 2018) and *Elymus* species

(Zhang et al., 2019). The EST-derived SSR markers generated in this study can be used for

functional analysis of *R* genes, once verified by wet lab experiments. Moreover, these markers

may also be used in genotyping, genetic diversity, linkage mapping, gene-based association

studies and marker-assisted selection in mango breeding. The sequences of the designed

forward and reverse primers derived from mango RGAs are found in Supplemental Table 4.

Summary and Conclusions

In this paper, 747 core mango RGAs were identified and broadly categorized into four

major families based on their conserved structural features: NBS-encoding proteins (155

proteins), RLPs (158), RLKs (362) and TM-CC (72). In addition, the mango RGAs have been

functionally well-annotated revealing the involved biological processes, molecular functions

and cellular components. This provides an important insight to the overall functional response

of mango against insect pests and diseases. Moreover, the evolutionary relationships,

diversity, and expression profiles across different mango varieties were presented as an

invaluable reference in the design of effective framework for the actual mango resistance

breeding programs. A marker resource, comprising of 134 RGA-linked EST-SSR markers,

was also developed to aid in the discovery of associated RGA/s to a specific disease or insect

pest through population genetics analyses (e.g., QTL mapping, association mapping). The

basic information from the mango RGAs and the linked EST-SSR markers established in this

paper will facilitate in the development of outstanding mango varieties with resistance to

various pests and diseases.

Acknowledgements

The authors would like to thank the Department of Science and Technology-Philippine

Council for Agriculture, Aquatic and Natural Resources Research and Development (DOST-

PCAARRD) for funding the project entitled "Characterization of 'Carabao' and other Mango

Varieties with Resistance to Fruit Fly and Anthracnose". Also, the University of the Philippines

Los Baños for the core-funded project, "Genome-wide analysis of resistance gene analogs

(RGAs) and development of RGA-linked DNA markers in Philippine Important Crops".

References

Aboul-Soud, M. A., Chen, X., Kang, J. G., Yun, B. W., Raja, M. U., Malik, S. I., & Loake, G. J. (2009). Activation tagging of ADR2 conveys a spreading lesion phenotype and resistance to biotrophic pathogens. *New phytologist*, *183*(4), 1163-1175.

Acevedo, F. E., Rivera-Vega, L. J., Chung, S. H., Ray, S., & Felton, G. W. (2015). Cues from chewing insects—the intersection of DAMPs, HAMPs, MAMPs and effectors. *Current Opinion in Plant Biology*, *26*, 80-86.

Anderson, J. P., Gleason, C. A., Foley, R. C., Thrall, P. H., Burdon, J. B., & Singh, K. B. (2010). Plants versus pathogens: an evolutionary arms race. *Functional plant biology*, *37*(6), 499-512.

Babicki, S., Arndt, D., Marcu, A., Liang, Y., Grant, J. R., Maciejewski, A., & Wishart, D. S. (2016). Heatmapper: web-enabled heat mapping for all. *Nucleic acids research*, *44*(W1), W147-W153.

Bailey, T. L., & Elkan, C. (1994). Fitting a mixture model by expectation maximization to discover motifs in bipolymers. Proc. Int. Conf. Intell. Syst. Mol. Biol. 1994, 2, 28–36.

Bally, I. S. (2006). Mangifera indica (mango). *Species profiles for pacific island agroforestry*, 1-25.

Bittner-Eddy, P. D., Crute, I. R., Holub, E. B., & Beynon, J. L. (2000). RPP13 is a simple locus in Arabidopsis thaliana for alleles that specify downy mildew resistance to different avirulence determinants in Peronospora parasitica. *The Plant Journal*, *21*(2), 177-188.

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, *30*(15), 2114-2120.

Boller, T., and He, S. Y. (2009). Innate immunity in plants: an arms race between pattern recognition receptors in plants and effectors in microbial pathogens. *Science*, *324*(5928), 742-744.

Borhan, M. H., Gunn, N., Cooper, A., Gulden, S., Tör, M., Rimmer, S. R., & Holub, E. B. (2008). WRR4 encodes a TIR-NB-LRR protein that confers broad-spectrum white rust resistance in Arabidopsis thaliana to four physiological races of Albugo candida. *Molecular Plant-Microbe Interactions*, *21*(6), 757-768.

Cai, K., Zhu, L., Zhang, K., Li, L., Zhao, Z., Zeng, W., & Lin, X. (2019). Development and characterization of EST-SSR markers from RNA-Seq data in Phyllostachys violascens. *Frontiers in plant science*, *10*, 50.

Chagné, D. (2015). Whole genome sequencing of fruit tree species. In *Advances in botanical research* (Vol. 74, pp. 1-37). Academic Press.

Chang, J. H., Tai, Y. S., Bernal, A. J., Lavelle, D. T., Staskawicz, B. J., & Michelmore, R. W. (2002). Functional analyses of the Pto resistance gene family in tomato and the identification of a minor resistance determinant in a susceptible haplotype. *Molecular plant-microbe interactions*, *15*(3), 281-291.

Choi, H. W., & Klessig, D. F. (2016). DAMPs, MAMPs, and NAMPs in plant innate immunity. *BMC plant biology*, *16*(1), 232.

Conesa, A., Götz, S., García-Gómez, J. M., Terol, J., Talón, M., & Robles, M. (2005).
Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics
research. *Bioinformatics*, *21*(18), 3674-3676.

Cui, H., Tsuda, K., & Parker, J. E. (2015). Effector-Triggered Immunity: From Pathogen
Perception to Robust Defense. Annual Review of Plant Biology, 66(1), 487–511.
doi:10.1146/annurev-arplant-050213-040012

Dangl, J. L., & Jones, J. D. (2001). Plant pathogens and integrated defence responses to
infection. *Nature*, *411*(6839), 826–833. doi:10.1038/35081161

Davidson, N. M., Hawkins, A. D., & Oshlack, A. (2017). SuperTranscripts: a data driven
reference for analysis and visualisation of transcriptomes. *Genome biology*, *18*(1), 148.

Dharmasiri, N., Dharmasiri, S., & Estelle, M. (2005). The F-box protein TIR1 is an auxin
receptor. *Nature*, *435*(7041), 441-445.

Dinesh-Kumar, S. P., Tham, W. H., & Baker, B. J. (2000). Structure–function analysis of the
tobacco mosaic virus resistance gene N. *Proceedings of the National Academy of*
Sciences, *97*(26), 14789-14794.

Edger, P. P., Heidel-Fischer, H. M., Bekaert, M., Rota, J., Glöckner, G., Platts, A. E., ... &
Hofberger, J. A. (2015). The butterfly plant arms-race escalated by gene and genome
duplications. *Proceedings of the National Academy of Sciences*, *112*(27), 8362-8366.

Eitas, T. K., Nimchuk, Z. L., & Dangl, J. L. (2008). Arabidopsis TAO1 is a TIR-NB-LRR protein
that contributes to disease resistance induced by the Pseudomonas syringae effector
AvrB. *Proceedings of the National Academy of Sciences*, *105*(17), 6475-6480.

Feuillet, C., Reuzeau, C., Kjellbom, P., & Keller, B. (1998). Molecular characterization of a
new type of receptor-like kinase (wlrk) gene family in wheat. *Plant molecular biology*, *37*(6),
943-953.

Finn, R. D., Mistry, J., Tate, J., Coggill, P., Heger, A., Pollington, J. E., ... & Holm, L. (2010).
The Pfam protein families database. *Nucleic acids research*, *38*(suppl 1), D211-D222.

Fu, L., Niu, B., Zhu, Z., Wu, S., & Li, W. (2012). CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics*, *28*(23), 3150-3152.

Gadissa, F., Tesfaye, K., Dagne, K., & Geleta, M. (2018). Genetic diversity and population
structure analyses of Plectranthus edulis (Vatke) Agnew collections from diverse agro-
ecologies in Ethiopia using newly developed EST-SSRs marker system. *BMC*
genetics, *19*(1), 92.

Glover, N. M., Redestig, H., & Dessimoz, C. (2016). Homoeologs: what are they and how do
we infer them?. *Trends in plant science*, *21*(7), 609-621.

Goodwin, S., Mcpherson, J. D., & Mccombie, W. R. (2016). Coming of age: ten years of next-
generation sequencing technologies. *Nature Reviews Genetics*, *17*(6), 333-351.

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., ... & Chen,
Z. (2011). Trinity: reconstructing a full-length transcriptome without a genome from RNA-Seq
data. *Nature biotechnology*, *29*(7), 644.

Gururani, M.A.; Venkatesh, J.; Upadhyaya, C.P.; Nookaraju, A.; Pandey, S.K.; Park, S.W.
Plant disease resistance genes: Current status and future directions. *Physiol. Mol. Plant*
Pathol. 2012, *78*, 51–65.
526

527 Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden, J., ... &
528 MacManes, M. D. (2013). De novo transcript sequence reconstruction from RNA-seq using
529 the Trinity platform for reference generation and analysis. *Nature protocols*, *8*(8), 1494.
530

531 Hiruma, K., Nishiuchi, T., Kato, T., Bednarek, P., Okuno, T., Schulze-Lefert, P., & Takano, Y.
532 (2011). Arabidopsis ENHANCED DISEASE RESISTANCE 1 is required for pathogen-induced
533 expression of plant defensins in nonhost resistance, and acts through interference of MYC2-
534 mediated repressor function. *The Plant Journal*, *67*(6), 980-992.
535

536 Hoang, D. T., Chernomor, O., Von Haeseler, A., Minh, B. Q., & Vinh, L. S. (2018). UFBoot2:
537 improving the ultrafast bootstrap approximation. *Molecular biology and evolution*, *35*(2), 518-
538 522.
539

540 Hong, K., Gong, D., Zhang, L., Hu, H., Jia, Z., Gu, H., & Song, K. (2016). Transcriptome
541 characterization and expression profiles of the related defense genes in postharvest mango
542 fruit against Colletotrichum gloeosporioides. *Gene*, *576*(1), 275-283.
543

544 Hu, Q., Min, L., Yang, X., Jin, S., Zhang, L., Li, Y., ... & Lindsey, K. (2018). Laccase GhLac1
545 modulates broad-spectrum biotic stress tolerance via manipulating phenylpropanoid pathway
546 and jasmonic acid synthesis. *Plant physiology*, *176*(2), 1808-1823.
547

548 Hulbert, S. H., Webb, C. A., Smith, S. M., & Sun, Q. (2001). Resistance gene complexes:
549 evolution and utilization. *Annual review of phytopathology*, *39*(1), 285-312.
550

551 Jones, J. D. G., & Dangl, J. L. (2006). The plant immune system. Nature, 444(7117), 323–
552 329.
553

554 Käll, L., Krogh, A., & Sonnhammer, E. L. (2004). A combined transmembrane topology and
555 signal peptide prediction method. *Journal of molecular biology*, *338*(5), 1027-1036.
556

557 Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K., von Haeseler, A., & Jermiin, L. S. (2017).
558 ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature*
559 *methods*, *14*(6), 587-589.
560

561 Karthika, R., Prasath, D., & Anandaraj, M. (2019). Transcriptome-wide identification and
562 characterization of resistant gene analogs (RGAs) of ginger (Zingiber officinale Rosc.) and
563 mango ginger (Curcuma amada Roxb.) under stress induced by pathogen. *Scientia*
564 *Horticulturae*, *248*, 81-88.
565

566 Kim, S. H., Kwon, S. I., Saha, D., Anyanwu, N. C., & Gassmann, W. (2009). Resistance to the
567 Pseudomonas syringae effector HopA1 is governed by the TIR-NBS-LRR protein RPS6 and
568 is enhanced by mutations in SRFR1. *Plant physiology*, *150*(4), 1723-1732.
569

570 Kong, Q., Qu, N., Gao, M., Zhang, Z., Ding, X., Yang, F., ... & Zhang, Y. (2012). The MEKK1-
571 MKK1/MKK2-MPK4 kinase cascade negatively regulates immunity mediated by a mitogen-
572 activated protein kinase kinase kinase in Arabidopsis. *The Plant Cell*, *24*(5), 2225-2236.
573

574 Koressaar, T., & Remm, M. (2007). Enhancements and modifications of primer design
575 program Primer3. *Bioinformatics*, *23*(10), 1289-1291.
576

Kuang, H., Wei, F., Marano, M. R., Wirtz, U., Wang, X., Liu, J., ... & Lobst, S. (2005). The R1 resistance gene cluster contains three groups of independently evolving, type I R1 homologues and shows substantial structural variation among haplotypes of Solanum demissum. *The Plant Journal*, *44*(1), 37-51.

Kumar, B., Kumar, U., & Yadav, H. K. (2015). Identification of EST–SSRs and molecular diversity analysis in Mentha piperita. *The Crop Journal*, *3*(4), 335-342.

Langmead, B., Salzberg, S. (2012). Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9, 357–359.

Lantican, D. V., Strickler, S. R., Canama, A. O., Gardoce, R. R., Mueller, L. A., & Galvez, H. F. (2019). De novo genome sequence assembly of dwarf coconut (Cocos nucifera L.'Catigan Green Dwarf') provides insights into genomic variation between coconut types and related palm species. *G3: Genes, Genomes, Genetics*, *9*(8), 2377-2393.

Li, B., & Dewey, C. N. (2011). RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC bioinformatics*, *12*(1), 323.

Li, P., Quan, X., Jia, G., Xiao, J., Cloutier, S., & You, F. M. (2016). RGAugury: a pipeline for genome-wide prediction of resistance gene analogs (RGAs) in plants. *BMC genomics*, *17*(1), 852.

Lokossou, A. A., Rietman, H., Wang, M., Krenek, P., van der Schoot, H., Henken, B., ... & Jacobsen, E. (2010). Diversity, distribution, and evolution of Solanum bulbocastanum late blight resistance genes. *Molecular Plant-Microbe Interactions*, *23*(9), 1206-1216.

Lolle, S., Greeff, C., Petersen, K., Roux, M., Jensen, M. K., Bressendorff, S., ... & Petersen, M. (2017). Matching NLR immune receptors to autoimmunity in camta3 mutants using antimorphic NLR alleles. *Cell host & microbe*, *21*(4), 518-529.

Lupas, A., Van Dyke, M., & Stock, J. (1991). Predicting coiled coils from protein sequences. *Science*, 1162-1164.

Mackey, D., Holt III, B. F., Wiig, A., & Dangl, J. L. (2002). RIN4 interacts with Pseudomonas syringae type III effector molecules and is required for RPM1-mediated resistance in Arabidopsis. *Cell*, *108*(6), 743-754.

Maria Cristina Suarez Rodriguez, Petersen, M., & Mundy, J. (2010). Mitogen-activated protein kinase signaling in plants. *Annual review of plant biology*, *61*, 621-649.

Masle, J., Gilmore, S. R., & Farquhar, G. D. (2005). The ERECTA gene regulates plant transpiration efficiency in Arabidopsis. *Nature*, *436*(7052), 866-870.

Meyers, B. C., Kozik, A., Griego, A., Kuang, H., & Michelmore, R. W. (2003). Genome-wide analysis of NBS-LRR–encoding genes in Arabidopsis. *The Plant Cell*, *15*(4), 809-834.

Michelmore, R. W., & Meyers, B. C. (1998). Clusters of resistance genes in plants evolve by divergent selection and a birth-and-death process. *Genome research*, *8*(11), 1113-1130.

Müller, T., & Vingron, M. (2004). Modeling amino acid replacement. *Journal of Computational Biology*, *7*(6), 761-776.

Narusaka, M., Shirasu, K., Noutoshi, Y., Kubo, Y., Shiraishi, T., Iwabuchi, M., & Narusaka, Y.
(2009). RRS1 and RPS4 provide a dual Resistance-gene system against fungal and bacterial
pathogens. *The Plant Journal*, *60*(2), 218-226.

Neupane, S., Andersen, E. J., Neupane, A., & Nepal, M. P. (2018). Genome-Wide
Identification of NBS-Encoding Resistance Genes in Sunflower (Helianthus annuus
L.). *Genes*, *9*(8), 384.

Nguyen, L. T., Schmidt, H. A., Von Haeseler, A., & Minh, B. Q. (2015). IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular
biology and evolution*, *32*(1), 268-274.

Okuyama, Y., Kanzaki, H., Abe, A., Yoshida, K., Tamiru, M., Saitoh, H., ... & Undan, J. (2011).
A multifaceted genomics approach allows the isolation of the rice Pia-blast resistance gene
consisting of two adjacent NBS-LRR protein genes. *The Plant Journal*, *66*(3), 467-479.

Parker, J. E., Coleman, M. J., Szabò, V., Frost, L. N., Schmidt, R., van der Biezen, E. A., ... &
Jones, J. D. (1997). The Arabidopsis downy mildew resistance gene RPP5 shares similarity
to the toll and interleukin-1 receptors with N and L6. *The Plant Cell*, *9*(6), 879-894.

Perazzolli, M., Malacarne, G., Baldo, A., Righetti, L., Bailey, A., Fontana, P., ... & Malnoy, M.
(2014). Characterization of resistance gene analogues (RGAs) in apple (Malus× domestica
Borkh.) and their evolutionary history of the Rosaceae family. *PloS one*, *9*(2).

Pieterse, C. M., Van der Does, D., Zamioudis, C., Leon-Reyes, A., & Van Wees, S. C. (2012).
Hormonal modulation of plant immunity. *Annual review of cell and developmental biology*, *28*,
489-521.

Piofczyk, T., Jeena, G., & Pecinka, A. (2015). Arabidopsis thaliana natural variation reveals
connections between UV radiation stress and plant pathogen-like defense responses. *Plant
Physiology and Biochemistry*, *93*, 34-43.

Pitzschke, A. (2013). Agrobacterium infection and plant defense—transformation success
hangs by a thread. Frontiers in Plant Science, 4(December), 1–12.

Rambaut,   A.   (2018)   FigTree   v1.4.4.   Available
at: http://tree.bio.ed.ac.uk/software/figtree/ (accessed February 7, 2020).

Rody, H. V., Bombardelli, R. G., Creste, S., Camargo, L. E., Van Sluys, M. A., & Monteiro-
Vitorello, C. B. (2019). Genome survey of resistance gene analogs in sugarcane: genomic
features and differential expression of the innate immune system from a smut-resistant
genotype. *BMC genomics*, *20*(1), 809.

Sánchez-Rodríguez, C., Estévez, J. M., Llorente, F., Hernández-Blanco, C., Jordá, L., Pagán,
I., ... & Molina, A. (2009). The ERECTA receptor-like kinase regulates cell wall–mediated
resistance to pathogens in Arabidopsis thaliana. *Molecular plant-microbe interactions*, *22*(8),
953-963.

Saucet, S. B., Ma, Y., Sarris, P. F., Furzer, O. J., Sohn, K. H., & Jones, J. D. (2015). Two
linked pairs of Arabidopsis TNL resistance genes independently confer recognition of bacterial
effector AvrRps4. *Nature communications*, *6*(1), 1-12.

Sekhwal, M. K., Li, P., Lam, I., Wang, X., Cloutier, S., & You, F. M. (2015). Disease resistance
gene analogs (RGAs) in plants. *International journal of molecular sciences*, *16*(8), 19248-
19290.

Sela, H., Cheng, J., Jun, Y., Nevo, E., & Fahima, T. (2009). Divergent diversity patterns of
NBS and LRR domains of resistance gene analogs in wild emmer wheat
populations. *Genome*, *52*(6), 557-565.

Shiu, S. H., & Bleecker, A. B. (2003). Expansion of the receptor-like kinase/Pelle gene family
and receptor-like proteins in Arabidopsis. *Plant physiology*, *132*(2), 530-543.

Sinapidou, E., Williams, K., Nott, L., Bahkt, S., Tör, M., Crute, I., ... & Beynon, J. (2004). Two
TIR: NB: LRR genes are required to specify resistance to Peronospora parasitica isolate Cala2
in Arabidopsis. *The Plant Journal*, *38*(6), 898-909.

Soltis, P. S., & Soltis, D. E. (2009). The role of hybridization in plant speciation. *Annual review*
of plant biology, *60*, 561-588.

Soubrier, J., Steel, M., Lee, M. S., Der Sarkissian, C., Guindon, S., Ho, S. Y., & Cooper, A.
(2012). The influence of rate heterogeneity among sites on the time dependence of molecular
rates. *Molecular Biology and Evolution*, *29*(11), 3345-3358.

Staal, J., Kaliff, M., Bohman, S., & Dixelius, C. (2006). Transgressive segregation reveals two
Arabidopsis TIR-NB-LRR resistance genes effective against Leptosphaeria maculans, causal
agent of blackleg disease. *The Plant Journal*, *46*(2), 218-230.

Staal, J., Kaliff, M., Dewaele, E., Persson, M., & Dixelius, C. (2008). RLM3, a TIR domain
encoding gene involved in broad-range immunity of Arabidopsis to necrotrophic fungal
pathogens. *The Plant Journal*, *55*(2), 188-200.

Tarr, D. E. K., & Alexander, H. M. (2009). TIR-NBS-LRR genes are rare in monocots: evidence
from diverse monocot orders. *BMC research notes*, *2*(1), 197.

Thatikunta, R., Sankar, A. S., Sreelakshmi, J., Palle, G., Leela, C., Rani, C. V. D., ... &
Dudhe, M. Y. (2016). Utilization of in silico EST–SSR markers for diversity studies in castor
(Ricinus communis L.). *Physiology and molecular biology of plants*, *22*(4), 535-545.

Thompson, J. D., Higgins, D. G., & Gibson, T. J. (1994). CLUSTAL W: improving the sensitivity
of progressive multiple sequence alignment through sequence weighting, position-specific
gap penalties and weight matrix choice. *Nucleic acids research*, *22*(22), 4673-4680.

Thompson, J. N., and Burdon, J. J. (1992). Gene-for-gene coevolution between plants and
parasites. *Nature*, *360*(6400), 121-125.

Tör, M., Lotze, M. T., & Holton, N. (2009). Receptor-mediated signalling in plants: molecular
patterns and programmes. *Journal of experimental botany*, *60*(13), 3645-3654.

Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B. C., Remm, M., & Rozen,
S. G. (2012). Primer3—new capabilities and interfaces. *Nucleic acids research*, *40*(15), e115-
e115.

Wan, L., Essuman, K., Anderson, R. G., Sasaki, Y., Monteiro, F., Chung, E. H., ... & Nishimura,
M. T. (2019). TIR domains of plant immune receptors are NAD+-cleaving enzymes that
promote cell death. *Science*, *365*(6455), 799-803.

Wang, X., & Wang, L. (2016). GMATA: an integrated software package for genome-scale SSR
mining, marker development and viewing. *Frontiers in plant science*, *7*, 1350.

Warren, R. F., Henk, A., Mowery, P., Holub, E., & Innes, R. W. (1998). A mutation within the
leucine-rich repeat domain of the Arabidopsis disease resistance gene RPS5 partially
suppresses multiple bacterial and downy mildew resistance genes. *The Plant Cell*, *10*(9),
1439-1452.
743
744 Williams, M.E. (June 16, 2011). Introduction to Phytohormones. Teaching Tools in Plant
745 Biology: Lecture Notes. The Plant Cell (online), doi/10.1105/tpc.110.tt0310.
746
747 Xie, D. X., Feys, B. F., James, S., Nieto-Rostro, M., & Turner, J. G. (1998). COI1: an
748 Arabidopsis gene required for jasmonate-regulated defense and fertility. *Science*, *280*(5366),
749 1091-1094.
750
751 Yang, Z. (1995). A space-time process model for the evolution of DNA
752 sequences. *Genetics*, *139*(2), 993-1005.
753
754 Zbierzak, A. M., Porfirova, S., Griebel, T., Melzer, M., Parker, J. E., & Dörmann, P. (2013). A
755 TIR–NBS protein encoded by A rabidopsis C hilling S ensitive 1 (CHS 1) limits chloroplast
756 damage and cell death at low temperature. *The Plant Journal*, *75*(4), 539-552.
757
758 Zdobnov, E. M., & Apweiler, R. (2001). InterProScan–an integration platform for the signature-
759 recognition methods in InterPro. *Bioinformatics*, *17*(9), 847-848.
760
761 Zhai, C., Lin, F., Dong, Z., He, X., Yuan, B., Zeng, X., ... & Pan, Q. (2011). The isolation and
762 characterization of Pik, a rice blast resistance gene which emerged after rice
763 domestication. *New Phytologist*, *189*(1), 321-334.
764
765 Zhang, X., Liang, P., & Ming, R. (2016). Genome-wide identification and characterization of
766 nucleotide-binding site (NBS) resistance genes in pineapple. *Tropical Plant Biology*, *9*(3), 187-
767 199.
768
769 Zhang, Z., Wu, Y., Gao, M., Zhang, J., Kong, Q., Liu, Y., ... & Zhang, Y. (2012). Disruption of
770 PAMP-induced MAP kinase cascade by a Pseudomonas syringae effector activates plant
771 immunity mediated by the NB-LRR protein SUMM2. *Cell host & microbe*, *11*(3), 253-263.
772
773 Zhang, Z., Xie, W., Zhao, Y., Zhang, J., Wang, N., Ntakirutimana, F., ... & Wang, Y. (2019).
774 EST-SSR marker development based on RNA-sequencing of E. sibiricus and its application
775 for phylogenetic relationships analysis of seventeen Elymus species. *BMC plant
776 biology*, *19*(1), 235.
777
778 Zipfel, C., Kunze, G., Chinchilla, D., Caniard, A., Jones, J. D. G., Boller, T., & Felix, G. (2006).
779 Perception of the Bacterial PAMP EF-Tu by the Receptor EFR Restricts Agrobacterium-
780 Mediated Transformation. Cell, 125(4), 749–760.
781

785 Table 1. List of source transcriptome sequences and their predicted number of RGAs.
786

Source SRA files	Mango variety	Tissue	NBS coding								RLP	RLK	TM-CC	Total
			NBS	CNL	TNL	CN	TN	NL	TX	Other				
Hong et al., 2016	'ZILL'	Fr	78	9	1	15	3	17	11	0	45	150	14	343
SRR1448302	'KEITT'	Fr	33	9	2	17	6	22	10	1	43	136	6	285
SRR1561197	'KENT'	Fr	20	0	0	3	1	1	4	0	15	68	12	124
SRR1562177	'KENT'	Fr	15	0	0	3	0	1	9	0	10	41	8	87
SRR1956775	'AMRAPALI'	L	40	1	0	9	1	2	11	0	10	21	6	101
SRR2159471	'TOMMY ATKINS'	Fl, L, Fr, S	74	7	1	15	3	18	12	2	78	223	21	454
SRR2162836	'AMIN ABRAHIMPUR'	Fl, L, Fr, S	71	10	1	11	4	23	18	1	76	216	22	453
SRR2162878	'BURMA'	Fl, L, Fr, S	24	0	0	3	0	2	11	0	37	112	11	200
SRR2162889	''M. CASTURI "PURPLE"	Fl, L, Fr, S	85	6	1	7	2	14	19	1	67	177	20	399
SRR2162908	'NEELUM'	Fl, L, Fr, S	66	7	1	14	5	18	4	1	72	214	20	422
SRR2162919	'THAI EVERBEARING'	Fl, L, Fr, S	80	10	2	19	1	13	15	2	71	210	24	447
SRR2162929	'TURPENTINE'	Fl, L, Fr, S	67	6	0	10	5	15	14	2	71	186	23	399
SRR2162953	'TURPENTINE', 'THAI EVERBEARING', 'NEELUM', 'M. CASTURI "PURPLE"', 'BURMA', 'AMIN ABRAHIMPUR' (Pooled)	Fl	64	11	1	17	1	24	20	4	66	225	17	450
SRR2165756	'TOMMY ATKINS'	S	8	0	0	0	0	1	7	0	8	28	4	56
SRR2736811	'ATAULFO'	Fr	36	4	0	11	0	5	8	0	19	69	18	170
SRR3192873	'AMRAPALI'	L	89	7	1	18	4	15	15	4	49	193	16	411
SRR3288569	'KEITT'	B	75	4	1	17	2	13	17	0	61	157	21	368
SRR3319054	'CHAUSA'	L	68	7	1	19	6	15	12	2	60	187	17	394
SRR3359450	'DASHEHARI'	Fr	38	3	1	7	0	10	7	1	19	130	14	230
SRR5966284	'AMRAPALI'	Fl	32	6	0	16	5	11	14	1	45	212	21	363
SRR5966306	'AMRAPALI'	Fl	45	1	0	16	3	11	18	1	52	246	25	418

SRR8449851	'CHOKANAN'	Fr	53	1	0	5	2	8	10	0	20	88	12	199
SRR8449858	'GOLDEN PHOENIX' and 'WATER LILY' (Pooled)	Fr	103	10	2	19	6	23	15	2	56	168	12	416
SRR8926025	MANGO cv. 1243	R; L; St, B, Fl; Fr	29	0	0	8	2	3	7	1	31	113	8	202
Total			1293	119	16	279	62	285	288	26	1081	3570	372	**7,391**

787 Fr: Fruit; Fl: Flower, L: Leaf, S: Seed; B: Buds; St: Stem; R: Roots

796   Table 2. The distribution of core RGAs of mango and its comparison to other major crops.

RGA codes		Mango (*Mangifera indica*)[a]	Cacao (*Theobroma cacao*)[b]	Papaya (*Carica papaya*)[b]	Tomato (*Solanum lycopersicum*)[b]	Corn (*Zea mays*)[b]	Rice (*Oryza sativa*)[b]	Banana (*Musa acuminata*)[b]
NBS	NBS	53	16	11	64	37	47	11
CC-NBS-LRR	CNL	17	173	10	70	52	222	50
TIR-NBS-LRR	TNL	2	14	4	19	0	0	0
CC-NBS	CN	29	13	5	8	23	38	7
TIR-NBS	TN	4	3	1	7	2	3	2
NBS-LRR	NL	27	66	21	91	49	196	44
TIR with unknown domain	TX	17	8	5	12	3	3	3
Other NBS-encoding	Other	6	0	3	1	1	0	0
RLP	RLP	158	167	27	86	63	118	107
RLK	RLK	362	640	299	512	671	867	530
TM-CC	TM-CC	72	181	81	133	161	158	138
Total RGAs		**747**	**1,281**	**467**	**1,003**	**1,062**	**1,652**	**892**

797   [a] inferred from this paper based on transcriptome data; [b] from Li et al. (2016) based on genome-wide RGA analysis

799 Table 3. Some mango RGAs with homology to known Resistance (*R*)/defense proteins

800 against pathogens and insects with homology to the entries in Uniprot/SwissProt database

801 based on BLASTp hits.

Mango *R*/defense protein	RGA domain	e-Value	R/defense protein name/description	Resistance/ defense response to
Zill_394_c0_g1.p2	NBS	2.41E-42		
SRR1448302_13791_c0_g1.p1	NL	3.90E-36		
SRR1448302_13356_c0_g1.p2	TN	2.76E-31	Putative disease resistance protein At4g11170	Pathogen and insect
SRR2159471_30596_c0_g1.p1	TX	1.18E-41		
SRR2162908_17959_c0_g1.p1	OTHER	7.44E-33		
SRR2162889_655_c0_g2.p3	NBS	1.22E-16	Disease resistance protein RRS1; Disease resistance protein RCH2; Probable WRKY transcription factor 52; Resistance to Colletotrichum higginsianum 2 protein; Resistance to Ralstonia solanacearum 1 protein	Pathogen
SRR2162878_10232_c0_g1.p1	NBS	8.95E-35	Disease resistance protein RRS1B; Probable WRKY transcription factor 16; WRKY DNA-binding protein 16	Pathogen
Zill_10069_c0_g1.p1	NBS	1.55E-20		
Zill_15696_c0_g1.p1	NL	6.92E-48		
Zill_16228_c0_g1.p1	TN	2.52E-65		
Zill_4100_c0_g1.p1	TX	2.37E-31	TMV resistance protein N	Pathogen
SRR1448302_8769_c0_g1.p1	CN	5.82E-43		
SRR1448302_10123_c0_g2.p1	TNL	0		
SRR2159471_13808_c0_g1.p1	OTHER	1.55E-53		
SRR3359450_14591_c0_g1.p1	RLP	9.35E-39		
Zill_7264_c0_g1.p1	TX	4.74E-39	Disease resistance protein TAO1; Protein TARGET OF AVRB OPERATION 1	Pathogen
SRR1448302_10424_c0_g1.p1	CN	2.98E-43	Disease resistance protein SUMM2; Disease resistance protein At1g12280; Protein SUPPRESSOR OF MKK1 MKK2 2	Pathogen
SRR1448302_12885_c0_g1.p1	NBS	1.28E-10		
SRR2162929_3892_c0_g1.p1	NL	1.04E-98		
Zill_3443_c0_g2.p1	NBS	5.51E-13	Disease resistance protein RFL1; RPS5-like protein 1; pNd13/pNd14	Pathogen
SRR2162919_15595_c0_g2.p2	CN	9.65E-21		
SRR2162929_6841_c0_g1.p1	NL	1.80E-28		
SRR5966306_7909_c0_g1.p1	TX	1.09E-21	Disease resistance protein RPS6; Resistance to Pseudomonas syringae 6	Pathogen
SRR8449858_33757_c0_g1.p1	NBS	1.44E-17	Disease resistance protein RPS6; Resistance to Pseudomonas syringae 6	Pathogen
SRR1448302_7186_c0_g1.p1	NBS	3.86E-36	Disease resistance protein RPS5; Resistance to Pseudomonas syringae protein 5; pNd3/pNd10	Pathogen
SRR2162889_11315_c0_g1.p1	CNL	0		
SRR2162953_17628_c0_g1.p1	NL	2.01E-42		
Zill_10085_c0_g1.p1	NBS	9.29E-55		
SRR2162908_5437_c0_g1.p1	NL	4.04E-62	Disease resistance protein RPS2; Resistance to Pseudomonas syringae protein 2	Pathogen
SRR2162919_3439_c1_g1.p1	CNL	9.91E-78		
SRR2162919_4996_c0_g1.p2	RLP	5.55E-14		
Zill_5583_c0_g1.p1	CN	3.98E-21		
Zill_10742_c0_g1.p1	NBS	1.36E-23	Disease resistance protein RPM1; Resistance to Pseudomonas syringae protein 3	Pathogen
Zill_30931_c0_g1.p1	NL	1.80E-50		
SRR1448302_9780_c0_g1.p1	CNL	4.05E-96		
SRR2159471_6253_c0_g1.p1	TX	2.23E-28	Disease resistance protein RPP5; Protein RECOGNITION OF PERONOSPORA PARASITICA 5	Pathogen and insect
SRR3319054_18081_c0_g1.p1	OTHER	1.42E-35		
Zill_4463_c0_g1.p2	RLP	4.77E-07	Probable disease resistance protein At4g19530	Pathogen
SRR2162919_8948_c0_g2.p1	TN	3.95E-30	Probable disease resistance protein At4g19520	Pathogen

ID	Type	E-value	Protein	Category
SRR1448302_1069_c0_g1.p1	RLP	9.17E-55	Disease resistance protein RPP2B	Pathogen
SRR2159471_12447_c0_g2.p1	NL	5.12E-101		
SRR2162953_222_c0_g1.p2	NL	7.60E-45	Disease resistance protein RPP2A	Pathogen
Zill_19434_c0_g1.p1	NBS	1.93E-53	Disease resistance protein RPP13; Resistance to Peronospora parasitica protein 13	Pathogen
Zill_10742_c0_g1.p3	NL	2.70E-20		
SRR2159471_23664_c0_g1.p1	CN	3.57E-29		
SRR2736811_8634_c0_g1.p1	CNL	4.58E-170		
SRR2162889_14411_c0_g1.p1	NBS	2.47E-24	Probable disease resistance protein RPP1; Protein RECOGNITION OF PERONOSPORA PARASITICA 1	Pathogen
SRR2162953_10793_c0_g1.p1	TX	2.59E-33		
SRR3288569_10876_c0_g3.p1	CN	2.07E-19	Probable disease resistance RPP8-like protein 2	Pathogen
SRR2162929_192_c0_g1.p3	TX	4.57E-38	Disease resistance protein RLM3; Protein RESISTANCE TO LEPTOSPHAERIA MACULANS 3	Pathogen
SRR2162878_1969_c0_g1.p1	RLP	5.14E-23	Disease resistance protein RML1B; Protein RESISTANCE TO LEPTOSPHAERIA MACULANS 1B	Pathogen
SRR2162919_4480_c1_g1.p1	NBS	2.22E-20		
SRR2162953_20108_c0_g1.p1	TX	3.93E-32		
SRR2162878_10232_c0_g1.p2	TX	3.23E-26	Disease resistance protein RML1A; Protein RESISTANCE TO LEPTOSPHAERIA MACULANS 1A	Pathogen
SRR2159471_14993_c0_g1.p1	OTHER	1.15E-32	Disease resistance-like protein DSC2; Protein DOMINANT SUPRESSOR OF CAMTA3 NUMBER 2	Pathogen and insect
SRR2159471_10112_c0_g1.p1	TX	5.42E-43		
Zill_8229_c0_g1.p2	NBS	9.39E-19	Disease resistance-like protein DSC1; Protein DOMINANT SUPPRESSOR OF CAMTA3 NUMBER 1	Pathogen and insect
SRR1448302_13356_c0_g1.p1	TX	1.50E-47		
SRR2159471_16556_c0_g1.p1	RLP	5.16E-27		
SRR2162836_1731_c0_g2.p1	NL	1.04E-101		
SRR2162889_14475_c0_g2.p1	TN	2.61E-56		
SRR2162919_11945_c0_g1.p1	TNL	0		
SRR2162953_14082_c0_g1.p2	OTHER	1.51E-29		
SRR2736811_2638_c0_g1.p1	CNL	6.55E-123		
SRR3192873_14562_c0_g2.p1	OTHER	0		
SRR2159471_15747_c0_g1.p2	NBS	4.36E-19	Disease resistance protein Pik-2; Pik-2 blast resistance protein	Pathogen
SRR2159471_15747_c0_g1.p1	NBS	2.00E-26	Disease resistance protein RGA5; Os11gRGA5; SasRGA5	Pathogen
Zill_11941_c0_g1.p1	NBS	5.81E-29	Putative disease resistance protein RGA4; RGA4-blb	Pathogen
SRR1956775_13322_c0_g1.p1	CN	4.06E-28		
SRR2162908_1538_c0_g1.p1	CNL	4.56E-115		
SRR3288569_1324_c0_g1.p1	NL	3.37E-158		
Zill_2822_c0_g1.p1	CN	3.84E-91	Putative disease resistance protein RGA3; Blight resistance protein B149; RGA1-blb	Pathogen
Zill_2317_c0_g1.p5	NBS	1.35E-26		
SRR1448302_19774_c0_g1.p1	CNL	0		
SRR1448302_12653_c0_g1.p1	NL	1.12E-77		
Zill_40171_c0_g1.p1	NBS	2.32E-24	Disease resistance protein RGA2; Blight resistance protein RPI; RGA2-blb	Pathogen
Zill_5557_c0_g1.p1	NL	2.55E-22		
Zill_12068_c0_g1.p2	RLP	1.75E-05		
SRR1956775_2968_c1_g2.p1	CN	4.43E-57		
Zill_11208_c0_g1.p1	CN	3.21E-88	Putative disease resistance protein RGA1; RGA3-blb	Pathogen
Zill_2928_c3_g1.p1	NBS	1.21E-50		
Zill_4058_c0_g1.p1	NL	8.99E-92		
SRR2159471_1162_c2_g3.p1	CNL	5.33E-138		
SRR5966284_6355_c0_g1.p2	NBS	7.60E-35	Putative late blight resistance protein homolog R1A-10	Pathogen
SRR1448302_30603_c0_g1.p1	NL	0		Pathogen

SRR3359450_19530_c0_g1.p1	NBS	6.35E-94	Disease resistance RPP13-like protein 4; Disease resistance protein ZAR1; Protein HOPZ-ACTIVATED RESISTANCE 1	
SRR2159471_29188_c0_g1.p1	NBS	1.87E-64		
SRR2162836_25848_c0_g1.p1	CN	1.77E-26	Putative disease resistance RPP13-like protein 3	Pathogen
SRR2162836_22262_c0_g1.p1	NL	5.61E-100		
Zill_8566_c0_g1.p1	NBS	2.82E-22		
SRR2159471_14284_c0_g1.p1	CNL	1.84E-167	Putative disease resistance RPP13-like protein 2	Pathogen
SRR3192873_20475_c0_g1.p1	RLP	5.06E-42		
SRR3319054_34638_c0_g1.p1	NL	1.14E-20		
Zill_11317_c0_g1.p1	NBS	8.38E-21		
Zill_2317_c0_g1.p2	RLP	8.51E-48		
SRR1448302_10269_c0_g2.p1	CN	7.02E-63	Putative disease resistance RPP13-like protein 1	Pathogen
SRR1448302_1339_c0_g1.p1	CNL	4.16E-158		
SRR1448302_2626_c0_g1.p1	NL	0		
SRR2162836_19327_c0_g1.p1	TM-CC	4.13E-25		
SRR2159471_25754_c0_g1.p1	RLK	1.28E-92	LEAF RUST 10 DISEASE-RESISTANCE LOCUS RECEPTOR-LIKE PROTEIN KINASE-like 2.5; Probable receptor-like serine/threonine-protein kinase LRK10L-2.5	Pathogen
SRR1448302_15170_c0_g1.p1	RLK	5.94E-71	LEAF RUST 10 DISEASE-RESISTANCE LOCUS RECEPTOR-LIKE PROTEIN KINASE-like 2.4; Probable receptor-like serine/threonine-protein kinase LRK10L-2.4	Pathogen
SRR3319054_22570_c0_g1.p1	RLK	7.38E-41	LEAF RUST 10 DISEASE-RESISTANCE LOCUS RECEPTOR-LIKE PROTEIN KINASE-like 2.2; Probable receptor-like serine/threonine-protein kinase LRK10L-2.2	Pathogen
SRR2159471_9027_c0_g2.p1	RLK	5.60E-151	LEAF RUST 10 DISEASE-RESISTANCE LOCUS RECEPTOR-LIKE PROTEIN KINASE-like 2.1; Probable receptor-like serine/threonine-protein kinase LRK10L-2.1	Pathogen
SRR2162836_4545_c0_g1.p1	RLK	0	LEAF RUST 10 DISEASE-RESISTANCE LOCUS RECEPTOR-LIKE PROTEIN KINASE-like 1.5; Probable receptor-like serine/threonine-protein kinase LRK10L-1.5	Pathogen
SRR1448302_13187_c0_g1.p1	RLK	1.62E-159	LEAF RUST 10 DISEASE-RESISTANCE LOCUS RECEPTOR-LIKE PROTEIN KINASE-like 1.3; Probable receptor-like serine/threonine-protein kinase LRK10L-1.3	Pathogen
Zill_2156_c0_g1.p1	RLK	4.69E-14	LEAF RUST 10 DISEASE-RESISTANCE LOCUS RECEPTOR-LIKE PROTEIN KINASE-like 1.2; Probable receptor-like serine/threonine-protein kinase LRK10L-1.2	Pathogen
Zill_5127_c0_g2.p1	RLK	1.43E-111	LEAF RUST 10 DISEASE-RESISTANCE LOCUS RECEPTOR-LIKE PROTEIN KINASE-like 1.1; Probable receptor-like serine/threonine-protein kinase LRK10L-1.1	Pathogen
Zill_13036_c0_g1.p1	RLK	1.86E-104	Rust resistance kinase Lr10; Probable receptor-like serine/threonine-protein kinase LRK10	Pathogen
SRR2162878_1544_c0_g2.p1	RLP	1.72E-119	LRR receptor-like serine/threonine-protein kinase ERECTA; Protein QUANTITATIVE RESISTANCE TO PLECTOSPHAERELLA 1; Protein QUANTITATIVE RESISTANCE TO RALSTONIA SOLANACEARUM 1; Protein TRANSPIRATION EFFICIENCY 1	Pathogen
SRR3288569_12948_c0_g1.p2	RLK	5.39E-133		
SRR2162878_2968_c0_g1.p1	RLK	2.12E-122	Serine/threonine-protein kinase EDR1; MAPKK kinase EDR1; Protein ENHANCED DISEASE RESISTANCE 1; AtEDR1; Serine/threonine/tyrosine-protein kinase 10	Pathogen
SRR2162889_24778_c0_g2.p1	NBS	1.15E-17	Disease resistance protein ADR2; Protein ACTIVATED DISEASE RESISTANCE 2; Protein WHITE RUST RESISTANCE 4	Pathogen
Zill_1197_c0_g1.p1	RLP	2.34E-85	Protein TRANSPORT INHIBITOR RESPONSE 1; Weak ethylene-insensitive protein 1	Pathogen and insect
SRR2162908_7086_c0_g1.p1	RLP	1.27E-144	Coronatine-insensitive protein 1; COI-1; F-box/LRR-repeat protein 2; AtCOI1; AtFBL2	Pathogen and insect

803 Table 4. Other mango RGAs with homology to proteins in Uniprot/SwissProt database based

804 on BLASTp hits and gene ontology analysis with "response to insects" annotation.

Mango contig ID	RGA domain	e-Value	Description
SRR2162908_12681_c0_g1.p1	RLK	0	Calmodulin-binding receptor kinase CaMRLK;
SRR8449851_23663_c0_g1.p1	RLP	6.40E-66	Calmodulin-binding receptor-like kinase; AtCaMRLK; Protein MATERNAL EFFECT EMBRYO ARREST 62
SRR5966284_6489_c0_g1.p1	RLK	4.04E-54	Cysteine-rich receptor-like protein kinase 18; Cysteine-rich RLK18
SRR2162953_2879_c0_g1.p2	RLK	7.43E-20	Putative cysteine-rich receptor-like protein kinase 43; Cysteine-rich RLK43
SRR5966306_10141_c0_g1.p1	RLK	3.68E-52	Cold-responsive protein kinase 1
SRR2162836_4413_c0_g3.p1	RLK	0	Putative receptor protein kinase ZmPK1
SRR2162836_3219_c0_g1.p1	RLK	0	Probable LRR receptor-like serine/threonine-protein kinase RFK1; Receptor-like kinase in flowers 1
SRR8449851_14359_c0_g1.p1	RLK	0	Probable LRR receptor-like serine/threonine-protein kinase RKF3; Receptor-like kinase in flowers 3
SRR2159471_39050_c1_g1.p1	RLK	0	Probable receptor-like protein kinase At1g11050
SRR3192873_14159_c0_g1.p1	RLK	0	C-type lectin receptor-like tyrosine-protein kinase At1g52310
SRR8449858_41086_c0_g1.p1	RLK	9.52E-66	Probable LRR receptor-like serine/threonine-protein kinase At1g53420
SRR1561197_1147_c0_g1.p1	RLK	8.21E-87	Probable LRR receptor-like serine/threonine-protein kinase At1g56130
SRR3192873_4970_c0_g1.p1	RLK	0	Probable LRR receptor-like serine/threonine-protein kinase At1g07650
SRR1448302_14204_c1_g1.p1	RLK	0	Probable leucine-rich repeat receptor-like serine/threonine-protein kinase At3g14840
Zill_8965_c0_g1.p1	RLK	1.47E-47	Probable LRR receptor-like serine/threonine-protein kinase At1g53430
SRR2159471_11990_c0_g1.p1	RLK	8.55E-62	Probable LRR receptor-like serine/threonine-protein kinase At5g45780
Zill_7246_c0_g2.p2	RLK	2.96E-117	Probable LRR receptor-like serine/threonine-protein kinase At1g56140

839 Figure 1. Distribution of the classifications based on conserved domains and motifs of core

840 resistance gene analogs (RGA) detected in mango transcriptome *de novo* assemblies.

Figure 2. Resistance gene analogs (RGAs) expression profile from mango transcriptomes

visualized as heatmap. Count values were normalized within samples as transcripts per million

(TPM) using RSEM. Y-axis correspond to RGA gene IDs. X-axis correspond to RNA-seq raw

read samples: *SRR_1. Hong et al., 2016 (ZILL), 2. SRR1448302 (KEITT), 3. SRR1561197 (KENT), 4 SRR1562177 (KENT),*

5. SRR1956775 (AMRAPALI), 6. SRR2159471 (TOMMY ATKINS), 7. SRR2162836 (AMIN ABRAHIMPUR), 8. SRR2162878

(BURMA), 9. SRR2162889 (M. CASTURI "PURPLE"), 10. SRR2162908 (NEELUM), 11. SRR2162919 (THAI EVERBEARING),

12. SRR2162929 (TURPENTINE), 13. SRR2162953 (Pooled – TURPENTINE, THAI EVERBEARING, NEELUM, M. CASTURI

"PURPLE", BURMA, AMIN ABRAHIMPUR), 14. SRR2165756 (TOMMY ATKINS), 15. SRR2736811 (ATAULFO), 16.

SRR3192873 (AMRAPALI), 17. SRR3288569 (KEITT), 18. SRR3319054 (CHAUSA), 19. SRR3359450 (DASHEHARI), 20.

SRR5966284 (AMRAPALI), 21. SRR5966306 (AMRAPALI), 22. SRR8449851 (CHOKANAN) 23. SRR8449858 (GOLDEN

PHOENIX and WATER LILY), 24. SRR8926025 (MANGO cv. 1243)

Figure 3. Gene ontology (GO) distribution of the expressed resistance gene analogs (RGAs) in mango based on molecular function (MF).

934 Figure 4. Gene ontology (GO) distribution of the expressed resistance gene analogs

935 (RGAs) in mango based on biological processes (BP).

975 Figure 5. Gene ontology (GO) distribution of the expressed resistance gene analogs (RGAs)

976 in mango based on cellular component (CC).

Figure 6. Maximum likelihood phylogenetic tree constructed   using IQ-TREE from the sequence alignment of the core resistance gene analogs (RGAs) generated from *de novo* transcriptome assemblies of mango. Best-fit model was selected according to Bayesian Information Criterion (BIC) using ModelFinder (Kalyaanamoorthy et al., 2017). General 'Variable Time' (VT) matrix (Mueller and Vingron, 2004) with empirical amino acid frequencies (+F) and FreeRate (+R5) rate heterogeneity across sites (Yang 1995; Soubrier et al. 2012) was used to generate the tree, tested with 1000 replicates of ultrafast bootstrapping (Hoang et al. 2017). The numbers correspond to the clades.

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

Figure 7. Distribution of the different repeat motifs that were detected in the mango RGA transcriptome.

1024

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039   Figure 8. Paired-repeat SSR motif distribution of the mango RGA transcriptome.

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055 Figure 9. Distribution of EST-SSR markers designed across the different RGA domains

1056 identified in *de novo* mango transcriptomes.