1
2
3
4    **Collective sperm movements are shaped by post-copulatory sexual**
5    **selection and phylogenetic history in *Peromyscus* mice**
6
7

8    Kristin A. Hook, W. David Weber, Heidi S. Fisher*
9
10
11
12
13   Department of Biology, University of Maryland, 1200 Biology-Psychology Building,
14   4094 Campus Drive, College Park, MD 20742, U.S.A.
15
16
17   * Heidi S. Fisher
18   Email: hsfisher@umd.edu
19
20
21
22

23   *Keywords:*
24
25   mating systems; sexual selection; sperm competition; sperm conjugation; sperm motility
26
27

28   *Author Contributions:*
29
30   KAH and HSF conceived of the study, designed experiments, and interpreted results;
31   KAH and WWD collected the data, KAH carried out the statistical analyses; all authors
32   wrote the manuscript, gave final approval for publication, and agree to be held
33   accountable for the work presented.
34
35

36   *Funding:*
37
38   This work was supported by a Eunice Kennedy Shriver National Institute of Child Health
39   and Human Development K99/R00 Pathway to Independence Award to HSF
40   [R00HD071972] and a National Science Foundation Postdoctoral Research Fellowship to
41   KAH [1711817].

**Abstract**

Sperm of some species form motile, coordinated groups as they migrate through the female

reproductive tract to the site of fertilization. This collective motion is predicted to improve sperm

swimming performance and therefore may be beneficial in a competitive context, but limited

evidence supports this theory. Here we examine sperm aggregates across closely-related species of

*Peromyscus* mice that naturally vary by mating system, and thus sperm competition intensity. We

find that phylogenetic history predicts the likelihood that sperm will aggregate, and that relative

testis size is negatively associated with variation in number of aggregated cells, suggesting that

sperm competition has a stabilizing effect on sperm group size. Moreover, we show that sperm

aggregates are kinematically beneficial for some species but costly for others, and these

differences are largely dependent on the orientation and composition of sperm within the groups.

In addition, when we compared sperm of the two sister-species that aggregate most frequently, we

find that sperm from the species that evolved under intense sperm competition forms aggregates

with more efficient geometry more frequently than sperm from its monogamous congener. These

results are consistent with the prediction that sperm aggregation evolved to improve motility in a

competitive context; however, when monogamy evolved secondarily, relaxed sexual selection

allowed for less efficient strategies to persist. Together, our findings in *Peromyscus* reveal that

collective sperm behavior is likely to evolve rapidly and is shaped by changes in the selective

regime.

**Introduction**

Sperm cells are one of the most diverse cell types in nature and exhibit striking variation both

within and across taxa (Pitnick et al. 2009). In addition to being morphologically diverse, sperm

may exhibit complex, emergent behaviors, including sperm conjugation, which occurs when two

or more cells join together for motility or transport through the female reproductive tract before

dissociating prior to fertilization (Pitnick et al. 2009; Higginson and Pitnick 2011). Although

relatively rare, these sperm-sperm interactions have evolved multiple times across independent

lineages of internally fertilizing species (Immler 2008; Pitnick et al. 2009; Higginson and Pitnick

2011), yet the adaptive significance of these gametic interactions remains unclear for many taxa.

One functional hypothesis posits that sperm aggregates may be advantageous if the combined

force generated by multiple flagella enable sperm conjugates to swim faster than single cells

(Moore et al. 2002; Immler et al. 2007). Improved sperm motility can be beneficial if it enables

cells to move quickly through hostile environments within the female tract (Birkhead et al. 1993;

Pizzari and Foster 2008), and in competitive environments in which females mate with multiple

males, as sperm velocity is often correlated with competitive fertilization success (e.g., Birkhead et

al. 1999; Gage et al. 2004; Pizzari and Foster 2008). While some studies that have quantified the

motility of sperm aggregates have found empirical support for this hypothesis (e.g., Hayashi 1998;

Moore et al. 2002; Woolley et al. 2009; Fisher and Hoekstra 2010; Pearcy et al. 2014), others have

not (e.g., Ishijima et al. 1999; Tung et al. 2017). For example, sperm groups swim faster than

solitary cells in the Norway rat (*Rattus norvegicus*), yet house mouse (*Mus musculus*) sperm swim

slower as groups under identical experimental conditions (Immler et al. 2007). Another non-

mutually exclusive hypothesis is that sperm aggregation facilitates migration through viscous or

viscoelastic secretions of the female reproductive tract (Moore and Taggart 1995; Suarez 2016),

including cervical mucus (Hanson and Overstreet 1981), which has received some empirical

support (e.g., Moore and Taggart 1995; Hayashi 1998; Moore et al. 2002; but see Tung et al.

2017). In the grey short-tailed opossum (*Monodelphis domestica*), for example, sperm pairs swim

with greater motility than single sperm in viscous fluids; however, these sperm pairs were

artificially induced to uncouple (Moore and Taggart 1995) and therefore the comparison may not

be biologically informative (Pitnick et al. 2009). Additionally, sperm aggregation may protect

sensitive regions of the sperm, such as the acrosome, from damage and preserve sperm

functionality during passage through the male or female reproductive tracts (Phillips 1970; Tung et

al. 1980; Bedford et al. 1984; Higginson and Henn 2012), or enhance egg penetration during

fertilization (Mackie and Walker 1974; but see McGlinn et al. 1979). While it is often assumed

that coordinated sperm movements are adaptive, inconsistent findings and a wide diversity of

naturally variable sperm behavior have limited our understanding of the functional advantages of

sperm aggregation.

Multiple independent origins of sperm conjugation suggests that the functional consequences,

as well as mechanisms that regulate these cellular interactions, are likely to vary throughout nature

(reviewed in Higginson and Pitnick 2011). Indeed, the formation of sperm aggregates and the

number of grouped cells varies widely across taxa (Pizzari and Foster 2008). In mammals alone,

sperm of some species assemble during epididymal transport and are molecularly "glued" to one

another as bi-flagellate pairs in grey short-tailed opossums (Taggart et al. 1993), as bundles of

roughly 100 cells in monotremes (Nixon et al. 2016), or as organized rouleaux of five or more

cells in guinea pigs (*Cavia porcellus*; McGlinn et al. 1979). Conversely, mammalian sperm may

also assemble after ejaculation to form variably-sized groups. For instance, sperm may form

temporary clusters of up to sixteen cells in bulls (*Bos taurus*; Tung et al. 2017) or more fixed

groups of up to 30 cells in house mice (Immler et al. 2007), 50 cells in the Norway rat (Immler et

al. 2007), or thousands of cells in the wood mouse (*Apodemus sylvaticus*; Moore et al. 2002)

whereby the hook-shaped heads interlock or attach to the flagella of other sperm. For these latter

species in which sperm groups form after ejaculation, both single and aggregated sperm typically

co-occur, thus allowing for direct comparisons between collective and solitary sperm movements

within an ejaculate while controlling for within-male variability.

Closely-related species in the rodent genus, *Peromyscus,* produce sperm that naturally vary in

their collective behavior. Sperm of some species assemble temporary groups after ejaculation by

adhering to one another at their head region (Pearce et al. 2018) and disassemble prior to

fertilization (Fisher and Hoekstra 2010). In *P. maniculatus*, sperm selectively group with the most

closely-related cells to form motile groups of up to 30 cells (Fisher and Hoekstra 2010), but there

is a non-monotonic association between group size and swimming velocity, indicating that some

groups are faster than single cells but others are not (Fisher et al. 2014). Conversely, sperm

produced by their sister-species, *P. polionotus*, also form aggregates but do so indiscriminately

with related and unrelated cells (Fisher and Hoekstra 2010), are less likely to be optimally-sized,

and overall move in a less linear trajectory (Fisher et al. 2014). Intriguingly, sperm competition is

predicted to be more intense in *P. maniculatus* than in *P. polionotus* due to their different mating

systems. In *P. maniculatus*, both sexes mate with multiple partners, often in overlapping series just

minutes apart (Dewsbury 1985), and females frequently carry multiple-paternity litters in the wild

(Birdsall and Nash 1973), whereas both behavioral (Dewsbury 1981) and genetic data (Foltz 1981)

indicate that *P. polionotus* is strictly monogamous. Evidence suggests that monogamy has evolved

at least twice within the *Peromyscus* lineage (reviewed in Bedford and Hoekstra 2015; Turner et

al. 2010), thus enabling us to investigate if post-copulatory sexual selection, driven by female

mating behavior, has shaped the evolution of sperm aggregation (Moore et al. 2002) more broadly

across the *Peromyscus* lineage. In this study, we quantify intra- and inter-specific differences in

the size and performance of sperm aggregates under consistent, controlled conditions to examine

the evolution of collective sperm behavior and empirically test whether sperm aggregation

improves swimming performance.

**Materials and Methods**

(a) Sperm collection

We obtained captive *Peromyscus maniculatus bairdii*, *P. polionotus subgriseus, P. leucopus*,

*P. eremicus*, and *P. californicus insignis* from the Peromyscus Genetic Stock Center at the

University of South Carolina, and *P. gossypinus gossypinus* from Dr. Hopi Hoekstra at Harvard

University and housed them in same-sex cages at 22ºC on a 16L:8D cycle in accordance with

guidelines established by the Institutional Animal Care and Use Committee at the University of

Maryland in College Park (protocol R-Jul-18-38). We sought samples from all available captive

*Peromyscus* species and avoided wild-caught specimens to control for variation due to life

experience. We obtained sperm samples from sexually mature males and accounted for relatedness

among the focal males by assigning siblings a unique 'Family' ID. We euthanized males via

isoflurane overdose and cervical dislocation, then weighed each male and both testes (Mettler

Toledo, Switzerland). Next, we removed a single caudal epididymis, made several small incisions

in the tissue, and submersed it in sperm medium (Modified Sperm Washing Medium, Irvine

Scientific, USA) that was pre-warmed at 37ºC; to reduce differences in sperm density despite

natural variation in epididymal sizes, we varied the volume (50μl - 1000μl) based on tissue size

and accounted for these differences when estimating final sperm counts for each male. To collect

sperm, we agitated the tissue at 300rpm (ThermoMixer F1.5, Eppendorf, Germany) at 37ºC for ten

minutes, inverting the tube at the five- and ten-minute mark, then incubated the tissue undisturbed

for two minutes. Using pipette tips cut to create a wider opening, we collected live sperm cells for

analysis from just below the meniscus of the solution to enrich for the most motile sperm

(Magdanz et al. 2019). Next we estimated sperm density using a computer-assisted sperm analysis

(CASA) system (Ceros II Animal, Hamilton Thorne, USA) and verified with a Neubauer-

improved hemocytometer (Marienfeld, Germany) then diluted samples with pre-warmed medium

to reach a standard concentration of 300-400 cells summed across the five 5-second videos at

100X magnification for cell tracking optimality and efficiency.

(b) Live sperm analysis

To conduct live sperm observations, we gently reverse pipetted 4µl of the sperm solution into

12µl of pre-warmed medium on a plastic slide within a 9mm x 0.12mm imaging spacer (Grace

Bio-Labs, USA) and covered by a plastic cover slip. This set-up served as a control and represents

a 'low-viscosity' environment. To test sperm motility in a 'high-viscosity' environment, we

followed the same procedures except that we mixed 4µl of sperm solution with 12µl of pre-

warmed medium enriched with methylcellulose (Sigma Aldrich M 7140; 15cP, 2% in water;

Suarez and Dai 1992). We then recorded 5-second videos at 60 frames/sec on the CASA system,

capturing at least five videos per male but recorded additional videos for samples with lower

sperm density ($n = 57$) to ensure an adequate number of observed cells per male. Videos were

recorded at 59±16 minutes post-harvest from the epididymal tissue, dependent on dilutions. The

number of cells analyzed are reported in Table 1.

We characterized sperm aggregation by scoring CASA videos using direct observations

because the system tracks particles, and thus each track may represent a single cell or an

aggregate. We counted the number of cells represented by each track on at least three different

frames/track. From these data, we calculated the proportion of cells that aggregated for each male

by dividing the total number of aggregates by the total number of motile cells across all tracks

(Crawley 2013). Then we calculated the mean number of cells in aggregate (i.e., aggregate size)

by dividing the sum of cells in aggregate by the sum of aggregates, both across all tracks for each

male and across all males for each species. Finally, we calculated the coefficient of variation (CV)

for both the proportion and number of cells that aggregated within each species using the

following formula: (standard deviation/mean) x 100. In addition, to further characterize

differences in sperm aggregation for the two species whose sperm were observed to aggregate the

most extensively, *P. maniculatus* and *P. polionotus*, we qualitatively scored the composition and

orientation of cells within the sperm aggregates. For males within these species, we calculated the

proportion of aggregates that were: 'aligned' (all sperm adhered to one another in a head-to-

flagella orientation and included no immotile, morphologically abnormal or damaged cells),

'defective' (one or more sperm were abnormal, immotile, or stuck to the slide), or 'opposed' (all

sperm are normal and motile but were not oriented in the same direction), the latter of which

included star-shaped aggregates (Fisher et al. 2014).

We recorded the following metrics for each recorded track (i.e., single sperm and aggregates):

*straight-line velocity* (VSL; calculated using the straight-line distance between the first and last

detected positions of the sperm head, divided by the time taken to swim the track; also known as

average velocity), *curvilinear velocity* (VCL; calculated using the summed distance between the

sperm head positions in each frame divided by the time taken to travel the track; also known as

speed), *average path velocity* (VAP; the time-averaged velocity of the sperm head along its

average path), and *linearity* (LIN; the ratio of VSL to VCL to measure the straightness of the

trajectory; World Health Organization 2010). We calculated the mean of each kinematic parameter

for both single cells and sperm aggregates separately for each of three populations of sperm cells:

all cells, motile cells (i.e., devoid of visually inspected tracks in which cells were unmoving, stuck,

or featured an obvious morphological abnormality such as a kinked midpiece), and progressively

motile cells (i.e., motile cells with a VSL $\geq$ 25μm/sec). We used these data to calculate the

proportion of motile aggregates by dividing the sum of motile aggregates by the total number of

sperm aggregates, and the proportion of progressively motile aggregates by dividing the sum of

progressive aggregates by the total number of sperm aggregates for each male (Crawley 2013). For

our kinematic analyses, we focused on the motile sperm dataset to eliminate artifacts from

damaged or dead cells, and the total sperm population to quantify frequency of aggregation and

aggregate size in an effort to capture natural aggregation rates (results from other cell populations

are reported in Table S1, Figure S1).

(c) Statistical Analyses

We performed all statistical analyses using R version 3.4.2 (R Core Team 2016) and visually

inspected diagnostic plots (qqplots and plots of the distribution of the residuals against fitted

values) to validate model normality. Only the best fitting models are reported here. We created all

figures using the 'ggplot2' package with R (Wickham 2016). One *P. californicus* male was

excluded from the aggregate analysis dataset because their measurements represented clear

outliers. All means are presented ± 1 standard error.

To compare species differences in the proportion of aggregated cells, we used the mean values

for each male and a generalized linear mixed model (GLMM) using the glmer function from the

"lme4" R package and a logit link function (Bates et al. 2015). The binomial response was the

number of sperm cells in aggregate, and the total number of sperm cells was the binomial

denominator. In the initial statistical model, we observed the residual deviance to be larger than

the residual degrees of freedom, which is an indication of overdispersion (Crawley 2013). We thus

used an observation-level random effect (OLRE) as a random factor in all subsequent analyses to

control for overdispersion (Harrison 2014). We considered family ID as a random factor in the

initial model and both random factors were then used in bivariate analyses for predictors of

interest that could potentially explain differences in the proportion of aggregated cells. These

predictors included male age, pairing status, the timing of video recordings relative to harvest of

the epididymal tissue, and the number of videos recorded. Only predictors that had a *p*-value at or

below 0.20 were considered for the final model. We further screened these predictors for

collinearity with other significant predictors using linear models and removed collinear predictors,

so that only the one with greater relative significance was included in the final GLMM. The

remaining model included pairing status and species as fixed factors. We calculated the variance

inflation factor (VIF) values and found evidence of collinearity due to two VIF values above the

recommended threshold of three (Zuur et al. 2010). Thus, we removed pairing status as a fixed

factor and family ID as a random factor, leaving only species as a fixed factor within our final

model. Post-hoc pairwise comparisons were performed using Tukey HSD adjustments for multiple

comparisons from the "LSmeans" R package (Lenth 2016). We additionally analyzed whether the

proportion of aggregated cells correlates with sperm swimming speed using the same methods

outlined above with VCL as the main explanatory variable within both inter- and intra-species

GLMMs. Our inter-species model included pairing status as a fixed factor and both family ID and

an OLRE as random factors, whereas the intra-species models reported here included only an

OLRE as a random factor.

To compare species differences in the number of aggregated cells, we used the mean values

for each male and initially used a linear mixed model (LMM) using the lmer function from the

"lme4" R package, but eventually reverted to using a linear model (LM) because the family ID

random factor did not significantly contribute to the residual variability in the response variable.

Predictors that were considered for the initial LMM included male age, pairing status, the timing

of video recordings relative to harvest of the epididymal tissue, total sperm cells, number of

recorded videos, an interaction between the latter two variables, and the ratio of total sperm cells

to the number of videos recorded. We considered predictors with $p$-values $< 0.20$ for the final

model, but first screened each for collinearity with other significant predictors using a linear

model. Whenever collinearity was present, only the predictor with the greater relative significance

was included in the LM. We dropped non-significant explanatory variables one at a time based on

model comparisons using an analyses of variance test to determine the minimal adequate model,

but were unable to meet the normality assumptions for this model. We also assessed species

differences in the proportion of aggregates that were motile or progressively motile from the total

population of aggregates using binomial GLMM. The binomial response was the number of motile

sperm aggregates, and the total number of sperm aggregates was the binomial denominator. Our

final model contained both an OLRE due to detected overdispersion and family ID as random

factors.

To determine if sperm aggregates have motility or force benefits over single cells, we initially

performed a principal component analysis (PCA) using three related swimming performance

measures (VSL, VCL, and VAP) to reduce dimensionality and obtain a composite measure for

motile solitary and aggregated sperm cells in both low- and high-viscosity media. Because we

found that species distributions overlapped for this composite measure for both single sperm and

aggregates in low- and high-viscosity media (Figures S2 and S3), we focused on individual

kinematic parameters (VSL, VCL, VAP, LIN) within each species separately. Using our dataset of

mean values per male, we used a paired student's t-test to compare each kinematic parameter

between solitary sperm cells and sperm aggregates within males for each separate species to

determine if sperm aggregates have motility benefit over single cells. To determine if aggregates

have a force benefit over single sperm, we conducted these same analyses in high-viscosity media.

We then combined these two datasets in low- and high- viscosity media and compared sperm

aggregate kinematics in both low- and high- viscosity media at the intra-male level using a paired

student's t-test within each species.

To assess the structure of *P. maniculatus* and *P. polionotus* aggregates, we used generalized

linear models (GLM) to compare the proportions of 'defective' and 'opposed' aggregates to

'aligned' aggregates. For the composition GLM, the binomial response was the number of

'defective' sperm aggregates, and the total number of sperm aggregates that were either 'defective'

or 'aligned' was the binomial denominator. We used a paired student's t-test to compare the VCL

of these aggregates within males. For the orientation GLM, the binomial response was the number

of 'opposed' sperm aggregates, and the total number of sperm aggregates that were either

'opposed' or 'aligned' was the binomial denominator. We used a paired student's t-test to compare

the VCL of these aggregates within males.

Finally, to account for variation in phylogenetic relationships among of the species used in this

study, we adopted a phylogenetic generalized least squares approach (Pagel 1999; Freckleton et al.

2002) using the "caper" (Orme et al. 2013) and "APE" (Paradis et al. 2004) packages in R and

using an ultra-metric phylogenetic tree of *Peromyscus* (provided by Dr. Roy Neal Platt II, Texas

Biomedical Research Institute), based on sequence variation in the mitochondrial gene,

cytochrome B. The species' relationships within this tree matched those from other previously

established phylogenies of *Peromyscus* (Bradley et al. 2007; Turner et al. 2010). We used this

phylogeny as a covariate in regression analyses to investigate the effect of relative testis weight on

sperm aggregation, including the proportion of aggregated cells and aggregate size, and the within-

species CV for each of these parameters. To control for differences in male body when examining

testis mass, we included body mass as a separate fixed factor within our analyses, a method better

suited to estimating relative testis weight is size rather than using the ratio of testis to body mass or

residuals (García-Berthou 2001; Lüpold et al. 2009).

**Results**

We investigated the frequency of sperm aggregation in each species and found significant

differences in the proportion of aggregated cells among species (binomial GLMM: $n = 134$, $P <$

$0.001$; Table 1, 2; Figure 1), with more variance across species than within species (variance

across species $= 0.10$; variance within each species $< 0.10$). Specifically, sperm from *P.*

*maniculatus* and *P. polionotus* similarly aggregate the most, whereas *P. leucopus* and *P.*

*gossypinus* sperm similarly aggregate the least among all the species. Pairwise comparisons

adjusted for multiple comparisons using LSmeans revealed that species within these pairs do not

significantly differ from one another, whereas all other pairwise species comparisons do (Table 1,

2). The coefficient of variation (CV) for the proportion of cells aggregated within each species are

as follows: 68.8% for *P. californicus*, 50.6% for *P. eremicus*, 14.4% for *P. polionotus*, 10.8% for

*P. maniculatus*, 125.7% for *P. leucopus*, and 60.5% for *P. gossypinus*. We found the sperm of

males with greater VCL were less likely to form aggregates (binomial GLMM: $n = 130$, $p =$

$0.1435$), specifically within *P. gossypinus* (binomial GLMM: $n = 21$, $p = 0.0292$), *P. leucopus*

(binomial GLMM: $n = 22$, $p = 0.000193$), and *P. polionotus* (binomial GLMM: $n = 24$, $p = 1.46E-$

$05$). Controlling for phylogenic relationships and body mass, we found no effect of testis size ($F_{2,3}$

$= 2.606$, $P = 0.15816$) or the within-species CV ($F_{2,3} = 0.3603$, $P = 0.5604$) on the proportion of

aggregated cells.

Moreover, we found significant differences in the mean number of cells aggregated among

species (LM: $F_{6,126}$ = 56.37, $P < 0.001$; Table 1, 2; Figure 2A), with more variance observed across

species ($s^2$ = 1.96) than within species ($s^2 < 1.00$, except for *P. polionotus* [$s^2$ = 2.15]). Post-hoc

pairwise comparisons revealed that both *P. maniculatus* and *P. polionotus* produce the largest

sperm aggregates ($P < 0.05$ for all pairwise comparisons), whereas *P. gossypinus*, *P. leucopus*, and

*P. californicus* produce the smallest aggregates ($P < 0.05$ for pairwise comparisons), the latter of

which produces sperm cell aggregates that are statistically similar in size to those produced by *P.*

*eremicus* ($P = 0.9203$; Figure 2A). Controlling for phylogenic relationships and body mass, we

found a significant effect of testis weight on the within-species CV of aggregate size ($F_{2,3}$ = 8.398,

$P = 0.02655$, Figure 2B), but found less of an effect on the direct measure of aggregate size ($F_{2,3}$ =

3.058, $P = 0.1522$). The CV for the number of cells aggregated within each species were as

follows: 26.5% for *P. californicus*, 19.2% for *P. eremicus*, 27.5% for *P. polionotus*, 18.4% for *P.*

*maniculatus*, 13.3% for *P. leucopus*, and 7.2% for *P. gossypinus*.

By comparing sperm aggregate composition and orientation within the species that produce

the largest and most frequent sperm aggregates, we found that there are significantly more sperm

aggregates in *P. maniculatus* in which all sperm are aligned in a head-to-flagella orientation

(99.1%, 731/738) than in *P. polionotus* (87.5%, 720/826; GLM: $n = 42$, $P = 3.92e-12$) and that

these aggregates have significantly faster VCL compared to aggregates with unaligned cell

orientations in both species (Figure 3; paired t-tests: *P. maniculatus* $t = 5.9627$, $df = 4$, $P =$

0.003972; *P. polionotus* $t = 11.247$, $df = 11$, $P = 2.257e-07$). We also found that there are

significantly fewer sperm aggregates in *P. maniculatus* with immotile, stuck, or morphologically

abnormal cells (8.1%, 64/795) than in *P. polionotus* (11.5%, 94/814; binomial GLM: $n = 40$, $P =$

0.00503). Importantly, aggregates with these defects had significantly lower VCL compared to

aggregates without in both species (*P. maniculatus* $VCL_{aligned} = 179.60 \pm 6.88$ $\mu$m/sec, $VCL_{defective}$

$= 132.47 \pm 6.18$ $\mu$m/sec, paired t-test: $t = 20.627$, $df = 14$, $P = 7.075e-12$; *P. polionotus* $VCL_{aligned} =$

$135.48 \pm 3.51$ $\mu$m/sec, $VCL_{defective} = 115.62 \pm 5.22$ $\mu$m/sec, paired t-test: $t = 16.312$, $df = 20$, $P =$

5.079e-13).

When comparing the proportion of motile and progressively motile aggregates across species,

our pairwise comparisons revealed that *P. eremicus* produced a significantly smaller proportion of

motile aggregates than all other species ($P < 0.05$ for all *P. eremicus* pairwise comparisons; $P >$

0.05 for all other pairwise comparisons). Fitted values of the proportion of motile aggregates using

LSmeans were $0.77 \pm 0.04$ for *P. eremicus*, $0.92 \pm 0.02$ for *P. polionotus*, $0.91 \pm 0.02$ for *P.*

*gossypinus*, $0.95 \pm 0.01$ for *P. maniculatus*, $0.93 \pm 0.02$ for *P. leucopus*, and $0.91 \pm 0.02$ for *P.*

*californicus*. Moreover, post-hoc comparisons revealed that *P. eremicus, P. polionotus,* and *P.*

*gossypinus* all had the smallest proportions of progressively motile aggregates ($P < 0.05$ for all

pairwise comparisons; fitted values using LSmeans were $0.64 \pm 0.06$, $0.65 \pm 0.05$, and $0.75 \pm 0.05$,

respectively), the latter species of which did not significantly differ from *P. californicus* ($P =$

0.1981; $0.87 \pm 0.02$ for LSmeans fitted values). Conversely, *P. maniculatus* and *P. leucopus* had

the largest proportions of progressively motile aggregates ($P = 0.3417$; fitted values using

LSmeans were $0.94 \pm 0.02$ and $0.91 \pm 0.02$, respectively), the latter of which did not differ from *P.*

*californicus* ($P = 0.9109$).

Overall, we found species-specific differences in the effect of sperm aggregation on motility,

regardless of environmental complexity (Table 3, Figure 4). In low-viscosity medium, we found

that sperm aggregates have a significantly greater VCL in *P. maniculatus*, VSL in *P. maniculatus*,

*P. leucopus*, and *P. californicus*, LIN in *P. leucopus*, and VAP in *P. californicus* compared to

single cells. Conversely, sperm aggregates had a significantly lower VCL, VSL, and VAP velocity

in *P. polionotus* and *P. gossypinus* than single sperm in the low-viscosity medium (Figure 4). In

the high-viscosity medium, we found that sperm aggregates have a significantly greater VCL in *P.*

*maniculatus* and *P. californicus* and a higher VSL and VAP in *P. californicus* (Figure 4) compared

to single sperm. Conversely, sperm aggregates in the high viscosity medium had a significantly

lower LIN in *P. californicus*, *P. eremicus*, *P. polionotus*, and *P. maniculatus* as well as a reduced

VSL and VAP in *P. polionotus* than single cells (Figure 4).

**Discussion**

While it is known that collective sperm behaviors have evolved independently in a number of

taxa (Higginson and Pitnick 2011), it remains unclear how sperm aggregation evolves among

closely related species. Our comparative study reveals that sperm aggregating behaviors vary

across mice in the genus *Peromyscus*. We observed an effect of phylogenetic history on the

frequency of sperm aggregation, indicating that collective sperm behavior likely evolved prior to

the divergence of present-day species. Additionally, we find a negative association between

relative testis weight, a robust proxy for intensity of sperm competition in rodents (Ramm et al.

2005), and the coefficient of variation for the number of cells in aggregate, suggesting that sexual

selection has a stabilizing effect on sperm aggregate size. We then compared the motility of single

sperm and sperm aggregates across all species under low- and high-viscosity conditions, and show

that aggregation is kinematically beneficial for some species yet costly for others, regardless of

environmental complexity. When examining sperm from the species that aggregate the most

profoundly, we find that the formation of the aggregates and the orientation of the cells within the

group are critical to kinematics. Moreover, we observe more aggregates with efficient geometry in

the species that has evolved under strong post-copulatory sexual selection compared to its

monogamous sister-species. These findings support the prediction that sperm aggregation evolved

in *Peromyscus* to improve motility in a competitive context but reveal that relaxed selection may

have enabled less efficient strategies to persist, thereby generating diversity in collective sperm

behaviors within these closely-related species.

Our results reveal distinct species-specific differences across *Peromyscus* mice in the

frequency of sperm aggregation and the average size of these cellular groups. Multiple

*Peromyscus* species produce sperm that aggregate more extensively than other studied muroid

rodents (Tourmente et al. 2016), with the notable exception of the wood mouse (*Apodemus*

*sylvaticus*; Moore et al. 2002). In general, the proportion of sperm cells that aggregate is most

similar within each sister-species pair. Specifically, *P. maniculatus* and *P. polionotus* produce

sperm that aggregate the most (>80% of sperm), but the pair they are most closely-related to, *P.*

*gossypinus* and *P. leucopus*, are the species least likely to produce sperm that aggregate (<10% of

sperm); the most distantly-related species pair we assessed, *P. californicus* and *P. eremicus*, both

produce sperm with a moderate propensity to aggregate (~25% of sperm). These findings support

one of two possibilities for the evolution of sperm aggregates within these species: (a) a genus-

wide ancestral trait of moderate sperm aggregation with subsequent diversification leading to an

increase in *P. maniculatus* and *P. polionotus* and a decrease in *P. gossypinus* and *P. leucopus*, or

(b) the independent evolution of aggregation in the ancestors of the *P. californicus*-species pair

and the *P. maniculatus*-species pair. Such complex evolutionary histories with losses and

recurrences of sperm conjugation, and subsequent species divergence, have also been

demonstrated in diving beetles (Dytiscidae; Higginson et al. 2012a), consistent with the

evolutionary lability that we observe. Our experimental results may explain selection against

sperm aggregation if forming groups reduces sperm swimming performance, which we find in at

least one species, *P. gossypinus*. However, we found that sperm produced by their sister-species,

*P. leucopus*, also rarely aggregate, despite our observation that these rare collective groups have a

greater average velocity and are more linear than single cells. Interestingly, we observed that these

two promiscuous species that rarely aggregate, *P. gossypinus* and *P. leucopus*, have the largest

relative testes of the species studied, suggesting that they may have evolved increased sperm

production to improve competitive fertilization success (sensu Parker 1982), rather than

adaptations that influence motility (Snook 2005). Moreover, within these species (and *P.*

*polionotus*) we found a negative association between sperm speed and aggregation, even after

controlling for sperm density, indicating that differences in aggregate formation are not simply a

by-product of increased encounter rates. Together these results suggest different strategies

employed by divergent species across the *Peromyscus* genus in response to sperm competition.

In contrast to our results on the frequency of sperm aggregation within species, we found that

the average size of sperm groups does not align as closely with phylogenetic relationships. In

addition, we found that species with relatively larger testes, which is positively associated with

increased sperm competition (Ramm et al. 2005), exhibit less variation (CV) in aggregate size.

This result supports the prediction that relaxed sperm competition allows for greater intermale

variation to persist in a population (Calhim et al. 2007) and suggests that this post-copulatory

sexual selection may be stabilizing sperm aggregate size for a species-specific 'optimum' (Fisher

et al. 2014). Similarly, other studies have shown that the strength of sexual selection regulates

variance in sperm morphology across taxa and at multiple levels of organization, including within-

and between-males (Immler et al. 2008; Fitzpatrick and Baer 2011; Carballo et al. 2019) and

within- and between-species (Calhim et al. 2007; Rowley et al. 2019). A study on sperm bundles

across ten *Carabus* ground beetles also found intense selection on bundle size, which are

dimorphic and either small or large; the large, but not small, sperm bundles are positively

correlated with measures of sperm competition risk, including copulatory piece length and mate

guarding, suggesting that diversity of large sperm bundles is associated with sexual selection

(Takami and Sota 2007). In contrast to these findings that competition drives sperm-sperm

interactions, a study on the evolution of such sperm traits in diving beetles found that variation in

sperm conjugation is more associated with female reproductive tract architecture (Higginson et al.

2012b,a). Therefore, while our results suggest that stabilizing selection on sperm aggregate size is

associated with an increase in sperm competition given the correlation with relative testis weight,

mechanisms of female control (Eberhard 1996) may also play an important evolutionary role.

We compared the motility of single and aggregated sperm sampled from the same male to test

whether sperm aggregates swim faster or more efficiently than single sperm, which is predicted if

the combined force of multiple flagella enhances their motility (Moore et al. 2002). We found

improved kinematic measures in half of the species studied (*P. maniculatus, P. californicus*, and

*P. leucopus*), thus supporting this functional hypothesis; however, in several species we found

aggregation had some negative (*P. polionotus* and *P. gossypinus*) or no (*P. eremicus*) impact on

motility relative to single sperm cells. While we found support for a theoretical prediction that

sperm aggregates achieve greater straight-line velocity because they move in a more linear path of

travel rather than at a faster speed (i.e., curvilinear velocity; Fisher et al. 2014) in two of our six

focal species (*P. californicus* and *P. leucopus*), we did not find this kinematic benefit for sperm

aggregation in all *Peromyscus* species. These results corroborate other studies in more disparate

taxonomic groups that have quantified sperm aggregation motility and found inconsistent results.

For example, sperm trains exhibit greater swimming progressive motility in the wood mouse

(Moore et al. 2002), and greater velocity than individual sperm in the Norway rat, but not the

house mouse (Immler et al. 2007). In invertebrates, the swimming velocity of fishfly sperm

increases with number of sperm in a bundle (Hayashi 1998), but in a marine snail, there is no

differences in swimming speed between paired and single sperm (Ishijima et al. 1999). One

possible explanation for these differences across taxa is that cell orientation within an aggregate is

critical for its collective motility. Sperm cells are predicted to be faster if they generate increased

force with proportionally less drag (see Higginson and Pitnick 2011 and references therein); such

effects may be true for sperm aggregations as well in which cells conjoin head-to-tail, thereby

increasing the length of the collective unit, or in which flagella within the group beat

synchronously (Higginson and Pitnick 2011). Our results support that even sperm aggregates that

conjoin head-to-head and are thus wider, as they are in *Peromyscus,* can offer a motility

advantage.

Our results suggest that relaxed sexual selection may allow the persistence of less optimal

strategies based on the quantitative and qualitative differences we observed among sperm

aggregates of different species. The most interesting motility results are those of the sister-species

pair with divergent mating systems that both form the most frequent and largest sperm aggregates.

In *P. maniculatus*, a promiscuous species, sperm aggregates exhibit greater straight-line and

curvilinear velocity compared to single cells, but the opposite was true for its monogamous

congener, *P. polionotus*. We find that these kinematic difference are associated with differences in

aggregate geometry; when sperm heads and flagella are not oriented in the same direction, the

cells within an aggregate exert opposing forces on one another, thereby reducing the overall

motility of the group (Fisher et al. 2014; Pearce et al. 2018). Indeed, we found that sperm from the

monogamous *P. polionotus* males are less likely to form aggregates with all sperm aligned and

more likely to include immotile or morphologically abnormal sperm, consequently resulting in

slower aggregates than those of *P. maniculatus* (Pearce et al. 2018). This finding is consistent with

previous reports that *P. polionotus* sperm tend to form optimal-sized aggregates less often than in

*P. maniculatus* (Fisher et al. 2014). Together, these observations further support the hypothesis

that sperm aggregation evolved prior to the divergence of the species pair (Fisher and Hoekstra

2010), and when monogamy evolved secondarily in *P. polionotus* (Greenbaum et al. 1978; Turner

et al. 2010), relaxed sexual selection allowed for the persistence of less motile sperm traits. In line

with this prediction, we observed the smallest proportion of motile and progressively motile sperm

aggregates in another monogamous species, *P. eremicus,* but the largest proportion of

progressively motile aggregates in two promiscuous species, *P. maniculatus* and *P. leucopus*.

Similar results have been reported in house mice, in which males evolving under intense sperm

competition produced a greater proportion of motile sperm, compared to males from lineages

subject to relaxed selection (Firman and Simmons 2011). Together these findings support that the

motility benefits conferred by sperm aggregation are associated with variation in post-copulatory

sexual selection.

When we compared single and aggregated sperm in a viscous environment to test if

aggregation improves motility in more complex fluids, similar to the secretions or cervical mucus

(Hanson and Overstreet 1981) of the female reproductive tract (Suarez 2016; Simons and Olson

2018), we found that collective motion is beneficial for some species but costly for others.

Compared to single sperm, aggregates swim with greater curvilinear velocity in *P. maniculatus*

and *P. californicus*, and greater straight-line and average path velocities in *P. californicus,* but

aggregates were less linear than single cells in all species except for *P. leucopus* and *P.*

*gossypinus*. While other studies have found kinematic benefits for sperm conjugates compared to

single cells in higher viscosities in the gray short-tailed opossum (Moore and Taggart 1995), the

wood mouse (Moore et al. 2002), and the fishfly (Hayashi 1998), a study in bulls found that sperm

were slower, exhibited less organized swimming patterns, and were less likely to cluster in viscous

fluids (Tung et al. 2017), both of which are consistent with our analysis of *Peromyscus* aggregates.

Ultimately, the benefit of sperm aggregation depends on the relative importance of each kinematic

parameter during sperm migration *in vivo*. Although beyond the scope of this study, we predict

that improved linearity afforded by collective motion may help to direct the sperm through the

dynamic fluids of the female reproductive tract (Ishikawa et al. 2016) and that increased velocity

will reduce the time it takes for the sperm to arrive at the fertilization site.

In conclusion, our study highlights the diversity of sperm aggregation within a single

taxonomic lineage and how selection has shaped the formation and performance of these cellular

groups. We show that both evolutionary history and varying levels of post-copulatory sexual

selection influence the frequency and size of sperm groups. Moreover, we find that sperm

aggregation can improve sperm motility in both simple and complex fluids, but this is not

consistent across all species. Theoretical predictions (Fisher et al. 2014; Pearce et al. 2018) and

emerging empirical evidence suggests that motility benefits may only be realized if cells maintain

optimal alignment within the groups and, if achieved, may provide these sperm with a competitive

advantage in the female reproductive tract (Higginson et al. 2012b). Future work investigating

sperm aggregates *in vivo* (e.g., Ishikawa et al. 2016; Wang and Larina 2018) will shed light on the

co-evolution of these unique gametic behaviors and the enormously variable and dynamic female

reproductive tracts through which sperm must navigate.

**Acknowledgments:**

We are grateful to Erica Glasper for providing *P. californicus* males and Hopi Hoekstra for

providing *P. gossypinus* males. Thanks to Harrison Arsis, Madeline-Sophie Dang, Catherine Liu,

and Audrey Mvemba for their assistance with video analysis and to Philip Johnson, Danielle

Adams, Sam Church, and Shelby Wilson for statistical advice. We thank Neal Platt for providing

the ultra-metric *Peromyscus* tree for use in our statistical analysis.

**References:**

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects models using

lme4. Journal of Statistical Software 67:1–48.

Bedford, J. M., J. C. Rodger, and W. G. Breed. 1984. Why so many mammalian spermatozoa - a

clue from marsupials? Proceedings of the Royal Society of London B: Biological Sciences

221:221–233.

Bedford, N. L., and H. E. Hoekstra. 2015. The natural history of model organisms: Peromyscus

mice as a model for studying natural variation. eLife Sciences 4:e06813.

Birdsall, D. A., and D. Nash. 1973. Occurrence of successful multiple insemination of females in

natural populations of deer mice (*Peromyscus maniculatus*). Evolution 27:106–110.

Birkhead, T. R., J. G. Martinez, T. Burke, and D. P. Froman. 1999. Sperm mobility determines the

outcome of sperm competition in the domestic fowl. Proceedings of the Royal Society of

London. Series B: Biological Sciences 266:1759–1764.

Birkhead, T. R., A. P. Møller, and W. J. Sutherland. 1993. Why do females make it so difficult for

males to fertilize their eggs? Journal of Theoretical Biology 161:51–60.

Bradley, R. D., N. D. Durish, D. S. Rogers, J. R. Miller, M. D. Engstrom, and C. W. Kilpatrick.

2007. Toward a molecular phylogeny for *Peromyscus*: evidence from mitochondrial

cytochrome-*b* sequences. Journal of Mammalogy 88:1146–1159.

Calhim, S., S. Immler, and T. R. Birkhead. 2007. Postcopulatory sexual selection is associated

with reduced variation in sperm morphology. PLoS ONE 2:e413.

Carballo, L., A. Battistotti, K. Teltscher, M. Lierz, A. Bublat, M. Valcu, and B. Kempenaers.

2019. Sperm morphology and evidence for sperm competition among parrots. Journal of

Evolutionary Biology 32:856–867.

Crawley, M. J. 2013. The R Book Second Edition. John Wiley & Sons, Ltd., England.

Dewsbury, D. A. 1981. An exercise in the prediction of monogamy in the field from laboratory

data on 42 species of muroid rodents. Biologist 63:138–162.

Dewsbury, D. A. 1985. Interactions between males and their sperm during multi-male copulatory

episodes of deer mice (*Peromyscus maniculatus*). Animal Behaviour 33:1266–1274.

Eberhard, W. 1996. Female Control: Sexual Selection by Cryptic Female Choice. Princeton

University Press.

Firman, R. C., and L. W. Simmons. 2011. Experimental evolution of sperm competitiveness in a

mammal. BMC Evolutionary Biology 11:19.

Fisher, H. S., L. Giomi, H. E. Hoekstra, and L. Mahadevan. 2014. The dynamics of sperm

cooperation in a competitive environment. Proceedings of the Royal Society of London B:

Biological Sciences 281:20140296.

Fisher, H. S., and H. E. Hoekstra. 2010. Competition drives cooperation among closely related

sperm of deer mice. Nature 463:801.

Fitzpatrick, J. L., and B. Baer. 2011. Polyandry reduces sperm length variation in social insects.

Evolution 65:3006–3012.

Foltz, D. W. 1981. Genetic evidence for long-term monogamy in a small rodent, *Peromyscus*

*polionotus*. The American Naturalist 117:665–675.

Freckleton, R. P., P. H. Harvey, and M. Pagel. 2002. Phylogenetic analysis and comparative data:

a test and review of evidence. The American Naturalist 160:712–726.

Gage, M. J. G., C. P. Macfarlane, S. Yeates, R. G. Ward, J. B. Searle, and G. A. Parker. 2004.

Spermatozoal traits and sperm competition in Atlantic salmon: relative sperm velocity is

the primary determinant of fertilization success. Current Biology 14:44–47.

García-Berthou, E. 2001. On the misuse of residuals in ecology: testing regression residuals vs. the

analysis of covariance. Journal of Animal Ecology 70:708–711.

Greenbaum, I. F., R. J. Baker, and P. R. Ramsey. 1978. Chromosomal evolution and the mode of

speciation in three species of *Peromyscus*. Evolution 32:646–654.

Hanson, F. W., and J. W. Overstreet. 1981. The interaction of human spermatozoa with cervical

mucus *in vivo*. American Journal of Obstetrics and Gynecology 140:173–178.

Harrison, X. A. 2014. Using observation-level random effects to model overdispersion in count

data in ecology and evolution. PeerJ 2:e616.

Hayashi, F. 1998. Sperm co-operation in the Fishfly, *Parachauliodes japonicus*. Functional

Ecology 12:347–350.

Higginson, D. M., and K. R. H. Henn. 2012. Effects of sperm conjugation and dissociation on

sperm viability *in vitro*. PloS one 7:e34190.

Higginson, D. M., K. B. Miller, K. A. Segraves, and S. Pitnick. 2012a. Convergence, recurrence

and diversification of complex sperm traits in diving beetles (Dytiscidae). Evolution

66:1650–1661.

Higginson, D. M., K. B. Miller, K. A. Segraves, and S. Pitnick. 2012b. Female reproductive tract

form drives the evolution of complex sperm morphology. Proceedings of the National

Academy of Sciences 109:4538–4543.

Higginson, D. M., and S. Pitnick. 2011. Evolution of intra-ejaculate sperm interactions: do sperm

cooperate? Biological Reviews 86:249–270.

Immler, S. 2008. Sperm competition and sperm cooperation: the potential role of diploid and

haploid expression. Reproduction 135:275–283.

Immler, S., S. Calhim, and T. R. Birkhead. 2008. Increased postcopulatory sexual selection

reduces the intramale variation in sperm design. Evolution 62:1538–1543.

Immler, S., H. D. Moore, W. Breed, and T. Birkhead. 2007. By hook or by crook? Morphometry,

competition and cooperation in rodent sperm. PloS one 2:e170.

Ishijima, S., S. A. Ishijima, and B. A. Afzelius. 1999. Movement of Turritella spermatozoa:

Direction of propagation and chirality of flagellar bends. Cell Motility 44:85–95.

Ishikawa, Y., T. Usui, M. Yamashita, Y. Kanemori, and T. Baba. 2016. Surfing and swimming of

ejaculated sperm in the mouse oviduct. Biology of Reproduction 94:89, 1–9.

Lenth, R. V. 2016. Least-squares means: The R package 'lsmeans'. Journal of Statistical Software

69:1–33.

Lüpold, S., G. M. Linz, J. W. Rivers, D. F. Westneat, and T. R. Birkhead. 2009. Sperm

competition selects beyond relative testes size in birds. Evolution 63:391–402.

Mackie, J. B., and M. H. Walker. 1974. A study of the conjugate sperm of the dytiscid water

beetles *Dytiscus marginalis* and *Colymbetes fuscus*. Cell and Tissue Research 148:505–

519.

Magdanz, V., S. Boryshpolets, C. Ridzewski, B. Eckel, and K. Reinhardt. 2019. The swim-up

technique separates bovine sperm by metabolic rates, motility and tail length. bioRxiv

624502.

McGlinn, S. M., B. A. Shepherd, and J. Martan. 1979. Rouleau Formation and fertility of

spermatozoa in guinea pigs treated with cyproterone acetate. Archives of Andrology

2:197–202.

Moore, H. D., K. Dvoráková, N. Jenkins, and W. Breed. 2002. Exceptional sperm cooperation in

the wood mouse. Nature 418:174–177.

Moore, H. D. M., and D. A. Taggart. 1995. Sperm pairing in the opossum increases the efficiency

of sperm movement in a viscous environment. Biology of Reproduction 52:947–953.

Nixon, B., H. Ecroyd, J.-L. Dacheux, F. Dacheux, V. Labas, S. D. Johnston, and R. C. Jones.

2016. Formation and dissociation of sperm bundles in monotremes. Biol Reprod 95:1–11.

Orme, D., R. Freckleton, G. Thomas, and T. Petzoldt. 2013. The caper package: comparative

analysis of phylogenetics and evolution in R. R package version 5:1–36.

Pagel, M. 1999. Inferring the historical patterns of biological evolution. Nature 401:877–884.

Paradis, E., J. Claude, and K. Strimmer. 2004. APE: Analyses of phylogenetics and evolution in R

language. Bioinformatics 20:289–290.

Parker, G. A. 1982. Why are there so many tiny sperm? Sperm competition and the maintenance

of two sexes. Journal of Theoretical Biology 96:281–294.

Pearce, D. J. G., L. A. Hoogerbrugge, K. A. Hook, H. S. Fisher, and L. Giomi. 2018. Cellular

geometry controls the efficiency of motile sperm aggregates. Journal of The Royal Society

Interface 15:20180702.

Pearcy, M., N. Delescaille, P. Lybaert, and S. Aron. 2014. Team swimming in ant spermatozoa.

Biology Letters 10:20140308.

Phillips, D. M. 1970. Ultrastructure of spermatozoa of the wooly opossum *Caluromys philander*.

Journal of Ultrastructure Research 33:381–397.

Pitnick, S., D. J. Hosken, and T. R. Birkhead. 2009. Sperm Morphological Diversity. Pp. 69–149

*in* T. R. Birkhead, D. J. Hosken, and S. Pitnick, eds. Sperm Biology: An Evolutionary

Perspective. Academic Press, London.

Pizzari, T., and K. R. Foster. 2008. Sperm sociality: cooperation, altruism, and spite. PLOS

Biology 6.

R Core Team. 2016. R: A Language and Environment for Statistical Computing. R Foundation for

Statistical Computing, Vienna, Austria.

Ramm, S. A., G. A. Parker, and P. Stockley. 2005. Sperm competition and the evolution of male

reproductive anatomy in rodents. Proceedings of the Royal Society B: Biological Sciences

272:949–955.

Rowley, A., L. Locatello, A. Kahrl, M. Rego, A. Boussard, E. Garza-Gisholt, R. M. Kempster, S.

P. Collin, E. Giacomello, M. C. Follesa, C. Porcu, J. P. Evans, F. Hazin, F. Garcia-

Gonzalez, T. Daly-Engel, C. Mazzoldi, and J. L. Fitzpatrick. 2019. Sexual selection and

the evolution of sperm morphology in sharks. Journal of Evolutionary Biology 32:1027–

1035.

Simons, J. E., and S. D. Olson. 2018. Sperm motility: models for dynamic behavior in complex

environments. Pp. 169–209 *in* M. Stolarska and N. Tarfulea, eds. Cell Movement.

Springer International Publishing, Cham.

Snook, R. R. 2005. Sperm in competition: not playing by the numbers. Trends in Ecology &

Evolution 20:46–53.

Suarez, S. S. 2016. Mammalian sperm interactions with the female reproductive tract. Cell and

Tissue Research 363:185–194.

Suarez, S. S., and X. Dai. 1992. Hyperactivation enhances mouse sperm capacity for penetrating

viscoelastic media. Biology of Reproduction 46:686–691.

Taggart, D. A., J. L. Johnson, H. P. O'Brien, and H. D. M. Moore. 1993. Why do spermatozoa of

the American marsupial form pairs? A clue from the analysis of sperm-pairing in the

epididymis of the grey short-tailed opossum, *Monodelphis domestica*. Anatomical Record

236:465–478.

Takami, Y., and T. Sota. 2007. Sperm competition promotes diversity of sperm bundles in

*Ohomopterus* ground beetles. Naturwissenschaften 94:543–550.

Tourmente, M., D. Zarka-Trigo, and E. R. S. Roldan. 2016. Is the hook of muroid rodent's sperm

related to sperm train formation? Journal of Evolutionary Biology 29:1168–1177.

Tung, C., C. Lin, B. Harvey, A. G. Fiore, F. Ardon, M. Wu, and S. S. Suarez. 2017. Fluid

viscoelasticity promotes collective swimming of sperm. Scientific Reports 7:1–9.

Tung, K. S. K., A. Okada, and R. Anagimachi. 1980. Sperm autoantigens and fertilization. I.

effects of antisperm autoantibodies on rouleaux formation, viability, and acrosome

reaction of guinea pig spermatozoa. Biology of Reproduction 23:877–886.

Turner, L. M., A. R. Young, H. Römpler, T. Schöneberg, S. M. Phelps, and H. E. Hoekstra. 2010.

Monogamy evolves through multiple mechanisms: evidence from V1aR in deer mice.

Molecular Biology and Evolution 27:1269–1278.

Wang, S., and I. V. Larina. 2018. *In vivo* three-dimensional tracking of sperm behaviors in the

mouse oviduct. Development 145:dev157685.

Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer.

Woolley, D. M., R. F. Crockett, W. D. I. Groom, and S. G. Revell. 2009. A study of

synchronisation between the flagella of bull spermatozoa, with related observations.

Journal of Experimental Biology 212:2215–2223.

World Health Organization. 2010. WHO laboratory manual for the examination and processing of

human semen.

Zuur, A. F., E. N. Ieno, and C. S. Elphick. 2010. A protocol for data exploration to avoid common

statistical problems. Methods in Ecology and Evolution 1:3–14.

# FIGURES AND TABLES

**Figure 1.**

The observed natural variation in the proportion of aggregated sperm cells for six closely related species of *Peromyscus* mice (phylogeny adapted from Bradley et al. 2007). Box-plots represent median and interquartile ranges with raw data overlaid. Statistically significant differences at the $P = 0.05$ level are denoted by differing letters; shared letters denote no statistical difference.

**Figure 2.**

Sperm aggregate size varies among species and is negatively associated with relative testes size. (A) The observed natural variation in the number of aggregated sperm cells for six closely related species of *Peromyscus* mice. Box-plots represent median and interquartile ranges with raw data overlaid. Statistically significant differences at the $P = 0.05$ level are denoted by differing letters; shared letters denote no statistical difference. (B) When controlling for phylogenetic relationships, the coefficient of variation (CV) for the number of aggregated sperm cells negatively correlates with relative testis mass across these species. Note truncated y-axes.

**Figure 3.**

The effect of sperm aggregate geometry on the curvilinear velocity (*μ*m/sec) of sperm aggregates for two species that aggregated most – *Peromyscus maniculatus* and *Peromyscus polionotus*. Circles represent mean values per male within each species, and black lines represent the mean value within each category. Note truncated y-axis.

**Figure 4.**

Kinematic parameters of sperm aggregates (closed circles) and solitary sperm cells (open circles) for six species of *Peromyscus* mice in low- and high-viscosity conditions. Circles represent mean values per male, and black lines represent species means. Statistical significance levels comparing aggregated and solitary cells within each species are indicated by * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. Note truncated y-axes.

**TABLE 1.**

Summary of live sperm aggregate results for mice in the genus *Peromyscus*

| *Peromyscus* Species | No. males | No. sperm | % motile sperm | % progressive sperm | No. sperm aggregates | % cells in aggregate | % motile aggregates | % progressive aggregates |
|---|---|---|---|---|---|---|---|---|
| *californicus* | 28 | 8235 | 81.2% | 78.9% | 823 | 29.7% (2446/8235) | 90.5% (745/823) | 86.6% (713/823) |
| *eremicus* | 21 | 4906 | 67.4% | 58.2% | 513 | 30.2% (1481/4906) | 77.8% (399/513) | 66.7% (342/513) |
| *polionotus* | 24 | 6360 | 82.4% | 57.3% | 949 | 79.5% (5059/6360) | 90.9% (863/949) | 66.1% (627/949) |
| *maniculatus* | 18 | 4991 | 85.0% | 84.3% | 822 | 81.3% (4059/4991) | 93.4% (768/822) | 92.3% (759/822) |
| *leucopus* | 22 | 6341 | 87.6% | 84.5% | 361 | 15.0% (949/6341) | 90.3% (326/361) | 88.1% (318/361) |
| *gossypinus* | 21 | 5970 | 82.2% | 70.3% | 254 | 9.6% (576/5970) | 88.6% (225/254) | 70.1% (178/254) |

Motile sperm cells include only those that exhibited movement patterns. Progressive sperm cells are motile cells with a straight-line velocity $\geq 25$ μm/sec.

**TABLE 2.**

Fixed effects from a binomial generalized linear mixed model examining differences in the proportion of aggregated sperm cells and a linear model examining differences in the number of aggregated sperm cells across six species of *Peromyscus* mice

| GLMM: PROPORTION OF CELLS IN AGGREGATE | | | | | |
|---|---|---|---|---|---|
| Model Term | Beta (SE) | Exp (beta) | 95% CI | z | Pr(>\|z\|) |
| Intercept | 1.52 (0.22) | | | | |
| *eremicus* | -2.47 (0.30) | 0.07 | (0.04, 0.13) | -8.14 | **< 0.001** |
| *gossypinus* | -3.94 (0.31) | 0.02 | (0.01, 0.03) | -12.89 | **< 0.001** |
| *californicus* | -2.58 (0.29) | 0.07 | (0.04, 0.12) | -9.06 | **< 0.001** |
| *leucopus* | -3.79 (0.30) | 0.02 | (0.01, 0.04) | -12.46 | **< 0.001** |
| *polionotus* | -0.25 (0.29) | 0.44 | (0.30, 0.58) | -0.85 | 0.393 |

| LM: NUMBER OF CELLS IN AGGREGATE | | | | | |
|---|---|---|---|---|---|
| Model Term | Beta (SE) | Exp (beta) | 95% CI | t | Pr(>\|z\|) |
| Intercept | 3.90 (0.26) | | | | |
| Total Sperm Cells | 0.00 (0.00) | 0.50 | (0.50, 0.50) | 5.83 | **< 0.001** |
| *eremicus* | -1.97 (0.24) | 0.12 | (0.08, 0.18) | -8.15 | **< 0.001** |
| *gossypinus* | -2.77 (0.24) | 0.06 | (0.04, 0.09) | -11.53 | **< 0.001** |
| *californicus* | -2.19 (0.23) | 0.10 | (0.07, 0.15) | -9.68 | **< 0.001** |
| *leucopus* | -2.68 (0.24) | 0.06 | (0.04, 0.10) | -11.18 | **< 0.001** |
| *polionotus* | 0.04 (0.23) | 0.51 | (0.40, 0.62) | 0.16 | 0.877 |

For both models, all rows are being compared with the intercept – *Peromyscus maniculatus*. 95% confidence intervals (CI) were calculated for each effect size.

**TABLE 3.**

Results from an intra-male analysis comparing motile solitary and aggregated sperm kinematics in low- and high-viscosity conditions for six species of *Peromyscus* mice to test whether sperm aggregates confer kinematic benefits (shaded in gray)

| *PEROMYSCUS* SPECIES | df | KINEMATIC VARIABLE | | | |
|---|---|---|---|---|---|
| | | Curvilinear Velocity (µm/sec) | Linearity (VSL/VCL) | Straight-Line Velocity (µm/sec) | Average Path Velocity (µm/sec) |
| LOW-VISCOSITY CONDITIONS | | | | | |
| *californicus* | 28 | $t = -1.0545$, $p = 0.3007$ | $t = -0.56153$, $p = 0.5789$ | $t = -2.2982$, **$p = 0.02923$** | $t = -2.8162$, **$p = 0.008805$** |
| *eremicus* | 20 | $t = 1.6225$, $p = 0.1204$ | $t = 0.73517$, $p = 0.4708$ | $t = 0.1312$, $p = 0.8969$ | $t = 1.0464$, $p = 0.3079$ |
| *polionotus* | 23 | $t = 9.4575$, **$p = 2.1566e\text{-}09$** | $t = 1.699$, $p = 0.1028$ | $t = 5.8355$, **$p = 6.026e\text{-}06$** | $t = 6.0729$, **$p = 3.408e\text{-}06$** |
| *maniculatus* | 17 | $t = 2.2482$, **$p = 0.03812$** | $t = 0.48075$, $p = 0.6368$ | $t = -2.2335$, **$p = 0.03924$** | $t = -1.9206$, $p = 0.07172$ |
| *leucopus* | 20 | $t = 0.23337$, $p = 0.8178$ | $t = -4.8385$, **$p = 9.973e\text{-}05$** | $t = -5.5521$, **$p = 1.959e\text{-}05$** | $t = -1.838$, $p = 0.08096$ |
| *gossypinus* | 20 | $t = 5.4048$, **$p = 2.73e\text{-}05$** | $t = -1.0247$, $p = 0.3177$ | $t = 3.715$, **$p = 0.001369$** | $t = 4.479$, **$p = 0.0002298$** |
| HIGH-VISCOSITY CONDITIONS | | | | | |
| *californicus* | 9 | $t = -3.7465$, **$p = 0.003357$** | $t = 2.682$, **$p = 0.02512$** | $t = -2.7743$, **$p = 0.0216$** | $t = -4.1106$, **$p = 0.002634$** |
| *eremicus* | 9 | $t = -0.71739$, $p = 0.4913$ | $t = 4.2087$, **$p = 0.002277$** | $t = 0.62918$, $p = 0.5449$ | $t = 0.56505$, $p = 0.5858$ |
| *polionotus* | 13 | $t = 2.0349$, $p = 0.06278$ | $t = 3.9242$, **$p = 0.001745$** | $t = 4.8314$, **$p = 0.0003279$** | $t = 4.18$, **$p = 0.001079$** |
| *maniculatus* | 11 | $t = -2.9397$, **$p = 0.01345$** | $t = 3.9259$, **$p = 0.002369$** | $t = -0.30835$, $p = 0.7636$ | $t = -1.0543$, $p = 0.3114$ |
| *leucopus* | 9 | $t = 0.32429$, $p = 0.7531$ | $t = -0.31635$, $p = 0.759$ | $t = -0.34159$, $p = 0.7405$ | $t = 0.95055$, $p = 0.3667$ |
| *gossypinus* | 10 | $t = -0.63835$, $p = 0.95376$ | $t = -0.076451$, $p = 0.9406$ | $t = -0.60402$, $p = 0.5593$ | $t = -0.71643$, $p = 0.4901$ |

Statistical results are based on paired-student t-tests.