Collective sperm movements in *Peromyscus* mice are shaped by phylogenetic

history and post-copulatory sexual selection

Kristin A. Hook, W. David Weber, Heidi S. Fisher

Department of Biology, University of Maryland, College Park, MD 20742, U.S.A.

Abstract

In some species, sperm form motile, coordinated groups as they migrate through the female

reproductive tract. Collective sperm migration is posited to have evolved to improve sperm

swimming performance, and thus may be beneficial in a competitive context, but limited evidence

supports this theory. Here we investigate sperm aggregation across closely-related species of

Peromyscus mice that naturally vary by mating system. We find that phylogenetic history predicts

the likelihood that sperm will aggregate but that variation in aggregate size negatively associates

with relative testis size, suggesting that sperm competition has a stabilizing effect on this trait.

Moreover, we show that sperm aggregation is not kinematically beneficial for all species, and we

hypothesize that swimming performance is dependent on the orientation and composition of sperm

groups. To test this, we compared sperm from the two sister-species that aggregate most frequently

and find that sperm of the species that evolved under intense sperm competition forms aggregates

with efficient geometry more frequently than sperm from its monogamous congener. Together, our

results are consistent with the hypothesis that sperm aggregation evolved to improve motility in a

competitive context; however, when monogamy evolved secondarily, relaxed sexual selection

allowed for less efficient strategies to persist.

Introduction

Sperm cells are one of the most diverse cell types in nature and exhibit striking variation both within and across taxa [1]. In addition to being morphologically diverse, sperm may exhibit complex, emergent behaviors, including sperm conjugation, which occurs when two or more cells join together for motility or transport through the female reproductive tract before dissociating prior to fertilization [1,2]. Although relatively rare, these sperm-sperm interactions have evolved multiple times across independent lineages of internally fertilizing species [1–3], yet the adaptive significance of these gametic interactions remains unclear for many taxa.

One functional hypothesis posits that sperm aggregates may be advantageous if the combined force generated by multiple flagella enable sperm conjugates to swim faster than single cells [4,5]. Improved sperm motility can be beneficial if it enables cells to move quickly through hostile environments within the female tract [6,7], and in competitive environments in which females mate with multiple males, as sperm velocity is often correlated with competitive fertilization success [7–9]. While some studies that have quantified the motility of sperm aggregates have found empirical support for this hypothesis [4,10–13], others have not [14,15]. For example, sperm groups swim faster than solitary cells in the Norway rat (*Rattus norvegicus*), yet house mouse (*Mus musculus*) sperm swim slower as groups under identical experimental conditions [5]. Another non-mutually exclusive hypothesis is that sperm aggregation facilitates migration through viscous or viscoelastic secretions of the female reproductive tract [16,17], including cervical mucus [18], which has received some empirical support [4,10,16; but see 15]. In the grey short-tailed opossum (*Monodelphis domestica*), for example, sperm pairs swim with greater motility than single sperm in viscous fluids; however, these sperm pairs were artificially induced to uncouple [16] and therefore the comparison may not be biologically informative [1]. Additionally, sperm aggregation may protect sensitive regions of the sperm, such as the acrosome, from damage and preserve sperm functionality during passage through the male or female reproductive tracts [19–22], or enhance egg penetration during fertilization [23; but see 24]. While it is often assumed

that coordinated sperm movements are adaptive, inconsistent findings and a wide diversity of

naturally variable sperm behavior have limited our understanding of the functional advantages of

sperm aggregation.

Multiple independent origins of sperm conjugation suggest that the functional consequences,

as well as mechanisms that regulate these cellular interactions, are likely to vary throughout nature

[2]. Indeed, the formation of sperm aggregates and the number of grouped cells varies widely

across taxa [7]. In mammals alone, sperm of some species assemble during epididymal transport

and are molecularly "glued" to one another as bi-flagellate pairs in grey short-tailed opossums

[25], as bundles of roughly 100 cells in monotremes [26], or as organized rouleaux of five or more

cells in guinea pigs (*Cavia porcellus*; [24]). Conversely, mammalian sperm may also assemble

after ejaculation to form variably-sized groups. For instance, sperm may form temporary clusters

of up to sixteen cells in bulls (*Bos taurus*; [15]) or more fixed groups of up to 30 cells in house

mice [5], 50 cells in the Norway rat [5], or thousands of cells in the wood mouse (*Apodemus*

sylvaticus; [4]) whereby the hook-shaped heads interlock or attach to the flagella of other sperm.

For these latter species in which sperm groups form after ejaculation, both single and aggregated

sperm typically co-occur, thus allowing for direct comparisons between collective and solitary

sperm movements within an ejaculate while controlling for within-male variability.

Closely-related species in the rodent genus, *Peromyscus,* produce sperm that naturally vary in

their collective behavior. Sperm of some species assemble temporary groups after ejaculation by

adhering to one another at their head region [27] and disassemble prior to fertilization [12]. In *P.*

maniculatus, sperm selectively group with the most closely-related cells to form motile groups of

up to 30 cells [12], but there is a non-monotonic association between group size and swimming

velocity, indicating that some groups are faster than single cells but others are not [28].

Conversely, sperm produced by their sister-species, *P. polionotus*, also form aggregates but do so

indiscriminately with related and unrelated cells [12], are less likely to be optimally-sized, and

overall move in a less linear trajectory [28]. Intriguingly, sperm competition is predicted to be

more intense in *P. maniculatus* than in *P. polionotus* due to their different mating systems. In *P.*

maniculatus, both sexes mate with multiple partners, often in overlapping series just minutes apart

[29], and females frequently carry multiple-paternity litters in the wild [30], whereas both

behavioral [31] and genetic data [32] indicate that *P. polionotus* is strictly monogamous. Evidence

suggests that monogamy has evolved at least twice within the *Peromyscus* lineage [33,34], thus

enabling us to investigate if post-copulatory sexual selection, driven by female mating behavior,

has shaped the evolution of sperm aggregation [4] more broadly across the *Peromyscus* lineage. In

this study, we quantify intra- and inter-specific differences in the size and performance of sperm

aggregates under consistent, controlled conditions to examine the evolution of collective sperm

behavior and empirically test whether sperm aggregation improves swimming performance.

Materials and Methods

(a) Sperm collection

We obtained captive *Peromyscus maniculatus bairdii, P. polionotus subgriseus, P. leucopus*,

P. eremicus, and *P. californicus insignis* from the Peromyscus Genetic Stock Center at the

University of South Carolina, and *P. gossypinus gossypinus* from Dr. Hopi Hoekstra at Harvard

University and housed them in same-sex cages at 22ºC on a 16L:8D cycle in accordance with

guidelines established by the Institutional Animal Care and Use Committee at the University of

Maryland in College Park (protocol R-Jul-18-38). We sought samples from all available captive

Peromyscus species and avoided wild-caught specimens to control for variation due to life

experience. We obtained sperm samples from sexually mature males and accounted for relatedness

among the focal males by assigning siblings a unique 'Family' ID. We euthanized males via

isoflurane overdose and cervical dislocation, then weighed each male and both testes (Mettler

Toledo, Switzerland). Next, we removed a single caudal epididymis, made several small incisions

in the tissue, and submersed it in sperm medium (Modified Sperm Washing Medium, Irvine

Scientific, USA) that was pre-warmed at 37ºC; to reduce differences in sperm density despite

natural variation in epididymal sizes, we varied the volume (50μl - 1000μl) based on tissue size

and accounted for these differences when estimating final sperm counts for each male. To collect

sperm, we agitated the tissue at 300rpm (ThermoMixer F1.5, Eppendorf, Germany) at 37ºC for ten

minutes, inverting the tube at the five- and ten-minute mark, then incubated the tissue undisturbed

for two minutes. Using pipette tips cut to create a wider opening, we collected live sperm cells for

analysis from just below the meniscus of the solution to enrich for the most motile sperm [35].

Next we estimated sperm density using a computer-assisted sperm analysis (CASA) system (Ceros

II Animal, Hamilton Thorne, USA) and verified with a Neubauer-improved hemocytometer

(Marienfeld, Germany) then diluted samples with pre-warmed medium to reach a standard

concentration of 300-400 cells summed across the five 5-second videos at 100X magnification for

cell tracking optimality and efficiency.

(b) Live sperm analysis

To conduct live sperm observations, we gently reverse pipetted 4μl of the sperm solution into

12μl of pre-warmed medium on a plastic slide within a 9mm x 0.12mm imaging spacer (Grace

Bio-Labs, USA) and covered by a plastic cover slip. This set-up served as a control and represents

a 'low-viscosity' environment. To test sperm motility in a 'high-viscosity' environment, we

followed the same procedures except that we mixed 4μl of sperm solution with 12μl of pre-

warmed medium enriched with methylcellulose (Sigma Aldrich M 7140; 15cP, 2% in water; [36]).

We recorded a minimum of five 5-second videos at 60 frames/second per male, except in one case

we were only able to record four videos yet were able to acquire an adequate number of observed

cells per male. Videos were recorded at 59±1 minutes post-harvest from the epididymal tissue,

dependent on dilutions. The number of cells analyzed are reported in Table 1.

We characterized sperm aggregation by scoring videos using direct observations because

CASA tracks particles, and thus each track may represent a single cell or an aggregate. We

counted the number of cells represented by each track on at least three different frames/track.

From these data, we calculated the proportion of cells that aggregated for each male by dividing

the total number of aggregates by the total number of motile cells across all tracks [37]. Then we

calculated the mean number of cells in aggregate (i.e., aggregate size) by dividing the sum of cells

in aggregate by the sum of aggregates, both across all tracks for each male and across all males for

each species. Finally, we calculated the coefficient of variation (CV) for both the proportion and

number of cells that aggregated within each species using the following formula: (standard

deviation/mean) x 100.

During video scoring, we identified aggregates that contained cells that were facing different

directions within the group, immotile, morphologically abnormal, or stuck to the microscope slide.

We used this dataset to characterize differences in sperm aggregation for *P. maniculatus* and *P.*

polionotus, the two species whose sperm were observed to aggregate the most extensively. We

characterized aggregates as: 'aligned' if all sperm adhered to one another in a head-to-flagella

orientation and contained no immotile, morphologically abnormal or damaged cells; 'defective' if

one or more sperm cell was marked as morphologically abnormal (kinked or broken), immotile, or

stuck to the slide; or 'opposed' if sperm were otherwise morphologically normal and motile but

were not oriented in the same direction, which included star-shaped aggregates [28]. We then used

these data to calculate the proportion of aggregates scored for these criteria within each male for

these two species.

We recorded the following metrics for each recorded track (i.e., single sperm and aggregates):

straight-line velocity (VSL; calculated using the straight-line distance between the first and last

detected positions of the sperm head, divided by the time taken to swim the track; also known as

average velocity), *curvilinear velocity* (VCL; calculated using the summed distance between the

sperm head positions in each frame divided by the time taken to travel the track; also known as

speed), *average path velocity* (VAP; the time-averaged velocity of the sperm head along its

average path), and *linearity* (LIN; the ratio of VSL to VCL to measure the straightness of the

trajectory; [38]). We calculated the mean of each kinematic parameter for both single cells and

sperm aggregates separately for each of three populations of sperm cells: all cells, motile cells

(i.e., devoid of visually inspected tracks in which cells were unmoving, stuck, or featured an

obvious morphological abnormality such as a kinked midpiece), and progressively motile cells

(i.e., motile cells with a VSL $\geq$ 25μm/sec). We used these data to calculate the proportion of

motile aggregates by dividing the sum of motile aggregates by the total number of sperm

aggregates, and the proportion of progressively motile aggregates by dividing the sum of

progressive aggregates by the total number of sperm aggregates for each male [37]. For our

kinematic analyses, we focused on the motile sperm dataset to eliminate artifacts from damaged or

dead cells, and the total sperm population to quantify frequency of aggregation and aggregate size

in an effort to capture natural aggregation rates (results from other cell populations are reported in

Table S1, Figure S1).

(c) Statistical Analyses

We performed all statistical analyses using R version 3.4.2 [39] and created all figures using

the 'ggplot2' package with R [40]. Generalized linear mixed models (GLMM) were analyzed

using the glmer function from the "lme4" R package and a logit link function [41]. We also used

this package to analyze linear mixed models (LMM) using the lmer function and generalized linear

models (GLM) using the glm function. We screened for collinearity using linear models and

variance inflation factor (VIF) values and used model selection to drop non-significant explanatory

variables one at a time based on model comparisons (using Akaike information criterion changes <

2 for GLMMs or an analysis of variance test for LMMs) to determine the minimal adequate

model. To validate model normality, we visually inspected diagnostic plots (qqplots and plots of

the distribution of the residuals against fitted values). Only the best fitting models are reported

here. Post-hoc pairwise comparisons of final models concerning aggregate frequency and size

were performed using Tukey HSD adjustments for multiple comparisons from the "LSmeans" R

package [42].

We used multiple data sets and statistical approaches to analyze the frequency and size of

sperm aggregates. For the mean proportion of aggregated cells by male, we used a GLMM in

which the binomial response was the number of sperm cells in aggregate, and the total number of

sperm cells was the binomial denominator. We used an observation-level random effect (OLRE) to

control for overdispersion [37,43] as well as family ID as random factors in these analyses. For the

mean proportion of aggregated cells by species, we included an ultra-metric phylogenetic tree of

Peromyscus as a covariate in our regression analyses using a phylogenetic generalized least

squares (PGLS) approach [44,45] using the "caper" [46] and "APE" [47] packages in R. For the

mean number of aggregated cells by male, excluding one *P. californicus* male whose measurement

represented a clear outlier, we initially used a LMM but then reverted to using a linear model (LM)

because the family ID random factor did not significantly contribute to the residual variability in

the response variable. For the mean number of aggregated cells by species, we used a PGLS

approach. We also conducted similar PGLS analyses using the within-species CV for each of these

parameters. When assessing the relationship between relative testis size and aggregation, our

models included testis mass with body mass as a separate fixed factor, a method better suited to

estimating relative testis weight is size rather than using the ratio of testis to body mass or

residuals [48,49].

We used multiple data sets and statistical approaches to further analyze sperm aggregation.

We used generalized linear models (GLM) to analyze the composition and orientation of sperm

aggregates in *P. maniculatus* and *P. polionotus,* paired student's t-test to compare the VCL of

aggregates based on their composition or orientation within species, and GLMMs to assess species

differences in the proportion of aggregates that were motile or progressively motile. To determine

if sperm aggregates have motility or force benefits over single cells, we initially performed a

principal component analysis (PCA) using three related swimming performance measures (VSL,

VCL, and VAP) to reduce dimensionality and obtain a composite measure for motile solitary and

aggregated sperm cells in both low- and high-viscosity media. Because we found that species

distributions overlapped for this composite measure for both single sperm and aggregates in low-

and high-viscosity media (Figures S2 and S3), we focused on individual kinematic parameters

(VSL, VCL, VAP, LIN) within each species separately. To determine if sperm aggregates have

motility benefit over single cells in either low- or high-viscosity media, we used paired student's t-

tests to compare each kinematic parameter between solitary and aggregated sperm cells within

males for each species within each environment.

Results

We found significant differences in the proportion of aggregated cells among *Peromyscus*

species (binomial GLMM: n = 134, P < 0.001; Table 1, 2), with more variance across species than

within species (s^2 across species = 0.10; s^2 within species < 0.10). Pairwise comparisons adjusted

for multiple comparisons revealed that species that are most closely related within the phylogeny

do not significantly differ from one another, whereas all other pairwise species comparisons do

(Figure 1). *P. maniculatus* and *P. polionotus* both have the highest frequencies of aggregation,

despite having different mating systems, which corresponds with our finding that relative testis

mass does not significantly associate with the proportion of cells aggregated (binomial GLMM: n

= 127, P = 0.163286). We also found no association between the frequency of sperm aggregation

and sperm velocity (VCL: binomial GLMM: n = 130, P = 0.1435). In examining the coefficient of

variation (CV) for the proportion of cells aggregated within each species, we found 68.8% for *P.*

californicus, 50.6% for *P. eremicus*, 14.4% for *P. polionotus*, 10.8% for *P. maniculatus*, 125.7%

for *P. leucopus*, and 60.5% for *P. gossypinus*. In a cross-species analysis controlling for

phylogenic relationships and body mass, we found no effect of testis size on the variation present

within species for the proportion of aggregated cells ($F_{2,3}$ = 0.3603, P = 0.5604).

For the number of aggregated cells, we found significant differences among species (LM:

$F_{6,126}$ = 56.37, P < 0.001; Table 1, 2), with more variance observed across species (s^2 = 1.96) than

within species ($s^2 < 1.00$, except for *P. polionotus* [$s^2 = 2.15$]). *P. maniculatus* and *P. polionotus*

produce the largest sperm aggregates, whereas *P. gossypinus*, *P. leucopus*, and *P. californicus*

produce the smallest aggregates ($P < 0.05$ for pairwise comparisons), the latter of which produces

sperm cell aggregates that are statistically similar in size to those produced by *P. eremicus* ($P =$

0.9203; Figure 2A). In a cross-species analysis controlling for phylogenic relationships and body

mass, we found a significant negative effect of a species' mean testis weight on the species'

variation in the number of cells in aggregate ($F_{2,3} = 8.398$, $P = 0.02655$; Figure 2B). The CV for

the number of cells aggregated within each species were as follows:: 26.5% for *P. californicus*,

19.2% for *P. eremicus*, 27.5% for *P. polionotus*, 18.4% for *P. maniculatus*, 13.3% for *P. leucopus*,

and 7.2% for *P. gossypinus*.

Within the two species that produce the largest and most frequent aggregates, sperm cells

within aggregates are significantly more aligned in a head-to-flagella orientation in *P. maniculatus*

(99.1%, 731/738) than in *P. polionotus* (87.5%, 720/826; GLM: $n = 42$, $P = 3.92e-12$), a feature

that significantly positively correlates with VCL in both species (Figure 3; paired t-tests: *P.*

maniculatus $t = 5.9627$, $df = 4$, $P = 0.003972$; *P. polionotus* $t = 11.247$, $df = 11$, $P = 2.257e-07$).

We also found that there are significantly fewer sperm aggregates in *P. maniculatus* with

immotile, stuck, or morphologically abnormal cells (8.1%, 64/795) than in *P. polionotus* (11.5%,

94/814; binomial GLM: $n = 40$, $P = 0.00503$), a feature that we found significantly reduces VCL

in both species (*P. maniculatus* $VCL_{aligned} = 179.60 \pm 6.88$ μm/sec, $VCL_{defective} = 132.47 \pm 6.18$

μm/sec, paired t-test: $t = 20.627$, $df = 14$, $P = 7.075e-12$; *P. polionotus* $VCL_{aligned} = 135.48 \pm 3.51$

μm/sec, $VCL_{defective} = 115.62 \pm 5.22$ μm/sec, paired t-test: $t = 16.312$, $df = 20$, $P = 5.079e-13$).

When comparing the proportion of motile and progressively motile aggregates across all

species, we found *P. eremicus* produced the smallest proportion of motile aggregates ($P < 0.05$ for

all *P. eremicus* pairwise comparisons; $P > 0.05$ for all other pairwise comparisons). Fitted values

of the proportion of motile aggregates using LSmeans were 0.77 ± 0.04 for *P. eremicus*, $0.91 \pm$

0.02 for *P. californicus*, 0.91 ± 0.02 for *P. gossypinus*, 0.92 ± 0.02 for *P. polionotus*, 0.93 ± 0.02

for *P. leucopus*, and 0.95 ± 0.01 for *P. maniculatus*. Similarly, *P. eremicus, P. polionotus,* and *P.*

gossypinus all had the smallest proportions of progressively motile aggregates ($P < 0.05$ for all

pairwise comparisons; fitted values using LSmeans were 0.64 ± 0.06, 0.65 ± 0.05, and 0.75 ± 0.05,

respectively), the latter species of which did not significantly differ from *P. californicus* ($P =$

0.1981; 0.87 ± 0.02 for LSmeans fitted values). Conversely, *P. maniculatus* and *P. leucopus* had

similarly greater proportions of progressively motile aggregates ($P = 0.3417$; fitted values using

LSmeans were 0.94 ± 0.02 and 0.91 ± 0.02, respectively), the latter of which did not differ from *P.*

californicus ($P = 0.9109$).

Overall, we found species-specific differences in the effect of sperm aggregation on motility,

regardless of environmental complexity (Table 3, Figure 4). In low-viscosity medium, we found

that sperm aggregates have a significantly greater VCL in *P. maniculatus*, VSL in *P. maniculatus*,

P. leucopus, and *P. californicus*, LIN in *P. leucopus*, and VAP in *P. californicus* compared to

single cells but a significantly lower VCL, VSL, and VAP velocity in *P. polionotus* and *P.*

gossypinus than single sperm (Figure 4). In the high-viscosity medium, we found that sperm

aggregates have a significantly greater VCL in *P. maniculatus* and *P. californicus* and a higher

VSL and VAP in *P. californicus* compared to single sperm but a significantly lower LIN in *P.*

californicus, *P. eremicus*, *P. polionotus*, and *P. maniculatus* as well as a reduced VSL and VAP in

P. polionotus than single cells (Figure 4).

Discussion

While it is known that collective sperm behaviors have evolved independently in a number of

taxa [2], it remains unclear how sperm aggregation evolves among closely-related species. Our

comparative study reveals that sperm aggregating behaviors vary across mice in the genus

Peromyscus, and that the evolution of this trait is complex. We observed an effect of phylogenetic

history on the frequency of sperm aggregation, indicating that collective sperm behavior likely

evolved prior to the divergence of present-day species. Additionally, we find a negative

association between relative testis weight, a robust proxy for intensity of sperm competition in

rodents [50], and the coefficient of variation for the number of cells in aggregate, suggesting that

sexual selection has a stabilizing effect on sperm aggregate size. We then compared the motility of

single and aggregated sperm across all species under low- and high-viscosity conditions, and show

that aggregation is not kinematically beneficial for all species, regardless of environmental

complexity. Next, by examining sperm from the species that aggregate the most frequently, we

found support for the theoretical hypothesis [28] that cellular orientation within the group is

critical to its kinematics. Moreover, we observed more aggregates with efficient geometry in the

species that evolved under strong post-copulatory sexual selection compared to its monogamous

sister-species. These findings support the prediction that sperm aggregation likely evolved in

Peromyscus to improve motility in a competitive context but that relaxed selection may have

enabled less efficient strategies to persist, thereby generating diversity in collective sperm

behaviors within these closely-related species.

Our results reveal distinct species-specific differences across *Peromyscus* mice in the

frequency of sperm aggregation and the average size of these cellular groups. Multiple

Peromyscus species produce sperm that aggregate more extensively than other studied muroid

rodents [51], with the notable exception of the wood mouse (*Apodemus sylvaticus*; [4]). In general,

the proportion of sperm cells that aggregate is most similar within each sister-species pair.

Specifically, *P. maniculatus* and *P. polionotus* produce sperm that aggregate the most (>80% of

sperm), but the pair they are most closely-related to, *P. gossypinus* and *P. leucopus,* are the species

least likely to produce sperm that aggregate (<10% of sperm); the most distantly-related species

pair we assessed, *P. californicus* and *P. eremicus*, both produce sperm with a moderate propensity

to aggregate (~25% of sperm). These findings support one of two possibilities for the evolution of

sperm aggregates within these species: (a) a genus-wide ancestral trait of moderate sperm

aggregation with subsequent diversification leading to an increase in *P. maniculatus* and *P.*

polionotus and a decrease in *P. gossypinus* and *P. leucopus*, or (b) the independent evolution of

aggregation in the ancestors of the *P. californicus*-species pair and the *P. maniculatus*-species pair.

Such complex evolutionary histories with losses and recurrences of sperm conjugation, and

subsequent species divergence, have also been demonstrated in diving beetles (Dytiscidae [52]),

consistent with the evolutionary lability that we observe. Our experimental results may explain

selection against sperm aggregation if forming groups reduces sperm swimming performance,

which we find in at least one species, *P. gossypinus*. However, we found that sperm produced by

their sister-species, *P. leucopus*, also rarely aggregate, despite our observation that these rare

collective groups have a greater average velocity and are more linear than single cells.

Interestingly, we observed that these two promiscuous species that rarely aggregate, *P. gossypinus*

and *P. leucopus*, have the largest relative testes of the species studied, suggesting that they may

have evolved increased sperm production to improve competitive fertilization success [53], rather

than adaptations that influence motility [54]. Moreover, within these species (and *P. polionotus*)

we found a negative association between sperm speed and aggregation, even after controlling for

sperm density, indicating that differences in aggregate formation are not simply a by-product of

increased encounter rates. Together these results suggest different strategies employed by

divergent species across the *Peromyscus* genus in response to sperm competition.

In contrast to our results on the frequency of sperm aggregation within species, we found that

the average size of sperm groups does not align as closely with phylogenetic relationships. In

addition, we found that species with relatively larger testes, which is positively associated with

increased sperm competition [50], exhibit less variation (CV) in aggregate size. This result

supports the prediction that relaxed sperm competition allows for greater intermale variation to

persist in a population [55] and suggests that this post-copulatory sexual selection may be

stabilizing sperm aggregate size for a species-specific 'optimum' [28]. Similarly, other studies

have shown that the strength of sexual selection regulates variance in sperm morphology across

taxa and at multiple levels of organization, including within- and between-males [56–58] and

within- and between-species [55,59]. A study on sperm bundles across ten *Carabus* ground beetles

also found intense selection on bundle size, which are dimorphic and either small or large; the

large, but not small, sperm bundles are positively correlated with measures of sperm competition

risk, including copulatory piece length and mate guarding, suggesting that diversity of large sperm

bundles is associated with sexual selection [60]. In contrast to these findings that competition

drives sperm-sperm interactions, a study on the evolution of such sperm traits in diving beetles

found that variation in sperm conjugation is more associated with female reproductive tract

architecture [61]. Therefore, while our results suggest that stabilizing selection on sperm aggregate

size is associated with an increase in sperm competition given the correlation with relative testis

weight, mechanisms of female control [62] may also play an important evolutionary role.

We compared the motility of single and aggregated sperm sampled from the same male to test

whether sperm aggregates swim faster or more efficiently than single sperm, which is predicted if

the combined force of multiple flagella enhances their motility [4]. We found improved kinematic

measures in half of the species studied (*P. maniculatus, P. californicus*, and *P. leucopus*), thus

supporting this functional hypothesis; however, in several species we found aggregation had a

negative (*P. polionotus* and *P. gossypinus*) or no (*P. eremicus*) impact on motility relative to single

sperm cells. While we found support for a theoretical prediction that sperm aggregates achieve

greater straight-line velocity because they move in a more linear path of travel rather than at a

faster speed (i.e., curvilinear velocity; [28]) in two of our six focal species (*P. californicus* and *P.*

leucopus), we did not find this kinematic benefit for sperm aggregation in all *Peromyscus* species.

These results corroborate other studies in more disparate taxonomic groups that have quantified

sperm aggregation motility and found inconsistent results. For example, sperm trains exhibit

greater swimming progressive motility in the wood mouse [4], and greater velocity than individual

sperm in the Norway rat, but not the house mouse [5]. In invertebrates, the swimming velocity of

fishfly sperm increases with number of sperm in a bundle [10], but in a marine snail, there is no

differences in swimming speed between paired and single sperm [14]. One possible explanation

for these differences in sperm aggregate swimming performance across taxa is that cell orientation

within a group is critical for its collective motility.

Sperm cells are predicted to be faster if they generate increased force with proportionally less

drag; such effects may also be true for sperm aggregations in which cells conjoin head-to-tail,

thereby increasing the length of the collective unit [2]. In *Peromyscus,* sperm conjoin head-to-head

to form aggregates and are thus wider, not longer*,* yet can still offer a motility advantage for some

species. This kinematic benefit, however, was not observed in all species, which led us to

hypothesize that aggregate motility is dependent on the orientation of sperm within a group and

whether it contains abnormal or immotile cells. When sperm heads and flagella are not oriented in

the same direction, the cells within an aggregate are expected to exert opposing forces on one

another, thereby reducing the overall motility of the group [27,28]. To test this hypothesis, we

quantified aggregate geometry and composition in the two species that form the most frequent and

largest sperm aggregates. We found that in *P. maniculatus*, a promiscuous species, sperm were

more likely form aggregates with optimal geometry, compared to sperm from the monogamous *P.*

polionotus, which frequently included misaligned, opposing or morphologically abnormal cells.

Consistent with our prediction, *P. maniculatus* aggregates exhibit greater straight-line and

curvilinear velocity compared to single cells, but the opposite was true for *P. polionotus*. This

finding is consistent with a previous report that *P. polionotus* sperm tend to form optimal-sized

aggregates less often than in *P. maniculatus* [28]. Together, these observations further support the

hypothesis that sperm aggregation evolved prior to the divergence of the species pair [12], and

when monogamy evolved secondarily in *P. polionotus* [34,63], relaxed sexual selection allowed

for the persistence of less motile sperm traits. In line with this prediction, we observed the smallest

proportion of motile and progressively motile sperm aggregates in another monogamous species,

P. eremicus, but the largest proportion of progressively motile aggregates in two promiscuous

species, *P. maniculatus* and *P. leucopus*. Similar results have been reported in house mice, in

which males evolving under intense sperm competition produced a greater proportion of motile

sperm, compared to males from lineages subject to relaxed selection [64]. Based on these findings,

we suggest that relaxed sexual selection may allow the persistence of less optimal strategies in

Peromyscus.

When we compared single and aggregated sperm in a viscous environment to test if

aggregation improves motility in more complex fluids, similar to the secretions or cervical mucus

[18] of the female reproductive tract [17,65], we found that collective motion is only beneficial in

some species. Compared to single sperm, aggregates swim with greater curvilinear velocity in *P.*

maniculatus and *P. californicus*, and greater straight-line and average path velocities in *P.*

californicus, but aggregates were less linear than single cells in all species except for *P. leucopus*

and *P. gossypinus*. While other studies have found kinematic benefits for sperm conjugates

compared to single cells in higher viscosities in the gray short-tailed opossum [16], the wood

mouse [4], and the fishfly [10], a study in bulls found that sperm were slower, exhibited less

organized swimming patterns, and were less likely to cluster in viscous fluids [15], both of which

are consistent with our analysis of *Peromyscus* aggregates. Ultimately, the benefit of sperm

aggregation depends on the relative importance of each kinematic parameter during sperm

migration *in vivo*. Although beyond the scope of this study, we predict that improved linearity

afforded by collective motion may help to direct the sperm through the dynamic structures and

fluids of the female reproductive tract [66] and that increased velocity will reduce the time it takes

for sperm to arrive at the fertilization site.

In conclusion, our study highlights the diversity of sperm aggregation within a single

taxonomic lineage and how selection has shaped the formation and performance of these cellular

groups. We show that both evolutionary history and varying levels of post-copulatory sexual

selection influence the frequency and size of sperm groups. Moreover, we find that sperm

aggregation can improve sperm motility in both simple and complex fluids, but this is not

consistent across all species. Theoretical predictions [27,28] and emerging empirical evidence

suggests that motility benefits may only be realized if cells maintain optimal alignment within the

groups and, if achieved, may provide these sperm with a competitive advantage in the female

reproductive tract [61]. Future work investigating sperm aggregates *in vivo* [e.g., 66,67] will shed

light on the co-evolution of these unique gametic behaviors with the remarkably variable and

dynamic female reproductive tracts through which sperm must navigate.

References:

1. Pitnick S, Hosken DJ, Birkhead TR. 2009 Sperm Morphological Diversity. In *Sperm Biology:*
An Evolutionary Perspective (eds TR Birkhead, DJ Hosken, S Pitnick), pp. 69–149. London:
Academic Press. (doi:10.1016/B978-0-12-372568-4.00003-3)

2. Higginson DM, Pitnick S. 2011 Evolution of intra-ejaculate sperm interactions: do sperm
cooperate? *Biological Reviews* **86**, 249–270. (doi:10.1111/j.1469-185X.2010.00147.x)

3. Immler S. 2008 Sperm competition and sperm cooperation: the potential role of diploid and
haploid expression. *Reproduction* **135**, 275–283. (doi:10.1530/REP-07-0482)

4. Moore HD, Dvoráková K, Jenkins N, Breed W. 2002 Exceptional sperm cooperation in the
wood mouse. *Nature* **418**, 174–177.

5. Immler S, Moore HD, Breed W, Birkhead T. 2007 By hook or by crook? Morphometry,
competition and cooperation in rodent sperm. *PloS one* **2**, e170.

6. Birkhead TR, Møller AP, Sutherland WJ. 1993 Why do females make it so difficult for males
to fertilize their eggs? *Journal of Theoretical Biology* **161**, 51–60.
(doi:10.1006/jtbi.1993.1039)

7. Pizzari T, Foster KR. 2008 Sperm sociality: cooperation, altruism, and spite. *PLOS Biology* **6**,
0925–0931. (doi:10.1371/journal.pbio.0060130)

8. Birkhead TR, Martinez JG, Burke T, Froman DP. 1999 Sperm mobility determines the outcome
of sperm competition in the domestic fowl. *Proceedings of the Royal Society of London.*
Series B: Biological Sciences **266**, 1759–1764. (doi:10.1098/rspb.1999.0843)

9. Gage MJG, Macfarlane CP, Yeates S, Ward RG, Searle JB, Parker GA. 2004 Spermatozoal
traits and sperm competition in Atlantic salmon: relative sperm velocity is the primary
determinant of fertilization success. *Current Biology* **14**, 44–47.
(doi:10.1016/j.cub.2003.12.028)

10. Hayashi F. 1998 Sperm co-operation in the Fishfly, *Parachauliodes japonicus*. *Functional*
Ecology **12**, 347–350. (doi:10.1046/j.1365-2435.1998.00205.x)

11. Woolley DM, Crockett RF, Groom WDI, Revell SG. 2009 A study of synchronisation
between the flagella of bull spermatozoa, with related observations. *Journal of Experimental*
Biology **212**, 2215–2223. (doi:10.1242/jeb.028266)

12. Fisher HS, Hoekstra HE. 2010 Competition drives cooperation among closely related sperm
of deer mice. *Nature* **463**, 801.

13. Pearcy M, Delescaille N, Lybaert P, Aron S. 2014 Team swimming in ant spermatozoa.
Biology Letters **10**, 20140308. (doi:10.1098/rsbl.2014.0308)

14. Ishijima S, Ishijima SA, Afzelius BA. 1999 Movement of *Turritella* spermatozoa: Direction of
propagation and chirality of flagellar bends. *Cell Motility* **44**, 85–95.
(doi:10.1002/(SICI)1097-0169(199910)44:2<85::AID-CM1>3.0.CO;2-#)

15. Tung C, Lin C, Harvey B, Fiore AG, Ardon F, Wu M, Suarez SS. 2017 Fluid viscoelasticity
promotes collective swimming of sperm. *Scientific Reports* **7**, 1–9. (doi:10.1038/s41598-017-
03341-4)

16. Moore HDM, Taggart DA. 1995 Sperm pairing in the opossum increases the efficiency of
sperm movement in a viscous environment. *Biology of Reproduction* **52**, 947–953.
(doi:10.1095/biolreprod52.4.947)

17. Suarez SS. 2016 Mammalian sperm interactions with the female reproductive tract. *Cell and
Tissue Research* **363**, 185–194. (doi:10.1007/s00441-015-2244-2)

18. Hanson FW, Overstreet JW. 1981 The interaction of human spermatozoa with cervical mucus
in vivo. *American Journal of Obstetrics and Gynecology* **140**, 173–178. (doi:10.1016/0002-
9378(81)90105-8)

19. Phillips DM. 1970 Ultrastructure of spermatozoa of the wooly opossum *Caluromys philander.
Journal of Ultrastructure Research* **33**, 381–397.

20. Tung KSK, Okada A, Anagimachi R. 1980 Sperm autoantigens and fertilization. I. effects of
antisperm autoantibodies on rouleaux formation, viability, and acrosome reaction of guinea
pig spermatozoa. *Biology of Reproduction* **23**, 877–886. (doi:10.1095/biolreprod23.4.877)

21. Bedford JM, Rodger JC, Breed WG. 1984 Why so many mammalian spermatozoa - a clue
from marsupials? *Proceedings of the Royal Society of London B: Biological Sciences* **221**,
221–233. (doi:10.1098/rspb.1984.0031)

22. Higginson DM, Henn KRH. 2012 Effects of sperm conjugation and dissociation on sperm
viability *in vitro. PloS one* **7**, e34190.

23. Mackie JB, Walker MH. 1974 A study of the conjugate sperm of the dytiscid water beetles
Dytiscus marginalis and *Colymbetes fuscus. Cell Tissue Res.* **148**, 505–519.
(doi:10.1007/BF00221935)

24. McGlinn SM, Shepherd BA, Martan J. 1979 Rouleau Formation and fertility of spermatozoa
in guinea pigs treated with cyproterone acetate. *Archives of Andrology* **2**, 197–202.
(doi:10.3109/01485017908987313)

25. Taggart DA, Johnson JL, O'Brien HP, Moore HDM. 1993 Why do spermatozoa of the
American marsupial form pairs? A clue from the analysis of sperm-pairing in the epididymis
of the grey short-tailed opossum, *Monodelphis domestica. Anatomical Record* **236**, 465–478.

26. Nixon B, Ecroyd H, Dacheux J-L, Dacheux F, Labas V, Johnston SD, Jones RC. 2016
Formation and dissociation of sperm bundles in monotremes. *Biol Reprod* **95**, 1–11.
(doi:10.1095/biolreprod.116.140491)

27. Pearce DJG, Hoogerbrugge LA, Hook KA, Fisher HS, Giomi L. 2018 Cellular geometry
controls the efficiency of motile sperm aggregates. *Journal of The Royal Society Interface* **15**,
20180702. (doi:10.1098/rsif.2018.0702)

28. Fisher HS, Giomi L, Hoekstra HE, Mahadevan L. 2014 The dynamics of sperm cooperation in
a competitive environment. *Proceedings of the Royal Society of London B: Biological*
Sciences **281**, 20140296.

29. Dewsbury DA. 1985 Interactions between males and their sperm during multi-male copulatory
episodes of deer mice (*Peromyscus maniculatus*). *Animal Behaviour* **33**, 1266–1274.
(doi:10.1016/S0003-3472(85)80186-X)

30. Birdsall DA, Nash D. 1973 Occurrence of successful multiple insemination of females in
natural populations of deer mice (*Peromyscus maniculatus*). *Evolution* **27**, 106–110.
(doi:10.2307/2407124)

31. Dewsbury DA. 1981 An exercise in the prediction of monogamy in the field from laboratory
data on 42 species of muroid rodents. *Biologist* **63**, 138–162.

32. Foltz DW. 1981 Genetic evidence for long-term monogamy in a small rodent, *Peromyscus*
polionotus. *The American Naturalist* **117**, 665–675. (doi:10.1086/283751)

33. Bedford NL, Hoekstra HE. 2015 The natural history of model organisms: *Peromyscus* mice as
a model for studying natural variation. *eLife Sciences* **4**, e06813. (doi:10.7554/eLife.06813)

34. Turner LM, Young AR, Römpler H, Schöneberg T, Phelps SM, Hoekstra HE. 2010
Monogamy evolves through multiple mechanisms: evidence from V1aR in deer mice.
Molecular Biology and Evolution **27**, 1269–1278. (doi:10.1093/molbev/msq013)

35. Magdanz V, Boryshpolets S, Ridzewski C, Eckel B, Reinhardt K. 2019 The swim-up
technique separates bovine sperm by metabolic rates, motility and tail length. *bioRxiv* ,
624502. (doi:10.1101/624502)

36. Suarez SS, Dai X. 1992 Hyperactivation enhances mouse sperm capacity for penetrating
viscoelastic media. *Biol Reprod* **46**, 686–691. (doi:10.1095/biolreprod46.4.686)

37. Crawley MJ. 2013 *The R Book Second Edition*. England: John Wiley & Sons, Ltd.

38. World Health Organization. 2010 WHO laboratory manual for the examination and processing
of human semen.

39. R Core Team. 2016 *R: A Language and Environment for Statistical Computing*. Vienna,
Austria: R Foundation for Statistical Computing. See https://www.R-project.org.

40. Wickham H. 2016 *ggplot2: Elegant Graphics for Data Analysis*. Springer.

41. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting linear mixed-effects models using
lme4. *Journal of Statistical Software* **67**, 1–48.

42. Lenth RV. 2016 Least-squares means: The R package 'lsmeans'. *Journal of Statistical
Software* **69**, 1–33. (doi:10.18637/jss.v069.i01)

43. Harrison XA. 2014 Using observation-level random effects to model overdispersion in count
data in ecology and evolution. *PeerJ* **2**, e616. (doi:10.7717/peerj.616)

44. Pagel M. 1999 Inferring the historical patterns of biological evolution. *Nature* **401**, 877–884.
(doi:10.1038/44766)

45. Freckleton RP, Harvey PH, Pagel M. 2002 Phylogenetic analysis and comparative data: a test
and review of evidence. *The American Naturalist* **160**, 712–726.

46. Orme D, Freckleton R, Thomas G, Petzoldt T. 2013 The caper package: comparative analysis
of phylogenetics and evolution in R. *R package version* **5**, 1–36.

47. Paradis E, Claude J, Strimmer K. 2004 APE: Analyses of phylogenetics and evolution in R
language. *Bioinformatics* **20**, 289–290. (doi:10.1093/bioinformatics/btg412)

48. García-Berthou E. 2001 On the misuse of residuals in ecology: testing regression residuals vs.
the analysis of covariance. *Journal of Animal Ecology* **70**, 708–711. (doi:10.1046/j.1365-
2656.2001.00524.x)

49. Lüpold S, Calhim S, Immler S, Birkhead TR. 2009 Sperm morphology and sperm velocity in
passerine birds. *Proceedings of the Royal Society B: Biological Sciences* **276**, 1175–1181.
(doi:10.1098/rspb.2008.1645)

50. Ramm SA, Parker GA, Stockley P. 2005 Sperm competition and the evolution of male
reproductive anatomy in rodents. *Proceedings of the Royal Society B: Biological Sciences*
272, 949–955. (doi:10.1098/rspb.2004.3048)

51. Tourmente M, Zarka-Trigo D, Roldan ERS. 2016 Is the hook of muroid rodent's sperm
related to sperm train formation? *Journal of Evolutionary Biology* **29**, 1168–1177.
(doi:10.1111/jeb.12857)

52. Higginson DM, Miller KB, Segraves KA, Pitnick S. 2012 Convergence, recurrence and
diversification of complex sperm traits in diving beetles (Dytiscidae). *Evolution* **66**, 1650–
1661.

53. Parker GA. 1982 Why are there so many tiny sperm? Sperm competition and the maintenance
of two sexes. *Journal of Theoretical Biology* **96**, 281–294. (doi:10.1016/0022-5193(82)90225-
9)

54. Snook RR. 2005 Sperm in competition: not playing by the numbers. *Trends in Ecology &
Evolution* **20**, 46–53. (doi:10.1016/j.tree.2004.10.011)

55. Calhim S, Immler S, Birkhead TR. 2007 Postcopulatory sexual selection is associated with
reduced variation in sperm morphology. *PLoS ONE* **2**, e413.
(doi:10.1371/journal.pone.0000413)

56. Immler S, Calhim S, Birkhead TR. 2008 Increased postcopulatory sexual selection reduces the
intramale variation in sperm design. *Evolution* **62**, 1538–1543. (doi:10.1111/j.1558-
5646.2008.00393.x)

57. Fitzpatrick JL, Baer B. 2011 Polyandry reduces sperm length variation in social insects.
Evolution **65**, 3006–3012. (doi:10.1111/j.1558-5646.2011.01343.x)

58. Carballo L, Battistotti A, Teltscher K, Lierz M, Bublat A, Valcu M, Kempenaers B. 2019
Sperm morphology and evidence for sperm competition among parrots. *Journal of
Evolutionary Biology* **32**, 856–867. (doi:10.1111/jeb.13487)

59. Rowley A *et al.* 2019 Sexual selection and the evolution of sperm morphology in sharks.
Journal of Evolutionary Biology **32**, 1027–1035. (doi:10.1111/jeb.13501)

60. Takami Y, Sota T. 2007 Sperm competition promotes diversity of sperm bundles in
Ohomopterus ground beetles. *Naturwissenschaften* **94**, 543–550. (doi:10.1007/s00114-007-
0225-3)

61. Higginson DM, Miller KB, Segraves KA, Pitnick S. 2012 Female reproductive tract form
drives the evolution of complex sperm morphology. *Proceedings of the National Academy of
Sciences* **109**, 4538–4543.

62. Eberhard W. 1996 *Female Control: Sexual Selection by Cryptic Female Choice*. Princeton
University Press.

63. Greenbaum IF, Baker RJ, Ramsey PR. 1978 Chromosomal evolution and the mode of
speciation in three species of *Peromyscus*. *Evolution* **32**, 646–654. (doi:10.2307/2407730)

64. Firman RC, Simmons LW. 2011 Experimental evolution of sperm competitiveness in a
mammal. *BMC Evolutionary Biology* **11**, 19. (doi:10.1186/1471-2148-11-19)

65. Simons JE, Olson SD. 2018 Sperm motility: models for dynamic behavior in complex
environments. In *Cell Movement* (eds M Stolarska, N Tarfulea), pp. 169–209. Cham: Springer
International Publishing. (doi:10.1007/978-3-319-96842-1_7)

66. Ishikawa Y, Usui T, Yamashita M, Kanemori Y, Baba T. 2016 Surfing and swimming of
ejaculated sperm in the mouse oviduct. *Biology of Reproduction* **94**.
(doi:10.1095/biolreprod.115.135418)

67. Wang S, Larina IV. 2018 In vivo three-dimensional tracking of sperm behaviors in the mouse
oviduct. *Development* **145**, dev157685. (doi:10.1242/dev.157685)

68. Bradley, RD, Durish ND, Rogers DS, Miller JR, Engstrom MD, Kilpatrick CW. 2007 Toward
a molecular phylogeny for *Peromyscus*: evidence from mitochondrial cytochrome-*b*
sequences. *Journal of Mammalogy* **88**, 1146–1159. (doi: 10.1644/06-MAMM-A-342R.1)

FIGURES AND TABLES

Figure 1.

The observed natural variation in the proportion of aggregated sperm cells for six closely related species of *Peromyscus* mice (phylogeny adapted from Bradley et al. 2007). Box-plots represent median and interquartile ranges with raw data overlaid. Statistically significant differences at the *P* = 0.05 level are denoted by differing letters; shared letters denote no statistical difference.

Figure 2.

Sperm aggregate size varies among species and is negatively associated with relative testes size. (A) The observed natural variation in the number of aggregated sperm cells for six closely related species of *Peromyscus* mice. Box-plots represent median and interquartile ranges with raw data overlaid. Statistically significant differences at the $P = 0.05$ level are denoted by differing letters; shared letters denote no statistical difference. (B) When controlling for phylogenetic relationships, the coefficient of variation (CV) for the number of aggregated sperm cells negatively correlates with relative testis mass across these species. Note truncated y-axes.

Figure 3.

The effect of sperm aggregate geometry on the curvilinear velocity (*μ*m/sec) of sperm aggregates for two species that aggregated most – *Peromyscus maniculatus* and *Peromyscus polionotus*. Circles represent mean values per male within each species, and black lines represent the mean value within each category. Note truncated y-axis.

Figure 4.

Kinematic parameters of sperm aggregates (closed circles) and solitary sperm cells (open circles) for six species of *Peromyscus* mice in low- and high-viscosity conditions. Circles represent mean values per male, and black lines represent species means. Statistical significance levels comparing aggregated and solitary cells within each species are indicated by * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. Note truncated y-axes.

TABLE 1.

Summary of live sperm aggregate results for mice in the genus *Peromyscus*

Peromyscus Species	No. males	No. sperm	% motile sperm	% progressive sperm	No. sperm aggregates	% cells in aggregate	% motile aggregates	% progressive aggregates
californicus	28	8235	81.2%	78.9%	823	29.7% (2446/8235)	90.5% (745/823)	86.6% (713/823)
eremicus	21	4906	67.4%	58.2%	513	30.2% (1481/4906)	77.8% (399/513)	66.7% (342/513)
polionotus	24	6360	82.4%	57.3%	949	79.5% (5059/6360)	90.9% (863/949)	66.1% (627/949)
maniculatus	18	4991	85.0%	84.3%	822	81.3% (4059/4991)	93.4% (768/822)	92.3% (759/822)
leucopus	22	6341	87.6%	84.5%	361	15.0% (949/6341)	90.3% (326/361)	88.1% (318/361)
gossypinus	21	5970	82.2%	70.3%	254	9.6% (576/5970)	88.6% (225/254)	70.1% (178/254)

Motile sperm cells include only those that exhibited movement patterns. Progressive sperm cells are motile cells with a straight-line velocity ≥ 25 μm/sec.

TABLE 2.

Fixed effects from a binomial generalized linear mixed model examining differences in the proportion of aggregated sperm cells and a linear model examining differences in the number of aggregated sperm cells across six species of *Peromyscus* mice

GLMM: PROPORTION OF CELLS IN AGGREGATE					
Model Term	Beta (SE)	Exp (beta)	95% CI	z	Pr(>|z|)
Intercept	1.52 (0.22)				
eremicus	-2.47 (0.30)	0.07	(0.04, 0.13)	-8.14	**< 0.001**
gossypinus	-3.94 (0.31)	0.02	(0.01, 0.03)	-12.89	**< 0.001**
californicus	-2.58 (0.29)	0.07	(0.04, 0.12)	-9.06	**< 0.001**
leucopus	-3.79 (0.30)	0.02	(0.01, 0.04)	-12.46	**< 0.001**
polionotus	-0.25 (0.29)	0.44	(0.30, 0.58)	-0.85	0.393

LM: NUMBER OF CELLS IN AGGREGATE					
Model Term	Beta (SE)	Exp (beta)	95% CI	t	Pr(>|z|)
Intercept	3.90 (0.26)				
Total Sperm Cells	0.00 (0.00)	0.50	(0.50, 0.50)	5.83	**< 0.001**
eremicus	-1.97 (0.24)	0.12	(0.08, 0.18)	-8.15	**< 0.001**
gossypinus	-2.77 (0.24)	0.06	(0.04, 0.09)	-11.53	**< 0.001**
californicus	-2.19 (0.23)	0.10	(0.07, 0.15)	-9.68	**< 0.001**
leucopus	-2.68 (0.24)	0.06	(0.04, 0.10)	-11.18	**< 0.001**
polionotus	0.04 (0.23)	0.51	(0.40, 0.62)	0.16	0.877

For both models, all rows are being compared with the intercept – *Peromyscus maniculatus*. 95% confidence intervals (CI) were calculated for each effect size.

TABLE 3.

Results from an intra-male analysis comparing motile solitary and aggregated sperm kinematics in low- and high-viscosity conditions for six species of *Peromyscus* mice to test whether sperm aggregates confer kinematic benefits (shaded in gray)

PEROMYSCUS SPECIES	df	KINEMATIC VARIABLE			
		Curvilinear Velocity (μm/sec)	Linearity (VSL/VCL)	Straight-Line Velocity (μm/sec)	Average Path Velocity (μm/sec)
LOW-VISCOSITY CONDITIONS					
californicus	28	$t = -1.0545, p = 0.3007$	$t = -0.56153, p = 0.5789$	$t = -2.2982, \boldsymbol{p = 0.02923}$	$t = - 2.8162, \boldsymbol{p = 0.008805}$
eremicus	20	$t = 1.6225, p = 0.1204$	$t = 0.73517, p = 0.4708$	$t = 0.1312, p = 0.8969$	$t = 1.0464, p = 0.3079$
polionotus	23	$t = 9.4575, \boldsymbol{p = 2.1566e\text{-}09}$	$t = 1.699, p = 0.1028$	$t = 5.8355, \boldsymbol{p = 6.026e\text{-}06}$	$t = 6.0729, \boldsymbol{p = 3.408e\text{-}06}$
maniculatus	17	$t = 2.2482, \boldsymbol{p = 0.03812}$	$t = 0.48075, p = 0.6368$	$t = -2.2335, \boldsymbol{p = 0.03924}$	$t = -1.9206, p = 0.07172$
leucopus	20	$t = 0.23337, p = 0.8178$	$t = -4.8385, \boldsymbol{p = 9.973e\text{-}05}$	$t = -5.5521, \boldsymbol{p = 1.959e\text{-}05}$	$t = -1.838, p = 0.08096$
gossypinus	20	$t = 5.4048, \boldsymbol{p = 2.73e\text{-}05}$	$t = -1.0247, p = 0.3177$	$t = 3.715, \boldsymbol{p = 0.001369}$	$t = 4.479, \boldsymbol{p = 0.0002298}$
HIGH-VISCOSITY CONDITIONS					
californicus	9	$t = -3.7465, \boldsymbol{p = 0.003357}$	$t = 2.682, \boldsymbol{p = 0.02512}$	$t = -2.7743, \boldsymbol{p = 0.0216}$	$t = -4.1106, \boldsymbol{p = 0.002634}$
eremicus	9	$t = -0.71739, p = 0.4913$	$t = 4.2087, \boldsymbol{p = 0.002277}$	$t = 0.62918, p = 0.5449$	$t = 0.56505, p = 0.5858$
polionotus	13	$t = 2.0349, p = 0.06278$	$t = 3.9242, \boldsymbol{p = 0.001745}$	$t = 4.8314, \boldsymbol{p = 0.0003279}$	$t = 4.18, \boldsymbol{p = 0.001079}$
maniculatus	11	$t = -2.9397, \boldsymbol{p = 0.01345}$	$t = 3.9259, \boldsymbol{p = 0.002369}$	$t = -0.30835, p = 0.7636$	$t = -1.0543, p = 0.3114$
leucopus	9	$t = 0.32429, p = 0.7531$	$t = -0.31635, p = 0.759$	$t = -0.34159, p = 0.7405$	$t = 0.95055, p = 0.3667$
gossypinus	10	$t = -0.63835, p = 0.95376$	$t = -0.076451, p = 0.9406$	$t = -0.60402, p = 0.5593$	$t = -0.71643, p = 0.4901$
Statistical results are based on paired-student t-tests.					