1
2
3
4 **Collective sperm movements are shaped by post-copulatory sexual**
5 **selection and phylogenetic history in *Peromyscus* mice**
6
7

8 Kristin A. Hook, W. David Weber, Heidi S. Fisher*
9
10
11
12
13 Department of Biology, University of Maryland, 1200 Biology-Psychology Building,
14 4094 Campus Drive, College Park, MD 20742, U.S.A.
15
16
17 * Heidi S. Fisher
18 Email: hsfisher@umd.edu

19
20
21
22
23 *Keywords:*
24
25 mating systems; sexual selection; sperm competition; sperm conjugation; sperm motility
26
27
28 *Author Contributions:*
29
30 KAH and HSF conceived of the study, designed experiments, and interpreted results;
31 KAH and WWD collected the data, KAH carried out the statistical analyses; all authors
32 wrote the manuscript, gave final approval for publication, and agree to be held
33 accountable for the work presented.
34
35

36 *Funding:*
37
38 This work was supported by a Eunice Kennedy Shriver National Institute of Child Health
39 and Human Development K99/R00 Pathway to Independence Award to HSF
40 [R00HD071972] and a National Science Foundation Postdoctoral Research Fellowship to
41 KAH [1711817].

Abstract

Sperm of some species form motile, coordinated groups as they migrate through the female

reproductive tract to the site of fertilization. This collective motion is predicted to improve sperm

swimming performance and therefore may be beneficial in a competitive context, but limited

evidence supports this theory. Here we examine sperm aggregates across closely-related species of

Peromyscus mice that naturally vary by mating system, and thus sperm competition intensity. We

find that phylogenetic history predicts the likelihood that sperm will aggregate, and that relative

testis size is negatively associated with variation in number of aggregated cells, suggesting that

sperm competition has a stabilizing effect on sperm group size. Moreover, we show that

aggregates are kinematically beneficial for some species but costly for others, and these

differences are largely dependent on the orientation and composition of sperm within the groups.

In addition, when we compared sperm of the two sister-species that aggregate most frequently, we

find that sperm from the species that evolved under intense sperm competition forms aggregates

with more efficient geometry more frequently than sperm from its monogamous congener. These

results are consistent with the prediction that sperm aggregation evolved to improve motility in a

competitive context; however, when monogamy evolved secondarily, relaxed sexual selection

allowed for less efficient strategies to persist. Together, our findings in *Peromyscus* reveal that

collective sperm behavior is likely to evolve rapidly and is shaped by changes in the selective

regime.

Introduction

Sperm cells are one of the most diverse cell types in nature and exhibit striking variation both

within and across taxa (1). In addition to being morphologically diverse, sperm may exhibit

complex, emergent behaviors, including sperm conjugation, which occurs when two or more cells

join together for motility or transport through the female reproductive tract before dissociating

prior to fertilization (1, 2). Although relatively rare, these sperm-sperm interactions have evolved

multiple times across independent lineages of internally fertilizing species (3, 1, 2), yet the

adaptive significance of these gametic interactions remains unclear for many taxa.

One functional hypothesis posits that sperm aggregates may be advantageous if the combined

force generated by multiple flagella enable sperm conjugates to swim faster than single cells (4, 5).

Improved sperm motility can be beneficial if it enables cells to move quickly through hostile

environments within the female tract (6, 7), and in competitive environments in which females

mate with multiple males, as sperm velocity is often correlated with competitive fertilization

success (e.g., 7–9). While some studies that have quantified the motility of sperm aggregates have

found empirical support for this hypothesis (e.g., 4, 10–13), others have not (e.g., 14, 15). For

example, sperm groups swim faster than solitary cells in the Norway rat (*Rattus norvegicus*), yet

house mouse (*Mus musculus*) sperm swim slower as groups under identical experimental

conditions (5). Another non-mutually exclusive hypothesis is that sperm aggregation facilitates

migration through viscous or viscoelastic secretions of the female reproductive tract (16, 17),

including cervical mucus (18), which has received some empirical support (e.g., 4, 12, 16), but see

(15). In the grey short-tailed opossum (*Monodelphis domestica*), for example, sperm pairs swim

with greater motility than single sperm in viscous fluids; however, these sperm pairs were

artificially induced to uncouple (16) and therefore the comparison may not be biologically

informative (1). Additionally, sperm aggregation may protect sensitive regions of the sperm, such

as the acrosome, from damage and preserve sperm functionality during passage through the male

or female reproductive tracts (19–22), or enhance egg penetration during fertilization (23) but see

(24). While it is often assumed that coordinated sperm movements are adaptive, inconsistent

findings and a wide diversity of naturally variable sperm behavior have limited our understanding

of the functional advantages of sperm aggregation.

Multiple independent origins of sperm conjugation suggest that the functional consequences,

as well as mechanisms that regulate these cellular interactions, are likely to vary throughout nature

(reviewed in 2). Indeed, the formation of sperm aggregates and the number of grouped cells varies

widely across taxa (7). In mammals alone, sperm of some species assemble during epididymal

transport and are molecularly "glued" to one another as bi-flagellate pairs in grey short-tailed

opossums (25), as bundles of roughly 100 cells in monotremes (26), or as organized rouleaux of

five or more cells in guinea pigs (*Cavia porcellus*, 24). Conversely, mammalian sperm may also

assemble after ejaculation to form variably-sized groups. For instance, sperm may form temporary

clusters of up to sixteen cells in bulls (*Bos Taurus*, 15) or more fixed groups of up to 30 cells in

house mice (5), 50 cells in the Norway rat (5), or thousands of cells in the wood mouse (*Apodemus*

sylvaticus, 4) whereby the hook-shaped heads interlock or attach to the flagella of other sperm. For

these latter species in which sperm groups form after ejaculation, both single and aggregated

sperm typically co-occur, thus allowing for direct comparisons between collective and solitary

sperm movements within an ejaculate while controlling for within-male variability.

Closely-related species in the rodent genus, *Peromyscus,* produce sperm that naturally vary in

their collective behavior. Sperm of some species assemble temporary groups after ejaculation by

adhering to one another at their head region (27) and disassemble prior to fertilization (10). In *P.*

maniculatus, sperm selectively group with the most closely-related cells to form motile groups of

up to 30 cells (10), but there is a non-monotonic association between group size and swimming

velocity, indicating that some groups are faster than single cells but others are not (28).

Conversely, sperm produced by their sister-species, *P. polionotus*, also form aggregates but do so

indiscriminately with related and unrelated cells (10), are less likely to be optimally-sized, and

overall move in a less linear trajectory (28). Intriguingly, sperm competition is predicted to be

more intense in *P. maniculatus* than in *P. polionotus* due to their different mating systems. In *P.*

maniculatus, both sexes mate with multiple partners, often in overlapping series just minutes apart

(29), and females frequently carry multiple-paternity litters in the wild (30), whereas both

behavioral (31) and genetic data (32) indicate that *P. polionotus* is strictly monogamous. Evidence

suggests that monogamy has evolved at least twice within the *Peromyscus* lineage (reviewed in 33;

34), thus enabling us to investigate if post-copulatory sexual selection, driven by female mating

behavior, has shaped the evolution of sperm aggregation (4) more broadly across the *Peromyscus*

lineage. In this study, we quantify intra- and inter-specific differences in the size and performance

of sperm aggregates under consistent, controlled conditions to examine the evolution of collective

sperm behavior and empirically test whether sperm aggregation improves swimming performance.

Results

We investigated the frequency of sperm aggregation in each species and found significant

differences in the proportion of aggregated cells among species (binomial GLMM: $n = 134$, $P <$

0.001; Table 1, 2; Figure 1), with more variance across species than within species (variance

across species = 0.10; variance within each species < 0.10). Specifically, sperm from *P.*

maniculatus and *P. polionotus* similarly aggregate the most, whereas *P. leucopus* and *P.*

gossypinus sperm similarly aggregate the least among all the species. Pairwise comparisons

adjusted for multiple comparisons using LSmeans revealed that species within these pairs do not

significantly differ from one another, whereas all other pairwise species comparisons do (Table 1,

2). The coefficient of variation (CV) for the proportion of cells aggregated within each species are

as follows: 68.8% for *P. californicus*, 50.6% for *P. eremicus*, 14.4% for *P. polionotus*, 10.8% for

P. maniculatus, 125.7% for *P. leucopus*, and 60.5% for *P. gossypinus*. Controlling for phylogenic

relationships and body mass, we found no effect of testis size ($F_{2,3} = 2.606$, $P = 0.15816$) or the

within-species CV ($F_{2,3} = 0.3603$, $P = 0.5604$) on the proportion of aggregated cells.

Moreover, we found significant differences in the mean number of cells aggregated among species

(LM: $F_{6,126} = 56.37$, $P < 0.001$; Table 1, 2; Figure 2A), with more variance observed across species

($s^2 = 1.96$) than within species ($s^2 < 1.00$, except for *P. polionotus* [$s^2 = 2.15$]). Post-hoc pairwise

comparisons revealed that both *P. maniculatus* and *P. polionotus* produce the largest sperm

aggregates ($P < 0.05$ for all pairwise comparisons), whereas *P. gossypinus*, *P. leucopus*, and *P.*

californicus produce the smallest aggregates ($P < 0.05$ for pairwise comparisons), the latter of

which produces sperm cell aggregates that are statistically similar in size to those produced by *P.*

eremicus ($P = 0.9203$; Figure 2A). Controlling for phylogenic relationships and body mass, we

found a significant effect of testis weight on the within-species CV of aggregate size ($F_{2,3} = 8.398$,

$P = 0.02655$, Figure 2B), but found less of an effect on the direct measure of aggregate size ($F_{2,3} =$

3.058, $P = 0.1522$). The CV for the number of cells aggregated within each species were as

follows: 26.5% for *P. californicus*, 19.2% for *P. eremicus*, 27.5% for *P. polionotus*, 18.4% for *P.*

maniculatus, 13.3% for *P. leucopus*, and 7.2% for *P. gossypinus*.

By comparing sperm aggregate composition and orientation within the species that produce

the largest and most frequent sperm aggregates, we found that there are significantly more sperm

aggregates in *P. maniculatus* in which all sperm are aligned in a head-to-flagella orientation

(99.1%, 731/738) than in *P. polionotus* (87.5%, 720/826; GLM: $n = 42$, $P = 3.92e-12$) and that

these aggregates have significantly faster speeds (VCL) compared to aggregates with unaligned

cell orientations in both species (Figure 3; paired t-tests: *P. maniculatus* $t = 5.9627$, $df = 4$, $P =$

0.003972; *P. polionotus* $t = 11.247$, $df = 11$, $P = 2.257e-07$). We also found that there are

significantly fewer sperm aggregates in *P. maniculatus* with immotile, stuck, or morphologically

abnormal cells (8.1%, 64/795) than in *P. polionotus* (11.5%, 94/814; binomial GLM: $n = 40$, $P =$

0.00503). Importantly, aggregates with these defects had significantly lower speeds (VCL)

compared to aggregates without in both species (*P. maniculatus* $VCL_{aligned} = 179.60 \pm 6.88\,\mu$m/sec,

$VCL_{defective} = 132.47 \pm 6.18\,\mu$m/sec, paired t-test: $t = 20.627$, $df = 14$, $P = 7.075e-12$; *P. polionotus*

$VCL_{aligned} = 135.48 \pm 3.51\,\mu$m/sec, $VCL_{defective} = 115.62 \pm 5.22\,\mu$m/sec, paired t-test: $t = 16.312$, df

$= 20$, $P = 5.079e-13$).

When comparing the proportion of motile and progressively motile aggregates across species,

our pairwise comparisons revealed that *P. eremicus* produced a significantly smaller proportion of

motile aggregates than all other species ($P < 0.05$ for all *P. eremicus* pairwise comparisons; $P >$

0.05 for all other pairwise comparisons). Fitted values of the proportion of motile aggregates using

LSmeans were 0.77 ± 0.04 for *P. eremicus*, 0.92 ± 0.02 for *P. polionotus*, 0.91 ± 0.02 for *P.*

gossypinus, 0.95 ± 0.01 for *P. maniculatus*, 0.93 ± 0.02 for *P. leucopus*, and 0.91 ± 0.02 for *P.*

californicus. Moreover, post-hoc comparisons revealed that *P. eremicus, P. polionotus,* and *P.*

gossypinus all had the smallest proportions of progressively motile aggregates ($P < 0.05$ for all

pairwise comparisons; fitted values using LSmeans were 0.64 ± 0.06, 0.65 ± 0.05, and 0.75 ± 0.05,

respectively), the latter species of which did not significantly differ from *P. californicus* ($P =$

0.1981; 0.87 ± 0.02 for LSmeans fitted values). Conversely, *P. maniculatus* and *P. leucopus* had

the largest proportions of progressively motile aggregates ($P = 0.3417$; fitted values using

LSmeans were 0.94 ± 0.02 and 0.91 ± 0.02, respectively), the latter of which did not differ from *P.*

californicus ($P = 0.9109$).

Overall, we found species-specific differences in the effect of sperm aggregation on motility,

regardless of environmental complexity (Table 3, Figure 4). In low-viscosity medium, we found

that sperm aggregates have a significantly greater VCL in *P. maniculatus*, VSL in *P. maniculatus*,

P. leucopus, and *P. californicus*, LIN in *P. leucopus*, and VAP in *P. californicus* compared to

single cells. Conversely, sperm aggregates had a significantly lower VCL, VSL, and VAP velocity

in *P. polionotus* and *P. gossypinus* than single sperm in the low-viscosity medium (Figure 4). In

the high-viscosity medium, we found that sperm aggregates have a significantly greater VCL in *P.*

maniculatus and *P. californicus* and a higher VSL and VAP in *P. californicus* (Figure 4) compared

to single sperm. Conversely, sperm aggregates in the high viscosity medium had a significantly

lower LIN in *P. californicus*, *P. eremicus*, *P. polionotus*, and *P. maniculatus* as well as a reduced

VSL and VAP in *P. polionotus* than single cells (Figure 4).

Discussion

While it is known that collective sperm behaviors have evolved independently in a number of

taxa (2), it remains unclear how sperm aggregation evolves among closely related species. Our

comparative study reveals that sperm aggregating behaviors vary across mice in the genus

Peromyscus. We observed an effect of phylogenetic history on the frequency of sperm

aggregation, indicating that collective sperm behavior likely evolved prior to the divergence of

present-day species. Additionally, we find a negative association between relative testis weight, a

robust proxy for intensity of sperm competition in rodents (35), and the coefficient of variation for

the number of cells in aggregate, suggesting that sexual selection has a stabilizing effect on sperm

aggregate size. We then compared the motility of single sperm and sperm aggregates across all

species under low- and high-viscosity conditions, and show that aggregation is kinematically

beneficial for some species yet costly for others, regardless of environmental complexity. When

examining sperm from the species that aggregate the most profoundly, we find that the formation

of the aggregates and the orientation of the cells within the group are critical to kinematics.

Moreover, we observe more aggregates with efficient geometry in the species that has evolved

under strong post-copulatory sexual selection compared to its monogamous sister-species. These

findings support the prediction that sperm aggregation evolved in *Peromyscus* to improve motility

in a competitive context but reveal that relaxed selection may have enabled less efficient strategies

to persist, thereby generating diversity in collective sperm behaviors within these closely-related

species.

Our results reveal distinct species-specific differences across *Peromyscus* mice in the

frequency of sperm aggregation and the average size of these cellular groups. Multiple

Peromyscus species produce sperm that aggregate more extensively than other studied muroid

rodents (36), with the notable exception of the wood mouse (4). In general, the proportion of

sperm cells that aggregate is most similar within each sister-species pair. Specifically, *P.*

maniculatus and *P. polionotus* produce sperm that aggregate the most (>80% of sperm), but the

pair they are most closely-related to, *P. gossypinus* and *P. leucopus*, are the species least likely to

produce sperm that aggregate (<10% of sperm); the most distantly-related species pair we

assessed, *P. californicus* and *P. eremicus*, both produce sperm with a moderate propensity to

aggregate (~25% of sperm). These findings support one of two possibilities for the evolution of

sperm aggregates within these species: (a) a genus-wide ancestral trait of moderate sperm

aggregation with subsequent diversification leading to an increase in *P. maniculatus* and *P.*

polionotus and a decrease in *P. gossypinus* and *P. leucopus*, or (b) the independent evolution of

aggregation in the ancestors of the *P. californicus*-species pair and the *P. maniculatus*-species pair.

Such complex evolutionary histories with losses and recurrences of sperm conjugation, and

subsequent species divergence, have also been demonstrated in diving beetles (Dytiscidae, 37),

consistent with the evolutionary lability that we observe. Our experimental results may explain

selection against sperm aggregation if forming groups reduces sperm swimming performance,

which we find in at least one species, *P. gossypinus*. However, we found that sperm produced by

their sister-species, *P. leucopus*, also rarely aggregate, despite our observation that these rare

collective groups have a greater average velocity and are more linear than single cells.

Interestingly, we observed that these two promiscuous species that rarely aggregate, *P. gossypinus*

and *P. leucopus*, have the largest relative testes of the species studied, suggesting that they may

have evolved increased sperm production to improve competitive fertilization success (sensu 38),

rather than adaptations that influence motility (39). Together these results suggest different

strategies employed by divergent species across the *Peromyscus* genus in response to sperm

competition.

In contrast to our results on the frequency of sperm aggregation within species, we found that

the average size of sperm groups does not align as closely with phylogenetic relationships. In

addition, we found that species with relatively larger testes, which is positively associated with

increased sperm competition (35), exhibit less variation (CV) in aggregate size. This result

supports the prediction that relaxed sperm competition allows for greater intermale variation to

persist in a population (40) and suggests that this post-copulatory sexual selection may be

stabilizing sperm aggregate size for a species-specific 'optimum' (28). Similarly, other studies

have shown that the strength of sexual selection regulates variance in sperm morphology across

taxa and at multiple levels of organization, including within- and between-males (41–43) as well

as within- and between-species (40, 44). A study on sperm bundles across ten *Carabus* ground

beetles also found intense selection on bundle size, which are dimorphic and either small or large;

the large, but not small, sperm bundles are positively correlated with measures of sperm

competition risk, including copulatory piece length and mate guarding, suggesting that diversity of

large sperm bundles is associated with sexual selection (45). In contrast to these findings that

competition drives sperm-sperm interactions, a study on the evolution of such sperm traits in

diving beetles found that variation in sperm conjugation is more associated with female

reproductive tract architecture (37, 46). Therefore, while our results suggest that stabilizing

selection on sperm aggregate size is associated with an increase in sperm competition given the

correlation with relative testis weight, mechanisms of female control (47) may also play an

important evolutionary role.

We compared the motility of single and aggregated sperm sampled from the same male to test

whether sperm aggregates swim faster or more efficiently than single sperm, which is predicted if

the combined force of multiple flagella enhances their motility (4). We found improved kinematic

measures in half of the species studied (*P. maniculatus, P. californicus*, and *P. leucopus*), thus

supporting this functional hypothesis; however, in several species we found aggregation had some

negative (*P. polionotus* and *P. gossypinus*) or no (*P. eremicus*) impact on motility relative to single

sperm cells. While we found support for a theoretical prediction that sperm aggregates achieve

greater straight-line velocity because they move in a more linear path of travel rather than at a

faster speed (i.e., curvilinear velocity, 28) in two of our six focal species (*P. californicus* and *P.*

leucopus), we did not find this kinematic benefit for sperm aggregation in all *Peromyscus* species.

These results corroborate other studies in more disparate taxonomic groups that have quantified

sperm aggregation motility and found inconsistent results. For example, sperm trains exhibit

greater swimming progressive motility in the wood mouse (4), and greater velocity than individual

sperm in the Norway rat, but not the house mouse (5). In invertebrates, the swimming velocity of

fishfly sperm increases with number of sperm in a bundle (12), but in a marine snail, there is no

differences in swimming speed between paired and single sperm (14). One possible explanation

for these differences across taxa is that cell orientation within an aggregate is critical for its

collective motility. Sperm cells are predicted to be faster if they generate increased force with

proportionally less drag (see 2 and references therein); such effects may be true for sperm

aggregations as well in which cells conjoin head-to-tail, thereby increasing the length of the

collective unit, or in which flagella within the group beat synchronously (2). Our results support

that even sperm aggregates that conjoin head-to-head and are thus wider, as they are in

Peromyscus, can offer a motility advantage.

Our results suggest that relaxed sexual selection may allow the persistence of less optimal

strategies based on the quantitative and qualitative differences we observed among sperm

aggregates of different species. The most interesting motility results are those of the sister-species

pair with divergent mating systems that both form the most frequent and largest sperm aggregates.

In *P. maniculatus*, a promiscuous species, sperm aggregates exhibit greater straight-line and

curvilinear velocity compared to single cells, but the opposite was true for its monogamous

congener, *P. polionotus*. We find that these kinematic differences are associated with differences

in aggregate geometry; when sperm heads and flagella are not oriented in the same direction, the

cells within an aggregate exert opposing forces on one another, thereby reducing the overall

motility of the group (27, 28). Indeed, we found that sperm from the monogamous *P. polionotus*

males are less likely to form aggregates with all sperm aligned and more likely to include immotile

or morphologically abnormal sperm, consequently resulting in slower aggregates than those of *P.*

maniculatus (27). This finding is consistent with previous reports that *P. polionotus* sperm tend to

form optimal-sized aggregates less often than in *P. maniculatus* (28). Together, these observations

further support the hypothesis that sperm aggregation evolved prior to the divergence of the

species pair (10), and when monogamy evolved secondarily in *P. polionotus* (34, 48), relaxed

sexual selection allowed for the persistence of less motile sperm traits. In line with this prediction,

we observed the smallest proportion of motile and progressively motile sperm aggregates in

another monogamous species, *P. eremicus,* but the largest proportion of progressively motile

aggregates in two promiscuous species, *P. maniculatus* and *P. leucopus*. Similar results have been

reported in house mice, in which males evolving under intense sperm competition produced a

greater proportion of motile sperm, compared to males from lineages subject to relaxed selection

(49). Together these findings support that the motility benefits conferred by sperm aggregation are

associated with variation in post-copulatory sexual selection.

When we compared single and aggregated sperm in a viscous environment to test if

aggregation improves motility in more complex fluids, similar to the secretions or cervical mucus

(18) of the female reproductive tract (17, 50), we found that collective motion is beneficial for

some species but costly for others. Compared to single sperm, aggregates swim with greater

curvilinear velocity in *P. maniculatus* and *P. californicus*, and greater straight-line and average

path velocities in *P. californicus,* but aggregates were less linear than single cells in all species

except for *P. leucopus* and *P. gossypinus*. While other studies have found kinematic benefits for

sperm conjugates compared to single cells in higher viscosities in the gray short-tailed opossum

(16), the wood mouse (4), and the fishfly (12), a study in bulls found that sperm were slower,

exhibited less organized swimming patterns, and were less likely to cluster in viscous fluids (15),

both of which are consistent with our analysis of *Peromyscus* aggregates. Ultimately, the benefit of

sperm aggregation depends on the relative importance of each kinematic parameter during sperm

migration *in vivo*. Although beyond the scope of this study, we predict that improved linearity

afforded by collective motion may help to direct the sperm through the dynamic fluids of the

female reproductive tract (51) and that increased velocity will reduce the time it takes for the

sperm to arrive at the fertilization site.

In conclusion, our study highlights the diversity of sperm aggregation within a single

taxonomic lineage and how selection has shaped the formation and performance of these cellular

groups. We show that both evolutionary history and varying levels of post-copulatory sexual

selection influence the frequency and size of sperm groups. Moreover, we find that sperm

aggregation can improve sperm motility in both simple and complex fluids, but this is not

consistent across all species. Theoretical predictions (27, 28) and emerging empirical evidence

suggests that motility benefits may only be realized if cells maintain optimal alignment within the

groups and, if achieved, may provide these sperm with a competitive advantage in the female

reproductive tract (46). Future work investigating sperm aggregates *in vivo* (e.g., 51, 52) will shed

light on the co-evolution of these unique gametic behaviors and the enormously variable and

dynamic female reproductive tracts through which sperm must navigate.

Materials and Methods

(a) Sperm collection

We obtained captive *Peromyscus maniculatus bairdii, P. polionotus subgriseus, P. leucopus,*

P. eremicus, and *P. californicus insignis* from the Peromyscus Genetic Stock Center at the

University of South Carolina, and *P. gossypinus gossypinus* from Dr. Hopi Hoekstra at Harvard

University and housed them in same-sex cages at 22ºC on a 16L:8D cycle in accordance with

guidelines established by the Institutional Animal Care and Use Committee at the University of

Maryland in College Park (protocol R-Jul-18-38). We sought samples from all available captive

Peromyscus species and avoided wild-caught specimens to control for variation due to life

experience. We obtained sperm samples from sexually mature males and accounted for relatedness

among the focal males by assigning siblings a unique 'Family' ID. We euthanized males via

isoflurane overdose and cervical dislocation, then weighed each male and both testes (Mettler

Toledo, Switzerland). Next, we removed a single caudal epididymis, made several small incisions

in the tissue, and submersed it in sperm medium (Modified Sperm Washing Medium, Irvine

Scientific, USA) that was pre-warmed at 37ºC; to reduce differences in sperm density despite

natural variation in epididymal sizes, we varied the volume (50µl - 1000µl) based on tissue size

and accounted for these differences when estimating final sperm counts for each male. To collect

sperm, we agitated the tissue at 300rpm (ThermoMixer F1.5, Eppendorf, Germany) at 37ºC for ten

minutes, inverting the tube at the five- and ten-minute mark, then incubated the tissue undisturbed

for two minutes. Using pipette tips cut to create a wider opening, we collected live sperm cells for

analysis from just below the meniscus of the solution to enrich for the most motile sperm (53).

Next we estimated sperm density using a computer-assisted sperm analysis (CASA) system (Ceros

II Animal, Hamilton Thorne, USA) and verified with a Neubauer-improved hemocytometer

(Marienfeld, Germany) then diluted samples with pre-warmed medium to reach a standard

concentration of 300-400 cells summed across the five 5-second videos at 100X magnification for

cell tracking optimality and efficiency.

(b) Live sperm analysis

To conduct live sperm observations, we gently reverse pipetted 4μl of the sperm solution into

12μl of pre-warmed medium on a plastic slide within a 9mm x 0.12mm imaging spacer (Grace

Bio-Labs, USA) and covered by a plastic cover slip. This set-up served as a control and represents

a 'low-viscosity' environment. To test sperm motility in a 'high-viscosity' environment, we

followed the same procedures except that we mixed 4μl of sperm solution with 12μl of pre-

warmed medium enriched with methylcellulose (Sigma Aldrich M 7140; 15cP, 2% in water; 54).

We then recorded 5-second videos at 60 frames/sec on the CASA system, capturing at least five

videos per male but recorded additional videos for samples with lower sperm density ($n = 57$) to

ensure an adequate number of observed cells per male. Videos (see example, Movie S1) were

recorded at 59±16 minutes post-harvest from the epididymal tissue, dependent on dilutions The

number of videos, tracks, and cells analyzed are reported in Table 1.

We characterized sperm aggregation by scoring CASA videos using direct observations

because the system tracks particles, and thus each track may represent a single cell or an

aggregate. We counted the number of cells represented by each track on at least three different

frames/track. From these data, we calculated the proportion of cells that aggregated for each male

by dividing the total number of aggregates by the total number of motile cells across all tracks

(55). Then we calculated the mean number of cells in aggregate (i.e., aggregate size) by dividing

the sum of cells in aggregate by the sum of aggregates, both across all tracks for each male and

across all males for each species. Finally, we calculated the coefficient of variation (CV) for both

the proportion and number of cells that aggregated within each species using the following

formula: (standard deviation/mean) x 100. In addition, to further characterize differences in sperm

aggregation for the two species whose sperm were observed to aggregate the most extensively, *P.*

maniculatus and *P. polionotus*, we qualitatively scored the composition and orientation of cells

within the sperm aggregates. For males within these species, we calculated the proportion of

aggregates that were: 'aligned' in which all sperm adhered to one another in a head-to-flagella

orientation and included no immotile, morphologically abnormal or damaged cells, 'defective',

which included one or more sperm that was abnormal, immotile, or stuck to the slide, and

'opposed' if all sperm are normal and motile, but were not oriented in the same direction, the latter

of which included star-shaped aggregates (28).

We recorded the following metrics for each recorded track (i.e., single sperm and aggregates):

straight-line velocity (VSL; calculated using the straight-line distance between the first and last

detected positions of the sperm head, divided by the time taken to swim the track; also known as

average velocity), *curvilinear velocity* (VCL; calculated using the summed distance between the

sperm head positions in each frame divided by the time taken to travel the track; also known as

speed), *average path velocity* (VAP; the time-averaged velocity of the sperm head along its

average path), and *linearity* (LIN; the ratio of VSL to VCL to measure the straightness of the

trajectory; 56). We calculated the mean of each kinematic parameter for both single cells and

sperm aggregates separately for each of three populations of sperm cells: all cells, motile cells

(i.e., devoid of visually inspected tracks in which cells were unmoving, stuck, or featured an

obvious morphological abnormality such as a kinked midpiece), and progressively motile cells

(i.e., motile cells with a VSL $\geq$25μm/sec). We used these data to calculate the proportion of motile

aggregates by dividing the sum of motile aggregates by the total number of sperm aggregates, and

the proportion of progressively motile aggregates by dividing the sum of progressive aggregates

by the total number of sperm aggregates for each male (55). For our kinematic analyses, we

focused on the motile sperm dataset to eliminate artifacts from damaged or dead cells, and the total

sperm population to quantify frequency of aggregation and aggregate size in an effort to capture

natural aggregation rates (results from other cell populations are reported in Table S1, Figure S1).

(c) Statistical analyses

We performed all statistical analyses using R version 3.4.2 (57) and visually inspected

diagnostic plots (qqplots and plots of the distribution of the residuals against fitted values) to

validate model normality. Only the best fitting models are reported here. We created all figures

using the 'ggplot2' package with R (58). One *P. californicus* male was excluded from the

aggregate analysis dataset because their measurements represented clear outliers. All means are

presented ± 1 standard error.

To compare species differences in the proportion of aggregated cells, we used the mean values

for each male and a generalized linear mixed model (GLMM) using the glmer function from the

"lme4" R package and a logit link function (59). The binomial response was the number of sperm

cells in aggregate, and the total number of sperm cells was the binomial denominator. In the initial

statistical model, we observed the residual deviance to be larger than the residual degrees of

freedom, which is an indication of overdispersion (55). We thus used an observation-level random

effect (OLRE) as a random factor in all subsequent analyses to control for overdispersion (60). We

considered family ID as a random factor in the initial model and both random factors were then

used in bivariate analyses for predictors of interest that could potentially explain differences in the

proportion of aggregated cells. These predictors included male age, pairing status, the timing of

video recordings relative to harvest of the epididymal tissue, and the number of videos recorded.

Only predictors that had a *p*-value at or below 0.20 were considered for the final model. We

further screened these predictors for collinearity with other significant predictors using linear

models and removed collinear predictors, so that only the one with greater relative significance

was included in the final GLMM. The remaining model included pairing status and species as

fixed factors. We calculated the variance inflation factor (VIF) values and found evidence of

collinearity due to two VIF values above the recommended threshold of three (61). Thus, we

removed pairing status as a fixed factor and family ID as a random factor, leaving only species as

a fixed factor within our final model. Post-hoc pairwise comparisons were performed using Tukey

HSD adjustments for multiple comparisons from the "LSmeans" R package (62).

To compare species differences in the number of aggregated cells, we used the mean values

for each male and initially used a linear mixed model (LMM) using the lmer function from the

"lme4" R package, but eventually reverted to using a linear model (LM) because the family ID

random factor did not significantly contribute to the residual variability in the response variable.

Predictors that were considered for the initial LMM included male age, pairing status, the timing

of video recordings relative to harvest of the epididymal tissue, total sperm cells, number of

recorded videos, an interaction between the latter two variables, and the ratio of total sperm cells

to the number of videos recorded. We considered predictors with p-value<0.20 for the final model,

but first screened each for collinearity with other significant predictors using a linear model and

removed whenever collinearity was present so that only the one with the greater relative

significance was included in the final LM. We dropped non-significant explanatory variables one

at a time based on model comparisons using an analyses of variance test to determine the minimal

adequate model, but were unable to meet the normality assumptions for this model. We also

assessed species differences in the proportion of aggregates that were motile or progressively

motile from the total population of aggregates using binomial GLMM. The binomial response was

the number of motile sperm aggregates, and the total number of sperm aggregates was the

binomial denominator. Our final model contained both an OLRE due to detected overdispersion

and family ID as random factors.

To determine if sperm aggregates have motility or force benefits over single cells, we initially

performed a principal component analysis (PCA) using three related swimming performance

measures (VSL, VCL, and VAP) to reduce dimensionality and obtain a composite measure for

motile solitary and aggregated sperm cells in both low- and high-viscosity media. Because we

found that species distributions overlapped for this composite measure for both single sperm and

aggregates in low- and high-viscosity media (Figures S2 and S3), we focused on individual

kinematic parameters (VSL, VCL, VAP, LIN) within each species separately. Using our dataset of

mean values per male, we used a paired student's t-test to compare each kinematic parameter

between solitary sperm cells and sperm aggregates within males for each separate species to

determine if sperm aggregates have motility benefit over single cells. To determine if aggregates

have a force benefit over single sperm, we conducted these same analyses in high-viscosity media.

We then combined these two datasets in low- and high- viscosity media and compared sperm

aggregate kinematics in both low- and high- viscosity media at the intra-male level using a paired

student's t-test within each species.

To assess the structure of *P. maniculatus* and *P. polionotus* aggregates, we used generalized

linear models (GLM) to compare the proportions of 'defective and 'opposed' aggregates to

'aligned' aggregates. For the composition GLM, the binomial response was the number of

'defective' sperm aggregates, and the total number of sperm aggregates that were either 'defective'

or 'aligned' was the binomial denominator. We used a paired student's t-test to compare the VCL

of these aggregates within males. For the orientation GLM, the binomial response was the number

of 'opposed' sperm aggregates, and the total number of sperm aggregates that were either

'opposed' or 'aligned' was the binomial denominator. We used a paired student's t-test to compare

the VCL of these aggregates within males.

Finally, to account for variation in phylogenetic relationships among of the species used in this

study, we adopted a phylogenetic generalized least squares approach (63, 64) using the "caper"

(65) and "APE" (66) packages in R and using an ultra-metric phylogenetic tree of *Peromyscus*

(provided by Dr. Roy Neal Platt II, Texas Biomedical Research Institute), based on sequence

variation in the mitochondrial gene, cytochrome B. The species' relationships within this tree

matched those from other previously established phylogenies of *Peromyscus* (34, 67). We used

this phylogeny as a covariate in regression analyses to investigate the effect of relative testis

weight on sperm aggregation, including the proportion of aggregated cells and aggregate size, and

the within-species CV for each of these parameters. Finally, to control for differences in male

body when examining testis mass, we included body mass as a separate fixed factor within our

analyses, a method better suited to estimating relative testis weight is size rather than using the

ratio of testis to body mass or residuals (68, 69).

Acknowledgments:

We are grateful to Erica Glasper for providing *P. californicus* males and Hopi Hoekstra for

providing *P. gossypinus* males. Thanks to Harrison Arsis, Madeline-Sophie Dang, Catherine Liu,

and Audrey Mvemba for their assistance with video analysis and to Philip Johnson, Danielle

Adams, Sam Church, and Shelby Wilson for statistical advice. We thank Neal Platt for providing

the ultra-metric *Peromyscus* tree for use in our statistical analysis.

References:

1. S. Pitnick, D. J. Hosken, T. R. Birkhead, "Sperm Morphological Diversity" in Sperm Biology,

An Evolutionary Perspective, T. R. Birkhead, D. J. Hosken, S. Pitnick, Eds. (Academic Press,

2009), pp. 69–149.

2. D. M. Higginson, S. Pitnick, Evolution of intra-ejaculate sperm interactions: do sperm

cooperate? *Biological Reviews* 86, 249–270 (2011).

3. S. Immler, Sperm competition and sperm cooperation: the potential role of diploid and haploid

expression. *Reproduction* 135, 275–283 (2008).

4. H. D. Moore, K. Dvoráková, N. Jenkins, W. Breed, Exceptional sperm cooperation in the

wood mouse. *Nature* 418, 174–177 (2002).

5. S. Immler, H. D. Moore, W. Breed, T. Birkhead, By hook or by crook? Morphometry,

competition and cooperation in rodent sperm. *PloS one* 2, e170 (2007).

6.   T. R. Birkhead, A. P. Møller, W. J. Sutherland, Why do females make it so difficult for males

to fertilize their eggs? *Journal of Theoretical Biology* 161, 51–60 (1993).

7.   T. Pizzari, K. R. Foster, Sperm sociality: cooperation, altruism, and spite. *PLoS Biology* 6

(2008).

8.   T. R. Birkhead, J. G. Martinez, T. Burke, D. P. Froman, Sperm mobility determines the

outcome of sperm competition in the domestic fowl. *Proceedings of the Royal Society of*

London. Series B: Biological Sciences 266, 1759–1764 (1999).

9.   M. J. G. Gage, C. P. Macfarlane, S. Yeates, R. G. Ward, J. B. Searle, G. A. Parker,

Spermatozoal traits and sperm competition in Atlantic salmon: relative sperm velocity is the

primary determinant of fertilization success. *Current Biology* 14, 44–47 (2004).

10.   H. S. Fisher, H. E. Hoekstra, Competition drives cooperation among closely related sperm of

deer mice. *Nature* 463, 801 (2010).

11.   M. Pearcy, N. Delescaille, P. Lybaert, S. Aron, Team swimming in ant spermatozoa. *Biology*

Letters 10, 20140308 (2014).

12.   F. Hayashi, Sperm co-operation in the fishfly, *Parachauliodes japonicus*. *Functional Ecology*

12, 347–350 (1998).

13.   D. M. Woolley, R. F. Crockett, W. D. I. Groom, S. G. Revell, A study of synchronisation

between the flagella of bull spermatozoa, with related observations. *Journal of Experimental*

Biology 212, 2215–2223 (2009).

14.   S. Ishijima, S. A. Ishijima, B. A. Afzelius, Movement of Turritella spermatozoa: Direction of

propagation and chirality of flagellar bends. *Cell Motility* 44, 85–95 (1999).

15.   C. K. Tung, C. Lin, B. Harvey, A.G. Fiore, F. Ardon, M. Wu, S. S. Suarez, Fluid

viscoelasticity promotes collective swimming of sperm. *Scientific Reports* 7, 1–9 (2017).

16.   H. D. M. Moore, D. A. Taggart, Sperm pairing in the opossum increases the efficiency of

sperm movement in a viscous environment. *Biol Reprod* 52, 947–953 (1995).

17. S. S. Suarez, Mammalian sperm interactions with the female reproductive tract. *Cell and*

Tissue Research 363, 185–194 (2016).

18. F. W. Hanson, J. W. Overstreet, The interaction of human spermatozoa with cervical mucus in

vivo. *American Journal of Obstetrics and Gynecology* 140, 173–178 (1981).

19. D. M. Phillips, Ultrastructure of spermatozoa of the wooly opossum *Caluromys philander*.

Journal of Ultrastructure Research 33, 381–397 (1970).

20. J. M. Bedford, J. C. Rodger, W. G. Breed, Why so many mammalian spermatozoa - a clue

from marsupials? *Proceedings of the Royal Society of London B: Biological Sciences* 221,

221–233 (1984).

21. D. M. Higginson, K. R. H. Henn, Effects of sperm conjugation and dissociation on sperm

viability in vitro. *PloS one* 7, e34190 (2012).

22. K. S. K. Tung, A. Okada, R. Anagimachi, Sperm autoantigens and fertilization. I. effects of

antisperm autoantibodies on rouleaux formation, viability, and acrosome reaction of guinea

pig spermatozoa. *Biology of Reproduction* 23, 877–886 (1980).

23. J. B. Mackie, M. H. Walker, A study of the conjugate sperm of the dytiscid water beetles

Dytiscus marginalis and *Colymbetes fuscus. Cell Tissue Res.* 148, 505–519 (1974).

24. S. M. McGlinn, B. A. Shepherd, J. Martan, Rouleau Formation and fertility of spermatozoa in

guinea pigs treated with cyproterone acetate. *Archives of Andrology* 2, 197–202 (1979).

25. D. A. Taggart, J. L. Johnson, H. P. O'Brien, H. D. M. Moore, Why do spermatozoa of the

American marsupial form pairs? A clue from the analysis of sperm-pairing in the epididymis

of the grey short-tailed opossum, *Monodelphis domestica. Anatomical Record* 236, 465–478

(1993).

26. B. Nixon, H. Ecroyd, J. L. Dacheux, F. Dacheux, V. Labas, S. D. Johnston, R. C. Jones,

Formation and dissociation of sperm bundles in monotremes. *Biology of Reproduction* 95

(2016).

27. D. J. G. Pearce, L. A. Hoogerbrugge, K. A. Hook, H. S. Fisher, L. Giomi, Cellular geometry

controls the efficiency of motile sperm aggregates. *Journal of The Royal Society Interface* 15,

20180702 (2018).

28. H. S. Fisher, L. Giomi, H. E. Hoekstra, L. Mahadevan, The dynamics of sperm cooperation in

a competitive environment. *Proceedings of the Royal Society of London B: Biological*

Sciences 281, 20140296 (2014).

29. D. A. Dewsbury, Interactions between males and their sperm during multi-male copulatory

episodes of deer mice (*Peromyscus maniculatus*). *Animal Behaviour* 33, 1266–1274 (1985).

30. D. A. Birdsall, D. Nash, Occurrence of successful multiple insemination of females in natural

populations of deer mice (*Peromyscus maniculatus*). *Evolution* 27, 106–110 (1973).

31. D. A. Dewsbury, An exercise in the prediction of monogamy in the field from laboratory data

on 42 species of muroid rodents. *Biologist* 63, 138–162 (1981).

32. D. W. Foltz, Genetic evidence for long-term monogamy in a small rodent, *Peromyscus*

polionotus. *The American Naturalist* 117, 665–675 (1981).

33. N. L. Bedford, H. E. Hoekstra, The natural history of model organisms: Peromyscus mice as a

model for studying natural variation. *eLife Sciences* 4, e06813 (2015).

34. L. M. Turner, *et al.*, Monogamy evolves through multiple mechanisms: evidence from V1aR

in deer mice. *Molecular Biology and Evolution* 27, 1269–1278 (2010).

35. S. A. Ramm, G. A. Parker, P. Stockley, Sperm competition and the evolution of male

reproductive anatomy in rodents. *Proceedings of the Royal Society B: Biological Sciences* 272,

949–955 (2005).

36. M. Tourmente, D. Zarka-Trigo, E. R. S. Roldan, Is the hook of muroid rodent's sperm related

to sperm train formation? *Journal of Evolutionary Biology* 29, 1168–1177 (2016).

37. D. M. Higginson, K. B. Miller, K. A. Segraves, S. Pitnick, Convergence, recurrence and

diversification of complex sperm traits in diving beetles (Dytiscidae). *Evolution* 66, 1650–

1661 (2012).

38. G. A. Parker, Why are there so many tiny sperm? Sperm competition and the maintenance of

two sexes. *Journal of Theoretical Biology* 96, 281–294 (1982).

39. R. R. Snook, Sperm in competition: not playing by the numbers. *Trends in Ecology &*

Evolution 20, 46–53 (2005).

40. S. Calhim, S. Immler, T. R. Birkhead, Postcopulatory sexual selection is associated with

reduced variation in sperm morphology. *PLoS ONE* 2, e413 (2007).

41. J. L. Fitzpatrick, B. Baer, Polyandry reduces sperm length variation in social insects. *Evolution*

65, 3006–3012 (2011).

42. S. Immler, S. Calhim, T. R. Birkhead, Increased postcopulatory sexual selection reduces the

intramale variation in sperm design. *Evolution* 62, 1538–1543 (2008).

43. L. Carballo, A. Battistotti, K. Teltscher, M. Lierz, A. Bublat, M. Valcu, B. Kempenaers,

Sperm morphology and evidence for sperm competition among parrots. *Journal of*

Evolutionary Biology 32, 856–867 (2019).

44. A. Rowley, L. Locatello, A. Kahrl, M. Rego, A. Boussard, E. Garza-Gisholt, R. M. Kempster,

S. P. Collin, E. Giacomello, M. C. Follesa, C. Porcu, Sexual selection and the evolution of

sperm morphology in sharks. *Journal of Evolutionary Biology* 32, 1027–1035 (2019).

45. Y. Takami, T. Sota, Sperm competition promotes diversity of sperm bundles in *Ohomopterus*

ground beetles. *Naturwissenschaften* 94, 543–550 (2007).

46. D. M. Higginson, K. B. Miller, K. A. Segraves, S. Pitnick, Female reproductive tract form

drives the evolution of complex sperm morphology. *Proceedings of the National Academy of*

Sciences 109, 4538–4543 (2012).

47. W. Eberhard, *Female Control: Sexual Selection by Cryptic Female Choice* (Princeton

University Press, 1996).

48. I. F. Greenbaum, R. J. Baker, P. R. Ramsey, Chromosomal evolution and the mode of

speciation in three species of Peromyscus. *Evolution* 32, 646–654 (1978).

49. R. C. Firman, L. W. Simmons, Experimental evolution of sperm competitiveness in a

mammal. *BMC Evolutionary Biology* 11 (2011).

50. J. E. Simons, S. D. Olson, "Sperm motility: models for dynamic behavior in complex

environments" in Cell Movement, M. Stolarska, N. Tarfulea, Eds. (Springer International

Publishing, 2018), pp. 169–209.

51. Y. Ishikawa, T. Usui, M. Yamashita, Y. Kanemori, T. Baba, Surfing and swimming of

ejaculated sperm in the mouse oviduct. *Biology of Reproduction* 94, 89, 1–9 (2016).

52. S. Wang, I. V. Larina, In vivo three-dimensional tracking of sperm behaviors in the mouse

oviduct. *Development* 145, dev157685 (2018).

53. V. Magdanz, S. Boryshpolets, C. Ridzewski, B. Eckel, K. Reinhardt, The swim-up technique

separates bovine sperm by metabolic rates, motility and tail length. *bioRxiv*, 624502 (2019).

54. S. S. Suarez, X. Dai, Hyperactivation enhances mouse sperm capacity for penetrating

viscoelastic media. *Biology of Reproduction* 46, 686–691 (1992).

55. M. J. Crawley, *The R Book Second Edition* (John Wiley & Sons, Ltd., 2013).

56. World Health Organization, "WHO laboratory manual for the examination and processing of

human semen" (2010).

57. R Core Team, *R: A Language and Environment for Statistical Computing* (R Foundation for

Statistical Computing, 2016).

58. H. Wickham, *ggplot2: Elegant Graphics for Data Analysis* (Springer, 2016).

59. D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting linear mixed-effects models using lme4.

Journal of Statistical Software 67, 1–48 (2015).

60. X. A. Harrison, Using observation-level random effects to model overdispersion in count data

in ecology and evolution. *PeerJ* 2, e616 (2014).

61. A. F. Zuur, E. N. Ieno, C. S. Elphick, A protocol for data exploration to avoid common

statistical problems. *Methods in Ecology and Evolution* 1, 3–14 (2010).

62. R. V. Lenth, Least-squares means: The R package lsmeans. *Journal of Statistical Software* 69,

1–33 (2016).

63. M. Pagel, Inferring the historical patterns of biological evolution. *Nature* 401, 877–884

(1999).

64. R. P. Freckleton, P. H. Harvey, M. Pagel, Phylogenetic analysis and comparative data: a test

and review of evidence. *The American Naturalist* 160, 712–726 (2002).

65. D. Orme, R. Freckleton, G. Thomas, T. Petzoldt, The caper package: comparative analysis of

phylogenetics and evolution in R. *R package version* 5, 1–36 (2013).

66. E. Paradis, J. Claude, K. Strimmer, APE: Analyses of phylogenetics and evolution in R

language. *Bioinformatics* 20, 289–290 (2004).

67. R. D. Bradley, N.D. Durish, D. S. Rogers, J. R. Miller, M. D. Engstrom, C. W. Kilpatrick,

Toward a molecular phylogeny for Peromyscus: evidence from mitochondrial cytochrome- *b*

sequences. *Journal of Mammalogy* 88, 1146–1159 (2007).

68. E. García-Berthou, On the misuse of residuals in ecology: testing regression residuals vs. the

analysis of covariance. *Journal of Animal Ecology* 70, 708–711 (2001).

69. S. Lüpold, G. M. Linz, J. W. Rivers, D. F. Westneat, T. R. Birkhead, Sperm competition

selects beyond relative testes size in birds. *Evolution* 63, 391–402 (2009)

Figure 1.

The observed natural variation in the proportion of aggregated sperm cells for six closely related species of *Peromyscus* mice (phylogeny adapted from Bradley et al. 2007). Box-plots represent median and interquartile ranges with raw data overlaid. Statistically significant differences at the *P* = 0.05 level are denoted by differing letters; shared letters denote no statistical difference.

Figure 2.

Sperm aggregate size varies among species and is negatively associated with relative testes size. (A) The observed natural variation in the number of aggregated sperm cells for six closely related species of *Peromyscus* mice. Box-plots represent median and interquartile ranges with raw data overlaid. Statistically significant differences at the $P = 0.05$ level are denoted by differing letters; shared letters denote no statistical difference. (B) When controlling for phylogenetic relationships, the coefficient of variation (CV) for the number of aggregated sperm cells negatively correlates with relative testis mass across these species. Note truncated y-axes.

Figure 3.

The effect of sperm aggregate geometry on the curvilinear velocity (*μ*m/sec) of sperm aggregates for two species that aggregated most – *Peromyscus maniculatus* and *Peromyscus polionotus*. Circles represent mean values per male within each species, and black lines represent the mean value within each category. Note truncated y-axis.

Figure 4.

Kinematic parameters of sperm aggregates (closed circles) and solitary sperm cells (open circles) for six species of *Peromyscus* mice in low- and high-viscosity conditions. Circles represent mean values per male, and black lines represent species means. Statistical significance levels comparing aggregated and solitary cells within each species are indicated by * *P* < 0.05, ** *P* < 0.01, *** *P* < 0.001. Note truncated y-axes.

TABLE 1.

Summary of live sperm aggregate results for mice in the genus *Peromyscus*

Peromyscus Species	No. males	No. sperm	% motile sperm	% progressive sperm	No. sperm aggregates	% cells in aggregate	% motile aggregates	% progressive aggregates
californicus	28	8235	81.2%	78.9%	823	29.7% (2446/8235)	90.5% (745/823)	86.6% (713/823)
eremicus	21	4906	67.4%	58.2%	513	30.2% (1481/4906)	77.8% (399/513)	66.7% (342/513)
polionotus	24	6360	82.4%	57.3%	949	79.5% (5059/6360)	90.9% (863/949)	66.1% (627/949)
maniculatus	18	4991	85.0%	84.3%	822	81.3% (4059/4991)	93.4% (768/822)	92.3% (759/822)
leucopus	22	6341	87.6%	84.5%	361	15.0% (949/6341)	90.3% (326/361)	88.1% (318/361)
gossypinus	21	5970	82.2%	70.3%	254	9.6% (576/5970)	88.6% (225/254)	70.1% (178/254)

Motile sperm cells include only those that exhibited movement patterns. Progressive sperm cells are motile cells with a straight-line velocity ≥ 25 μm/sec.

TABLE 2.

Fixed effects from a binomial generalized linear mixed model examining differences in the proportion of aggregated sperm cells and a linear model examining differences in the number of aggregated sperm cells across six species of *Peromyscus* mice

GLMM: PROPORTION OF CELLS IN AGGREGATE					
Model Term	Beta (SE)	Exp (beta)	95% CI	z	Pr(>|z|)
Intercept	1.52 (0.22)				
eremicus	-2.47 (0.30)	0.07	(0.04, 0.13)	-8.14	**< 0.001**
gossypinus	-3.94 (0.31)	0.02	(0.01, 0.03)	-12.89	**< 0.001**
californicus	-2.58 (0.29)	0.07	(0.04, 0.12)	-9.06	**< 0.001**
leucopus	-3.79 (0.30)	0.02	(0.01, 0.04)	-12.46	**< 0.001**
polionotus	-0.25 (0.29)	0.44	(0.30, 0.58)	-0.85	0.393
LM: NUMBER OF CELLS IN AGGREGATE					
Model Term	Beta (SE)	Exp (beta)	95% CI	t	Pr(>|z|)
Intercept	3.90 (0.26)				
Total Sperm Cells	0.00 (0.00)	0.50	(0.50, 0.50)	5.83	**< 0.001**
eremicus	-1.97 (0.24)	0.12	(0.08, 0.18)	-8.15	**< 0.001**
gossypinus	-2.77 (0.24)	0.06	(0.04, 0.09)	-11.53	**< 0.001**
californicus	-2.19 (0.23)	0.10	(0.07, 0.15)	-9.68	**< 0.001**
leucopus	-2.68 (0.24)	0.06	(0.04, 0.10)	-11.18	**< 0.001**
polionotus	0.04 (0.23)	0.51	(0.40, 0.62)	0.16	0.877

For both models, all rows are being compared with the intercept – *Peromyscus maniculatus*. 95% confidence intervals (CI) were calculated for each effect size.

TABLE 3.

Results from an intra-male analysis comparing motile solitary and aggregated sperm kinematics in low- and high-viscosity conditions for six species of *Peromyscus* mice to test whether sperm aggregates confer kinematic benefits (shaded in gray)

PEROMYSCUS SPECIES	df	KINEMATIC VARIABLE			
		Curvilinear Velocity (µm/sec)	Linearity (VSL/VCL)	Straight-Line Velocity (µm/sec)	Average Path Velocity (µm/sec)
LOW-VISCOSITY CONDITIONS					
californicus	28	t = -1.0545, p = 0.3007	t = -0.56153, p = 0.5789	t = -2.2982, ***p* = 0.02923**	t = - 2.8162, ***p* = 0.008805**
eremicus	20	t = 1.6225, p = 0.1204	t = 0.73517, p = 0.4708	t = 0.1312, p = 0.8969	t = 1.0464, p = 0.3079
polionotus	23	t = 9.4575, ***p* = 2.1566e-09**	t = 1.699, p = 0.1028	t = 5.8355, ***p* = 6.026e-06**	t = 6.0729, ***p* = 3.408e-06**
maniculatus	17	t = 2.2482, ***p* = 0.03812**	t = 0.48075, p = 0.6368	t = -2.2335, ***p* = 0.03924**	t = -1.9206, p = 0.07172
leucopus	20	t = 0.23337, p = 0.8178	t = -4.8385, ***p* = 9.973e-05**	t = -5.5521, ***p* = 1.959e-05**	t = -1.838, p = 0.08096
gossypinus	20	t = 5.4048, ***p* = 2.73e-05**	t = -1.0247, p = 0.3177	t = 3.715, ***p* = 0.001369**	t = 4.479, ***p* = 0.0002298**
HIGH-VISCOSITY CONDITIONS					
californicus	9	t = -3.7465, ***p* = 0.003357**	t = 2.682, ***p* = 0.02512**	t = -2.7743, ***p* = 0.0216**	t = -4.1106, ***p* = 0.002634**
eremicus	9	t = -0.71739, p = 0.4913	t = 4.2087, ***p* = 0.002277**	t = 0.62918, p = 0.5449	t = 0.56505, p = 0.5858
polionotus	13	t = 2.0349, p = 0.06278	t = 3.9242, ***p* = 0.001745**	t = 4.8314, ***p* = 0.0003279**	t = 4.18, ***p* = 0.001079**
maniculatus	11	t = -2.9397, ***p* = 0.01345**	t = 3.9259, ***p* = 0.002369**	t = -0.30835, p = 0.7636	t = -1.0543, p = 0.3114
leucopus	9	t = 0.32429, p = 0.7531	t = -0.31635, p = 0.759	t = -0.34159, p = 0.7405	t = 0.95055, p = 0.3667
gossypinus	10	t = -0.63835, p = 0.95376	t = -0.076451, p = 0.9406	t = -0.60402, p = 0.5593	t = -0.71643, p = 0.4901
Statistical results are based on paired-student t-tests.					