Sustainable DNA data storage with low biased manipulation

Yanmin Gao[1,2], Xin Chen[3], Jianye Hao[4], Chengwei Zhang[5], Hongyan Qiao[1,2] and Hao Qi[1,2*]

[1] School of Chemical Engineering and Technology, Tianjin University, Tianjin, China.

[2] Key Laboratory of Systems Bioengineering (Ministry of Education), Tianjin University, Tianjin, China.

[3] Center for Applied Mathematics, Tianjin University, Tianjin, China.

[4] College of intelligence and computing, Tianjin University, Tianjin, China.

[5] College of Information Science and Technology, Dalian Maritime University, Dalian, China.

* Correspondence should be addressed to H.Q. (haoq@tju.edu.cn)

Abstract:

In DNA data storage, the huge sequence complexity is challenging repeatable and efficient information reading. Here, we demonstrated that biased amplification is serious matter for stable information retrieve from synthetic oligo pool comprising over ten thousand strands encoding over 2.85 MB digital information. Therefore, we adapted isothermal DNA amplification as a new biochemical framework, termed as isothermal DNA reading (iDR) for low biased manipulation of DNA pool with huge sequence complexity. iDR was built to work in a priming-free and error-spreading proof manner and achieved low-biased amplification. Finally, we demonstrated that immobilized DNA materials read by iDR is able to stably read data deeply and repeatedly, a practical and sustainable DNA storage system. Necessary amount of sequencing reads for perfect

21   oligos retrieve significantly decreased. These advanced features enable the iDR chemical

22   framework being ideal alternative for manipulating the huge sequence complexity and building

23   the sustainable and efficient DNA storage "hardware".

Introduction:

In DNA data storage, technology originally developed for bioengineering approach including array oligo synthesis, PCR and DNA sequencing were integrated to construct the "hardware" for DNA storage(*1-5*). Array synthesized DNA oligo pool comprising of from thousands up to millions of oligo strands with several hundred bases in length has been utilized in many applications, e.g., probe blend, DNA origami assembly, and genome synthesis(*1*). Because of both the location on microchip and DNA sequence context, the copy number for each oligo strand from array synthesis is distinct(*5*). Furthermore, the sheer number of synthesized oligos from array on microchip is very small, roughly from 10^5 to 10^{12} at the concentration of femtomolar depending on the synthesis platform(*6, 7*). Generally, amount of DNA materials, over a few hundred nanograms, at the concentration of micromolar is necessary for high quality DNA sequencing covering all oligos in the pool on commercial DNA sequencing platform, e.g., Illumina(*8*). Therefore, amplification is crucial to boost the signal for the subsequent DNA sequencing for stably reading data.

The oligo pool size for DNA data storage is much larger at least several orders of magnitude than that in other bioengineering applications(*9-11*). The unevenness of copy number generated a huge complexity that caused serious problem for DNA molecule retrieval and data decoding(*4, 5, 12, 13*). Thus far, in reported systems the information reading was achieved by PCR amplification and NGS, but the copy number unevenness originally stemmed from microchip synthesis was further skewed by highly biased amplification process(*5, 12*). Therefore, more amplified DNA materials was necessary to fetch the minor oligo strands from the skewed oligo pool. The length and sequence context, GC content and secondary structure of DNA molecule are well known to introduce large amplification bias in PCR for amplification of multiple templates in parallel(*14*). Minor oligos could be excluded easily in a few amplification cycles because of its replication

disadvantage. However, DNA storage requires amplification of from tens of thousands up to

million distinct DNA strands at the same time. Thus far, this problem was dealt by great effort to

develop high encoding, physical and sequencing redundancy(*13, 15*). However, high redundancy

sacrificed storage density and increased cost in synthesis, sequencing and decoding calculation. In

previous studies, it has been demonstrated that deep PCR amplification (over 60 thermal

amplification cycles) increased the unevenness of copy number largely and required several orders

of magnitude more sequencing reads for decoding(*12*). For extreme case, only a few oligos (less

than 10 % of the total oligos) wiped out almost all others after amplification(*16*). Moreover,

successive repeated amplification significantly skewed the copy number distribution and caused

large dropout(*12*). Even the sequences were carefully designed to minimize the sequence context

difference, biased amplification still caused chaos from the huge sequence complexity of oligo

pool for DNA storage and then imperiled the information decoding. And for this pure engineering

approach of DNA storage, it is valuable to develop alternative technology with distinct features.

For practical data storage, crucial issues have to been addressed. Information reading method

should be redesigned to handle amplification of oligo pool with high complexity of DNA sequence

at low bias and also support the repeated reading for long-term storage. The current thermal cycling

process in DNA amplification not only consumes energy but also will trouble the operation of

storage device. It has been reported that long thermal treatment at around 65°C resulted to oligo

decay and higher temperature caused more damage to longer DNA molecule(*16, 17*). All these

issues come down to developing biosystem which could stably and repeatedly handle oligo pool

with high sequence complexity and uneven copy number under more simple thermal condition.

Here, we adapted a BASIC code system(*18, 19*) recently developed for digital distributed file

system for DNA storage, in which a high information density was achieved with a low encoding

redundancy of 1.56%. For amplification of oligo pool with high sequence complexity, we

challenge to develop alternative system for low biased amplification and construction of repeatable

DNA storage. Statistics analysis demonstrated that oligo pool was more prone to dropout than base

error and oligo dropout significantly increased with the amplification depth and repeated times in

biased amplification of PCR. Therefore, we recruited the isothermal DNA replication to challenge

the stable manipulation of DNA pool with huge sequence complexity. Isothermal DNA reading

(termed as iDR) was specifically built to address the crucial issues of inefficient-priming, product-

as-template that caused the replication bias and error-spreading and oligo dropout remained very

stable at low level. Finally, we demonstrated that immobilized DNA pool read by iDR as an

efficient and repeated DNA system for sustainable information storage. For the development of

practical DNA "hardware", and iDR could work with PCR together in the different process, e.g.,

material preparation and stably reading respectively. To the best of our knowledge, this is the first

sustainable DNA storage system successfully achieved the stable information "deep reading".

Results:

BASIC code for DNA mediated distributed storage.

In current DNA storage, the digital file was divided and written into a large group of small piece

DNA oligos. Every single oligo function as individual information carrying unit. And then, the

entire file can be read out from sequencing all the oligos (Fig. 1A). Therefore, DNA storage can

be considered as distributed storage system built on biomaterial. Due to this nature, we adapted a

BASIC coding system that is one well optimized regenerating code with reducing computational

complexity for distributed storage (*18, 19*). Generally, the encoding process started by dividing

the target file into non-overlapping groups and the split information was encoded into DNA

sequence following an optimized encoding process (fig. S1 and Supplementary Note 1) with two

adjustable parameter K (corresponding to oligo number in one non-overlapping group, K was set

as constant value of 256) and L (corresponding to the length of oligo), which is flexible for various

DNA oligo length from different commercial synthesis platform. Besides the normal file types, for

the first time we tested the storage of genome sequence including human mitochondrion and one

artificial bacteria cell (Supplementary Note 2) with a different encoding strategy. Genome

nucleotide sequence was converted to binary file following a simple principle of T to 00, G to 01,

C to 10 and A to 11, by which higher information storage density could be obtained as one

nucleotide base to 2-bits binary in comparing with simply treating nucleotide letter as text contents.

Because in general text computer system, e.g., ASCII, one letter is encoded by 8-bit binary. In

nature genome sequence, there are many complicate structures hard for correct sequencing.

Through the encoding process, the genome sequence was rewritten into nucleotide sequence again

but with complicated sequence avoided and accurate sequencing guaranteed by error correcting.

Furthermore, oligo pools with various payload length have been designed to store 2.85 MB files

in totally 109,568 oligo strands (Fig. 1A, fig. S2A and Table S1). Reed-Solomon code was used

for error correction and coding redundancy for tolerance of missing entire oligo, causing dropout.

Relative high information density was 1.65 bits/nt with a 1.56% of coding redundancy, which was

calculated by previous reported method without modification (Supplementary Note 3)(*13*).

Success decoding could be achieved as long as oligo dropout less than 1.56%, randomly losing 4

strands from 256 (fig. S1B). Technically, higher encoding redundancy tolerates losing more oligos

and success decoding can be achieved from lower sequencing coverage. However, higher

redundancy requires more DNA synthesis. Considering the synthesis cost is higher than

sequencing, represented over 90% of the total cost of DNA storage in our experiment, it is more

practical to trade-off encoding redundancy for synthesis cost.

Chip-based synthesis only produces very small amount of oligo, roughly from 10^5 to 10^{12} at the

concentration of femtomolar depending on the synthesis platform(*7*). In the current workflow of

DNA data storage, the quality of oligo pool majorly impacted the storage performance(*4*).

Particularly, both the heterogenous oligo sequences and the unevenness of oligo copy number

generated a huge sequence complexity and then caused the PCR amplification bias. Minor oligo

molecules are prone to drop-out and more sequencing coverage was required for decoding oligo

pool with largely skewed copy number distribution (Fig. 1B). The material of skewed oligo pool

makes stable and repeatable information decoding a huge challenge.

Low-bias and error-spreading proof isothermal amplification

It well known that PCR generated biased amplification from its inherent mechanism, i.e., product-

as-template, priming and thermal cycling dependent amplification(*3, 14, 20, 21*). In order to

address these problems that easily skewed the oligo copy number distribution, we recruited

isothermal DNA replication, which are majorly used in systems for biomolecular analysis or

complex DNA circuit network (*22-24*). To the best of our knowledge, strand displacement

replication has not been tested for stable amplification of large DNA library, not the size of library

in DNA storage. It is still unknown if huge sequence complexity could be stably amplified by this

novel DNA replication mechanism. Therefore, we challenge to build low biased isothermal

amplification of DNA pool with high sequence complexity. After systematical optimization of

reaction temperature, reagent component, replicon length and nickase recognition site sequence

(figs. S2B-E), Nt.BbvCI nickase and exonuclease deficient DNAP I Klenow fragment $(3'{\rightarrow}5'$ exo^{-})

were used and the amplification efficiency of iDR is around 70% of PCR (fig. S2F). The

immobilized isothermal DNA replication system was designated as iDR, isothermal DNA reading

for large DNA pool amplification (Fig. 2A). The Nt.BbvCI nickase recognition site is only 7nts in

length and its sequence has been avoided from the payload part of oligo pool in the file encoding

step. By comparison, it will not cause much extra burden on synthesis.

The intrinsic features of iDR specifically facilitate DNA storage application. In theory, for one 50

ul reaction, 10 PCR thermal cycles required about 177.8J energy only for thermal regulation at

least two orders of magnitude more than 30 mins iDR of 2.52J (Fig. 2B and Supplementary Note

4) and it may be a huge issue at large scale of operation. Real time monitoring indicated that 30

mins is the optimal amplification time (fig. S2G) and the amplification time was set as 30 mins for

all experiments in this manuscript. Single-stranded or double-stranded DNA can be produced in a

controlled manner (Fig. 2C). Specially, ssDNA was amplified in a primer-free manner and allows

iDR to be a universal method for reading information with no sequence information needed in

advance. The nickase mediated site-specific phosphodiester bond cleavage initiated the iDR

amplification and generated a 5' terminal phosphate group, its function was verified by direct

ligation to a FAM labeled probe (Fig. 2D). This inborn phosphate group is very convenient for

subsequent functional adapter linking.

Amplified oligos were sequenced on commercial Illumina Hiseq 4000 platform with 150 paired-

end cycles (Supplementary Note 5) and then deeply analyzed by a set of statistics methods

developed from bioinformatic BLAST program (fig. S3 and Supplementary Note 6). Sequenced

reads with various number of letter error including substitution and indel were different. Among

them, single letter error accounted for the majority in mutant sequenced reads for both PCR (80.1%)

and iDR (81.7%) (Fig. 2E and fig. S4). The total indel (0.03%) and substitution (0.2%) rate were

consistent with previous studies(*13*). Although, RS code embedded in BASIC code is able to

correct multiple errors in same DNA strand but will increase the computation complexity

largely(*17, 25*). Therefore, both sequenced reads with no or single substitution/indel error were

collected as valid reads for further analysis and decoding (fig. S1B).

The distribution of the number of reads per each given sequence were different between iDR and

PCR of various thermal cycles of from 10 to 60. The distribution normality was quantified by a

modified mathematical statistics function (Supplementary Note 6). In theory, oligo pool of higher

value distribution normality require less sequencing reads to recover all synthesized oligos(*12*).

The coverage distribution of both PCR and iDR were positively skewed with a long tail, which

comprising of high copy number oligos, the normality of PCR decreased from 0.50 to 0.18 with

thermal cycles increased from 10 to 60 (Fig. 2F). However, 0.60 was obtained for iDR, it indicated

that most balanced DNA pool was amplified by iDR. The proportion of oligos with high copy

number, the top 30% high coverage, enlarged with the PCR thermal cycles increased (Fig. 2G). In

comparison, the dropout rate increased significantly from 4.18% to 13.80% with PCR

amplification becoming deeper, much higher than 1.33% of iDR (Fig. 2G). No obvious difference

was observed between the coverage distribution of iDR amplification from free oligo pool and

oligo pool immobilized on magnetic beads (fig. S5).

Deep errors, reads with mutant in high copy number, impeded information decoding. Sequenced

reads were further sorted out as group of M0G0 (with no letter error) and M1G1 (with single letter

substitution or indel error) by developed BLAST programs. Therefore, for same oligo, if the

coverage in M0G0 is smaller than the coverage in M1G1, it indicated that for this oligo sequencing

reads with mutant is more than correct one in the pool. It is hard to identify the correct oligo from

sequencing reads by principle of law of large number-makers and will tend to influence the correct

decoding. We identified this as number of deep-error, it means how many oligos with more mutant

sequencing reads than correct reads. It was observed that PCR with various thermal cycles gave

average number of 177 for 10 cycles, 350 for 30 cycles and 1175 cycles for 60 cycles, all of them

is larger than iDR of 46 (Fig. 2H). It indicated that biased amplification spread and accumulated

error reads.

Low replication fidelity of DNA polymerases and around 1% miss reading coming along with

NGS sequencing process(*26*) largely contributed to the massive error reads with low copy number,

e.g. 1-2 copy number, but it is relatively easy to identify them from correct reads in high copy

number. Considering the mechanism, product-as-template and inefficient priming process caused

the high amplification bias and made PCR amplification prone to error-spreading and oligo

dropout. In contrast, iDR was designed to synthesize new oligo only from the original templates

without priming process for replication initiation and therefore iDR achieved a low biased and

error-spreading proof amplification and will require much less calculation resource in decoding

process. Additionally, high temperature treatment resulted to DNA oligo decay especially for long

strands(*16, 17*). The decay lost rate in a 10 cycles PCR was calculated as 21.8‰ and 0.035‰ for

iDR (Fig. 2B and Supplementary Note 7). Finally, all the files in DNA pool was successfully

decoded following the BASIC decoding process and the minimal mean coverage depth of

sequencing reads were calculated (Table S2)

Sustainable DNA storage for repeated reading

Repeatable reading is crucial for practical DNA storage system. So far, different repeated reading

systems have been proposed and achieved, in which generally DNA pool was amplified from

aliquot of previous amplified product (Fig. 3A). However, error rate and oligo dropout were

significantly accumulated through this repeated amplification(*12, 13, 16*). Due to the low biased

amplification performance, we believe that DNA storage for stable repeated reading could be

achieved from iDR system. In this isothermal system (Fig. 3B), both the low biased iDR

amplification performance and DNA pool immobilization on microbeads could contribute to the

stability in repeated reading, but either of them has not been systematically tested so far. Therefore,

we propose the isothermal reading of immobilized DNA pool as stable and robust DNA storage

system.

The amplification capability for repeated deep reading oligo pool containing from 11,520 to 89,088

DNA strands was examined. PCR amplification was successively performed 10 times from aliquot

of previous reaction and iDR amplification was repeated 10 times from immobilized oligos pool.

In #1, #5 and #10 of successive PCR, the proportion of amplified oligos with up to total 10

substitution or indel letter error decreased from 89.09% ($\pm$10 of PCR #1) to 49.97% ($\pm$10 of PCR

#10) and from 86.77% ($\pm$1 of PCR #1) to 48.58% ($\pm$1 of PCR #10). In comparison, the repeated

iDR remained very consistent over 90% (Fig. 3C). These statistics results indicated that large noise

was introduced during PCR procedure and amplified oligo with imperfect length increased from

inefficient replication or miss priming. However, no obvious difference was observed in

proportion of M1G1 in total sequenced reads between PCR and iDR, but the mean copy number

in M1G1 reads increased from 1.10 of #1 PCR to 1.95 of #10 PCR, but only slightly changed for

iDR, 1.08 of #1, 1.08 of #5 and 1.05 of #10 (Fig. 3D). This result was in agreement with previous

experiment and indicated that error accumulated to high copy number during PCR and iDR

achieved error-spreading proof amplification, in which amplified DNA strand with error was not

further amplified as template.

Interestingly, it was observed that the distribution of the number of reads per each given sequence

changed differently. Normality of #1, #5 and #10 PCR decreased significantly from 0.50 to 0.15.

In contrast, #1, #5 and #10 iDR gave a consistent normality of 0.58 (Fig. 3E). For successive PCR,

the coverage distribution was largely positively skewed and the proportion of both low copy

number and high copy number oligos significantly increased. It indicated that large enrichment

driven by amplification bias for part of oligos efficiently occurred with the successive PCR, but

not in repeated iDR. Furthermore, it was observed that only top 1% oligo of high coverage

increased its proportion significantly and both 1% oligos of middle and low coverage decreased

while the oligo pool was successively read by PCR. In contrast, all the 1% oligos remained steady

in repeated iDR (fig. S6). The dropout rate in random valid reads set with 10x coverage was

quantified to further assess the amplification bias. For PCR, the dropout rate increased sharply

from 4.18% (#1 PCR) to 53.19% (#10 PCR) (Fig.3F), but iDR remained steady at about 2%. Due

to the tolerance for 1.56% dropout in BASIC encoding algorithms, we also calculated the coverage

depth for random valid reads set with 1.56% dropout (Supplementary Note 3), the crucial

parameter for the theoretical minimum decoding coverage. For PCR, the minimum coverage depth

was quantified as 17.2x (#1 PCR) and 167x (#5 PCR) and 426x (#10 PCR). For iDR, the minimum

coverage depth was quantified as 11 for #1, 12 for #5 and 12.5 for #10. Therefore, for 445KB file

encoded in 11520 DNA oligo pool, 10,101,935 and 149, 919 minimal sequencing reads were

necessary for perfect decoding (Fig. 4 and Supplementary Note 8).

Thus far, for PCR based sustainable DNA media reading, one strategy is deep amplification from

trace DNA material with large number of thermal cycles and another is successive amplification

from aliquot of previous reaction. However, we demonstrated that both strategies are not practical.

The nature of PCR including especially inefficient priming and product-as-template generated

large amplification bias that significantly skewed the copy number distribution and resulted huge

dropout in both 60 cycles deep amplification and 10 times successive amplification, especially for

oligo pool with huge sequence complexity. Based on these deep statistical analyses, we pointed

out that iDR amplification of immobilized DNA pool was stable, robust and suitable for repeated

reading in DNA data storage.

Discussion:

DNA data storage is one very artificial application and actually the concept has been proposed for a long time(*27*). But till lately, the significantly increasing capability of DNA synthesis and sequencing start making it possible. In many respects, practical DNA data storage requires more powerful capability for all related biotechnologies, including synthesis, sequencing and manipulation of oligo pool with high sequence complexity. The size of oligo pool used in DNA storage is already several orders of magnitude larger than that in other applications. However, there is still short of hardware technology which is designed for DNA storage, especially manipulation of DNA oligo pool with huge sequence complexity and unevenness of molecule copy number. In studies to date, PCR is still the only method for amplification of oligo pool for reading information. However, with deep bioinformatic and statistics analysis, we demonstrated that PCR amplification skewed the oligo pool even after a very few thermal cycles (10 cycles) and the skewness largely increased, deep error spreading and massive dropout, as the function of amplification cycle numbers. Deep error, mutant sequenced reads in high coverage, interfered decoding and caused significant increased calculation, but decoding fatally crashed due to massive dropout. We contend that the features of low temperature, priming-free, and enzyme-mediated double helix unzipping in iDR amplification presented here, overcame the major amplification bias related issues of PCR including the inefficient priming, product-as-template, sequence context dependent, and high temperature heating. iDR achieved very stable amplification performance, which efficiently prevented the mutant error from spreading, decreased over 70% deep error, and achieved successive repeated deep reading with consistent outcome quality. Actually, the physical storage density, the most significant advantage of DNA storage with a calculation of a few kilograms DNA material for storage of data from the whole human society(*28*), highly depend on the quality of

DNA reading. Therefore, higher physical storage density could be achieved by iDR system, which

could store information with at least two orders of magnitude less DNA material and sequencing

resource than the current PCR method (Fig. 4 and Supplementary Note 8).

Although, there is many variants of PCR, such as emulsion PCR, digital PCR, and multiplex PCR,

but none of them can avoid the crucial issues of product-as-template, inefficient priming, and

complex thermal regulation. Each of these will raised the cost for handling the large DNA pool.

Although, iDR is adapted from stand displacement replication, but this is the first system applying

this amplification mechanism on DNA pool with high sequence complexity for artificial

information storage and demonstrated that immobilized DNA pool read by iDR is able to stably

read data deeply and repeatedly. Originally, the core thing of DNA storage is adapting current

biotechnology for artificial purpose of storing information at a large scale and accuracy which is

distinct for bioengineering approaches. Undeniably, PCR possesses power in DNA amplification,

and iDR could work with PCR together in the different steps of DNA storage, e.g., material

preparation and stably reading respectively.

Furthermore, Replication fidelity of DNAP I Klenow fragment used in iDR system is much lower

than Q5 DNA polymerase in PCR. There is still much room for improvement. Additionally,

practical DNA storage system must move out of biochemical test tube to build device by highly

integrating all related biochemical processes. Prototypes of DNA storage hardware have already

been proposed with high density DNA material in microfluidic device(*29, 30*). We contend that

besides some crucial features of iDR make it more fit for hardware construction. In iDR, only low

temperature, slightly higher than room temperature, was required. Considering the operation of

DNA storage up to large scale, there will be a huge difference in energy consumption. Furthermore,

iDR is able to work in a primer-free fashion, only defined protein enzyme mix is required no matter

what information was encoded, which makes it possible for universal information reading without

any sequence information required in advance. To the best of our knowledge, iDR is the first

system for stable repeated DNA information reading which is crucial and necessary feature for

practical and sustainable storage hardware. Therefore, we believed that this new biochemical

framework with stable repeated reading lays a foundation for development of practical and

sustainable DNA storage.

ACKNOWLEDGMENTS

We would like to thank Professor Hanxu Hou from Dongguan University of Technology for advice and assistance with designing algorithm for BASIC code. We also thank Yixi Wang from College of intelligence and computing at Tianjin University for her help in test of encoding system. This work was supported by the National Science Foundation of China (Grant No.21476167, No.21778039 and No.21621004).

AUTHOR CONTRIBUTIONS

Y.G. and H.Q. designed and performed all experiments. X.C., J.H. and C.Z. designed and developed the encoding and decoding program. X.C. developed program for the bioinformatics statistics analysis. Y.G., H.Qiao and H.Q. collected and analyzed all experiment data. G.Y. T.H. and H.Q. wrote the manuscript. H.Q. designed experiments, analyzed data and supervised this work.

COMPETING FINANCIAL INTERESTS

H.Q. is the inventor of two patents application for the biochemical method described in this article. The initial filings were assigned Chinese patent application (201911086860.0 and 201911087247.0) and international patent application (PCT/CN2019/123916). The remaining authors declare no competing financial interests.

References:

329   1.   S. Kosuri, G. M. Church, Large-scale de novo DNA synthesis: technologies and
330   applications. *Nat Methods* **11**, 499-507 (2014).

331   2.   S. Goodwin, J. D. McPherson, W. R. McCombie, Coming of age: ten years of next-
332   generation sequencing technologies. *Nat Rev Genet* **17**, 333-351 (2016).

333   3.   I. Kozarewa *et al.*, Amplification-free Illumina sequencing-library preparation facilitates
334   improved mapping and assembly of (G+C)-biased genomes. *Nat Methods* **6**, 291-295
335   (2009).

336   4.   L. Organick *et al.*, Experimental Assessment of PCR Specificity and Copy Number for
337   Reliable Data Retrieval in DNA Storage. (2019).

338   5.   Y.-J. Chen *et al.*, Quantifying Molecular Bias in DNA Data Storage. (2019).

339   6.   J. C. Klein *et al.*, Multiplex pairwise assembly of array-derived DNA oligonucleotides.
340   *Nucleic Acids Res* **44**, e43 (2016).

341   7.   H. G. Jingdong Tian, Nijing Sheng, Xiaochuan Zhou, Erdogan Gulari, Xiaolian Gao &
342   George Church, Accurate multiplex gene synthesis from programmable DNA microchips.
343   *Nature* **432**, 1050-1054 (2004).

344   8.   S. Linnarsson, Recent advances in DNA sequencing methods - general principles of sample
345   preparation. *Exp Cell Res* **316**, 1339-1343 (2010).

346   9.   S. Kosuri *et al.*, Scalable gene synthesis by selective amplification of DNA pools from
347   high-fidelity microchips. *Nat Biotechnol* **28**, 1295-1299 (2010).

348   10.   R. J. Lipshutz, S. P. Fodor, T. R. Gingeras, D. J. Lockhart, High density synthetic
349   oligonucleotide arrays. *Nat Genet* **21**, 20-24 (1999).

350   11.   M. T. Bonde *et al.*, Direct mutagenesis of thousands of genomic targets using microarray-
351   derived oligonucleotides. *ACS Synth Biol* **4**, 17-22 (2015).

352   12.   Y. Erlich, D. Zielinski, DNA Fountain enables a robust and efficient storage architecture.
353   *Science* **355**, 950-954 (2017).

354   13.   L. Organick *et al.*, Random access in large-scale DNA data storage. *Nat Biotechnol* **36**,
355   242-248 (2018).

356   14.   D. Aird *et al.*, Analyzing and minimizing PCR amplification bias in Illumina sequencing
357   libraries. *Genome Biol* **12**, R18 (2011).

358   15.   N. Goldman *et al.*, Towards practical, high-capacity, low-maintenance information storage
359   in synthesized DNA. *Nature* **494**, 77-80 (2013).

360   16.   R. Heckel, G. Mikutis, R. N. Grass, A Characterization of the DNA Data Storage Channel.
361   *Sci Rep* **9**, 9663 (2019).

362   17.   R. N. Grass, R. Heckel, M. Puddu, D. Paunescu, W. J. Stark, Robust chemical preservation
363   of digital information on DNA in silica with error-correcting codes. *Angewandte Chemie*
364   **54**, 2552-2555 (2015).

365   18.   H. Hou, K. W. Shum, M. Chen, H. Li, BASIC Codes: Low-Complexity Regenerating
366   Codes for Distributed Storage Systems. *IEEE Transactions on Information Theory* **62**,
367   3053-3069 (2016).

368   19.   H. Hou, Y. S. Han, K. W. Shum, H. Li, A Unified Form of EVENODD and RDP Codes
369   and Their Efficient Decoding. *IEEE Transactions on Communications* **66**, 5053-5066
370   (2018).

371   20.   E. L. van Dijk, Y. Jaszczyszyn, C. Thermes, Library preparation methods for next-
372   generation sequencing: tone down the bias. *Exp Cell Res* **322**, 12-20 (2014).

21. J. Dabney, M. Meyer, Length and GC-biases during sequencing library amplification: a
comparison of various polymerase-buffer systems with ancient and modern DNA
sequencing libraries. *Biotechniques* **52**, 87-94 (2012).
22. Y. Zhao, F. Chen, Q. Li, L. Wang, C. Fan, Isothermal Amplification of Nucleic Acids.
Chem Rev **115**, 12491-12545 (2015).
23. M. S. Reid, X. C. Le, H. Zhang, Exponential Isothermal Amplification of Nucleic Acids
and Assays for Proteins, Cells, Small Molecules, and Enzyme Activities: An EXPAR
Example. *Angewandte Chemie* **57**, 11856-11866 (2018).
24. D. Y. Zhang, G. Seelig, Dynamic DNA nanotechnology using strand-displacement
reactions. *Nat Chem* **3**, 103-113 (2011).
25. L. Anavy, I. Vaknin, O. Atar, R. Amit, Z. Yakhini, Data storage in DNA with fewer
synthesis cycles using composite DNA letters. *Nat Biotechnol*, (2019).
26. S. M. Yazdi, Y. Yuan, J. Ma, H. Zhao, O. Milenkovic, A Rewritable, Random-Access
DNA-Based Storage System. *Sci Rep* **5**, 14138 (2015).
27. M. R. Wallace, Molecular Cybernetics: The Next Step? *Kybernetes* **7**, 265-268 (1978).
28. A. Extance, How DNA could store all the world's data. *Nature* **537**, 22-24 (2016).
29. S. Newman *et al.*, High density DNA data storage library via dehydration with digital
microfluidic retrieval. *Nat Commun* **10**, 1706 (2019).
30. C. N. Takahashi, B. H. Nguyen, K. Strauss, L. Ceze, Demonstration of End-to-End
Automation of DNA Data Storage. *Sci Rep* **9**, 4998 (2019).

Figure 1: Overview of the DNA data storage.

(**A**) Schematic of the DNA mediated distributed data storage and totally 2.85 MB data, including texts, images, videos, and block chain file, computer virus code, genetic information, were encoded into total 109,568 synthesis oligos using an adapted BASIC code system.

(**B**) Illustration of data storage in synthesis oligo pool. The huge sequence complexity and unevenness of oligo copy number is challenging the stable storage and precise decoding.

Figure 2: Designed isothermal DNA reading (iDR) for DNA data storage.

(A) Illustration of the DNA amplification of PCR (upper) and iDR (lower). PCR depend on thermal cycling to drive replication. By contrast, isothermal DNA reading was developed from systematically optimized strand displacement mediated replication, in which nickase and specific DNA polymerase collaborate to drive amplification under consistent low temperature.

(B) Calculation of energy consumption and the oligo degradation decay for one 50ul liquid

reaction amplified by 10 thermal cycles PCR or iDR. PCR and iDR required 177.8 J and 2.52 J

respectively only for thermal regulation theoretically. Thermal treatment caused oligo decay rate

was calculated as 21.8‰ for 10 cycles PCR and 0.035‰ for iDR respectively.

(C) Single-stranded and double-stranded oligo amplified by iDR from a 218nt dsDNA template

were analyzed on 10% native PAGE gel. M: 20 bp DNA Ladder; C: negative control of

amplification without input template; ss: single-stranded DNA product; ds: double-stranded DNA

product.

(D) A 5' terminal FAM labeled 30nt single-stranded probe ligated to 198nt single-stranded DNA

with an inborn 5' terminal phosphate group directly from iDR amplification was analyzed and

imaged on a 12% UREA denature PAGE. M: 20 bp DNA Ladder; LP: ligation product; ss: ssDNA

product of iDR.

(E) Sequenced reads with single letter error, substitution or indel, accounted for the vast majority

in total mutant sequenced reads for both PCR (80.1%) and iDR (81.7%). Error bars represent the

mean ±s.d., where n = 3.

(F) Coverage depth distribution of sequenced reads for 10, 30 and 60 cycles PCR and iDR

amplification were positively skewed but with 0 oligo dropout. The distribution normality was

0.50 (10c), 0.40 (30c), 0.28 (60c), 0.60 (iDR) respectively.

(G) The percentage of reads with having a large coverage (the top 30% of coverage). The

percentage increased significantly from 11.31% to 34.52% as the number of PCR cycles increased

from 10 to 60 while the number of these reads stabilized at 207. Meanwhile the number of these

reads with a large coverage was 197 and these reads accounted for 7.77% of per million sequenced

reads. Oligo dropout rate calculated from random reads set with a mean 10x coverage depth was

plotted. For 10, 30 and 60 thermal cycles PCR (red line), it was counted as 4.18%, 6.41% and

13.90% respectively, and 1.33% for iDR (blue dot). Error bars represent the mean ±s.d., where n

= 10.

(H) The number of reads with mutant in a high copy number. The number increased significantly

from 177±9 to 1175±26 as the number of PCR cycles increased from 10 to 60. The number of iDR

was 46±10.

Figure 3: Multiple information reading from DNA storage.

(**A**) Illustration of successive DNA reading. Repeated PCR amplification was successively performed 10 times with aliquot of previous reaction as template.

(**B**) the oligos pool immobilized on magnetic beads was successive amplified 10 times by iDR. The same master oligo pool was used as the initial template and #1, #5 and #10 amplified oligos were analyzed.

(**C**) The proportion of reads with up to 10 substitution or indel letter error per million sequenced reads were counted as 90.65% (#1 PCR, light red), 84.26% (#5 PCR, red), 49.97% (#10 PCR, dark red), 96.35% (#1 iDR, light blue), 96.14% (#5 iDR, blue), and 94.12% (#10 iDR, dark blue), and

error reads with up to 1 error were counted as 88.18% (#1 PCR), 82.05% (#5 PCR), 48.58% (#1

PCR), 94.18% (#1 iDR), 94.04% (#5 iDR), and 92.21% (#10 iDR). Error bars represent the mean

±s.d., where n = 3.

(D) The proportion of M1G1 reads per million valid reads were, PCR (red column), 14.65% (#1

PCR), 14.82% (#5 PCR), 15.10% (#10 PCR), and iDR (blue column) 15.46% (#1 iDR), 14.55%

(#5 iDR), and 14.86% (#10 iDR). The average error reads depth of PCR (red line, 0.5 for #1, 0.22

for #5, 0.15 for #10) increased and iDR (blue), but iDR (blue line, 1.08 for #1), 1.08 for #5), 1.05

for #10) remained stable. Error bars represent the mean ±s.d., where n = 3.

(E) The distribution of reads number per each given sequence per million sequenced reads of #1,

#5, and #10 of PCR and iDR. The distribution normality was 0.5 (#1 PCR), 0.22 (#5 PCR), 0.15

(#10 PCR), 0.58 (#1, #5, #10 iDR) respectively.

(F) The dropout rate for random sequenced reads set with 10x coverage depth was plotted, PCR

(red column, 4.18% for #1, 22.89% for #5 and 53.19% for #10) and iDR (blue column, 1.86% for

#1, 2.08% for #5 and 2.09% for #10). Error bars represent the mean ±s.d., where n = 3. The

coverage depth for random sequenced reads with 1.56% dropout was calculated as 17.2 (#1 PCR),

167 (#5 PCR), 426 (#10 PCR) and 11 (#1 iDR), 12 (#5 iDR), and 12.5 (#10 iDR) respectively.

Error bars represent the mean ±s.d., where n = 10.

Figure 4. Sustainable and repeated DNA storage.

Chip-synthesized DNA oligo pool, in which digital information including image, video and text,

were stored, was repeated read by system based on PCR or iDR respectively. For current reported

DNA system with repeated reading based on PCR, 10,101,935 NGS noise reads was necessary for

perfect decoding 445KB digital files, but only 149,919 NGS noise reads were necessary for iDR.

ONLINE METHODS:

DNA Master Oligo Pool. The pool (Pool 1/2--Twist Bioscience; Pool 3--CustomArray) was resuspended in 1x TE buffer for a final concentration of 2 ng/uL. PCR was performed using Q5 High-Fidelity DNA Polymerase (NEB). Mix 10 ng of ssDNA pool (5 uL) with 2 μL of 100 μM of the forward primer and 2 μL of 100 μM of the Adaptor 2 (Adaptor 2-1), 10 μL 5x Q5 Reaction Buffer, 4 μL of 2.5 mM dNTPs, 0.5 μL Q5 High-Fidelity DNA Polymerase. Thermocycling conditions were as follows: 5 min at 98℃; 10 cycles of: 30 s at 98℃, 30 s at 56℃, 15 s at 72℃; followed by a 5 min extension at 72℃. The reaction was then purified according to the instructions in the Eastep Gel and PCR Cleanup Kit and eluted in 50 μL DNase/RNase-free water. This library was considered the master pool. All primers we used are in table S3.

iDR reaction. 10 ng of DNA oligo out of the master pool was attached to 1 μL of Streptavidin Magnetic Beads (NEB). The iDR reaction mixtures contained 1 μL of the DNA template attached to the beads (10 ng/μL), 0.25 mM dNTPs, 2.5 μL 10x NEBuffer 2, 0.08 U/μL Nt.BbvCI (NEB), 0.16 U/μL KF polymerase (exo⁻) (Vazyme), 4 μM T4 Gene 32 Protein, 0.2 mg/mL BSA, 0.5 μM Adaptor 2 (For production of ssDNA, adaptor 2 was not added.). The mixtures were incubated at 37℃ for 30 min. After amplification, the specific amplicon and the template was isolated through magnetic pull-down. In the process of repeated iDR, the template attached to magnetic beads was washed by Wash/Binding buffer (0.5 M NaCl, 20 mM Tris-HCl (pH 7.5), 1 mM EDTA) twice and mixed with the above mentioned iDR reaction mixtures except for the DNA template. The process was proceeded for 10 times. To retrieve the information, the amplified products were purified and then sequenced on one Illumina Hiseq 4000 platform with 150 paired-end cycles in Novogene (Supplementary Note 5).

PCR reaction. PCR was performed using Q5 High-Fidelity DNA Polymerase and forward

primer/adaptor 2 (10ng DNA master pool, 2 µL of forward primer (100 µM); 2 µL of adaptor 2

(100 µM)), 10 µL 5x Q5 reaction buffer in a 50 µL reaction. Thermocycling conditions were as

follows: 5 min at 98℃; 10 cycles of: 30 s at 98, 30 s at 58℃, 10 s at 72℃, followed by extension

at 72℃ for 5 min. In the repeated PCR, each subsequent PCR reaction consumed 1 µL of the prior

PCR reaction and employed 10 cycles in each 50-µL reaction. The PCR product was purified by

Eastep Gel and PCR Cleanup Kit and eluted in 50 µL DNase/RNase-free water. Then we

sequenced the PCR product on Illumina Hiseq 4000 platform.

Data availability. The BASIC code for encoding and decoding for both Linux and Windows and

bioinformatic analysis programs may be obtained via (https://github.com/xiaomingao/DNA-

information-storage). Furthermore, the original sequencing FASTQ file and the designed sequence

file   may   be   obtained   via

(http://pan.tju.edu.cn:80/#/link/627DB5C9EB819984F1183D8D4A0B72E3, Code: oZ3D).