Septins coordinate cell wall integrity and lipid metabolism in a sphingolipid-dependent process

Alexander Mela[1+] and Michelle Momany[1*]

[1]Fungal Biology Group and Plant Biology Department, University of Georgia,

2502 Miller Plant Science Building, Athens, GA 30602

[*] Corresponding Author:

Email: mmomany@uga.edu (MM)

[+] Current address:

Department of Plant & Microbial Biology, University of California, Berkeley

Keywords: Septins; MAPK Signaling; Cell wall integrity; Sterol Rich Domains

Abstract

During normal development and response to environmental stress, fungi must coordinate synthesis of the cell wall and plasma membrane. Septins, small cytoskeletal GTPases, colocalize with membrane sterol-rich regions and facilitate recruitment of cell wall synthases during dynamic wall remodeling. In this study we show that null mutants missing an *Aspergillus nidulans* core septin present in hexamers and octamers (Δ*aspA^{cdc11}*, Δ*aspB^{cdc3}*, or Δ*aspC^{cdc12}*) are sensitive to multiple cell wall-disturbing agents known to activate the cell wall integrity MAPK pathway and that this sensitivity can be remediated by osmotic support. The null mutant missing the octamer-exclusive core septin (Δ*aspD^{cdc10}*) showed similar osmotic-remedial sensitivity, but only to a single cell wall-disturbing agent and the null mutant missing the noncore septin (Δ*aspE*) showed very mild osmotic-remedial sensitivity to a different single agent. Representative core septin null mutants showed changes in cell wall polysaccharide composition, organization, and chitin synthase localization. Double mutant analysis with Δ*mpkA* suggested core septins interact with the cell wall integrity pathway. Null mutants missing any of the five septins were resistant to ergosterol-disrupting agents. The Δ*aspA^{cdc11}*, Δ*aspB^{cdc3}*, and Δ*aspC^{cdc12}* mutants showed increased sensitivity to sphingolipid-disrupting agents that was remediated by addition of exogenous phytosphingosine. Representative core septins were mislocalized after treatment with sphingolipid-disrupting agents, but not after treatment with ergosterol-disrupting agents. When challenged with both sphingolipid-disturbing and cell wall-disturbing agents in combination, remediation of the lipid defect restored proper growth to Δ*aspA^{cdc11}*, Δ*aspB^{cdc3}*, and Δ*aspC^{cdc12}*, but remediation of the cell wall defect did not. Our data suggest that the core hexamer and octamer septins are involved in cell wall integrity signaling with the noncore septin playing a minor role; that all five septins are involved in monitoring ergosterol metabolism; that the hexamer septins are required for sphingolipid metabolism; and that septins require sphingolipids to coordinate the cell wall integrity response.

Introduction

The cell wall and plasma membrane (PM) are the primary lines of defense against environmental insults

for fungal cells. With the large surface area of hyphal networks and intimate contact with the surrounding

media, fungi encounter many stresses, ranging from ion imbalance to predation. The cell wall contains

several polysaccharide constituents: chitin provides the rigid framework of the cell wall, β-glucan

maintains the shape, and mannans form the outermost, protective layer (1-6). Precise regulation of cell

wall and plasma membrane architecture is attained by tightly coordinated signaling pathways which

control genes responsible for maintaining homeostasis between the intracellular and extracellular

environments.

Septins are highly conserved small GTPase cytoskeletal proteins that function as molecular scaffolds for

dynamic cell wall and plasma membrane-remodeling, as well as diffusion barriers restricting movement

of membrane and cell wall-associated molecules (7-14). The *Saccharomyces cerevisiae* septins Cdc3,

Cdc10, Cdc11, and Cdc12 have been termed 'core septins' because they are monomers which assemble

into non-polar heterooligomers and micrometer-scale higher order structures in the form of bars, rings, or

gauzes (15-18). *A. nidulans* contains four orthologous septin proteins: AspA[Cdc11], AspB[Cdc3], AspC[Cdc12],

and AspD[Cdc10], as well as AspE, which has no *S. cerevisiae* orthologue (19). *A. nidulans* contains two

distinct subpopulations of heterooligomers; an octameric oligomer consisting of all four core septins in

the same order as proposed in *S. cerevisiae* (AspA[Cdc11]-AspC[Cdc12]-AspB[Cdc3]- AspD[Cdc10]-AspD[Cdc10]-AspB

[Cdc3]- AspC[Cdc12]- AspA[Cdc11]), and a second hexameric oligomer, with the proposed order (AspA[Cdc11]-AspC

[Cdc12]-AspB[Cdc3]- AspB[Cdc3]- AspC[Cdc12]- AspA[Cdc11])(12, 13, 20). For clarity, we will refer to AspA[Cdc11],

AspB[Cdc3], and AspC[Cdc12] as 'core hexamer septins'; AspA[Cdc11], AspB[Cdc3], AspC[Cdc12], and AspD[Cdc10] as

'core octamer septins'; and AspE as the 'noncore septin' because it does not assemble into oligomeric

structures, though it is required for higher order structure assembly at the multicellular stage (20, 21).

Previous studies in *Candida albicans* have shown that septins provide the scaffolding for cell wall

proteins via the **C**ell **W**all **I**ntegrity (CWI) MAPK signaling pathway (7, 9, 22-24). The cell wall

integrity pathway, along with the other major MAPK signaling pathways (**H**igh **O**smolarity **G**lycerol

(HOG), cAMP-PKA, **T**arget **o**f **R**apamycin (TOR), Calcineurin/Calcium, and Mating/Pheromone

response pathways) are highly conserved across the Kingdom Fungi, and have been shown to undergo

extensive cross-talk to coordinate virtually all biological functions in the cell, from expansion and

division to asexual reproduction (25-32).

Sphingolipids are long chain base-containing lipids (33) that are metabolized in a highly conserved

pathway in plants, animals, and fungi; sphingolipid metabolism shares direct connections to other major

metabolic pathways, such as sterol metabolism and fatty acid and phospholipid synthesis (34).

Sphingolipid pathway intermediates, such as phytosphingosine, have been shown to be involved in CWI

pathway signaling in *S. cerevisiae* (33, 35). **S**terol **r**ich **d**omains (SRDs), also called 'lipid rafts' or 'lipid

microdomains,' are regions of the plasma membrane enriched in specific classes of lipids, including

sterols (ergosterol, the major sterol found in most fungi (36)), sphingolipids, and phosphoinositides (37,

38), which have been shown to be functionally important for maintenance of cell polarity (39).

Membrane organization, plasticity, and overall integrity have been attributed to sphingolipid, sterol, and

glycerophospholipid interactions (40). *In vitro* work has shown that yeast septins bind directly to the

phosphoinositides PIP2, PI(4,5)P2, PI(5)P, and PI(4)P (41-44). Septins have also been shown to

genetically interact with Sur2, a sphinganine C4-hydroxylase which catalyzes the conversion of

sphinganine to phytosphingosine in sphingolipid biosynthesis (45), and to physically interact with

GTPases, Bud1 and Cdc42 to maintain sphingolipid-dependent diffusion barriers at cell membranes in

yeast (46). Interdependent colocalization of septins and sterol rich microdomains have been described in

a number of *in vivo* and *in vitro* systems (47, 48). In the dimorphic fungus *U. maydis,* septins and sterol

rich domains depend on each other to localize properly at the hyphal tip for cell signaling and cell polarity

establishment (49). In the budding yeast *S. cerevisiae*, long-chain sphingolipids have been implicated in

maintaining asymmetric distribution and mobility of some membrane-spanning molecules, such as

multidrug resistance transporters (50). In contrast to these findings, a more recent comprehensive study

of protein segregation during cell division in budding yeast found that the majority of ER and PM

proteins with transmembrane domains were actually symmetrically segregated, however septins were

shown to be responsible for partitioning of the PM-associated ER at the bud neck, thereby restricting

diffusion of ER lumen and a particular set of membrane-localized proteins (51).

Recent work has started to unravel the functional connections between the septins, cell wall integrity

MAPK pathway signaling, and lipid metabolism, however most studies have focused on a small sub-set

of septin monomers and/or were conducted in primarily yeast-type fungi (52-57). Here we show in the

filamentous fungus *A. nidulans* that the core septins, AspA^{Cdc11}, AspB^{Cdc3}, AspC^{Cdc12}, and AspD^{Cdc10} are

required for proper coordination of the cell wall integrity pathway, that all five septins are involved in

lipid metabolism, and that these roles require sphingolipids.

Results

Mutants missing core hexamer septins are hypersensitive to multiple cell wall-disturbing agents. The

mutant missing the octamer-exclusive core septin AspD^{Cdc10} is sensitive to a single agent and the

mutant missing the noncore septin AspE is mildly sensitive to a single agent. To determine whether *A.*

nidulans septins are important for cell wall integrity, we used spore dilution assays to test the growth of

septin deletion mutants on media containing a variety of known cell wall-disturbing agents. Wild type and

septin null mutants were tested on Calcofluor White (CFW) and Congo Red (CR), cell wall polymer-

intercalating agents that perturb chitin and β-1,3-glucan, respectively; Caspofungin (CASP), an inhibitor

of β-1,3-glucan synthase; and Fludioxonil (FLU), a phenylpyrrol fungicide that antagonizes the group III

histidine kinase in the osmosensing pathway and consequently affects cell wall integrity pathway

signaling (**Fig 1**)(58-67). Septin mutants Δ*aspA^{cdc11},* Δ*aspB^{cdc3}*, and Δ*aspC^{cdc12}* showed hypersensitivity to

CFW, CR, CASP, and FLU; Δ*aspD^{cdc10}* displayed no sensitivity to CFW, CR, and FLU, and sensitivity to

CASP (See **S1 Table** for list of strains used in this study). The Δ*aspE* mutant showed no sensitivity to

CFW, CASP, and FLU, and very mild sensitivity to CR. (**Fig 1A**).

Septin mutant sensitivity to cell wall-disturbing agents is remediated by osmotic stabilization. One

hallmark of cell wall integrity defects is rescue by the addition of an osmotic stabilizer, such as sorbitol or

sucrose. The addition of exogenous 1.2M sorbitol partially remediated the hypersensitivity to CASP and

fully remediated the hypersensitivity to CFW, CR, and FLU for Δ*aspA* ^{cdc11}, Δ*aspB* ^{cdc3}, and Δ*aspC* ^{cdc12}

(**Fig 1B**). Sorbitol also remediated the sensitivity of Δ*aspD*^{cdc10} to CASP and the mild sensitivity of

Δ*aspE* to CR. The osmotic remediation of growth defects in the mutants indicates that their sensitivity to

cell wall-disturbing agents is likely the result of a cell wall integrity defect.

Fig 1. Septin null mutants exhibit sensitivity to cell wall-disturbing agents, which can be remediated
by osmotic support. (Top Row) Solid media spotting assay. WT and septin null mutants Δ*aspA*^{cdc11},
Δ*aspB*^{cdc3}, Δ*aspC*^{cdc12}, Δ*aspD*^{cdc10}, and Δ*aspE* were tested for sensitivity by spotting decreasing spore
concentrations on complete medium plates with or without cell wall-disturbing agents Calcofluor White
(CFW), Congo Red (CR), Caspofungin (CASP), and Fludioxonil (FLU) at the indicated final
concentrations. (Bottom Row) WT and septin null mutants were tested for osmotic remediation of
hypersensitivity to cell wall-disturbing agents, by spotting decreasing spore concentrations on media

amended with exogenous 1.2M sorbitol. Spore concentrations were 10^7 conidia/mL – 10^4 conidia/mL for

all assays. Differences in colony color result from changes in spore production, spore pigment, and

production of secondary metabolites under stress. N≥5

Core septin null mutant ∆aspB^{cdc3} has altered chitin and β-1,3-glucan localization. Because the core

septin mutants showed sensitivity to cell wall-disturbing agents consistent with action via the cell wall

integrity pathway, we predicted that there might be differences in cell wall polymer localization in the

mutants. To examine cell wall polymer localization, we did live-cell imaging of WT and ∆aspB^{cdc3} since

it is a member of both hexamers and octamers. Conidia were incubated on coverslips in liquid media,

stained with CFW or aniline blue to observe chitin and β-1,3-glucan, respectively, and immediately

observed by fluorescence microscopy. Z-stack images were analyzed one-by-one, compressed into

maximum projection images to visualize the fluorescence signal in the entire hyphal structure, and by line

scans of aniline blue and calcofluor white staining patterns (**Fig 2, S1 Fig**). As previously reported,

∆aspB^{cdc3} showed more presumptive branch initials per hyphal compartment than WT (21). The aniline

blue staining showed a reduction of labeling at all ∆aspB^{cdc3} hyphal tips. Smaller, less intense aniline blue

labeling also occurred at presumptive branch initials (**Fig 2A, S1A**). The CFW staining of ∆aspB^{cdc3}

showed a shift of the tip band of staining closer to the hyphal apex and a less well-defined endocytic

collar zone (**Fig 2B, S1B**). These data show that cell wall organization is altered in ∆aspB^{cdc3} and raise

the possibility that it might be altered in other core hexamer septin null mutants as well.

Core septin null mutant ∆aspB^{cdc3}has higher levels of chitin. Previous studies have shown that

perturbations to one cell wall component often trigger compensatory changes to others via the cell wall

integrity pathway (28, 68, 69). To analyze the cell wall composition of septin mutants, two independent

biological replicates of a diagnostic glycosyl linkage analysis (70) were conducted to quantify the cell

wall polysaccharide composition of WT, ∆aspB^{cdc3} (a core septin null mutant which showed

hypersensitivity to cell wall-disturbing agents), and ∆aspE (non-core septin mutant)(**S2 Fig**). All samples

contained 3- and 4-linked glucose as the primary cell wall components, as well as 4-linked N-

acetylglucosamine (the monomer of chitin) and a relatively minor amount of mannan. The septin mutants

Δ*aspB*^{cdc3} and Δ*aspE* showed 7% and 2% increases in the average area of the detected linkage peak for 4-

GlcNAc compared to WT, respectively; the amount of 4-Glc and of mannan (data not shown) did not

show significant differences between samples. These data indicate an increase in chitin content of the

Δ*aspB*^{cdc3} mutant and raise the possibility that it might be altered in other core hexamer septin null

mutants as well.

163
164
Fig 2. Cell wall staining reveals unique patterns of chitin and β-1,3-glucan deposition in the core
septin hexamer null mutant Δ*aspB*^{cdc3}. (A) WT and Δ*aspB*^{cdc3} cells were incubated for approximately
14h and (A) stained with aniline blue to visualize β-1,3-glucan or (B) stained with Calcofluor White
(CFW) to visualize chitin. Representative images are shown from at least three independent biological
replicates, with ≥100 cells observed each. Magenta arrows in panel A denote hyphal tips with
representative β-1,3-glucan deposition. Green arrows in panel B denote hyphal tips with representative
chitin deposition. Insets show enlargement of area around the arrows. Scale bars = 10μm. N≥3
172

Chitin synthase localization is altered in the core septin null mutant Δ*aspA* [cdc11]. Membrane-spanning

cell wall synthases are the ultimate effectors of the cell wall integrity pathway. *A. nidulans* contains six

genes for chitin synthases: *chsA, chsB, chsC, chsD, csmA*, and *csmB*. Chitin synthase B localizes to sites

of polarized growth in hyphal tips, as well as developing septa in vegetative hyphae and conidiophores, a

pattern very similar to septin localization. Deletion of chitin synthase B shows severe defects in most

filamentous fungi analyzed thus far, and repression of the chitin synthase B gene expression in *chsA,*

chsC, and *chsD* double mutants exacerbated growth defects from a number of developmental states

observed in each single mutant, suggesting it plays a major role in chitin synthesis at most growth stages

(22). For these reasons, we chose chitin synthase B as a candidate to observe in a septin mutant

background for possible defects in localization. We hypothesized that recruitment or maintenance of cell

wall synthases, such as chitin synthase could be disrupted in septin null mutants which showed sensitivity

to cell wall-disturbing agents and altered cell wall staining patterns. To determine the localization of

synthases, a chitin synthase B-GFP (*chsB-GFP*) strain was crossed with strains in which core hexamer

septins were deleted. After repeated attempts, the only successful cross was with core hexamer deletion

strain Δ*aspA* [cdc11]. The Δ*aspA* [cdc11], *chsB-GFP* strain showed conspicuous differences in chitin synthase

localization patterns compared to WT (**Fig 3 and Fig. S3**). In WT, the chitin synthase-GFP signal was at

the tips of >90% of branches, presumptive branch initials, and apical hyphal tips (SFig 3A). In the septin

null mutant Δ*aspA* [cdc11], GFP signal was absent in most presumptive branch initials, but present in longer

side branches that were at least 10 µm long and in the apical hyphal tip(s) (**Fig 3 and Fig. S3A**).

Intriguingly, the chitin label calcofluor white was more intense in the septin deletion strain consistent with

the increase in 4-GlcNAc levels seen in polysaccharide analysis (Fig. S3B). Calcofluor white labeling

also showed that the polymer chitin was present throughout the hyphae in both the WT and septin

deletion strains (Fig. 2) suggesting that chitin synthases had been active in both, but perhaps were not

retained as long in the presumptive branch initials of the septin deletion mutant. These data support the

hypothesis that core hexamer septins play a role in recruitment and/or retention of cell wall synthases to

the proper locations at the plasma membrane during growth and development.

Fig 3. Chitin synthase is mislocalized in core septin null mutant Δ*aspA* ^{cdc11}. (A) Chitin synthase B-GFP in WT background (Columns 1-2). (B) chitin synthase B-GFP in a Δ*aspA* ^{cdc11} mutant background. Calcofluor White labeling shows the presence of the polymer chitin at septa, main hyphal tips, branches, and putative branch initials (Column 3). Representative images are shown from at least three independent biological replicates, with ≥100 cells observed. White arrows highlight hyphal branches and tips. White arrow heads highlight putative branch initials. Scale bars = 10μm. N≥3.

Double mutant analyses suggest the core septin $aspB^{cdc3}$ ***modulates the cell wall integrity pathway in***

the ***Δmpk A background under cell wall stress.*** To determine whether there are genetic interactions

between the septins and the cell wall integrity pathway kinases, double mutants were generated by sexual

crosses and analyzed. The first cell wall integrity pathway kinase, PkcA^{Pkc1} (ANID_00106), is essential in

A. nidulans, and so null alleles could not be utilized for double mutant analysis. A null allele of the final

kinase in the CWI MAPK signaling cascade, MpkA^{Slt2} (ANID_11786), was crossed with $\Delta aspB^{cdc3}$ and

$\Delta aspE$ and progeny were challenged with cell wall-disturbing treatments. If the septins are directly in the

CWI pathway, we would expect double mutants to show the same phenotypes as the parental null mutant

that acts earliest. If the septins are in a parallel pathway or alternate node which also affects cell wall

integrity, we would expect a novel/synergistic phenotype in the double mutants.

Spore dilution assays were conducted, challenging the double mutants and the parental strains with cell

wall-disturbing treatments. The double mutants $\Delta aspB^{cdc3}\Delta mpkA^{slt2}$ and $\Delta aspE\,\Delta mpkA^{slt2}$ displayed a

colony-level radial growth defect and reduced conidiation which phenocopied the $\Delta mpkA^{slt2}$ single

mutant. When challenged with low concentrations of CASP and CFW, the $\Delta aspB^{cdc3}\Delta mpkA^{slt2}$ and

$\Delta aspE\,\Delta mpkA^{slt2}$ mutants were more sensitive than $\Delta aspB^{cdc3}$ and $\Delta aspE$ single mutants, but suppressed

the colony growth defects of $\Delta mpkA^{slt2}$. The novel phenotype of the double mutants shows that septins are

involved in cell wall integrity and raises the possibility that they act in a bypass or parallel node for

remediation of cell wall defects **(Fig 4)**.

Fig 4. Double mutant analyses suggest core septins modulate the cell wall integrity pathway. (A) Simplified schematic diagram of the *A. nidulans* cell wall integrity MAPK signaling pathway (71). (B) Solid media spotting assays. (Top) WT, $\Delta aspB^{cdc3}$, $\Delta mpkA^{slt2}$, and two $\Delta aspB^{cdc3}\Delta mpkA^{slt2}$ double mutant strains, were tested for sensitivity by spotting decreasing spore concentrations on complete media plates with or without cell wall-disturbing agents. (Bottom) WT, $\Delta aspE$, $\Delta mpkA^{slt2}$, and two $\Delta aspE\Delta mpkA^{slt2}$ double mutant strains were tested for sensitivity by spotting decreasing spore concentrations on complete media plates with or without cell wall-disturbing agents. Differences in colony color result from changes in spore production, spore pigment, and production of secondary metabolites under stress. Transcription Factors, (TFs); Calcofluor White, (CFW); Caspofungin (CASP). Spore concentrations were [10^6 conidia/mL – 10^3 conidia/mL] for all assays in figure. N=3

Core septin null mutants are insensitive to treatments which disrupt the Ca2+/Calcineurin, cAMP-

PKA, or TOR Pathways. Noncore septin null mutant ∆aspE is sensitive to some TOR pathway

inhibitors. One possible explanation for the observed sensitivity to cell wall-disturbing agents could be

that septins are involved in 'cross-talk' with other MAPK signaling pathways that have been shown to

interact with cell wall integrity pathway signaling such as the Calcium/Calcineurin (CAMK) signaling

pathway. To test this possibility, calcium chloride, EGTA (calcium chelating agent), and FK-506

(calcineurin inhibitor) were added to each treatment (**S4 Fig**) (72-74). The treatments showed no obvious

colony growth defects, suggesting that Ca2+/Calcineurin signaling pathway crosstalk does not

significantly contribute to the observed sensitivity of septin null mutants to cell wall-disturbing agents.

Another pathway closely associated with cell wall integrity, lipid biosynthesis, and lipid signaling is the

TOR MAPK signaling pathway (75). To test the involvement of septins in this signaling pathway, septin

null mutants were challenged with rapamycin (a potent inhibitor of the TOR pathway), as well as

methylxanthine derivatives and TOR pathway inhibitors, caffeine and theophylline (76-78). There were

no observable growth defects in the presence of rapamycin (final concentration was 600ng/mL, which is

approximately 3-fold higher than inhibitory concentration for known TOR pathway mutants) in the septin

null mutants compared to WT (**S4 Fig**) (79, 80). Caffeine and theophylline have been shown to interfere

with phosphodiesterase activity in the cAMP-PKA and TOR pathways, and $\Delta pkaA^{tpk1}$ and $\Delta tor1^{torA}$

mutants show hypersensitivity to caffeine treatment which cannot be remediated by exogenous sorbitol

(58, 78, 81-83). If the septins were involved in crosstalk between the cAMP-PKA or TOR pathways and

the CWI pathway, we would predict that septin null mutants would be hypersensitive to caffeine and

theophylline treatments. To our surprise, only $\Delta aspE$ showed hypersensitivity to both caffeine and

theophylline, and the hypersensitivity was not remediated by an osmotic stabilizer (**S4 Fig and data not**

shown). These data suggest that TOR, cAMP-PKA, and Calcium/Calcineurin signaling pathways do not

contribute to the cell wall sensitivity or plasma membrane resistance phenotypes in the core septin null

mutants, but the cAMP-PKA or TOR pathways may contribute to the phenotypes of the septin $\Delta aspE$ null

mutant.

Core septin null mutants show increased resistance to disruption of ergosterol biosynthesis and $\Delta aspA$

^{cdc11} and $\Delta aspE$ show increased resistance to the ergosterol-binding drug Natamycin. We hypothesized

that septins could indirectly modulate MAPK pathways, particularly the CWI pathway, through

interactions with plasma membrane lipids within sterol rich domains (SRDs). Recent work has shown

that the cell wall integrity pathway in *S. cerevisiae* is regulated by sphingolipids and ergosterol,

facilitating proper deposition of cell wall polymers at actively growing regions and sites of septation (38,

55). It is well-established that septins localize to sites of polarized growth and septation, where highly

dynamic remodeling of the plasma membrane and cell wall via membrane-bound synthases and polarity-

associated proteins takes place (84). These highly dynamic plasma membrane SRDs in fungi often

contain ergosterol and sphingolipids. To determine whether loss of septins might affect one of the major

SRD-associated lipids, each septin mutant was challenged by drug treatments disrupting the ergosterol

biosynthesis pathway in spore dilution assays (**Fig 5**). These assays included an azole treatment

(Itraconazole), an allylamine treatment (Terbinafine), and a polyene treatment (Natamycin), which each

impact a different step in ergosterol biosynthesis (**Fig 5A**). Allylamines (Terbinafine) inhibit the

conversion of squalene to squalene epoxide, azoles (Itraconazole) inhibit the conversion of lanosterol to

4,4-dimethylcholesta-8,14,24-trienol, and polyenes bind directly to ergosterol (85).

The $\Delta aspA^{cdc11}$, $\Delta aspB^{cdc3}$, $\Delta aspC^{cdc12}$, and $\Delta aspD^{cdc10}$ mutants were more resistant to Itraconazole and

Terbinafine than WT or $\Delta aspE$. Only $\Delta aspA^{cdc11}$ and $\Delta aspE$ showed strong resistance to Natamycin

treatment (**Fig 5B**). The addition of exogenous 50μM ergosterol was able to fully remediate the

sensitivity of all null mutant strains and WT to Itraconazole, Terbinafine, and Natamycin, suggesting

ergosterol was indeed the primary lipid component disrupted by these treatments (**Fig 5C**). These data

suggest that all five septins are involved in monitoring ergosterol metabolism and/or deposition.

Fig 5. Core septin null mutants show increased resistance to disruption of ergosterol biosynthesis and Δ*aspA* ^{cdc11} and Δ*asp*E show increased resistance to the ergosterol-binding polyene drug Natamycin. (A) A simplified schematic diagram of the ergosterol biosynthesis pathway, showing where allylamines, azoles, and polyene antifungal agents affect each step (85). (B) WT and septin null mutants Δ*aspA* ^{cdc11}, Δ*aspB* ^{cdc3}, Δ*aspC* ^{cdc12}, Δ*aspD* ^{cdc10}, and Δ*aspE*, were tested for sensitivity by spotting decreasing spore concentrations on solid media with or without the ergosterol biosynthesis-disturbing agents Terbinafine, Itraconazole, and Natamycin at concentrations shown. (C) Remediation of sensitivity to ergosterol biosynthesis-inhibiting treatments. WT and septin null mutants were tested for the ability of exogenous ergosterol (50μM) to remediate sensitivity to ergosterol biosynthesis-disturbing agents. Differences in colony color result from changes in spore production, spore pigment, and production of secondary metabolites under stress. Spore concentrations were [10^7 conidia/mL – 10^4 conidia/mL] for all assays in figure. N=3

Core hexamer septin null mutants show altered sensitivity to disruption of sphingolipid biosynthesis.

Sphingolipids are a class of plasma membrane lipids which has been shown to be associated with sterol-

rich domains, along with sterols and phosphoinositides, and therefore are possible targets for septin-

mediated interactions at the membrane. To determine whether loss of septins impacts sphingolipid

metabolism in *A. nidulans*, the septin null mutants were challenged with sphingolipid biosynthesis-

disrupting agents Myriocin and Aureobasidin A (AbA) (**Fig 6**). Myriocin disrupts the first committed

step of the biosynthetic pathway, converting palmitoyl-coa and serine to 3-ketodihydrosphingosine,

preventing the accumulation of downstream intermediates, such as ceramides and sphingoid bases like

phytosphingosine (PHS), as well as complex sphingolipids at the final steps of the pathway (**Fig 6A**) (37,

47). Aureobasidin A inhibits IPC synthase, disrupting the conversion of inositolphosphorylceramide

from phytoceramide, and consequently causing the accumulation of intermediates such as

phytosphingosine, which has been shown to be toxic at high concentrations (**Fig 6A**) (86).

The core hexamer null mutants $\Delta aspA^{cdc11}$, $\Delta aspB^{cdc3}$, and $\Delta aspC^{cdc12}$ were more sensitive to Myriocin

than the other septin null mutants or WT (**Fig 6B**). $\Delta aspA^{cdc11}$, $\Delta aspB^{cdc3}$, and $\Delta aspC^{cdc12}$ were also more

resistant to AbA than null mutants of the octamer-exclusive septin $\Delta aspD^{cdc10}$, the noncore septin $\Delta aspE$,

or WT (**Fig 6B**). Strikingly, the addition of exogenous PHS (15µM) to the Myriocin treatment fully

remediated the sensitivity of $\Delta aspA^{cdc11}$, $\Delta aspB^{cdc3}$, and $\Delta aspC^{cdc12}$ (**Fig 6C**). $\Delta aspA^{cdc11}$, $\Delta aspB^{cdc3}$, and

$\Delta aspC^{cdc12}$ were also more resistant to higher concentrations (20µM) of the phytosphingosine

intermediate (PHS), which has been shown to be toxic at high concentrations (**Fig 6C**).

To address whether cell wall and plasma membrane defects might be associated with one another in

septin null mutants, a combinatory drug treatment approach was taken. Sublethal concentrations of CFW

(2µg/mL) and Myriocin (10µg/mL), in which all strains grew at every spore concentration, were

combined (**Fig 6D**). When combined, the two drugs resulted in additive, colony-level growth defects for

$\Delta aspA^{cdc11}$, $\Delta aspB^{cdc3}$, and $\Delta aspC^{cdc12}$. This sensitivity to both drugs in combination was remediated by

the addition of exogenous PHS. Surprisingly the addition of exogenous sorbitol, which had fully

remediated the hypersensitivity of the septin null mutants to all previously tested cell wall-disturbing

agents, resulted in a more dramatic growth defect in media containing only Myriocin, or in combination

with CFW (**Fig 6C and D**). These data together suggest that the core hexamer septins AspA[Cdc11],

AspB[Cdc3], and AspC [Cdc12] may monitor sphingolipid metabolism. They further suggest that the hexamer

septins may signal sphingolipid status to the cell wall integrity pathway, and that this signal is required

for proper cell wall integrity function.

Fig 6. Septin core hexamer null mutants show altered sensitivity to agents which disrupt sphingolipid biosynthesis. (A) A simplified diagram of the sphingolipid biosynthesis pathway in *A. nidulans* (87). (B) *Solid media spotting assay.* WT and septin null mutants, $\Delta aspA^{cdc11}$, $\Delta aspB^{cdc3}$, $\Delta aspC^{cdc12}$, $\Delta aspD^{cdc10}$, and $\Delta aspE$ were tested for sensitivity by spotting decreasing spore concentrations on solid media with or without sphingolipid biosynthesis-disturbing agents Myriocin and Aureobasidin A. (C) *Remediation of sensitivity to sphingolipid biosynthesis-disturbing agents.* WT and septin null mutants were tested for remediation of sensitivity to sphingolipid biosynthesis-disturbing agents by spotting decreasing spore concentrations on solid media amended with exogenous phytosphingosine (PHS) intermediate (15μM) or 1.2M sorbitol, with or without Myriocin. (D) Combinatory treatment with cell wall and sphingolipid biosynthesis-disturbing agents. WT and septin null mutants were tested for sensitivity to cell wall and sphingolipid biosynthesis-disturbing agents in combination, by spotting decreasing spore concentrations on solid media with or without 'sub-lethal' concentrations of Calcofluor White (CFW), Myriocin, or CFW + Myriocin, amended with either exogenous phytosphingosine intermediate (15uM) or 1.2M sorbitol. Differences in colony color result from changes in spore production, spore pigment, and production of secondary metabolites under stress. Spore concentrations were [10^7 conidia/mL – 10^4 conidia/mL] for all assays in figure. N=3

Localization of core hexamer septins AspA^{Cdc11} and AspB^{Cdc3} is disrupted by sphingolipid inhibitors.

We predicted that since core septin hexamer null mutants $\Delta aspA^{cdc11}$, $\Delta aspB^{cdc3}$, and $\Delta aspC^{cdc12}$ were

sensitive to drugs which inhibit sphingolipid biosynthesis the localization of these septins might be

altered when exposed to the same treatments. We examined whether disruption of sphingolipid

biosynthesis causes changes in septin localization using live-cell imaging with fluorescence microscopy.

A strain carrying AspA^{Cdc11}-GFP was grown in liquid complete medium overnight, treated with

exogenous PHS (15µM), and imaged for 3 hours post-treatment. There was a dynamic shift in septin

localization under PHS treatment over the course of the experiment, compared to the vehicle control. The

septin-GFP signal shifted from a relatively homogenous cortical localization along the hyphal tips to a

more stochastic, punctate localization along the entire length of hyphae (**Fig 7, middle panel**).

In contrast to these results, treatments with the ergosterol-binding polyene, Filipin III (25µg/mL) and

phosphatidylinositol 3-kinase inhibitor, Wortmannin (20µg/mL), did not affect the localization pattern of

septins as dramatically as the PHS treatment (**Fig 7, left- and right-most panels**). Similar patterns of

aberrant localization were observed under PHS treatments for AspB^{Cdc3}-GFP (**S5 Fig**). Myriocin

treatment also resulted in a similar pattern of mislocalization in AspA^{Cdc11}-GFP cells when compared to

treatment with PHS (**S6 Fig**). These results suggest sphingolipid content and/or distribution within the

plasma membrane contributes to the localization and stability of core hexamer septins at the plasma

membrane and that ergosterol and phosphoinositides may not be as vital for this process.

Fig 7. Septin AspA^{Cdc11}-GFP localization disrupted by sphingolipid biosynthesis inhibitors.
AspA^{Cdc11}-GFP strain incubated in liquid media for approximately 16h and imaged 180 minutes after
replacing with fresh media containing sterol, sphingolipid, and phosphoinositide-disturbing agents,
Filipin, phytosphingosine, and Wortmannin, respectively. Representative images are shown from three
independent biological replicates with ≥100 cells observed. (Top Row) Vehicle controls for each
treatment. (Bottom Row) Filipin (25µg/mL), phytosphingosine (15µM), and Wortmannin (20µg/mL)
treatments respectively. Insets show enlarged section of micrographs from each picture to better visualize
pattern of fluorescence signal. Imaging conducted with Deltavision I deconvolution inverted fluorescence
microscope. White arrows denote hyphae which are highlighted in enlarged images. Scale bars = 10µm.
N=3

Discussion

Our data show that *A. nidulans* septins play roles in both plasma membrane and cell wall integrity and

that distinct subgroups of septins carry out these roles. Previous work has shown that the five septins of *A.*

nidulans septins form hexamers (AspA^{Cdc11}, AspB^{Cdc3}, and AspC^{Cdc12}) and octamers (AspA^{Cdc11},

AspB^{Cdc3}, AspC^{Cdc12}, and AspD^{Cdc10}) and that the noncore septin AspE does not appear to be a stable

member of a heteropolymer (20). The current work suggests that though all septins are involved in

coordinating cell wall and membrane integrity, the roles of hexamers, octamers, and the noncore septin

are somewhat different. Core hexamer septins appear to be most important for sphingolipid metabolism,

all five septins appear to be involved in ergosterol metabolism, and core septins are most important for

cell wall integrity pathway with the noncore septin possibly playing a minor role. As summarized in

Figure 8 and discussed in more detail below, our previous and current data are consistent with a model in

which: (A) All five septins assemble at sites of membrane and cell wall remodeling in a sphingolipid-

dependent process; (B) All five septins recruit and/or scaffold ergosterol and the core hexamer septins

recruit and/or scaffold sphingolipids and associated sensors at these sites, triggering changes in lipid

metabolism; and (C) The core septins recruit and/or scaffold cell wall integrity machinery to the proper

locations and trigger changes in cell wall synthesis. The noncore septin might play a minor role in this

process.

Septins assemble at sites of membrane and cell wall remodeling in a sphingolipid-dependent process.

We hypothesize that sterol rich domain-associated lipids (ergosterol, sphingolipids, and

phosphoinositides) recruit or facilitate binding and assembly of septins on the membrane, with

sphingolipids contributing more to the stabilization of septins than other SRD-associated lipids in *A.*

nidulans (**Fig 8A**). Previous studies showed preferential *in vitro* binding of yeast core septin orthologues

to the phosphoinositides PIP2, PI(4,5)P2, PI(5)P, and PI(4)P, however our treatments with Wortmannin

and Filipin III, known disruptors of phosphoinositides and ergosterol respectively (88-92), did not affect

septin localization as dramatically as sphingolipid-disturbing treatments (**Fig 7**,**S3, and S4 Fig**). The

marked septin mislocalization we observed upon treatment with phytosphingosine and Myriocin strongly

supports the idea that sphingolipids at the membrane contribute to core septin localization and help

maintain septin assemblies at the proper locations. Previous studies on composition of lipid

411 microdomains showed that relatively minor differences in sphingolipid structure can have significant

412 effects on sterol and phospholipid interactions, consequently resulting in major changes in membrane

413 properties (93, 94). Perhaps sphingolipids (and ergosterol to a lesser extent) help to stabilize SRDs in a

414 way that facilitates assembly of septin filaments and higher order structures via diffusion, collision, and

415 annealing as proposed by Bridges et al (2014) (41-43, 95). Consistent with this idea, *A. nidulans* AspB

416 filaments have been shown to move along the plasma membrane, break apart, and 'snap' together in a

417 way that suggests collision and annealing (21).

418 ***Septins recruit and/or scaffold lipids and associated sensors, triggering changes in lipid metabolism***.

419 Consistent with a role for septins in modulating membrane composition, all septin null mutants were

420 resistant to ergosterol-disrupting treatments and core hexamer septin mutants were affected by disruption

421 of sphingolipid biosynthesis (**Fig 5A-B, Fig 6**). Based on proposed mechanisms for septins as diffusion

422 barriers or organizational scaffolds of membrane-associated proteins in yeast and smut fungi, we propose

423 septins monitor lipid microdomain composition and/or organization in filamentous ascomycetes (49,

424 51)(**Fig 8B**). Septins do not have a transmembrane domain, a feature that often defines established

425 membrane 'sensors' that monitor local lipid environments (96, 97); however, septins share a highly

426 conserved polybasic domain proposed to facilitate septin-membrane interactions (44, 98). In addition to

427 the polybasic domain, septins have recently been shown to contain an amphipathic helix motif which has

428 been implicated in septin sensing of membrane curvatures (99). Given that septins have been shown to

429 assemble into non-polar higher order structures along the plasma membrane (95), septin assembly itself

430 might be the mechanism by which lipid composition and protein organization is monitored at the

431 cytosolic face of membranes. Perhaps septin assemblies that pass specific size or geometric thresholds

432 trigger signaling through MAPK and other pathways.

Figure 8. Model for septin modulation of cell wall and plasma membrane integrity through
interactions with sterol rich domains (SRDs). As described in the text, our data suggest that all five
septins are involved in cell wall and membrane integrity coordination. The core septins that participate in
hexamers appear to be most important for sphingolipid metabolism while all septins appear to be involved
in ergosterol metabolism and cell wall integrity. Because SRDs contain both sphingolipids and ergosterol
and because it is not yet clear how subgroups of septins interact with each other at SRDs, we show all
core septins in our model without distinguishing hexamers and octamers. In this model, septins are
proposed to colocalize with SRDs in a manner which promotes: (A) assembly of septins into higher order
structures along the membrane at sites of polarized growth or cell wall/PM remodeling; (B) the
recruitment and/or scaffolding of lipids and associated membrane-bound sensors to monitor membrane
composition and/or stress. The status of membrane composition is relayed to the nucleus where it triggers
changes in expression of genes responsible for lipid metabolism; and (C) the recruitment and/or
scaffolding of cell wall integrity pathway machinery to monitor cell wall composition and stress, followed
by the recruitment and/or scaffolding of cell wall synthases, possibly with the help of other septin-
interacting proteins such as formins. Question marks (**?**) denote speculative processes or interactions
which have not been characterized.
451
452

Core septins recruit and/or scaffold cell wall integrity machinery to the proper locations and trigger

changes in cell wall synthesis. In addition to monitoring and relaying information about the membrane,

septins clearly have a role in normal cell wall growth and remediation of cell wall stress via the cell wall

integrity pathway. We propose a major role of the core septins is to recruit and/or organize integral

proteins to sites of polarized growth or remodeling at the cell cortex to ensure cell wall integrity pathway

functions are carried out (**Fig 8C**). The sensitivity to cell wall-disturbing agents, altered cell wall

composition, and altered polysaccharide deposition in the core septin null mutants (**Fig 1-2** and **S1-S2**

Fig) are consistent with phenotypes of cell wall integrity pathway mutants in previous studies (100-102).

Our glycosyl linkage analysis showed that cell wall chitin content is increased in the septin mutants

compared to WT (**S2 Fig**). Hyper-synthesis of chitin has been shown to occur during cell wall stress via

the cell wall integrity pathway in *S. cerevisiae* and *Candida albicans* (103-110). Double mutant analyses

between septins and CWI pathway kinases also support a role for core septins in maintaining cell wall

integrity under stress (**Fig 4**). Suppression of cell wall defects under cell wall stress by deletion of septins

in an Δ*mpkA* ^{slt2} background suggests a parallel node by which septins negatively regulate cell wall

integrity pathway sensors or kinases could exist. This interpretation is consistent with studies in yeast

showing Bni4 (ANID_00979), a formin which is phosphorylated by and functions downstream of MAPK

$Slt2^{mpkA}$, directly interacts with the core septin orthologues in order to recruit chitin synthases to the bud

neck (111-113). Though the involvement of AspE in the cell wall integrity pathway appears to be minor

based on the very mild sensitivity of the null mutant to Congo Red (**Fig 1**), sensitivity to TOR and cAMP-

PKA pathway inhibitors suggests that AspE might participate indirectly through MAPK pathway cross-

talk (**Fig S4**). It could be that in the septin null mutants, septin filaments are not able to properly

assemble and monitor lipid composition and protein organization at the cytosolic face of membranes,

resulting in defects in plasma membrane and cell wall integrity (Fig 8).

Though we have discussed membrane and cell wall integrity separately, it is possible that that membrane

defects in the septin null mutants contribute to the observed cell wall changes or that cell wall defects

contribute to the observed changes in lipid metabolism. When septin deletion mutants were challenged

with both membrane-disturbing and cell wall-disturbing agents in combination, remediation of the lipid

defect (via PHS) restored proper growth, but remediation of the cell wall defect (via sorbitol) did not

remediate lethality (**Fig 6C-D**). This suggests that there is a synergistic effect of disrupting sphingolipid

metabolism and cell wall architecture in septin null mutants and that septin-sphingolipid interactions are

required for roles in maintaining cell wall integrity.

Because *A. nidulans* septin deletion mutants are viable, we were able to systematically analyze the roles

of all septins in this organism. Our data show that all septins are required for proper coordination of lipid

metabolism, with core hexamer septins most important for sphingolipid metabolism and all septins

involved in ergosterol metabolism. Our data also show that the core septins are most important for cell

wall integrity pathway, but that these roles require sphingolipids. Based on our data we propose that

septins are critical for tight coordination of plasma membrane metabolism and cell wall synthesis during

normal development and response to exogenous stress and that the site of this coordination is likely

SRDs. Future work will address how subgroups of septins interact with each other and with sphingolipids

and ergosterol in SRDs to coordinate the cell wall integrity response.

Methods

Spore Dilution Drug Sensitivity Assays

Strains used in this study are listed in **S1 Table**. Media used were previously described (114).

Aspergillus nidulans strains were harvested in ddH20, spores normalized to $1X10^{7}$ or $1X10^{6}$

conidia/mL, and then serial diluted 4-fold into separate Eppendorf tubes. All strains were inoculated in

10uL droplets in a grid pattern on petri plates containing 25mL of solid (1.8% agar) complete medium

(CM) (1% glucose, 0.2% peptone, 0.1% Casamino Acids, 0.1% yeast extract, trace elements, nitrate salts,

and 0.01% vitamins, pH 6.5; with amino acid supplements as noted) or minimal medium (MM)(1%

glucose, trace elements, 1% thiamine,0.05% biotin, pH 6.5; with amino acid supplements as noted) with

or without amended supplements. All incubations were conducted at 30°C as indicated for 3-4 days

before images were taken. Sorbitol, NaCl, and KCl were added at 2M, 1M, and 1.5M respectively to

media before autoclaving. Stock solutions were prepared as follows: Blankophor BBH/Calcofluor White

(Bayer Corporation; standard-SV-2460; 25mg/mL in ddH20 and adjusted pH with 1M KOH until

solubilized), Congo Red (Fisher Scientific; Lot No.8232-6; 10mg/mL in ddH20), Caspofungin acetate

(1mg/mL in ddH20), Fludioxonil/Pestanal (2mg/mL in DMSO), and caffeine monohydrate (63.66mg/mL

in ddH20 and gently heated with stirring until solubilized); calcium chloride (2M in ddH20), EGTA

(0.5M in ddH20), Rapamycin/Sirolimus (1mg/mL in acetonitrile), FK-506 (5mg/mL in DMSO),

Natamycin (1.5mg/mL in MeOH), Itraconazole (10mg/mL in DMSO), Terbinafine (10mg/mL in DMSO).

Stock solutions of Myriocin (5mg/mL), phytosphingosine (1mg/mL), and Aureobasidin A (1mg/mL)

were prepared in DMSO, EtOH, and MeOH respectively and stored at -20C in the dark. Images of plates

were captured using a cellular device with an 8.0 Megapixel camera, and subsequently processed using

Photoshop CS5 Version 12.0 X32.

Generating double mutants and other strains by crossing

Parental strains were coinoculated in CM liquid at 1×10^5 conidia/mL supplemented with all parental

auxotrophic markers and allowed to incubate at 30°C for up to one week or until a thick mycelial mat had

formed. Mycelial mats were transferred to solid MM plates containing only shared auxotrophic markers

from the genetic backgrounds of each parental strain.   Plates containing mycelial mats were parafilmed

and incubated in the dark at room temperature for up to 2 weeks or until mature cleistothecia form on the

mycelial mats. Multiple cleistothecia from each genetic cross were collected in water, diluted, and plated

onto solid media containing all auxotrophic supplements from each parental strain to allow growth of all

resulting progeny. Approximately 50 progenies were collected from each dissected cleistothecium, and

each colony was transferred to master plates, and replica-plated onto minimal media without any

supplements, to isolate prototrophic progeny. Five to ten progenies from each cross were then 3-phased

streaked to obtain single colony isolates for PCR verification. All progenies of genetic crosses were

verified by diagnostic PCR using KOD XTREME Hot Start DNA Polymerase (71975-3, EMD Millipore)

or OneTaq® Hot Start Quick-Load® 2X Master Mix with Standard Buffer (M0488L, New England

BioLabs inc.) according to manufacturer's instructions. Double mutant strains of *ΔmpkA^slt2^* were verified

for deletion of *mpkA^slt2^* by amplification of entire gene using primers, MpkA-806-F' and MpkA-3779-R',

followed by SacI HF restriction enzyme digestion of PCR product to better visualize band sizes. All

progeny from chitin synthase and septin null mutant crosses were determined to be virtually identical to

each septin mutant parental strain in growth/morphology on a colony level and by microscopy, PCR-

verified for the deletion of each septin gene, and screened visually by fluorescence microscopy for the

presence of chitin synthase-GFP signal. Chitin synthase-GFP strains in a *ΔaspA^cdc11^* genetic background

were verified for deletion of *aspA^cdc11^* by amplification of the entire gene using primers, AspA-KO-F' and

AspA-KO-R', followed by XhoI restriction enzyme digestion for verification of band sizes. Strains and

primer sets used in this study can be found in S1 and S2 Tables respectively.

Growth conditions and microscopy

Growth and preparation of cells were as previously reported (115). Conidia were inoculated on sterile

coverslips in 10mL liquid complete or minimal media at $1X10^5$ conidia/mL and incubated at 30C in a small

petri dish for the specified amount of time. Cell walls were stained for chitin with Blankophor BBH (CFW)

(American Cyanamid, Wayne, NJ; 25mg/mL stock solution in ddH2O and pH adjusted by 1M KOH until

solubilized; working solution made by diluting stock solution by 100X and 8ul dissolved in 5mL ddH20

prepared fresh for working solution and used immediately), β-(1,3)-glucans were stained with aniline blue

(stock solution prepared fresh to 10 mg/mL final concentration in ddH20; working solution prepared at

135.55μM in 50mM phosphate buffer, pH adjusted to 9.5 with 5M KOH, and used immediately; coverslips

stained for 5 minutes in the dark at RT). Live cell imaging experiments tracking septin localization were

conducted using Filipin III (stock solution prepared at 5mg/mL in DMSO; working solution was used at

25μg/mL final concentration in liquid complete media), phytosphingosine (working solution was used at

15μM in liquid complete media), and Wortmannin (stock solution prepared to 2mg/mL in DMSO; working

solution was used at 20ug/mL in liquid complete media). Vehicle controls and conducted at ≤1% w/v in

liquid media. Imaging was performed in the Biomedical Microscopy Core at the University of Georgia.

Microscopy was carried out using Zeiss Axioplan microscope and Zeiss Axiocam MRc charge-coupled

device camera and software, as well as Deltavision I Deconvolution Inverted Microscope and LSM880

Confocal Fluorescent microscope with Diode laser (405nm), Argon (458, 488, 514nm) and HeNe (543,

633nm) laser lines. All micrograph comparisons between treatments imaged with identical microscope

settings. Subsequent image analysis and scale bars added to micrographs, using ImageJ software 1.48v,

Java 1.6.0_20 (64-bit) or Zen 2.3 imaging software, and final figures compiled in Photoshop CS5

software version 12.0 x32.

Quantification of aniline blue and CFW staining patterns by line scans

Max projection images and scale bars were made in each micrograph in Zen (Black) Edition.

Micrographs were false colored black and white, line scans were drawn by hand through hyphal tips, line

scan profiles were generated in Zen 3.1 (Blue) imaging software, and associated date points for each

profile were exported to Microsoft Excel version 2004 (Build 12730.20270) to create graphs and conduct

statistical analysis. Figures were generated using Adobe Photoshop 2020 Version: 21.1.2.

Quantification of chitin synthase-GFP and CFW fluorescence signal by line scans

Max projection images were made for each micrograph, line scans were drawn by hand through hyphal

branch tips in ImageJ software 1.48v, and data points were exported to Microsoft Excel version 2004

(Build 12730.20270) to create graphs and conduct statistical analysis. Figures were assembled using

Adobe Photoshop 2020 Version: 21.1.2.

Cell Wall Extraction

Cell walls were isolated from a protocol based on (Bull, 1970); cell wall extraction and lyophilization

were conducted as previously described in (Guest and Momany, 2000) with slight modifications listed in

full procedure here. A single batch of complete media (recipe described above) was autoclaved and

supplements added: Arginine, Methionine, Pyridoxine, and Riboflavin, were added to single flask and

then distributed to individual flasks to be inoculated. 1×10^4 conidia/mL were inoculated in 2 flasks each

of 100mL liquid Complete Media. Flasks were incubated at 30C in orbital shaker at 200rpm for 48 hours.

Mycelia was gravity filtrated, then vacuum-filtrated through #42 Whatman filter paper, subsequently

washed with 50mL each of chilled ddH20 to remove residual media and stored at -80C until completely

frozen. Mycelium from each sample was allowed to thaw on ice and then washed sequentially with 50mL

chilled ddH20 and 0.5M NaCl. Fungal hyphal mats were transferred to mortar and pestle, then

subsequently flash frozen in liquid nitrogen and ground in chilled Tris/EDTA Disruption Buffer (DB;

20mM Tris, 50mM EDTA, pH 8.0) with pre-chilled mortar and pestle. Samples were monitored by

microscopy under 60X or 100X objective until hyphal ghosts were evident. Cell walls were separated by

centrifugation at 13,800g for 10 min at 4C. Cell pellet was placed in a beaker with 40-100mL of chilled

Tris/EDTA buffer and stirred at 4C for 12 hours. Cell pellet was collected by centrifugation as above and

stirred again with 100mL chilled ddH20 at 4C for 4 hours. Cell wall materials was collected by vacuum

filtration, frozen at -80C, lyophilized to dryness, and stored at room temperature (25°C) for further

analysis.

Cell Wall Glycosyl Linkage Analysis

To determine the glycosyl linkages, the samples were acetylated using pyridine and acetic anhydride in

order to get better solubility, before two rounds of permethylation using sodium hydroxide (15 min) and

methyl iodide (45 min). The permethylated material was hydrolyzed using 2M TFA (2 h in sealed tube at

121°C), reduced with $NaBD_4$, and acetylated using acetic anhydride/pyridine. The resulting PMAAs

(Neutral sugars) were analyzed on an Agilent 7890A GC interfaced to a 5975C MSD (mass selective

detector, electron impact ionization mode); separation was performed on a 30 m Supelco SP-2331 bonded

phase fused silica capillary column using Supelco SP-2331 fused silica capillary column (30 m x 0,25 mm

ID). The PMAAs of amino sugars were separated on Supelco Equity-1 column (30 m x 0.25 mm ID).

Further, the relative quantities of respective glycosyl linkages were calculated by integrating the peak area

of respective peak. Since the neutral and amino sugars were analyzed on different instruments, the peak

area of amino sugars was normalized with 4-Glc peak, which is prominent in both the instruments and the

integrated and normalized peak areas were pooled together to calculate the relative percentage of individual

linkages. Two independent, biological replicates were conducted for this analysis and processed in tandem.

The average area (%) of detected linkages of one representative data set is included in the graphs to show

relative differences between cell wall polysaccharides between samples.

Acknowledgements

These studies were supported by the University of Georgia Franklin College of Arts and Sciences support

to MM and Plant Biology Department support to AM. Chitin synthase strains were generously provided

by Dr. Hiroyuki Horiuchi in the Department of Biotechnology at the University of Tokyo. The glycosyl

linkage analysis was supported by a Chemical Sciences, Geosciences and Biosciences Division, Office of

Basic Energy Sciences, U.S. Department of Energy grant (DE-SC0015662) to Parastoo Azadi at the

Complex Carbohydrate Research Center at UGA.

References

1.   Shaw JA, Mol PC, Bowers B, Silverman SJ, Valdivieso MH, Duran A, et al. The function of chitin synthases 2 and 3 in the Saccharomyces cerevisiae cell cycle. J Cell Biol. 1991;114(1):111-23.

2.   Bulawa CE, Miller DW, Henry LK, Becker JM. Attenuated virulence of chitin-deficient mutants of Candida albicans. Proc Natl Acad Sci U S A. 1995;92(23):10570-4.

3.   Sudoh M, Yamazaki T, Masubuchi K, Taniguchi M, Shimma N, Arisawa M, et al. Identification of a novel inhibitor specific to the fungal chitin synthase. Inhibition of chitin synthase 1 arrests the cell growth, but inhibition of chitin synthase 1 and 2 is lethal in the pathogenic fungus Candida albicans. J Biol Chem. 2000;275(42):32901-5.

4.   Munro CA, Gow NA. Chitin synthesis in human pathogenic fungi. Med Mycol. 2001;39 Suppl 1:41-53.

5.   Schmidt M. Survival and cytokinesis of Saccharomyces cerevisiae in the absence of chitin. Microbiology. 2004;150(Pt 10):3253-60.

6.   Orlean P. Architecture and biosynthesis of the Saccharomyces cerevisiae cell wall. Genetics. 2012;192(3):775-818.

7.   Gladfelter AS, Pringle JR, Lew DJ. The septin cortex at the yeast mother-bud neck. Curr Opin Microbiol. 2001;4(6):681-9.

8.   Longtine MS, Bi E. Regulation of septin organization and function in yeast. Trends Cell Biol. 2003;13(8):403-9.

9.   Caudron F, Barral Y. Septins and the lateral compartmentalization of eukaryotic membranes. Dev Cell. 2009;16(4):493-506.

10.   Takizawa PA, DeRisi JL, Wilhelm JE, Vale RD. Plasma membrane compartmentalization in

yeast by messenger RNA transport and a septin diffusion barrier. Science. 2000;290(5490):341-4.

11.   Dobbelaere J, Barral Y. Spatial coordination of cytokinetic events by compartmentalization of the

cell cortex. Science. 2004;305(5682):393-6.

12.   Oh Y, Bi E. Septin structure and function in yeast and beyond. Trends Cell Biol. 2011;21(3):141-

8.

13.   Glomb O, Gronemeyer T. Septin Organization and Functions in Budding Yeast. Front Cell Dev

Biol. 2016;4:123.

14.   Trimble WS, Grinstein S. Barriers to the free diffusion of proteins and lipids in the plasma

membrane. J Cell Biol. 2015;208(3):259-71.

15.   McMurray MA, Bertin A, Garcia G, 3rd, Lam L, Nogales E, Thorner J. Septin filament formation

is essential in budding yeast. Dev Cell. 2011;20(4):540-9.

16.   Sirajuddin M, Farkasovsky M, Hauer F, Kuhlmann D, Macara IG, Weyand M, et al. Structural

insight into filament formation by mammalian septins. Nature. 2007;449(7160):311-5.

17.   Bertin A, McMurray MA, Grob P, Park SS, Garcia G, 3rd, Patanwala I, et al. Saccharomyces

cerevisiae septins: supramolecular organization of heterooligomers and the mechanism of filament

assembly. Proc Natl Acad Sci U S A. 2008;105(24):8274-9.

18.   McMurray MA, Thorner J. Turning it inside out: the organization of human septin hetero-

oligomers. Cytoskeleton (Hoboken). 2019.

19.   Pan F, Malmberg RL, Momany M. Analysis of septins across kingdoms reveals orthology and

new motifs. BMC Evol Biol. 2007;7:103.

20.   Hernandez-Rodriguez Y, Masuo S, Johnson D, Orlando R, Smith A, Couto-Rodriguez M, et al.

Distinct septin heteropolymers co-exist during multicellular development in the filamentous fungus

Aspergillus nidulans. PLoS One. 2014;9(3):e92819.

21.   Hernandez-Rodriguez Y, Hastings S, Momany M. The septin AspB in Aspergillus nidulans forms

bars and filaments and plays roles in growth emergence and conidiation. Eukaryot Cell. 2012;11(3):311-

23.

22.   Fukuda K, Yamada K, Deoka K, Yamashita S, Ohta A, Horiuchi H. Class III chitin synthase

ChsB of Aspergillus nidulans localizes at the sites of polarized cell wall synthesis and is required for

conidial development. Eukaryot Cell. 2009;8(7):945-56.

23.   Blankenship JR, Fanning S, Hamaker JJ, Mitchell AP. An extensive circuitry for cell wall

regulation in Candida albicans. PLoS Pathog. 2010;6(2):e1000752.

24.   Mostowy S, Cossart P. Septins: the fourth component of the cytoskeleton. Nat Rev Mol Cell Biol.

2012;13(3):183-94.

25.   Wiederhold NP, Kontoyiannis DP, Prince RA, Lewis RE. Attenuation of the activity of

caspofungin at high concentrations against candida albicans: possible role of cell wall integrity and

calcineurin pathways. Antimicrob Agents Chemother. 2005;49(12):5146-8.

26.   Munro CA, Selvaggini S, de Bruijn I, Walker L, Lenardon MD, Gerssen B, et al. The PKC, HOG

and Ca2+ signalling pathways co-ordinately regulate chitin synthesis in Candida albicans. Mol Microbiol.

2007;63(5):1399-413.

27.   Fujioka A, Terai K, Itoh RE, Aoki K, Nakamura T, Kuroda S, et al. Dynamics of the Ras/ERK

MAPK cascade as monitored by fluorescent probes. J Biol Chem. 2006;281(13):8917-26.

28. Walker LA, Munro CA, de Bruijn I, Lenardon MD, McKinnon A, Gow NA. Stimulation of chitin

synthesis rescues Candida albicans from echinocandins. PLoS Pathog. 2008;4(4):e1000040.

29. Rodriguez-Pena JM, Garcia R, Nombela C, Arroyo J. The high-osmolarity glycerol (HOG) and

cell wall integrity (CWI) signalling pathways interplay: a yeast dialogue between MAPK routes. Yeast.

2010;27(8):495-502.

30. Colabardini AC, Ries LN, Brown NA, Savoldi M, Dinamarco TM, von Zeska Kress MR, et al.

Protein kinase C overexpression suppresses calcineurin-associated defects in Aspergillus nidulans and is

involved in mitochondrial function. PLoS One. 2014;9(8):e104792.

31. Garcia R, Bravo E, Diez-Muniz S, Nombela C, Rodriguez-Pena JM, Arroyo J. A novel

connection between the Cell Wall Integrity and the PKA pathways regulates cell wall stress response in

yeast. Sci Rep. 2017;7(1):5703.

32. Manfiolli AO, Mattos EC, de Assis LJ, Silva LP, Ulas M, Brown NA, et al. Aspergillus

fumigatus High Osmolarity Glycerol Mitogen Activated Protein Kinases SakA and MpkC Physically

Interact During Osmotic and Cell Wall Stresses. Front Microbiol. 2019;10:918.

33. Dickson RC. Thematic review series: sphingolipids. New insights into sphingolipid metabolism

and function in budding yeast. J Lipid Res. 2008;49(5):909-21.

34. Sims KJ, Spassieva SD, Voit EO, Obeid LM. Yeast sphingolipid metabolism: clues and

connections. Biochem Cell Biol. 2004;82(1):45-61.

35. Roelants FM, Torrance PD, Bezman N, Thorner J. Pkh1 and Pkh2 differentially phosphorylate

and activate Ypk1 and Ykr2 and define protein kinase modules required for maintenance of cell wall

integrity. Mol Biol Cell. 2002;13(9):3005-28.

36.   Weete JD, Abril M, Blackwell M. Phylogenetic distribution of fungal sterols. PLoS One.

2010;5(5):e10899.

37.   Alvarez FJ, Douglas LM, Konopka JB. Sterol-rich plasma membrane domains in fungi. Eukaryot

Cell. 2007;6(5):755-63.

38.   Tanaka S, Tani M. Mannosylinositol phosphorylceramides and ergosterol coodinately maintain

cell wall integrity in the yeast Saccharomyces cerevisiae. FEBS J. 2018;285(13):2405-27.

39.   Bagnat M, Simons K. Lipid rafts in protein sorting and cell polarity in budding yeast

Saccharomyces cerevisiae. Biol Chem. 2002;383(10):1475-80.

40.   van Meer G, Voelker DR, Feigenson GW. Membrane lipids: where they are and how they

behave. Nat Rev Mol Cell Biol. 2008;9(2):112-24.

41.   Tanaka-Takiguchi Y, Kinoshita M, Takiguchi K. Septin-mediated uniform bracing of

phospholipid membranes. Curr Biol. 2009;19(2):140-5.

42.   Bertin A, McMurray MA, Thai L, Garcia G, 3rd, Votin V, Grob P, et al. Phosphatidylinositol-4,5-

bisphosphate promotes budding yeast septin filament assembly and organization. J Mol Biol.

2010;404(4):711-31.

43.   Krokowski S, Lobato-Marquez D, Chastanet A, Pereira PM, Angelis D, Galea D, et al. Septins

Recognize and Entrap Dividing Bacterial Cells for Delivery to Lysosomes. Cell Host Microbe.

2018;24(6):866-74 e4.

44.   Zhang J, Kong C, Xie H, McPherson PS, Grinstein S, Trimble WS. Phosphatidylinositol

polyphosphate binding to the mammalian septin H5 is modulated by GTP. Curr Biol. 1999;9(24):1458-

67.

45.   Costanzo M, Baryshnikova A, Bellay J, Kim Y, Spear ED, Sevier CS, et al. The genetic

landscape of a cell. Science. 2010;327(5964):425-31.

46.   Clay L, Caudron F, Denoth-Lippuner A, Boettcher B, Buvelot Frei S, Snapp EL, et al. A

sphingolipid-dependent diffusion barrier confines ER stress to the yeast mother cell. Elife.

2014;3:e01883.

47.   Martin SW, Konopka JB. Lipid raft polarization contributes to hyphal growth in Candida

albicans. Eukaryot Cell. 2004;3(3):675-84.

48.   Beh CT, Alfaro G, Duamel G, Sullivan DP, Kersting MC, Dighe S, et al. Yeast oxysterol-binding

proteins: sterol transporters or regulators of cell polarization? Mol Cell Biochem. 2009;326(1-2):9-13.

49.   Canovas D, Perez-Martin J. Sphingolipid biosynthesis is required for polar growth in the

dimorphic phytopathogen Ustilago maydis. Fungal Genet Biol. 2009;46(2):190-200.

50.   Singh P, Ramachandran SK, Zhu J, Kim BC, Biswas D, Ha T, et al. Sphingolipids facilitate age

asymmetry of membrane proteins in dividing yeast cells. Mol Biol Cell. 2017;28(20):2712-22.

51.   Sugiyama S, Tanaka M. Distinct segregation patterns of yeast cell-peripheral proteins uncovered

by a method for protein segregatome analysis. Proc Natl Acad Sci U S A. 2019;116(18):8909-18.

52.   Bridges AA, Gladfelter AS. Septin Form and Function at the Cell Cortex. J Biol Chem.

2015;290(28):17173-80.

53.   Merlini L, Bolognesi A, Juanes MA, Vandermoere F, Courtellemont T, Pascolutti R, et al. Rho1-

and Pkc1-dependent phosphorylation of the F-BAR protein Syp1 contributes to septin ring assembly. Mol

Biol Cell. 2015;26(18):3245-62.

54.   Badrane H, Nguyen MH, Clancy CJ. Highly Dynamic and Specific Phosphatidylinositol 4,5-

Bisphosphate, Septin, and Cell Wall Integrity Pathway Responses Correlate with Caspofungin Activity

against Candida albicans. Antimicrob Agents Chemother. 2016;60(6):3591-600.

55.   de Almeida RFM. A route to understanding yeast cellular envelope - plasma membrane lipids

interplaying in cell wall integrity. FEBS J. 2018;285(13):2402-4.

56.   Wu JQ, Ye Y, Wang N, Pollard TD, Pringle JR. Cooperation between the septins and the

actomyosin ring and role of a cell-integrity pathway during cell division in fission yeast. Genetics.

2010;186(3):897-915.

57.   Zhang Y, Gao T, Shao W, Zheng Z, Zhou M, Chen C. The septins FaCdc3 and FaCdc12 are

required for cytokinesis and affect asexual and sexual development, lipid metabolism and virulence in

Fusarium asiaticum. Mol Plant Pathol. 2017;18(9):1282-94.

58.   Kuranda K, Leberre V, Sokol S, Palamarczyk G, Francois J. Investigating the caffeine effects in

the yeast Saccharomyces cerevisiae brings new insights into the connection between TOR, PKC and

Ras/cAMP signalling pathways. Mol Microbiol. 2006;61(5):1147-66.

59.   Kovacs Z, Szarka M, Kovacs S, Boczonadi I, Emri T, Abe K, et al. Effect of cell wall integrity

stress and RlmA transcription factor on asexual development and autolysis in Aspergillus nidulans.

Fungal Genet Biol. 2013;54:1-14.

60.   Roncero C, Duran A. Effect of Calcofluor white and Congo red on fungal cell wall

morphogenesis: in vivo activation of chitin polymerization. J Bacteriol. 1985;163(3):1180-5.

61.   Pancaldi S, Poli F, Dall'Olio G, Vannini GL. Morphological anomalies induced by Congo red in

Aspergillus niger. Arch Microbiol. 1984;137(3):185-7.

62.   Kopecka M, Gabriel M. The influence of congo red on the cell wall and (1----3)-beta-D-glucan

microfibril biogenesis in Saccharomyces cerevisiae. Arch Microbiol. 1992;158(2):115-26.

63.   Onishi J, Meinz M, Thompson J, Curotto J, Dreikorn S, Rosenbach M, et al. Discovery of novel
antifungal (1,3)-beta-D-glucan synthase inhibitors. Antimicrob Agents Chemother. 2000;44(2):368-77.

64.   Kojima K, Bahn YS, Heitman J. Calcineurin, Mpk1 and Hog1 MAPK pathways independently
control fludioxonil antifungal sensitivity in Cryptococcus neoformans. Microbiology. 2006;152(Pt
3):591-604.

65.   Yun Y, Liu Z, Zhang J, Shim WB, Chen Y, Ma Z. The MAPKK FgMkk1 of Fusarium

graminearum regulates vegetative differentiation, multiple stress response, and virulence via the cell wall

integrity and high-osmolarity glycerol signaling pathways. Environ Microbiol. 2014;16(7):2023-37.

66.   Dongo A, Bataille-Simoneau N, Campion C, Guillemette T, Hamon B, Iacomi-Vasilescu B, et al.

The group III two-component histidine kinase of filamentous fungi is involved in the fungicidal activity

of the bacterial polyketide ambruticin. Appl Environ Microbiol. 2009;75(1):127-34.

67.   Hohmann S. Osmotic stress signaling and osmoadaptation in yeasts. Microbiol Mol Biol Rev.

2002;66(2):300-72.

68.   Garcia-Rodriguez LJ, Trilla JA, Castro C, Valdivieso MH, Duran A, Roncero C. Characterization

of the chitin biosynthesis process as a compensatory mechanism in the fks1 mutant of Saccharomyces

cerevisiae. FEBS Lett. 2000;478(1-2):84-8.

69.   Fortwendel JR, Juvvadi PR, Perfect BZ, Rogg LE, Perfect JR, Steinbach WJ. Transcriptional

regulation of chitin synthases by calcineurin controls paradoxical growth of Aspergillus fumigatus in

response to caspofungin. Antimicrob Agents Chemother. 2010;54(4):1555-63.

70.   Pettolino FA, Walsh C, Fincher GB, Bacic A. Determining the polysaccharide composition of

plant cell walls. Nat Protoc. 2012;7(9):1590-607.

71.   Yoshimi A, Miyazawa K, Abe K. Cell wall structure and biogenesis in Aspergillus species.

Biosci Biotechnol Biochem. 2016;80(9):1700-11.

72.   Yoshimoto H, Saltsman K, Gasch AP, Li HX, Ogawa N, Botstein D, et al. Genome-wide analysis

of gene expression regulated by the calcineurin/Crz1p signaling pathway in Saccharomyces cerevisiae. J

Biol Chem. 2002;277(34):31079-88.

73.   Sio SO, Suehiro T, Sugiura R, Takeuchi M, Mukai H, Kuno T. The role of the regulatory subunit

of fission yeast calcineurin for in vivo activity and its relevance to FK506 sensitivity. J Biol Chem.

2005;280(13):12231-8.

74.   Wang S, Liu X, Qian H, Zhang S, Lu L. Calcineurin and Calcium Channel CchA Coordinate the

Salt Stress Response by Regulating Cytoplasmic Ca2+ Homeostasis in Aspergillus nidulans. Appl

Environ Microbiol. 2016;82(11):3420-30.

75.   Harvald EB, Olsen AS, Faergeman NJ. Autophagy in the light of sphingolipid metabolism.

Apoptosis. 2015;20(5):658-70.

76.   Barbet NC, Schneider U, Helliwell SB, Stansfield I, Tuite MF, Hall MN. TOR controls

translation initiation and early G1 progression in yeast. Mol Biol Cell. 1996;7(1):25-42.

77.   Cardenas ME, Cutler NS, Lorenz MC, Di Como CJ, Heitman J. The TOR signaling cascade

regulates gene expression in response to nutrients. Genes Dev. 1999;13(24):3271-9.

78.   Scott PH, Lawrence JC, Jr. Attenuation of mammalian target of rapamycin activity by increased

cAMP in 3T3-L1 adipocytes. J Biol Chem. 1998;273(51):34496-501.

79.   Fitzgibbon GJ, Morozov IY, Jones MG, Caddick MX. Genetic analysis of the TOR pathway in

Aspergillus nidulans. Eukaryot Cell. 2005;4(9):1595-8.

80.   Soulard A, Cremonesi A, Moes S, Schutz F, Jeno P, Hall MN. The rapamycin-sensitive

phosphoproteome reveals that TOR controls protein kinase A toward some but not all substrates. Mol

Biol Cell. 2010;21(19):3475-86.

81.   Butcher FR, Potter VR. Control of the adenosine 3',5'-monophosphate-adenyl cyclase system in

the livers of developing rats. Cancer Res. 1972;32(10):2141-7.

82.   Ni M, Rierson S, Seo JA, Yu JH. The pkaB gene encoding the secondary protein kinase A

catalytic subunit has a synthetic lethal interaction with pkaA and plays overlapping and opposite roles in

Aspergillus nidulans. Eukaryot Cell. 2005;4(8):1465-76.

83.   Binder U, Oberparleiter C, Meyer V, Marx F. The antifungal protein PAF interferes with

PKC/MPK and cAMP/PKA signalling of Aspergillus nidulans. Mol Microbiol. 2010;75(2):294-307.

84.   Foderaro JE, Douglas LM, Konopka JB. MCC/Eisosomes Regulate Cell Wall Synthesis and

Stress Responses in Fungi. J Fungi (Basel). 2017;3(4).

85.   Onyewu C, Blankenship JR, Del Poeta M, Heitman J. Ergosterol biosynthesis inhibitors become

fungicidal when combined with calcineurin inhibitors against Candida albicans, Candida glabrata, and

Candida krusei. Antimicrob Agents Chemother. 2003;47(3):956-64.

86.   Cheng J, Park TS, Fischl AS, Ye XS. Cell cycle progression and cell polarity require sphingolipid

biosynthesis in Aspergillus nidulans. Mol Cell Biol. 2001;21(18):6198-209.

87.   Li S, Bao D, Yuen G, Harris SD, Calvo AM. basA regulates cell wall organization and

asexual/sexual sporulation ratio in Aspergillus nidulans. Genetics. 2007;176(1):243-53.

88.   Walker EH, Pacold ME, Perisic O, Stephens L, Hawkins PT, Wymann MP, et al. Structural

determinants of phosphoinositide 3-kinase inhibition by wortmannin, LY294002, quercetin, myricetin,

and staurosporine. Mol Cell. 2000;6(4):909-19.

89.   Wymann MP, Bulgarelli-Leva G, Zvelebil MJ, Pirola L, Vanhaesebroeck B, Waterfield MD, et

al. Wortmannin inactivates phosphoinositide 3-kinase by covalent modification of Lys-802, a residue

involved in the phosphate transfer reaction. Mol Cell Biol. 1996;16(4):1722-33.

90.   Santos NC, Ter-Ovanesyan E, Zasadzinski JA, Prieto M, Castanho MA. Filipin-induced lesions

in planar phospholipid bilayers imaged by atomic force microscopy. Biophys J. 1998;75(4):1869-73.

91.   Takeo K. A correlation between mode of growth and regional ultrastructure of the plasma

membrane of Schizosaccharomyces pombe as revealed by freeze-fracturing before and after filipin

treatment. J Gen Microbiol. 1985;131(2):309-16.

92.   de Kruijff B, Demel RA. Polyene antibiotic-sterol interactions in membranes of Acholeplasma

laidlawii cells and lecithin liposomes. 3. Molecular structure of the polyene antibiotic-cholesterol

complexes. Biochim Biophys Acta. 1974;339(1):57-70.

93.   Bjorkbom A, Rog T, Kaszuba K, Kurita M, Yamaguchi S, Lonnfors M, et al. Effect of

sphingomyelin headgroup size on molecular properties and interactions with cholesterol. Biophys J.

2010;99(10):3300-8.

94.   Ramstedt B, Slotte JP. Sphingolipids and the formation of sterol-enriched ordered membrane

domains. Biochim Biophys Acta. 2006;1758(12):1945-56.

95.   Bridges AA, Zhang H, Mehta SB, Occhipinti P, Tani T, Gladfelter AS. Septin assemblies form by

diffusion-driven annealing on membranes. Proc Natl Acad Sci U S A. 2014;111(6):2146-51.

96.   Cho H, Stanzione F, Oak A, Kim GH, Yerneni S, Qi L, et al. Intrinsic Structural Features of the

Human IRE1alpha Transmembrane Domain Sense Membrane Lipid Saturation. Cell Rep.

2019;27(1):307-20 e5.

97.   Nyholm TK, Ozdirekcan S, Killian JA. How protein transmembrane segments sense the lipid

environment. Biochemistry. 2007;46(6):1457-65.

98.   Valadares NF, d' Muniz Pereira H, Ulian Araujo AP, Garratt RC. Septin structure and filament

assembly. Biophys Rev. 2017;9(5):481-500.

99.   Cannon KS, Woods BL, Crutchley JM, Gladfelter AS. An amphipathic helix enables septins to

sense micrometer-scale membrane curvature. J Cell Biol. 2019;218(4):1128-37.

100.   Valiante V, Macheleidt J, Foge M, Brakhage AA. The Aspergillus fumigatus cell wall integrity

signaling pathway: drug target, compensatory pathways, and virulence. Front Microbiol. 2015;6:325.

101.   Valiante V, Jain R, Heinekamp T, Brakhage AA. The MpkA MAP kinase module regulates cell

wall integrity signaling and pyomelanin formation in Aspergillus fumigatus. Fungal Genet Biol.

2009;46(12):909-18.

102.   Valiante V, Heinekamp T, Jain R, Hartl A, Brakhage AA. The mitogen-activated protein kinase

MpkA of Aspergillus fumigatus regulates cell wall signaling and oxidative stress response. Fungal Genet

Biol. 2008;45(5):618-27.

103.   Ram AF, Wolters A, Ten Hoopen R, Klis FM. A new approach for isolating cell wall mutants in

Saccharomyces cerevisiae by screening for hypersensitivity to calcofluor white. Yeast. 1994;10(8):1019-

30.

104.   Ram AF, Kapteyn JC, Montijn RC, Caro LH, Douwes JE, Baginsky W, et al. Loss of the plasma

membrane-bound protein Gas1p in Saccharomyces cerevisiae results in the release of beta1,3-glucan into

the medium and induces a compensation mechanism to ensure cell wall integrity. J Bacteriol.

1998;180(6):1418-24.

105.   Kapteyn JC, Ram AF, Groos EM, Kollar R, Montijn RC, Van Den Ende H, et al. Altered extent

of cross-linking of beta1,6-glucosylated mannoproteins to chitin in Saccharomyces cerevisiae mutants

with reduced cell wall beta1,3-glucan content. J Bacteriol. 1997;179(20):6279-84.

106.   Kapteyn JC, Hoyer LL, Hecht JE, Muller WH, Andel A, Verkleij AJ, et al. The cell wall

architecture of Candida albicans wild-type cells and cell wall-defective mutants. Mol Microbiol.

2000;35(3):601-11.

107.   Popolo L, Gilardelli D, Bonfante P, Vai M. Increase in chitin as an essential response to defects

in assembly of cell wall polymers in the ggp1delta mutant of Saccharomyces cerevisiae. J Bacteriol.

1997;179(2):463-9.

108.   Dallies N, Francois J, Paquet V. A new method for quantitative determination of polysaccharides

in the yeast cell wall. Application to the cell wall defective mutants of Saccharomyces cerevisiae. Yeast.

1998;14(14):1297-306.

109.   Osmond BC, Specht CA, Robbins PW. Chitin synthase III: synthetic lethal mutants and "stress

related" chitin synthesis that bypasses the CSD3/CHS6 localization pathway. Proc Natl Acad Sci U S A.

1999;96(20):11206-10.

110.   Lagorce A, Le Berre-Anton V, Aguilar-Uscanga B, Martin-Yken H, Dagkessamanskaia A,

Francois J. Involvement of GFA1, which encodes glutamine-fructose-6-phosphate amidotransferase, in

the activation of the chitin synthesis pathway in response to cell-wall defects in Saccharomyces

cerevisiae. Eur J Biochem. 2002;269(6):1697-707.

111.   Perez J, Arcones I, Gomez A, Casquero V, Roncero C. Phosphorylation of Bni4 by MAP kinases

contributes to septum assembly during yeast cytokinesis. FEMS Yeast Res. 2016;16(6).

112.   Heinisch JJ, Rodicio R. Protein kinase C in fungi-more than just cell wall integrity. FEMS

Microbiol Rev. 2018;42(1).

113.   Larson JR, Kozubowski L, Tatchell K. Changes in Bni4 localization induced by cell stress in

Saccharomyces cerevisiae. J Cell Sci. 2010;123(Pt 7):1050-9.

114.   Harris SD, Morrell JL, Hamer JE. Identification and characterization of Aspergillus nidulans

mutants defective in cytokinesis. Genetics. 1994;136(2):517-32.

115.   Momany M, Westfall PJ, Abramowsky G. Aspergillus nidulans swo mutants show defects in

polarity establishment, polarity maintenance and hyphal morphogenesis. Genetics. 1999;151(2):557-67.

Supporting information

907
S1 Fig. Line scans of cell wall staining reveals unique patterns of chitin and β-1,3-glucan deposition
in core septin null mutants. (A) Line scan profiles (Left column) of aniline blue staining patterns,

corresponding to the red arrows drawn through the hyphal tips of WT and $\Delta aspB^{cdc3}$ (Right column). (B)
Line scan profiles (Left column) of calcofluor white staining patterns, corresponding to the red arrows
drawn through the hyphal tips of WT and $\Delta aspB^{cdc3}$ (Right column). Asterisk in Panel A line scan profile
denotes saturation of fluorescence signal.

S2 Fig. Cell wall glycosyl linkage analysis shows increased chitin content in $\Delta aspB^{cdc3}$ septin null
mutant. Results of cell wall polysaccharide glycosyl linkage analysis using GC MS/MS showing the
average area (%) of detected linkages of 4-linked glucose and 4-linked N-acetyl glucosamine. Two
independent biological replicates gave similar results. A representative data set is shown. Samples: WT,
$\Delta aspB^{cdc3}$, and $\Delta aspE$.

S3 Fig. Quantification of chitin synthase B-GFP and CFW fluorescence signal. (A) Quantification of
percentage (%) of hyphal branch tips with chitin synthase B-GFP present vs. absent by line scans in WT
and Δ*aspA*^{cdc11}. Mean values shown for line scans of ≥85 hyphal branches over two independent
biological replicates. Asterisks indicate different standard error of the mean between sample sets. (B)
Quantification of maximum fluorescence intensity by line scans of CFW stained WT and Δ*aspA*^{cdc11}

hyphae (subapical region). Mean values shown for line scans of ≥65 hyphal branches over two
independent biological replicates. Asterisks indicate different standard error of the mean between sample
sets and statistically different values by two-tailed Ttest (p < 0.05) (N=50).

S4 Fig. Core septin null mutants are not sensitive to treatments which disrupt the Ca2+/Calcineurin
or TOR pathways. AspE is hypersensitive to treatments which disrupt the TOR pathway. WT and
septin null mutants $\Delta aspA^{cdc11}$, $\Delta aspB^{cdc3}$, $\Delta aspC^{cdc12}$, $\Delta aspD^{cdc10}$, and $\Delta aspE$ were tested for sensitivity by
spotting decreasing spore concentrations on complete media plates with or without Calcium/Calcineurin
pathway-disturbing agents (EGTA, CaCl2, and FK-506) or Target of Rapamycin (TOR) pathway-
disturbing agents (Rapamycin, caffeine, and theophylline). Differences in colony color result from
changes in spore production, spore pigment, and production of secondary metabolites under stress. Spore
concentrations were [1^7 conidia/mL – 10^4 conidia/mL] for all assays in figure. N=3

S3

S5 Fig. Septin AspB^{Cdc3}-GFP localization disrupted by sphingolipid biosynthesis intermediate
phytosphingosine. AspB^{Cdc3}-GFP strain incubated in liquid media for approximately 16h and imaged
180 minutes after replacing with fresh media containing sphingolipid biosynthesis-inhibiting agents.

Representative images are shown from three independent biological replicates, with ≥100 cells observed.
(Left Panel) AspB-GFP in vehicle control treatments and (Right Panel) phytosphingosine (15µM)
treatment. Enlarged section of micrographs from each picture to better visualize pattern of fluorescence.
White arrows denote hyphae which are highlighted in enlarged images. Imaging conducted with
Deltavision I deconvolution inverted fluorescence microscope. Scale bars = 10µm. N=3

S6 Fig. Septin AspA^{Cdc11}-GFP localization disrupted by sphingolipid biosynthesis inhibitors.
AspA^{Cdc11}-GFP strain incubated in liquid media for approximately 16h and imaged 180 minutes after
replacing with fresh media containing sphingolipid biosynthesis-inhibiting agents. Representative images
are shown from three independent biological replicates, with ≥100 cells observed. (Left Panel) AspA-
GFP in vehicle control treatment, (Middle Panel) Myriocin (17.5µg/mL) and (Right Panel)
phytosphingosine (15µM) treatment. Enlarged section of micrographs from each picture to better
visualize pattern of fluorescence signal. White arrows denote hyphae which are highlighted in enlarged
images. Imaging conducted with Deltavision I deconvolution inverted fluorescence microscope. Scale
bars = 10µm. N=3

971   **S1 Table.** List of fungal strains used in this study.

Strain Number	Genotype	Source
FGSC A850 (WT)	biA1; _argB::trpC_B; methG1; veA1; trpC801	FGSC
ASH5 (ΔaspA)	aspA::argB2 biA1 argB::trpC_B methG1 veA1 trpC801	Lindsey and Momany, 2010
AYR32 (ΔaspB)	aspB::AfpyrG; pyroA4; argB2	Hernandez-Rodriguez et al., 2012
ARL161 (ΔaspC)	aspC::AfpyrG pyrG89 biA1 argB::trpC_B methG1 veA1 trpC801	Lindsey and Momany, 2010
AKK3 (ΔaspD)	aspD::AfpyrG; pyroA4; argB2	Hernandez-Rodriguez et al., 2014
ASH41 (ΔaspE)	aspE::AfpyrG; riboB2	Hernandez-Rodriguez et al., 2012
ANID_05666 (ΔmpkA)	pyrG89; argB2;ΔnkuA(ku70)::argB;ΔmpkA::pyrG;pyroA	CP De Souza et al., 2013
AAM016 (ΔmpkAΔaspB)	aspB::AfpyrG; pyroA4; argB2 + pyrG89; argB2;ΔnkuA(ku70)::argB;ΔmpkA::pyrG;pyroA	This study
AAM017 (ΔmpkAΔaspB)	aspB::AfpyrG; pyroA4; argB2+ pyrG89; argB2;ΔnkuA(ku70)::argB;ΔmpkA::pyrG;pyroA	This study

AAM019 (ΔmpkAΔaspE)	aspE::AfpyrG; riboB2+ pyrG89;argB2;ΔnkuA(ku70)::argB;ΔmpkA::pyrG;pyroA	This study
AAM020 (ΔmpkAΔaspE)	aspE::AfpyrG; riboB2+ pyrG89; argB2;ΔnkuA(ku70)::argB;ΔmpkA::pyrG;pyroA	This study
ARL141	*aspA-GFP-AfpyrG pyrG89 biA1 argB::trpC_B methG1 veA1 trpC801*	Lindsey and Momany, 2010
EB-5	*biA1 pyrG89 argB2 pyroA4 wA3 ΔchsB::pyr-4-alcA(p)-chsB argB::chsB(p)-egfp-chsB*	Fukuda et al., 2009
AAM022	*aspA::argB2 biA1 pyrG89 argB2 wA3 ΔchsB::pyr-4-alcA(p)-chsB argB::chsB(p)-egfp-chsB*	This study

973   **S2 Table.** List of primers and sequences used in this study.

Primer Name	Primer Sequence
PyrG-Af-R'	**5'**-CAG AGC CCA CAG AGC GCC TTG AG-**3'**
AspB-KO-F'	**5'**-GGT CAT TCC TGG TGT GAC AGT ACC-**3'**
AspE-KO-F'	**5'**-GAT CCA AAT TCC AGG TTC GAT GAC-**3'**
MpkA-806-F'	**5'**-ATC CTA GAC TCG ACG CCT CA-**3'**
MpkA-3779-R'	**5'**-ACA AAA ACC CCA TCG TCC GA-**3'**
AspA-KO-F'	**5'**-TAG ATC AAG CTC CGC CGG AA-**3'**
AspA-KO-R'	**5'**-TGA CTC CAG CGA CGA TGA GT-**3'**