Metastable GPCR dimers trigger the basal signal by recruiting G-proteins

Rinshi S. Kasai[1], Takahiro K. Fujiwara[2], and Akihiro Kusumi[3]*

[1]Institute for Frontier Life and Medical Sciences and [2]Institute for Advanced Study, Institute for Integrated Cell-Material Sciences (KUIAS-iCeMS), Kyoto University, Kyoto 606-8507, Japan. [3]Membrane Cooperativity Unit, Okinawa Institute of Science and Technology Graduate University (OIST), Onna-son, Okinawa, 904-0495 Japan.

*To whom all correspondence should be addressed:
Akihiro Kusumi
e-mail: akihiro.kusumi@oist.jp
Ph: 011-81-98-966-1373, Fx: 011-81-98-966-1062

G-protein-coupled receptors (GPCRs) constitute the largest family of integral membrane proteins in the human genome and are responsible for various important signaling pathways for vision, olfaction, gustation, emotion, cell migration, etc. A distinct feature of the GPCR-family proteins is that many GPCRs, including the prototypical GPCR, β2-adrenergic receptor (β2AR), elicit low levels of basal constitutive signals without agonist stimulation, which function in normal development and various diseases[1-3]. However, how the basal signals are induced is hardly known. Another general distinctive feature of GPCRs is to form metastable homo-dimers, with lifetimes on the order of 0.1 s, even in the resting state. Here, our single-molecule-based quantification[4] determined the dissociation constant of β2AR homo-dimers in the PM (1.6 ± 0.29 copies/μm^2) and their lifetimes (83.2 ± 6.4 ms), and furthermore found that, in the resting state, trimeric G-proteins were recruited to both β2AR monomers and homo-dimers. Importantly, inverse agonists, which suppress the GPCR's basal constitutive activity, specifically blocked the G-protein recruitment to GPCR homo-dimers, without affecting that to monomers. These results indicate that the G-proteins recruited to transient GPCR homo-dimers are responsible for inducing their basic constitutive signals. These results suggest novel drug development strategies to enhance or suppress GPCR homo-dimer formation.

β2AR forms metastable homo-dimers

Single-molecule imaging-tracking revealed that β2AR (tagged with ACP at the N-terminus and labelled with ATTO594; **Extended Data Fig. 1**) undergoes thermal diffusion, and often forms transient homo-dimers, undergoing rapid interconversions between monomers and dimers continually, as found for other GPCRs (**Fig. 1a, Extended Data Fig. 2a**)[4-8]. Every time we found a β2AR dimer, we measured the dimer duration, and after observing many dimers, we obtained the distribution of dimer durations (**Fig. 1b, Extended Data Fig. 2d**). The histogram could be fitted by a single exponential function, providing the exponential dimer lifetime of 99 ms (throughout this report, all statistical parameters, including SEM, and the number of experiments are summarised in

Extended Data Tables 1-4; all observations were performed at 37°C). The lifetime value obtained

in this manner, termed $\tau_{observed}$, was corrected for the photobleaching lifetime and incidental

colocalisation lifetime (this method was used for evaluating all of the lifetimes investigated in the

present research; see **Methods**), giving a dimer lifetime τ of 83.2 ± 6.4 ms (**Fig. 1c**). The dimer

lifetime obtained at a time resolution of 4 ms (250 Hz) was 78.8 ± 10.7 ms, after corrections for

photobleaching and incidental colocalisation lifetime (**Extended Data Fig. 3, Fig. 1c**), which is

statistically indistinguishable from the dimer lifetime of 83.2 ms evaluated from 30-Hz observations.

This result indicates that a frame rate of 30 Hz is sufficiently fast for evaluating the homo-dimer

lifetime of β2AR. Together, these results demonstrate that the β2AR homo-dimers are indeed quite

metastable, turning into monomers in a matter of 80 ms, and then quite readily forming another

dimer with a different partner molecule (**Extended Data Fig. 2a**).

2D-dissociation constant for β2AR dimers

Using our single-molecule-based "super-quantification" method[4], we obtained the 2-dimensional

dissociation constant (2D-K_D) of β2AR homo-dimers in the PM of living cells, which was 1.6 ± 0.29

copies/μm^2 (**Fig. 2a, Extended Data Fig. 2b**; see **Methods**). This value is comparable to the

2D-K_D of another GPCR, formyl-peptide receptor (FPR), which was 3.6 copies/μm^2 (in fact, this was

the first determination of the 2D-K_D in the PM of living cells)[4].

Under physiological conditions, β2AR is expressed at number densities of 16 ~ 260 copies/μm^2,

depending on the cell type[9,10]. From the 2D-K_D determined here, we conclude that 12 ~ 240

copies/μm^2 exist as dimers and 4 ~ 20 copies/μm^2 exist as monomers at any moment; *i.e.*, 75 ~

92% of β2AR molecules exist as homo-dimers at any moment, although dimers and monomers are

interconverting rapidly (with a dimer lifetime of ~80 ms). In contrast, FPR is mainly expressed in

neutrophils, and an average of about 6,000 copies of FPR copies exist in the PM of a single

neutrophil[11]. Assuming that the neutrophil surface area is 1,300 μm^2 (approximated by a sphere with

a radius of 10 μm, providing a number density of ~4.6 FPR copies/ μm^2) and based on the 2D-K_D,

3,200 copies of FPR would exist as monomers and 2,800 copies as dimers; namely, 54% of FPR

molecules are dimers, which is about two-thirds of the dimer fraction of β2AR. Note that, even under

the conditions where the dimer population dominates (*i.e.*, at higher expression levels), the dimer

lifetimes do not change and remain on the order of 0.1 s. Only the monomer lifetimes are shortened

at higher expression levels.

Direct detection of β2AR dimers by the BiFC method

Thus far, we have detected dimers by the colocalisation of two single molecules, which reveals the

close encounters of molecules in the space scale of 220 nm, a distance much greater than the

molecular scale. Therefore, in this method, the incidental approaches of two molecules were

evaluated and subtracted. To ascertain the molecular-level interactions (dimers), we employed

YFP-based bimolecular fluorescence complementation (BiFC; meanwhile, our attempt to detect

FRET did not work, perhaps because the distance between the probes attached to two C-termini was

too far for FRET)[4,12]. In BiFC, two potentially interacting proteins are fused to the N- and C-terminal

half-molecules of YFP. If the proteins of interest form dimers, then YFP may be reconstituted and

emit fluorescence signals (see **Methods**). Thus, if BiFC occurs, it would lend strong support for the

formation of β2AR dimers.

To quantitatively examine BiFC, we varied the expression levels of β2AR. With an increase in

the number density of expressed molecules (determined by the signal from ATTO594 bound to ACP;

the ATTO594/ACP ratio is >0.95[13]), the number density of the BiFC spots of β2AR increased,

whereas the BiFC-spot number density of the monomer reference molecule, ACP-TM, increased only

slightly (**Fig. 2b; Extended Data Fig. 2c**). This result indicates that β2AR forms directly-bound

dimers.

Inverse agonists reduce the β2AR dimer affinity

Treatments with the inverse agonists ICI-118,551 (ICI) and Timolol reduced the β2AR dimer lifetime

(0.74x and 0.69x, respectively; **Fig. 1c, Extended Data Fig. 2d**), whereas ICI increased the 2D-K_D

of β2AR by a factor of 2.1 (lowered the β2AR dimer affinity; **Fig. 1c**), indicating that inverse agonists

reduce the β2AR dimer affinity. Meanwhile, the addition of the agonist isoproterenol (Iso) prolonged

the dimer lifetime significantly (1.55x) (**Fig. 1c; Extended Data Fig. 2d**). Cholesterol reportedly

modulates GPCR functions[14], but in our experiments, mild cholesterol depletion by

methyl-β-cyclodextrin (MβCD) treatment did not affect the dimer lifetime (**Fig. 1c**).

Since the inverse agonists suppressed the basal constitutive GPCR signaling, and we found

that they also reduced the β2AR dimer lifetime, we next examined the dimer lifetimes of the β2AR

mutants with high and low basal activities in the resting state (HBA and LBA mutants, representing

E268A and C327R mutants, respectively[15,16]). The HBA mutant exhibited a 1.78x longer dimer

lifetime, and interestingly, after the inverse-agonist addition, it was reduced to the dimer lifetime of

the wild-type receptor (**Fig. 1c; Extended Data Fig. 2d**). Meanwhile, the LBA mutant exhibited a

0.74x shorter dimer lifetime than that of the wild-type receptor, and the agonist addition did not

affect it (**Fig. 1c; Extended Data Fig. 2d**).

Strikingly, the dimer lifetimes of WT-β2AR after the addition of inverse agonists were quite

comparable to those of the LBA mutant, and the dimer lifetime of WT-β2AR after the addition of the

agonist isoproterenol was comparable to that of the HBA mutant. These results suggest that the

β2AR conformations would be quite different after the binding of the inverse agonists and the

agonist isoproterenol, as compared with those before their additions, possibly resembling the

conformations of the LBA and HBA mutants, respectively, and detectable as changes in the

homo-dimer lifetimes.

The basal constitutive activity of β2AR

We confirmed the basal constitutive activity of β2AR, as well as the effects of the β2AR agonist

isoproterenol and the inverse agonists ICI-118,551 and Timolol, by measuring the changes in the

cytoplasmic cAMP levels in the L-cell line stably expressing β2AR at higher levels (**Extended Data**

Fig. 4, Extended Data Table 4). The addition of the agonist isoproterenol to the cells increased

the cAMP level by a factor of ~4, relative to the control. Meanwhile, the addition of the inverse

agonists ICI-118,551 and Timolol reduced the cAMP levels by factors of 1.9 and 1.3 (53 and 76% of

the control), respectively, showing the presence of the basal constitutive activity of β2AR and the

effectiveness of the inverse agonists, ICI-118,551 and Timolol, for strongly reducing this basal

activity.

Trimeric G proteins are recruited to both β2AR monomers and dimers

The presence of the basal constitutive activity of β2AR suggests that a stimulatory trimeric G-protein might be recruited to β2AR in the resting state. Therefore, we examined whether Gαs and Gβ (conjugated with mGFP) are recruited to β2AR monomers or dimers, by examining their colocalisations. Using L cells expressing both β2AR (ATTO594-ACP-β2AR) and Gαs or Gβ, we found that, even in resting cells, Gαs and Gβ were recruited to both β2AR monomers and dimers (**Fig. 3a; Supplementary Videos 1 and 2**).

The recruitment frequencies of Gαs and Gβ to β2AR monomers and dimers were measured (per sec per μm^2) and then normalised by the number densities of Gαs (or Gβγ) and β2AR protomers in the PM (normalised frequencies; **Fig. 3b, c; Extended Data Table 3**)[17]. In resting cells, both Gαs and Gβ molecules were recruited to β2AR monomers or dimers equally well, suggesting that (non-stimulated) β2AR monomers and/or dimers might induce intracellular Gs-dependent signals even in the resting state, which would lead to the basal constitutive activity of β2AR.

The exponential dwell lifetimes of G proteins on β2AR were on the order of 50 ~ 60 ms for all of the combinations of Gαs and Gβ molecules vs β2AR monomers and dimers (**Extended Data Fig. 5**). However, these durations are only the maximal estimates of the dwell lifetimes of Gαs and Gβ molecules on β2AR monomers and dimers, due to the present instrumental limitations of detecting short binding durations.

Notably, the treatment of the cells with the agonist, isoproterenol, did not change the recruitment frequencies in any combinations of Gαs and Gβ vs β2AR monomers and dimers, although it significantly enhanced the cytoplasmic cAMP level (**Extended Data Fig. 4, Extended Data Table 4**). This result is quite surprising, but it suggests that the isoproterenol-induced β2AR conformation enhanced its GEF activity, rather than increasing the on-rate of G protein binding to β2AR.

Inverse agonists specifically block G-protein recruitment to β2AR dimers

Next, we examined the effects of the inverse agonists. The recruitment of Gαs and Gβ to β2AR

dimers was almost completely blocked, whereas that to β2AR monomers was unaffected (**Fig. 3b, c**). This result unequivocally shows that β2AR dimers are responsible for the basal constitutive activity of β2AR. Furthermore, it demonstrates that the inverse agonists work by blocking the recruitment of stimulatory G-proteins to β2AR dimers, perhaps by inducing β2AR conformational changes in the binding sites for G-proteins, rather than blocking the β2AR dimerisation (although, as already described, the dimer affinity of the inverse agonist-bound β2AR is lower, resembling the LBA mutant, and thus inverse agonists might change the GPCR conformations in two ways: lowering the GPCR dimer affinity and modulating the G-protein binding sites).

To further confirm that β2AR dimers trigger downstream signaling without an agonist, and to detect the basal dimer signal at the β2AR expression levels used for single-molecule experiments, we examined whether artificially-induced stable β2AR dimers could elicit the downstream signal. To detect the downstream signal at very low expression levels of β2AR, which are necessary for detecting dimers by single-molecule imaging, and also to detect the signal in a different manner, rather than observing the cAMP level again (**Extended Data Fig. 4**), we examined the intracellular Ca^{2+} mobilisation, which could occur further downstream from the cAMP amplification[18]. The induction of the artificially-induced stable β2AR (conjugated to FKBP) dimers by the FKBP-A20187 dimerisation system indeed elicited Ca^{2+} mobilisation (**Extended Data Fig. 6**). This result further confirmed that β2AR dimers could trigger the downstream signals without agonist stimulation.

Discussion

All of the class-A GPCRs examined thus far, including β2AR examined here and by others[5], FPR, D2 dopamine receptor, M1 muscarinic receptor, and α1AR, form metastable dimers with lifetimes on the order of 0.1 s in the non-stimulated state[4-8]. Therefore, we propose that the dynamic equilibrium between monomers and homo-dimers is a common characteristic property of the class-A GPCRs. The homo-dimerisation of GPCRs without any noticeable amino-acid homologies suggests that, through evolution, GPCRs changed almost all of their amino acids, but conserved the physical ability to form transient homo-dimers. This infers that these dynamic dimers are critical for some of the GPCR functions.

GPCRs are distinct among other receptors, in that under non-stimulated conditions GPCRs

maintain the low basal constitutive activity, which can be inhibited by inverse agonists. The basal

activities of GPCRs are implied in normal development and various diseases[1-3]. Therefore, we

examined the possibility that the function of the GPCR dimers is to induce the basal constitutive

activity in resting cells (without agonists), and obtained three unexpected, surprising findings. First,

both β2AR monomers and dimers continually recruit G proteins, one molecule after another, even in

the resting state. Second, the G-protein recruitment frequencies in the resting state are about the

same in the presence of the agonist. Third, more importantly, the inverse agonists specifically and

almost completely blocked the recruitment of Gαs and Gβ to β2AR dimers, without affecting their

recruitment frequencies to β2AR monomers (**Fig. 3b, c**). The third result unequivocally shows that

the G proteins bound to β2AR dimers, and not monomers, in the resting state are responsible for

triggering the β2AR's basal constitutive activity, which is sensitive to inverse agonism. It further

suggests that the binding of the inverse agonists to β2AR induces conformational changes that

decrease the β2AR dimer affinities for G proteins (**Fig. 2a**) and at the same time, hide the binding

site(s) for G proteins in the dimer conformations.

Although the function of the GPCR dimers is to maintain the basal constitutive activity, it is

critically important to realise that the GPCR dimers are metastable with lifetimes on the order of 0.1

s. Therefore, the GPCR dimers are dispersing continually, forming new dimers with other partner

molecules all the time. In cells, this allows the basal constitutive activities of GPCRs to be maintained

at certain levels, and yet the levels of the basal GPCR signals (the average number density of GPCR

dimers at any time) are regulated globally, by modulating the GPCR number density by varying the

expression levels of the GPCR, and locally, by temporarily and spatially varying the actin-based

membrane skeleton and the membrane curvature[5,19-21]. In addition, the metastable GPCR dimers

might play important roles in the activation of G proteins after the agonist addition.

The present results suggest that the major functions of some of the orphan GPCRs might be

performed by the basal activities induced by their homo-dimers, as the GPCR family includes many

orphan receptors[3,22]. Based on the results described in this report, we strongly suggest the pursuit of

drug development strategies that emphasise the generation of more inverse agonists for various

GPCRs, as well as drugs that enhance or suppress GPCR homo-dimer formation.

Online content

Any methods, additional references, Nature Research reporting summaries, source data, extended

data, supplementary information, acknowledgements, peer review information; details of author

contributions and competing interests; and statements of data and code availability are available.

Acknowledgements

We thank Prof. Hajime Fujisawa of Nagoya University for providing mouse L cells, Prof. Tatsuya Haga of Gakushuin University for providing the cDNAs encoding β2AR and Gα, Prof. Tobias Meyer of Stanford University for providing the cDNA encoding YFP-FKBP through Addgene (plasmid 201751), Prof. Michiyuki Matsuda of Kyoto University for the excellent suggestion about the method for establishing cell lines stably expressing target proteins at high levels, Assoc. Prof. Akira Hattori for the use of the PerkinElmer time-resolved fluorescence resonance energy transfer reader, EnVision Xcite, for the cAMP measurement, Drs. Taka-Aki Tsunoyama of OIST, Ko Hirosawa of Gifu University, Prof. Kenichi Suzuki of Gifu University, and all of the members of the Kusumi laboratory for valuable discussions, suggestions, and support, and Mr. Koji Kanemasa of incomings for assistance with figure preparation. This work was supported by Grants-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (Kiban C to R.S.K. [17K07333], Wakate B to R.S.K. [26870292], Kiban S to R.S.K. [Co-PI, 18H05269], and Kiban B to R.S.K. [Co-PI, 17H03666], Kiban B to T.K.F. [16H04775], and Kiban S to A.K. [16H06386]), Grants-in-Aid for Innovative Areas (Molecular Engine) from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) of Japan to R.S.K. [18H05424], and the Joint Research Program of Biosignal Research Center of Kobe University to R.S.K. [301002]. WPI-iCeMS of Kyoto University is supported by the World Premier Research Center Initiative (WPI) of the MEXT.

Author contributions

R.S.K. performed the single fluorescent-molecule tracking experiments and biochemical experiments. R.S.K., T.K.F., and A.K. developed and built the single-molecule imaging station. T.K.F. developed the analysis software. R.S.K. and A.K. conceived and formulated the project, evaluated and discussed the data, and wrote the manuscript, and all authors participated in revising the manuscript.

Methods

Cell culture, cDNA construction and expression in L cells, and treatment of cells with agonists and inverse agonists.

L cells (a kind gift from H. Fujisawa of Nagoya University), which do not express β2AR [23], were confirmed to be free of mycoplasma contamination using MycoAlert (Lonza), and were routinely cultured in Dulbecco's Minimum Essential Medium (DMEM; Sigma-Aldrich) supplemented with 10% (v/v) FBS (Sigma-Aldrich), 100 units/ml penicillin (Wako), and 0.1 mg/ml streptomycin (Wako). Human β2AR was fused at its N-terminus to the ACP- or Halo7-tag protein (NEB and Promega, respectively) at the cDNA level, and cloned into the expression plasmid pEGFP-N1 (Clontech) after the removal of the cDNA region encoding EGFP. The ACP- and Halo7-tag proteins do not induce dimerisation[12,24]. For the genetic conjugation of mGFP to Gαs, the mGFP cDNA was inserted between the region encoding the 71st and 82nd amino acids of mouse Gαs (Origene) in the pcDNA3.1 (+) vector (Invitrogen)[25], and for that to Gα, mGFP was fused to the N terminus of bovine Gα1 (a kind gift from T. Haga of Gakushuin University) [26] in the pEGFP-C1 vector (Clontech), after the removal of the cDNA region encoding EGFP. The details of the constructed cDNAs are shown in **Extended Data Fig. 1**. The L cells were transfected with these cDNAs using LipofectAMINE LTX (Life Technologies), according to the manufacturer's recommendations. Transfected cells were seeded in glass-base dishes (35 mmϕ with a glass window of 12 mmϕ, 0.12~0.17-mm-thick glass; Iwaki), and cultured in Ham's F12 medium (Sigma-Aldrich), supplemented with 10 % (v/v) FBS (Sigma-Aldrich), 100 units/ml penicillin (Wako), and 0.1 mg/ml streptomycin (Wako), for 24~48 h before microscope observations. The F12 medium was employed because it has a lower fluorescence background as compared with DMEM, which was used for routine culturing.

The agonist, isoproterenol (Sigma-Aldric), and the inverse agonists, ICI-118,551 (Tocris) and timolol (Wako), were dissolved in Hank's balanced salt solution (Nissui) buffered with 2 mM piperazine-N,N'-bis(ethanesulfonic acid) (PIPES, Dojindo) at pH 7.4

(P-HBSS), at concentrations of 20, 200, and 200 µM, respectively. For the additions to the

cells, 1 ml solutions of these reagents were added to the cells in 1 ml of the Ham's F12

medium described above, providing the final concentrations of 10 µM isoproterenol, 100 µM

ICI-118,551, and 100 µM timolol.

Labelling of β2AR with the fluorescent dye ATTO594

ATTO594-Coenzyme A was custom-synthesised by Shinsei Kagaku, by conjugating

ATTO594-maleimide (ATTO-TECH) to Coenzyme A-SH (New England Biolabs), followed by

HPLC purification. The chemical structure of ATTO594-Coenzyme A was confirmed by liquid

chromatography-mass spectrometry (LC-MS; LCMS-2010A EV, Shimadzu). The ACP-tagged

β2AR expressed in the PM of the L cells was labelled with ATTO594, by incubating the cells

with 2 µM ATTO594-Coenzyme A and 1 µM phosphopantetheine transferase (New England

Biolabs) in Ham's F12 medium without FBS at ~25˚C for 20 min. These conditions have

been known to achieve ~95% labelling of ACP[24]. The cells were washed with P-HBSS three

times, and then subjected to microscope observations. The cells exhibiting the presence of

~3 ATTO594-ACP-B2AR spots/µm^2 in the PM were used for single-molecule imaging

experiments.

Single fluorescent-molecule video imaging of live cells

The fluorescently-labelled ACP-β2AR expressed in the bottom PM was observed at the level

of single molecules at 37˚C, using home-built objective-lens-type total internal reflection

fluorescence microscopes based on Olympus IX-70, Olympus IX-81, or Nikon TE-2000

inverted microscopes[4,7,27,28], with a 100x objective lens, as optimised for the present

research. Fluorescence images were projected onto a two-stage microchannel plate

intensifier (C8600-03, Hamamatsu Photonics), coupled to an electron bombardment CCD

camera (C7190-23; Hamamatsu Photonics) or an sCMOS camera (ORCA-Flash 4.0;

Hamamatsu Photonics) operated at 30 Hz. For observations at a frame rate of 250 Hz, a

CMOS camera (FASTCAM 1024PCI-II, Photron) was employed[29]. Fluorescent spots were

identified by using an in-house computer program, as described previously[4]. The positions

(x and y coordinates) of all of the observed single fluorescent-molecules were determined

by a computer program that employed the method developed previously[30,31]. For the

simultaneous dual-color single-molecule imaging, the images were spectrally split into two

emission arms, each containing the same image intensifier and camera.

To maintain the cells at 37˚C during the microscope observations, the entire

microscope, except for the two detection arms and the excitation arm, was placed in a

home-built microscope environment chamber made with thermo-insulating transparent

vinyl sheets, and three heating circulators (SKH0-112-OT, Kokensya) were appropriately

placed in the chamber to supply warmed air with slow circulation[28,32,33]. For further

stabilisation, a plate heater (INUG2, Tokai Hit) was mounted on the microscope stage. The

temperature of the surrounding air just above the stage was continually monitored during

the observations with a digital thermometer (DT-312, Intermedical).

Detection of colocalisation-codiffusion, and evaluations of β2AR dimer lifetimes

and two-dimensional dissociation constants of β2AR dimers

The number densities of β2AR molecules existing as (true) dimers and the number density

of the expressed β2AR molecules were determined by single-molecule imaging of

ATTO594-ACP-β2AR[4]. Colocalisations of two fluorescent spots (molecules) of the same color

(the same ATTO594 probe) or those of two different colors (YFP for BiFC and ATTO594)

were detected when the two fluorescent molecules became localised within 220 nm of each

other[4,28]. The number densities of incidental colocalisations at given molecular densities

were subtracted[4]. The two-dimensional dissociation constant (2D-K_D) of β2AR in the PM was

determined by using the method previously developed by us[4].

To directly detect the actual binding of two β2AR molecules, forming homo-dimers, we

employed the bimolecular fluorescence complementation (BiFC) method[4,34]. The details of

the BiFC method are described in the next subsection.

336   To determine the lifetime of β2AR dimers using single color observations (with

337 ATTO594-ACP-β2AR), the duration of each colocalisation-codiffusion event was

338 measured[4,12], and after the durations of many events were determined, histograms showing

339 the distribution of colocalisation durations were obtained. All of the histograms obtained

340 from the image sequences obtained at a time resolution of 33 ms could be fitted by single

341 exponential functions, which provided decay time constants (lifetime; $\tau_{observed}$). $\tau_{observed}$ was

342 then corrected for the photobleaching lifetime of the fluorescent probe ATTO594 (τ_{bleach} =

343 7.21 ± 0.81 s for 30 Hz [n = 300] and 0.812 ± 0.041 s for 250 Hz [n = 539]), to obtain the

344 duration of apparent colocalisation ($\tau_{apparent\ colocalisation}$), using the equation [$\tau_{observed}^{-1}$ -

345 $2\tau_{bleach}^{-1}$]$^{-1}$ [4,7,12]. The effective dimer lifetime or the effective binding lifetime of proteins, τ,

346 was evaluated by subtracting the duration of incidental colocalisation (without specific

347 molecular interactions; $\tau_{incidental}$) from $\tau_{apparent\ colocalisation}$. For details, see our previous

348 publications[4,7]. $\tau_{incidental}$ was evaluated by using a monomer reference molecule, ACP linked

349 to the transmembrane domain of the low-density lipoprotein receptor, called ACP-TM[7], and

350 was found to be 19 ± 0.46 ms (exponential lifetime; n = 241 examined dimers; 39 examined

351 movies; No. of Freedoms = 97).

353 **Observations of the recruitment of Gαs and Gβγ to βAR monomers and dimers**

354 The recruitment of Gαs and Gβγ to βAR monomers and dimers was detected by observing

355 the colocalisation-codiffusion of mGFP-conjugated Gαs and Gβγ with ATTO594-labelled

356 β2AR monomers and dimers, at the level of single molecules. The colocalisation events

357 lasting for longer than 66 ms (2 video frames) were analysed. The frequency of incidental

358 colocalisation was estimated by superimposing an image sequence with one color on

359 another image sequence with a different color that was artificially shifted by ~500 nm

360 (horizontally), and was subtracted. The recruitment frequencies of Gαs and Gβγ to βAR

361 monomers and dimers (per sec per μm^2) were normalised by the number densities of Gαs

362 (or Gβγ) and β2AR protomers in the PM.

The BiFC method for examining the formation of β2AR dimers (the interaction of two β2AR molecules at the molecular level)

The YFP-based BiFC(N) and BiFC(C) moieties employed previously[4,35] were respectively fused to the C terminus of ACP-β2AR (**Extended Data Fig. 1**) in the pEGFP-N1 vector (Invitrogen; the cDNA region encoding EGFP was removed). These proteins were co-expressed in L cells, by transfection with LipofectAMINE LTX (Life Technologies) according to the manufacturer's recommendations, and then ATTO594 was conjugated to the ACP moieties of ACP-β2AR-BiFC(N) and BiFC(C). As the monomer reference molecule for the BiFC data, ACP-TM was employed and the YFP-based BiFC(N) and BiFC(C) moieties were fused to the C terminus of ACP-TM, as previously described[4]. Using dual-color single-molecule imaging (ATTO594 and mGFP), the number densities of the expressed molecules (the sum of the number densities of ACP-conjugated molecules) and those of the fluorescence spots of reconstituted BiFC molecules (spots/μm^2) were determined.

Evaluation of the changes of the cytosolic cAMP levels

The amounts of cAMP in the cytosol were measured by using the time-resolved fluorescence resonance energy transfer (TR-FRET) kit from Perkin Elmer (EnVision Xcite), according to the manufacturer's recommendations. Since the L cells used for single-molecule experiments expressed very low levels of β2AR, and thus the basal level of cAMP in the resting state was below the detection limit, we first established a stable L-cell line expressing β2AR at high levels[36], and used these cells to evaluate the changes of the cAMP levels in the cytosol before and after the additions of the agonist and inverse agonists. The drug treatments were performed as described at the end of the subsection, "Cell culture, cDNA construction and expression in L cells, and treatment of cells with agonists and inverse agonists".

Generation of artificial β2AR dimers and observations of Ca^{2+} mobilisation

To artificially generate stable homo-dimers of β2AR, we employed a chemical dimerising

system based on the FK506-binding protein (FKBP)-ligand interface[37]. First, the cDNA

encoding Halo7-β2AR-FKBP_F36V (β2AR-FKBP) was cloned into the expression plasmid

vector pEGFP-N1 (Clontech) after the removal of the cDNA region encoding EGFP, and then

the cDNA encoding FKBP_F36V was replaced by the cDNA encoding FKBP (obtained from

the YFP-FKBP cDNA, the Addgene plasmid 20175 donated by T. Meyer[38]). Halo7-β2AR-FKBP

expressed in L cells was fluorescently labelled by incubating the cells for 24 - 48 h with 5 nM

Halo-tag conjugated with SaraFluor 650T (Goryo-Kayaku;

SaraFluor650T-Halo7-β2AR-FKBP). The crosslinker for FKBP, AP20187 (Takara), was added

at a final concentration of 5 nM during the image acquisition, and dimer formation was

detected at the level of single molecules.

For Ca^{2+} mobilisation assays, a freshly-prepared 1 mg/ml DMSO solution of Fluo4-AM

(Dojindo) was added to HBSS buffered with 10 mM

(4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid) at pH 7.4 (H-HBSS), containing 1.25

mM Probenecid (Dojindo) and 0.04 % (w/v) Pluronic F127 (Dojindo) (loading buffer), to

obtain a final Fluo4-AM concentration of 5 μg/ml (4.6 μM), basically following the

manufacturer's recommendations. The L cells prepared as above for detecting dimers of

SaraFluor650T-Halo7-β2AR-FKBP at the single-molecule level were incubated with the

loading buffer at 37°C for 1 h and washed with H-HBSS three times, and then 500 μl

H-HBSS was added for epi-fluorescence microscopic observations. Fluo4 fluorescence was

recorded at 2 Hz. The stable dimers of SaraFluor650T-Halo7-β2AR-FKBP were induced by

the addition of 5 nM AP20187, as described above. To determine the saturation levels of the

fluorescence signal intensity of Fluo4 at higher concentrations of Ca^{2+}, 1 μM (final

concentration) ionomycin (Wako) was added to the cells to equalise the intracellular Ca^{2+}

concentrations with that outside the cells (1.3 mM).

Software and statistical analysis

Superimpositions of image sequences obtained from dual-color single fluorescent molecule

observations and single fluorescent spot tracking were performed using C++-based

computer programs produced in-house, as described previously[28,30]. Statistical analysis was

performed by the two-sided Welch's t test (except for the data shown in **Extended Data**

Fig. 5c, where the one-sided t test was used) or the Brunner-Munzel statistical test, by

using the R Studio Software (R Foundation for Statistical Computing,

https://www.R-project.org/). P values less than 0.05 were considered to be statistically

significant. Linear and non-linear curve fittings were performed by using OriginPro 9.1 for

Windows, with appropriate scripts.

Data and code availability

All data generated or analysed for this study are available within the paper and its

associated Extended Data Figures and Tables. All other data presented and software codes

are available upon request from the corresponding author. A **Reporting Summary** for this

paper is available.

References

1   Seifert, R. & Wenzel-Seifert, K. Constitutive activity of G-protein-coupled receptors: cause of

disease and common property of wild-type receptors. *Naunyn-Schmiedeberg's Arch.*

Pharmacol. **366**, 381-416 (2002).

2   Nakashima, A. *et al.* Agonist-independent GPCR activity regulates anterior-posterior

targeting of olfactory sensory neurons. *Cell* **154**, 1314-1325 (2013).

3   Wacker, D., Stevens, R.C. & Roth, B.L. How Ligands Illuminate GPCR Molecular

Pharmacology. *Cell* **170**, 414-427 (2017).

4   Kasai, R.S. *et al.* Full characterization of GPCR monomer-dimer dynamic equilibrium by single

molecule imaging. *J. Cell Biol.* **192**, 463-480 (2011).

5   Calebiro, D. *et al.* Single-molecule analysis of fluorescently labeled G-protein-coupled

receptors reveals complexes with distinct dynamics and organization. *Proc. Natl. Acad. Sci.*

USA **110**, 743-748 (2013).

6   Hern, J.A. *et al.* Formation and dissociation of M1 muscarinic receptor dimers seen by total

internal reflection fluorescence imaging of single molecules. *Proc. Natl. Acad. Sci. USA* **107**,

2693-2698 (2010).

7   Kasai, R.S., Ito, S.V., Awane, R.M., Fujiwara, T.K. & Kusumi, A. The Class-A GPCR Dopamine

D2 Receptor Forms Transient Dimers Stabilized by Agonists: Detection by Single-Molecule

Tracking. *Cell Biochem. Biophys.* **76**, 29-37 (2018).

8   Tabor, A. *et al.* Visualization and ligand-induced modulation of dopamine receptor

dimerization at the single molecule level. *Sci. Rep.* **6**, 33233 (2016).

9   James, J.R., Oliveira, M.I., Carmo, A.M., Iaboni, A. & Davis, S.J. A rigorous experimental

framework for detecting protein oligomerization using bioluminescence resonance energy

transfer. *Nat. Methods* **3**, 1001-1006 (2006).

10   Mercier, J.F., Salahpour, A., Angers, S., Breit, A. & Bouvier, M. Quantitative assessment of

beta 1- and beta 2-adrenergic receptor homo- and heterodimerization by bioluminescence

resonance energy transfer. *J. Biol. Chem.* **277**, 44925-44931 (2002).

11   Tennenberg, S.D., Zemlan, F.P. & Solomkin, J.S. Characterization of

N-formyl-methionyl-leucyl-phenylalanine receptors on human neutrophils. Effects of isolation

and temperature on receptor expression and functional activity. *J. Immunol.* **141**,

3937-3944 (1988).

12 Suzuki, K.G. *et al.* Transient GPI-anchored protein homodimers are units for raft organization

and function. *Nat. Chem. Biol.* **8**, 774-783 (2012).

13 Meyer, B.H. *et al.* FRET imaging reveals that functional neurokinin-1 receptors are

monomeric and reside in membrane microdomains of live cells. *Proc. Natl. Acad. Sci. USA*

103, 2138-2143 (2006).

14 Katritch, V., Cherezov, V. & Stevens, R.C. Structure-function of the G protein-coupled

receptor superfamily. *Annu. Rev. Pharmacool. Toxicol.* **53**, 531-556 (2013).

15 Ballesteros, J.A. *et al.* Activation of the beta 2-adrenergic receptor involves disruption of an

ionic lock between the cytoplasmic ends of transmembrane segments 3 and 6. *J. Biol. Chem.*

276, 29171-29177 (2001).

16 O'Dowd, B.F. *et al.* Site-directed mutagenesis of the cytoplasmic domains of the human beta

2-adrenergic receptor. Localization of regions involved in G protein-receptor coupling. *J. Biol.*

Chem. **263**, 15985-15992 (1988).

17 Sungkaworn, T. *et al.* Single-molecule imaging reveals receptor-G protein interactions at cell

surface hot spots. *Nature* **550**, 543-547 (2017).

18 Schmidt, M. *et al.* A new phospholipase-C-calcium signalling pathway mediated by cyclic AMP

and a Rap GTPase. *Nat. Cell Biol.* **3**, 1020-1024 (2001).

19 Kalay, Z., Fujiwara, T.K. & Kusumi, A. Confining domains lead to reaction bursts: reaction

kinetics in the plasma membrane. *PloS one* **7**, e32948 (2012).

20 Kusumi, A. *et al.* Dynamic organizing principles of the plasma membrane that regulate signal

transduction: commemorating the fortieth anniversary of Singer and Nicolson's fluid-mosaic

model. *Annu. Rev. Cell Dev. Biol.* **28**, 215-250 (2012).

21 Kusumi, A. *et al.* Paradigm shift of the plasma membrane concept from the two-dimensional

continuum fluid to the partitioned fluid: high-speed single-molecule tracking of membrane

molecules. *Annu. Rev. Biophys. Biomol. Struct.* **34**, 351-378 (2005).

22 Davenport, A.P. *et al.* International Union of Basic and Clinical Pharmacology. LXXXVIII. G

protein-coupled receptor list: recommendations for new pairings with cognate ligands.

Pharmacol. Rev. **65**, 967-986 (2013).

23 Dixon, R.A. *et al.* Ligand binding to the beta-adrenergic receptor involves its rhodopsin-like

core. *Nature* **326**, 73-77 (1987).

24 George, N., Pick, H., Vogel, H., Johnsson, N. & Johnsson, K. Specific labeling of cell surface

proteins with chemically diverse compounds. *J. Am. Chem. Soc.* **126**, 8896-8897 (2004).

25 Yu, J.Z. & Rasenick, M.M. Real-time visualization of a fluorescent G(alpha)(s): dissociation of

the activated G protein from plasma membrane. *Mol. Pharmacol.* **61**, 352-359 (2002).

26 Janetopoulos, C., Jin, T. & Devreotes, P. Receptor-mediated activation of heterotrimeric

G-proteins in living cells. *Science* **291**, 2408-2411 (2001).

27 Iino, R., Koyama, I. & Kusumi, A. Single molecule imaging of green fluorescent proteins in

living cells: E-cadherin forms oligomers on the free cell surface. *Biophys. J.* **80**, 2667-2677

(2001).

28 Koyama-Honda, I. *et al.* Fluorescence imaging for monitoring the colocalization of two single

molecules in living cells. *Biophys. J.* **88**, 2126-2136 (2005).

29 Hiramoto-Yamaki, N. *et al.* Ultrafast diffusion of a fluorescent cholesterol analog in

compartmentalized plasma membranes. *Traffic* **15**, 583-612 (2014).

30 Fujiwara, T., Ritchie, K., Murakoshi, H., Jacobson, K. & Kusumi, A. Phospholipids undergo

hop diffusion in compartmentalized cell membrane. *J. Cell Biol.* **157**, 1071-1081 (2002).

31 Gelles, J., Schnapp, B.J. & Sheetz, M.P. Tracking kinesin-driven movements with

nanometre-scale precision. *Nature* **331**, 450-453 (1988).

32 Murakoshi, H. *et al.* Single-molecule imaging analysis of Ras activation in living cells. *Proc.*

Natl. Acad. Sci. USA **101**, 7317-7322 (2004).

33 Tsunoyama, T.A. *et al.* Super-long single-molecule tracking reveals

dynamic-anchorage-induced integrin function. *Nat. Chem. Biol.* **14**, 497-506 (2018).

34 Hu, C.D., Chinenov, Y. & Kerppola, T.K. Visualization of interactions among bZIP and Rel

family proteins in living cells using bimolecular fluorescence complementation. *Mol. Cell* **9**,

789-798 (2002).

35   Briddon, S.J. & Hill, S.J. Pharmacology under the microscope: the use of fluorescence

correlation spectroscopy to determine the properties of ligand-receptor complexes. *Trends*

Pharmacol. Sci. **28**, 637-645 (2007).

36   Yusa, K., Zhou, L., Li, M.A., Bradley, A. & Craig, N.L. A hyperactive piggyBac transposase for

mammalian applications. *Proc. Natl. Acad. Sci. USA* **108**, 1531-1536 (2011).

37   Clackson, T. *et al.* Redesigning an FKBP-ligand interface to generate chemical dimerizers with

novel specificity. *Proc. Natl. Acad. Sci. USA* **95**, 10437-10442 (1998).

38   Inoue, T., Heo, W.D., Grimley, J.S., Wandless, T.J. & Meyer, T. An inducible translocation

strategy to rapidly activate and inhibit small GTPase signaling pathways. *Nat. Methods* **2**,

415-418 (2005).

Fig. 1

Fig. 1. β2AR dynamically interconverts between monomers and transient homodimers.

Mean, SEM, the number of experiments, and other statistical parameters are summarised in

Extended Data Table 1.

(a) A typical snapshot sequence (among 13 movies) showing transient homo-dimerisation of β2AR

molecules. Two molecules became colocalised at frame 2 (67 ms), diffused together for 134 ms

(colocalisation), and then separated.

(b) Histogram showing the distribution of the durations of the WT-β2AR colocalisation events

(homo-dimers). Note that the $\tau_{observed}$ is the value before corrections for photobleaching and

incidental colocalisations.

(c) Summary of the dimer lifetimes (after corrections) of WT and mutant β2AR obtained before and

after the additions of various reagents. ** and *n.s.*: statistically significant and non-significant with *P*

values smaller or greater than 0.05, respectively. This convention was used throughout this report.

Fig. 2

Fig. 2. Determining the 2D-*K*D of β2AR dimers with/without an inverse agonist and the BiFC detection of β2AR dimers. Mean, SEM, the number of experiments, and other statistical parameters are summarised in **Extended Data Table 2**.

The number density of β2AR molecules in true dimers (after subtraction of incidental overlap of fluorescent spots; ± inverse agonist; fitted with Eq. 5 in Ref4; red curve) **(a)** and the number densities of the BiFC spots for ACP-β2AR-YFP(N/C) and ACP-TM-YFP(N/C) (monomer reference molecule) **(b)**, plotted as a function of the number density of expressed molecules (*NDmol*). In **b**, the lines are provided to facilitate visualisation.

Fig. 3

Fig. 3. Non-stimulated β2AR homo-dimers recruit trimeric G-proteins: a process

suppressed by inverse agonists. Mean, SEM, the number of experiments, and other statistical

parameters are summarised in **Extended Data Table 3**.

(a) A typical snapshot image sequence (among 40 movies) showing that a trimeric G-protein Gαs

subunit transiently binds to a β2AR monomer.

(b, c) The inverse agonists, ICI-118,551 and Timolol, inhibitors of β2AR basal constitutive activities,

blocked the recruitment of trimeric G-proteins to β2AR homo-dimers, but not that to monomers. Gαs

(b) and Gβγ **(c)** colocalisation frequencies with β2AR monomers and homo-dimers. The normalised

frequencies of colocalisations = the number of colocalisation events per sec per μm² normalised by

the number densities of Gαs (or Gβγ) and β2AR protomers.