Association of gut microbiota with metabolism in juvenile Atlantic Salmon

Dvergedal, H.[1], Sandve, S.R.[1*], Angell, I.L.[2], Klemetsdal, G.[1], Rudi, K.[2]

[1]Department of Animal and Aquacultural Sciences, Faculty of Biosciences, Norwegian University

of Life Sciences, P. O. Box 5003, NO-1433, Aas, Norway

[2]Faculty of Chemistry, Biotechnology and Food Science, Norwegian University of Life Sciences,

P. O. Box 5003, NO-1433, Aas, Norway

*Correspondence: Simen Rød Sandve, Norwegian University of Life Sciences, NO-1433 Aas, Norway**. simen.sandve@nmbu.no**

Abstract

The gut microbiome plays a key role in animal health and metabolism through the intricate functional interconnection between the feed, gut microbes, and the host. Unfortunately, in aquaculture, the links between gut microbes and fish genetics and production phenotypes are not well understood.

In this study, we investigate the associations between gut microbial communities, fish feed conversion, and fish genetics in the domestic Atlantic salmon. Microbial community composition was determined for 230 juvenile fish from 23 full-sib families and was then regressed on growth, carbon and nitrogen metabolism, and feed efficiency. We only found weak associations between host genetics and microbial composition. However, we did identify significant ($p < 0.05$) associations between the abundance of three microbial operational taxonomical units (OTUs) and fish metabolism phenotypes. Two OTUs were associated with both carbon metabolism in adipose tissue and feed efficiency, while a third OTU was associated with weight gain.

In conclusion, this study demonstrates an intriguing association between host lipid metabolism and the gut microbiota composition in Atlantic salmon.

Keywords: microbiome, Atlantic salmon, genetics, metabolism, feed efficiency, carbon turnover

Background

Efficient and environmentally sustainable animal production systems are urgently required to

ensure long-term food security, especially as global aquaculture consumption is projected to

double by 2050 (www.fao.org). One important aspect of improving sustainability is to improve

feed conversion and growth. In humans and other vertebrate systems, the gut microbiome plays a

central role in the path from "feed-to-animal" [1–4], and recent studies have also shown that host-

genetic factors can modulate microbiome composition. Such functional interconnection between

feed, microbes, and host (i.e. the feed-microbiome-host axis), opens up intriguing avenues for

optimizing aquaculture production systems, for example by breeding for 'optimized' microbiome

composition[5].

Yet, even though the dietary composition is known to impact the gut microbiome in aquaculture

species[1], almost nothing is known about the link between the gut microbiota and important

production phenotypes, or to what extent microbiota composition itself could be a new breeding

target for aquaculture breeding programs[5].

To address this pressing knowledge gap we use a family-based experimental design to test if

variation in the gut microbiome composition in juvenile Atlantic salmon is associated with key

phenotypes related to host metabolism as well as variation in host genetics. Our results identified

phenotypic associations between host gut microbiome and lipid metabolism, growth, as well as to

feed efficiency, which open the possibility for metabolic modulation through the gut microbiota.

Materials and methods

Experimental setup

A family experiment with Atlantic salmon was carried out at the fish laboratory, Norwegian

University of Life Sciences (NMBU), Aas, Norway, according to the laws and regulations

controlling experiments on live animals in EU (Directive 2010/637EU) and Norway (FOR-2015-

06-18-761). The experiment was approved by the Norwegian Food Safety Authority (FOTS ID

11676). The experiment included 23 full-sib families from AquaGen's breeding population

comprising of 230 fish, with each family being distributed on two tanks. Five fish, representing 10

fish from each family was investigated (Fig. 1).

Figure 1. Schematic outline of the experimental setup. Twenty-three families were distributed on 46 tanks (two
tanks per family). Dominant live bacteria were identified prior to association analyses to fish genetics and metabolism.

Microbiota analyses

Distal intestinal samples (n = 230) were obtained by squeezing out the gut segment content using

sterile tweezers in 1 ml phosphate buffered saline (PBS) and put on ice until further processing.

To distinguish between DNA from dead and alive bacteria, the samples were treated with

propidium monoazide (PMA) within eight hours post sampling. Samples were pulse centrifuged

up to 1200 rpm and split in two, where one part (n = 230, PMA treated samples) was added PMA

dye (Biotium, USA to a final concentration of 50 µM, and the other part was kept as a control (n

= 230, non-PMA treated samples) with no added PMA. The samples were then kept dark for 5 min

before exposure to light for 30 min in a lightbox from Geniul. DNA extraction (n = 460) was done

using mag midi DNA extraction kit (LGC Genomics, UK) following the manufacturer's

recommendations.

The 16S rRNA amplicon library was prepared and sequenced as previously described [6]. The

sequence reads were processed using USEARCH v8 [7] where the sequences were paired-end

joined, demultiplexed, and quality filtered (maxxee = 1.0, minlength = 350, singeltons discarged),

before operational taxonomic unit (OTU) clustering with 97% identity threshold was performed

using the UPARSE pipeline [8]. Taxonomy assigment was done using SILVA database [9]. Diversity

analysis was done using a sequence depth of 10 000 sequences per sample.

To filter out OTUs from dead bacteria and bacteria considered as contaminants, filtering was done

using the following criteria on each individual fish gut microbiome; OTUs which showed a more

than 3-fold reduction in the PMA treated sample was considered dead, while OTUs that showed a

more than 6-fold increase in the non PMA treated sample were considered contaminants because

there were no other alternative explanations. Out of the 230 fish gut microbiomes, 188 passed the

sequence quality control filtering criteria, including rarefaction at 10 000 sequences and

live/dead/contaminantion screening. Correlations between OTUs were determined using

Spearman's rank correlation coefficient.

Raw 16S rRNA sequence data are deposited in the SRA database under the accession number

PRJNA590084.

Phenotypic data

A diet labeled with the stable isotopes [15]N and [13]C was fed during the experimental period of 12

days. The host metabolism related traits analyzed are listed in Table 1. Details for phenotypic data

for growth and metabolic traits are explained in Dvergedal et al. [10].

Table 1. Description of the 13 variables phenotyped.

No.	Variables	Description
1	IW	Initial weight [1]
2	FW	Final weight [1]
3	WG	Weight gain [1] (FW-IW)
4	RG	Relative weight gain (%) (((FW-IW)/FW)100)
5	AMC	Atom % [13]C in muscle
6	AMN	Atom % [15]N in muscle
7	ALC	Atom % [13]C in liver
8	ALN	Atom % [15]N in liver
9	AAC	Atom % [13]C in adipose tissue
10	IFCR_AMC	Individual isotope-based indicator of feed conversion ratio, from Atom % [13]C in muscle
11	IFCR_AMN	Individual isotope-based indicator of feed conversion ratio, from Atom % [15]N in muscle
12	IFER_AMC	Individual isotope-based indicator of feed efficiency ratio, from Atom % [13]C in muscle
13	IFER_AMN	Individual isotope-based indicator, of feed efficiency ratio, from Atom % [15]N in muscle
14	OTU1	Operational taxonomic unit 1, classified as *Caulobacteriaceae*
15	OTU2	Operational taxonomic unit 2 classified as *Pseudomonas fluorescens*
16	OTU3	Operational taxonomic unit 3 classified as *Sphingomonas*
17	OTU5	Operational taxonomic unit 5 classified as *Bradyrhizobium*
18	OTU6	Operational taxonomic unit 6 classified as *Ralstonia* sp
19	OTU7	Operational taxonomic unit 7 classified as *Pseudoalteromonas*

Outlier detection

To obtain approximate normality of the relative abundances of OTUs we transformed the OTU

data using the natural logarithm (Ln). Influence statistic was used for outlier detection by

regressing Ln (OTU) on all the phenotypes (Table 1) using PROC REG in SAS®. The cutoff value

for outliers was calculated as $3p/n$ (> 0.10), where n is the number of samples (i.e. animals) used

to fit the model (n = 188), and *p* is the number of parameters in the model. A total of 16 outliers

were detected and deleted.

Estimation of heritability

To estimate heritabilities of the microbiota at the level of each OTU we first did a single-trait

analysis of variance of the Ln (OTUs). In each analysis, the model was:

$[y] = [X][b] + [Z_a][a] + [Z_t][t] + [e],$ **[1]**

where $[y]$ is a vector of individual OUT 'phenotypes' (i.e. the trait), $[b]$ is a vector of fixed effects,

including sampling *day$_i$* (i = 1-4), $[a] \sim N(0, \mathbf{G}\sigma_a^2)$ is a vector of random additive genetic effects

for the trait, $[t] \sim N(0, \mathbf{I}\sigma_t^2)$ is a vector of random tank effects for the trait, and $[e] \sim N(0, \mathbf{I}\sigma_e^2)$, is a

vector of random residuals for the trait. The **X** and **Z** matrices are corresponding incidence

matrices, $\mathbf{G}\sigma_a^2$ is the genomic (co)variance matrix, $\mathbf{I}\sigma_t^2$ is the (co)variance matrix due to tank

effects, and $\mathbf{I}\sigma_e^2$ denotes the error (co)variance matrix. The number of phenotyped individuals was

rather low (n = 172), and the genomic relationship matrix was generated according to VanRaden's

first method[11]. The matrix **G** (2282x2282) was calculated based on a subset of 51,543 SNPs of

high genotype quality, covering all autosomal chromosomes (AquaGen's custom Axiom®SNP

genotyping array from Thermo Fisher Scientific (San Diego, CA, USA) includes 56,177 single-

nucleotide polymorphisms).

Heritabilities of the OTUs were estimated as: $h^2 = \frac{\sigma_a^2}{\sigma_a^2 + \sigma_t^2 + \sigma_e^2}$, where σ_a^2, σ_t^2, and σ_e^2 are the

estimates of the individual additive genetic, tank environmental, and individual residual variance,

respectively, of the trait. The fraction of variance explained by the tank was estimated as: $c^2 =$

$\frac{\sigma_t^2}{\sigma_a^2 + \sigma_t^2 + \sigma_e^2}$. Significance of the genetic effect was tested using a likelihood-ratio (LR) test-statistic,

comparing a single-trait model with genetic effects (H_1) to a model without genetic effects (H_0):

$LR = 2\left(\left(log\, L\, |\hat{\theta}_{H_1}\right) - \left(log\, L\, |\hat{\theta}_{H_0}\right)\right).$

The genetic effect was considered significant if $LR > \chi^2_{(\alpha=0.05;\, df=1)}.$

Genome-wide association analysis

To associate variation in microbial composition with host genetics a genome-wide association

study was done using Ln (OTUs) as response variables. The analysis was carried out by a linear

mixed-model algorithm implemented in a genome-wide complex trait analysis (GCTA) [12]. The

leave one chromosome out option (*--mlm-loco*) was used, meaning that the chromosome harboring

the SNP tested for was left out when building the genetic relationship matrix (GRM). The linear

mixed model can be written:

$Y_i = a + bx + g_i^- + \varepsilon_i,$ **[2]**

where Y_i is one of the Ln (OTUs) of individual i, a is the intercept, b is the fixed regression of the

candidate SNP to be tested for association, x is the SNP genotype indicator variable coded as 0, 1

or 2, g_i^- is the random polygenic effect for individual $i \sim N\,(0,\, \mathbf{G}\sigma_g^2\,)$ where $\mathbf{G}$ is the GRM and σ_g^2

is the variance component for the polygenic effect, and ε_i is the random residual. In this algorithm,

σ_g^2 is re-estimated each time a chromosome is left out from the calculation of the GRM. The dataset

was filtered, and individuals with < 10% missing genotypes were kept ($n = 2279$). Further, it was

required that SNPs should have minor allele frequency (MAF) $\geq$ 1% and a call rate > 90%. After

filtering 54,200 SNPs could be included in the analysis. The level of significance for SNP was

evaluated with a built-in likelihood-ratio test, and the threshold value for genome-wide

significance was calculated by the use of Bonferroni correction $(0.05/54200) = 9.23 \times 10^{-7}$,

corresponding to a -$\log_{10}$ *p-value (p)* of 6.03.

Association between OTUs and fish phenotypes

We examined the association between microbiota and several individual fish phenotypes related

to the metabolism of the fish, including growth, nutrient turnover, and feed efficiency parameters

(Table 1). This phenotypic association were tested with a linear mixed-effect model:

$$[y] = [X][b] + [Z_a][a] + [Z_t][t] + [e], \qquad\qquad [3]$$

where, $[b]$ is a vector of fixed effects for the trait containing the regressions of sampling day (Day)

and the Ln (OTUs), while the remaining are described with **[1]**. In preceding analyses, we

experienced a strong co-linearity between the tank and the family effects, which in consequence

led us to analyze for the phenotypic association between production variables and the OTUs with

a model accounting for the genomic relationships between individuals (also known as a genomic

selection model).

The genetic analyses in **[1]** and **[3]** were carried out using the ASReml4 software package[13].

Results

Overall microbiota composition

A total of 704 OTUs were defined, with the dead bacterial fraction representing 9.1% of the

sequencing reads (contained in 254 OTUs). In addition, a fraction of 0.003% of the sequencing

reads (contained in 146 OTUs) were considered as contaminations (Fig. 2A). Among the 304

OTUs passing the live/dead/contamination filtering we identified a clear over-representation of

Proteobacteria, both with respect to OTU prevalence and quantity (Fig. 2B). OTU1

Caulobacteraceae and OTU2 *Pseudomonas fluorescens* dominated, with mean abundances of

32.9% and 34.8%, respectively. There were 6 OTUs with a mean abundance > 1 % in all the

samples. We detected an overall negative correlation between OTU1 *Caulobacteraceae* to the

other OTUs (Fig. 2C).

Figure 2. General characteristics of gut microbiota. (A) Fractions of dead, live, and contamination OTUs. Fractions
calculated based on the ratio of OTU counts in PMA treated samples versus untreated samples. (B) Prevalence
(number of OTUs) and quantity (number of sequencing reads) for the different phyla in the most abundant OTUs. (C)
Spearman correlations between the dominant live bacteria (high-abundant OTUs). The high abundant OTUs were
identified as those that have an abundance of > 1% on average in all the samples.

Effects of rearing tank and host genetics on the gut microbiome composition

To assess the contribution of genetics (i.e. heritability) and environment (i.e. tanks effect) in

driving the variation in microbial composition between individual fish we applied analyses of

variance, using tank as a covariate. Although we classified 304 OTUs across all gut microbiomes,

the top 6 most abundant OTUs represented 85% of the 16S sequences in our dataset. We, therefore,

conducted these analyses using only these OTUs. The results (Table 2) showed small genetic

components for OTUs 1 and 3 and only a small non-significant tank effect for OTU1 (0.03).

However, neither the tank nor genetic effects were significantly different from zero ($p > 0.05$).

However, the standard errors were large, meaning that the experiment did not have the power to

estimate these components precisely. This is supported by the estimates of variance components

for genetic or tank effects (or both) of the remaining OTUs being zero (i.e. restricted to the

boundary of the parameter space) (Table 2).

Table 2. Estimates of the genetic, tank and residual variance components (σ_a^2, σ_t^2 and σ_e^2, respectively), the
fraction of phenotypic variance explained by the environmental tank effect (c^2), heritability (h^2), as well
as the χ^2 statistics for the additive genetic family effect, with the corresponding level of significance (p).

	σ_a^2	σ_t^2	σ_e^2	c^2	h^2	χ^2	p
OTU1	0.007	0.009	0.25	0.03 ± 0.08	0.03 ± 0.09	0.13	0.72
OTU2	0.00_1	0.00_1	0.14	0	0	0	1
OTU3	0.007	0.00_1	0.58	0	0.10 ± 0.09	1.72	0.19
OTU5	0.00_1	0.00_1	0.13	0	0	0	1
OTU6	0.00_1	0.00_1	0.07	0	0	0	1
OTU7	0.00_1	0.00_1	0.18	0	0	0	1

$_1$Restricted on the boundary of parameter space

Finally, we utilize the existing genotyping data for these fish[10] to perform a genome-wide

association analyses for the OTU abundances. No genome-significant associations between SNPs

and OTUs were identified, however, the Manhattan plots show clear peaks at chromosomes 14,

24, 3, and 5 (Suppl. Fig. 1), with some SNPs having significant associations to OTU1 and OTU2

at the chromosome level (Suppl. Fig. 2, Suppl. Table 1).

Community structure is associated with fish growth and metabolism

Linear regressions were used to examine the phenotypic relationship between the gut microbiome

and host metabolism traits (i.e. growth, nutrient turnover, and feed efficiency, see table 1). Indeed,

these analyses (Table 3) do indicate a link between the production variables and the gut

microbiome (significant associations in Table 3, for all results see Suppl. Table 2).

Table 3. Regression estimates, standard errors, F- and p-values when regressing OTUs on growth,
metabolism, and feed efficiency variables. The model also contained regression on day and random effects
of animal (utilizing genomic relationships) and tank, for which variance components are included.

215
216
217

Dependent variable	Variables	Estimate	Stderr	F-value	p-value	Variance component
WG	Day	3.338	0.177	355.73	<0.005	
	OTU1	-0.324	0.376	0.15	NS	
	OTU2	2.191	1.199	0.04	NS	
	OTU3	-1.352	0.624	4.66	<0.05	
	OTU5	-0.743	0.535	1.78	NS	
	OTU6	-0.440	0.866	0.26	NS	
	OTU7	0.179	0.492	0.30	NS	
	Tank					13.84 ± 3.25
	Animal					5.54 ± 0.69
AAC	Day	0.082	0.019	17.95	<0.005	
	OTU1	0.007	0.003	1.39	NS	
	OTU2	-0.016	0.009	2.15	NS	
	OTU3	-0.001	0.005	1.31	NS	
	OTU5	0.006	0.004	4.63	<0.05	
	OTU6	-0.001	0.007	0.01	NS	
	OTU7	0.010	0.004	8.53	<0.005	
	Tank					0.90 ± 0.20
	Animal					$0.87 \times 10^{-4} \pm 2.21 \times 10^{-4}$
IFER_AMC	Day	0.365	0.021	297.49	<0.005	
	OTU1	0.043	0.026	5.55	<0.03	
	OTU2	-0.101	0.083	0.41	NS	
	OTU3	0.018	0.043	1.34	NS	
	OTU5	-0.023	0.037	0.03	NS	
	OTU6	0.035	0.060	0.34	NS	
	OTU7	0.081	0.034	5.22	<0.03	
	Tank					0.20 ± 0.04
	Animal					$0.43 \times 10^{-2} \pm 1.51 \times 10^{-3}$
IFER_AMN	Day	0.149	0.009	273.21	<0.005	
	OTU1	0.020	0.009	3.89	<0.05	
	OTU2	-0.026	0.027	2.86	NS	
	OTU3	0.001	0.014	0.64	NS	
	OTU5	-0.016	0.012	0.79	NS	
	OTU6	0.008	0.020	0.16	NS	
	OTU7	0.027	0.011	5.83	<0.03	
	Tank					$0.04 \pm 7.86 \times 10^{-3}$
	Animal					$0.29 \times 10^{-3} \pm 1.51 \times 10^{-4}$

OTU3 (*Sphingomonas)* regressed negatively on weight gain ($p < 0.05$), OTU1

(*Caulobacteriaceae)* and OTU7 (*Pseudoalteromonas)* regressed positively on feed efficiency

indicators ($p < 0.05$), while OTU7 ($p < 0.005$) and OTU5 (*Bradyrhizobium)* ($p < 0.05$) regressed

positively on carbon metabolism in adipose tissue (AAC variable).

Discussion

A major strength of this experiment is that each individual fish microbiome can be linked to

detailed individual-level phenotypes of growth, feed efficiency, and nutrient turnover as measured

by the use of stable-isotope profiling in the liver, muscle, and adipose tissues[10]. Intriguingly, the

phenotypic associations between OTUs and fish production-related phenotypes revealed several

significant relationships (Table 3). We observed significant positive associations between lipid

carbon metabolism (the AAC phenotype) and OTU5 ($p < 0.05$) and OTU7 ($p < 0.005$). These

OTUs belong to the genera *Bradyrhizobium,* and *Pseudoalteromonas*, respectively.

Pseudoalteromonas is known to have the capacity to produce a range of biologically active

extracellular compounds, ranging from antimicrobial compounds and proteases to compounds

important for host metamorphosis[14]. This genus has also been used as probiotics in fish farming[15].

Bradyrhizobium, on the other hand, is a widespread environmental bacterium capable of degrading

aromatic compounds and nitrogen fixation[16]. However, the potential mechanisms for the bacterial

associations with lipid metabolism in salmon are completely unknown. The *Sphingomonas* OTU3

showed a significant negative association with weight gain. This genus has previously been

associated with antibiotic resistance connected to disease treatment of juvenile salmon in fresh-

water[17], which may indicate that reduced weight gain could be connected to the opportunistic

properties of *Sphingomonas*[18]. Lastly, OTU1 belonging to *Caulobacteriacea* showed a strong

negative correlation with the other OTU (Fig. 2C), and a significant positive association with two

of the feed utilization efficiency metrics (Table 3), which also had a positive association to OTU7.

One interpretation of this is that *Caulobacteriacea* has a mutualistic association with juvenile

salmon, possibly by protecting juvenile salmon against opportunistic infections.

The association between OTUs and the lipid carbon metabolism (AAC) in Atlantic salmon means

that fish with higher abundances of these microbes also convert the feed protein fraction to lipids

at a higher rate. However, the mechanisms driving this association remains elusive. One possibility

is that the Atlantic salmon gut microbiota has a direct impact on the production of biomolecules

in the distal intestine which are readily absorbed and deposited as fat in adipose tissues. It is worth

noting that Dvergedal et al. [10] have reported that fish with a higher turnover of carbon in lipid

tissues also has improved feed efficiency (see IFER variable in Tables 1 and 3) and fast growth.

In other words, the fish with high carbon turnover in lipid tissues will likely have a positive energy

balance and therefore also the opportunity to convert more surplus energy into lipids for storage.

We did however not observe associations between gut microbes and nutrient turnover in muscle

or liver. This lack of association with nitrogen turnover could be because the majority of the protein

fraction is digested and absorbed before the distal intestine where our microbial samples were

collected from. It is also possible that the associations between microbial composition and fish

metabolism are driven by indirect factors. Since growth is positively correlated with feed intake[10],

these fish might also have increased passage rate in the gastrointestinal tract due to high feed

intake. This could indirectly affect the competition and balance among microbes and thereby shift

the community structure. It is thus critical that future studies include functional meta-omics data

that can also demonstrate shifts in activities in microbial metabolic pathways.

Host genetic effects on gut microbiomes have been identified in recent studies in a wide range of

animals, including invertebrates[19], mammals[20,21], and fish[22]. Here we only found weak (and non-

significant) associations between host genetics and relative OTU abundance (Table 2,

Supplementary Fig. 1 and 2). Neither did we find rearing tank effects, in correspondence with a

study in tilapia[23]. However, it is clear from the standard errors (Table 2) that our OTU heritability

estimates were very unprecise. Hence, to assess the importance of host genetics on gut microbiome

composition in Atlantic salmon, future studies must increase sample size significantly and apply

'common rearing' experimental designs to avoid confounding tank and family effects.

Conclusion

In conclusion, our results demonstrate an association between the microbial composition in the

distal gut and a key aspect of Atlantic salmon metabolism. This association could be a direct effect

of microbes contributing to improved nutrient availability and absorption for the host.

Alternatively, these associations could be uncoupled from the microbiota function and instead

driven by feeding behavior and passage rates. Future experiments should, therefore, aim to

measure changes in microbial metabolic pathways to separate causal from correlative microbe-

host associations.

Funding

This study was supported by The Norwegian University of Life Sciences, AquaGen AS and

Foods of Norway, a Centre for Research-based Innovation (the Research Council of Norway;

grant no. 237841/O30), GenoSysFat (the Research Council of Norway Havbruk; grant no.

244164/E40), and DigiSal (the Research Council of Norway grant no. 248792).

Acknowledgments

We thank Margareth Øverland, Liv Torunn Mydland and Jørgen Ødegård for the planning of the

experiment and Jørgen Ødegård for creating the genomic relationship matrics used in this study.

SUPPLEMENTARY TABLES

Suppl. Table 1. Single-nucleotide polymorphisms (SNP) associated with the OTU variables.

Trait	Chr	SNP	bp	A1	A2	Freq	b	se	*p*	log10(*p*)
OTU1	5	ctg7180001925291_748_SCT	27518103	C	T	0.10	-0.51	0.12	2.49E-05	-4.6
	5	ctg7180001311517_3981_SAG	29290387	G	A	0.10	-0.46	0.11	2.51E-05	-4.6
	5	ctg7180001833723_2640_SAG	56739234	A	G	0.01	-1.44	0.28	4.53E-07	-6.3
	14	ctg7180001903206_2590_SCT	5592915	C	T	0.15	-0.45	0.10	1.81E-06	-5.7
	19	ctg7180001841090_969_SAG	42066887	G	A	0.02	-1.03	0.23	8.21E-06	-5.1
	19	ctg7180001800309_4785_SCT	67320228	C	T	0.09	-0.55	0.13	1.70E-05	-4.8
	19	ctg7180001645427_1176_SAG	75579801	A	G	0.02	-1.16	0.27	1.57E-05	-4.8
OTU2	1	ctg7180001908955_3685_SAG	4697317	G	A	0.06	-0.46	0.11	2.11E-05	-4.7
	5	ctg7180001722113_3376_SGT	25219965	G	T	0.01	-1.05	0.21	7.26E-07	-6.1
	5	ctg7180001672687_10612_SAG	28647949	A	G	0.01	-1.05	0.21	7.26E-07	-6.1
	6	ctg7180001847439_3403_SAC	32454164	A	C	0.03	-0.59	0.14	2.32E-05	-4.6
	6	ctg7180001828921_349_SAG	64415855	G	A	0.01	-0.96	0.20	1.30E-06	-5.9
	11	ctg7180001900871_1769_SCT	85379984	T	C	0.08	-0.39	0.09	1.13E-05	-4.9
	12	ctg7180001666651_1965_SAC	12523402	A	C	0.03	-0.70	0.16	1.56E-05	-4.8
	12	ctg7180001840586_4279_SCT	43076320	C	T	0.07	-0.47	0.10	1.92E-06	-5.7
	12	ctg7180001839373_5142_SAC	54225053	C	A	0.01	-1.14	0.21	8.12E-08	-7.1
	12	ctg7180001484203_1236_SAC	68394895	C	A	0.02	-1.09	0.20	9.88E-08	-7.0
	15	ctg7180001298386_4207_SGT	20950022	T	G	0.02	-0.95	0.21	3.77E-06	-5.4
	15	ctg7180001878210_7094_SAG	36237432	A	G	0.02	-0.78	0.18	1.97E-05	-4.7
	22	ctg7180001807745_17982_SAC	34245759	C	A	0.02	-1.08	0.21	1.40E-07	-6.9
	24	ctg7180001301776_10058_SAG	26185685	A	G	0.10	-0.39	0.09	8.01E-06	-5.1
	29	ctg7180001835668_5175_SAC	35464325	A	C	0.04	-0.76	0.13	6.34E-09	-8.2
	29	ctg7180001540342_979_SAC	37759078	C	A	0.13	-0.32	0.07	1.51E-05	-4.8
OTU5	7	ctg7180001798708_11367_SCT	14075405	C	T	0.51	0.27	0.06	2.56E-05	-4.6
	7	ctg7180001798708_5931_SAG	14080841	G	A	0.40	-0.28	0.07	3.62E-05	-4.4
	15	ctg7180001298386_4207_SGT	20950022	T	G	0.02	-1.15	0.26	1.41E-05	-4.9
OTU6	3	ctg7180001338507_1900_SCT	12338431	C	T	0.36	0.17	0.04	1.47E-05	-4.8
	3	ctg7180001916975_10480_SGT	26091556	T	G	0.43	0.18	0.04	8.24E-06	-5.1
OTU7	5	ctg7180001841488_1545_SCT	8907693	C	T	0.15	0.38	0.08	5.82E-06	**-5.2**

Suppl. Table 2. Regression estimates, standard errors, *F*- and *p*-values when regressing OTUs on growth, metabolism, and feed efficiency variables. The model also contained regression on day and random effects of animal (utilizing genomic relationships), and tank for which variance components are included.

Dependent variable	Variables	Estimate	Stderr	F-value	*p*-value	Variance component
WG	Day	3.338	0.177	355.73	<0.005	
	OTU1	-0.324	0.376	0.15	NS	
	OTU2	2.191	1.199	0.04	NS	
	OTU3	-1.352	0.624	4.66	<0.05	
	OTU5	-0.743	0.535	1.78	NS	
	OTU6	-0.440	0.866	0.26	NS	
	OTU7	0.179	0.492	0.30	NS	
	Tank					13.84 ± 3.25
	Animal					5.54 ± 0.69
RG	Day	9.969	0.522	365.90	<0.005	
	OTU1	0.983	0.854	3.64	NS	
	OTU2	-3.500	2.715	0.01	NS	
	OTU3	1.118	1.410	2.04	NS	
	OTU5	-0.149	1.214	0.06	NS	
	OTU6	1.094	1.963	0.31	NS	
	OTU7	2.121	1.117	3.29	NS	
	Tank					122.00 ± 26.75
	Animal					14.56 ± 2.56
AMC	Day	0.270	0.027	101.92	<0.005	
	OTU1	0.004	0.006	0.01	NS	
	OTU2	-0.020	0.020	0.00	NS	
	OTU3	0.006	0.010	0.42	NS	
	OTU5	0.001	0.009	0.08	NS	
	OTU6	0.010	0.014	0.49	NS	
	OTU7	0.007	0.008	0.55	NS	
	Tank					0.48 ± 0.11
	Animal					$0.41 \times 10_{-3} \pm 1.03 \times 10_{-4}$
AMN	Day	0.288	0.019	230.96	<0.005	
	OTU1	0.002	0.014	0.66	NS	
	OTU2	-0.048	0.045	0.59	NS	
	OTU3	0.018	0.023	0.99	NS	
	OTU5	0.015	0.020	0.85	NS	
	OTU6	0.020	0.032	0.37	NS	
	OTU7	0.017	0.018	0.57	NS	
	Tank					0.17 ± 0.04
	Animal					$0.41 \times 10_{-2} \pm 7.08 \times 10_{-4}$
ALC	Day	0.088	0.028	9.78	<0.005	
	OTU1	0.004	0.004	0.29	NS	
	OTU2	-0.014	0.013	0.08	NS	
	OTU3	-0.004	0.007	0.34	NS	
	OTU5	0.005	0.006	1.22	NS	
	OTU6	0.009	0.010	0.89	NS	
	OTU7	0.007	0.005	1.21	NS	
	Tank					1.83 ± 0.41
	Animal					$0.18 \times 10_{-3} \pm 4.52 x_{-4}$

313 *Suppl. Table 2. Continued*

Dependent variable	Variables	Estimate	Stderr	F-value	p-value	Variance component
ALN	Day	0.445	0.032	191.05	<0.005	
	OTU1	0.013	0.015	0.05	NS	
	OTU2	-0.061	0.047	0.00	NS	
	OTU3	0.007	0.024	0.52	NS	
	OTU5	0.012	0.021	0.88	NS	
	OTU6	0.024	0.034	0.49	NS	
	OTU7	0.034	0.019	2.58	NS	
	Tank					0.51 ± 0.11
	Animal					$0.37 \times 10^{-2} \pm 7.42 \times 10^{-4}$
AAC	Day	0.082	0.019	17.95	<0.005	
	OTU1	0.007	0.003	1.39	NS	
	OTU2	-0.016	0.009	2.15	NS	
	OTU3	-0.001	0.005	1.31	NS	
	OTU5	0.006	0.004	4.63	<0.05	
	OTU6	-0.001	0.007	0.01	NS	
	OTU7	0.010	0.004	8.53	<0.005	
	Tank					0.90 ± 0.20
	Animal					$0.87 \times 10^{-4} \pm 2.21 \times 10^{-4}$
IFCR_AMC	Day	0.231	0.020	130.26	<0.005	
	OTU1	-0.024	0.023	2.34	NS	
	OTU2	0.053	0.072	0.21	NS	
	OTU3	-0.020	0.037	1.32	NS	
	OTU5	0.019	0.032	0.09	NS	
	OTU6	-0.009	0.052	0.03	NS	
	OTU7	-0.044	0.030	2.25	NS	
	Tank					0.19 ± 0.04
	Animal					$0.24 \times 10^{-2} \pm 1.08 \times 10^{-3}$
IFCR_AMN	Day	0.554	0.047	138.65	<0.005	
	OTU1	-0.086	0.049	1.71	NS	
	OTU2	0.149	0.153	1.61	NS	
	OTU3	-0.049	0.079	1.18	NS	
	OTU5	0.083	0.069	0.85	NS	
	OTU6	-0.057	0.111	0.26	NS	
	OTU7	-0.105	0.063	2.47	NS	
	Tank					1.01 ± 0.21
	Animal					$0.64 \times 10^{-2} \pm 4.43 \times 10^{-3}$
IFER_AMC	Day	0.365	0.021	297.49	<0.005	
	OTU1	0.043	0.026	5.55	<0.03	
	OTU2	-0.101	0.083	0.41	NS	
	OTU3	0.018	0.043	1.34	NS	
	OTU5	-0.023	0.037	0.03	NS	
	OTU6	0.035	0.060	0.34	NS	
	OTU7	0.081	0.034	5.22	<0.03	
	Tank					0.20 ± 0.04
	Animal					$0.43 \times 10^{-2} \pm 1.51 \times 10^{-3}$
IFER_AMN	Day	0.149	0.009	273.21	<0.005	
	OTU1	0.020	0.009	3.89	<0.05	
	OTU2	-0.026	0.027	2.86	NS	
	OTU3	0.001	0.014	0.64	NS	
	OTU5	-0.016	0.012	0.79	NS	
	OTU6	0.008	0.020	0.16	NS	
	OTU7	0.027	0.011	5.83	<0.03	
	Tank					$0.04 \pm 7.86 \times 10^{-3}$
	Animal					$0.29 \times 10^{-3} \pm 1.51 \times 10^{-4}$

315   **SUPPLEMENTARY FIGURES**

316
317   **Suppl. Fig. 1. Genome-wide Manhattan plot for the different OTUs**. The horizontal line
318   represents the genome-wide Bonferroni -$\log_{10}$ (p) = 6.03 threshold.

Suppl. Fig. 2. Q-Q plots from genome-wide association analyses of the different OTUs.

Bibliography

1. Egerton, S., Culloty, S., Whooley, J., Stanton, C. & Ross, R. P. The gut microbiota of marine fish. *Front. Microbiol.* **9,** 873 (2018).

2. den Besten, G. *et al.* The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *J. Lipid Res.* **54,** 2325–2340 (2013).

3. Smith, C. C. R., Snowberg, L. K., Gregory Caporaso, J., Knight, R. & Bolnick, D. I. Dietary input of microbes and host genetic variation shape among-population differences in stickleback gut microbiota. *ISME J.* **9,** 2515–2526 (2015).

4. Kogut, M. H. & Arsenault, R. J. Editorial: gut health: the new paradigm in food animal production. *Front. Vet. Sci.* **3,** 71 (2016).

5. Limborg, M. T. *et al.* Applied hologenomics: feasibility and potential in aquaculture. *Trends Biotechnol.* **36,** 252–264 (2018).

6. Rudi, K. *et al.* Stable Core Gut Microbiota across the Freshwater-to-Saltwater Transition for Farmed Atlantic Salmon. *Appl. Environ. Microbiol.* **84,** (2018).

7. Edgar, R. C. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26,** 2460–2461 (2010).

8. Edgar, R. C. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat. Methods* **10,** 996–998 (2013).

9. Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res.* **41,** D590-6 (2013).

10. Dvergedal, H., Ødegård, J., Øverland, M., Mydland, L. T. & Klemetsdal, G. Selection for feed efficiency in Atlantic salmon using individual indicator traits based on stable isotope profiling. *Genet. Sel. Evol.* **51,** 13 (2019).

11. VanRaden, P. M. Efficient methods to compute genomic predictions. *J. Dairy Sci.* **91,** 4414–4423 (2008).

12. Yang, J., Zaitlen, N. A., Goddard, M. E., Visscher, P. M. & Price, A. L. Advantages and pitfalls in the application of mixed-model association methods. *Nat. Genet.* **46,** 100–106 (2014).

13. Gilmour, A. R., Gogel, B. J., Cullis, B. R., Welham, S. J. & Thomson, R. *ASReml user guide*. (VSN International LTD, 2015).

14. Holmström, C. & Kjelleberg, S. Marine Pseudoalteromonas species are associated with higher organisms and produce biologically active extracellular agents. *FEMS Microbiol. Ecol.* **30,** 285–293 (1999).

15. Wesseling, W. *et al.* Adverse Effects of Immobilised*Pseudoalteromonas* on the Fish
Pathogenic*Vibrio anguillarum:* An*In Vitro* Study. *J. Mar. Biol.* **2016,** 1–11 (2016).

16. VanInsberghe, D. *et al.* Non-symbiotic Bradyrhizobium ecotypes dominate North American
forest soils. *ISME J.* **9,** 2435–2441 (2015).

17. Miranda, C. D. & Zemelman, R. Bacterial resistance to oxytetracycline in Chilean salmon
farming. *Aquaculture* **212,** 31–47 (2002).

18. Koskinen, R. *et al.* Characterization of Sphingomonas isolates from Finnish and Swedish
drinking water distribution systems. *J. Appl. Microbiol.* **89,** 687–696 (2000).

19. Rudman, S. M. *et al.* Microbiome composition shapes rapid genomic adaptation of
Drosophila melanogaster. *Proc. Natl. Acad. Sci. USA* **116,** 20025–20032 (2019).

20. Wallace, R. J. *et al.* A heritable subset of the core rumen microbiome dictates dairy cow
productivity and emissions. *Sci. Adv.* **5,** eaav8391 (2019).

21. Goodrich, J. K., Davenport, E. R., Clark, A. G. & Ley, R. E. The Relationship Between the
Human Genome and Microbiome Comes into View. *Annu. Rev. Genet.* **51,** 413–433 (2017).

22. Sevellec, M., Laporte, M., Bernatchez, A., Derome, N. & Bernatchez, L. Evidence for host
effect on the intestinal microbiota of whitefish (*Coregonus* sp.) species pairs and their
hybrids. *Ecol. Evol.* (2019). doi:10.1002/ece3.5676

23. Giatsis, C., Sipkema, D., Smidt, H., Verreth, J. & Verdegem, M. The colonization dynamics
of the gut microbiota in tilapia larvae. *PLoS One* **9,** e103641 (2014).

378