1 **Intron retention as a new marker of the pre-disease state and its recovery to the**
2 **normal state by a traditional Japanese multi-herbal medicine**

3
4 Running title: Recovery of alternative splicing from the senescence to healthy type by
5 an herbal medicine
6
7

8 **Norihiro Okada[a#], Kenshiro Oshima[a$], Yuki Iwasaki[a$], Akiko Maruko[a], Erica**
9 **Iioka[a], Trieu-Duc Vu[a], Naoki Fujitsuka[b], Akinori Nishi[b], Aiko Sugiyama[b], Mitsue**
10 **Nishiyama[b], Atsushi Kaneko[b], Masahiro Yamamoto[b] and Susumu Nishimura[ac]**
11

[a] Foundation for Advancement of International Science, 3-24-16 Kasuga, Tsukuba,
Ibaraki 305-0821 Japan,
[b] Tsumura Kampo Research Laboratories, Tsumura & CO., 3586 Yoshiwara, Ami-
machi, Ibaraki 300-1192 Japan,
[c] Laboratory Animal Resource Center, University of Tsukuba, 1-1-1 Tennodai, Tsukuba,
Ibaraki 305-8575 Japan
18
19 [$] These two authors contributed equally to this work.
20 [#] To whom correspondence should be addressed.
21 nokada@fais.or.jp

**Abstract**

Intron retention (IR) is an important regulatory mechanism that affects gene expression

and protein functions. Using klotho mice as a model, we proposed that retained introns

are an excellent marker for the pre-disease state. Surprisingly, among widespread

retained introns that accumulated during aging in the liver, a subset was recovered to the

normal state by a Japanese traditional herbal medicine. IR-recovered genes fell into two

categories: (1) those involved in the spliceosome and (2) those involved in liver-specific

metabolism. By integrating data for splicing patterns, transcriptomes, and metabolomes,

we hypothesize that this medicine-related IR recovery under the pre-disease state

reflects the actual recovery of liver-specific function to the healthy state. Accordingly,

the study provides proof-of-concept evidence related to the ancient Chinese statement

proposing the medicine's usefulness for treating the pre-disease state. This approach

lays out a method for elucidating unknown molecular mechanisms of an herbal

medicine with multiple ingredients. (149 words)

**Introduction**

The vast majority of precursor mRNAs (pre-mRNAs) in mammalian cells consists of

exons separated by introns. Introns are normally removed by a mechanism called

splicing, which leaves the joined exons that form the mature mRNA (Padgett et al.,

1986, Patel and Steitz, 2003). The process of removing an intron from pre-mRNA is

highly dynamic, during which a complex rearrangement of protein-protein, RNA-RNA,

and RNA-protein interactions takes place in the spliceosome (Mayeda et al., 1994).

Because of such complexity, the process is highly vulnerable to external stimuli

(Stegeman and Weake, 2017). Dysregulation of RNA splicing has been identified as a

causative defect in several diseases, especially in cancer (Dvinge et al., 2016).

It has recently become widely recognized among molecular biologists that aging

reflects the gradual deterioration of the molecular components of the cell, the concerted

functioning of which are vital for cell viability and proliferation (Lopez-Otin et al.,

2013). In this sense, aging is regarded as a kind of disease, the extent of which varies

among individuals. The complexity of alternative splicing of pre-mRNAs makes this

process vulnerable even to aging, as suggested by many instances in which aging is

accompanied by a change in the expression level of spliceosomal proteins or by a

change in their splicing patterns (Zane et al., 2014). For example, levels of SF3B1 and

SRSF family proteins can be affected by changes in alternative splicing during aging

(Deschenes and Chabot, 2017). As these changes in turn may alter the alternative

splicing of numerous transcripts, a change in the expression of spliceosomal and

associated proteins is primarily important.

Intron retention (IR) is one of the forms of alternative splicing and is considered to

be harmful to the organism by (1) slowing down splicing kinetics to delay the onset of

gene expression (Braunschweig et al., 2014), (2) increasing pre-mRNA degradation in

the nucleus by nuclear exosomes (Niemela et al., 2014), and (3) increasing (pre-)mRNA

degradation in the cytoplasm by nonsense-mediated decay (Lejeune and Maquat, 2005).

IR accumulation has even been proposed as a post-transcriptional signature of aging

(Adusumalli et al., 2019). Analyses of the transcriptome from mouse, human brain, and

Drosophila head have shown a global increase in the level of IR during aging, indicating

that this process may be evolutionarily conserved (Adusumalli et al., 2019).

In the present study, we used the klotho mouse, which is a model for senescence and

exhibits a syndrome resembling human aging that includes a reduced life span,

decreased activity, infertility, osteoporosis, atherosclerosis, atrophy of the skin, and

emphysema (Kuro-o et al., 1997). This phenotype is caused by the disruption of a single

gene, klotho (*Kl*). In a normal mouse, the klotho protein and FGF23 function as a

receptor in the kidney to exert a diuretic action on phosphorus (Razzaque, 2009). In a

klotho mouse, however, the blood level of phosphorus is very high because of a

deficiency in the klotho protein, which leads to rapid senescence in this mouse.

Japanese multi-herbal medicines (Kampo) originated in ancient China and are

widely used in Japan for various disorders. In contrast to chemically synthesized

Western drugs, whose effects on humans are carefully evaluated during various stages

of testing and whose mechanisms of actions are generally rigorously evaluated, it has

been difficult to identify the molecular substances in the Kampo that are responsible for

their potency and to analyze the mechanisms by which they exert their effects. It has

long been said that thousands of chemical compounds in Kampo work together,

sometimes synergistically, and that their side-effects, if any, are relatively minor,

although their mechanism of action remains a total mystery (Zhou et al., 2016).

Juzen-taiho-to (JTT), being composed of 10 crude drugs as one of the formulations

approved as a medicine in Japan, was first described in a drug textbook in the Song

Dynasty (AD 1151) in China and was introduced in Japan during the Kamakura period

(Saiki, 2000) (AD 1185−1333). It has been traditionally administered to patients with

deficiency syndrome, consisting of disturbances and imbalances in the homeostatic

condition of the body (Onishi et al., 1998). More specifically, it is prescribed for

patients with fatigue, anemia, night sweats, anorexia, or circulatory problems. JTT has

stimulatory effects on the immune response including enhancement of phagocytosis,

cytokine induction, induction of mitogenic activity of spleen cells, and activation of

macrophage and natural killer cells (Matsumoto et al., 2000). JTT can also prolong life

when combined with the surgical excision of tumors and has a protective effect against

the deleterious effects of chemotherapy drugs (Wang et al., 2018).

Knowing that aging can lead to an increase in retained introns, we used klotho mice

that had not yet undergone extensive aging and examined the effects of JTT on IR

patterns in genes expressed in their organs by using an RNA sequencing (RNA-seq)

analysis (Vu et al., 2019). We discovered that IR occurred even during this earlier stage

in the aging process when no change in protein level and no pathological changes in the

organs were observed and can thus be used as a new marker of the pre-disease state.

Detection of the pre-disease state is important for human health, because the state can

be reversible when it is appropriately treated. Here we provide definitive evidence that

JTT can specifically decrease the retained introns of genes for liver metabolic functions

among those that show an increase during the process of aging. With the support of both

transcriptome data related to differentially expressed genes (DEGs) and changes in the

metabolome, we suggest that JTT has a beneficial effect on metabolism in the liver at an

earlier stage of aging. We thus propose that the decrease in liver IR at this earlier stage

of aging by JTT is an indication of the effective actions of this herbal medicine at the

time corresponding to the pre-disease stage.

**Results**

**Looking for a marker of the pre-disease state in klotho mice at 7 weeks of age**

We used klotho mice, a model mouse for aging, for our experiment to look for a new

marker for the pre-disease state as well as a possible effect of a Japanese herbal

medicine on this state. Because aging is currently regarded as a kind of disease (see

Introduction), the pre-aging state can be regarded as a pre-disease state. We have chosen

the time of 7 weeks after birth for the date of sampling (Fig. 1AB), which is less than

half of the half-life of these mice (14.7 weeks; Fig. 1B). We expected that markers, if

any, isolated from these mice at 7 weeks would represent the pre-disease state. We also

designed our experiments to examine the effect of JTT on the improvement of the pre-

disease state (Fig. 1A). We designated klotho mice as KL and wild-type mice as WT,

each of which was fed with (+) or without (−) 0.5% JTT from 3.5 weeks of age until the
mice were 7 weeks old.
First, to examine the extent of aging of liver tissue in klotho mice, we looked for
histological changes at 7 weeks of age relative to liver tissue in WT mice. Fig. 1C
shows that there was no such alteration, indicating that the liver did not exhibit a
senescent phenotype at this age. Second, we chose several proteins and examined their
protein level by performing western blotting analysis (Fig. 1D). The representative
proteins are DDX5, SRSF6, and NXF1, all of which are splicing-related genes, and
PPARD and SIRT7, both of which are involved in metabolic pathways of the liver.
NDUFS2 is the core subunit of the mitochondrial Complex1, and LXRβ is encoded by
*Nr1h2* and regulates macrophage function. GAPDH was used as the loading control.
Furthermore, we checked the expression of RGN/SMP30, a marker of senescence
whose expression decreases with aging (Maruyama et al., 2010) (Fig. 1E). There was no
observable difference in the expression of these proteins between KL− and WT− (Fig.
1DE), and thus we can conclude that the klotho mouse liver at 7 weeks of age does not
show clear signs of aging.
159
160    **Metabolome analysis shows that klotho mice at 7 weeks of age express starvation-**
161    **like characteristics**
162
To look for possible internal changes in the liver that could describe the pre-disease
state in this earlier stage of aging, we investigated 110 metabolites of the liver using
CE-TOFMS (CE: capillary electrophoresis, TOFMS: time-of-flight mass spectrometry),
and compared metabolite levels between KL− and WT−, the comparison of which is a
measure of senescence. Among 12 metabolites that showed such differences, the data
for 7 metabolites ($P < 0.05$) are shown in Fig. 2. Remarkably, 3-hydroxybutyric acid (3-
HBA) was significantly increased in KL− (Fig. 2A), to a level 2.5 times higher than that
in WT−. 3-HBA is synthesized in the liver from fatty acid as one of the ketone bodies
under conditions of starvation and can be used as an energy source when blood glucose
is low (Fukao et al., 2014). Consistent with this, certain intermediate metabolites of
glycolysis, namely glucose 1-phosphate, fructose 1,6-diphosphate, dihydroxyacetone
phosphate, lactic acid, CoA divalent, and alanine in KL− were significantly decreased as
compared with their levels in WT− (Fig. 2B). These data suggest that klotho mice are
undergoing a fasting state, in spite of the presence of ample food, during which the
synthesis of 3-HBA is activated. Accordingly, an increase in 3-HBA and a reduction in
glycolysis in KL− might be representative of senescence in klotho mice.

**DEG analysis shows the alteration of gene expression during aging**

Knowing that the liver of klotho mice is under the starvation-like condition, we performed RNA-seq analysis to look for a possible alteration of the gene expression level in the liver during the pre-disease state. From liver collected from KL+, KL−, WT+ and WT−, we isolated the mRNAs and then sequenced them. Multi-dimensional scaling confirmed the consistency of each dataset (Fig. S1A). In addition, transcriptomes from individual organs from KL+ and KL− indicated that JTT induced extensive transcriptome changes (Fig. S1B).

We then extracted a down-regulated gene set and an up-regulated gene set in KL− from the comparison of KL− and WT− (1228 genes and 1352 genes, respectively; Fig. 3A). The former represents genes whose expression decreased in the liver probably due to aging-related deterioration, and the latter represents genes whose expression increased possibly reflecting an adaptive response to aging. To characterize enriched cell types in these genes, we performed a cell type enrichment (Cten) analysis (Shoemaker et al., 2012) by using the software provided by these authors, in which, for each cell type, a transcript was defined as a highly expressed cell-specific gene in that cell if it had an expression value at least 15× greater than its median expression value across all cells examined. The software can accurately identify the appropriate cell type in a highly heterogeneous environment and provides insight into the suggested level of enrichment to minimize the number of false discoveries. Fig. 3C shows that liver-specific genes were especially enriched among the genes whose expression decreased during aging, suggesting that liver function was reduced during aging. In contrast, a variety of non-liver-specific cell types were enriched among those genes whose expression increased during aging (Fig. 3D). It should be noted that several macrophage-related cells were enriched in these increased genes, consistent with the generally believed notion that the aging process activates innate immunity in spite of its severe deterioration of adaptive immunity (Solana et al., 2006).

**DEG analysis shows the improvement in liver function as a result of JTT administration**

To examine the influence of administration of JTT on the gene expression level in the livers of klotho mice, we extracted the set of genes that were up-regulated and down-regulated in KL+ after administration of JTT from the comparison of KL+ and KL−

(532 genes and 412 genes, respectively; Fig. 3B). Cten analysis of the up-regulated

genes demonstrated that liver-specific genes were highly enriched (Fig. 3E), whereas

that of down-regulated did not exhibit any enrich https://orcid.org/0000-0002-3655-

8060

ent. A comparison of the up-regulated gene set from the comparison of KL+ and KL–

(532 genes) and the down-regulated gene set in KL– from the comparison of KL– and

WT– (1228 genes) resulted in 213 DEGs, each of which exhibits the clear recovery

pattern from the decreased level of KL– to the level of WT– in the KL+ (Fig. 3F). Fig.

3G shows a boxplot summarizing the expression of 213 genes. Each data point was

normalized to the average expression level of each gene, and all the data were summed.

By using these genes, we performed a Cten analysis as shown in Fig. 3H and showed

that liver-specific functions are highly enriched in the 213 genes. We then extracted GO

terms enriched for these genes. Fig. 3I shows that genes related to liver-specific

functions, including metabolic process, flavonoid biosynthetic process, flavonoid

glucuronidation, and lipid metabolic process, were enriched, suggesting that the liver

functions were recovered by administration of JTT.

**IR and its recovery after administration of JTT is most prominent in the liver**

Knowing that klotho mice at 7 weeks of age represent a starvation-like state and that

their liver-specific functions at that time might be reduced due to the decreased

expression of liver-specific genes, we looked for possible changes in alternative splicing

at this age because it is known that several stresses can change alternative splicing

patterns (Boutz et al., 2015, Ninomiya et al., 2011, Shalgi et al., 2014; see later

discussion). We performed RNA-seq analysis in additional three organs, namely, blood,

kidney and heart for comparison.

To identify what type of alternative splicing is most prominent in the four organs

examined here, we first analyzed five types of alternative splicing, namely alternative 5'

splice site, alternative 3' splice site, skipped exon (SE), IR, and mutually exclusive

exons, between KL+ and KL– by using rMATS (Table 1-1). SE and IR were the most

prominent types of alternative splicing among the four organs. Noticeably, a change in

the frequency of IR in the liver was the most prominent effect followed by that in the

blood (Table 1-1).

We then calculated the difference in the frequency of IR in comparisons among

KL+, KL–, WT+, and WT– with the criteria of $P < 0.01$ (Table 1-2). We first noticed a

difference in the frequency of retained introns for the comparison between KL+ and

WT– was noted for only 59 loci, whereas that for KL– and WT– was noted for 212 loci.
The comparison between KL+ and WT– is a measure of the effect of JTT over
senescence, whereas the comparison between KL– and WT– measures the effect of the
senescence level in terms of the extent of retained introns accumulated during aging in
klotho mice. The above data suggest that the administration of JTT shifted the AS
pattern in klotho mice to be more similar to that of WT mice.
The 142 IR loci (Table 1-1 & -2) that were differentially affected between KL+ and
KL– in the liver included those genes for which the frequency of IR was decreased with
JTT in KL+ (132 loci; designated as "DecIR" type) and those genes for which the
frequency of IR was increased with JTT in KL+ (10 loci; designated as "IncIR" type)
(Fig. 4A). Likewise, the comparison between KL+ and KL– for the other three organs
showed a number of differential IR loci that could be divided into these two types. The
liver and blood exhibited higher values in the DecIR type as compared with the IncIR
type, whereas the other two organs, kidney and heart, had comparable values for DecIR
and IncIR (Fig. 4A). From these data, we designated liver and blood as two target
organs for JTT. Since we showed that the liver-specific function was recovered by
administration of JTT (see above), we focused in particular on the analyses of genes that
underwent a decrease in IR in the liver in response to JTT.
269
270 **Analysis of IR-recovered genes in the liver**
271
To demonstrate our strategy for evaluating the extent of IR, we first show an example
based on the data for *Anxa11* expression in the liver (Fig. 4C). *Anxa11* plays an
important role in cell division, $Ca^{2+}$ signaling, vesicle trafficking, and apoptosis (Han et
al., 2017). Sequenced *Anxa11* transcripts showed extensive accumulation of a retained
intron in KL–, whereas this intron was retained to a lesser extent for KL+ and WT–
(Fig. 4Ci).
Then, to analyze IR of *Anxa11* mathematically, we calculated the read counts of
skipping junctions (SJCs) and those of inclusion junctions (IJCs) for KL+, KL–, and
WT– (Fig. 4Cii) according to the strategy illustrated by rMATS (Fig. 4B). For KL+, we
then took the base-2 logarithm of the SJC/IJC ratio, which gives the fold change
(hereafter designated as FC) between the SJC and IJC at the intron junction (the thick
black bars in Fig. 4Ci). The same process was carried out for KL– and WT–, and the FC
values for KL+, KL–, and WT– were compared. The lower FC value for KL– was
recovered in KL+ to the same level as that of WT– (Fig. 4Ciii). The V shape of this bar

plot indicated a recovery from the senescence-type IR to the healthy-type IR after JTT

administration (see below).

As was described above, we first obtained the DecIR type (132 loci, 124 genes) in

klotho mice caused by JTT administration by comparing KL+ vs KL–. This simply

represents the recovery of retained introns in KL+. Then, we obtained the IncIR type

(190 loci, 180 genes) in klotho mice by comparing KL– vs WT–. This represents a

measure of the senescence-related changes in retained introns. Accordingly, the

comparison between these two types allowed us to highlight the effect of JTT at 7

weeks of age as shown in a Venn diagram (Fig. 5A). The overlapping 70 loci (67 genes)

represent the "complete recovery" loci, which are defined as those with significant

differences between KL+ and KL– and between KL– and WT–, but with FC values for

KL+ and WT– that were similar. The non-overlapping 62 (61 genes) and 120 loci (115

genes) represent "partial recovery" and "no recovery" loci, respectively. Fig. 5B shows a

heatmap of the 70 "complete recovery" loci, each of which represents a clear recovery

pattern. Fig. 5C is a boxplot of the FC values of KL+, KL–, and WT– for these 70 loci,

which shows the typical V-shaped pattern of recovery.  Fig. 5D & E show boxplots of

the "partial recovery" loci and the "no recovery" loci, respectively. The former

represents loci with a significant difference between KL+ and KL– and without a

significant difference between KL– and WT–. The latter represents loci with a

significant difference between KL– and WT– and without a significant difference

between KL+ and KL–.

**Characteristics of loci with retained introns in the liver**

We next characterized introns among the 70 "complete recovery" loci, the 120 "no

recovery" loci and all other genes (~250,000 loci) expressed in the liver. The average

length of introns from the "complete recovery" and "no recovery" loci was significantly

shorter than that of all other introns (Fig. 6A). There was, however, no significant

difference in intron length between the "complete recovery" and "no recovery" loci,

suggesting that the "complete recovery" loci were not selectively discriminated from the

"no recovery" loci during the recovery process associated with JTT administration. The

GC content of introns associated with the "complete recovery" and "no recovery"

groups was significantly higher than that of all other liver-related introns (Fig. 6B).

Also, there was no significant difference between "complete recovery" and "no

recovery" introns in this respect, suggesting that GC content was not linked to recovery

status among IR loci. Then, we examined the strength of 5' and 3' splice sites by

calculating scores for these splice sites using the software MaxEntScan. The 5' splice

sites of introns of the "complete recovery" and "no recovery" loci were slightly but

significantly weaker than those of all other liver-related introns (Fig. 6C), but we found

no such differences at the 3' splice sites (Fig. 6D). These characteristics of introns at

IR loci were observed in the three additional organs (see below and Fig. S2A−D)

and are consistent with previous data (Braunschweig et al., 2014). Thus, loci with

shorter intron lengths, a higher GC content, and a weaker splicing score at the 5' site

have a tendency to have retained introns, but these are not characteristics that make

retained introns more likely to undergo recovery in the presence of JTT in the klotho

mice. We also confirmed the susceptibility of introns to IR in genes in terms of their

shorter intron length, higher GC content, and weaker intron score at the 5' splice site in

the other three organs (Fig. S2A−D) as was shown in the case of liver (Fig. 6).

Next, we searched for possible enrichment of binding sites for RNA-binding

proteins in the introns, as these are known to influence splice site selection and are

required for splicing (Padgett et al., 1986,Patel and Steitz, 2003). Binding sites for five

RNA-binding proteins were especially enriched in introns among the 70 "complete

recovery" loci but not among the 120 "no recovery" loci (Fig. 6E). This suggests that

genes expressed in the liver were more likely to recover from IR after JTT treatment if

they had these RNA-binding sites. Likewise, we examined the tendency for enrichment

for transcription factors (TFs) that bind to promoter regions in liver and in blood.

Binding sites for four TFs (Hoxa10, Hoxd12, Hoxd10, and Hoxd13) were especially

enriched in "complete recovery" genes in comparison with "no recovery genes" in liver

(Fig. 6F).

**Validation of IR-recovered genes in the liver by using RT-PCR**

Eleven genes were chosen for Integrative Genomics Viewer (IGV) mapping and RT-

PCR, which confirmed the recovery of IR upon JTT administration (Fig. 7, Fig. S3).

These genes are involved in RNA splicing (*Ddx5, Srsf5*) and typical liver functions such

as glucose and fatty acid metabolic pathways (*Sirt7, Acadm, Decr2, Gnmt*), lipid

metabolic processes (*Acadm*), heme biosynthesis and bile acid production (*Cyp27a1,*

*Hsd3b7*), and oxidation-reduction processes (*Acadm, Cyp27a1, Hsd3b7*).

**Biological function of IR-recovered genes in the liver**

Possible effects of JTT on the selection of liver retained intron loci for recovery during

aging were analyzed (Fig. 8). The results obtained from the Gene Ontology (GO)

analysis using Biological Process terms in Fig. 8A & B suggest two conclusions. First,

in the comparison between DecIR type (132 loci, 124 genes) of KL+ vs KL– and IncIR

type (190 loci, 180 genes) of KL– vs WT–, genes involved in RNA splicing and mRNA

processing are susceptible to IR during the aging process, and these retained introns

tend to be recovered by JTT (i.e., the red and blue bars are comparable; Fig. 8A). The

tendency of genes involved in these processes to be susceptible to IR during aging is

consistent with a previous study (Zane et al., 2014). These splicing-related genes can

control the expression of downstream genes through the regulation of their alternative

splicing. We defined 250 genes as potential splicing-related genes to highlight their

importance in the regulation of alternative splicing (see Methods for the definition of

splicing-related genes). They included almost all the spliceosomal proteins and

spliceosome regulatory proteins. Among 368 gene loci that exhibited a difference in

alternative splicing for the comparison of KL+ and KL– in the liver (Table 1-2), 19

genes (data not shown) were detected among the list of 250 potential splicing-related

genes. Among these 19 genes, 9 potential splicing-related genes in the liver were

subjected to IR, all of which exhibited the recovery pattern in terms of the comparison

in the FC among KL+, KL–, and WT– (Table 2). The patterns of recovery of the seven

potential splicing-related genes in liver associated with JTT administration together with

their FC values are shown in Fig. 7, Fig. 8D(i–iii), and Fig. S5B. Accordingly, JTT

appears to work in the direction of recovery, especially for potential splicing-related

genes.

Second, in addition to the recovery of IR for genes related to RNA splicing and

RNA processing, IR recovery was not evenly distributed among genes that accumulated

retained introns during aging but instead occurred selectively among liver-specific

genes. This conclusion was more obvious when loci of "complete recovery" (70 loci)

and "no recovery" (120 loci) were compared (Fig. 8B). The GO terms metabolic

process, oxidation-reduction process, carbohydrate metabolic process, and regulation of

gluconeogenesis, which represent major functions of the liver, were enriched among

genes that underwent "complete recovery" (i.e., red bars), whereas genes in the "no

recovery" group are associated with more general functions such as translation, protein

transport, and apoptosis (i.e., blue bars). These data suggest that, among genes

susceptible to IR during aging, the liver-specific genes selectively underwent a shift in

retained introns from the senescent state to the normal state after administration of JTT.

To further confirm that the recovery of retained introns by administration of JTT
occurred in liver-specific genes, Cten analysis was performed as shown in Fig. 8C.
Genes with "complete recovery" were significantly enriched among liver-specific genes
($-\log_{10} P = 6.040$), confirming that the IR recovery from aging specifically occurred in
liver-specific genes.

**Biological function of IR-recovered genes in the blood**

To examine the characteristics of recovered genes among the other three organs, GO
analysis was performed for these organs, but only in the case of blood was the
significant enrichment of pathways with $P < 0.05$ observed (data not shown). GO
analysis for genes expressed in the blood that recovered from IR after JTT
administration indicated that the terms for immune system process and adaptive
immune response (Fig. S4B) were significantly enriched.

**Overlapping IR-recovered genes between liver and blood**

In addition to organ-specific genes with recovered IR, 11 genes that underwent IR
recovery were expressed in both the liver and blood (Fig. 8E, Fig. S5A).  Fig. S5B
shows the patterns of IR recovery of genes that occurred in both of these organs.
Independent occurrence of recovery in different organs strongly suggests the
importance of the recovery of these genes for the organism. Among these 11 genes,
four splicing-related genes (*Sf3b1*, *Ddx5*, *Srsf5*, and *Srsf6*) were included, again
highlighting the importance of the recovery of these genes. Among the remaining
seven genes, we paid attention in particular to the following five genes, namely *Nr1h2*
(LXRβ), *Faah*, *Gdi1*, *Idh3g*, and *Sh3glb2* (EB2). *Nr1h2* is a key regulator gene in
macrophage function (Zelcer and Tontonoz, 2006). *Faah* is connected with
macrophage function through lipid metabolism (Xu et al., 2017), and *Gdi1* is
connected with interferon-γ activity through innate immunity (Ohshima et al.,
2015). *Idh3g* and *Sh3glb2* (EB2) are involved in mitochondrial function (Fino et al.,
2017). Considering that one of the prominent functions of JTT has been the activation
of the innate immune system, especially macrophage function, the recovery of IR
of *Nr1h2* (LXRβ), *Faah*, and *Gdi1* is especially interesting.

**Possible involvement of TFs in the increase in retained introns during aging and**
**their decrease by administration of JTT**

We first examined enrichment for TF-binding sites in the promoter region spanning

500 bp upstream from the transcription start site in the IR-increased genes in the liver

(180 genes, 190 loci) and in the blood (12 genes, 12 loci), categorized as IncIR.KL− vs

WT− (Fig. 9A). This type of data shows which TFs were possibly involved in the

generation of retained introns that increased during progression of the pre-disease

process. Most TFs were organ specific, as only two TFs, SPDEF and Sp1, were shared

among genes in both organs, suggesting that they play a more general role in aging. We

then examined enrichment for the binding sites of TFs in the IR-recovered genes in the

liver (124 genes, 132 loci) and in the blood (39 genes, 40 loci), categorized as

DecIR.KL+ vs KL− (Fig. 9B). These data clearly showed that different sets of TFs were

used for transcription in IR-recovered genes in liver and in blood. Fig. 9C shows data

from Fig. 9A & B for the liver, showing that some TFs were likely involved in both the

increase and decrease of retained introns, whereas others were uniquely used in one case

or the other.

This observation prompted us to examine how many IR-increased genes or IR-

recovered genes in the respective organs have a common TF-binding site in the

promoter region spanning 500 bp upstream from the transcription start site. In the case

of an increase in IR during aging in the liver, 171 of 180 genes (190 loci) had at least

one of four TFs, namely KLF14, SCL, THRb, and Zac1, as shown in Fig. 9D(i)(ii),

which represents 95% coverage. In the case of a decrease in retained introns by

administration of JTT, 120 of 124 genes (132 loci) had at least one of four TFs, namely

SP2, Tgif2, KLF14, and Hoxd12, as shown in Fig. 9E(i)(ii), which represents 96.8%

coverage. These data clearly show that a distinct set of genes sharing the same TF-

binding sites was chosen during the IR-increasing process of aging and during the IR-

decreasing process after administration of JTT and were networked. The significance of

this observation will be discussed below.

A similar analysis was performed for the blood data, the results of which are shown

in Fig. S6.

**Discussion**

**IR is a marker of the pre-disease state**

It has been proposed that retained intron accumulation is a signature for aging
(Adusumalli et al., 2019, Huan Li, 2017, Stegeman and Weake, 2017). In the present
study, we clearly demonstrated that IR occurs during the pre-disease state when neither
pathological alterations nor a difference in protein level could be observed. We
speculated that some disruption of homeostasis induced by a change in metabolites
triggered IR as a result of a stress during this earlier stage of aging. As shown in Fig.
2A, a significant increase in 3-HBA and decreases in intermediate metabolites of
glycolysis in KL– (Fig. 2B) suggest that the liver at this stage is subjected to starvation-
like conditions, providing the possibility that this starvation-like state is a stress signal
that triggers intron retention. Also, the decrease in 3-HBA in KL+, even if not
statistically significant, suggests at least a partial recovery from a starvation-like
condition. In addition, metabolome analysis also showed that the concentrations of
several nucleotides that are precursors of RNA were lower in KL– (Fig. 2C) relative to
that in WT–, suggesting that this also triggered intron retention. It is interesting to note
that the concentrations of some of these precursors recovered in KL+ to WT levels.
Accordingly, it is possible that the recovery of these precursors after the administration
of JTT affected the 180 IR-increased genes to reduce some of these retained introns to
the normal state, although this does not necessarily explain the reason why retained
introns of only a specific subset of genes were recovered to the normal state by JTT (see
the later discussion below).
483
484  **IR-recovered genes can be classified according to their functions**
485
Fig. 10A illustrates two major categories among the 132 IR-recovered genes, namely
alternative splicing and lipid/glucose metabolism. Splicing-related genes such as *Ddx5*,
*Sf3b1*, *Srsf5*, and *Srsf6* were included in these IR-recovered genes (Table 2). Splicing-
related genes are vulnerable to disruption of homeostasis in many systems, including
cancer (Dvinge and Bradley, 2015, Zhang and Manley, 2013), aging (Deschenes and
Chabot, 2017), and several stresses (Monteuuis et al., 2019, Shalgi et al., 2014). It is
possible that the IR recovery of some of these splicing-related genes reflects the
recovery of the normal splicing ability in KL+ after administration of JTT (see the later
discussion below).
In addition to genes involved in splicing, retained introns of genes involved in
lipid/glucose metabolism such as *Sirt7*, *Decr2*, *Acadm*, *Adipor1*, *Ppard*, *Nr1h2* (*LXRβ*),
and *Faah* were recovered (Fig. 10A). They are relevant to hepatic metabolic pathways
involving glucose and fatty acids. Under fasting or starvation conditions, endogenous

glucose is mainly produced by a breakdown of glycogen and gluconeogenesis in the
liver and is metabolized into pyruvate, which is used to generate ATP through the
tricarboxylic acid cycle and oxidative phosphorylation in mitochondria (Nishi et al.,
2017). In addition, fatty acids are oxidized through hepatic mitochondrial β-oxidation to
generate ketone bodies. Thus, liver-produced glucose and ketone bodies provide
essential sources of energy for extrahepatic tissues during fasting and starvation (Rui,
2014). Metabolome analysis showed that the condition of the liver in KL– resembled
starvation and that the condition of the liver in KL+ was improved, suggesting that the
improvement in the condition of KL+ triggered a cue for the recovery of IR of these
genes (see the model proposed below).
In addition to genes belonging to these two categories, a few genes involving
mitochondrial processes and immune system and macrophage processes are also
included in these IR-recovered genes. These are listed and referenced in the legend of
Fig. 10A, in which the possibility is discussed that the IR recovery of several nuclear
receptor genes such as *Nr1h2* (LXRβ) might reflect the effect of JTT on macrophage
function.
515
516   **Some liver metabolic functions were improved to the normal state by**
517   **administration of JTT**
518
Before discussing a model concerning the mechanism by which only certain genes
undergo IR recovery in response to JTT, we highlight the results of 213 genes from the
DEG analysis, the expression levels of which were down-regulated in KL– and were
recovered to the level of WT– in KL+ (Fig.3G). GO analysis of these genes showed that
metabolic processes and glucuronidation in the liver likely were up-regulated in KL+
and had recovered after administration of JTT from the down-regulated state in KL–
(Fig. 3I).
There are many clinical and basic scientific reports on the improvement of liver
functions by JTT, typically exemplified by reports that JTT can protect the liver from
injury by chemicals administered to patients with cancer or other diseases. A few
examples are listed below (others are listed in the legend for Fig. 10A). JTT protects
against isoniazid/rifampicin-induced hepatic injury by modulating oxidative stress and
inflammatory responses (Yoshioka et al., 2019). JTT exerts protective effects against
alcohol-induced liver disease by modulating oxidative stress and the inflammatory
response (Fukaya et al., 2018). JTT has the potential to protect against bromobenzene-
induced hepatotoxicity and to modulate oxidative stress (Yoshioka et al., 2016). The

presence of such a variety of reports convinced us that an improvement in liver

function, especially metabolic activities, is one of the major effects of JTT. These

published data are consistent with our DEG data described above.

**A model to explain the reason why IR in a specific subset of genes involving liver**

**metabolism was recovered**

As was mentioned in the Introduction, IR has been considered to be harmful to the

organism (Weischenfeldt et al., 2005). Recent global screens across many cells and

tissue types from humans and mice, however, have gradually revealed the role of IR as

a negative regulator of gene expression that is integrated into a regulatory network of

RNA processing and has functional significance (Braunschweig et al., 2014, Jacob and

Smith, 2017, Liu et al., 2017). Recent studies have shown that programmed IR is a

critical regulatory pathway in normal development and differentiation, such as in

granulocyte differentiation (Wong et al., 2013), terminal erythroid differentiation

(Pimentel et al., 2016), male germ cell differentiation (Naro et al., 2017) and B cell

development (Ullrich & Guigo, 2019). The fact that IR within the cyclin A2 gene

induces G1/S arrest in differentiated human tissues suggests that IR may play a role in

cell differentiation and senescence through regulation of the cell cycle (Honda et al.,

2012).

During the heat shock response in mice, there is widespread retention of introns in

transcripts from 1,700 genes, which are enriched for tRNA synthetase, nuclear pore, and

spliceosome functions. These transcripts with retained introns are, for the most part,

nuclear and untranslated. These splicing-inhibited transcripts induced by heat shock

stress are mostly spliced post-transcriptionally, in contrast to other normal splicing that

occurs co-transcriptionally (Shalgi et al., 2014). These nuclear retained transcripts are

presumed to await a signal for splicing that is cued by the release from the stress (Jacob

and Smith, 2017). The presence of other similar examples (Boutz et al., 2015, Ninomiya

et al., 2011) of stress-induced regulation of intron retention suggests that this regulatory

mechanism contributes to a wide range of gene expression.

As aging is also a kind of stress, we considered intron retention during aging as

analogous to that which occurs during the heat shock response as described above. Fig.

10C(i) shows a model for the case of KL− in that transcripts with IR are retained in the

nucleus upon the stress signal of aging such that global metabolic activity in liver cells

is repressed to save energy. After administration of JTT, as liver-specific metabolic

functions are recovered to some extent as described above, improvement in metabolic

activity results in a signal to the liver cells to recover these retained introns by sending a
cue for post-transcriptional splicing (Fig. 10C(ii)). If this model is correct, we can
determine which pathways in the cells are improved by knowing which functions the
IR-recovered genes are involved in. In the present case, the recovery of retained introns
of genes involved in lipid/glucose metabolic pathways (Fig. 10A) suggests that the
liver-specific metabolic functions were recovered. Genes detected by DEG data (Fig. 3F
G), the Cten analysis (Fig. 3H), and its GO analysis (Fig. 3I) confirmed that this is the
case. Also, the recovery of retained introns of genes involved in splicing pathways (Fig.
8A) suggests that splicing functions were recovered, at least in part. In addition, since a
few genes involving mitochondrial processes and immune system and macrophage
processes are also included in the IR-recovered genes, we speculate that these functions
were also recovered by administration of JTT.
583
584 **The mechanism for aging-related intron retention in transcripts and possible gene**
585 **specificity of JTT-related recovery**
586
Regarding the mechanisms related to how transcripts retained their introns and to
how a specific group of transcripts was recovered, we suggested above that IR-
susceptible genes during aging and IR-recovered genes after JTT administration are
transcriptionally controlled by a very few TFs that are specific to the liver and blood
(Fig. 9). There is emerging evidence that mRNA splicing is controlled by multi-layered
mechanisms that involve transcription and epigenetics (Braunschweig et al., 2014, Lev
Maor et al., 2015, Luco et al., 2011). Therefore, it is possible that TFs, chromatin
constituents, and epigenetic factors such as those related to histone modification and
DNA methylation not only control transcription but also regulate splicing (Lev Maor et
al., 2015, Luco et al., 2011). These factors can affect the rate of RNA polymerase II
elongation, which in turn affects the pattern of alternative splicing. Accordingly, it is
possible that accumulation of retained introns during aging in klotho mice was caused
by a change in the elongation rate of transcription. As shown in Fig. 9, there are several
TF-binding sites that are enriched in the IR-susceptible genes during aging (190 loci,
180 genes). Accordingly, an aging signal such as fasting may lead specific TFs to retard
the transcription rate, resulting in IR. Also, it is possible that some active ingredients of
JTT in cooperation with these TFs could contribute to the recovery of IR.
Also, it is interesting to note that the concentration of certain nucleotide diphosphates
(UDP and GDP) was decreased in KL− and recovered to WT levels in KL+ (Fig. 3C). If
a low concentration of nucleotides triggers a change in the elongation rate of

transcription (Kwak and Lis, 2013), it is also possible that this may have induced the
accumulation of retained introns. The recovery of such concentrations to the normal
level in KL+ is consistent with the recovery of retained introns in KL+.

Although there are instances in which chemical compounds influence AS by directly
interacting with splicing factors (Kataoka, 2017), there has been no report of changes in
AS patterns that result from an interaction with a TF. It is of interest to examine whether
active ingredients of a Kampo medicine can interact with TFs to influence AS patterns
co-transcriptionally or post-transcriptionally. The current results regarding the reduction
of retained introns by JTT will provide a rich resource for the subsequent studies of the
regulation of AS.

**Perspective**

This study is the first to shed light on the comprehensive mechanism of a multi-herbal
medicine from the viewpoint of systems biology. It is interesting to note that the ancient
Chinese medical textbook *Inner Canon of the Yellow Emperor* stated that the pre-
disease state should be treated early with Kampo medicine (UNESCO, 2011). The
present data represent a molecular proof of this historical statement. Kampo medicine is
a precious heritage in Japan, and there are similar medicines used around the world such
as traditional Chinese medicines (Patwardhan et al., 2005), traditional medicines in
India (Ayurveda) (Patwardhan et al., 2005), and ancient Greek medicines (Unani
medicine). To elucidate the mechanisms of these medicines, which are the result of
human wisdom that has accumulated over several thousand years, and to pass on this
knowledge to subsequent generations are the main goals of our research.

**Materials and Methods**

**Juzen-taiho-to (JTT)**

JTT was supplied by Tsumura & Co. (Tokyo, Japan) in the form of a powdered
extract. It was obtained by spray-drying a hot water extract mixture of the following 10
crude drugs in the ratios provided in parentheses: Astragali radix (10.52), Cinnamomi
cortex (10.52), Rehmanniae radix (10.52), Paeoniae radix (10.52), Cnidii rhizome
(10.52), Atractylodis lanceae rhizome (10.52), Angelicae radix (10.52), Ginseng radix
(10.52), Poria (10.52), and Glycyrrhizae radix (5.32). The origins and species of each
component, the contents of characteristic ingredients, and other pharmaceutical-grade

qualities of JTT are strictly controlled as it is an ethical drug approved by the Ministry
of Health, Welfare and Labor of Japan.
645

**RNA extraction and RNA-seq**

After removal of organs, they were subjected to RNA extraction after being soaked
with RNALater (Thermo Fisher Scientific, Rockford, IL, USA). RNA extraction was
performed on individual tissue samples with the Pure Link RNA Mini kit (Invitrogen,
MA, USA). Briefly, 600 μL of Lysis Buffer and 900 μL of TRIzol (Thermo Fisher
Scientific) were added to 0.03 g of tissue, and the tissue was homogenized. After the
sample was incubated for 10 min at room temperature and centrifuged at $12,000 \times g$ for
15 min, the supernatant was treated with DNase and purified by column cartridge. The
quality of RNA was checked by Qubit (Thermo Fisher Scientific) and TapeStation
(Agilent Technologies, CA, USA). RNA library construction was performed by using a
TruSeq Stranded mRNA Sample Prep kit (Illumina, CA, USA). Paired-end (150 base
pairs × 2) sequencing with the NovaSeq 6000 platform (Illumina) was outsourced to
Takara Bio, Shiga, Japan.

**Quality check and filtering of RNA-seq data and mapping analysis**

For purification of the sequencing data, cutadapt v.1.16 (Deschenes and Chabot, 2017)
was used to remove Illumina adapter sequences, followed by removal of the poly(A)
sequence using fastx_clipper software in the fastx toolkit software package v.0.0.14
(http://hannonlab.cshl.edu/fastx_toolkit/). To remove low-quality bases or sequences,
we trimmed the sequences using fastq_quality_trimmer software (parameters: -t 20 -l 30
-Q 33) and fastq_quality_filter software (parameters: -q 20 -p 80 -Q 33), both of which
are included in the fastx toolkit. During the above processing, any read in which one of
the pairs was missing was removed using Trimmomatic v.0.38 (Bolger et al., 2014).
Then, reads containing mouse rRNA, tRNA, or phiX sequences as the control sequence
from Illumina were removed using Bowtie 2 v. 2.3.4.1 (Langmead and Salzberg, 2012).
We then carried out the second round of removal of any unpaired reads using
bam2fastq. After completion of these filtering steps, 20 million reads of each of the
forward and reverse sequences per sample were mapped to the mouse genome sequence
build GRCm38 using Tophat v2.1.1 (Trapnell et al., 2009). The mouse genome
sequence was downloaded from iGenomes of Illumina
(http://jp.support.illumina.com/sequencing/sequencing_software/igenome.html).

Multiple mapped reads were removed using samtools (parameters: samtools view -q 4).

Uniquely mapped reads were counted by gene annotation (Ensembl release 81) using

featureCounts v.1.6.2. The counted values were normalized with the TMM method

using EdgeR library in R v.3.5.0 and used for expression analysis.

**Analysis of alternative splicing**

Loci with significantly different splicing patterns with $P < 0.01$ were detected within

six combinations: KL+/KL–, KL+/WT+, KL+/WT–, KL–/WT+, KL–/WT–, and

WT+/WT–, where '+' and '–' refer to treatment with and without, respectively, JTT,

using rMATS v.4.0.2 (Shen et al., 2014). Statistical significance was tested using the

skipping junction counts (SJCs) and the inclusion junction counts (IJCs) as calculated

by rMATS at the corresponding loci. We checked the mapping status and generated the

mapping result view using IGV (http://software.broadinstitute.org/software/igv/)

(Robinson et al., 2011).

**Analysis of characteristics for the IR gene group**

We analyzed enriched gene functions and pathways at 142 loci (134 genes) at which

significantly different IR events were detected between KL+ and KL– using DAVID ver

6.8 (https://david.ncifcrf.gov/) (Huang da et al., 2009) with the following criteria: Count

$\geq 1$, EASE $\leq 1$.

**Splice site score**

For the evaluation of splice site strength, maximum entropy scores for 5' and 3' splice

sites were calculated using MaxEntScan (Yeo and Burge, 2004).

**Motif analysis**

For the comparison of protein-binding motifs in the mRNA, we compared motifs with

the ATtRACT database of RNA-biding proteins and their associated motifs (Giudice et

al., 2016). We compared TF-binding sites using HOMER (Heinz et al., 2010) with

default parameters. All detected motifs were then compared using Fisher's test.

**Definition of potential splicing-related genes**

The KEGG pathway lists ~140 proteins as spliceosomal proteins, and we then added
109 genes from the Gene Ontology (GO) list (GO:0000398) of mRNA splicing via
spliceosome. We also added 66 genes from the GO list (GO:0003729) of mRNA
binding and 180 genes from the GO list (GO:0006397) of mRNA processing, both of
which were chosen to overlap with genes listed by Han et al. (2017) that were selected
based on the criterion that a knock-down experiment influenced their alternative
splicing. After removing duplicates, we ultimately defined 250 genes as potential
splicing-related genes in this analysis. They included almost all the spliceosomal
proteins and spliceosome regulatory proteins.

**Reverse transcription PCR amplification of IR loci**

The extracted RNA was treated with Recombinant DNase I (Takara Bio, Japan) to
digest the remaining genomic DNA and was purified by phenol/chloroform/isoamyl
alcohol (25:24:21) and ethanol precipitation. The purified RNA was reverse transcribed
using High-Capacity cDNA Reverse Transcription kits (Thermo Fisher Scientific).
Primers were prepared in exons adjacent to the IR locus, and PCR amplification was
performed using the cDNA. Reaction conditions were as follows: 5 µL 10× PCR buffer
(Takara Bio), 5 µL dNTPs (25 mM; Takara Bio), 1 µL primers (10 pmol/µL each
primer), 0.2 µL ExTaq DNA polymerase (5 U/µL; Takara Bio), 40 ng of template DNA,
and DNase-free water added to a final volume of 50 µL. PCR was performed using
GeneAmp PCR system 9700 (Thermo Fisher Scientific) with the following conditions:
initial annealing at 96°C for 5 min, followed by 25 or 30 cycles (96ºC for 30 sec, 55ºC
for 45 sec, 72ºC for 2 min). After a final extension at 72°C for 5 min, the PCR mixtures
were held at 4°C. The amplicon was confirmed based on size by TapeStation. The
primer set used is listed in supplementary information (SI). Primer sequences were
listed in Table S1.

**Analysis of DEG**

Genes that had significantly differential expression with $P < 0.01$ in two comparisons,
namely KL+/KL– and KL–/WT–, were detected using edgeR (Robinson et al., 2010).
The recovered genes that were significantly down-regulated in KL– as compared with
WT– and significantly up-regulated in KL+ as compared with KL– were extracted.

**Western blot analysis**

Mouse liver samples were homogenized in ice-cold RIPA buffer (50 mM Tris, 150 mM NaCl, 1% NP-40, 0.5% deoxycholic acid sodium monohydrate, 0.1% SDS, 10 mM NaF, pH 7.4) with a Disposable homogenizer (Nippi BioMasher) (30 strokes), and the homogenates were centrifuged at $10,000 \times g$ and 4°C for 20 min. The protein concentration of each supernatant was measured by the BCA protein assay (Thermo Fisher Scientific), and supernatants were diluted to equal protein concentrations, combined with 2 M DTT (final concentration, 0.2 M) and 4× SDS sample buffer (6% SDS, 40% glycerol, 0.4% bromophenol blue, 250 mM Tris, pH 6.8), and boiled for 5 min at 95°C.

Protein samples (30 μg/well) were resolved by SDS-PAGE and then transferred onto PVDF membranes. The membranes were blocked with 5% milk in TBST (20 mM Tris, 150 mM NaCl, containing 0.05% Tween-20, pH 7.4) and incubated with primary antibodies at 4°C overnight. Primary antibodies used in this study were anti-RGN/SMP30 (17947-1-AP, Proteintech), anti-DDX5,p68 (10804-1-AP, Proteintech), anti-NDUFS2 (GTX114924, GeneTex), anti-PPARD (10156-2-AP, Proteintech), anti-SIRT7 (12994-1-AP, Proteintech), anti-NXF1 (10328-1-AP, Proteintech), anti-SRSF6 (11772-1-AP, Proteintech), anti-LXR beta (ab28479, Abcam), and anti-GAPDH (sc-32233, Santa Cruz Biotechnology). After incubation with secondary antibodies peroxidase-conjugated anti–rabbit IgG (SA00001-2, Proteintech) or anti–mouse IgG (sc-516102, Santa Cruz Biotechnology), protein bands were detected using ECL Prime Western Blotting Detection Reagents (GE Healthcare). Some of the membranes were probed with anti-GAPDH , which was used as the loading control for other blots in each experiment. The signal intensity was quantified using ImageJ (NIH) or a ChemiDoc system (Bio-Rad). Western blots were repeated a minimum of three times with different animals, and representative blots are shown.

**Data availability**

The original RNA-seq datasets have been deposited in the DDBJ Sequence Read Archive under accession numbers DRR167982–DRR167990, which are linked to the BioProject accession number PRJDB7898.

**Acknowledgements**

The present work was financially supported by FAIS and Zenick Co. represented by

Mr. Y. Otake and Mr. T. Sugino, respectively. We thank Mr. K. Onodera for

management and discussions and Ms. A. Koizumi for making illustrations.

**Competing interests**

N.O., K.O., Y.I., A.M., E.I., T.V., and S.N., received a research grant from Zenick Co.

and Tsumura & Co. N.F., A.N., A.S., M.N., A.K., and M.Y. are employees of Tsumura

& Co.

**References**

1.    Adusumalli, S., Ngian, Z.K., Lin, W.Q., Benoukraf, T., and Ong, C.T. (2019).

Increased intron retention is a post-transcriptional signature associated with

progressive aging and Alzheimer's disease. *Aging Cell* **18**(3), e12928. DOI:

10.1111/acel.12928.

2.    Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer

for Illumina sequence data. *Bioinformatics* **30**(15), 2114-2120. DOI:

10.1093/bioinformatics/btu170.

3.    Boutz, P.L., Bhutkar, A., and Sharp, P.A. (2015). Detained introns are a novel,

widespread class of post-transcriptionally spliced introns. *Genes Dev* **29**(1), 63-

80. DOI: 10.1101/gad.247361.114.

4.    Braunschweig, U., Barbosa-Morais, N.L., Pan, Q., Nachman, E.N., Alipanahi, B.,

Gonatopoulos-Pournatzis, T., Frey, B., Irimia, M., and Blencowe, B.J. (2014).

Widespread intron retention in mammals functionally tunes transcriptomes.

*Genome Res* **24**(11), 1774-1786. DOI: 10.1101/gr.177790.114.

5.    Deschenes, M., and Chabot, B. (2017). The emerging role of alternative splicing in

senescence and aging. *Aging Cell* **16**(5), 918-933. DOI: 10.1111/acel.12646.

6.    Dvinge, H., and Bradley, R.K. (2015). Widespread intron retention diversifies most

cancer transcriptomes. *Genome Med* **7**(1), 45. DOI: 10.1186/s13073-015-0168-9.

7.    Dvinge, H., Kim, E., Abdel-Wahab, O., and Bradley, R.K. (2016). RNA splicing

factors as oncoproteins and tumour suppressors. *Nat Rev Cancer* **16**(7), 413-430.

DOI: 10.1038/nrc.2016.51.

8.    Fino, K.K., Yang, L., Silveyra, P., Hu, S., Umstead, T.M., DiAngelo, S., Halstead,
E.S., Cooper, T.K., Abraham, T., Takahashi, Y*., et al.* (2017).
SH3GLB2/endophilin B2 regulates lung homeostasis and recovery from severe
influenza A virus infection. *Scientific Reports* **7**(1), 7262. DOI: 10.1038/s41598-
017-07724-5.

9.    Fouache, A., Zabaiou, N., De Joussineau, C., Morel, L., Silvente-Poirot, S., Namsi,
A., Lizard, G., Poirot, M., Makishima, M., Baron, S*., et al.* (2019). Flavonoids
differentially modulate liver X receptors activity-Structure-function relationship
analysis. *J Steroid Biochem Mol Biol* **190**, 173-182. DOI:
10.1016/j.jsbmb.2019.03.028.

10.    Fukao, T., Mitchell, G., Sass, J.O., Hori, T., Orii, K., and Aoyama, Y. (2014).
Ketone body metabolism and its defects. *J Inherit Metab Dis* **37**(4), 541-551. DOI:
10.1007/s10545-014-9704-9.

11.    Fukaya, S., Nagatsu, A., and Yoshioka, H. (2018). The Kampo formula "Juzen-
taiho-to" exerts protective effects on ethanol -induced liver injury mice.
*Fundamental Toxicological Sciences* **5**(3), 105-112. DOI: doi:10.2131/fts.5.105

12.    Giudice, G., Sanchez-Cabo, F., Torroja, C., and Lara-Pezzi, E. (2016). ATtRACT-
a database of RNA-binding proteins and associated motifs. *Database (Oxford)*
**2016**. DOI: 10.1093/database/baw035.

13.    Han, H., Braunschweig, U., Gonatopoulos-Pournatzis, T., Weatheritt, R.J., Hirsch,
C.L., Ha, K.C.H., Radovani, E., Nabeel-Shah, S., Sterne-Weiler, T., Wang, J*., et*
*al.* (2017). Multilayered Control of Alternative Splicing Regulatory Networks by
Transcription Factors. *Mol Cell* **65**(3), 539-553 e537. DOI:
10.1016/j.molcel.2017.01.011.

14.    Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X.,
Murre, C., Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-
determining transcription factors prime cis-regulatory elements required for
macrophage and B cell identities. *Mol Cell* **38**(4), 576-589. DOI:
10.1016/j.molcel.2010.05.004.

15.    Honda, A., Valogne, Y., Bou Nader, M., Brechot, C., and Faivre, J. (2012). An
intron-retaining splice variant of human cyclin A2, expressed in adult
differentiated tissues, induces a G1/S cell cycle arrest in vitro. *PLoS One* **7**(6),
e39249. DOI: 10.1371/journal.pone.0039249.

16.    Huan Li, Z.W., Tianyi Ma, Gang Wei, Ting Ni (2017). Alternative splicing in aging
and age-related diseases. *Translational Medicine of Aging* **1**, 32-40. DOI,
https://www.sciencedirect.com/science/article/pii/S246850111730007X.

17.   Huang da, W., Sherman, B.T., and Lempicki, R.A. (2009). Systematic and
integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat*
*Protoc* **4**(1), 44-57. DOI: 10.1038/nprot.2008.211.

18.   Ishikawa, S., Ishikawa, T., Tezuka, C., Asano, K., Sunagawa, M., and Hisamitsu,
T. (2017). Efficacy of Juzentaihoto for Tumor Immunotherapy in B16 Melanoma
Metastasis Model. *Evid Based Complement Alternat Med* **2017**, 6054706. DOI:
10.1155/2017/6054706.

19.   Jacob, A.G., and Smith, C.W.J. (2017). Intron retention as a component of
regulated gene expression programs. *Hum Genet* **136**(9), 1043-1057. DOI:
10.1007/s00439-017-1791-x.

20.   Kaneko, A., Matsumoto, T., Matsubara, Y., Sekiguchi, K., Koseki, J., Yakabe, R.,
Aoki, K., Aiba, S., and Yamasaki, K. (2017). Glucuronides of phytoestrogen
flavonoid enhance macrophage function via conversion to aglycones by beta-
glucuronidase in macrophages. *Immun Inflamm Dis* **5**(3), 265-279. DOI:
10.1002/iid3.163.

21.   Kataoka, N. (2017). Modulation of aberrant splicing in human RNA diseases by
chemical compounds. *Hum Genet* **136**(9), 1237-1245. DOI: 10.1007/s00439-017-
1789-4.

22.   Kitamura, K., Iino, Y., Kamide, Y., Kudo, F., Nakayama, T., Suzuki, K., Taiji, H.,
Takahashi, H., Yamanaka, N., and Uno, Y. (2015). Clinical practice guidelines for
the diagnosis and management of acute otitis media (AOM) in children in Japan -
2013 update. *Auris Nasus Larynx* **42**(2), 99-106. DOI: 10.1016/j.anl.2014.09.006.

23.   Ko, J.H., Lee, J.H., Choi, B., Park, J.Y., Kwon, Y.W., Jeon, S., Park, S.D., and Kim,
S.N. (2018). Neuroprotective Effects of Gagam-Sipjeondaebo-Tang, a Novel
Herbal Formula, against MPTP-Induced Parkinsonian Mice and MPP(+)-Induced
Cell Death in SH-SY5Y Cells. *Evid Based Complement Alternat Med* **2018**,
2420809. DOI: 10.1155/2018/2420809.

24.   Kuro-o, M., Matsumura, Y., Aizawa, H., Kawaguchi, H., Suga, T., Utsugi, T.,
Ohyama, Y., Kurabayashi, M., Kaname, T., Kume, E.*, et al.* (1997). Mutation of
the mouse klotho gene leads to a syndrome resembling ageing. *Nature* **390**(6655),
45-51. DOI: 10.1038/36285.

25.   Kwak, H., and Lis, J.T. (2013). Control of transcriptional elongation. *Annu Rev*
*Genet* **47**, 483-508. DOI: 10.1146/annurev-genet-110711-155440.

26.   Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie
2. *Nat Methods* **9**(4), 357-359. DOI: 10.1038/nmeth.1923.

27.   Lejeune, F., and Maquat, L.E. (2005). Mechanistic links between nonsense-

mediated mRNA decay and pre-mRNA splicing in mammalian cells. *Curr Opin*
*Cell Biol* **17**(3), 309-315. DOI: 10.1016/j.ceb.2005.03.002.

28.  Lev Maor, G., Yearim, A., and Ast, G. (2015). The alternative role of DNA
methylation in splicing regulation. *Trends Genet* **31**(5), 274-280. DOI:
10.1016/j.tig.2015.03.002.

29.  Liu, Y., Gonzalez-Porta, M., Santos, S., Brazma, A., Marioni, J.C., Aebersold, R.,
Venkitaraman, A.R., and Wickramasinghe, V.O. (2017). Impact of Alternative
Splicing on the Human Proteome. *Cell Rep* **20**(5), 1229-1241. DOI:
10.1016/j.celrep.2017.07.025.

30.  Lopez-Otin, C., Blasco, M.A., Partridge, L., Serrano, M., and Kroemer, G. (2013).
The hallmarks of aging. *Cell* **153**(6), 1194-1217. DOI: 10.1016/j.cell.2013.05.039.

31.  Luco, R.F., Allo, M., Schor, I.E., Kornblihtt, A.R., and Misteli, T. (2011).
Epigenetics in alternative pre-mRNA splicing. *Cell* **144**(1), 16-26. DOI:
10.1016/j.cell.2010.11.056.

32.  Maruyama, N., Ishigami, A., and Kondo, Y. (2010). Pathophysiological
significance of senescence marker protein-30. *Geriatr Gerontol Int* **10 Suppl 1**,
S88-98. DOI: 10.1111/j.1447-0594.2010.00586.x.

33.  Matsumoto, T., Sakurai, M.H., Kiyohara, H., and Yamada, H. (2000). Orally
administered decoction of Kampo (Japanese herbal) medicine, "Juzen-Taiho-To"
modulates cytokine secretion and induces NKT cells in mouse liver.
*Immunopharmacology* **46**(2), 149-161. DOI: 10.1016/S0162-3109(99)00166-6.

34.  Mayeda, A., Munroe, S.H., Caceres, J.F., and Krainer, A.R. (1994). Function of
conserved domains of hnRNP A1 and other hnRNP A/B proteins. *EMBO J* **13**(22),
5483-5495. DOI: 10.1002/j.1460-2075.1994.tb06883.x

35.  Monteuuis, G., Wong, J.J.L., Bailey, C.G., Schmitz, U., and Rasko, J.E.J. (2019).
The changing paradigm of intron retention: regulation, ramifications and recipes.
*Nucleic Acids Res* **47**(22), 11497-11513. DOI: 10.1093/nar/gkz1068.

36.  Munakata, K., Takashima, K., Nishiyama, M., Asano, N., Mase, A., Hioki, K.,
Ohnishi, Y., Yamamoto, M., and Watanabe, K. (2012). Microarray analysis on
germfree mice elucidates the primary target of a traditional Japanese medicine
juzentaihoto: acceleration of IFN-alpha response via affecting the ISGF3-IRF7
signaling cascade. *BMC Genomics* **13**, 30. DOI: 10.1186/1471-2164-13-30.

37.  Naro, C., Jolly, A., Di Persio, S., Bielli, P., Setterblad, N., Alberdi, A.J., Vicini, E.,
Geremia, R., De la Grange, P., and Sette, C. (2017). An Orchestrated Intron
Retention Program in Meiosis Controls Timely Usage of Transcripts during Germ
Cell    Differentiation.    *Dev    Cell*    **41**(1),    82-93    e84.    DOI:

10.1016/j.devcel.2017.03.003.

38.    Niemela, E.H., Oghabian, A., Staals, R.H., Greco, D., Pruijn, G.J., and Frilander,
M.J. (2014). Global analysis of the nuclear processing of transcripts with
unspliced U12-type introns by the exosome. *Nucleic Acids Res* **42**(11), 7358-7369.
DOI: 10.1093/nar/gku391.

39.    Ninomiya, K., Kataoka, N., and Hagiwara, M. (2011). Stress-responsive
maturation of Clk1/4 pre-mRNAs promotes phosphorylation of SR splicing factor.
*J Cell Biol* **195**(1), 27-40. DOI: 10.1083/jcb.201107093.

40.    Nishi, A., Ohbuchi, K., Kushida, H., Matsumoto, T., Lee, K., Kuroki, H.,
Nabeshima, S., Shimobori, C., Komokata, N., Kanno, H.*, et al.* (2017).
Deconstructing the traditional Japanese medicine "Kampo": compounds,
metabolites and pharmacological profile of maoto, a remedy for flu-like
symptoms. *NPJ Syst Biol Appl* **3**, 32. DOI: 10.1038/s41540-017-0032-1.

41.    Ohshima, J., Sasai, M., Liu, J., Yamashita, K., Ma, J.S., Lee, Y., Bando, H.,
Howard, J.C., Ebisu, S., Hayashi, M.*, et al.* (2015). RabGDIalpha is a negative
regulator of interferon-gamma-inducible GTPase-dependent cell-autonomous
immunity to Toxoplasma gondii. *Proc Natl Acad Sci U S A* **112**(33), E4581-4590.
DOI: 10.1073/pnas.1510031112.

42.    Okumi, H., and Koyama, A. (2014). Kampo medicine for palliative care in Japan.
*Biopsychosoc Med* **8**(1), 6. DOI: 10.1186/1751-0759-8-6.

43.    Onishi, Y., Yamaura, T., Tauchi, K., Sakamoto, T., Tsukada, K., Nunome, S.,
Komatsu, Y., and Saiki, I. (1998). Expression of the anti-metastatic effect induced
by Juzen-taiho-to is based on the content of Shimotsu-to constituents. *Biol Pharm*
*Bull* **21**(7), 761-765. DOI: 10.1248/bpb.21.761.

44.    Padgett, R.A., Grabowski, P.J., Konarska, M.M., Seiler, S., and Sharp, P.A. (1986).
Splicing of messenger RNA precursors. *Annu Rev Biochem* **55**, 1119-1150. DOI:
10.1146/annurev.bi.55.070186.005351.

45.    Park, S.M., Kim, S.W., Jung, E.H., Ko, H.L., Im, C.K., Lee, J.R., Byun, S.H., Ku,
S.K., Kim, S.C., Park, C.A.*, et al.* (2018). Sipjeondaebo-tang Alleviates Oxidative
Stress-Mediated Liver Injury through Activation of the CaMKK2-AMPK
Signaling Pathway. *Evid Based Complement Alternat Med* **2018**, 8609285. DOI:
10.1155/2018/8609285.

46.    Patel, A.A., and Steitz, J.A. (2003). Splicing double: insights from the second
spliceosome. *Nat Rev Mol Cell Biol* **4**(12), 960-970. DOI: 10.1038/nrm1259.

47.    Patwardhan, B., Warude, D., Pushpangadan, P., and Bhatt, N. (2005). Ayurveda
and traditional Chinese medicine: a comparative overview. *Evid Based*

*Complement Alternat Med* **2**(4), 465-473. DOI: 10.1093/ecam/neh140.

48.   Pimentel, H., Parra, M., Gee, S.L., Mohandas, N., Pachter, L., and Conboy, J.G.
(2016). A dynamic intron retention program enriched in RNA processing genes
regulates gene expression during terminal erythropoiesis. *Nucleic Acids Res* **44**(2),
838-851. DOI: 10.1093/nar/gkv1168.

49.   Razzaque, M.S. (2009). The FGF23-Klotho axis: endocrine regulation of
phosphate homeostasis. *Nat Rev Endocrinol* **5**(11), 611-619. DOI:
10.1038/nrendo.2009.196.

50.   Robinson, J.T., Thorvaldsdottir, H., Winckler, W., Guttman, M., Lander, E.S., Getz,
G., and Mesirov, J.P. (2011). Integrative genomics viewer. *Nat Biotechnol* **29**(1),
24-26. DOI: 10.1038/nbt.1754.

51.   Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.
*Bioinformatics* **26**(1), 139-140. DOI: 10.1093/bioinformatics/btp616.

52.   Rui, L. (2014). Energy metabolism in the liver. *Compr Physiol* **4**(1), 177-197. DOI:
10.1002/cphy.c130024.

53.   Saiki, I. (2000). A Kampo medicine "Juzen-taiho-to"--prevention of malignant
progression and metastasis of tumor cells and the mechanism of action. *Biol*
*Pharm Bull* **23**(6), 677-688. DOI: 10.1248/bpb.23.677.

54.   Saiki, I., Koizumi, K., Goto, H., Inujima, A., Namiki, T., Raimura, M., Kogure, T.,
Tatsumi, T., Inoue, H., Sakai, S.*, et al.* (2013). The long-term effects of a kampo
medicine, juzentaihoto, on maintenance of antibody titer in elderly people after
influenza vaccination. *Evid Based Complement Alternat Med* **2013**, 568074. DOI:
10.1155/2013/568074.

55.   Shalgi, R., Hurt, J.A., Lindquist, S., and Burge, C.B. (2014). Widespread inhibition
of posttranscriptional splicing shapes the cellular transcriptome following heat
shock. *Cell Rep* **7**(5), 1362-1370. DOI: 10.1016/j.celrep.2014.04.044.

56.   Shen, S., Park, J.W., Lu, Z.X., Lin, L., Henry, M.D., Wu, Y.N., Zhou, Q., and Xing,
Y. (2014). rMATS: robust and flexible detection of differential alternative splicing
from replicate RNA-Seq data. *Proc Natl Acad Sci U S A* **111**(51), E5593-5601.
DOI: 10.1073/pnas.1419161111.

57.   Shoemaker, J.E., Lopes, T.J.S., Ghosh, S., Matsuoka, Y., Kawaoka, Y., and Kitano,
H. (2012). CTen: a web-based platform for identifying enriched cell types from
heterogeneous microarray data. *BMC Genomics* **13**(1), 460. DOI: 10.1186/1471-
2164-13-460.

58.   Solana, R., Pawelec, G., and Tarazona, R. (2006). Aging and innate immunity.

*Immunity* **24**(5), 491-494. DOI: 10.1016/j.immuni.2006.05.003.

59.  Stegeman, R., and Weake, V.M. (2017). Transcriptional Signatures of Aging. *J Mol*
*Biol* **429**(16), 2427-2437. DOI: 10.1016/j.jmb.2017.06.019.

60.  Takeno, N., Inujima, A., Shinohara, K., Yamada, M., Shibahara, N., Sakurai, H.,
Saiki, I., and Koizumi, K. (2015). Immune adjuvant effect of Juzentaihoto, a
Japanese traditional herbal medicine, on tumor vaccine therapy in a mouse model.
*Int J Oncol* **47**(6), 2115-2122. DOI: 10.3892/ijo.2015.3208.

61.  Trapnell, C., Pachter, L., and Salzberg, S.L. (2009). TopHat: discovering splice
junctions with RNA-Seq. *Bioinformatics* **25**(9), 1105-1111. DOI:
10.1093/bioinformatics/btp120.

62.  Ullrich, S., and Guigo, R. (2019). Dynamic changes in intron retention are tightly
associated with regulation of splicing factors and proliferative activity during B-
cell development. *Nucleic Acids Res*. DOI: 10.1093/nar/gkz1180.

63.  UNESCO (2011). Huang Di Nei Jing (Yellow Emperor's Inner Canon) (Beijing:
National Library of China ).

64.  Vu, T.-D., Iwasaki, Y., Shigenobu, S., Maruko, A., Oshima, K., Iioka, E., Huang,
C-L. Abe, T., Tamaki, S., Lin, Y-W., Chen, C-K., Lu, M-Y., Hojo, M., Wang, H-
V., Tzeng, S-F., Huang, H-J., Kanai, A,, Gojobori, T., Chiang, T-Y., Sun, H.S., Li,
W-H., Okada, N. (2019). Behavioral and neurotranscriptomic synchronization
between the two opponents of a fighting pair of the fish Betta splendens. *PLOS*
*Biology*.  submitted.

65.  Wang, Z., Qi, F., Cui, Y., Zhao, L., Sun, X., Tang, W., and Cai, P. (2018). An update
on Chinese herbal medicines as adjuvant treatment of anticancer therapeutics.
*Biosci Trends* **12**(3), 220-239. DOI: 10.5582/bst.2018.01144.

66.  Weischenfeldt, J., Lykke-Andersen, J., and Porse, B. (2005). Messenger RNA
surveillance: neutralizing natural nonsense. *Curr Biol* **15**(14), R559-562. DOI:
10.1016/j.cub.2005.07.002.

67.  Wong, J.J., Ritchie, W., Ebner, O.A., Selbach, M., Wong, J.W., Huang, Y., Gao, D.,
Pinello, N., Gonzalez, M., Baidya, K.*, et al.* (2013). Orchestrated intron retention
regulates normal granulocyte differentiation. *Cell* **154**(3), 583-595. DOI:
10.1016/j.cell.2013.06.052.

68.  Xu, W., Wei, Z., Dong, J., Duan, F., Chen, K., Chen, C., Liu, J., Yang, X., Chen,
L., Xiao, H.*, et al.* (2017). Global Metabolomics Reveals the Metabolic
Dysfunction in Ox-LDL Induced Macrophage-Derived Foam Cells. *Front*
*Pharmacol* **8**, 586. DOI: 10.3389/fphar.2017.00586.

69.  Yeo, G., and Burge, C.B. (2004). Maximum entropy modeling of short sequence

motifs with applications to RNA splicing signals. *J Comput Biol* **11**(2-3), 377-394. DOI: 10.1089/1066527041410418.

70. Yoshioka, H., Fukaya, S., Miura, N., Nagatsu, A., and Nonogaki, T. (2016). Hepatoprotective effect of kampo formula "Juzen-taiho-to" on bromobenzene-induced toxicity in mice. *Fundamental Toxicological Sciences* **3**(5), 233-236. DOI: 10.2131/fts.3.233.

71. Yoshioka, H., Fukaya, S., Tominaga, S., Nagatsu, A., Miura, N., and Maeda, T. (2019). Protective effect of the Kampo formula "Juzen-taiho-to" on isoniazid- and rifampicin-induced hepatotoxicity in mice. *Fundamental Toxicological Sciences* **6**(1), 25-29. DOI: 10.2131/fts.6.25.

72. Zane, L., Sharma, V., and Misteli, T. (2014). Common features of chromatin in aging and cancer: cause or coincidence? *Trends Cell Biol* **24**(11), 686-694. DOI: 10.1016/j.tcb.2014.07.001.

73. Zelcer, N., and Tontonoz, P. (2006). Liver X receptors as integrators of metabolic and inflammatory signaling. *J Clin Invest* **116**(3), 607-614. DOI: 10.1172/JCI27883.

74. Zhang, J., and Manley, J.L. (2013). Misregulation of pre-mRNA alternative splicing in cancer. *Cancer Discov* **3**(11), 1228-1237. DOI: 10.1158/2159-8290.CD-13-0253.

75. Zheng, H.C., Noguchi, A., Kikuchi, K., Ando, T., Nakamura, T., and Takano, Y. (2014). Gene expression profiling of lens tumors, liver and spleen in alpha-crystallin/SV40 T antigen transgenic mice treated with Juzen-taiho-to. *Mol Med Rep* **9**(2), 547-552. DOI: 10.3892/mmr.2013.1854.

76. Zhou, X., Seto, S.W., Chang, D., Kiat, H., Razmovski-Naumovski, V., Chan, K., and Bensoussan, A. (2016). Synergistic Effects of Chinese Herbal Medicine: A Comprehensive Review of Methodology and Current Research. *Front Pharmacol* **7**, 201. DOI: 10.3389/fphar.2016.00201.

1068

**Figure 1. Klotho mice at 7 weeks of age represent the pre-disease state**

(A) Schematic illustration of animal experiments. Three-week-old male α-klotho

knockout ($Kl^{-/-}$/Jcl) mice (klotho mice) (n = 8) and WT (C57BL/6JJcl) mice (n = 8)

were purchased from CLEA Japan, Tokyo, Japan. The mice were acclimated for 0.5

weeks in a vinyl isolator, during which they were given radiation-sterilized water and

CE-2 diet (CLEA Japan) ad libitum. Thereafter, the klotho mice and WT mice were
each divided into two groups, with four animals in each. In each case, one group of
mice was fed CE-2 containing 0.5% (w/w) JTT, whereas the other group was fed CE-2
only from 3.5 weeks of age until the mice were 7 weeks old. Ope stands for operation.
(B) Estimation of the half-life of klotho mice. N = 5. (C) Histological observations of
liver tissues from KL+ and KL− showed that they did not exhibit any signs of aging
relative to WT mice. (D) The expression of seven proteins, each of which showed IR,
was examined by western blotting. Expression levels did not change in KL+, KL−, and
WT− at 7 weeks of age. (E) Representative western blotting data (upper) of a
senescence marker protein, RGN/SMP30, together with its quantified expression
(lower) show that KL− are not senescent. Statistical significance was estimated by one-
way ANOVA. Error bars indicate mean ±standard deviations of triplicate
measurements.
1087
1088

1089

1090

**Figure 2. A metabolome analysis showed that the klotho mice are undergoing a**

**starvation-like condition at 7 weeks of age**

Based on a metabolome analysis of (A) 3-HBA and (B) other glycolysis-related

metabolites, KL− data indicate that these mice are undergoing a starvation-like

condition. *$P \leq 0.05$, **$P \leq 0.001$, unpaired Student's t-test. Based on the metabolome

analysis about metabolites related to RNA precursors, the concentrations of some

nucleotides were recovered in KL+ in comparison with KL− and WT−. *$P \leq 0.05$, **$P$

$\leq 0.001$, unpaired Student's t-test.

1099

1100

1103

**Figure 3. DEG analysis showed that liver function in klotho mice improved after administration of JTT**

Number of up-regulated genes and down-regulated genes from the DEG data for the comparison (A) of KL– vs WT– and (B) of KL+ vs KL–. Cten analysis of (C) 1228 down-regulated genes in KL– in the comparison of KL– vs WT–, (D) 1352 up-regulated

genes in KL–in the comparison of KL– vs WT–, and (E) 532 up-regulated genes in KL+

in the comparison of KL+ vs KL–. (F) A Venn diagram of transcriptome comparisons

from KL+ vs KL– and KL– vs WT–. Two hundred thirteen genes were shared between

the comparisons. (G) Box plots of 213 gene expression data show a significant decrease

in KL– in comparison with that in WT– and a significant increase in KL+ in

comparison with KL–. Values along the vertical axis are normalized based on the

overall average. (H) Cten analysis (http://www.influenza-x.org/~jshoemaker/cten/) for

213 genes that were differentially expressed in the comparison between KL+ vs KL–

and between KL– vs WT–. The horizontal line indicates $P = 0.01$. (I) Bar graphs

showing $-\log_{10} P$ of GO terms enriched for the 213 genes shown in (F). Only those

terms with $P < 0.01$ are shown.

1120

1121

1122

1123

1124

1125

1126

**Figure 4. Quantification of IR events**

(A) Among the four organs examined, liver had the most notable recovery of retained

introns after administration of JTT. Bar graphs show the number of retained introns that

decreased (in orange) or increased (in blue) in the four organs as a result of

administration of JTT. Based on the rMATS analysis, genes with a significant change in

IR in the comparison of KL+ vs KL− were divided into two groups. In the liver, the first

group includes genes that showed a decrease in IR in KL+ (132 loci; DecIR type,

implying that the fold change of KL+ was larger than that of KL−), and the second

includes genes with an increase in IR in KL+ (10 loci; IncIR type, implying that the fold change of KL+ was smaller than that of KL–). (B) The quantification of IR events from mRNA-seq data using rMATS. The IJCs represent the reads containing the intron sequence at the junction. The SJCs represent the reads without intron sequences at the junction. (C) (i) The mapping results from KL+, KL–, and WT– are shown for a single gene, *Anxa11*, using Integrative Genomics Viewer (IGV). Thick black bars indicate the sites of junctions in the reads where SJCs and IJCs were analyzed. (ii) Read counts from KL+, KL–, and WT– for IJCs and SJCs. (iii) FC in the number of SJCs relative to IJCs for KL+, KL–, and WT–.

**Figure 5. Seventy loci of complete recovery show a typical V-shaped pattern of recovery**

(A) A Venn diagram of two types of loci, the DecIR type (blue circle) in the comparison of KL+ and KL– and the IncIR type (red circle) in the comparison of KL– and WT–. Three categories, namely "partial recovery", "complete recovery", and "no recovery" loci, are distinguishable. (B) Heat map illustration of FC values of loci in KL+, KL–, and WT– for the 70 "complete recovery" loci. (C–E) Boxplots of the FC values of (C) the 70 "complete recovery" loci, (D) the 62 "partial recovery" loci, and (E) the 120 "no-recovery" loci in KL+, KL–, and WT–.

1160

1161

**Figure 6. Retained intron loci have distinguishing characteristics**

(A–D) Boxplots showing (A) intron lengths, (B) the GC percentage in intron sequences,

and the strength score of the (C) 5' and (D) 3' splice sites compared among three groups

of introns, namely "no recovery", "complete recovery", and all introns from liver-

expressed genes (254,005 loci). *$P \leq 0.05$, ***$P \leq 0.001$, unpaired Student's *t*-test; ns,

not significant. (E, F) Bar graphs showing percentage of loci in the "complete recovery"

and "no recovery" groups that have (E) motifs for RNA-binding proteins within the

intron or (F) motifs for DNA-binding proteins in the promoter region. **$P \leq 0.01$, ***$P$

$\leq 0.001$, Fisher's test.

1171

1172

1173

1174

**Figure 7. RT-PCR validation of retained introns and their recovery after**

**administration of JTT**

(A–H) Eight genes, (A) *Sirt7*, (B) *Decr2*, (C) *Ddx5*, (D) *Acadm*, (E) *Cyp27a1*, (F)

*Hsd3b7*, (G) *Pcyt2*, and (H) *Mug1*, were subjected to the following analyses. (i) IGV

mapping of reads from KL+, KL–, and WT–. The numbers shown to the left of each

map represent read counts. (ii) Bar graphs showing fold changes in SJCs relative to IJCs

for KL+, KL–, and WT–. (iii) RT-PCR validation of RNA expression from KL+, KL–,

and WT–. A and B indicate exons, with the intervening intron indicated for the higher-

molecular-weight product. (iv) The ratio of each transcript as determined using

TapeStation. Data from KL+, KL–, and WT– are shown in red, blue, and green,

respectively.

1186

1187

**(A)**

**(B)**

**(C)**

1188

1189

**Figure 8. Genes with recovered retained introns in liver have the characteristics of liver-specific genes.**

(A) Bar graphs showing $-\log_{10} P$ of GO terms enriched for genes whose IR events were significantly changed in the liver. Red bars show enrichment of GO terms in genes with the DecIR type in KL+ in the KL+ and KL– comparison, and blue bars show enrichment of GO terms in genes with the IncIR type in KL– in the KL– and WT– comparison. (B) Bar graphs showing $-\log_{10} P$ of GO terms enriched for genes defined as "complete recovery" (red bars) and "no recovery" (blue bars) loci. (C) Cten analysis for genes expressed in the liver with the DecIR type in red and those with the IncIR type in blue as shown in Fig. 5A. Horizontal lines indicate $P = 0.05$ (lower line) and 0.01 (upper line). (D) The mapping results from IGV for five gene loci, (i) *Nxf1*, (ii) *Srrm1*, (iii) *Thoc2*, (iv) *Nr1i3*, and (v) *Ppard*, are shown (upper). Bar graphs (lower) show the FC using the IJC and SJC values for KL+, KL–, and WT–. (E) Eleven genes were shared in common between the liver and blood in the recovery process of IR after administration of JTT. The Venn diagram (left) of the DecIR genes that were shared in the liver and the blood. The overlapping genes between blood and liver are listed (right), among which splicing-related genes are shown in red.

1207

1208

1209
1210
1211

**(D)**

(i)

(ii)

| TF | num of genes |
|---|---|
| KLF14(Zf) | 121 |
| Sp2(Zf) | 120 |
| Sp5(Zf) | 104 |
| KLF5(Zf) | 102 |
| SCL(bHLH) | 94 |
| THRb(NR) | 92 |
| Zac1(Zf) | 84 |
| KLF6(Zf) | 83 |
| ETV1(ETS) | 82 |
| AR-halfsite(NR) | 81 |

**(E)**

(i)

(ii)

| TF | num of genes |
|---|---|
| Sp2(Zf) | 93 |
| KLF14(Zf) | 92 |
| Sp5(Zf) | 79 |
| Tgif2(Homeobox) | 65 |
| Sp1(Zf) | 52 |
| NFY(CCAAT) | 45 |
| Hoxd11(Homeobox) | 43 |
| Hoxa13(Homeobox) | 42 |
| Hoxa11(Homeobox) | 40 |
| Hoxd13(Homeobox) | 31 |

**Figure 9. A very few TFs may control genes with DecIR as well as those with IncIR**

(A) Bar graphs showing $-\log_{10}P$ of enrichment of the TF-binding motif in genes with IncIR from blood and liver. (B) Bar graphs showing $-\log_{10}P$ of enrichment of the TF-binding motif in genes with DecIR from blood and liver. (C) Bar graphs showing $-\log_{10}P$ of enrichment of the TF-binding motif in genes with IncIR and DecIR from liver. (D) Genes with IncIR may be controlled by a very few TFs. (i) Venn diagrams of the IncIR genes that have TF-binding motifs in the liver and (ii) the top 10 motifs represented in these genes. (E) Genes with DecIR may be controlled by a very few TFs. (i) Venn diagrams of the DecIR genes that have TF-binding motifs in the liver and (ii) the top 10 motifs represented in these genes.

**Figure 10. IR is a marker of the pre-disease state and the recovery of retained introns suggests the recovery to the normal state.**

(A) Illustration of the functional orchestration of IR-recovered genes under surveillance of homeostasis within the host. Two categories of genes that are involved are shown: alternative splicing and the biological function lipid/glucose metabolism. We hypothesize that the surveillance of homeostasis affects which genes undergo IR recovery after JTT administration during the earlier stage of aging (7 weeks of age) from the pre-disease state to the normal state (see the Discussion). Except for genes for alternative splicing and lipid/glucose metabolism, several genes involved in mitochondria function and the immune system including macrophage function were included among 124 IR-recovered genes (Fig. 5A; 132 loci). *Polrmt, Uros, Sh3glb2, Idh3, Cyp27a1*, and *Hsd3b7* are involved in mitochondrial function. JTT protects mitochondrial function (Park et al., 2018), and a JTT-related multi-herbal medicine prevents mitochondrial dysfunction, providing a possible treatment for the prevention of Parkinson disease (Ko et al., 2018). In addition, the anti-tumor effect of JTT may rely partly on activation of mitochondria (Zheng et al., 2014). *Gdi1, Sh3glb2, Faah*, and LXRβ are involved in the immune system including macrophage function and interferon activity. JTT has been widely used as a palliative to treat cancer patients (Okumi and Koyama, 2014) and also to treat patients with bacterial infections such as acute otitis media (Kitamura et al., 2015). JTT also contributes to the maintenance of antibody titer in elderly people after influenza vaccination (Saiki et al., 2013). A number of animal studies have demonstrated an adjuvant effect of JTT, especially with respect to activation of innate immune cells such as macrophages, dendritic cells, natural killer cells, and natural killer T cells (Ishikawa et al., 2017). Microarray analysis indicates that the target of JTT might be the transcription machinery regulating the steady-state level of the type 1 interferon-related genes (Munakata et al., 2012). Oral administration of JTT can elevate the phagocytic activity of macrophage-like cells in the various tissues (Saiki, 2000, Takeno et al., 2015). Similarly, the phytoestrogen flavonoids found in Glycyrrhizae radix, one of the components of JTT, can enhance phagocytosis in a murine macrophage cell line, as we described previously (Kaneko et al., 2017). JTT contains various types of flavonoids, some of which have been identified as selective agonists of LXR, according to molecular-docking analysis *in silico* (Fouache et al., 2019). LXRs, PPARs, and CARs are nuclear receptors and regulate genes related to lipid/glucose metabolism and macrophage function. (B) Schematic representation of the disease process from the healthy state through the pre-disease state. Possible changes that occur in the body are shown with arrows. (C) A model to explain why a specific

subset of genes with IR was recovered in KL+ by administration of JTT. (i) In the case

of KL−, it is presumed that ~200 pre-mRNAs containing retained introns were stored in

the nucleus because of the stress signal induced by aging so that the cells can save

energy. (ii) In the case of KL+, as liver metabolic stress was recovered after JTT

administration, the improved condition of the liver cells led to enhanced post-

transcriptional splicing among pre-mRNAs with retained introns involved in

metabolism.

1272

1273

1274 **Table 1-1.** Number of alternative splicing loci between KL+ and KL–

| | A3SS | | A5SS | | SE | | IR | | MXE | | Total | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | $P < 0.01$ | FDR <0.05 | $P < 0.01$ | FDR < 0.05 | $P < 0.01$ | FDR <0.05 | $P < 0.01$ | FDR <0.05 | $P < 0.01$ | FDR <0.05 | $P < 0.01$ | FDR <0.05 |
| **Blood** | 16 | 1 | 13 | 4 | 58 | 6 | 69 | 10 | 4 | 0 | 160 | 21 |
| **Kidney** | 17 | 2 | 13 | 5 | 96 | 18 | 33 | 9 | 3 | 1 | 162 | 35 |
| **Heart** | 16 | 0 | 3 | 3 | 90 | 2 | 38 | 5 | 7 | 1 | 154 | 11 |
| **Liver** | 63 | 16 | 35 | 9 | 121 | 46 | 142 | 80 | 7 | 2 | 368 | 153 |

FDR: false discovery rate

A5SS: alternative 5' splice site

A3SS: alternative 3' splice site

SE: skipped exon

IR: Intron retention

MXE: mutually exclusive exons

1281

1282

1283 **Table 1-2.** Number of alternative splicing loci with significant difference in liver

| | SE | A5SS | A3SS | MXE | IR | Total |
|---|---|---|---|---|---|---|
| **KL+/KL-** | 121 | 35 | 63 | 7 | 142 | 368 |
| **KL+/WT+** | 222 | 37 | 59 | 6 | 90 | 414 |
| **KL+/WT-** | 199 | 31 | 43 | 7 | 59 | 339 |
| **KL-/WT+** | 294 | 67 | 121 | 16 | 382 | 880 |
| **KL-/WT-** | 205 | 36 | 77 | 9 | 212 | 539 |
| **WT+/WT-** | 90 | 23 | 25 | 5 | 56 | 199 |

1284

1285

1286

1287

1288

1289

1290 **Table 2.** Increase or decrease in IR ratio of splicing-related genes in blood, heart, kidney and liver

1291 by administration of JTT

| Gene Symbol | IR locus | Blood | Heart | Kidney | Liver |
|---|---|---|---|---|---|
| *Cdk11b* | 4:155625654-155625726 | | | | ↓ |
| *Cdk16* | X:20696836-20696921 | ↓ | | | |
| *Cdk5* | 5:24422388-24422497 | | | | ↓ |
| *Celf1* | 2:91013402-91016565 | ↓ | | | |
| *Ddx5* | 11:106782693-106783958 | ↓ | | | ↓ |
| *Hnrnpl* | 7:8818532-28818643 | | ↑ | | |
| *Nxf1* | 19:8765031-8766796 | | | | ↓ |
| *Sf3b1* | 1:55014562-55016375 | ↓ | | | ↓ |
| *Srrm1* | 4:135344669-135346734 | | | | ↓ |
| *Srsf5* | 12:80947865-80949094 | ↓ | | ↓ | ↓ |
| *Srsf6* | 2:162932142-162933425 | ↓ | | | ↓ |
| *Thoc2* | X:41822871-41823361 | | | | ↓ |
| *U2af1l4* | 7:30564101-30564202 | | | ↑ | |

1292

1293

1294

1295

1296