sciCNV: High-throughput paired profiling of transcriptomes and DNA copy

number variations at single cell resolution

Authors: Ali Mahdipour-Shirayeh, Natalie Erdmann, Chungyee Leung-Hagesteijn and Rodger E.

Tiedemann[*]

Princess Margaret Cancer Centre, University Health Network, Toronto, Ontario, Canada

Departments of Medicine and Medical Biophysics, University of Toronto, Toronto, Ontario,

Canada

*Senior and corresponding author: Rodger E. Tiedemann: rodger.tiedemann@uhn.ca

SUMMARY

Chromosome copy number variations (CNVs) are a near-universal feature of cancer however

their effects on cellular function are incompletely understood. Single cell RNA sequencing

(scRNA-seq) can reveal cellular gene expression however cannot directly link this to CNVs.

Here we report new normalization methods (RTAM1 and -2) for scRNA-seq that improve gene

expression alignment between cells, enhancing gene expression comparisons and the application

of scRNA-seq to CNV detection. We also report sciCNV, a pipeline for inferring CNVs from

RTAM-normalized data. Together, these tools provide dual profiling of transcriptomes and

CNVs at single-cell resolution, enabling exploration of the effects of cancer CNVs on cellular

programs. We apply these tools to multiple myeloma (MM) and examine the cellular effects of

cancer CNVs +8q. Consistent with prior reports, MM cells with +8q22-24 upregulate MYC,

MYC-target genes, mRNA processing and protein synthesis, verifying the approach. Overall, we

provide new tools for scRNA-seq that enable matched profiling of the CNV landscape and

transcriptome of single cells, facilitate deconstruction of the effects of cancer CNVs on cellular

reprogramming within single samples.

INTRODUCTION

Aneuploidy and focal CNVs are a pervasive feature of cancer[1]. However, the specific influence

of many cancer CNVs on the transcription programs and molecular processes within cells

remains only partially understood[2,3]. While may studies have sought to integrate cancer CNVs

with gene expression effects by examining differences across tumor samples, such studies are

often impeded by the extensive genetic heterogeneity that exists between tumors, confounding

the attribution of transcriptional variances to single genetic loci. As a consequence, integrative

inter-tumor studies commonly require the profiling of large numbers of samples to minimize

confounding variations, and their application to less common cancer types or CNVs may be

problematic.

To assess the transcriptional effects of specific CNVs, examination of the intra-clonal effects of

emergent CNV at single cell resolution would have the advantage that subclonal cells with and

without CNV are otherwise virtually isogenic, and that gene expression changes could be more

readily attributed to the CNV. Therefore, simultaneous sequencing of single cell genomes and

transcriptomes should in theory enable direct pairing of CNVs with their transcriptional

outcomes with high precision. Notably, sequencing of both genomic DNA and RNA within

single cells has been reported[4-6]. Prior techniques, however, provided profiling of only a few

cells and thus afford only a limited view of the heterogeneity within any cancer. For the

identification of CNV function, sequencing of much larger numbers of cells is required to

capture intra-tumor subclones and to adequately profile subclonal populations for the gene

expression pattern that aligns with an individual CNV.

Whilst scRNA-seq can reveal the transcription state of single cells, it cannot directly relate this to

DNA lesions. CNVs can be inferred from scRNA-seq, which could thus be leveraged to provide

both layers of omics information within individual cells. However, previously reported

approaches[7-9] reveal constraints imposed by the sparsity of single-cell data and have not reported

quantitative measures of accuracy. In particular, inconsistencies in the detection of lowly-

expressed genes within single cells causes stochastic noise that influences transcriptome

distribution and interferes with RNA-based CNV detection. Normalization is thus critical for

accurate scRNA-seq interpretation[10-14] and for secondary CNV detection.

Here we report RTAM1 and RTAM2 normalization methods for scRNA-seq that improve gene

expression alignment between cells and thus enhance the sensitivity of scRNA-seq for the

detection of small expression changes arising from gene copy number differences. We also

report sciCNV, a new pipeline for inferring CNVs in single cells from RTAM-normalized

scRNA-seq data. Together, these methods enable high-throughput paired profiling of both DNA

copy number and RNA in the same cells, facilitating direct intra-clonal examination of the

effects of cancer CNVs on gene expression programs at a cellular level.

RESULTS

Enhanced single-cell RNA-seq normalization methods: RTAM1 and -2

Single-cell RNA-seq enables gene expression comparisons between cells. However, the accuracy

of these comparisons depends critically upon data normalization. Normalization is essential to

convert raw RNA transcript data from single cells into gene expression values that are

objectively comparable between cells, eradicating biases related to cellular sequencing depth,

transcriptome size, technical factors or batch effect. As the best methods for normalizing

scRNA-seq remain controversial, each with its own drawbacks, we developed RTAM1 and -2

(described in the **online methods** and **supplementary figures S1-3**) and compared the RTAM

methods with other normalization strategies currently in use.

To compare the methods for their control of systemic and stochastic variations between cells due

to size or sequence depth, we generated scRNA-seq data for normal and malignant cells

belonging uniformly to the B cell lineage (n>15,000 cells) (**figure 1a**). We examined cells with a

similar ancestry in order to minimize confounding variations between them due to lineage

specification. However, we deliberately generated data from a mix of both small quiescent B

cells and large proliferating plasma cells to ensure that the normalization methods would be

challenged by cells embodying a full spectrum of sizes and transcriptional activities. The cells

were isolated from human bone marrow samples by FACS and were profiled using the 10X

Genomics single cell RNA-seq library kit. Cell- and gene-specific transcripts were enumerated

using barcoded unique molecular identifiers (UMI).

The raw scRNA-seq data from one of the test samples is depicted in **figure 1b**. As shown, the

distributions of transcript numbers per gene varied significantly from cell to cell, predominantly

reflecting differences in the cellular transcriptome sizes and demonstrating a clear need for

normalization. The test samples were next normalized using either TPM[15], SCRAN[11], SCONE[12]

or Seurat's SCTransform function[16] (**figure 1b** and **supplementary figures S4-S20**). To

compare the alignments of the normalized transcriptomes, we examined the mean and median

expression in each cell of a curated list of housekeeping genes (HKG) known to be broadly

expressed with low variation[9]. We also examined the average expression in each cell of all of the

ubiquitously-expressed genes (UEG) detected in >95% of the cells in the sample. For each

sample tested, the UEG represent the largest possible set of genes that are commonly expressed

across the test cells. Whilst the expression of any individual gene is expected to vary between

cells for both biological and technical reasons, the average expression per cell of a large set of

ubiquitous genes should be similar, particularly amongst cells of the same lineage, and its

variance between cells provides a metric of normalization effectiveness.

As shown in **figures 1c, 2** and **supplementary figures S4-20**, TPM, which normalizes cellular

transcriptomes primarily by their total transcript count, produced a very large variance in the

average expression of HKG or UEG between individual cells, suggesting significant limitations

for scRNA-seq application. By comparison, SCRAN and SCONE produced superior alignment

of gene expression averages across cells. However, SCONE, which produced the better

alignment, achieved this only by implementing quantile normalization – exchanging the actual

distribution of transcript counts in each cell for a standardized distribution – which caused a

significant loss of inter-cellular variation, particularly in key highly-expressed genes. The

expression of IGH or IGL genes, for example, a critical feature of plasma cells, was reduced by

quantile normalization into a virtual constant across cells (**supplementary figure S21**).

As each of these scRNA-seq normalization methods has limitations, we developed RTAM1 and -

2. The RTAM methods originate from a consideration of the strengths and weaknesses of

scRNA-seq. Whereas lowly expressed genes are detected within single cells with low resolution

(due to low integer transcript counts) and show significant stochastic variation, highly expressed

genes are robustly detected and show finer quantisation of variation relative to intensity. RTAM1

and -2 thus utilize highly-expressed genes, whose expression is resolved with greater accuracy,

to align cellular transcriptomes. Genes are ranked in each cell by their expression and the

summed intensities of the top-ranked genes is standardized in log-space using unique non-linear

cell- and gene-specific adjustments of gene expression determined either by cellular gene

expression rank (RTAM1) or by gene expression intensity (RTAM2) (see **methods**).

Importantly, compared to TPM, SCTransform or SCRAN, both RTAM1 and RTAM2 reduce the

cell-to-cell variance in the average (median or mean) expression of HKG and UEG sets (**figure 2**

and **supplementary figures S4-S20**). The coefficients of variation (CV) produced by each

normalization method for the "average" expression of HKG or UEG in individual cells is shown

in **figure 2**, for 3 independent patient samples. As shown, RTAM1 and RTAM2 reduce

variations in the average gene expression of single cells, irrespective of whether this average

expression is calculated by any of 3 different methods. By design, the RTAM methods also

standardize the average expression of highly-expressed genes, and thus overall these methods

produce superior alignments of cellular transcriptomes and of gene expression between cells. At

the same time, both RTAM1 and RTAM2 maintain the original variability observed between

cells in the expression of individual highly-transcribed genes, unlike the quantile normalization

implemented by SCONE (**supplementary figure S21**). Overall, therefore, the RTAM methods

represent useful new strategies for normalizing scRNA-seq data that can enhance the accuracy of

gene expression comparisons between cells.

Single-cell inferred chromosomal copy number variation: sciCNV

We next sought to develop a method for detecting single-cell chromosomal CNV from scRNA-

seq, leveraging the enhanced normalization provided by RTAM to increase the sensitivity of

single-cell transcriptomics for CNV detection. To optimize DNA copy number estimates from

gene expression, and to mitigate against data sparsity in single cells, we developed a two-

pronged approach, called sciCNV (described in the **methods**). Briefly, for each cell RTAM-

normalized gene expression was aligned with that of control diploid cells to develop expression

disparity scores, which were averaged in a moving window across the genome. The average gene

expression of the control cells was weighted by the probability of gene detection, enhancing its

comparison with single cell data, where signal '*dropout*' was common for many genes. In a

parallel method, the expression disparity values for each gene were exchanged for binary +1 or -

1 values, which were summed cumulatively in each cell as a function of genomic location; the

gradient of this function yielded a second estimate of the single cell CNV profile that was

sensitive to small concordant expression variations in contiguous genes and that was insensitive

to large single-gene variations. The CNV estimates of the two methods were then combined by

their geometric mean.

Figure 3 shows sciCNV applied to scRNA-seq data from primary multiple myeloma (MM) cells.

Significantly, the CNV profile of a single MM cell, inferred from its RNA, closely resembles the

average CNV profile of $>10^4$ tumor bulk cells, derived from whole exome DNA sequencing

(WES) of the same MM sample (R^2=0.72) (**figure 3a,b**). In addition, the CNV predictions

produced from a single cell by sciCNV also aligned with CNV estimates provided by FISH

(**figure 3c**). Furthermore, examination of >1,700 individual plasma cells from the same MM

patient biopsy using sciCNV revealed that the tumor clone CNVs were robustly detected in each

of the MM cells (**supplementary figure S22a**), despite biological and technical variations

between them; and were not detected in normal plasma cells (NPC). Thus, sciCNV can utilize

scRNA-seq data to reveal CNVs in single cells. Moreover, it can distinguish cancer cells and

normal cells on the basis of their CNV profile (**supplementary figure S22b,c**).

To further examine the accuracy and consistency of sciCNV in detecting tumor CNVs in single

cells, we next compared the sciCNV profiles of thousands of individual plasma cells from 4 MM

samples with the corresponding tumor CNV profiles generated from DNA sequencing of $>10^6$

tumor cells (**figure 4**). Remarkably, the single cell CNV profiles calculated by sciCNV from

RNA substantively recapitulated the complex CNV landscape of each tumor clone, defined by

tumor bulk DNA sequencing. Thus, for the MM199 sample, gains within 1q, 7q, 8q, 9, 17q, 18q,

19p and Xq, and losses within 1p, 2q, 4p, 6p, 6q, 13, 14q, 16p, 17p, 17q, and Xp were all

identified by sciCNV at a single cell level, and these corresponded to CNVs identified by tumor

bulk WES (**figure 4a**). Similarly, for sample MM241, trisomies of chromsomes 5, 7 and 19 and

monosomy of chromosome 13 were all identified by sciCNV in single MM cells, matching the

WES-defined CNV profile of pooled tumor bulk cells (**figure 4b**). Furthermore, sciCNV also

identified a subpopulation of MM241 cells with subclonal gain of 1q, and the existence of this

subclone appeared to be borne out by the tumor DNA WES profile which also identified a 1q-

arm copy gain with an average CNV peak height of approximately 35%. Similarly, sciCNV

profiling of copy number variations in individual cells from sample MM244, developed using

RNA-seq data, identified clonal gains within chromosome 7 and clonal losses within

chromosomes 1p, 10p, 13, 14q, 16q and 17p that were also identified by DNA sequencing of

pooled tumor bulk cells from the same patient; and additionally identified subclonal loss of the

distal arms of 1p and 2q, and subclonal gains of 11q and 19p (in subpopulations of cells) that

were also detected at subclonal CNV levels (0.15-0.50) by WES (**figure 4c**). Finally, sciCNV

identified clonal gains of 1q and 6p and clonal deletions within 1p, 2p, 6q, 13, 14q, 18 and 22 in

single cells from MM254 that were all also identified on tumor bulk DNA sequencing (**figure

4d**). Thus, these data demonstrate that the genomic CNV landscape of individual tumor cells can

largely be inferred from scRNA-seq transcriptome data by sciCNV, with results that show

reasonable correlation to DNA-based CNV profiling of pooled tumor bulk cells, although the

authors acknowledge that using RNA-based approaches small CNV regions may be missed.

Notably, several alternative methods exist for inferring CNVs from scRNA-seq[8,9,17,18]. Of these,

inferCNV[17] has been broadly used to calculate and visualize CNVs at single cell resolution.

Next, therefore, we compared our RTAM2-sciCNV pipeline with inferCNV, applying both

methods to the same set of normal and MM cells shown in figure 4 and comparing their single

cell outputs with the CNV profiles obtained from tumor bulk WES (**supplementary figures

S23-26**). As shown, sciCNV more reliably predicted tumor CNVs that were identified by WES,

and did so more consistently across individual cells. For example, for MM199 cells sciCNV

more readily detected gains of the distal arm of 1q, 3q, 7q, 9q and within 17q and 18q than

inferCNV (**supplementary figures S23**). At the same time sciCNV generally produced less false

positive CNV noise than inferCNV. Thus, for MM241 both methods detected subclonal gain of

1q and trisomies of 5,7 and 19, and deletion of 13 and 14q, but inferCNV also reported a number

of false positive deletions, particularly in chromosomes 2 and 4 (**supplementary figures S24)**.

Overall, therefore, whilst both methods provide CNV predictions that are substantially correct, in

general sciCNV offered a closer correlation with WES-based CNV findings.

Dissecting the effects of CNVs on gene expression within single cells: the transcriptional

consequences of +8q22-24 in MM

Simultaneous profiling of both DNA copy number and RNA in the same cell should enable

examination of the effect of a CNV on transcriptional programs. To test this, we used sciCNV to

study the effect of gain of the distal arm of 8q on cellular processes. We examined +8q as this is

one of the most recurrent abnormalities in human cancer[2,17] and is known to target MYC[18],

providing a benchmark result for testing our method.

Using sciCNV, sample MM199 was found to contain cells with subclonal gain of chromosome

8q22-24 (**figure 5a**). Importantly, MM199 also contained intra-clonal isogenic cells without +8q.

To facilitate gene set enrichment analyses (GSEA)[19] of these sibling cells, with and without +8q,

the subclones were next subsampled to obtain subpopulations of cells with matching

transcriptome depth (**figure 5b**). This eliminated biases in cell sequencing depths or in

transcriptome sizes between the subclones, and thus balanced the likelihood of gene-set detection

between subclones, at least in the absence of biological enrichment of specific genes. The gene

expression of the matched intra-clonal subpopulations, representing isogenic sibling cells with

and without +8q22-24, with equivalent transcriptome sizes, were then compared by GSEA using

RTAM2-normalized data. From an analysis of 215 cytoband gene sets (comprising genes defined

by chromosome location), +8q cells were strongly enriched for gene-sets located at 8q22-24,

with striking statistical confidence (p=0.000, q=0.000, FWER=0.000), compared to cells without

+8q (**figure 5d** and **supplemental figure S27).** In contrast, no other genomic regions were

significantly enriched in either subclone. Thus, sciCNV accurately resolved single MM cells into

intra-tumor subclones, isolating +8q22-24 as a unique variation distinguishing these.

We next used GSEA to explore the influence of +8q22-24 on cellular programming. Notably, the

primary MM cells from MM199 identified by sciCNV as having +8q22-24 showed increased

MYC expression (p<0.05) compared to sibling cells without +8q, as expected (**figure 5c**).

Furthermore, the +8q cells also showed broadly increased expression of hallmark MYC target

genes (p=0.000, q=0.000, FWER=0.000). Finally, from an analysis of 3303 curated gene sets,

primary MM199 cells with +8q22-24 also showed upregulation of gene-sets encoding the

machinery of mRNA translation and protein synthesis, including specifically genes involved in

3'UTR-mediated mRNA translation regulation (enrichment rank 5/3303), ribosome biogenesis

(rank 4/3303) and peptide chain elongation (rank 3/3303)(**figure 5d, supplemental figure S27**).

Conspicuously, these transcriptional effects of +8q22-24 in MM cells were remarkably close to

those of +8q23-24 in breast cancer (Nikolsky_Breast_Cancer_8q23-24 amplicon, FWER

p=0.000, enrichment rank 1/3303) (**figure 5d, supplemental figure S27**). Thus, overall our

analyses indicate that +8q22-24 in primary MM cells induces upregulation of MYC, MYC-target

genes and transcriptional programs that promote the upregulation of mRNA translation and

peptide synthesis. Analogous gene expression changes appear to occur across malignancies

including MM and breast cancer. Strikingly, we were able to identify these transcriptional

consequences of +8q22-24 using scRNA-seq analysis of a single MM patient sample. By

comparison, the Nikolsky_Breast_Cancer_8q23-24 amplicon signature was derived from the

combined genomic analyses of 191 breast cancer samples[20].

The cellular re-programming induced by +8q22-24 might be expected to promote an increase in

gene expression and in cell mass. Notably, however, in the MM sample examined the mTOR-

interacting gene, DEPTOR, encoded at 8q24, was also upregulated in +8q cells (**figure 5c**).

Upregulation of DEPTOR may limit cell size[21] and counter any tendency to increased cell mass

driven by MYC and ribosome biogenesis. Indeed, from our examination of +8q at a single cell

level we observed that the transcriptome sizes of +8q cells were in fact mildly reduced,

compared to sibling cells without the CNV (p<0.001) (**figure 5c**). Examination of cellular

transcriptome size is a capability unique to scRNA-seq but is generally not feasible from bulk

tumor gene expression studies. Thus, from single-cell analyses of +8q22-24 in primary MM cells

we find that copy number gain of this region acts to boost MYC, MYC target gene expression

and protein synthesis capacity (ribosomes, translation) without increasing cellular transcriptome

size. As the transcription of MYC-target genes is relatively increased, this likely leads to

enhanced expression of MYC-target genes as proteins. However, the increase in protein

synthesis capacity with reduced transcriptome size may also serve to improve the dynamics of

protein production by cells and may thus reduce the lag-time between gene expression and

proteome change, potentially enhancing cellular adaptability. This represents a unique

perspective on the function of +8q22-24 which is offered only by a multi-omic analysis of single

cells.

DISCUSSION

CNVs are critical drivers of cancer biology yet their specific effects on cellular processes often

remain only partially characterized. Here, we report bioinformatic methods for the dual profiling

of DNA copy number and RNA within the same cells, using scRNA-seq data, which enable

direct examination of the effect of intra-clonal CNVs on gene expression. We describe RTAM

normalization, which provides enhanced transcriptome alignment to improve gene expression

comparisons between cells, and sciCNV, which provides robust RNA-based CNV profiling at

single cell resolution. In order to demonstrate and validate these new tools, we examine the

transcriptional effects of copy number gain of chromosome region 8q22-24, which represents

one of the most common CNV in human cancer. From an examination of intra-clonal

subpopulations of cells with and without 8q gain we show using only a single MM patient

sample that +8q22-24 induces expression of MYC and MYC target genes, as expected,

demonstrating that intra-clonal CNVs can be linked to transcriptional consequences using only

scRNA-seq alone. This process can be compared to historical studies of cancer CNVs, in which

DNA and RNA profiling of hundreds of tumor samples must be performed to achieve similar

results. Moreover, single cell approaches potentially provide higher precision results due to the

ability to use intra-clonal isogenic cells as controls. Tumor registry studies, in contrast, are

encumbered by the immense genomic heterogeneity the exists between tumors, which persists

even when tumors are grouped into cohorts.

Although +8q22-24 upregulates the expression of a broad spectrum of MYC target genes, using

single cell profiling we demonstrate that the transcriptomes of MM cells with +8q are in fact

smaller than those of cells lacking +8q, at least in the sample examined by us, likely as a result of

DEPTOR overexpression from 8q24. At the same time, we find that +8q22-24 cells in primary

MM consistently upregulate gene sets involved in mRNA translation, ribosomal biogenesis and

peptide elongation. Thus +8q22-24 enhances protein synthesis capacity, without increasing

transcriptome size. This likely improves the dynamics of protein expression responses to gene

expression; and may enhance the malleability of MM cells with +8q24 to environmental

challenges.

Study of the effects of CNVs on cellular programming via single-cell transcriptomics offers a

number of advantages. As intra-clonal cells that diverge at a single CNV are virtually isogenic,

divergences in their gene expression can be precisely attributed to the CNV. Furthermore, as

cells with and without the CNV are derived from the same sample, differences in their gene

expression due to the microenvironment, clinical factors or sample processing are minimized. By

comparison, tumor cohort studies must instead rely upon the identification of correlations

between CNVs and gene expression across unrelated samples and suffer from the substantial

confounding genetic and clinical heterogeneity that exists between samples. As a result, the

effects of many cancer CNVs on cellular processes remain only partially characterized.

Defining the molecular mechanisms by which oncogenic chromosomal events influence cellular

behaviour is a critical step in understanding cancer pathogenesis. Indeed, it is essential

knowledge that underlies the development of therapeutic strategies to address the foundations of

malignant disease. Auspiciously, single cell multi omics offers the promise of a new era in

cancer genomics, in which the multi-faceted effects of CNVs on gene expression and cancer cell

programming can now be determined with precision.

METHODS

Samples

Bone marrow samples were collected from MM patients in heparinized tubes in compliance with

the Declaration of Helsinki and a protocol approved by the University Health Network Research

Ethics Board.

Cell isolation

Marrow cells were separated by Ficoll-Plaque (600g x 30 minutes) and treated with ACK red

blood cell lysis buffer (Sigma-Aldrich) for 2 minutes. Cells were then stained with fluorescently-

labelled antibodies (Biolegend) targeting CD2 (APC-cy7, RPA-2.10), CD3 (APC-cy7, HIT3a),

CD4 (APC-cy7, A161A1), CD11b (PEcy7, CBRM1/5), CD14 (PE-cy7, M5E2), CD34 (PE-

Dazzle, 581), CD235a (PE-cy7, H1264), CD38 (APC, HB-7), CD138 (FITC, MI15), BCMA

(PE, 19F2,) and CD20 (Pacific blue, 2H7) for 30 minutes in PBS 10% FBS 5mM EDTA at 4°C.

B cells and plasma cells were enriched on a MoFlo Astrio (Beckman Coulter) by excluding

duplets, non-viable cells (binding 7AAD) and cells expressing non-B lineage markers (CD2,

CD3, CD4, CD11b, CD124, CD235a, CD34), and by gating cells with lymphoid FSC-SSC

characteristics and expressing CD38, CD138 or CD20, as described[23] and illustrated in

Supplemental Fig. S29.

Single cell RNA-sequencing

Sequencing libraries were prepared from enriched cells using the 10X Genomics Chromium

Single Cell Library Kit chemistry v2 and Chromium Single Cell Controller; and were sequenced

on an Illumina HiSeq 2500 using paired-ends, targeting >50K mean reads per cell, >80%

sequencing saturation and phred-Q30 quality scores >80. Reads were de-multiplexed using

Illumina bcl2fastq, mapped to GRCh38 and were then consolidated into unique transcripts per

gene per cell using Cell Ranger 2.1.0 (10X Genomics). Poor quality cells with high

mitochondrial transcript content were excluded if this represented >5% of total cellular

transcripts or was >3 median absolute deviations from the median.

Non-RTAM scRNA-seq normalization

Normalization and data analysis were performed with R 3.6.1. scRNA-seq expression matrices

were normalized using RSEM's transcript per million (TPM) approach[15], SCRAN[11], SCONE[12]

or Suerat's (v3.1) SCTransform[16] function, employing *scran*, *scone* or *seurat V3.1* R packages.

To facilitate SCONE's quantile normalization, we limited the input matrix to only those genes

that had ≥1 read in ≥1 cell. When applying Seurat's SCTransform we used the 'corrected UMI'

count provided by this function. To compare the output data from each method on similar scales

we applied $\log_2(.+1)$ transformation as required.

RTAM scRNA-seq normalization

RTAM1: The normalized gene expression $GE_{i,j}$ of gene i in cell j was calculated from the raw

transcript count $T_{i,j}$, which was first normalized to the total transcript count per cell and then

scaled to a standard transcriptome library size (10^5) and log_2 transformed, mirroring the TPM

approach[15], yielding library-normalized values $L_{i,j}$:

364
$$L_{i,j} = log_2\left[\left(T_{i,j} \times \frac{10^5}{\sum_{i=o}^{i=Nj} T_{i,j}}\right) + 1\right]$$

N_j is the number of expressed genes (with ≥ 1 transcript) detected in cell j. Genes with non-zero

values were then assigned a rank in each cell $R_{i,j}$ according to descending $L_{i,j}$. Where multiple

genes had the same $L_{i,j}$ these were assigned an equal rank equivalent to the mean of the values

had these genes instead been assigned sequential ranks. M_j, the sum of $L_{i,j}$ of the top-ranked N_T

genes was calculated for each cell (from genes satisfying $1 \leq R_{i,j} \leq N_T$), where N_T was the number

of highly-expressed genes used to normalize the dataset.

372
$$M_j = \sum_{R_{i,j}=1}^{R_{i,j}=N_T} L_{i,j}$$

374 M_k, a target constant for correcting M_j values, was calculated as the mean of the initial M_j of all

375 cells. The adjustment variable, $h_{i,j}$, for gene i in cell j, was derived such that it adjusts the initial

376 M_j for each cell to the target M_k by differentially adjusting cellular gene expression on a log_2

377 scale according to $R_{i,j}$, and was calculated in the form:

378
$$h_{i,j} = C_j \cdot log_2\left[R_{i,j} + 1\right]$$

380 where C_j is a constant for each cell and equals:

382
$$C_j = \frac{M_k - M_j}{\left(N_T + \frac{3}{2}\right) \cdot log_2\left(N_T + \frac{3}{2}\right) - \left(\frac{N_T}{log_e 2}\right) - \frac{3}{2} \cdot log_2\left(\frac{3}{2}\right)}$$

Normalized $GE_{i,j}$ was calculated as: $GE_{i,j} = L_{i,j} + h_{i,j}$. Or overall,

$$GE_{i,j} = log_2\left[\left(T_{i,j} \times \frac{10^5}{\sum_{i=o}^{i=Nj} T_{i,j}}\right) + 1\right] + log_2[R_{i,j} + 1] \cdot \left[\frac{M_k - M_j}{\left(N_T + \frac{3}{2}\right) \cdot log_2\left(N_T + \frac{3}{2}\right) - \left(\frac{N_T}{log_e 2}\right) - \frac{3}{2} \cdot log_2\left(\frac{3}{2}\right)}\right]$$

RTAM2: N_j, $L_{i,j}$, $R_{i,j}$, M_j and M_k were calculated as for RTAM1. $G_{i,j}$ was defined as

$$G_{i,j} = T_{i,j} \times \frac{10^5}{\sum_{i=o}^{i=Nj} T_{i,j}}$$

and reflects the transcript count $T_{i,j}$ of gene i in cell j, corrected by scaling to a standard library

size (10^5). The RTAM2 adjustment variable, $h_{i,j}$, once again was derived to correct cellular M_j to

the constant M_k by adjusting gene expression on a non-linear log_2 scale. However, the RTAM2

adjustment differs from RTAM1 in that the per-gene adjustment is derived from the gene's

expression intensity rather than its expression rank and was calculated as:

$$h_{i,j} = \left(M_k - M_j\right) \cdot \frac{[G_{1,j} - G_{i,j} + 1]}{\sum_{p=1}^{p=N_T}[G_{1,j} - G_{p,j} + 1]}$$

RTAM2 normalized gene expression, $GE_{i,j}$, was calculated as: $GE_{i,j} = L_{i,j} + h_{i,j}$.

Or,

$$GE_{i,j} = log_2[G_{i,j} + 1] + \left(M_k - M_j\right) \cdot \frac{[G_{1,j} - G_{i,j} + 1]}{\sum_{p=1}^{p=N_T}[G_{1,j} - G_{p,j} + 1]}$$

400
The derivation of RTAM1 and RTAM2 functions are described in the supplemental materials.

sciCNV

Single-cell CNV were inferred from the RNA counts of test cells by comparing these normal

control cells of the same lineage. In brief, test and control cells were co-normalized using

RTAM2. The combined $GE_{i,j}$ matrix was then truncated to informative genes (N_{CNV}) by

excluding genes expressed in <2% cells; and was re-sorted by gene location. *i was* redefined by

the rank location of informative genes, numbered from chromosomes 1→Y, p-arm→q-arm, $i =$

1, 2, 3, ⋯, N_{CNV}. To mitigate against data sparsity within individual cells, a moving average of

gene expression (maGE_i) was calculated across chromosome segments extending $\pm n/2$ genes

from gene i, where $n = N_{CNV}/(S \times 50)$. The variable S defined the sharpness (default=1.0) of the

analysis resolution and was adjustable to offset sample data sparsity. Data from control cells was

merged into a single '*average cell*' with the expression of each gene calculated from its average

across all control cells, including zero values, thus integrating the probability of gene detection.

A moving average of gene expression across chromosome segments extending $\pm n/2$ genes from

gene i was calculated for the merged control cell, ma$GE_{i,control}$, mirroring that of test cells. A

weighted disparity score, $W_{i,j}$, comparing gene expression in test cell j and the merged control

cells was calculated as:

420
$$W_{i,j} = \frac{maGE_{i,j} - maGE_{i,control}}{maGE_{i,j} + maGE_{i,control} + \alpha}$$

422   To avoid division by zero, $\alpha = 0.00001$. $W_{i,j}$ was centred on zero in each cell by subtracting its

423   median, and then linearized by the equation:

425
$$W_{i,j}^{lin} = 3.3222(W_{i,j})^4 + 5.6399(W_{i,j})^3 + 4.2189(W_{i,j})^2 + 3.8956W_{i,j}$$

$W_{i,j}^{lin}$ provided a first-pass estimate of single cell CNVs. To further enhance CNV detection, we

deployed a second method of assessing gene expression disparity that was sensitive to small

concordant expression variations in contiguous genes and insensitive to large single-gene

variations. A moving average (*ma*) of gene expression, $U_{i,j}$, across chromosome segments was

calculated for both test and control cells using the $W_{i,j}$ function above but employing a smaller

window extending from $i \pm n/(5 \times S)$ genes. $U_{i,j}$ was centered on zero in each cell by subtracting

its median and then positive/negative values were digitized into +1/-1 values respectively. The

digital values were summed cumulatively across the genome as a function of i, generating a

digital moving score (*DMS*).

437
$$maDMS_{i,j} = \begin{cases} maDMS_{i-1,j} + 1, & U_{i,j} > 0, \\ maDMS_{i-1,j} - 1, & U_{i,j} < 0, \\ maDMS_{i-1,j}, & U_{i,j} = 0. \end{cases}$$

439   A second digital moving score, $V_{i,j}$, was calculated as for $U_{i,j}$ but employed gene expression

440   values without average smoothing in order to obtain enhanced resolution for small CNV. The

441   non-moving average DMS (*nmaDMS*) was then calculated as:

443
$$nmaDMS_{i,j} = \begin{cases} nmaDMS_{i-1,j} + 1 - d_j, & V_{i,j} > 0, \\ nmaDMS_{i-1,j} - 1 - d_j, & V_{i,j} < 0, \\ nmaDMS_{i-1,j} - d_j, & V_{i,j} = 0. \end{cases}$$

445   The *nmaDMS* function in each cell j was adjusted to sum cumulatively to zero at the last gene

446   (it's distal terminus) by cumulatively subtracting d_j at each gene, where:

448
$$d_j = \frac{\left|\{V_{i,j} : V_{i,j} > 0\}\right| - \left|\{V_{i,j} : V_{i,j} < 0\}\right|}{|N_{CNV}|}$$

nma*DMS* was scaled to ma*DMS* by the ratio of their respective ranges, (*maDMS_{MAX}* -

maDMS_{MIN}) / (nma*DMS_{MAX}* − nma*DMS_{MIN}*). maDMS and nmaDMS were then combined by

weighted average to provide the merged function, $DMS_{i,j}$, integrating both smooth low-resolution

and noisy high-resolution data. The gradient of this function, $dDMS_{i,j}/di$, measured over

chromosome segments $i \pm n/2$, provided a second measure of CNV.

The $W_{i,j}^{lin}$ and $dDMS_{i,j}$ CNV methods were combined by geometric mean. When the two methods

yielded conflicting polarity results, $sciCNV_{i,j} \rightarrow 0$. Using the mean $sciCNV_{i,j}$ result from all

clonal test cells a scale factor, s, was manually derived to ensure that CNV values optimally

centered on integers. For example, for MM samples, s was calculated so that clone mean sciCNV

for chromosome 13 deletion scaled to -1. s was then uniformly applied to all cells:

462
$$sciCNV_{i,j} = s. \sqrt[2]{W_{i,j}^{lin}.dDMS_{i,j}}$$

Heatmaps were generated using heatmap.3 R package. Cells were clustered iteratively to identify

subclones, using gene expression-based tSNE clustering and sciCNV-based Pearson correlation

coefficients. After subclones were identified, supervised clustering was applied. Stochastic noise

was reduced by standardizing each sciCNV profile against the median of its nearest neighbours.

To further reduce low-level sciCNV noise centered around zero, $|sciCNV_{i,j}|$ values less than

threshold, T, were set to zero. For tumors in which chromosome gains and losses were

substantially imbalanced (with respect to the numbers of genes involved) an optional sciCNV

baseline correction, b, was applied to correct the zero-set point and improve CNV detection. The

method is described further in the **supplemental materials**.

Whole exome sequencing

DNA was isolated from tumor plasma cells using a Micro kit (Qiagen) and then sheared into

150-200bp fragments using a Covaris E220. Exome libraries were prepared using a Hyper Prep

Kit (KAPA Biosystems), pre-capture PCR and SureSelectXT Human All Exon V5+UTR baits

(Agilent). Post-capture fragments were amplified by PCR, purified using AMPure-XP beads

(Beckman Coulter Genomics) and sequenced using paired-ends on an Illumina HiSeq. FASTQ

files were generated with Illumina bcl2fastq and reads were aligned to GRCh38 using BWA

v0.7.8[24] and SAMTOOLS v1.2[22].

Correlation of sciCNV and WES-derived CNV

MM199 scRNA-seq data was RTAM2 normalized and sciCNV profiles were calculated (S=1.4,

b=0.1) without a noise filter. MM199 WES was performed using 200ng DNA (>30,000

equivalent exomes) extracted from 1.9×10^6 pooled cells. WES-derived CNVs were identified

using DNAcopy R package v1.54.0. CNV estimates from the two methods were smoothed using

windows of identical genomic size and the datasets were paired by nearest genomic location.

Fluorescence in-situ hybridization (FISH)

Cells on cytospin slides were hybridized with Vysis LSI IGH/CCND1 (14q32:11q13),

IGH/FGFR3(14q32:4p16), D5S23, D5S721 SGn/CEP9 SA/CEP15 probes (Abbott Molecular) or

Cytocell Aquarius CKS1B/CDKN2C probes according to the manufacturer's instructions. At

least 100 cells were scored for each.

Tumor CNV score

For each sample, we identified cells with clonal CNVs by sciCNV and using these derived a tumor-specific average sciCNV profile. The sciCNV profiles of all individual cells were then assessed against the tumor average. The tumor score (TS) for cell j was summed cumulatively across the genome as:

$$TS_j = \sum_{i=1}^{|\mathcal{G}|} TS_{i,j}$$

where $\mathcal{G}$ is the set of all genes in sciCNV matrix and

$$TS_{i,j} = \begin{cases} +\sqrt{sciCNV_{i,j}}, & sciCNV_{i,j} \times \mathbb{E}[sciCNV_i] > 0, \\ -\sqrt{sciCNV_{i,j}}, & sciCNV_{i,j} \times \mathbb{E}[sciCNV_i] > 0, \\ -\sqrt{sciCNV_{i,j}}, & \mathbb{E}[sciCNV_i] = 0. \end{cases}$$

$\mathbb{E}[sciCNV_i]$ is the average tumor cell $sciCNV_i$ for gene i.

Immunoglobulin isotype score

To evaluate if plasma cells belonged to the tumor clone, we scored their immunoglobulin heavy and light chain (IGH, IGK, IGL) gene expression for restriction to the tumor isotype. Using RTAM2 data, a threshold was identified for IGHM, IGHD, IGHG1/2, IGHA1/2, IGHE and for IGKC and IGL(C2-7) which segregated non-expressing and expressing cells. The immunoglobulin isotype score was calculated for each cell by summing the supra-threshold expression of tumor immunoglobulin genes and subtracting the supra-threshold expression of non-tumor immunoglobulin genes.

Gene-set enrichment analysis

Intra-clonal cells with divergent CNV were segregated by sciCNV. Paired subpopulations of cells that diverged at a single CNV were then subsampled for cells that were matched for normalized cellular transcriptome size. The matched cells were then compared using gene-set enrichment analysis (GSEA v3.0)[19] of RTAM2-normalized transcriptomes.

DATA & CODE AVAILABILITY

522   The raw and processed scRNA-seq and WES data are deposited at NCBI GEO, GSE141299. The

523   microarray data (Affymetrix U133Plus2.0) for MM tumors with 1q-FISH status was obtained

524   from the NCBI GEO, GSE2658[22]. The code used in this study is available at

525   https://www.github.com/TiedemannLab/sciCNV.

REFERENCES

1 Beroukhim, R. *et al.* The landscape of somatic copy-number alteration across human
cancers. *Nature* **463**, 899-905, doi:10.1038/nature08822 (2010).
2 Tate, J. G. *et al.* COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids*
Res **47**, D941-D947, doi:10.1093/nar/gky1015 (2019).
3 Taylor, A. M. *et al.* Genomic and Functional Approaches to Understanding Cancer
Aneuploidy. *Cancer Cell* **33**, 676-689 e673, doi:10.1016/j.ccell.2018.03.007 (2018).
4 Dey, S. S., Kester, L., Spanjaard, B., Bienko, M. & van Oudenaarden, A. Integrated
genome and transcriptome sequencing of the same cell. *Nat Biotechnol* **33**, 285-289,
doi:10.1038/nbt.3129 (2015).
5 Han, K. Y. *et al.* SIDR: simultaneous isolation and parallel sequencing of genomic DNA
and total RNA from single cells. *Genome Res* **28**, 75-87, doi:10.1101/gr.223263.117
(2018).
6 Macaulay, I. C. *et al.* G&T-seq: parallel sequencing of single-cell genomes and
transcriptomes. *Nat Methods* **12**, 519-522, doi:10.1038/nmeth.3370 (2015).
7 Fan, J. *et al.* Linking transcriptional and genetic tumor heterogeneity through allele
analysis of single-cell RNA-seq data. *Genome Res* **28**, 1217-1227,
doi:10.1101/gr.228080.117 (2018).
8 Patel, A. P. *et al.* Single-cell RNA-seq highlights intratumoral heterogeneity in primary
glioblastoma. *Science* **344**, 1396-1401, doi:10.1126/science.1254257 (2014).
9 Tirosh, I. *et al.* Dissecting the multicellular ecosystem of metastatic melanoma by single-
cell RNA-seq. *Science* **352**, 189-196, doi:10.1126/science.aad0501 (2016).
10 Vallejos, C. A., Risso, D., Scialdone, A., Dudoit, S. & Marioni, J. C. Normalizing single-cell
RNA sequencing data: challenges and opportunities. *Nat Methods* **14**, 565-571,
doi:10.1038/nmeth.4292 (2017).
11 Lun, A. T., Bach, K. & Marioni, J. C. Pooling across cells to normalize single-cell RNA
sequencing data with many zero counts. *Genome Biol* **17**, 75, doi:10.1186/s13059-016-
0947-7 (2016).
12 Cole, M. B. *et al.* Performance Assessment and Selection of Normalization Procedures
for Single-Cell RNA-Seq. *bioRxiv*, doi:doi: https://doi.org/10.1101/235382 (2017).
13 Stegle, O., Teichmann, S. A. & Marioni, J. C. Computational and analytical challenges in
single-cell transcriptomics. *Nat Rev Genet* **16**, 133-145, doi:10.1038/nrg3833 (2015).
14 Vieth, B., Parekh, S., Ziegenhain, C., Enard, W. & Hellmann, I. A Systematic Evaluation of
Single Cell RNA-Seq Analysis Pipelines: Library preparation and normalisation methods
have the biggest impact on the performance of scRNA-seq studies. *bioRxiv*, 583013,
doi:10.1101/583013 (2019).
15 Li, B., Ruotti, V., Stewart, R. M., Thomson, J. A. & Dewey, C. N. RNA-Seq gene expression
estimation with read mapping uncertainty. *Bioinformatics* **26**, 493-500,
doi:10.1093/bioinformatics/btp692 (2010).

16   Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-
566   seq data using regularized negative binomial regression. *Genome Biol* **20**, 296,
567   doi:10.1186/s13059-019-1874-1 (2019).
568   17   Davoli, T., Uno, H., Wooten, E. C. & Elledge, S. J. Tumor aneuploidy correlates with
569   markers of immune evasion and with reduced response to immunotherapy. *Science* **355**,
570   doi:10.1126/science.aaf8399 (2017).
571   18   Misund, K. *et al.* MYC dysregulation in the progression of multiple myeloma. *Leukemia*,
572   doi:10.1038/s41375-019-0543-4 (2019).
573   19   Mootha, V. K. *et al.* PGC-1alpha-responsive genes involved in oxidative phosphorylation
574   are coordinately downregulated in human diabetes. *Nat Genet* **34**, 267-273,
575   doi:10.1038/ng1180 (2003).
576   20   Nikolsky, Y. *et al.* Genome-wide functional synergy between amplified and mutated
577   genes in human breast cancer. *Cancer Res* **68**, 9532-9540, doi:10.1158/0008-5472.CAN-
578   08-3082 (2008).
579   21   Peterson, T. R. *et al.* DEPTOR is an mTOR inhibitor frequently overexpressed in multiple
580   myeloma cells and required for their survival. *Cell* **137**, 873-886,
581   doi:10.1016/j.cell.2009.03.046 (2009).
582   22   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
583   2078-2079, doi:10.1093/bioinformatics/btp352 (2009).

SUPPLEMENTARY INFORMATION:

Supplementary methods, notes and figures can be found on-line.

ACKNOWLEDGEMENTS:

The authors thank the patients and physicians who made this study possible. They also thank N.

Winegarden, N. Khuu and G. Basi in the Princess Margaret Genomics Facility and Z. Lu in the

Princess Margaret Bioinformatics Core for technical assistance; and Drs. Gary Bader and Caleb

Stein for independent review and comments. This work was supported by funding from The

Princess Margaret Cancer Centre Foundation, the Terry Fox Foundation and the Canadian

Cancer Society Research Institute.

Author Contribution

A.M-S performed research and analyzed data. N.E., C.L-H. FISH, FACS and whole exome

sequencing, respectively. R.E.T. designed research, analyzed data and wrote the paper.

Competing interests

The authors declare that they have no competing interests.

Figure 1. Comparison scRNA-seq normalization strategies, including RTAM1 and RTAM2 methods.

a. Overview of workflow showing bone marrow sampling, FACS sorting of B cell lineage cells, scRNA-seq and data normalization. MM, multiple myeloma. FACS, fluorescence activated cell sorting.

b. Plot of scRNA-seq data from >6,000 B cells and plasma cells, isolated from the bone marrow

of a MM patient, depicting the raw (pre-normalized) transcript counts per gene per cell. Each dot

represents an integer transcript count for one or more genes in a single cell; cells (columns) are

ranked from left to right by their total transcript count.

c. The same data is shown following normalization using TPM, SCRAN, SCONE, SCTransform,

RTAM1 or RTAM2 methods (and following log transformation). To compare the methods, the

mean expression of a curated set of house-keeping genes (HKG) is plotted in each cell as a blue

dot.

See also **supplemental figures S4-S20**.

Figure 2. RTAM scRNA-seq normalization reduces cell to cell variation in average

expression of house-keeping genes (HKG) and ubiquitously-expressed genes (UEG).

Alignment of single cell transcriptomes is compared using 6 different methods for scRNA-seq

normalization.

a. The coefficient of variation (CV) across cells in the average expression of HKGs within each

cell is shown for 3 scRNA-seq samples (containing >15,000 cells), normalized by 6 different

methods. To avoid bias in the analysis, average expression was alternatively calculated as either

the mean of the detected HKG (top panel), the mean of all HKG [with imputation of null dropout

values] (bottom left panel), or the median of all HKG without imputation (right panel). As the

various normalization methods expand or compress the distribution of the overall gene

expression data to different extents, the CV of HKG averages is plotted against the CV of

expression of all genes.

b. The coefficient of variation (CV) for all genes ubiquitously expressed (UEGs)(in >95% cells)

was determined for the same patient samples. Once again, the average expression of UEG in

each cell was alternatively calculated as either the mean of the detected UEG (top panel), the

mean of all UEG imputing null dropouts (bottom left panel), or the median of all UEG without

imputation (right panel).

A lower CV in the mean expression of HKG/UEG implies better alignment of average gene

expression across cells. As the normalization methods expand or compress the distribution of the

overall gene expression data to different extents, the CV of HKG/UEG averages (y-axis) is

plotted against the CV of expression of all genes (x-axis).

Figure 3. sciCNV can predict DNA copy number variations (CNV) in single cells from scRNA-seq data.

a. The CNV profiles of 1,625 pooled tumor cells (middle panel) or of a single tumor cell (lower panel) from sample MM199 were inferred from scRNA-seq using RTAM2/sciCNV and are shown compared with the bulk tumor CNV profile (upper panel), which was obtained by whole exome sequencing (WES) of DNA purified from 1.9×10^6 cells. Cells were isolated from bone marrow by FACS.

b. The scRNA-seq-based sciCNV profiles from (**a**) are correlated with the tumor bulk CNV profile derived from WES.

48 **c.** FISH verification of sciCNV copy number predictions, for the genes highlighted in **a.**,

49 showing 3 copies of CKS1B (1q21), 1 copy of FGFR3 (4p16), 1 copy of SEC63 (6q21), 2 copies

50 of PNOC (8p21), 3 copies of MYC (8q24), 1 copy of RB1 (13q14,) and 1 copy of TP53 (17P13),

51 as predicted. Brightness and contrast were adjusted during figure construction to enhance probe

52 visualization.

Figure 4. Using scRNA-seq data, sciCNV consistently infers single cell CNVs across

thousands of cells that correlate with bulk tumor DNA-based CNV profiling

a.-d. The CNV profiles of thousands of individual plasma cells from MM patient samples

MM199 (n=1724)(a), MM241(n=2806) (b), MM244 (n=3569)(c) and MM254 (n=649)(d) were

inferred from scRNA-seq using RTAM2/sciCNV (lower panels) and are shown compared to the

bulk tumor CNV profiles generated by DNA whole exome sequencing (upper panels). The

sciCNV profiles of normal plasma cells (NPCs)(n=205) are shown as additional controls. For the

61   heatmaps revealing CNV profiles, at single cell resolution, the tumor cells in each sample are

62   grouped into inferred subclones (color bars at left) distinguished by divergent CNV.

Figure 5. The transcriptional effects of +8q22-24 in MM cells, revealed by dual DNA CNV and RNA profiling of single cells using scRNA-seq and sciCNV.

a. Heatmaps showing the sciCNV profiles of MM199 cells, showing intra-clonal subpopulations (subclones) that diverge at +8q.

b. The transcriptome size distribution for cells in each subclones is shown. The original subclones (left) were sampled for subpopulations of cells with comparable transcriptome sizes (right panel), which were then compared in subsequent analyses.

c. Plots showing the mRNA expression of MYC or DEPTOR genes, located on chromosome 8q24, in MM199 cells by +8q status. Expression data is compressed by normalization (RTAM2) and log scale. The pre-normalized transcriptome sizes of subclonal MM cells from MM199 are also shown, demonstrating fewer RNA transcripts in cells with +8q. P-values were calculated by t-test.

d. Results of gene set enrichment analysis (GSEA) performed on MM199 intra-clonal subpopulations, comparing cells with or without +8q. GSEA of chromosome position gene sets (n=215) showed enrichment of chromosome 8q22-24 gene sets in cells identified by sciCNV at single cell resolution as containing +8q (left panels). Key results from GSEA of hallmark (n=49) and curated (n=3303) gene sets are shown in the middle and right panels; demonstrating upregulation of MYC target genes in MM199 +8q cells, and enriched expression of the gene sets: Nikolsky_Breast_Cancer_8Q23-24_Amplicon, KEGG_Ribosome, Reactome_Peptide Elongation and Reactome_3'_UTR_mediated translation_regulation.