1 **Modeling How Community Assembly Alters the**

2 **Functioning of Ecosystems**

4 *Authors and Affiliations*

6 Koffel, Thomas, W.K. Kellogg Biological Station, Michigan State University, 3700 E
7 Gull Lake Dr., Hickory Corners, MI 49060. U.S.A. koffelth@msu.edu
8
9 Kremer, Colin T, W.K. Kellogg Biological Station, Michigan State University, 3700 E
10 Gull Lake Dr., Hickory Corners, MI 49060. U.S.A. kremerco@msu.edu
11
12 Bannar-Martin, K. kbannarm@gmail.com
13
14 Ernest, S.K. Morgan, Department of Wildlife Ecology and Conservation, University of
15 Florida, Gainesville, FL 32611, USA.
16
17 Eisenhauer, Nico, [1] German Centre for Integrative Biodiversity Research (iDiv)
18 Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany; [2] Institute of
19 Biology, Leipzig University, Deutscher Platz 5e, 04103 Leipzig, Germany.
20 nico.eisenhauer@idiv.de
21
22 Roscher, Christiane, [1] Helmholtz Centre for Environmental Research (UFZ),
23 Department Physiological Diversity, Permoserstrasse 15, 04318 Leipzig, Germany; [2]
24 German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig,
25 Deutscher Platz 5e, 04103 Leipzig, Germany. christiane.roscher@ufz.de
26
27
28 Cabral, Juliano Sarmento, Ecosystem Modeling, Center for Computational and
29 Theoretical Biology (CCTB), University of Würzburg, Emil-Fischer-Str. 32, 97074
30 Würzburg, Germany. juliano.sarmento_cabral@uni-wuerzburg.de
31
32
Leibold, Mathew A., Department of Biology, University of Florida, Gainesville, FL
32611, USA. mleibold@ufl.edu

Abstract:

Although the effects of species richness on ecosystem functioning have been

extensively studied, there is increased interest in understanding how community

assembly in a more general sense can alter the functioning of ecosystems. We

identify two complementary approaches for evaluating how community assembly

influences biomass production. The first quantifies the relative importance of

complementarity and selection by contrasting monocultures with polycultures. The

second identifies the effects of species replacements (losses and/or gains of species

relative to the baseline polyculture) as well as the indirect effects on other species'

productivity. We compare and contrast these two approaches using simulated

communities where species compete for different resource types and experience

different levels of environmental heterogeneity and find that these metrics can

jointly diagnose the mechanisms of competition driving productivity. We apply our

methods to data from a long-term biodiversity-ecosystem experiment (the Jena

Experiment) and find that the data do not correspond to any of the competition

scenarios we modeled. We address two additional possible complications:

facilitation by nitrogen fixing plants, and non-equilibrium behavior during

community assembly, and find that facilitation/competition by nitrogen fixing

plants is the more likely explanation for the results obtained at Jena.

 Introduction:

Community assembly drives shifts in community structure (i.e., species richness and
composition) which can have important consequences for the functioning of
ecosystems. While the impact of changes in species richness on ecosystem function
(commonly called biodiversity-ecosystem function, or BEF, effects; reviewed by
Cardinale et al. 2012, Hooper et al. 2012, Tilman et al. 2012) are the best-studied,
the importance of all of community assembly's impacts on community structure
(referred to here as community assembly-functioning of ecosystem, or CAFE,
effects) is increasingly recognized (e.g. Ptacnik et al. 2010, Thompson and Gonzalez
2016, Hodapp et al. 2016, Leibold et al. 2017, Bannar-Martin et al. 2018). Drivers of
community assembly include species interactions, disturbance, dispersal,
environmental heterogeneity, and landscape effects that can lead to changes in
community composition, without necessarily affecting species richness (Brown et al.
2001). Community assembly is thus complex and depends on multiple factors and
processes that may also consequently affect ecosystem function (e.g. Thompson and
Gonzalez 2016, Hodapp et al. 2016, Leibold et al. 2017). Therefore, it is community
assembly processes as a whole that ultimately link community composition,
including species richness, to ecosystem attributes.

How can we study these community assembly effects on ecosystem function? One
approach would be to comprehensively study the detailed interactions among the
species involved and theoretically analyze these interactions to fully disentangle
how community assembly affects ecosystem attributes. This approach, however, is
logistically demanding and likely to give answers that are system specific. An
alternative is to find general metrics that parse the effects of various components of
community assembly on ecosystem attributes. Several methods are available (e.g.
Kirwan et al. 2009, Bell et al. 2009, Hector et al. 2009, Bannar-Martin et al. 2018); in
this paper we focus on two of these. The first, which we call the Complementarity-
Selection Effects method (originally called the 'additive partitioning method'), is
based on comparing the contributions of species to ecosystem function when they

are in monocultures to those that emerge due to species interactions in polycultures

(Loreau and Hector 2001). Such comparisons allow total ecosystem functioning in a

community to be partitioned into the function that results from selection effects

(when species that are most productive in monocultures come to dominate

multispecies assemblages and their functioning) from what is due to

complementarity effects (when productivity in multispecies communities is higher

than expected from proportional monoculture contributions due to more complex

species interactions such as niche partitioning or facilitation). Both of these effects

can be positive (e.g. when selection favors the more productive monocultures or

when complementarity results in enhanced production relative to that expected or

when there is facilitation among species) or negative (when the reverse is true). It

is generally thought that selection effects occur due to environmental filtering which

selects for more productive species that are better suited to the local environment,

whereas complementarity is related to niche partitioning that results in enhanced

coexistence and increased resource-use efficiency. This approach is carefully

tailored to the way most BEF experiments are actually conducted because BEF

experiments require data on monocultures for all the component species and they

use null models to compare final outcomes of polycultures to these monocultures

instead of comparing them to other multispecies communities that may

alternatively occur.

Another approach for linking community assembly effects with ecosystem function

is based on pairwise comparisons of communities and quantifies how differences in

total function relate to changes in composition and individual species' functions by

calculating 'Price components' (Fox 2006, Fox and Kerr 2012, Bannar-Martin et al.

2018). Although this approach is constrained to ecosystem functions that can be

calculated as the sum of contributions of their constituent species (Fox 2006), it has

the advantage that it can be more generally used to look how any change in

community structure (i.e., species richness, taxonomic turnover, changes in resident

abundances or function) alters ecosystem functioning. Calculating Price

components can be used flexibly to make inferences based on complex pairwise

comparisons (Bannar-Martin et al. 2018) or on temporal effects related to

environmental change (Kremer et al. in prep). Here we will focus on comparisons

between the communities with the most diverse species pool and those with lower

levels of richness. There are many different ways the Price components can be

configured (Bannar-Martin et al. 2018), but we employ a three-part arrangement

that separately evaluate the functional effects of species gains ('SG'), species losses

('SL') and the differences in density/function among species present in both

communities being compared (the Context Dependent Effect or 'CDE'). In Bannar-

Martin et al. (2018) these are called the Community Assembly components; we use

this terminology in the rest of this paper.

The Complementarity-Selection Effects (Loreau and Hector 2001) and Community

Assembly components methods (Fox and Kerr 2012, Bannar-Martin et al. 2018)

provide complementary information about the impact of community assembly on

ecosystem function. The first focuses on mechanisms related to environmental

filtering and niche relations that affect dominance and coexistence, whereas the

second focuses on critical aspects of community assembly involving the loss and

gain of species as well as the consequent effects on species that remain in the

community. We study how the two approaches might be combined and compared to

allow us to understand underlying mechanisms by applying these methods to a

community model we constructed based on resource-competition theory (Tilman

1982, Tilman et al. 1997, Schreiber and Tobiason 2003, Gross and Cardinale 2007,

Hodapp et al. 2016, Koffel et al. 2016). To model the community assembly process,

we viewed a BEF experiment as having the following sequence of events: a) there is

a filter on the species pool (taken from the total species pool of the experiment), that

is imposed by the researchers via selective seeding and/or weeding, b) interactions

among the remaining species lead to some of the species going extinct, and c) there

are density-dependent responses among the remaining species that then determine

ecosystem attributes. We then used this mathematical model to solve the assembly

of communities from all possible subsets of a given total species pool, imitating a

standard BEF experiment. We applied the model to various assembly scenarios to

consider a wide range of possible ways resource competition may operate to affect

community assembly in a BEF experiment - including competition for

complementary vs. antagonistic resources with homogeneous or heterogeneous

resource supply at the local scale. We apply the Complementarity-Selection Effects

method (Loreau and Hector 2001) and the Community Assembly components

method (Bannar-Martin et al. 2018) to the results of the models to assess if these

methods can be used to distinguish among the various assembly scenarios. Finally,

we compare our theoretical findings to analyses of data from the long term BEF

experiment conducted in grasslands at Jena (Roscher et al. 2004, 2005; Weigelt et al.

2010) and find that the data do not conform to any of the scenarios we initially

modelled. We consequently modelled two other scenarios that we hypothesized

might be important, inadequate time to achieve steady-state competitive outcomes

and facilitation by nitrogen fixing plants to illustrate how we can improve our

understanding of how community assembly affects the functioning of ecosystems

and find that the data conform to results expected if facilitation is important but not

to lack of steady-state competition outcomes.

<u>Methods:</u>

To model community assembly in BEF experiments, we used basic resource

competition models for two resources as initially developed by MacArthur (1972)

and further elaborated by Tilman (1982):

$$\frac{dB_j}{dt} = [g_j(R_1, R_2) - m_j]B_j$$

171 (1)

$$\frac{dR_i}{dt} = l_i(S_i - R_i) - \sum_{j=1}^{n} C_{ij}(R_1, R_2)B_j$$

This model describes the population dynamics of n species j (whose biomasses are

denoted B_j) that compete for 2 resources, R_1 and R_2. Here $g_j(R_1, R_2)$ and m_j

respectively determine species j's per capita growth and mortality rates, S_i is the

supply of resource R_i, which is diluted at rate l_i, and $C_{ij}(R_1, R_2)$ is the per capita

consumptive impact of B_j on R_i. Species j's interactions with the two resources can

be divided in two parts: (1) its requirements (through the per capita growth rate g_j)

and; (2) its impacts on the resources (through the per capita consumptive impacts

C_{ij}).

Competitive Scenarios: We considered a range of competitive scenarios, classified

by the types of resources involved, including resources that are 'strongly

complementary' (virtually 'essential', *sensu* Tilman 1982) vs. 'weakly

complementary' and 'weakly antagonistic' (these two cases correspond closely to

'substitutable' in Tilman's terminology, the perfectly substitutable model has

undesirable mathematical properties for our purpose) vs. 'strongly antagonistic'. To

implement this variation in resource types we used a flexible form of the $g_j(R_1, R_2)$

function developed by Schreiber and Tobiason (2003):

$$g_j(R_1, R_2) = [(x_{1j}R_1)^\alpha + (x_{2j}R_2)^\alpha]^{\frac{1}{\alpha}} \qquad (2)$$

Here α is a tunable shape parameter that determines the type of resource (i.e.

complementary or antagonistic). The traits x_{1j} and x_{2j} are species j's acquisition

rates for R_1 and R_2, and control how efficient species j is at acquiring resources.

Thus we modelled the four different resource scenarios by picking four different

values for the shape parameter α : strongly complementary ($\alpha \ll 1$), strongly

antagonistic ($\alpha \gg 1$), weakly complementary ($\alpha \lesssim 1$) and weakly antagonistic

($\alpha \gtrsim 1$).

To fully specify the model we also had to tune species' consumptive impacts on their

resources, $\boldsymbol{C_{ij}(R_1, R_2)}$. To do so we followed Koffel et al. (2016) and model per

capita consumptive impacts separately for each resource scenario to better match

the biology of resource uptake.

For strongly complementary resources ($\alpha \ll 1$), we assume that both resources are
necessary to grow, which means that growth translates into proportional uptakes of
both resources through fixed stoichiometric coefficients q_{ij}:

$$C_{ij}(R_1, R_2) = q_{ij}\, g_j(R_1, R_2) \tag{3}$$

This is consistent with the way essential resources are usually modeled (Tilman
1982). We further assume that $q_{1j} = x_{2j}$ and $q_{2j} = x_{1j}$, which ensures that every
species has a greater impact on the resource that most limits its growth, a standard
necessary condition for coexistence to happen in resource competition theory
(Tilman 1982, Leibold 1995, Chase and Leibold 2003).

For strongly antagonistic resources ($\alpha \gg 1$), resources can be substituted for each
other, but a mixed uptake does not translate additively into growth because of the
antagonism between the two resources. Such antagonism can for example happen
when the two resources are patchily distributed. In this scenario, the consumers
focus their uptake on the most profitable resource (based on its abundance) and
avoid the least profitable (often rarest) one. Mixed uptake only occurs for a narrow
range of availabilities corresponding to equally profitable resources, where the
consumer is about to switch from consuming one resource to consuming the other.
Switching uptake can be modelled by the following mathematical expression:

$$C_{ij}(R_1, R_2) = \frac{(x_{ij}R_i)^\alpha}{(x_{1j}R_1)^\alpha + (x_{2j}R_2)^\alpha}\, g_j(R_1, R_2) \tag{4}$$

This choice of consumptive impact is consistent with mass balance of substitutable
resources, where all the resources taken up translate into growth by an equivalent
amount of biomass. This can formally be checked by summing the two uptake rates
of a species j given by Equation (4) and retrieving its growth rate ($g_j = C_{1j} + C_{2j}$).

Finally, in the weakly complementary and weakly antagonistic resource scenarios
($\alpha \lesssim 1$ and $\alpha \gtrsim 1$, respectively), resources are close to being consumed
proportionally to their availability, i.e $C_{ij}(R_1, R_2) \approx x_{ij}R_i$, consistent with their
close proximity to the perfectly substitutable case ($\alpha = 1$; Schreiber and Tobiason
2003). Such a close-to-proportional consumptive impact is obtained by borrowing

equation (4) from the antagonistic resource scenario and using it for these two

weakly interacting scenarios. Indeed, equation (4) tends towards linear

consumption when α tends towards 1.

Fig. 1 illustrates how each of these resource types affects trade-off relations among

minimum resource requirements for the two resources. Previous work has almost

exclusively focused on the strongly complementary scenario (e.g. Tilman et al. 1997,

Gross and Cardinale 2007, Hodapp et al. 2016); however, we found that different

types of competition lead to different outcomes of community assembly, associated

with distinctive signatures in terms of Complementarity-Selection Effects and

Community Assembly components (see Box 1 and `Results' section).

Figure 1. ZNGIs (colored, thick) and supply points (black dots) for each of the eight
resource scenarios. ZNGIs are curved upwards or downwards when resources are
complementary or antagonistic, respectively, and close to linear for the two weakly
interacting scenarios. Within each scenario, the positions of the ZNGIs of the
different species indicate where they lie along the generalist-specialist continuum:
generalists have symmetrical ZNGIs at equal distance from the two resources axes
while specialists on one resource have asymmetrical ZNGIs closer to that resource's
axes. The superimposed supply points helps visualize environmental heterogeneity
in each scenario. In the homogeneous case (top row), we also show how the
consumption by the single species or pair of species (colored or gray arrows &

points, respectively) that dominates after community assembly decreases resource
availability from the supply point to the resource levels at equilibrium, which is the
outcome of invasion analysis (See main text and Box 1). Similar arrows could have
been used to show the variation in outcome from each supply point in the
heterogeneous case (lower panels) but are omitted for clarity purposes.
260
Resource Supply: We also considered how adding environmental heterogeneity at
the local scale affects the results obtained from the Complementarity-Selection
(Loreau and Hector 2001) and Community Assembly components methods. We ran
our model assuming two different, spatially-implicit scenarios for the supply of
resources over the local community: homogeneous resource supply and
heterogeneous resource supply. The homogeneous environment was characterized
by a constant influx of both resources (i.e. a single 'supply point', *sensu* Tilman
1982). The second scenario examines a heterogeneous environment, where species
compete in an ecosystem characterized by 100 microhabitats, each with a separate
supply point (see Fig. 1).
271
Generating a species pool: BEF experiments consist of replicate communities where
species are selected from a common species pool. To generate this species pool, we
created a global pool of ten species whose acquisition traits for the two resources
are regularly spaced along a generalist-specialist continuum defined by a linear
trade-off between the two acquisition traits. We implemented this with the
Schreiber and Tobiason equation (equation 2) that we used to model species growth
rates. In equation (2), x_{1j} and x_{2j} control how efficient species j is at acquiring
resources. We assumed that these two traits are bound by a trade-off, which we
assumed to be linear for simplicity ($x_{1j} + x_{2j} = 1$). Varying these two traits between
0 and 1 generates a generalist-specialist continuum.
282
283 *Community Assembly Outcomes:*
284
For each resource competition scenario (complementary vs. antagonistic,
homogeneous vs. heterogeneous), we imitated the conditions used in standard BEF

experiments to generate multiple individual species pools with different subsets of

species from the global pool that differ in their species richness and composition.

Our species pool of ten species yields 1023 possible subsets of this species pool,

including the 10 different monocultures and the full species pool. We then

considered the equilibrium states of communities initialized with all possible

subsets of this global pool, arranged by their starting richness (e.g., 10 one-species

communities, 45 two-species communities, etc.). This was performed using a global

invasion analysis (Box 1), based on ZNGIs and their envelopes (Koffel et al. 2016). In

the homogeneous case (a single supply point), only one or two species can subsist at

equilibrium on two resources (Levin 1970) but more species can coexist if supply

points are heterogeneous. In the heterogeneous case, 100 supply points were drawn

from a multivariate normal distribution with the same mean as in the homogeneous

case and a variance of 0.5 (See Fig. 1). The equilibrium community state was

determined for each supply point. Because these equilibria for both the

homogeneous and heterogeneous cases are deterministic, only one or two species

will ultimately persist for any single supply point (more species can coexist for

multiple supply points), and the identities of coexisting species and their

abundances are predetermined by their growth functions and consumption vectors.

Each resulting equilibrium community was characterized by the identity of the

species remaining after community assembly and their equilibrium biomass. In the

homogenous case, total ecosystem function is simply the sum of biomass for all

species in the equilibrium community. For the heterogeneous case, equilibrium

biomasses were obtained by averaging over the set of supply points, enabling a

more direct comparison with the homogeneous case.

We then consider key properties of the communities that emerge under each

resource scenario (across diversity levels and all initial compositions). These

include the final biomass of each species, which together determine the total

biomass or function of the ecosystem, and realized species richness (the final

number of species present). This information for each of the subsets was then used

in the Complementarity-Selection and Community Assembly calculations.

Box 1: Mechanisms

In this box, we give a mechanistic explanation of how different competitive

scenarios determine the identity and function of persistent species. In turn, these

will determine the sign and magnitude of the Community Assembly components

that arise from comparing communities assembled from high and low diversity

species pools. We consider both strongly antagonistic and complementary resource

scenarios. We show a simplified example where the full species pool only includes

five species, three of which appear in the low diversity subset. We focus on a

homogenous environment (with one supply point; black dot) in the antagonistic

example and a heterogeneous environment (with three supply points) in the

complementary case.

These situations are represented and analyzed using the graphical approach to

resource competition (MacArthur 1972, Tilman 1982, Leibold 1995, Chase and

Leibold 2003, Koffel et al. 2016). We perform the invasion analysis by drawing the

ZNGIs of the species in the community all together. The species whose ZNGIs are

entirely located above other species' ZNGIs are systematically excluded (e.g. purple,

orange, and green in A). When the impact vectors of the remaining species (red and

blue in A) are combined with a supply point, we obtain the set of these species that

actually end up persisting in the equilibrium community (e.g. red and blue coexist in

A). In a heterogeneous environment (C and D), we perform the invasion analysis for

each supply point independently and average the productivity of the resulting

persisting species to obtain the equilibrium community.

In the antagonistic case, when all members of the full species pool are initially

present (A), the three intermediate species (purple, orange, green; dotted lines) are

excluded by the two extreme specialists (red and blue; solid lines), which coexist.

However, if one of these two specialists (e.g., blue) is randomly absent from a low-
diversity community (B), it is replaced by its closest functional equivalent among
the intermediate species (here, green). This species is not completely specialized on
resource 2, so it does not perform as well as the lost specialist. Due to weak
competition with this intermediate species, the performance of the remaining
specialist (red) is lower than in the full species pool.
356
In the complementary case, the most diverse species pool (C) experiences the
exclusion of the two most extreme species (orange and green), while the three
intermediate species (blue, red, and purple) coexist, as each one is locally adapted to
a particular supply point. If two of these intermediate species, which dominate the
full species pool, are absent in a smaller community (e.g., red and purple), an
extreme species (e.g., green) can survive and coexist with the remaining dominant
species (blue). This extreme species (green) only partially compensates for the loss
of function that occurs due to the absence of the purple species. Similarly, the
remaining dominant species (blue) partially compensates for the loss of function
due to the absence of the red species by increasing in abundance.

Complementarity-Selection analysis:
369
The outputs from our competition models serve as inputs to the Complementarity-
Selection approach (Loreau and Hector 2001). For the purposes of this calculation,
we assumed all species had equal 'initial' abundances. The decomposition divides
total function into the Selection effect (SE) and Complementarity effect (CE) for each
of the 1023 communities. SE and CE were averaged across all possible species pools
at each level of introduced richness regardless of how many species actually existed
at equilibrium.
377
378 *Community Assembly analysis:*
379

No supplementary assumption was needed to apply the Community Assembly

Components (Fox and Kerr 2012; Bannar-Martin et al. 2018). We compared the

community initialized with the entire pool of ten species (our baseline, *sensu*

Bannar-Martin et al. 2018) to each less diverse community. Using the Community

Assembly configuration of the Price equation as described in Bannar-Martin et al.

(2018), we calculated the impact on ecosystem function of species lost (SL), species

gained (SG) and the changes in function of species shared between the most diverse

(and hence, productive) community and each comparison community (CDE). We

then averaged each of these components across all possible species pools within

each level of introduced richness.

Comparison with experimental studies:

To illustrate how our approach can provide insights about actual patterns of

community assembly and the functioning of ecosystems in empirical systems, we

applied the same analyses to data from a well-studied, representative BEF

experiment conducted at Jena (Roscher et al. 2004; 2005; Weigelt et al. 2016).

Established in 2002, this experiment manipulated grassland communities,

controlling the seeded species richness (2, 4, 6, 8, and 16 species, as well as the full

60 species pool) and functional group representation (1 to 4 groups) of 66 plots;

additional small monocultures (3.5 x 3.5 m plots) were established for each species.

Randomly selected subplots were sampled in the spring and summer of each year:

above ground biomass was clipped, sorted to species, dried, and weighed (see

Roscher et al. 2004, 2005 for additional experimental details). Roscher et al. (2005)

found evidence of positive selection and complementarity effects across plots with 2

- 8 species, focusing on data from May of the second year of the experiment (2003).

Using data available from Weigelt et al. (2010), we expanded this Selection-

Complementarity analysis to cover 2003-2008 (focusing on spring data, as do

Roscher et al. 2005), and augmented it by applying the Community Assembly

approach. Prior to analysis, we averaged species' biomasses across replicate

subplots within each experimental plot. The results provide an empirical case study

for comparison with our modeling exercise.

Complementarity-Selection results for each year are based on monoculture biomass

data collected during the corresponding year. Generally, monoculture biomasses

declined over the six-year horizon of this analysis (Marquard et al. 2013; declines in

polyculture yield were weaker). Consequently, there is a tendency for the

magnitude of Complementarity-Selection components to grow more pronounced

over time, due to their mathematical definition (Loreau and Hector 2001). As the

Community Assembly calculations are independent of monoculture data, they were

not similarly affected.

Results:

Overall we studied eight different scenarios with our model. These varied in the

type of competition and the degree of resource heterogeneity as shown in Table 1

and Fig. 1. Although ecosystem function (here total biomass or yield) was always a

saturating function of initial species richness (size of the species pool), different

scenarios produced qualitatively distinct relationships between realized species

richness (at equilibrium) and initial species richness (Fig. 2). These and other

results were often similar for different combinations of resource type and/or

environmental heterogeneity. We focus on qualitatively distinct outcomes as

follows:

Table 1. The eight modeling scenarios in our study, which differ in the type of

resource competition and degree of environmental heterogeneity they feature, as

described in text and illustrated in Fig. 1. Our results showed that both competition

and heterogeneity produced variable effects on complementarity (CE), selection

(SE), species loss (SL), species gain (SG) and context dependence (CDE). We also

found that comparisons between SG and CDE were informative. The magnitude of

these effects is summarized qualitatively through the categories "exactly zero" (0),

"approximately zero" (≈ 0), "negative" (−), "positive" (+) and "strongly positive"

(++).

Scenario	Resource type	Environmental heterogeneity	Predicted effect (see Results) on:					
			CE	SE	SL	SG	SG vs CDE	CDE
1	strongly antagonistic	homogeneous	++	−	−	++	>	≈ 0
2	strongly antagonistic	heterogeneous	++	−	−	++	>	≈ 0
3	weakly antagonistic	homogeneous	++	−	−	++	>	−
4	weakly antagonistic	heterogeneous	++	−	−	++	>	−
5	weakly complementary	homogeneous	≈ 0	+	−	++	>	0
6	weakly complementary	heterogeneous	≈ 0	+	−	+	<	+
7	strongly complementary	homogeneous	≈ 0	+	−	++	>	0
8	Strongly complementary	heterogeneous	+	+	−	+	<	+

Figure 2. A) Realized richness obtained after species sorting, averaged across all possible species pools at each level of introduced richness. In the four antagonistic resource scenarios, realized richness quickly saturated at two species. In the complementary resource scenarios, average realized richness is hump-shaped in the homogeneous case, but increasing in the heterogeneous case. B) Total community yield obtained after species sorting, averaged across all possible species pools at each level of introduced richness; values are scaled relative to the yield of the most productive monoculture (solid horizontal line). All scenarios give saturating responses but in scenarios where resources are close to substitutable (central panels), moving from monocultures to polycultures causes weaker increases in yields.

Figure 3. Complementarity (CE) and selection effects (SE) vary in magnitude and
sign as a function of introduced richness, competitive scenario (columns), and
supply point heterogeneity (rows). Antagonistic resources lead to CE > 0 and SE < 0,
while complementary resources result in CE > 0 and SE > 0. As in Fig. 2, violin plots
indicate averages and variability across all possible species pools at each level of
introduced richness.

Figure 4. Community Assembly Components method, comparing the most diverse
polyculture (initial richness of 10) against less diverse mixtures. CDE is nearly
always zero or slightly negative, except when resources are complementary and
have a heterogeneous supply. As in Fig. 3, violin plots indicate averages and
variability across all possible species pools at each level of introduced richness.

Antagonistic resources:

Antagonistic resources, whether weak or strong, or supplied uniformly or

heterogeneously, produced similar results. First, the maximum yield of polycultures

was always greater than the maximum monoculture. Second the realized richness

was always two species or fewer. In these scenarios, the average complementarity

effect (CE) was always positive — setting aside the weakly antagonistic case with

introduced richness of two — whereas the average selection effect (SE) was always

negative (Fig. 3). Finally, the average Community Assembly Components always

showed positive species gains (SG) and negative species loss (SL) components

whereas the CDE was always small or zero (Fig. 4).

Strongly complementary resources in homogeneous environment:

In contrast to antagonistic resources, results varied substantially depending on just

how strongly resources are complementary and on the degree of environmental

heterogeneity. Starting with strongly complementary resources (almost equivalent

to essential resources), we found that maximum yield in polycultures did not differ

from that of the most productive monoculture. Both selection (SE) and

complementarity effects (CE) were positive, in contrast with antagonistic resources

(Fig. 3). The increase in yield with diversity was largely due to an increase in the

mean selection effect: as the size of species pools grew, more productive

competitors were more likely to be present. The relationship with complementarity

was unimodal and reflected the increased role of complementarity on yield in small

species pools where the better competitors were absent. The effects of

complementarity were also generally small. Community Assembly Components

showed that effects of CDE were zero and that all of the effects involved positive

effects of SG and negative effects of SL (Fig. 4). We also found that mean realized

species richness was a unimodal function of the size of the initial species pool in the

homogeneous scenario (but still limited to two or fewer species; Fig. 2.A and S1).

This means that the relationship between realized diversity and yield appears

complex (Figure S1).

Strongly complementary resources in heterogeneous environments:

The patterns differed from the homogenous environment results in several respects.

First, the maximum productivity was typically higher than that of the most

productive monoculture. Also, realized species richness was now positively related

to the size of the species pool. In contrast with the homogeneous case, the positive

complementarity effect (CE) persisted at high diversity levels (Fig. 3), while species

gains (SG) had much smaller effects and CDE was positive, counteracting a similar

SL effect (Fig. 4). We also examined an intermediate scenario, with smaller

variance in supply point heterogeneity, and found intermediate patterns between

the homogenous case and the full variance case: intermediate SE and CE, and CDE

and SG of comparable magnitudes (not shown).

Weakly complementary resources:

With resources that were only weakly complementary, the increase in yield with

diversity was less pronounced. Both CE and SE results were similar to the

complementary case (Fig. 3), except that environmental heterogeneity did not

support a sustained positive CE effect at high diversity levels.

Relations between realized species richness and biomass:

In each of our scenarios, we also examined how total biomass was related to the

realized species richness (the number of species coexisting at equilibrium). Over all

initial richness levels, there was generally a positive relationship between standing

biomass and realized species richness (Fig. S1). However, scenarios differed in the

strength of this relationship and in the structure of covariation between biomass

and realized species richness when the size of the species pool was accounted for.

In most scenarios (scenarios 1-6 and 8 in Table 1), we found that biomass was

positively related to realized species richness and that this was true both at the

global scale of analysis and within subsets of communities that differed in the size of

the introduced species pool. In contrast, when resources were strongly

complementary and environmental heterogeneity low (scenario 7) we found that

total biomass was negatively related to realized species richness for any given level

of initial richness (Fig. S2).

Mechanisms underlying relationships between resource scenarios and

Complementary-Selection and Community Assembly patterns

The striking differences between the antagonistic and complementary resource

competition scenarios captured by the Community Assembly components can be

linked to the influence of resource type on the outcomes of community assembly. In

the antagonistic case, a pair of extreme specialists dominates all the other species

for any supply point (Box 1, panels A, B). This strongly affects both the

complementarity and selection effects (CE and SE), and the Community Assembly

components (SL, SG, and CDE). Dominance by a pair of specialists leads to a realized

richness that cannot exceed two even in the heterogeneous case. In monoculture,

each specialist strategy generally performs worse than the intermediate generalist

strategies, which explains why the selection effect is negative. Yet, the two

specialists together perform better than any other set of one or two species, driving

the positive CE pattern. When one of the two dominant specialist strategies is absent

(causing a negative SL component), it is replaced by a less specialized species with

lower function than the missing specialist (which results in a positive SG component

but |SL| > |SG|). The less specialized species competes more strongly with the

remaining specialist due to greater niche overlap, reducing the performance of the

remaining specialist, which creates the negative CDE.

In the complementary cases, there is usually a unique generalist species from the

full species pool that dominates for a given supply point (Box 1, panels C, D). This

dominant species is also the one that performs the best – better than any other set

of one or two species – for this supply point, which explains why selection is

positive and complementarity weak and hump-shaped. The CDE of exactly zero in

the homogeneous case is a direct consequence of having a single dominant strategy:

either this dominant strategy is absent from a low diversity community, which will

then ultimately share no species with the high diversity community, or the

dominant strategy is present, producing an endpoint identical to the high diversity

community. Finally, the positive CDE in the heterogeneous case arises because in the

absence of one or more of the dominant strategies, other dominant species that still

occur compensate for functional losses. Such CDE is larger than the SG component

on average because the functional contributions of new species (able to join a

community due to the loss of at least one dominant species) are smaller than

increases in function occurring through compensation by the remaining dominants.

Comparisons with experimental data:

Although all of our scenarios produce broadly saturating BEF patterns, our results

indicate that we can qualitatively discriminate among most of these eight different

community assembly scenarios (Table 1) by combining the Community Assembly

(Fox and Kerr 2012, Bannar-Martin et al. 2018) and the Complementarity-Selection

(Loreau and Hector 2001) analyses. This suggests that applying these methods to

empirical data may offer insights into which competitive scenarios drive assembly,

giving rise to empirically-obtained BEF relationships (or at least provide evidence

about which mechanisms may be ruled out). To assess this, we analyzed data from a

representative BEF experiment, conducted at Jena, Germany, to determine if we

could identify the community assembly mechanisms underpinning the functioning

of this grassland ecosystem. We used six years of data from the Jena experiment,

starting in 2003 (a year after the experiment was established), to consider whether

results were stable over time.

We found that the data from Jena did not match any of the scenarios shown in Table

1. On the one hand, we see a divergence between increasingly negative selection

effects and increasingly positive complementarity effects from 2003 to 2008 (Fig. 5),

creating a pattern consistent with the antagonistic resource scenario using the

Complementarity-Selection approach (Fig. 3, see also Marquard et al. 2009). On the

other hand, we see persistent trends in the Community Assembly components,

including positive CDE effects across the entire time span (Fig. 6), which were only

observed in our model scenarios given competition for heterogeneous,

complementary resources (Fig. 4).

Consequently, the analysis of Complementarity-Selection effects (Fig. 5) is

consistent with antagonistic resources, be it in homogeneous or heterogeneous

supply points (compare with Fig. 3), whereas the Community Assembly

Components suggest a scenario of complementary resources in the presence of

environmental heterogeneity (comparing Fig. 5 with Fig. 4). This indicates, perhaps

unsurprisingly, that the eight simplistic scenarios we considered probably lack

additional mechanisms involved in regulating or modifying the effects of community

assembly on ecosystem function (and hence the metrics used to study it).

611
Figure 5. Complementarity (CE) and selection (SE) effects (top row), and
Community Assembly components comparing the most diverse (60 species)
community to less diverse communities (bottom row) over successive years at the
Jena Experiment, as a function of introduced species richness (log scale). Early in the
experiment, CE and SE were both positive, and weakly related to diversity (as found
by Roscher et al. 2005). However, by the end of the period we examined, the CE

effect was positive and increased strongly with diversity, while the SE effect was
negative and decreased with diversity. Although there is some variation in the
magnitude of the Community Assembly components through time, at all lower
diversity levels, CDE effects are consistently positive, SG is negligible, and SL
strongly contributes to declines in function. Trend lines show Generalized Additive
Models (GAMs) fit to the observed plot-level values; points jittered for visual clarity.
624

We thus considered other possible factors that might have affected community

assembly at Jena. One of these is that the observed dynamics may not have reached

steady state as assumed by our models above. To address this possibility, we

simulated the temporal dynamics of community assembly for the heterogeneous,

complementary resources scenario (because previous work has mostly emphasized

that this scenario is more likely than others for plants; see Tilman 1982, Tilman et

al. 1997, Gross and Cardinale 2007, Hodapp et al. 2016). Starting from an initial

condition where all species are seeded at an equally low density, we let their

populations grow and compete over time. When a species' density dropped under a

certain minimal threshold (set equal to 10^{-4}), we considered it extinct and set its

density equal to zero. We then extracted species densities at representative time

points and applied both partitions as previously described. Although this exercise

shows that there are important transient dynamics in the community assembly

dynamics of BEF experiments (Fig. 6), none of the resulting patterns are consistent

with the patterns we observed in Jena (Fig. 5). Intriguingly, we found that the

dynamics in our model (Fig. 6) showed a strong qualitative shift in the relative

importance of CE and SE through time and a somewhat subtler shift from linear to

unimodal CDE and an increase in SG through time. However, the data from Jena

show no such trends.

Figure 6: Transient dynamics in Complementarity-Selection effects and Community Assembly components as assembly proceeds towards equilibrium in the scenario where species compete for heterogeneous, strongly complementary resources.

We also considered that facilitation, perhaps due to the nitrogen (N) fixing effects of

legumes, may play an important role in assembly, and consequently affect

ecosystem functioning. To address this, we adapted the model of Koffel et al.

(2018), and again applied Complementarity-Selection and Community Assembly

analyses (see SI for details). For the case with high environmental heterogeneity

and strong facilitation under high initial N-limitation, our results (Fig. 7) correspond

at least qualitatively to our observations at Jena. Both the facilitation model and

Jena data demonstrate greater complementarity effects (CE) in comparison to

selection (SE) and, at least for Jena's 2007 and 2008 data, a negative SE in the

Complementary-Selection Effects analysis (Fig. 5,7A). This seems consistent with

observations from the Jena Experiment that showed that N fixation rates by

legumes, and thus potentially facilitation, increased from 2004 to 2008 (Roscher et

al. 2011). The Community Assembly Components also show similarities, with a

substantial and humped CDE effect that emerges from the model (Fig. 8B) also being

seen in the Jena data (Fig. 6). The addition of facilitation creates a combination of

Complementarity-Selection Effects and Community Assembly Components patterns

that otherwise appear to point to divergent resource competition models in the

absence of facilitation.

Figure 7: Model results from including nitrogen fixing plants that compete with non-fixers. Complementary-Selection Effects (panel A) and Community Assembly Components (panel B).

Discussion:

There are two major implications of our work. First, we find that the joint use of the Community Assembly (Bannar-Martin et al. 2018) and the Complementarity-

Selection (Loreau and Hector 2001) approaches can be a powerful tool in inferring

mechanisms involved in linking community dynamics to the functioning of

ecosystems. Second, we find that empirical results (at least in the case of the Jena

BEF experiment) do not match any of the scenarios we examined for purely

competitive mechanistic models but that they did when we incorporated facilitation

by N-fixing plants. This suggests that the joint evaluation of Complementarity-

Selection and Community Assembly components could be a powerful tool for

identifying plausible mechanisms that relate community assembly to the

functioning of ecosystems.

Validating the joint use of Community Assembly and Complementarity-Selection

partitions to interpret mechanisms involved in BEF dynamics:

Numerous studies in controlled and replicated conditions (such as those that prevail

in the large literature involving BEF studies) have shown that there is generally a

positive and directional effect between the size of the species pool with access to a

community and the function of at least some aspects of ecosystems, and in

particular the total biomass of the organisms involved. Previous work on

relationships between species pool diversity and ecosystem function has largely

focused on two very general effects that can explain this pattern: selection (the

species pool, when large, is more likely to include highly competitive and highly

productive species) and complementarity (the species pool, when large, is more

likely to include species that coexist and whose production is less mutually

inhibitory). To date, theoretical explanations for these two general effects have

focused on resource competition and used models based on complementary

resources (e.g., Tilman et al. 1997, Gross and Cardinale 2007, Leibold et al. 2017).

However, other mechanisms may also be involved. Here we explored the joint

consequences of manipulating the mechanisms governing species competition

(ranging from strongly antagonistic to strongly complementary resources), resource

heterogeneity and the size of the species pool on realized species richness,

composition, and production. We found that selection enhanced production when

resources were complementary, but undermined production when resources were
antagonistic. Complementarity between species had little effect on production
except when resources were antagonistic and/or heterogeneous.
713
Consequently, the mechanisms involved are not necessarily clearly resolved by the
study of complementarity and selection alone. We found that additional insights
could be gained by also considering the Community Assembly components (Bannar-
Martin et al. 2018). Our work shows that these components can provide additional
and complementary insights into the mechanisms structuring the relationships
among species pool size, ecosystem function, realized species richness and
community composition. In our case, the joint use of Community Assembly and
Complementarity-Selection approaches provided well-defined criteria by which we
could distinguish between competition for complementary vs. antagonistic
resources as well as evaluate the role of environmental heterogeneity and
facilitation.
725
726 *Insights about mechanisms that drive community-ecosystem function relations:*
727
Our study also highlights the fact that we do not necessarily understand all the
mechanisms that drive community-ecosystem relationships. Patterns in the data
from the Jena experiment did not correspond to any of the eight scenarios we
modeled and allowed us to reject all eight of these hypotheses. On the one hand,
this shows how powerful the joint application of Community Assembly and
Complementarity-Selection metrics can be. However, this also shows that the
mechanisms governing the diversity and function of communities must involve
other processes besides simple resource competition. This is not totally surprising
since our theoretical models are quite simplistic caricatures of how species interact.
Here we used the joint analysis of Community Assembly and Complementarity-
Selection metrics to evaluate two other possible explanations. In one, we evaluated
the possibility that community assembly at Jena was not in final equilibrium — a
hypothesis supported by the directional changes in Complementarity-Selection

components across years (Fig. 5) but found that this did not explain the data at Jena.

In the other, we derived predictions that included the role of N-fixing plants and

found that this model with facilitation did qualitatively correspond to the

observations at Jena. Critically, we note that a correspondence between theory and

observation is not definitive proof that a particular mechanism is operating. For

example, even though our model was explicitly based on N fixation, there is no

reason to believe that the facilitation mechanism that emerge in our model is

restricted to this ecological situation. Results similar to ours could arise from a

broader range of facilitation mechanisms at play in BEF experiments, such as

facilitation related to other abiotic factors (Wright et al. 2017) or based on soil

organisms (Eisenhauer et al. 2012). Of course, there are probably a number of other

mechanisms that could also be modeled, including interactions mediated via other

organisms such as grazers, pathogens and other mutualists.

Community assembly and the functioning of ecosystems in other scenarios:

How then does community assembly determine the functioning of ecosystems in

general? We focused on a very simple scenario for community assembly that

mimics the experimental design of 'classic' BEF experiments. This scenario consists

of looking at all possible subsets of introduced species combinations for a given

'total species pool'. This can be thought of as a filter on potential to 'arrive' in a

community via dispersal limitation. Assembly is then followed by species

interactions in which some species within this subset of potential species are

excluded by competition for resources. We find that this process can be related to

analysis by both the Complementarity-Selection and the Community Assembly

approaches. However, community assembly in real communities can involve

numerous factors not typically incorporated in BEF experiments, including species

invasions, disturbance and dispersal dynamics, environmental change, and

stochastic processes (e.g. Thompson and Gonzalez 2016, Hodapp et al. 2016, Leibold

et al. 2017, Bannar-Martin et al. 2018). The approach developed here on the well-

known BEF experimental setup could potentially be adapted to the study of these

other natural conditions. This would, however, require careful thinking about how

to model these mechanisms and their interaction with resource competition.

While using the Community Assembly and the Complementarity-Selection

approaches together can be a useful way to assess mechanisms, it is important to

note that Complementarity-Selection partitions require estimates of ecosystem

function in monocultures. These are readily available in most 'classic' BEF

experiments as a pro-forma aspect of the experimental design. However, these are

almost always absent in other experiments that examine how community assembly

affects ecosystem functions (e.g. disturbance, invasions, extinctions, environmental

change), let alone in observational data. The logistical challenge is even greater in

studies examining the effects of environmental change on ecosystem function, as

monoculture treatments would be required for all the relevant environmental

conditions. Additionally, many organisms are not easily grown in monocultures

due, for example, to dependencies on biotic resources, or movement. Thus, the

scenarios where the Complementarity-Selection approach can be used are limited.

In contrast, the Community Assembly approach can be calculated for any pair of

communities. Given the design of BEF studies, we opted to use the most diverse

species pool as our baseline for comparisons, but this is not required (see Bannar-

Martin et al. 2018 for examples). While the Complementarity-Selection approach is

constrained by the requirement of monoculture data, it did appear to more clearly

distinguish between antagonistic and complementary resource scenarios. This

ability in part may rest in the definition of these metrics: both CE and SE are affected

by changes in the abundance of a species. In contrast, of the three Community

Assembly components, SL and SG are mostly influenced by the actual loss or gain of

species from a community; all effects of less extreme changes in abundance are

relegated to the CDE component alone. For this reason, while it is possible to apply a

Community Assembly approach in a manner directly analogous to

Complementarity-Selection (contrasting an observed polyculture with an idealized

polyculture where species perform proportional to their monocultures), the results

are somewhat less informative.

Our work shows that, while both methods exhibit their own advantages and

limitations, together they provide a more powerful means of diagnosing

mechanisms. Future work could focus on continuing to refine these metrics. This

might include developing analogs of Complementarity-Selection partitions that do

not require monoculture data. Alternatively, new or existing extensions of the

Community Assembly approach (e.g., 5-part Price equation; Fox & Kerr 2012;

Bannar-Martin et al. 2018) could be applied to these or other model scenarios. For

example, while empirical applications of the trait-based version of the Price

equation (Fox & Harpole 2008) can be hampered by the availability of trait data, in

modeling exercises these traits are explicitly known (e.g., species' ZNGIs in our

scenarios). Further work could elucidate not only the mechanisms structuring

ecosystem function across communities of different diversity levels but relate

ecosystem properties to the functional composition of communities.

Our study also illustrates that one can discriminate between *a-priori* hypotheses and

ad-hoc hypotheses in ways that relate to the entire issue of 'replicability' in science

(e.g. Nichols et al. 2019). Here we set out to test 8 different hypotheses about how

resource competition affects the relations between community assembly and the

functioning of ecosystems. We generated these predictions with *a-priori* models

based on mechanistic resource competition theory (Tilman et al. 1997, Gross and

Cardinale 2007, Hodapp et al. 2016). We subsequently tested these predictions on

data from a previously conducted experiment that had not been previously studied

in this way and found that we rejected all of our a priori hypotheses. Of course, we

might have found that one or more of these hypotheses were not rejected and

stopped there with statistically valid inferences. However, the rejection of all

hypotheses begs the question 'what could the explanation instead be?'. Here we did

some *post-hoc* modeling based on two additional mechanisms and found that one of

these was consistent with the data. However, the statistical significance of this

hypothesis is not clear since we might have rejected it and continued on to testing

other hypotheses until we found one that was consistent with the data whereupon

we would have perhaps stopped. In our discussion we tried to indicate that other

mechanisms could also have been similarly successful. The point is not that we

don't believe that our tests are useful (they did in fact reject 9/10 hypotheses), but

rather that the strength of our inferences about the hypothesis that was supported

by the data should be appropriately weighed by the fact that it was an *ad-hoc*

hypothesis rather than an *a-priori* one.

Implications for future work:

These encouraging results suggest a number of ways whereby the analysis of

community assembly components could be improved:

a) Most obviously, one could continue to propose and test other mechanisms,

such as plant-soil feedbacks and multi-trophic interactions, that could have

produced the patterns we observed in the Jena data, eventually potentially

creating a 'periodic table' for how the Complementarity-Selection and

Community Assembly metrics map onto a diverse array of community

assembly scenarios. While there is some allure to this suggestion, there are a

considerable number of interacting mechanisms to test requiring careful

thinking about how to construct and interpret models (e.g., considering

transient dynamics), or how to parse varying contrasts in the data (e.g.,

considering facilitation by legumes).

b) Discriminating among the wide array of mechanisms driving community

assembly and its effects on ecosystem function likely requires moving

beyond simple dichotomous tests with one or two metrics since many

alternative models/mechanisms are capable of reproducing similar, low-

dimensional patterns. Here we demonstrated how the addition of several

carefully considered metrics could enhance our understanding of these

mechanisms, as well as our ability to differentiate them. It is worth

considering whether other additional metrics could similarly advance

understanding. The degree to which this is so would depend on finding

metrics that reveal distinct aspects of mechanisms and that, thus, are not

simply redundant with each other. It is also useful when the link between

mechanism and metrics is relatively transparent (as we explored in Box 1).

In cases where this is not the case this is perhaps less true and other

modeling or experimental studies would be needed to consolidate any

conclusions. The increase in the number of metrics, however, may be

necessary if theoretical models increase in complexity by including missing

processes or by assessing different assumptions in order to decrease

equifinality and increase inference.

c) Another reason to consider additional metrics is that not all data will be

suitable for the same metrics. In our case, while Complementarity-Selection

was particularly useful in discriminating between different types of resource

competition, the approach is constrained by the need for monoculture data.

We will need new techniques if we wish to apply these types of approaches

to a wider array of experimental conditions, with designs that do not closely

target BEF relations or that involve organisms where such monocultures are

logistically difficult (e.g. birds, migrating organisms, species-rich tropical

communities).

d) While we need to explore other metrics to use in studies other than BEF

experiments, it would still be interesting to apply the methods in this paper

to other BEF experiments. We suspect that this may likely show substantial

variation in results across locations, habitats, and organisms. For example, it

would be interesting to compare results at Jena with data from sites that are

not as likely to be nitrogen limited. Adequately considering such variation

would require a far more comprehensive theoretical and analytical effort

than is warranted in this paper.

e) Even more challenging is to consider how the size of the species pool jointly

affects biodiversity, community composition, and ecosystem functioning

under other scenarios than those at play in classical BEF experiments.

Numerous challenges to BEF research have focused on the degree to which

BEF experiments may be unrealistic representations of how biodiversity and

ecosystem function are likely to interrelate because the experimental setup

for community assembly in BEF experiments is so artificial (Wardle 2016).

We believe the Community Assembly approach can shed light on other

situations including arrivals and removals of invasive species as well as

environmental change (see Bannar-Martin et al. 2018, Kremer et al. in prep).

Linking these scenarios to explicit theoretical models (as we did here by

asking how resource types and environmental heterogeneity affect a BEF

experiment) represents an exciting avenue for future devoted work.

Software availability:

An R package is available to facilitate future work (including analyzing other

biodiversity experiments) https://github.com/ctkremer/priceTools

Acknowledgements:

This paper is a joint effort of the working group sCAFE and an outcome of a

workshop kindly supported by sDiv, the Synthesis Centre of the German Centre for

Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig (DFG FZT 118). The Jena

Experiment is funded by the German Research Foundation (FOR456/1451) with

additional support from the University of Jena and the Max Planck Society.

References

Bannar-Martin, K. H., Kremer, C. T., Ernest, S. K. M., Leibold, M. A., Auge, H., Chase, J.,

Declerck, S. A. J., Eisenhauer, N., Harpole, S., Hillebrand, H., Isbell, F., Koffel, T.,

Larsen, S., Narwani, A., Petermann, J. S., Roscher, C., Cabral, J. S., and Supp, S. R.

(2018). Integrating community assembly and biodiversity to better understand

ecosystem function: the Community Assembly and the Functioning of Ecosystems

(CAFE) approach. *Ecology Letters*, 21:167–180.

Bell, T., Lilley, A. K., Hector, A., Schmid, B., King, L., and Newman, J. A. (2009). A linear

model method for biodiversity-ecosystem functioning experiments. *The American*

*Naturalis*t, 174(6):836–849.

Cardinale, B.J., Duffy, J.E., Gonzalez, A., Hooper, D.U., Perrings, C., Venail, P., et al.

(2012). Biodiversity loss and its impact on humanity. *Nature*, 486, 59-67.

Chase, J. M. and Leibold, M. A. (2003). *Ecological Niches: Linking Classical and*

Contemporary Approaches. University of Chicago Press, Chicago.

Eisenhauer, N., Reich, P.B. and Scheu, S., 2012. Increasing plant diversity effects on

productivity with time due to delayed soil biota effects on plants. Basic and Applied

Ecology, 13(7), pp.571-578.

Fox, J.W. (2006). Using the Price equation to partition the effects of biodiversity loss

on ecosystem function. *Ecology*, 87, 2687-2696.

Fox, J.W., & Harpole, W.S. (2008). Revealing how species loss affects ecosystem

function: the trait-based Price equation partition. *Ecology*, 89, 269–279.

Fox, J.W., & Kerr, B. (2012). Analyzing the effects of species gain and loss on

ecosystem function using the extended Price equation partition. *Oikos*, 121, 290-

298.

Gross, K., & Cardinale, B.J. (2007). Does species richness drive community

production or vice versa? Reconciling historical and contemporary paradigms in

competitive communities. *The American Naturalis*t., 170, 207-220.

Hector, A., Bell, T., Connolly, J., Finn, J., Fox, J. Kirwan, L., Loreau, M., McLaren, J.,

Schmid, B., and Weigelt, A. (2009). The analysis of biodiversity experiments: from

pattern toward mechanism. pp. 94-104 in Naeem, S., Bunker, D.E., Hector, A.,

Loreau, M., and Perrings, C. (eds.). Biodiversity, Ecosystem Functioning, and Human

Wellbeing: An Ecological and Economic Perspective. Oxford University Press.

Oxford.

Hooper, D. U., Adair, E. C., Cardinale, B. J., Byrnes, J. E. K., Hungate, B. A., Matulich, K.

L., Gonzalez, A., Duffy, J. E., Gamfeldt, L., and O'Connor, M. I. (2012). A global

synthesis reveals biodiversity loss as a major driver of ecosystem change. *Nature*,

486(7401):105–108.

Kirwan, L., Connolly, J., Finn, J. A., Brophy, C., Lüscher, A., Nyfeler, D., and Sebastià,

M.-T. (2009). Diversity-interaction modeling: estimating contributions of species

identities and interactions to ecosystem function. *Ecology*, 90(8):2032–2038.

Koffel, T., Daufresne, T., Massol, F., and Klausmeier, C. A. (2016). Geometrical

envelopes: extending graphical contemporary niche theory to communities and eco-

evolutionary dynamics. *Journal of Theoretical Biology*, 407:271–289.

Leibold, M. A. (1995). The niche concept revisited: mechanistic models and

community context. *Ecology*, 76(5):1371–1382.

Leibold, M. A., Chase, J. M., & Ernest, S. K. M. (2017). Community assembly and the

functioning of ecosystems: how metacommunity processes alter ecosystem

attributes. *Ecology*, 98, 909–919.

León, J. A. and Tumpson, D. B. (1975). Competition between two species for two

complementary or substitutable resources. *Journal of Theoretical Biology*,

50(1):185–201.

Levin, S. A. (1970). Community equilibria and stability, and an extension of the

competitive exclusion principle. *The American Naturalist*, 104(939):413–423.

Loreau, M. & Hector, A. (2001). Partitioning selection and complementarity in
biodiversity experiments. *Nature*, 412, 72-76.

MacArthur, R. H. (1972). *Geographical Ecology: Patterns in the Distribution of Species*.
Harper & Row, New York.

Marquard, E., Weigelt, A., Temperton, V.M., Roscher, C., Schumacher J., Buchmann, N.,
Fischer, M., Weisser, W.W., Schmid, B., (2009). Plant species richness and functional
composition drive overyielding in a six-year grassland experiment. *Ecology* 90(12),
pp. 3290-3302.

Nichols, J.D., Kendall, W.L., Boomer, G.S. (2019) Accumulating evidence in ecology:
Once is not enough. *Ecology and Evolution* 2019; 00:1-14.

1001

1002 Ptacnik, R., Moorthi, S.D., & Hillebrand, H. (2010). *Hutchinson Reversed, or Why There*
1003 *Need to Be So Many Species*. In G. Woodward (Ed.), Advances in Ecological Research
1004 (Vol. 43, pp. 1–43). Burlington: Academic Press.

1005

1006 Roscher, C., Temperton, V.M., Scherer-Lorenzen, M., Schmitz, M., Schumacher, J.,
1007 Schmid, B., Buchmann, N., Weisser, W.W. and Schulze, E.D., (2005). Overyielding in
1008 experimental grassland communities–irrespective of species pool or spatial scale.
1009 *Ecology Letters*, *8*(4), pp.419-429.

1010

1011 Roscher, C., Schumacher, J., Baade, J., Wilcke, W., Gleixner, G., Weisser, W.W., Schmid,
1012 B. and Schulze, E.D., (2004). The role of biodiversity for element cycling and trophic
1013 interactions: an experimental approach in a grassland community. *Basic and Applied*
1014 *Ecology*, *5*(2), pp.107-121.

1015

1016 Roscher, C., Thein, S., Weigelt, A., Temperton, V.M., Buchmann, N. and Schulze, E.D.,
1017 2011. N 2 fixation and performance of 12 legume species in a 6-year grassland
1018 biodiversity experiment. Plant and Soil, 341(1-2), pp.333-348.

1019

1020 Rychtecká, T., Lanta, V., Weiterová, I., and Lepš, J. (2014). Sown species richness and

1021 realized diversity can influence functioning of plant communities differently.

1022 *Naturwissenschaften*, 101(8):637–644.

1023

1024 Schreiber, S. J. and Tobiason, G. A. (2003). The evolution of resource use. *Journal of*

1025 *Mathematical Biology*, 47:56–78.

1026

1027 Thompson, P.L., and Gonzalez, A. (2016). Ecosystem multifunctionality in

1028 metacommunities. *Ecology*, doi: 10.1002/ecy.1502.

1029

1030 Tilman, D. (1982). *Resource Competition and Community Structure*.

1031 Princeton University Press, Princeton, NJ.

1032

1033 Tilman, D., Lehman, C. L., and Thomson, K. T. (1997). Plant diversity and ecosystem

1034 productivity: theoretical considerations. *Proceedings of the National Academy of*

1035 *Sciences (USA),* 94:1857–1861.

1036

1037 Tilman, D., Reich, P. B., and Isbell, F. (2012). Biodiversity impacts ecosystem

1038 productivity as much as resources, disturbance, or herbivory. *Proceedings of the*

1039 *National Academy of Sciences*, 109(26), 10394-10397.

1040

1041 Wardle, D.A. (2016). Do experiments exploring plant diversity-ecosystem

1042 functioning relationships inform how biodiversity loss impacts natural ecosystems?

1043 *J. Veg. Sci.*, 27, 646-653.

1044

1045 Weigelt, A., Marquard, E., Temperton, V.M., Roscher, C., Scherber, C., Mwangi, P.N.,

1046 Felten, S., Buchmann, N., Schmid, B., Schulze, E.D. and Weisser, W.W. (2010). The

1047 Jena Experiment: six years of data from a grassland biodiversity experiment.

1048 *Ecology*, *91*(3), pp.930-931.

1049

1050   Weigelt, A., de Luca, E., Roscher, C., Temperton, V., Buchmann, N., Fischer, M., Scher-

1051   Lorenzen, M., Schmid, B., Schulze, E.-D., Weisser, W.W., Guo, L., Meyer, S.T. (2016).

1052   Collection of aboveground community and species-specific plant biomass from the

1053   Jena Experiment (time series since 2002). PANGAEA.

1054   https://doi.org/10.1594/PANGAEA.866358

1055

1056   Wright, A.J., Wardle, D.A., Callaway, R. and Gaxiola, A., 2017. The overlooked role of

1057   facilitation in biodiversity experiments. Trends in Ecology & Evolution, 32(5),

1058   pp.383-390.

1059

Supplementary Informations

1060

1061

1062

1063 **Figures**

1064

1065 **Figure S1**. Relationship between yield and realized richness for the 8 scenarios. The

1066 small light points represent the equilibrium community resulting from seeding all

1067 possible subsets of the global species pool. The large darker points give the average

1068 yield and realized richness, where the average is taken across all possible

1069 communities for a given level of introduced richness.

1070

1071

Figure S2. Relationship between realized richness and yield for the strongly

complementary case in a homogeneous environment (scenario 7, Table 1). Each

small point represents an equilibrium community resulting from seeding a given

subset of the global species pool, coded by levels of introduced richness (from 1 to

10). For each level of introduced richness, the large points give the average yield

across each level of realized richness. The lines connecting these points highlight the

fact that within scenario 7, total biomass is negatively related to realized species

richness for any given level of introduced richness.

1080

1081

1082

1083

1084 **Figure S3**. Jena yield (biomass) over time, relative to hypothetical community based

1085 on monoculture performance, as a function of introduced richness (log scale).

1086 Complementarity-Selection analysis (Fig. 4) partitions these responses.

1087

1088 **Facilitation model**

1089

To study the effects of facilitation on the Complementarity-Selection Effects and the

Community Assembly components, we developed a variant of the resource

competition model from the main text specifically tailored to model the interactions

between nitrogen-fixing and non-fixing terrestrial plants. This model is very similar

to the model of Koffel et al. (2018), with a few differences that will be highlighted

below. This model still fits the general form of Eq. (1), with now R_1 and R_2

respectively being soil Phosphorous (P) and Nitrogen (N). Yet, the expressions of

both the per capita growth rate and the consumptive impacts differ from the general

form from Schreiber and Tobiason (2003) presented in the main text. The growth

rate $g_j(R_1, R_2)$ here reads:

1100

$$g_j(R_1, R_2) = \text{Min}(x_{1j}R_1, x_{2j}R_2 + f_j)$$

1101

1102 As the two resources R_1 and R_2 correspond to P and N, they are modeled as being

1103 strictly essential ($\alpha = -\infty$ in the approach presented in the main text) which results

1104 in taking the minimum of the growth rates on the two resources via the "Min"

1105 function.

1106

The extra term that is not accounted for by the Schreiber and Tobiason (2003)

approach is the constant fixation rate f_j that boosts growth when N is limiting. The

fixation rate f_j is the focal trait here, and the species pool was generated by

selecting species with varying values of f_j. A species with $f_j = 0$ is a non-fixer, while

species with non-zero values of f_j describe N-fixers with various intrinsic abilities to

fix N. We included a cost to N-fixation in the form of a decreasing ability to acquire

phosphorous through x_{1j} as f_j increased, mathematically given by $x_{1j} = h(f_j)$, with

h a decreasing function. Conversely, soil nitrogen acquisition x_{2j} was assumed to be

the same for all species, and thus taken constant.

For facilitation to happen, N fixed by the N-fixers has to become available to the
other species. This is done by modifying the consumptive impacts to account for N
restitution through recycling, potentially leading to net negative consumptive
impacts when N restitution by an N-fixer is greater than N uptake. This leads to the
following consumptive impacts on N:
1121

$$C_{2j}(R_1, R_2) = q_{2j}\left[\mathrm{Min}(x_{1j}R_1, x_{2j}R_2) - \lambda_{2j}m\right]$$

1122

The positive term in $C_{2j}(R_1, R_2)$ accounts for soil N uptake through growth, and is
thus equal to $g_j(R_1, R_2)$ without the fixation term f_j because fixed N comes from the
atmosphere, not the soil. The negative term is N restitution through recycling from
dead plant biomass (generated at the rate m, assuming the same mortality rate for
all species for simplicity) with an efficiency given by λ_{2j}. We assumed that the
efficiency λ_{2j} was larger for species with large fixation rate f_j, translating the fact
that N-fixing species tend to see their dead biomass more readily recycled. The
coefficient q_{2j} is a stoichiometric coefficient similar to the ones introduced in the
models from the main text.

1132

1133 On the P side, the consumptive impact reads:

1134

$$C_{1j}(R_1, R_2) = q_{1j}[g_j(R_1, R_2) - \lambda_{1j}m]$$

1135

1136 The first term of the sum is here exactly equal to growth $g_j(R_1, R_2)$, which translates
1137 the fact that unlike N, all the P entering plant biomass has been taken up from the
1138 soil. We assume that P is also recycled, through an efficiency λ_{1j} assumed to be the
1139 same for all species for simplicity. The coefficient q_{1j} is again a stoichiometric
1140 coefficient.

1141

1142 The global species pool was generated by selecting 10 equally spaced values for the
1143 fixation rate f_j, and the corresponding phosphorous acquisition rates x_{1j} and N-

1144   recycling efficiencies λ_{2j} given by the trade-offs detailed above. All the other

1145   parameters were chosen to be the same for these 10 species.

1146

1147   For every given subset of the species pool, community assembly was simulated

1148   numerically by seeding the species present in that subset altogether and letting

1149   them grow until an equilibrium was reached.

1150

Because N-fixing species with large f_j can, under certain circumstances, have

negative consumptive impacts $C_{2j}(R_1, R_2)$, their presence leads to the net

enrichment of the N pool. This results in strong fixers facilitating non-fixers and

weak fixers (species that have small fixation rates f_j) through two mechanisms: 1) a

strong N-fixer can coexist at equilibrium with a weak fixer, supplying the latter with

limiting N, preventing the weak fixer from going extinct or enabling it to reach a

higher biomass than in the absence of the strong fixer, and 2) in a initially very N-

limited environment that prevents the weak-fixer from establishing, a strong N-fixer

can establish and lead to soil N accumulation. This makes establishment of the weak

fixer possible, but eventually leads the strong fixer to its own demise as the weak

fixer excludes it. This second mechanism is a pure effect of community assembly

that can generate facilitation-driven succession, as described in detail in Koffel et al.

(2018).