Modeling How Community Assembly Alters the Functioning of Ecosystems

Authors and Affiliations

Koffel, Thomas, W.K. Kellogg Biological Station, Michigan State University, 3700 E Gull Lake Dr., Hickory Corners, MI 49060. U.S.A. koffelth@msu.edu

Kremer, Colin T, [1] W.K. Kellogg Biological Station, Michigan State University, 3700 E Gull Lake Dr., Hickory Corners, MI 49060. U.S.A; [2] Department of Ecology and Evolutionary Biology, University of California Los Angeles, 612 Charle E. Young Drive East, PO Box 957246, Los Angeles, CA 90095-7246 USA. kremer@ucla.edu

Bannar-Martin, K. kbannarm@gmail.com

Ernest, S.K. Morgan, Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, FL 32611, USA.

Eisenhauer, Nico, [1] German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany; [2] Institute of Biology, Leipzig University, Deutscher Platz 5e, 04103 Leipzig, Germany. nico.eisenhauer@idiv.de

Roscher, Christiane, [1] Helmholtz Centre for Environmental Research (UFZ), Department Physiological Diversity, Permoserstrasse 15, 04318 Leipzig, Germany; [2] German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany. christiane.roscher@ufz.de

Cabral, Juliano Sarmento, Ecosystem Modeling, Center for Computational and Theoretical Biology (CCTB), University of Würzburg, Emil-Fischer-Str. 32, 97074 Würzburg, Germany. juliano.sarmento_cabral@uni-wuerzburg.de

Leibold, Mathew A., Department of Biology, University of Florida, Gainesville, FL 32611, USA. mleibold@ufl.edu

Abstract: Although the effects of species richness on ecosystem functioning have been

extensively studied, there is increased interest in understanding how community

assembly in general might alter the functioning of ecosystems. We focus on two

complementary approaches for evaluating how community assembly influences

ecosystem function (here, productivity). The first quantifies the relative importance of

complementarity and selection by contrasting monocultures with polycultures. The

second identifies the effects of species losses and/or gains relative to the baseline

polyculture, as well as the indirect effects on other species' productivity. We evaluate and

contrast these two approaches, using simulated communities structured by different,

known competition mechanisms, where species compete for different resources and

experience varying levels of environmental heterogeneity. We find that the metrics

provided by these approaches can jointly discriminate the mechanisms of competition

driving productivity. We then apply our methods to data from a long-term biodiversity-

ecosystem experiment (the Jena Experiment) and find that the data do not correspond to

any of the competition scenarios we modeled. We address two additional possible

complications: facilitation by nitrogen fixing plants, and non-equilibrium behavior during

community assembly, and find that a combination of resource competition and

facilitation by nitrogen fixing plants is the more likely explanation for the results

obtained at Jena.

Introduction:

Community assembly drives shifts in community structure (i.e., species richness and composition) which can have important consequences for the functioning of ecosystems. While the impact of changes in species richness on ecosystem function (commonly called biodiversity-ecosystem function, or BEF, effects; reviewed by Cardinale et al. 2012, Hooper et al. 2012, Tilman et al. 2012) are the best-studied, the importance of all of community assembly's impacts on community structure (referred to here as community assembly-functioning of ecosystem, or CAFE, effects) is increasingly recognized (e.g. Ptacnik et al. 2010, Thompson and Gonzalez 2016, Hodapp et al. 2016, Leibold et al. 2017, Bannar-Martin et al. 2018). Drivers of community assembly include species interactions, disturbance, dispersal, environmental heterogeneity, and landscape effects that can lead to changes in community composition, without necessarily affecting species richness (Brown et al. 2001). Community assembly is thus complex and depends on multiple factors and processes that may also consequently affect ecosystem function (e.g. Thompson and Gonzalez 2016, Hodapp et al. 2016, Leibold et al. 2017). Therefore, it is community assembly processes as a whole that ultimately link community composition, including species richness, to ecosystem attributes.

How can we study these community assembly effects on ecosystem function? One approach would be to comprehensively study the detailed interactions among the species involved and theoretically analyze these interactions to fully disentangle how community assembly affects ecosystem attributes. This approach, however, is logistically demanding and likely to give answers that are system specific. An alternative is to develop metrics that parse the effects of various components of community assembly on

ecosystem attributes. Useful measures should be general enough to allow cross-system

comparisons and simple enough that they reduce the full complexity of communities,

while remaining sufficiently connected to real community features to be informative. In

this paper we focus on two of several methods that provide complementary information,

the Complementarity-Selection method and the Price equation components (e.g. Kirwan

et al. 2009, Bell et al. 2009, Hector et al. 2009, Bannar-Martin et al. 2018).

The Complementarity-Selection method (originally called the 'additive

partitioning method') offers insights into the ecosystem function of polycultures

(communities) by comparing them with null models of hypothetical polycultures where

species' functions are proportional to their function in monoculture (Loreau and Hector

2001). Total ecosystem functioning is partitioned into the function arising from: (i)

selection effects (which are positive when polycultures are dominated by species that are

high functioning in monoculture), and (ii) complementarity effects (which are positive

when species contribute more function to polycultures than expected given their

functioning in monoculture, due to interactions like niche partitioning or facilitation).

Both of these effects can also be negative, if low-functioning species dominate or species

interactions reduce function. It is generally thought that selection effects occur due to

environmental filtering, which selects for more productive species that are better suited to

the local environment. Complementarity is typically related to niche partitioning that

results in enhanced coexistence and increased resource-use efficiency. This approach is

well-suited to the design of most BEF experiments, as they quantify the function

(productivity) of monocultures of species used to form higher-diversity polycultures.

However, this approach does not allow direct comparison of communities of varying

composition and structure.

Another approach for understanding how the ecosystem function of communities

(polycultures) relates to community assembly, the Price equation method, focuses

directly on comparing pairs of communities (Fox 2006). Here, differences in total

function between communities are related to changes in composition and individual

species' functions, as quantified by calculating 'Price components' (Fox 2006, Fox and

Kerr 2012, Bannar-Martin et al. 2018). Price components can be used flexibly to make

different types of inferences depending on choices about (i) which communities are

compared, and (ii) which of several arrangements of Price components are examined

(Bannar-Martin et al. 2018). For this paper, we focus on (i) comparing communities with

the most diverse species pool to communities with lower levels of richness, and (ii) using

a three-part Price component arrangement that separates the functional effects of species

gains ('SG'), species losses ('SL') and the differences in density/function among species

present in both communities being compared (the Context Dependent Effect or 'CDE').

In Bannar-Martin et al. (2018) these are called the Community Assembly components;

we use this terminology in the rest of this paper. Price equation approaches like the

Community Assembly analysis can be applied more generally than the Complementarity-

Selection method, and used to examine how <u>diverse</u> changes in community structure

(including species richness, taxonomic turnover, and species' abundances, function, or

traits; Fox & Harpole 2008) alter ecosystem functioning. However, their use is restricted

to ecosystem functions that can be calculated as the sum of contributions of their

constituent species (Fox 2006).

We study how these two approaches might be combined, and if jointly they can enhance

our understanding of the mechanisms underlying the functioning of communities. Our

ultimate goal is to better understand real-world communities. However, we begin by

applying these methods to predictions from a community model we constructed based on

resource-competition theory (Tilman 1982, Tilman et al. 1997, Schreiber and Tobiason

2003, Gross and Cardinale 2007, Hodapp et al. 2016, Koffel et al. 2016) that can emulate

typical BEF experiments. We applied the model to various assembly scenarios to

consider a wide range of possible ways resource competition may occur and affect

community assembly (including competition for complementary vs. antagonistic

resources with homogeneous or heterogeneous resource supply at the local scale). We

then apply the Complementarity-Selection approach (Loreau and Hector 2001) and the

Community Assembly approach (Bannar-Martin et al. 2018) to the results of the models

to assess if these methods can be used to distinguish among the various known assembly

scenarios. Next, turning from theoretical results, we consider data from the long-term

BEF experiment conducted in grasslands at Jena (Roscher et al. 2004, 2005; Weigelt et

al. 2010) and find that the data are not consistent with any of our initial assembly

scenarios. This led to a set of *a posteriori* hypotheses, which we identify and address

separately (consistent with recommendations for improving transparency and

reproducibility in biology; e.g. Ihle et al. 2017, Nichols et al. 2019). Specifically: in these

real-world communities there may be 1) inadequate time to achieve steady-state

competitive outcomes and/or 2) facilitation (e.g. due to the effects of nitrogen fixing

plants). We explore these possibilities by adapting our model, and find support for the

facilitation scenario. Put together, these results show how the joint use of the

Complementarity-Selection and Community Assembly approaches make it possible to

discriminate the mechanisms underlying the functioning of ecological communities.

Methods:

To model community assembly in simulated BEF experiments, we first

considered a flexible version of the basic resource competition model for two resources

as initially developed by MacArthur (1972) and further elaborated by Tilman (1982):

151
$$\frac{dB_j}{dt} = [g_j(R_1, R_2) - m_j]B_j$$

152   (1)

153
$$\frac{dR_i}{dt} = l_i(S_i - R_i) - \sum_{j=1}^{n} C_{ij}(R_1, R_2)B_j$$

This model describes the population dynamics of n species j (whose biomasses are

denoted B_j) that compete for two resources, R_1 and R_2. Here $g_j(R_1, R_2)$ and m_j

respectively determine species j's per capita growth and mortality rates, S_i is the supply

of resource R_i, which is diluted at rate l_i, and $C_{ij}(R_1, R_2)$ is the *per capita* consumptive

impact of B_j on R_i. Species j's interactions with the two resources can be divided in two

parts: (1) its requirements (through the *per capita* growth rate g_j) and; (2) its impacts on

the resources (through the *per capita* consumptive impacts C_{ij}).

Competitive scenarios: We considered a range of competitive scenarios, classified by the

types of resources involved, including resources that are 'strongly complementary'

(virtually 'essential', *sensu* Tilman 1982) vs. 'weakly complementary' and 'weakly

antagonistic' (these two cases correspond closely to 'substitutable' in Tilman's

terminology, the perfectly substitutable model has undesirable mathematical properties

for our purpose) vs. 'strongly antagonistic'. To implement this variation in resource types

we used a flexible form of the growth rate $g_j(R_1, R_2)$ that can generate all these scenarios

by tuning a single parameter α (Schreiber and Tobiason, 2003; See Appendix A).

Fig. 1 illustrates how each of these resource types affects trade-off relations

among minimum resource requirements for the two resources. Previous work has almost

exclusively focused on the strongly complementary scenario (e.g. Tilman et al. 1997,

Gross and Cardinale 2007, Hodapp et al. 2016); however, we found that different types of

competition lead to different outcomes of community assembly, associated with

distinctive signatures in terms of Complementarity-Selection effects and Community

Assembly components (see Box 1 and `Results' section).

We also considered how adding environmental heterogeneity at the local scale

affects the results obtained from the Complementarity-Selection (Loreau and Hector

2001) and Community Assembly components methods. We ran our model assuming two

different, spatially-implicit scenarios for the supply of resources over the local

community: homogeneous resource supply and heterogeneous resource supply. The

homogeneous environment was characterized by a constant influx of both resources (i.e.

a single 'supply point', *sensu* Tilman 1982). The second scenario examines a

heterogeneous environment, where species compete in an ecosystem characterized by

100 microhabitats, each with a separate supply point (see Fig. 1).

Community Assembly Outcomes:

To model the community assembly process, we viewed a BEF experiment as

having the following sequence of events: a) there is a filter on the species pool (taken as a

subset from the total species pool of the experiment), that is imposed by the researchers

via selective seeding and/or weeding, b) interactions among the remaining species lead to

some of the species going extinct, and c) there are density-dependent responses among

the remaining species that then determine ecosystem attributes.

For each resource competition scenario (complementary vs. antagonistic,

homogeneous vs. heterogeneous), we imitated these BEF conditions by generating

multiple individual species pools with different subsets of species from the global pool

that differ in their species richness and composition. In practice, we created a global pool

of ten species whose acquisition traits for the two resources are regularly spaced along a

generalist-specialist continuum defined by a trade-off between the two acquisition traits

that parametrize the growth rate $g_j(R_1, R_2)$ (Appendix A). Our species pool of ten

species yields 1023 possible subsets of this species pool, including the 10 different

monocultures and the full species pool. We then considered the equilibrium states of

communities initialized with all possible subsets of this global pool, arranged by their

starting richness (e.g., 10 one-species communities, 45 two-species communities, etc.).

This was performed using a global invasion analysis (Box 1), based on ZNGIs and their

envelopes (Koffel et al. 2016). In the homogeneous case (a single supply point), only one

or two species can subsist at equilibrium on two resources (Levin 1970) but more species

can coexist if supply points are heterogeneous. In the heterogeneous case, 100 supply

points were drawn from a multivariate normal distribution with the same mean as in the

homogeneous case and a variance of 0.5 (See Fig. 1). The equilibrium community state

was determined for each supply point. Because these equilibria for both the homogeneous

and heterogeneous cases are deterministic, only one or two species will ultimately persist

for any single supply point (more species can coexist for multiple supply points), and the

identities of coexisting species and their abundances are predetermined by their growth

functions and consumption vectors. Each resulting equilibrium community was

characterized by the identity of the species remaining after community assembly and their

equilibrium biomass. In the homogenous case, total ecosystem function is simply the sum

of biomass for all species in the equilibrium community. For the heterogeneous case,

equilibrium biomasses were obtained by averaging over the set of supply points, enabling

a more direct comparison with the homogeneous case.

We then consider key properties of the communities that emerge under each

resource scenario (across diversity levels and all initial compositions). These include the

final biomass of each species, which together determine the total biomass or function of

the ecosystem, and realized species richness (the final number of species present). This

information for each of the subsets was then used in the Complementarity-Selection and

Community Assembly calculations.

Complementarity-Selection analysis:

The outputs from our competition models serve as inputs to the Complementarity-

Selection approach (Loreau and Hector 2001). For the purposes of this calculation, we

assumed all species had equal 'initial' abundances. The decomposition divides total

function into the Selection effect (SE) and Complementarity effect (CE) for each of the

1023 communities. SE and CE were averaged across all possible species pools at each

level of introduced richness regardless of how many species actually existed at

equilibrium.

Community Assembly analysis:

No supplementary assumption was needed to apply the Community Assembly

components (Fox and Kerr 2012; Bannar-Martin et al. 2018). We compared the

community initialized with the entire pool of ten species (our baseline, *sensu* Bannar-

Martin et al. 2018) to each less diverse community. Using the Community Assembly

configuration of the Price equation as described in Bannar-Martin et al. (2018), we

calculated for each community the impact on ecosystem function of species lost (SL),

species gained (SG) and the changes in function of shared species (CDE) relative to the

final community obtained for the most diverse species pool. We then averaged each of

these components across all possible species pools within each level of introduced

richness.

Comparison with experimental studies:

To illustrate how our approach can provide insights about community assembly

processes driving patterns of community richness and the functioning of ecosystems in

empirical systems, we applied the same analyses to data from a well-studied,

representative BEF experiment conducted at Jena (Roscher et al. 2004; 2005; Weigelt et

al. 2016). Established in 2002, this experiment manipulated grassland communities,

controlling the seeded species richness (2, 4, 6, 8, and 16 species, as well as the full 60

species pool) and functional group representation (1 to 4 groups) of 66 plots; additional

small monocultures (3.5 x 3.5 m plots) were established for each species. Randomly

selected subplots were sampled in the spring and summer of each year: aboveground

biomass was clipped, sorted to species, dried, and weighed (see Roscher et al. 2004, 2005

for additional experimental details). Roscher et al. (2005) found evidence of positive

selection and complementarity effects across plots with 2 - 8 species, focusing on data

from May of the second year of the experiment (2003). Using data available from

Weigelt et al. (2010), we expanded this Selection-Complementarity analysis to cover

2003-2008 (focusing on spring data, as do Roscher et al. 2005), and augmented it by

applying the Community Assembly approach. Prior to analysis, we averaged species'

biomasses across replicate subplots within each experimental plot. The results provide an

empirical case study for comparison with our modeling exercise.

267   Complementarity-Selection results for each year are based on monoculture

biomass data collected during the corresponding year. Generally, monoculture biomasses

declined over the six-year horizon of this analysis (Marquard et al. 2013; declines in

polyculture yield were weaker). Consequently, there is a tendency for the magnitude of

Complementarity-Selection effects to grow more pronounced over time, due to their

mathematical definition (Loreau and Hector 2001). As the Community Assembly

calculations are independent of monoculture data, they were not similarly affected.

<u>Results:</u>

In these initial analyses, we studied eight different scenarios with our model by varying

the type of competition and the degree of resource heterogeneity as shown in Table S1

and Fig. 1. Different scenarios produced qualitatively distinct relationships between

realized species richness (at equilibrium) and initial species richness (Fig. 2A), although

ecosystem function (here total biomass or yield) was always a saturating function of

initial species richness (size of the species pool; Fig. 2B), These results were similar for

different combinations of resource type and/or environmental heterogeneity. Regarding

our two approaches for analyzing communities, we focus on qualitatively distinct

outcomes as follows:

Antagonistic resources:

287   Antagonistic resources, whether weak or strong, or supplied uniformly or

heterogeneously, produced similar results. First, the maximum yield of polycultures was

always greater than the maximum monoculture. Second the realized richness was always

two species or fewer. In these scenarios, the average complementarity effect (CE) was

always positive — setting aside the weakly antagonistic case with introduced richness of

two — whereas the average selection effect (SE) was always negative (Fig. 3A). Finally,

the average Community Assembly components always showed positive species gains

(SG) and negative species loss (SL) components, whereas the CDE was always small or

zero (Fig. 3B).

Strongly complementary resources in homogeneous environment:

298   In contrast to antagonistic resources, results varied substantially depending on just

how strongly resources are complementary and on the degree of environmental

heterogeneity. Starting with strongly complementary resources (almost equivalent to

essential resources), we found that maximum yield in polycultures did not differ from

that of the most productive monoculture. Both selection (SE) and complementarity

effects (CE) were positive, in contrast with antagonistic resources (Fig. 3A). The increase

in yield with diversity was largely due to an increase in the mean selection effect: as the

size of species pools grew, more productive competitors were more likely to be present.

The relationship with complementarity was unimodal and reflected the increased role of

complementarity on yield in small species pools where the better competitors were

absent. The effects of complementarity were also generally small. Community Assembly

components showed that effects of CDE were zero and that all of the effects involved

positive effects of SG and negative effects of SL (Fig. 3B). We also found that mean

realized species richness was a unimodal function of the size of the initial species pool in

the homogeneous scenario (but still limited to two or fewer species; Fig. 2.A and S1).

This means that the relationship between realized diversity and yield appears complex

(Figure S1).

Strongly complementary resources in heterogeneous environments:

The patterns differed from the homogenous environment results in several

respects. First, the maximum productivity was typically higher than that of the most

productive monoculture. Also, realized species richness was now positively related to the

size of the species pool. In contrast with the homogeneous case, the positive

complementarity effect (CE) persisted at high diversity levels (Fig. 3A), while species

gains (SG) had much smaller effects and CDE was positive, counteracting a similar SL

effect (Fig. 3B). We also examined an intermediate scenario, with smaller variance in

supply point heterogeneity, and found intermediate patterns between the homogenous

case and the full variance case: intermediate SE and CE, and CDE and SG of comparable

magnitudes (not shown).

 Weakly complementary resources:

With resources that were only weakly complementary, the increase in yield with

diversity was less pronounced. Both CE and SE results were similar to the

complementary case (Fig. 3A), except that environmental heterogeneity did not support a

sustained positive CE effect at high diversity levels.

Relations between realized species richness and biomass:

In each of our scenarios, we also examined how total biomass was related to the

realized species richness (the number of species coexisting at equilibrium). Over all

initial richness levels, there was generally a positive relationship between standing

biomass and realized species richness (Fig. S1). However, scenarios differed in the

strength of this relationship and in the structure of covariation between biomass and

realized species richness when the size of the species pool was accounted for. In most

scenarios (scenarios 1-6 and 8 in Table S1), we found that biomass was positively related

to realized species richness and that this was true both at the global scale of analysis and

within subsets of communities that differed in the size of the introduced species pool. In

contrast, when resources were strongly complementary and environmental heterogeneity

low (scenario 7), total biomass was negatively related to realized species richness for any

given level of initial richness (Fig. S2).

Mechanisms underlying relationships between competition scenarios and

Complementary-Selection and Community Assembly patterns

The striking differences between the antagonistic and complementary resource

competition scenarios captured by the Community Assembly components can be linked

to the influence of resource type on the outcomes of community assembly. In the

antagonistic case, a pair of extreme specialists dominates all the other species for any

supply point (Box 1, panels A, B). This strongly affects both the complementarity and

selection effects (CE and SE), and the Community Assembly components (SL, SG, and

CDE). Dominance by a pair of specialists leads to a realized richness that cannot exceed

two even in the heterogeneous case. In monoculture, each specialist strategy generally

performs worse than the intermediate generalist strategies, which explains why the

selection effect is negative. Yet, the two specialists together perform better than any other

set of one or two species, driving the positive CE pattern. When one of the two dominant

specialist strategies is absent (causing a negative SL component), it is replaced by a less

specialized species with lower function than the missing specialist (which results in a

positive SG component but |SL| > |SG|). The less specialized species competes more

strongly with the remaining specialist due to greater niche overlap, reducing the

performance of the remaining specialist, which creates the negative CDE.

In the complementary cases, there is usually a unique generalist species from the

full species pool that dominates for a given supply point (Box 1, panels C, D). This

dominant species is also the one that performs the best – better than any other set of one

or two species – for this supply point, which explains why selection is positive and

complementarity is weak and hump-shaped. The CDE of exactly zero in the

homogeneous case is a direct consequence of having a single dominant strategy: either

this dominant strategy is absent from a low diversity community, which will then

ultimately share no species with the high diversity community, or the dominant strategy

is present, producing an endpoint identical to the high diversity community. Finally, the

positive CDE in the heterogeneous case arises because in the absence of one or more of

the dominant strategies, other dominant species that still occur compensate for functional

losses. Such CDE is larger than the SG component on average because the functional

contributions of new species (able to join a community due to the loss of at least one

dominant species) are smaller than increases in function occurring through compensation

by the remaining dominants.

____

Box 1: Mechanisms

In this box, we give a mechanistic explanation of how different competitive scenarios

determine the identity and function of persistent species. In turn, these will determine the

sign and magnitude of the Community Assembly components that arise from comparing

communities assembled from high and low diversity species pools. We consider both

strongly antagonistic and complementary resource scenarios. We show a simplified

example where the full species pool only includes five species, three of which appear in

the low diversity subset. We focus on a homogenous environment (with one supply point;

black dot) in the antagonistic example and a heterogeneous environment (with three

supply points) in the complementary case.

These situations are represented and analyzed using the graphical approach to resource

competition (MacArthur 1972, Tilman 1982, Leibold 1995, Chase and Leibold 2003,

Koffel et al. 2016). We perform the invasion analysis by drawing the ZNGIs of the

species in the community all together. The species whose ZNGIs are entirely located

above other species' ZNGIs are systematically excluded (e.g. purple, orange, and yellow

in A). When the impact vectors of the remaining species (red and blue in A) are

combined with a supply point, we obtain the set of these species that actually end up

persisting in the equilibrium community (e.g. red and blue coexist in A). In a

heterogeneous environment (C and D), we perform the invasion analysis for each supply

point independently and average the productivity of the resulting persisting species to

obtain the equilibrium community.

In the antagonistic case, when all members of the full species pool are initially present

(A), the three intermediate species (purple, orange, yellow; dotted lines) are excluded by

the two extreme specialists (red and blue; solid lines), which coexist. However, if one of

these two specialists (e.g., blue) is randomly absent from a low-diversity community (B),

it is replaced by its closest functional equivalent among the intermediate species (here,

yellow). This species is not completely specialized on resource 2, so it does not perform

as well as the lost specialist. Due to weak competition with this intermediate species, the

performance of the remaining specialist (red) is lower than in the full species pool

situation.

In the complementary case, the most diverse species pool (C) experiences the exclusion

of the two most extreme species (orange and yellow), while the three intermediate species

(blue, red, and purple) coexist, as each one is locally adapted to a particular supply point.

If two of these intermediate species, which dominate the full species pool, are absent in a

smaller community (e.g., red and purple), an extreme species (e.g., yellow) can survive

and coexist with the remaining dominant species (blue). This extreme species (yellow)

only partially compensates for the loss of function that occurs due to the absence of the

purple species. Similarly, the remaining dominant species (blue) partially compensates

for the loss of function due to the absence of the red species by increasing in abundance.

Comparisons with experimental data:

Although most of our scenarios (with the exception of scenario 7) produce

broadly saturating BEF patterns, our results indicate that we can qualitatively

discriminate among most of these eight different community assembly scenarios (Table

S1) by combining the Community Assembly (Fox and Kerr 2012, Bannar-Martin et al.

2018) and the Complementarity-Selection (Loreau and Hector 2001) analyses. This

suggests that applying these methods to empirical data may offer insights into which

competitive scenarios drive assembly and give rise to empirically-obtained BEF

relationships (or at least provide evidence about which mechanisms may be ruled out). To

assess this, we analyzed data from a representative BEF experiment, conducted at Jena,

Germany, to determine if we could identify the resource competition mechanisms

underpinning the functioning of this grassland ecosystem. We used six years of data from

the Jena experiment, starting in 2003 (a year after the experiment was established), to

consider whether results were stable over time.

We found that the data from Jena did not match any of the scenarios shown in

Table S1. On the one hand, we see a divergence between increasingly negative selection

effects and increasingly positive complementarity effects from 2003 to 2008 (Fig. 4),

creating a pattern consistent with the antagonistic resource scenario using the

Complementarity-Selection approach (Fig. 3A, see also Marquard et al. 2009). On the

other hand, we see persistent trends in the Community Assembly components, including

positive CDE effects across the entire time span (Fig. 4), which were only observed in

our model scenarios given competition for heterogeneous, complementary resources (Fig.

3B).

Consequently, the analysis of Complementarity-Selection effects (Fig. 4) is

consistent with antagonistic resources, be it in homogeneous or heterogeneous supply

points (compare with Fig. 3A), whereas the Community Assembly components suggest a

scenario of complementary resources in the presence of environmental heterogeneity

(comparing Fig. 4 with Fig. 3B). This indicates, perhaps unsurprisingly, that the eight

simplistic scenarios we considered probably lack additional mechanisms involved in

regulating or modifying the effects of community assembly on ecosystem function (and

hence the metrics used to study it).

As a result, we considered other possible factors that might have affected

community assembly at Jena. One of these is that the empirically observed dynamics may

not have reached steady state as assumed by our models above, a hypothesis supported by

the directional changes in Complementarity-Selection effects across the first three to four

years of the experiment (Fig. 4). To address this possibility, we simulated the temporal

dynamics of community assembly for the heterogeneous, complementary resources

scenario (because previous work has mostly emphasized that this scenario is more likely

than others for plants; see Tilman 1982, Tilman et al. 1997, Gross and Cardinale 2007,

Hodapp et al. 2016). Starting from an initial condition where all species are seeded at an

equally low density, we let their populations grow and compete over time. When a

species' density dropped under a certain minimal threshold (set equal to 10^{-4}), we

considered it extinct and set its density equal to zero. We then extracted species densities

at representative time points and applied both partitions as previously described.

Although this exercise shows that there can be important transient dynamics in the

community assembly dynamics of BEF experiments (Fig. S4), none of the resulting

patterns are consistent with the patterns we observed in Jena (Fig. 4). In fact, the

qualitatively stable distribution of metric values for Jena data after the fourth year suggest

that the empirical system was near, if not at, equilibrium. Intriguingly, we found that the

dynamics in our model (Fig. S4) showed a strong qualitative shift in the relative

importance of CE and SE through time and a somewhat subtler shift from linear to

unimodal CDE and an increase in SG through time. However, the data from Jena show

no such trends.

We also considered that facilitation, perhaps due to the nitrogen (N) fixing effects

of legumes, may play an important role in assembly, and consequently affect ecosystem

functioning. To address this, we adapted the model of Koffel et al. (2018) that combines

strongly complementary resource competition with facilitation, and again applied

Complementarity-Selection and Community Assembly analyses (see Appendix B). With

high environmental heterogeneity and strong facilitation under high initial N-limitation,

our results (Fig. 5) correspond at least qualitatively to our observations at Jena. Both the

facilitation model and Jena data demonstrate greater complementarity effects (CE) in

comparison to selection (SE) and, at least for Jena's 2007 and 2008 data, a negative

(Compare Fig. 4 and Fig. 5A). This seems consistent with observations from the Jena

experiment that showed that N fixation rates by legumes, and thus potentially facilitation,

increased from 2004 to 2008 (Roscher et al. 2011). The Community Assembly

components also show similarities, with a substantial and humped CDE effect that

emerges from the model (Fig. 5B) also being seen in the Jena data (Fig. 4). The addition

of facilitation creates a combination of Complementarity-Selection effects and

Community Assembly components patterns that otherwise appear to point to divergent

resource competition models in the absence of facilitation.

To evaluate this hypothesis, we conducted separate Complementarity-Selection

and Community Assembly analyses for Jena plots that did and did not include N-fixing

plants (legumes) in their species pools (Fig. 6). Plots that contained legumes showed

results that qualitatively match the facilitation scenario (compare with Fig. 5) and

differed from plots without legumes as might have been predicted. However, plots that

lacked legumes exhibit patterns that are both qualitatively inconsistent with any of our

scenarios and relatively weak, perhaps due to subtler mechanisms. Further exploration of

these effects is beyond the scope of the present publication.

Discussion:

There are two major implications of our work. First, we find that the joint use of

the Community Assembly (Bannar-Martin et al. 2018) and the Complementarity-

Selection (Loreau and Hector 2001) approaches can be a powerful tool in inferring (or

excluding) mechanisms that drive community dynamics and consequently the functioning

of ecosystems. Second, we find that applying this joint approach to a real-world system

(the Jena BEF experiment) enabled us to rule out a series of purely competitive

community mechanisms – where using only a single approach would have led us to

inaccurate conclusions. This ultimately resulted in new hypotheses, and the conclusion

that facilitation by N-fixing plants is critical. This suggests that the joint evaluation of

Complementarity-Selection effects and Community Assembly components could be a

powerful tool for identifying plausible mechanisms that relate community assembly to

the functioning of ecosystems.

Validating the joint use of Complementarity-Selection and Community Assembly

approaches to interpret mechanisms underlying patterns in BEF studies:

518   Numerous studies in controlled and replicated conditions (such as those that

prevail in the large BEF literature) have shown that there is generally a positive and

directional effect between the size of the species pool with access to a community and the

function of at least some aspects of ecosystems (especially total biomass). Previous work

on relationships between species pool diversity and ecosystem function has largely

focused on two very general effects that can explain this pattern: selection (the species

pool, when large, is more likely to include highly competitive and highly productive

species) and complementarity (the species pool, when large, is more likely to include

species that coexist and whose production is less mutually inhibitory). To date,

theoretical explanations for these two general effects have focused on resource

competition and used models based on complementary resources (e.g., Tilman et al.

1997, Gross and Cardinale 2007, Leibold et al. 2017).

530   However, other competitive mechanisms may be involved, and there has been

little consideration of whether or not they would impart distinct complementarity and

selection effects. Our results show that indeed different mechanisms can produce similar

patterns of complementarity and selection, as the antagonistic resource and facilitation

scenarios both lead to increasing positive complementary effects (CE) and decreasing

negative selection effects (SE) with increased diversity (compare Fig. 3A with Fig. 5).

This means that attributing specific Complementarity-Selection patterns to particular

mechanisms may fall short. However, added insights from the Community Assembly

approach (Bannar-Martin et al. 2018) improved the situation, as these two scenarios

differed by the sign of the CDE and its relative magnitude compared to SG. Together, the

joint use of Community Assembly and Complementarity-Selection approaches provided

well-defined criteria by which we could distinguish between competition for

complementary vs. antagonistic resources as well as evaluate the role of environmental

heterogeneity and facilitation.

Challenges identifying mechanisms that drive community-ecosystem function relations:

Our study also reiterates the fact that we do not yet understand all the mechanisms

that drive community-ecosystem relationships. Patterns in the data from the Jena

experiment did not correspond to any of the eight scenarios we initially modeled, which

correspond to the classic scenarios of resource competition in community ecology, and

allowed us to reject all eight of these hypotheses. On the one hand, this shows how

powerful the joint application of Community Assembly and Complementarity-Selection

metrics can be. However, it also shows that the mechanisms governing the diversity and

function of communities must involve other processes besides simple resource

competition. This is not totally surprising since our theoretical models are caricatures of

how species interact in communities, where other factors including spatial structure,

grazers, pathogens, and mutualists may matter.

Further progress in this arena may require considering some of these other

scenarios, aided by Community Assembly and Complementarity-Selection approaches.

For example, we illustrated this by considering (1) non-equilibrium dynamics and (2)

facilitation by an N-fixer; the latter scenario qualitatively agreed with observed patterns

at Jena where other scenarios failed. This provides a new hypothesis, which may be

address by subsequent research – a critical step, as we note that a correspondence

between theory and observation is not definitive proof that a particular mechanism is

operating. For example, our model of facilitation was based on N-fixation, but results

similar to ours could arise from a broader range of facilitation mechanisms at play in BEF

experiments, perhaps related to other abiotic factors (Wright et al. 2017) or based on soil

organisms (Eisenhauer et al. 2012). Alternatively, other mechanisms besides facilitation

might be capable of generating similar patterns. Such additional mechanisms may play a

secondary role as revealed in the analysis of plots that did not contain legumes as shown

in Fig. 6. Ultimately, for Complementarity-Selection and Community Assembly

approaches to advance science, any given analysis and its results needs to be followed up

with additional research and experimentation to verify putative mechanisms.

Community assembly and the functioning of ecosystems outside of BEF designs:

The design of BEF experiments imposes constraints on the assembly of

communities that may not be present in real ecosystems. For example, species invasions,

disturbance and dispersal dynamics, environmental change, and stochastic processes are

all common in real communities (e.g. Thompson and Gonzalez 2016, Hodapp et al. 2016,

Leibold et al. 2017, Bannar-Martin et al. 2018). We have shown that Complementarity-

Selection and Community Assembly approaches offer useful insights when applied to

BEF-style scenarios and data that limit these kinds of effects. This utility may extend to

communities (and data) that differ from BEF designs, but requires careful thinking.

In particular, the Complementarity-Selection approach requires estimates of

ecosystem function in monocultures. These are readily available in most 'classic' BEF

experiments as a *pro-forma* aspect of their design. However, these are almost always

absent in other experiments that examine how community assembly affects ecosystem

functions (e.g. disturbance, invasions, extinctions, environmental change), let alone in

observational data. The logistical challenge is even greater in studies examining the

effects of environmental change on ecosystem function, as monoculture treatments would

be required for all the relevant environmental conditions. Additionally, many organisms

are not easily grown in monocultures due, for example, to dependencies on biotic

resources, mutualists, or movement. Thus, the scenarios where the Complementarity-

Selection approach can be used are limited.

In contrast, the Community Assembly approach can be calculated for any pair of

communities, without monoculture data. Given the design of BEF studies, we opted to

use the most diverse species pool as our baseline for comparisons, but this is not required

(see Bannar-Martin et al. 2018 for examples). While the Complementarity-Selection

approach is constrained by the requirement of monoculture data, it did appear to more

clearly distinguish between antagonistic and complementary resource scenarios. This

ability in part may rest in the definition of these metrics: both complementarity (CE) and

selection (SE) are affected by modest changes in the abundance of a species. In contrast,

of the three Community Assembly components, SL and SG are mostly influenced by the

actual loss or gain of species from a community; all effects of less extreme changes in

abundance are relegated to the CDE component alone. For this reason, while it is possible

to apply a Community Assembly approach in a manner directly analogous to

Complementarity-Selection (contrasting an observed polyculture with an idealized

polyculture where species perform proportional to their monocultures), the results are

somewhat less informative.

Our work shows that, while both methods exhibit their own advantages and

limitations, together they provide a more powerful means of diagnosing mechanisms.

Future work could focus on continuing to refine these metrics. This might include

developing analogs of Complementarity-Selection partitions that do not require

monoculture data. Alternatively, new or existing extensions of the Community Assembly

approach (e.g., 5-part Price equation; Fox & Kerr 2012; Bannar-Martin et al. 2018) could

be applied to these or other model scenarios, perhaps focusing on further dissecting the

CDE component. For example, while empirical applications of the trait-based version of

the Price equation (Fox & Harpole 2008) can be hampered by the availability of trait

data, in modeling exercises these traits are explicitly known (e.g., species' ZNGIs in our

scenarios). Further work could elucidate not only the mechanisms structuring ecosystem

function across communities of different diversity levels but relate ecosystem properties

to the functional composition of communities.

Implications for future work:

These encouraging results suggest a number of ways whereby the analysis of

community assembly and its effects on ecosystem function could be improved:

a) Most obviously, one could continue to propose and test other mechanisms, such

as plant-soil feedbacks and multi-trophic interactions, that could have produced

the patterns we observed in the Jena data, eventually potentially creating a

'periodic table' for how the Complementarity-Selection and Community

Assembly metrics map onto a diverse array of community assembly scenarios.

While there is some allure to this suggestion, there are a considerable number of

interacting mechanisms to test requiring careful thinking about how to construct

and interpret models (e.g., considering transient dynamics), or how to parse

varying contrasts in the data (e.g., considering facilitation by legumes).

b) Discriminating among the wide array of mechanisms driving community

assembly and its effects on ecosystem function likely requires moving beyond

simple dichotomous tests with one or two metrics since many alternative

models/mechanisms are capable of reproducing low-dimensional patterns. Here

we demonstrated how the addition of several carefully considered metrics could

enhance our understanding of these mechanisms, as well as our ability to

differentiate them. It is worth considering whether other additional metrics could

similarly advance understanding. The degree to which this is so would depend on

finding metrics that reveal distinct aspects of mechanisms and that, thus, are not

simply redundant with each other. It is also useful when the link between

mechanism and metrics is relatively transparent (as we explored in Box 1). More

complex relationships between mechanisms and metrics might require basic

modeling or experimental studies to help interpret the results. The increase in the

number of diagnostic metrics, however, may be necessary if theoretical models

649   increase in complexity by including missing processes or by assessing different

650   assumptions to increase inference resolution (e.g. Cabral et al. 2019).

651   c)   Another reason to consider additional metrics is that not all data will be suitable

652   for the same metrics. In our case, while the Complementarity-Selection approach

653   was particularly useful in discriminating between different types of resource

654   competition, the approach is constrained by the need for monoculture data. We

655   will need new techniques if we wish to apply these types of approaches to a wider

656   array of experimental conditions, with designs that do not closely target BEF

657   relations or that involve organisms where such monocultures are logistically

658   difficult (e.g. birds, migrating organisms, species-rich tropical communities).

659   d)   While we need to explore other metrics to use in studies other than BEF

660   experiments, it would still be interesting to apply the methods in this paper to

661   other BEF experiments. We suspect that this may likely show substantial variation

662   in results across locations, habitats, and organisms. For example, it would be

663   interesting to compare results at Jena with data from sites that are not as likely to

664   be nitrogen limited. Adequately considering such variation would require a far

665   more comprehensive theoretical and analytical effort than we could undertake in

666   this initial paper.

667   e)   Even more challenging is considering how the size of the species pool jointly

668   affects biodiversity, community composition, and ecosystem functioning under

669   other scenarios than those at play in classical BEF experiments. Numerous

670   challenges to BEF research have focused on the degree to which BEF experiments

671   may be unrealistic representations of how biodiversity and ecosystem function are

672 likely to interrelate because the experimental setup for community assembly in

673 BEF experiments is so artificial (Wardle 2016). We believe the Community

674 Assembly approach can shed light on other situations including arrivals and

675 removals of invasive species as well as environmental change (see Bannar-Martin

676 et al. 2018, Kremer et al. in prep). Linking these scenarios to explicit theoretical

677 models (as we did here by asking how resource types and environmental

678 heterogeneity affect a BEF experiment) represents an exciting avenue for future

679 devoted work.

681 *Software availability:*

682 An R package is available to facilitate future work (including analyzing other biodiversity

683 experiments) https://github.com/ctkremer/priceTools

685 *Acknowledgements:* This paper is a joint effort of the working group sCAFE and an

686 outcome of a workshop supported by sDiv, the Synthesis Centre of the German Centre

687 for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig (DFG FZT 118). The

688 Jena Experiment is funded by the German Research Foundation (FOR456/1451) with

689 additional support from the University of Jena and the Max Planck Society. MAL was

690 also supported by a research award from the Alexander Von Humboldt Foundation.

References

Bannar-Martin, K. H., Kremer, C. T., Ernest, S. K. M., Leibold, M. A., Auge, H., Chase,

J., Declerck, S. A. J., Eisenhauer, N., Harpole, S., Hillebrand, H., Isbell, F., Koffel, T.,

Larsen, S., Narwani, A., Petermann, J. S., Roscher, C., Cabral, J. S., and Supp, S. R.

(2018). Integrating community assembly and biodiversity to better understand ecosystem

function: the Community Assembly and the Functioning of Ecosystems (CAFE)

approach. *Ecology Letters*, 21:167–180.

Bell, T., Lilley, A. K., Hector, A., Schmid, B., King, L., and Newman, J. A. (2009). A

linear model method for biodiversity-ecosystem functioning experiments. *The American*

*Naturalis*t, 174(6):836–849.

Cabral, J.S., Wiegand, K. and Kreft, H., (2019). Interactions between eco-logical,

evolutionary, and environmental processes unveil complex dynamics of insular plant

diversity. *Journal of Biogeography*, 45, 1582-1597.

Cardinale, B.J., Duffy, J.E., Gonzalez, A., Hooper, D.U., Perrings, C., Venail, P., et al.

(2012). Biodiversity loss and its impact on humanity. *Nature*, 486, 59-67.

Chase, J. M. and Leibold, M. A. (2003). *Ecological Niches: Linking Classical and*

Contemporary Approaches. University of Chicago Press, Chicago.

Eisenhauer, N., Reich, P.B. and Scheu, S., 2012. Increasing plant diversity effects on

productivity with time due to delayed soil biota effects on plants. Basic and Applied

Ecology, 13(7), pp.571-578.

Fox, J.W. (2006). Using the Price equation to partition the effects of biodiversity loss on

ecosystem function. *Ecology*, 87, 2687-2696.

Fox, J.W., & Harpole, W.S. (2008). Revealing how species loss affects ecosystem

function: the trait-based Price equation partition. *Ecology*, 89, 269–279.

Fox, J.W., & Kerr, B. (2012). Analyzing the effects of species gain and loss on ecosystem

function using the extended Price equation partition. *Oikos*, 121, 290-298.

Gross, K., & Cardinale, B.J. (2007). Does species richness drive community production

or vice versa? Reconciling historical and contemporary paradigms in competitive

communities. *The American Naturalis*t., 170:207-220.

Hector, A., Bell, T., Connolly, J., Finn, J., Fox, J. Kirwan, L., Loreau, M., McLaren, J.,

Schmid, B., and Weigelt, A. (2009). The analysis of biodiversity experiments: from

pattern toward mechanism. pp. 94-104 in Naeem, S., Bunker, D.E., Hector, A., Loreau,

M., and Perrings, C. (eds.). Biodiversity, Ecosystem Functioning, and Human Wellbeing:

An Ecological and Economic Perspective. Oxford University Press. Oxford.

Hooper, D. U., Adair, E. C., Cardinale, B. J., Byrnes, J. E. K., Hungate, B. A., Matulich,

K. L., Gonzalez, A., Duffy, J. E., Gamfeldt, L., and O'Connor, M. I. (2012). A global

synthesis reveals biodiversity loss as a major driver of ecosystem change. *Nature*,

486(7401):105–108.

Ihle, M., Winnery, I.S., Kristalli, A., and Croucher, M. (2017). Striving for transparency

and credible research: practical guidelines for behavioral ecologists. *Behavioral Ecology,*

28:348-354.

Kirwan, L., Connolly, J., Finn, J. A., Brophy, C., Lüscher, A., Nyfeler, D., and Sebastià,

M.-T. (2009). Diversity-interaction modeling: estimating contributions of species

identities and interactions to ecosystem function. *Ecology*, 90(8):2032–2038.

Koffel, T., Daufresne, T., Massol, F., and Klausmeier, C. A. (2016). Geometrical

envelopes: extending graphical contemporary niche theory to communities and eco-

evolutionary dynamics. *Journal of Theoretical Biology*, 407:271–289.

Leibold, M. A. (1995). The niche concept revisited: mechanistic models and community

context. *Ecology*, 76(5):1371–1382.

Leibold, M. A., Chase, J. M., & Ernest, S. K. M. (2017). Community assembly and the

functioning of ecosystems: how metacommunity processes alter ecosystem attributes.

Ecology, 98, 909–919.

León, J. A. and Tumpson, D. B. (1975). Competition between two species for two

complementary or substitutable resources. *Journal of Theoretical Biology*, 50(1):185–

201.

Levin, S. A. (1970). Community equilibria and stability, and an extension of the

competitive exclusion principle. *The American Naturalist*, 104(939):413–423.

Loreau, M. & Hector, A. (2001). Partitioning selection and complementarity in

biodiversity experiments. *Nature*, 412, 72-76.

MacArthur, R. H. (1972). *Geographical Ecology: Patterns in the Distribution of Species*.

Harper & Row, New York.

Marquard, E., Weigelt, A., Temperton, V.M., Roscher, C., Schumacher J., Buchmann,

N., Fischer, M., Weisser, W.W., Schmid, B., (2009). Plant species richness and

functional composition drive overyielding in a six-year grassland experiment. *Ecology*

90(12), pp. 3290-3302.

Nichols, J.D., Kendall, W.L., Boomer, G.S. (2019) Accumulating evidence in ecology:

Once is not enough. *Ecology and Evolution* 2019; 00:1-14.

Ptacnik, R., Moorthi, S.D., & Hillebrand, H. (2010). *Hutchinson Reversed, or Why There*

Need to Be So Many Species. In G. Woodward (Ed.), Advances in Ecological Research

(Vol. 43, pp. 1–43). Burlington: Academic Press.

Roscher, C., Temperton, V.M., Scherer-Lorenzen, M., Schmitz, M., Schumacher, J.,

Schmid, B., Buchmann, N., Weisser, W.W. and Schulze, E.D., (2005). Overyielding in

experimental grassland communities–irrespective of species pool or spatial scale.

Ecology Letters, *8*(4), pp.419-429.

Roscher, C., Schumacher, J., Baade, J., Wilcke, W., Gleixner, G., Weisser, W.W.,

Schmid, B. and Schulze, E.D., (2004). The role of biodiversity for element cycling and

trophic interactions: an experimental approach in a grassland community. *Basic and*

Applied Ecology, *5*(2), pp.107-121.

Roscher, C., Thein, S., Weigelt, A., Temperton, V.M., Buchmann, N. and Schulze, E.D.,

2011. N 2 fixation and performance of 12 legume species in a 6-year grassland

biodiversity experiment. Plant and Soil, 341(1-2), pp.333-348.

Rychtecká, T., Lanta, V., Weiterová, I., and Lepš, J. (2014). Sown species richness and

realized diversity can influence functioning of plant communities differently.

Naturwissenschaften, 101(8):637–644.

Schreiber, S. J. and Tobiason, G. A. (2003). The evolution of resource use. *Journal of*

Mathematical Biology, 47:56–78.

Thompson, P.L., and Gonzalez, A. (2016). Ecosystem multifunctionality in

metacommunities. *Ecology*, doi: 10.1002/ecy.1502.

Tilman, D. (1982). *Resource Competition and Community Structure.*

Princeton University Press, Princeton, NJ.

Tilman, D., Lehman, C. L., and Thomson, K. T. (1997). Plant diversity and ecosystem

productivity: theoretical considerations. *Proceedings of the National Academy of*

Sciences (USA), 94:1857–1861.

Tilman, D., Reich, P. B., and Isbell, F. (2012). Biodiversity impacts ecosystem

productivity as much as resources, disturbance, or herbivory. *Proceedings of the National*

Academy of Sciences, 109(26), 10394-10397.

Wardle, D.A. (2016). Do experiments exploring plant diversity-ecosystem functioning

relationships inform how biodiversity loss impacts natural ecosystems? *J. Veg. Sci.*, 27,

646-653.

Weigelt, A., Marquard, E., Temperton, V.M., Roscher, C., Scherber, C., Mwangi, P.N.,

Felten, S., Buchmann, N., Schmid, B., Schulze, E.D. and Weisser, W.W. (2010). The

Jena Experiment: six years of data from a grassland biodiversity experiment. *Ecology*,

91(3), pp.930-931.

Weigelt, A., de Luca, E., Roscher, C., Temperton, V., Buchmann, N., Fischer, M., Scher-

Lorenzen, M., Schmid, B., Schulze, E.-D., Weisser, W.W., Guo, L., Meyer, S.T. (2016).

Collection of aboveground community and species-specific plant biomass from the Jena

Experiment (time series since 2002). PANGAEA.

https://doi.org/10.1594/PANGAEA.866358

Wright, A.J., Wardle, D.A., Callaway, R. and Gaxiola, A., 2017. The overlooked role of

facilitation in biodiversity experiments. Trends in Ecology & Evolution, 32(5), pp.383-

390.

 Figure Legends:

Figure 1. Zero Net Growth Isoclines (ZNGIs; colored, thick) and supply points (black

dots) for each of the eight resource scenarios. ZNGIs are curved upwards or downwards

when resources are complementary or antagonistic, respectively, and close to linear for

the two weakly interacting scenarios. Within each scenario, the positions of the ZNGIs of

the different species indicate where they lie along the generalist-specialist continuum:

generalists have symmetrical ZNGIs at equal distance from the two resources axes while

specialists on one resource have asymmetrical ZNGIs closer to that resource's axes. The

superimposed supply points help visualize environmental heterogeneity in each scenario.

In the homogeneous case (top row), we also show how the consumption by the single

species or pair of species (colored or gray arrows & points, respectively) that dominates

after community assembly decreases resource availability from the supply point to the

resource levels at equilibrium, which is the outcome of invasion analysis (See main text

and Box 1). Similar arrows could have been used to show the variation in outcome from

each supply point in the heterogeneous case (lower panels) but are omitted for clarity

purposes.

Figure 2. A) Realized richness obtained after species sorting (violin plots), averaged

across all possible species pools at each level of introduced richness (solid black line). In

the four antagonistic resource scenarios, realized richness quickly saturated at two

species. In the complementary resource scenarios, average realized richness is hump-

shaped in the homogeneous case, but increasing in the heterogeneous case. B) Total

community yield obtained after species sorting (violin plots), averaged across all possible

species pools at each level of introduced richness (solid black line); values are scaled

relative to the yield of the most productive monoculture (horizontal reference line). All

scenarios give saturating responses but in scenarios where resources are close to

substitutable (central panels), moving from monocultures to polycultures causes weaker

increases in yields.

Figure 3. A) Complementarity (CE) and selection effects (SE) vary in magnitude and

sign as a function of introduced richness, competitive scenario (columns), and supply

point heterogeneity (rows). Antagonistic resources lead to CE > 0 and SE < 0, while

complementary resources result in CE > 0 and SE > 0. B) Community Assembly

components method, comparing the most diverse polyculture (initial richness of 10)

against less diverse mixtures. CDE is nearly always zero or slightly negative, except

when resources are complementary and have a heterogeneous supply. For both A) and

B), violin plots and solid lines respectively indicate variability and average values across

all possible species pools at each level of introduced richness (as in Fig. 2). Violins are

jittered slightly horizontally for clarity.

Figure 4. Complementarity (CE) and selection (SE) effects (top row), and Community

Assembly components comparing the most diverse (60 species) community to less

diverse communities (bottom row) over successive years at the Jena Experiment, as a

function of introduced species richness (log scale). Early in the experiment, CE and SE

were both positive, and weakly related to diversity (as found by Roscher et al. 2005).

However, by the end of the period we examined, the CE effect was positive and increased

strongly with diversity, while the SE effect was negative and decreased with diversity.

Although there is some variation in the magnitude of the Community Assembly

components through time, at all lower diversity levels, CDE effects are consistently

positive, SG is negligible, and SL strongly contributes to declines in function. Trend lines

show Generalized Additive Models (GAMs) fit to the observed plot-level values,

including confidence bands; violins were jittered horizontally for visual clarity.

Figure 5: Model results after including nitrogen fixing plants that compete with non-

fixers, including Complementarity-Selection effects (top) and Community Assembly

components (bottom). Violin plots and solid lines respectively indicate variability and

average values across all possible species pools at each level of introduced richness (as in

Fig. 2).

Figure 6: Complementarity-Selection effects (upper panels) and Community Assembly

components (lower panels) for subsets of the Jena experiment that contained at least one

legume (LG, left panels) or none (right panels). Trend lines show Generalized Additive

Models (GAMs) fit to the observed plot-level values, including confidence bands.

Figure 1:

905
906

907   Figure 2*:*

908
909

Figure 3:

914   Figure 4:

918　　Figure 5:

923 Figure 6:

924
925
926
927

928   Figure Box 1: