Characterization of Primed Adaptation in the *Escherichia coli* type I-E CRISPR-Cas System

Anne M. Stringer[1], Lauren A. Cooper[2], Sujatha Kadaba[1], Shailab Shrestha[1] and Joseph T. Wade[1,2*]

[1]Wadsworth Center, New York State Department of Health, Albany, New York, USA

[2]Department of Biomedical Sciences, School of Public Health, University at Albany, Albany, New York, USA.

*Corresponding author:

joseph.wade@health.ny.gov

ABSTRACT

CRISPR-Cas systems are bacterial immune systems that target invading nucleic acid. The

hallmark of CRISPR-Cas systems is the CRISPR array, a genetic locus that includes short

sequences known as "spacers", that are derived from invading nucleic acid. Upon exposure to an

invading nucleic acid molecule, bacteria/archaea with functional CRISPR-Cas systems can add

new spacers to their CRISPR arrays in a process known as "adaptation". In type I CRISPR-Cas

systems, which represent the majority of CRISPR-Cas systems found in nature, adaptation can

occur by two mechanisms: naïve and primed. Here, we show that, for the archetypal type I-E

CRISPR-Cas system from *Escherichia coli*, primed adaptation occurs at least 1,000 times more

efficiently than naïve adaptation. By initiating primed adaptation on the *E. coli* chromosome, we

show that spacers can be acquired across distances of >100 kb from the initially targeted site, and

we identify multiple factors that influence the efficiency with which sequences are acquired as

new spacers. Thus, our data provide insight into the mechanism of primed adaptation.

INTRODUCTION

CRISPR-Cas systems are adaptive immune systems that are widespread throughout the bacterial

and archaeal kingdoms and protect cells from invading nucleic acid molecules. Although the

mechanistic details vary widely across different CRISPR-Cas systems, the basic mechanism is

shared across all systems (Wright et al., 2016). A short RNA, known as a CRISPR RNA

(crRNA) associates with a surveillance CRISPR-associated (Cas) protein, or complex of Cas

proteins. This protein-RNA complex binds to the invading nucleic acid molecule, at least in part

due to base-pairing between the crRNA and complementary sequence in the invading nucleic

acid (the "protospacer"). This leads to cleavage of the invading nucleic acid either by the

surveillance protein-RNA complex, or by an additional protein that is recruited to this complex.

The most abundant type of CRISPR-Cas system in nature is the type I system (Koonin et al.,

2017); the archetypal type I system is the type I-E system of *Escherichia coli*, which employs a

large complex of five Cas proteins (some proteins present in multiple copies), "Cascade", as the

surveillance complex, and recruits a separate nuclease protein, Cas3 (Brouns et al., 2008).

A hallmark of all CRISPR-Cas systems is the CRISPR array, a genetic locus consisting of

multiple units of short (~20-40 nt) "repeat" and "spacer" sequences (Wright et al., 2016). Within

a given CRISPR array, repeat sequences are identical, and each repeat is separated from the next

by a spacer. Spacer sequences are variable, and are derived from invading nucleic acid

molecules. The CRISPR array is typically transcribed as a single, long RNA, which is then

processed by a Cas protein to yield individual crRNAs. Each crRNA contains a single spacer

flanked by partial repeats, and the spacer sequence provides the base-pairing complementarity to

a protospacer in the nucleic acid target. For CRISPR-Cas systems targeting DNA, an additional

sequence in the DNA target is typically required for CRISPR-Cas immunity. This short sequence

(typically 2-4 bp), known as the protospacer-adjacent motif (PAM), is located close to the

protospacer, and serves as a binding site for one of the Cas proteins in the surveillance complex

(Leenay and Beisel, 2017).

A key feature of all CRISPR-Cas systems is their ability to acquire new immunity elements, in a

process known as adaptation (Jackson et al., 2017; Sternberg et al., 2016). During adaptation,

one or more spacer+repeat units are added to the end of a CRISPR array. For type I CRISPR-Cas

systems, there are two mechanisms of adaptation: naïve and primed. Naïve adaptation occurs

when an organism is invaded by a nucleic acid sequence that is not already matched by an

existing spacer. Naïve adaptation typically requires only the Cas1 and Cas2 proteins, and has

been proposed to occur by Cas1-2 acquisition of DNA fragments generated by resection of

double-strand breaks (Levy et al., 2015). By contrast, primed adaptation requires invasion of the

cell by a nucleic acid sequence for which there is already a partially or completely matching

spacer. The mechanism of primed adaptation is not well understood, but requires all Cas proteins

(Datsenko et al., 2012). Targeting of a protospacer by Cascade with a partially or fully

complementary spacer leads to recruitment of Cas3, which is both a nuclease and a helicase.

Primed adaptation likely involves translocation of Cas3 away from the protospacer-bound

Cascade (Dillard et al., 2018; Redding et al., 2015), with Cas3 generating the substrates for

Cas1-2 to integrate new spacers into the CRISPR array (Dillard et al., 2018; Künne et al., 2016;

Semenova et al., 2016). However, the mechanism by which Cas3 and/or other proteins generate

substrates for Cas1-2 is poorly understood (Musharova et al., 2017).

For the purposes of this study, we refer to the source of a spacer as the "pre-spacer" location, and we refer to the site of Cascade binding as the protospacer. The sequence context of spacers acquired during primed adaptation has a large impact on the efficiency with which a given sequence is acquired (Datsenko et al., 2012; Fineran et al., 2014; Li et al., 2014; Musharova et al., 2018, 2018; Rao et al., 2017; Richter et al., 2014; Savitskaya et al., 2013; Shmakov et al., 2014; Staals et al., 2016; Strotskaya et al., 2017; Swarts et al., 2012). Previous studies have shown that pre-spacers are located predominantly upstream of the protospacer, consistent with the unidirectional helicase activity of Cas3. For type I-F systems, pre-spacers are equally abundant on both the target and non-target strands (Richter et al., 2014; Staals et al., 2016), whereas for type I-B, I-C, and I-E systems, pre-spacers are located predominantly on the non-target strand (Datsenko et al., 2012; Fineran et al., 2014; Li et al., 2014; Rao et al., 2017; Savitskaya et al., 2013; Shmakov et al., 2014; Strotskaya et al., 2017; Swarts et al., 2012). For type I-F systems, pre-spacers are located over a ~5 kb region upstream of the protospacer (Staals et al., 2016), whereas for type I-E systems, acquisition has been reported over distances of up to ~10 kb (Strotskaya et al., 2017), although most studies involved plasmids of <10 kbp for which the distance of primed adaptation cannot be measured since it likely exceeds the size of the plasmid (Datsenko et al., 2012; Fineran et al., 2014; Savitskaya et al., 2013; Shmakov et al., 2014). For all type I systems tested to date, most pre-spacers are flanked by an optimal PAM sequence (Datsenko et al., 2012; Fineran et al., 2014; Li et al., 2014; Rao et al., 2017; Richter et al., 2014; Savitskaya et al., 2013; Shmakov et al., 2014; Staals et al., 2016; Strotskaya et al., 2017; Swarts et al., 2012); indeed, pre-spacers that are not immediately flanked by an optimal PAM sequence are often the result of (i) "slipping" errors, where an optimal PAM is located one

or two bases away from the expected location, suggesting incorrect processing of the DNA by

Cas1-2 (Li et al., 2014; Rao et al., 2017; Shmakov et al., 2014; Staals et al., 2016), or (ii)

"flipping" errors, where the PAM is on the opposite side and strand of the pre-spacer, suggesting

that Cas1-2 has incorrectly inserted the spacer into the CRISPR array in the reverse orientation

(Li et al., 2014; Rao et al., 2017; Shmakov et al., 2014; Staals et al., 2016).

Here, we further investigate the process of primed adaptation in the type I-E system of *E. coli*.

Previous studies have shown that primed adaptation is considerably more efficient than naïve

adaptation in type I-B, I-E and I-F systems (Datsenko et al., 2012; Li et al., 2014; Staals et al.,

2016). We quantify this difference for the *E. coli* system, showing that primed adaptation is at

least 1,000 times more efficient than naïve adaptation. We observe primed adaptation from the *E.*

coli chromosome over distances >100 kb. Using the rich source of pre-spacers inferred from

these data, we identify 7 features of pre-spacers that determine the efficiency with which they are

acquired, supporting and extending previous observations: (i) the DNA strand of the pre-spacer

relative to the protospacer (i.e. target vs non-target strand), (ii) the position of the pre-spacer

upstream or downstream of the protospacer, (iii) the presence and position of an AAG PAM

immediately adjacent to the pre-spacer, (iv) distance of the pre-spacer from the protospacer over

a ~100 kb region, (v) the presence of an AAG within the pre-spacer, (vi) whether or not the pre-

spacer is acquired from within ~200 nt of the protospacer, (vii) the sequence of positions 30-33

of the pre-spacer. Based on these factors, we propose a model of primed adaptation in which a

Cas3-containing complex travels large distances along the DNA from the protospacer, cutting

the DNA at or near every AAG, contributing to the generation of substrates for Cas1-2.

RESULTS AND DISCUSSION

Primed adaptation is >1,000-fold more efficient than naïve adaptation in *E. coli*

Studies of naïve adaptation in *E. coli* have involved high-level overexpression of Cas1 and Cas2

(Levy et al., 2015; Yosef et al., 2013). Hence, it is unclear how the efficiency of naïve adaptation

compares to that of primed adaptation. We previously used a highly sensitive, fluorescence-

based reporter (Amlinger et al., 2017) to precisely quantify primed adaptation in *E. coli* (Cooper

et al., 2018). For protospacers with mismatches in the PAM-proximal "seed" region, or extensive

mismatches in the PAM-distal region, we failed to detect adaptation, strongly suggesting that

naïve adaptation is extremely inefficient in these cells. To directly compare the efficiencies of

primed and naïve adaptation in genetically similar strains, we used a Δ*cas3* strain of *E. coli* in

which all other *cas* genes are constitutively expressed from their native locus. This strain also

contains a minimal CRISPR array (i.e. one repeat, and a leader sequence) associated with an out-

of-frame copy of *yfp*. Expansion of this artificial array by a single repeat/spacer unit puts *yfp* in

frame, leading to YFP fluorescence (Figure 1A). We then introduced a plasmid expressing *cas3*

from an arabinose-inducible promoter, or an equivalent empty vector. We also introduced a

plasmid expressing a crRNA that targets a protospacer on the same plasmid ("self-targeting

plasmid"; stp), or an equivalent empty vector. Thus, we were able to measure the level of

adaptation in four different strains: (i) $cas3^+$ $crRNA^+$, (ii) $cas3^-$ $crRNA^+$, (iii) $cas3^+$ $crRNA^-$, (iv)

$cas3^-$ $crRNA^-$. We detected robust adaptation in cells expressing *cas3* and containing the stp

($cas3^+$ $crRNA^+$). By contrast, we detected background levels of fluorescence in cells lacking

cas3 and/or the stp crRNA (Figure 1B; 0-2 YFP^+ cells from a total of ~80,000 each). Thus, we

detected robust primed adaptation, but no naïve adaptation in a genetically similar strain. We

cannot rule out the possibility that naïve adaptation occurs at a frequency below detection using

this highly sensitive assay; nonetheless, we can conclude that primed adaptation is at least 1,000-

fold more efficient than naïve adaptation in cells expressing *cas* genes at equivalent levels.

Studies of other type I systems suggest that naïve adaptation is similarly inefficient, with primed

adaptation being >500-fold more efficient than naïve adaptation in a type I-F system (Staals et

al., 2016), and naïve adaptation being undetectable in a type I-B system (Li et al., 2014). This

suggests that CRISPR immunity is extremely inefficient for prokaryotes with type I systems

when encountering an invading DNA molecule that is not already at least a partial match to an

existing spacer. Hence, CRISPR immunity in such a situation would likely only be effective in

the context of a large, clonal population of bacteria/archaea.

Mapping newly acquired spacers suggests primed adaptation can occur from chromosomal

sites

Previous studies have shown that the majority of pre-spacers acquired by primed adaptation in *E.*

coli are located on the non-target strand, and are associated with an AAG PAM (Datsenko et al.,

2012; Fineran et al., 2014; Savitskaya et al., 2013; Swarts et al., 2012). To determine whether the

same features are true for our experimental set-up, we induced primed adaptation using the stp,

and determined the location of pre-spacers by PCR-amplification and sequencing of the

expanded CRISPR arrays. As expected, the majority of acquired spacers mapped to the stp

(Figure 2A), and the distribution of pre-spacers on the stp was independent of the CRISPR array

in which the spacer was acquired (*E.* coli has two CRISPR arrays) (Figure 2B). 75.3% of pre-

spacers were located on the same DNA strand as the protospacer, and 94.4% of spacers were

associated with an AAG PAM. Thus, our data are consistent with earlier studies of primed

adaptation. Intriguingly, many pre-spacers did not map to the stp, suggesting another source for

the acquired spacers. We determined whether any of the unmapped pre-spacers map to the *cas3*-

expressing plasmid or to the chromosome. Many of the reads map uniquely to the *cas3*-

expressing plasmid (Figure 2C), although it is important to note that much of the sequence of this

plasmid is identical to that of the stp, and reads that match shared regions were mapped to the

stp. Unexpectedly, a small proportion of the reads mapped uniquely to the chromosome. We

speculated that pre-spacers mapping uniquely to the *cas3*-expressing plasmid or the chromosome

represent a second round of primed adaptation. This is possible because the strain we used

contains both the CRISPR-I and CRISPR-II arrays. Hence, even though we sequenced the first

acquired spacer in each array, this spacer may have been acquired as a result of primed

adaptation using a newly acquired spacer in the other array. Consistent with this hypothesis, pre-

spacers mapping uniquely to the chromosome were found adjacent to sequences shared with the

stp, i.e. the *araC* gene (Figure 2D), or the transcription terminators of rRNA loci. Thus, our data

indicate that primed adaptation from chromosomal pre-spacers can be readily detected using this

approach.

Primed adaptation occurs over a distance of >100 kb from the protospacer

Previous studies suggest that primed adaptation in *E. coli* occurs over relatively long distances,

exceeding the size of a typical plasmid (Strotskaya et al., 2017). Given that we observed primed

adaptation from chromosomal sites when targeting the stp with a crRNA, we reasoned that

targeting a chromosomal site with a crRNA would lead to extensive primed adaptation from the

chromosome, allowing us to infer the distance over which primed adaptation occurs. In two

independent experiments, we initiated primed adaptation from protospacers located (i)

immediately upstream of the *lacZ* gene, with the target strand of the protospacer on the plus

strand of the genome ("*lacZ*+" protospacer), and (ii) ~10 kb from the *lacZ* gene, inside the *mhpT*

gene, with the target strand of the protospacer on the minus strand of the genome ("*mhpT*-"

protospacer). Both protospacers have an AGG PAM, which can induce both interference and

primed adaptation (Cooper et al., 2018; Fineran et al., 2014; Musharova et al., 2019; Xue et al.,

2015). Moreover, we previously showed that targeting the *lacZ*+ protospacer with a crRNA

leads to robust association of Cascade (Cooper et al., 2018). Following induction of primed

adaptation for each of the two protospacers, we inferred the location of pre-spacers by PCR-

amplification and sequencing of the first acquired spacer in the expanded CRISPR-II array.

Thus, we observed robust primed adaptation on the chromosome, centered at each of the

protospacers (Figure 3). Consistent with primed adaptation studies using plasmids, the majority

of pre-spacers were located on the non-target strand, upstream of the protospacer (Figure 3).

Remarkably, we detected primed adaptation over a distance of >100 kb upstream of each of the

two protospacers, with the extent of primed adaptation decreasing as a function of distance from

the protospacer, albeit with considerable local variation (Figure 3). Of note, a recent study

described priming on the *E. coli* chromosome over a similar distance (Shiriaeva et al., 2019). Our

data strongly suggest that Cas3 can translocate >100 kb from the protospacer during primed

adaptation. Since the strain we used has both CRISPR arrays intact, it is possible that some

spacers acquired in CRISPR-II represent two rounds of primed adaptation, with the first round of

primed adaptation leading to spacer acquisition in CRISPR-I. However, we believe this is a rare

occurrence. There is an insertion element ~15 kb away from the *lacZ*+ protospacer that contains

many frequently used pre-spacers. This insertion element sequence is duplicated at several other

genomic locations, but we observed very few pre-spacers adjacent to the duplicated regions.

Hence, we conclude that very few spacers acquired in CRISPR-I from this insertion element led

to a second round of primed adaptation in CRISPR-II. Nonetheless, even if a substantial

proportion of the pre-spacers we observed result from two rounds of primed adaptation, a single

round of primed adaptation must still be able to occur over a distance of >50 kb.

The large distance over which we observed primed adaptation explains why the efficiency of

primed adaptation from a plasmid does not appear to decrease as a function of distance from the

protospacer; presumably, Cas3 translocates many times around the circular plasmid. It also

suggests that primed adaptation would facilitate acquisition of spacers from any region of even

large bacteriophage genomes or plasmid, which may increase the effectiveness of the immune

response. The large distance over which primed adaptation occurs on the chromosome also

means that there are many more unique pre-spacers than used during primed adaptation from a

plasmid. We reasoned that this much larger set of pre-spacers would allow for a more in-depth

analysis of the pre-spacer features that influence the efficiency of primed adaptation.

Off-target Cascade binding sites are not associated with primed adaptation

We previously showed that Cascade association with a DNA target requires only limited base-

pairing between the crRNA spacer and the protospacer. Hence, Cascade binds to many

chromosomal sites that typically have between 5 and 10 nt matches to the 5' end of a crRNA

spacer, immediately flanked by an optimal (AAG) PAM (Cooper et al., 2018). When targeting

the *lacZ+* protospacer with a crRNA, we observed Cascade binding to ~100 off-target sites

(Cooper et al., 2018). We determined the level of primed adaptation from the non-target strand in

the 10 kb upstream each of the 76 off-target sites associated with a seed sequence resembling

that of the on-target *lacZ+* protospacer. The level of Cascade association and the local pre-spacer

usage frequency were not significantly correlated (Figure 4; Spearman's Correlation Test, two-

tailed, $p = 0.72$), with pre-spacer usage near off-target sites typically being due to duplicated

sequences with identical copies close to the *lacZ+* protospacer. Moreover, the local pre-spacer

usage frequency in regions adjacent to off-target Cascade binding sites was not significantly

higher than that of 1000 randomly selected genomic locations (Mann-Whitney U Test $p = 0.054$).

Thus, our data suggest that off-target Cascade binding events do not lead to primed adaptation,

consistent with our previous observation that extensive PAM-distal mismatches between a

crRNA spacer and its cognate protospacer prevent primed adaptation (Cooper et al., 2018).

Uneven distribution of chromosomal pre-spacers relative to the protospacer

The vast majority of pre-spacers we identified are 33 nt long (98.6% for both the *lacZ+* and

mhpT- protospacers; note that pre-spacers include the last base of the PAM, which is known to

be added to the CRISPR array by Cas1-2; (Goren et al., 2012)). Unless specifically stated, the

analyses described below focus exclusively on the 33 nt pre-spacers. We also excluded genomic

regions covered by insertion elements, repetitive sequences that confound alignment of DNA

sequence reads. Lastly, since the independent replicate datasets for each protospacer were highly

similar (Figure 5A + B), all analyses described below use datasets where the two replicates were

combined.

Although the majority of pre-spacers were located on the non-target strand, upstream of the

protospacers, we also observed a low level of primed adaptation on the target strand, upstream of

the protospacer (Figure 3 + 6A). As for pre-spacers on the non-target strand, the frequency of

pre-spacer usage on the target-strand decreased as a function of distance from the protospacer,

albeit with considerable local variation. Strikingly, the fraction of pre-spacers on the target

strand, upstream of the protospacer, was ~4-fold higher for the *mhpT-* protospacer than for the

lacZ+ protospacer, suggesting that the protospacer sequence context determines the bias in

primed adaptation towards the non-target strand (Figure 3 + 6A). Since the two protospacers we

used are only ~10 kb apart, and are in opposite DNA orientations, many of the *target* strand pre-

spacers for the *mhpT-* protospacer are from the same genomic region and orientation as *non-*

target strand pre-spacers for the *lacZ+* protospacer (pink box in Figure 3). Indeed, we observed

very similar profiles of pre-spacer location and abundance across this region for the two datasets

(Figure 6B), indicating that primed adaptation on the target strand likely occurs by the same

mechanism as primed adaptation on the non-target strand.

The strong bias in pre-spacer usage to the region on the non-target strand, upstream of the

protospacer, presumably reflects a bias in the direction of Cas3 translocation away. We propose

that Cas3 translocation is strongly biased to the non-target strand because of the inherent

asymmetry in the Cascade-bound protospacer. In type I-F systems, the only characterized type I

systems where Cas3 translocation appears to occur with equal efficiency on both the target and

non-target strands, Cas2 is fused to Cas3, such that there are two Cas3 subunits in the Cas1-2-3

complex (Fagerlund et al., 2017). We suggest that this allows translocation of Cas2-3 in either

strand with roughly equal efficiency, presumably using a different Cas3 subunit for each of the

two strands. However, there are likely to be other factors that influence the degree to which Cas3

selects strand, since our data show that the ratio of target:non-target strand usage varies between

protospacers (Figure 6A).

Acquisition of spacers with non-AAG PAMs is frequently due to slipping

We examined all pre-spacers associated with the *lacZ+* and *mhpT-* protospacers on the non-target strand, in the 10 kb region upstream of the protospacer. 95.0% and 96.0% of pre-spacer usage was associated with an AAG PAM for the *lacZ+* and *mhpT-* protospacers, respectively. While the large majority of pre-spacer usage is associated with an AAG PAM, we were interested to determine the basis for selecting pre-spacers with a non-AAG PAM. Previous studies of type I-E, type I-B, type I-C and type I-F CRISPR-Cas systems have shown that pre-spacers with non-canonical PAMs can be acquired due to a phenomenon known as "slipping", whereby the pre-spacer with a non-AAG PAM is positioned one or two nucleotides from a canonical, AAG-associated pre-spacer (Li et al., 2017; Rao et al., 2017; Shmakov et al., 2014; Staals et al., 2016). To determine the frequency of slipping in our data, we calculated all pairwise distances for non-AAG pre-spacers to AAG pre-spacers on the non-target strand, in the 10 kb upstream of each of the protospacers. We observed a strong enrichment of pre-spacers that could be assigned to "-1", "+1", or "+2" slips (Figure 7A + B), with +1 slips being the most frequent, accounting for 37% of all non-AAG pre-spacer usage. Our data indicate that together, +1 slips and -1 slips represent the majority (60%) of primed spacer acquisition on the non-target strand for pre-spacers not associated with an AAG PAM. The remaining pre-spacers that lack an AAG PAM and are not associated with slipping events typically have suboptimal PAMs, with the third base of the PAM strongly enriched for G, and the second base moderately enriched for A or C (Figure 7C). Thus, our data suggest that while the specificity for an AAG PAM is high during primed adaptation, there is a hierarchy of selectivity within the PAM, with the third position being the most important, and the first position being the least important.

Analysis of slipping in a type I-F CRISPR-Cas system demonstrated that slipped pre-spacers are

associated with PAMs that lead to efficient primed adaptation (Jackson et al., 2019). Our data

indicate that the most common form of slipping is a +1 slip, which would lead to an AGN PAM

(Figure 7B). The next most common form of slipping is a -1 slip, which would lead to a NAA

PAM (Figure 7B). Previous studies of PAM sequences that lead to interference and/or primed

adaptation indicate that of all the possible AGN and NAA PAMs, only AGT, and possibly CAA,

fail to lead to interference and/or primed adaptation (Musharova et al., 2019; Xue et al., 2015).

Thus, slipping in the *E. coli* CRISPR-Cas system likely leads to acquisition of a functional

spacer in most cases.

Previous studies of naïve adaptation showed only ~25% of pre-spacer usage is associated with

AAG PAMs (Levy et al., 2015; Yosef et al., 2013), in contrast to the >95% observed for primed

adaptation in our study and other studies (Savitskaya et al., 2013). One possible explanation is

that the high level of Cas1-2 overexpression required to detect naïve adaptation causes a

reduction in PAM specificity. An alternative hypothesis is that Cas1-2 association with the

translocating Cas3-containing complex during priming leads to an increase in specificity for

AAG; Cas3 cuts DNA with some sequence specificity (Künne et al., 2016), although this low

level of specificity alone is unlikely to account for the large difference in PAM specificity

between naïve and primed adaptation. Cas8e, a subunit of Cascade, binds specifically to AAG

PAMs. It is possible that Cas8e translocates with Cas3 and facilitates nicking at AAG sequences.

In support of this idea, Cas8 is fused to Cas3 in some bacterial species (Westra et al., 2012).

Moreover, Cas8e associates only weakly with the rest of the Cascade complex, and frequently

dissociates from the complex (Jore et al., 2011), in particular when there are suboptimal bases in

the PAM or in the PAM-proximal region of the protospacer (Jung et al., 2017). However, *in vitro*

reconstitution of Cas3 translocation for the *Thermobifida fusca* Cas3 suggests that translocating

Cas3 does not associate with Cas8e (Dillard et al., 2018).

Acquisition of spacers from the target strand is frequently due to flipping, often following a

slipping event

We examined all pre-spacers associated with the *lacZ+* and *mhpT-* protospacers on the target

strand, in the 10 kb downstream of the protospacer, i.e. on the opposite strand to where the most

frequently used pre-spacers are located. 2.4% and 2.1% of pre-spacer usage within 10 kb of the

protospacer, on this side (i.e. upstream of the protospacer on the non-target strand, or

downstream of the protospacer on the target strand) is on the target strand for the *lacZ+* and

mhpT- protospacers, respectively. While the large majority of pre-spacer usage is on the non-

target strand, we were interested to determine the basis for selecting pre-spacers from the target

strand. Previous studies of type I-E, type I-B, type I-C and type I-F CRISPR-Cas systems have

shown that pre-spacers with non-canonical PAMs can be acquired due to a phenomenon known

as "flipping", whereby a pre-spacer is selected from the non-target strand but then inserted into

the CRISPR array in the opposite orientation (Figure 8A) (Li et al., 2017; Rao et al., 2017;

Shmakov et al., 2014; Staals et al., 2016). To determine the frequency of flipping in our data, we

calculated all pairwise distances for target-strand pre-spacers to AAG pre-spacers on the non-

target strand, in the 10 kb upstream of each of the protospacers. We reasoned that a distance of

32 nt would represent a flipping event involving a canonical pre-spacer with an AAG PAM. We

observed an enrichment of distances that support flipping events involving spacers with a -1, +1,

or +2 nt slip, in addition to flipping events involving canonical (i.e. no slip) pre-spacers (Figure

8B). Thus, relative to the frequency of pre-spacer usage for the canonical pre-spacers, flipped

pre-spacers with an additional slip occur more often. We presume that pre-spacers from the non-

target strand with an AAG PAM are unlikely to be inserted into the CRISPR array in the

incorrect orientation, whereas pre-spacers generated by slipping events are more likely to be

flipped because they have a suboptimal PAM.

Pre-spacers with non-canonical lengths are often associated with slipping

While the large majority of pre-spacers are 33 nt long, we were interested to determine the basis

for selecting pre-spacers with non-canonical lengths. 0.3% of both *lacZ+* and *mhpT-* pre-spacer

usage was associated with pre-spacers that are 32 nt long, and 1.1% of pre-spacer usage for both

lacZ+ and *mhpT-* pre-spacers was associated with pre-spacers that are 34 nt long. We chose to

focus on the region <10 kb upstream of the protospacer, on the non-target strand, where the

usage of pre-spacers of non-canonical lengths is highest (89.8% and 82.9% of all non-canonical

length pre-spacers for *lacZ+* and *mhpT-* protospacers, respectively). The majority of 32 nt pre-

spacer usage for both *lacZ+* and *mhpT-* was associated with AAG PAMs; hence, these pre-

spacers are equivalent to efficiently used 33 nt pre-spacers, but lack the most PAM-distal base.

These AAG-associated pre-spacers account for 87.7% (lacZ+) and 95.7% (*mhpT-*) of all 32 nt

pre-spacer usage. Similarly, the majority of 34 nt pre-spacer usage was associated with AAG

PAMs; hence, these pre-spacers are equivalent to efficiently used 33 nt pre-spacers, but with an

additional base at the PAM-distal end. These AAG-associated pre-spacers account for 79.7%

(*lacZ+*) and 86.7% (*mhpT-*) of all 34 nt pre-spacer usage. There is also a significant enrichment

for 32 nt pre-spacer usage associated with +1 slips, and for 34 nt pre-spacer usage associated

with -1 slips (Figure 9), indicating that slipping sometimes involves maintaining the PAM-distal

boundary while either shortening or lengthening the spacer. The equivalent observation has been

made for type I-B, type I-C and type I-F CRISPR-Cas systems (Li et al., 2017; Rao et al., 2017;

Staals et al., 2016).

The efficiency of primed adaptation is reduced for pre-spacers within 200 bp of the

protospacer

To determine if the pattern of pre-spacer usage is dependent upon the location of the protospacer,

we initiated primed adaptation by targeting a location ~10 kb downstream of the *lacZ+*

protospacer, within the *codA* gene; we refer to this as the *codA+* protospacer. Replicate datasets

for *codA+* were highly correlated (Figure 5C); hence we combined the two datasets for further

analysis. As expected, we detected pre-spacer usage over a ~100 kb distance from the *codA+*

protospacer, with the majority of pre-spacer positions overlapping those used when targeting the

lacZ+ protospacer (Figure 10A). We compared the frequency of pre-spacer usage for pre-spacers

<10 kb upstream of the *lacZ+* protospacer on the non-target strand when targeting either the

lacZ+ or *codA+* protospacer. In almost all cases, the relative usage of pre-spacers was similar for

each of the protospacers targeted (Figure 10B), strongly suggesting that pre-spacer usage

frequency is an inherent property of the pre-spacer, and is not related to the location of the

protospacer. However, the three pre-spacers located within 200 nt of the *lacZ+* protospacer were

used proportionally less frequently when targeting the *lacZ+* protospacer than when targeting the

codA+ protospacer (Figure 10B). Thus, our data suggest that pre-spacers very close to the

protospacer are used inefficiently. Biochemical studies of Cas3 suggested that Cas3 generates a

single-stranded region adjacent to the Cascade-bound protospacer of <300 nt (Redding et al.,

2015) or <500 nt (Dillard et al., 2018), for Cas3 from *E. coli* or *T. fusca*, respectively. We

propose that Cas3 generates a single-stranded region of a similar length *in vivo*, and that no

spacers can be acquired from the single-stranded DNA. Our data suggest that in the region <200

bp from the protospacer, the relative efficiency of pre-spacer usage frequency increases as a

function of increasing distance from the protospacer (Figure 10B). Hence, we propose that Cas3

transitions stochastically between a state where it nicks DNA frequently, generating an extended

region of single-stranded DNA, and a state where it nicks DNA infrequently, with the average

distance traveled until the transition being 100-200 bp.

The efficiency of primed adaptation is reduced for pre-spacers with an internal AAG

It is well established that the frequency of pre-spacer usage varies considerably between pre-

spacer sequences (Musharova et al., 2017, 2018; Savitskaya et al., 2013). However, the basis for

this variability is poorly understood. As described above, the frequency of pre-spacer usage

decreases globally as a function of increasing distance from the protospacer (Figure 3 + 11A+B).

We fitted these data to an exponential decay model, relating pre-spacer usage to distance from

the protospacer. We then selected "enriched" pre-spacers that have a >4-fold higher usage

frequency than predicted by the model, and "de-enriched" pre-spacers that have a >4-fold lower

usage frequency than predicted by the model. We determined the frequency of all tetranucleotide

and trinucleotide sequences within the sets of enriched and de-enriched pre-spacers (Figure 11C-

D). Almost all tetranucleotide and trinucleotide sequences were found at similar frequencies

within the enriched and de-enriched pre-spacers. However, AAG-containing pre-spacers were

found far less frequently in enriched pre-spacers than in de-enriched pre-spacers. We compared

the frequency of pre-spacer usage for AAG-containing and non-AAG-containing pre-spacers

(Figure 11A-B). The frequency was, on average, ~3.5-fold higher for non-AAG-containing pre-

spacers than for AAG-containing pre-spacers. We conclude that the presence of an AAG within

a pre-spacer substantially reduces the frequency with which that pre-spacer is selected during

primed adaptation.

To experimentally test the effect of an AAG within a pre-spacer, we generated a second self-

targeting plasmid, "stp2", that includes a site for easy introduction of potential pre-spacers

flanked by an AAG PAM (see Materials and Methods for a detailed description of the plasmid

and assessment of its ability to cause primed adaptation). The introduced pre-spacer in

unmodified stp2 has AAA at positions 19-21. We also constructed a derivative of stp2 in which

the introduced pre-spacer has been modified to contain AAG rather than AAA at positions 19-

21. We refer to this mutant stp2 plasmid as stp2-mut1 (Figure 12A). We initiated primed

adaptation in cells containing either stp2 or stp2-mut1. We determined the location of pre-

spacers for each of the two plasmids by PCR-amplification and sequencing of the expanded

CRISPR arrays. We compared the relative usage frequency for each pre-spacer on the non-target

strand that is associated with an AAG PAM (these represent the most frequently used pre-

spacers; Figure 2A) for each of the two plasmids. As expected, pre-spacers with the same

sequence were used at the same frequencies for each plasmid (Figure 12B). By contrast, the

AAG-containing pre-spacer in stp2-mut1 was used at a substantially lower frequency than the

equivalent AAA-containing pre-spacer in wild-type stp2 (Figure 12B). Thus, our experimental

data are consistent with the bioinformatic analysis, and show that pre-spacers with an internal

AAG are used at a reduced frequency during primed adaptation. We note also that a recent,

independent study reached a similar conclusion about the impact of AAG sequences within pre-

spacers (Musharova et al., 2018).

A previous study mapped pre-spacers for naïve adaptation in a strain of *E. coli* that is not able to

undergo primed adaptation (Yosef et al., 2013). The authors sequenced 934,202 newly acquired

spacers that came from the chromosome, covering 84,951 pre-spacer sites. Although the

frequency of pre-spacers with AAG PAMs is considerably lower for naïve adaptation than

primed adaptation, many pre-spacers are associated with an AAG PAM (Yosef et al., 2013).

Using these data, we determined the frequency of usage for all potential chromosomal pre-

spacers associated with an AAG PAM. 18,872 sequences were identified as pre-spacers at least

once, whereas 108,207 sequences were not. We then compared the number of AAG-containing

(i.e. an internal AAG) pre-spacers within each of these groups to determine whether the impact

of an internal AAG is limited to primed adaptation. Although the proportion of AAG-containing

pre-spacers was similar for each group, a significantly larger fraction (38.8%) of the unused pre-

spacers had an internal AAG than the fraction of used pre-spacers (34.4%; Chi Squared Test with

Yates' Continuity Correction $p < 1e^{-7}$). As a control, we analyzed the proportion of AGA-

containing pre-spacers for the same groups and found no significant difference (Chi Squared

Test with Yates' Continuity Correction $p = 0.08$). We then divided the 18,872 used pre-spacer

sequences into AAG-containing and non-containing groups and compared the usage frequencies

for each group. While the distribution of usage frequencies was similar between the two groups

(Figure 12C), it was slightly and significantly higher for the AAG-lacking group (Mann Whitney

U Test $p = 2.5e^{-4}$). We saw no significant difference when comparing AGA-containing and

AGA-lacking pre-spacers (Mann Whitney U Test $p = 0.63$). We conclude that AAG-containing

pre-spacers are acquired at a lower efficiency during naïve adaptation. However, the magnitude

of the effect appears very small, in contrast to the large effect we observed for primed adaptation

(Figures 11 + 12B). Consistent with this, a previous study did not detect an effect of AAG within

pre-spacers on the efficiency of naïve adaptation (Musharova et al., 2018).

Since AAG is the PAM associated with >95% of pre-spacer usage during primed adaptation, we

presume that the impact of AAGs within pre-spacers is connected to the selection of pre-spacers

with AAG PAMs. One possible explanation for the reduced frequency of usage for AAG-

containing pre-spacers is that multiple Cas3-containing complexes could translocate from a

single Cascade-bound protospacer, such that stable association of a Cas3-containing complex

with an AAG PAM sequence prevents association of a second Cas3-containing complex with a

nearby AAG PAM. However, this would not explain the size of the decrease in pre-spacer usage

frequency associated with an internal AAG (3.5-fold), or its impact over a >100 kb region; for

the effect of an internal AAG to be explained by competition within Cas3-continaing complexes,

most of the >1,000 AAG sequences within 100 kb of the protospacer would need to be occupied

at any given time. A more likely explanation is that AAG sequences are frequently nicked, which

would represent a semi-stable change in the DNA that can be caused by a transiently positioned

Cas3-containing complex. Recent studies suggest that pre-spacers are nicked within the PAM

during primed adaptation (Musharova et al., 2017; Shiriaeva et al., 2019). We propose that the

Cas3-containing complex nicks all AAG sequences that it encounters as it translocates away

from the Cascade-bound protospacer (after transitioning from the highly active state within 200

nt of the protospacer), and that pre-spacers with an internal nick cannot be used as substrates for

adaptation by Cas1-2.

Intriguingly, the impact of an AAG within a pre-spacer appears to diminish as the AAG is

positioned further from the PAM (Figure 13A) (Musharova et al., 2018). Based on our nicking

model, we hypothesized that closely positioned pairs of AAG sequences would lead to closely

spaced nicks, and hence single-stranded DNA gaps that may then prevent the ability of these

sequences to be integrated into a CRISPR array. To test this hypothesis, we repeated the stp2

primed adaptation assay described above (Figure 12B) using a derivative of the stp2-mut1

plasmid where the sequence between the PAM and the internal AAG of the modified pre-spacer

was changed to a more G/C-rich sequence with a higher melting temperature. We reasoned that

this would reduce the propensity for the DNA between the PAM and the internal AAG to

become single-stranded in the event that the PAM AAG and the internal AAG were both nicked.

We refer to this plasmid as stp2-mut2 (Figure 12A). We also constructed an equivalent plasmid,

stp2-mut3, where the AAG within the pre-spacer was changed to an AAA (Figure 12A).

Although the relative usage frequency of the modified pre-spacer was higher for stp2-mut2 than

for stp2-mut1 (Figure 13B), it was also higher for stp2-mut3 than for stp2 (Figure 13C),

suggesting that this effect is independent of the internal AAG, and arguing against our

hypothesis. Moreover, these data suggest that the sequence of the PAM-proximal region of a pre-

spacer impacts the frequency with which it is used during primed adaptation.

The efficiency of primed adaptation is modulated by a PAM-distal motif in the pre-spacer

Although the presence of an AAG within the pre-spacer is clearly an important factor in

determining the frequency of pre-spacer usage, there is considerable local variability in pre-

spacer usage for pre-spacers that lack an internal AAG (Figure 11A + B). We used the frequency

of pre-spacer usage for AAG-flanked pre-spacers in the 100 kb region upstream of the *lacZ+* and

mhpT- protospacers to fit exponential decay models, estimating pre-spacer usage as a function of

distance from the protospacer. Based on the modeled exponential decay, we estimate that the

frequency of pre-spacer usage drops by 50% every ~11,500 – 13,900 kb. Strikingly, an *in vitro*

study of *E. coli* Cas3 translocation indicated that 50% of Cas3 molecules dissociate from DNA

after ~12 kb (Redding et al., 2015), strongly suggesting that the decrease in pre-spacer usage we

observe as a function of distance from the protospacer reflects the extent of Cas3 translocation.

We compared the frequency of pre-spacer usage to the predicted frequency based on the

exponential decay model. We then selected all pre-spacers for which the frequency of usage was

>4-fold greater than that predicted by the model. Alignment of these pre-spacer sequences

revealed sequence bias at positions 28, 30, 32 and 33 (Figure 14A). We next selected all pre-

spacers for which the frequency of usage was >4-fold lower than that predicted by the model.

Alignment of these pre-spacer sequences revealed sequence bias at positions 30, 31, 32 and 33

(Figure 14B). Strikingly, the hierarchy of preferred bases at positions 30, 32 and 33 was

precisely inverted for the enriched and de-enriched pre-spacers. To determine whether the

identified sequence biases are meaningful, we determined the information content of the position

weight matrices (Schneider et al., 1986) for the enriched and de-enriched pre-spacers. We then

determined the information content of the same number of randomly selected pre-spacer

sequences, and repeated this analysis 100,000 times. Based on the distribution of information

content scores for the randomly selected sequences, the information content scores of the

position weight matrices from the enriched and de-enriched pre-spacers were significantly higher

than expected by chance ($Z = 10.4$ and $p < 1e^{-14}$ for enriched pre-spacers; $Z = 4.9$ and $p = 5e^{-7}$ for

de-enriched pre-spacers). We conclude that the sequence of base positions at the 3' end of the

pre-spacer influences usage frequency, with positions 30, 32 and 33 being the most important.

To experimentally test the impact of pre-spacer positions 30-33 on primed adaptation, we

constructed derivatives of stp2 that contain a pre-spacer where positions 30-31 were modified

from TAAA to CCAA (stp2-mut4), or positions 30-33 were modified to CCCG (stp2-mut5;

Figure 14C). Based on the enrichment/de-enrichment of pre-spacers in the chromosomal primed

adaptation experiment (Figure 14A-B), we reasoned that changing positions 30-33 to CCAA

would not have a large effect on the frequency of pre-spacer usage, since our data suggest that

only one of the two modified bases (position 30) would generate a sub-optimal pre-spacer. By

contrast, we reasoned that changing positions 30-33 to CCCG would have a substantial negative

impact on the frequency of pre-spacer usage, since it has the most disfavored base at three PAM-

distal positions. Consistent with our expectation, changing positions 30-31 to CC had no effect

on pre-spacer usage frequency (Figure 14D), whereas changing positions 30-33 to CCCG

substantially reduced pre-spacer usage frequency (Figure 14E).

A previous study suggested that the sequence of positions 32 and 33 of the pre-spacer influences

the efficiency of naïve adaptation. Specifically, a C at position 32 or T at position 33 resulted in

inefficient adaptation whereas an A at position 32 and or an A at position 33 resulted in efficient

adaptation (Yosef et al., 2013). Our data are consistent with the idea that the sequence of

positions 28-33 are important, and moreover are consistent with the specific sequence effects

observed for naïve adaptation. We conclude that the sequence of the PAM-distal portion of the

pre-spacer influences both naïve and primed adaptation, likely due to effects on Cas1-2

association and/or integration of the pre-spacer into the CRISPR array. Our data suggest that the

number of important positions at the PAM-distal end of the pre-spacer is larger than previously

suggested (Yosef et al., 2013), encompassing perhaps as many as six positions.

A model for primed adaptation

We propose that once recruited to a protospacer-bound Cascade, Cas3 translocates along the

DNA, upstream from the PAM, consistent with *in vitro* studies (Dillard et al., 2018; Redding et

al., 2015). The initial translocating state of Cas3 is likely to be different to its final translocating

state; we propose that this initial state is associated with frequent, sequence-independent nicking

of one strand of the DNA. Our data suggest that the transition from the frequent nicking state to

the infrequent nicking state is stochastic, and occurs in all cases within ~200 nt of the

protospacer. Biochemical data suggest that translocating Cas3 initially remains associated with

Cascade (Dillard et al., 2018; Redding et al., 2015). We propose that the interaction between

Cas3 and Cascade causes Cas3 to be in the frequent nicking state, and that it is the dissociation

of Cas3 from Cascade as tension builds in the DNA that causes Cas3 to transition to the

infrequent nicking state. After the transition, we propose that Cas3, or possibly Cas1 that is

associated with Cas3, cuts DNA at AAG sequences. Given the magnitude of the defect in primed

adaptation associated with the presence of an AAG within a spacer, it is likely that this nicking

occurs at almost every AAG encountered by Cas3. It is unclear why the degree to which an AAG

within a pre-spacer affects primed adaptation is dependent on the position of the AAG relative to

the PAM. Given that an AAG within a pre-spacer has a small but significant negative impact on

naïve adaptation, we suggest that Cas1 is responsible for the DNA cut at AAG sequences.

Presumably, during naïve adaptation a second cut by Cas1 is less likely than during primed

adaptation, because Cas1 would not be associated with a translocating Cas3, and hence is less

likely to be suitably positioned on the DNA.

It is unclear how substrates are generated for Cas1-2 during primed adaptation, although nicking

of at least one DNA strand by Cas3 is required (Datsenko et al., 2012). There is strong evidence

for a nick between the second and third positions of the PAM (Musharova et al., 2017; Shiriaeva

et al., 2019), consistent with our proposal that every AAG is cut by the translocating Cas3-

containing complex. However, this would only account for one of the four cuts required for

substrate generation. Future work is required to determine whether Cas3, Cas1, and/or non-Cas

proteins are required for this process. Once substrates have been generated, the efficiency of

integration into the array is dependent on the sequence. Our data indicate that the six most PAM-

distal positions of the pre-spacer contribute to the efficiency of primed adaptation (Figure 14).

Given that the importance of the two most PAM-distal positions has been recognized to be

important for naïve adaptation (Yosef et al., 2013), we propose that the impact of these positions

is solely on Cas1-2. These positions may be important for stable association of Cas1-2, or for the

process of integration into the CRISPR array. Our data also suggest that sequences within pre-

spacers, more proximal to the PAM, play an important role in primed adaptation (Figure 13), but

the mechanism for this is unknown.

MATERIALS AND METHODS

Strains and Plasmids

Strains, plasmids, and oligonucleotides are listed in Tables 1, 2, and 3, respectively. All strains

are derivatives of MG1655 (Blattner et al., 1997). CB386, AMD536 and AMD688 have been

previously described (Cooper et al., 2018; Luo et al., 2014). 1XDNAi, as described previously

(Caliando and Voigt, 2015), contains a chromosomally integrated actuator, pACT-01. AMD671

is a derivative of 1XDNAi that we constructed using FRUIT recombineering (Stringer et al.,

2012). Specifically, *thyA* was deleted in 1XDNAi, using oligonucleotides JW472 and JW473

and strain AMD052 (MG1655 Δ*thyA*) as a template. *thyA* was inserted downstream of the

pACT-01 actuator using FRUIT (Stringer et al., 2012) with oligonucleotides JW9016 and

JW9017. Oligonucleotides JW9009 and JW9010 were used to amplify *cas1-cas2* from MG1655

to replace *thyA* downstream of pACT-01. Lastly, the *lacZ* gene was replaced by *thyA* using

FRUIT (Stringer et al., 2012) with oligonucleotides JW9066 and JW9067.

The self-targeting plasmid (stp; pAMD189), the Cas3-expressing plasmid (pAMD191), and the

plasmid that expresses the crRNA targeting the *lacZ+* protospacer (pCB380), have been

described previously (Cooper et al., 2018; Luo et al., 2014). All other crRNA-expressing

plasmids are derivatives of pAMD179 (Cooper et al., 2018). To clone individual spacers, pairs of

oligonucleotides were annealed, extended, and cloned using In-Fusion (Clontech) into the <u>*Xho*</u>I

and <u>*Sac*</u>II sites of pAMD179 to generate pAMD211 (targets *codA+* protospacer; with

oligonucleotides JW8010 and JW6518), and pAMD212 (targets *mhpT-* protospacer; with

oligonucleotides JW8011 and JW6518).

The second self-targeting plasmid (stp2) is a derivative of pSDS009. pSDS009 was generated by

cloning a synthesized dsDNA fragment (gBlock JW9076; Integrated DNA Technologies, Inc.;

Table 3) using In-Fusion (Clontech) into the *Cla*I and *Hin*dIII sites of pBAD24 amp (Guzman et

al., 1995). stp2 and its derivatives were generated by annealing and extending pairs of

oligonucleotides, and cloning using In-Fusion (Clontech) into the _Hin_dIII site of pSDS009 to

generate pSK014 (stp2; cloned with oligonucleotides JW10011 and JW10012), pSK013 (stp2-

mut1; cloned with oligonucleotides JW10009 and JW10010), pSK015 (stp2-mut2; cloned with

oligonucleotides JW10013 and JW10014), pSK035 (stp2-mut3; cloned with oligonucleotides

JW10015 and JW10016), pSK017 (stp2-mut4; cloned with oligonucleotides JW10021 and

JW10022), and pSK016 (stp2-mut5; cloned with oligonucleotides JW10017 and JW10018).

Quantification of adaptation using a YFP reporter

AMD688 was transformed with two plasmids: the first plasmid was either pAMD189 or empty

pBAD24, while the second was either pAMD191 or empty pBAD33. Cells were grown

overnight in LB supplemented with 100 ug/mL ampicillin and 30 ug/mL chloramphenicol and

were sub-cultured the next day 1:100 for 6 hours in LB supplemented with 0.2% arabinose and

30 ug/mL chloramphenicol. Cells were pelleted by centrifugation and resuspended in M9

minimal medium in twice the original volume ($OD_{600} \approx 1.0$). Samples were transferred to 5 mL

polystyrene round-bottom tubes and analyzed by flow cytometry for single-cell detection of YFP

expression using the BD FACSAria IIU Cell Sorter. We recorded 100,000 events for each

sample.

PCR and Sequencing to Assess Primed Adaptation

To determine the set of pre-spacers acquired during priming, pAMD191 was transformed into AMD536 along with either pAMD189, pCB380, pAMD211 or pAMD212. Cells were grown overnight in LB supplemented with 100 μg/mL ampicillin, 30 μg/mL chloramphenicol, and 0.2% glucose at 37 °C with aeration, and sub-cultured the next day in LB supplemented with 0.2% arabinose at 37 °C with aeration for one hour.

To experimentally assess the impact of specific sequences on pre-spacer usage, pSK013, pSK014, pSK015, pSK035, pSK016, or pSK017 were transformed into AMD671. Cells were grown overnight in LB supplemented with 100 μg/mL ampicillin, and 0.2% glucose at 37°C with aeration, and sub-cultured the next day in LB supplemented with 0.2% arabinose at 37°C with aeration for six hours.

For both sets of cultures described above, cells were pelleted from 1 mL of culture by centrifugation, and cell pellets were frozen at -20 °C. PCRs were then performed on the cell pellets, amplifying the CRISPR arrays using oligonucleotides JW7816 and JW7817 for CRISPR-I and JW7818 and JW7819 for CRISPR-II. PCR products were visualized on acrylamide gels and the first expansion band was extracted and purified. A second round of amplification was performed using a universal forward oligonucleotide (JW7820 or JW8053), and one of a set of reverse oligonucleotides containing Nextera (Illumina) indices (JW8054, JW8057, JW8062, JW8476, JW8477, JW8478, JW8479, JW8480, JW8481 or JW8485). Samples were purified using a MinElute kit (Qiagen) and pooled. Samples were sequenced using an Illumina MiSeq Instrument (Wadsworth Center Applied Genomic Technologies Core).

Plasmids pSK013, pSK014, pSK015, pSK035, pSK016, or pSK017 serendipitously function as

self-targeting plasmids despite the lack of a cloned spacer. These plasmids are derivatives of

pSDS009, which is designed to express crRNAs from a cassette that includes a spacer with a full

repeat downstream, but only 8 nt of repeat sequence upstream. pSDS009 does not itself contain a

canonical spacer between the partial and complete repeat sequences. Similarly, pSK013,

pSK014, pSK015, pSK035, pSK016, and pSK017 do not include a canonical spacer sequence.

Nonetheless, multiple lines of evidence indicate that these plasmids drive efficient primed

adaptation in cells expressing *cas3*: (i) the majority of pre-spacer usage is from the plasmid; (ii)

96% of pre-spacer usage on the plasmid is associated with an AAG PAM; (iii) for AAG-flanked

pre-spacers that are shared with the stp, are found on the stp non-target strand, and are not shared

with pAMD191, pre-spacer usage from pSK014 correlates well with that from the stp

(correlation coefficient of 0.93). We suspect that the downstream repeat sequence in pSDS009

and its derivatives also functions as an upstream repeat sequence for a cryptic crRNA. In this

scenario, the sequence downstream of this repeat serves as a spacer, with the end of the crRNA

presumably being determined by the position of the downstream transcription terminator. We

previously showed that the exact same sequence results in expression of a cryptic crRNA from a

derivative of pAMD179 (Cooper et al., 2018). The presumed spacer sequence is duplicated in the

same orientation, a short distance downstream, which would facilitate interference and primed

adaptation.

Spacer sequence extraction and alignment to reference genomes

Newly acquired spacers found directly adjacent to the leader proximal endogenous spacer (the

first primed adaptation acquired spacer) were extracted from .fastq files using custom Python

scripts for each CRISPR array being analyzed. The scripts first selected sequences that included

the first spacer of the unexpanded CRISPR array (spacer #1) and at least one repeat sequence.

The script then removed sequences for which the gap between spacer #1 and the next upstream

repeat differed from the expected value. The script then extracted the sequence between the first

repeat and the repeat immediately upstream of spacer #1. This was presumed to consist of

complete spacer sequences, or combinations of spacers and repeats (i.e. the result of >1 spacer

being added to the CRISPR array). Sequences were discarded if they were shorter than 30 nt or

longer than 36 nt, ensuring that only arrays with a single expansion were analyzed. Spacers were

mapped to their respective genomes (stp, pAMD191, or *E. coli* K-12 MG1566 U00096.3) using

CLC Genomics Workbench v10.1.1, requiring perfect matches. For experiments involving

mapping reads to multiple genomes, we first mapped to stp, then mapped all unmapped reads to

pAMD191, and then mapped all unmapped reads to *E. coli* K-12 MG1655.

Analysis of primed adaptation in regions adjacent to off-target Cascade binding sites

We previously described a set of 76 off-target Cascade binding sites whose sequence is

consistent with Cascade binding in association with the crRNA targeting the *lacZ*+ protospacer

(Cooper et al., 2018). We counted the frequency of pre-spacer usage for each of these off-target

protospacers, searching on the non-target strand in the 10 kb upstream of the protospacer. We

repeated the analysis for 1,000 randomly selected genomic positions equally distributed across

the forward and reverse strands.

Analysis of slipping for 33 nt pre-spacers

We selected all 33 nt pre-spacers on the non-target strand, in the 10 kb regions upstream of the

lacZ+ and *mhpT-* protospacers, for which the PAM was not AAG. We calculated all pairwise

distances for these pre-spacers to AAG-associated pre-spacers on the non-target strand, in the 10

kb upstream of each of the protospacer. For the analysis represented in Figure 7A, only unique

pre-spacer positions were considered, so information on the frequency with which different pre-

spacers are used was lot. Values plotted in Figure 7A represent the number of instances of the

indicated slipping distance divided by the total number of comparisons between pre-spacer

positions and AAGs, multiplied by 1000. Values listed in Figure 7B take into account the fact

that many pre-spacer positions were detected more than once, and hence these numbers control

for the relative use of the different sequences. Pre-spacers with a non-AAG PAM that did not

derive from a -1, +1 or +2 slip were selected; pre-spacers that were sequenced multiple times

were represented at the frequency with which they were detected. The associated PAM

sequences were then converted into the logos shown in Figure 7C using Weblogo (Crooks et al.,

2004).

Analysis of flipping

We selected all 33 nt pre-spacers on the target strand, in the 10 kb regions downstream of the

lacZ+ and *mhpT-* protospacers. We calculated all pairwise distances for these pre-spacers to

AAG-associated pre-spacers on the non-target strand, in the 10 kb upstream of each of the

protospacer. For the analysis represented in Figure 8B, only unique pre-spacer positions were

considered, so information on the frequency with which different pre-spacers are used was lot.

Values plotted in Figure 8B represent the number of instances of the indicated slipping distance

divided by the total number of comparisons between pre-spacer positions and AAGs, multiplied

by 1000.

Analysis of slipping for spacers of non-canonical length

We selected all 32 and 34 nt pre-spacers on the non-target strand, in the 10 kb regions upstream

of the *lacZ+* and *mhpT-* protospacers. We calculated all pairwise distances for these pre-spacers

to AAG-associated pre-spacers on the non-target strand, in the 10 kb upstream of each of the

protospacer. For the analysis represented in Figure 9C-D, only unique pre-spacer positions were

considered, so information on the frequency with which different pre-spacers are used was lot.

Values plotted in Figure 9C-D represent the number of instances of the indicated slipping

distance divided by the total number of comparisons between pre-spacer positions and AAGs,

multiplied by 1000.

Identification of pre-spacers with higher-than-expected or lower-than-expected usage
frequency, independent of internal AAG sequences

We took the frequency of pre-spacer usage for all AAG-flanked 33 nt pre-spacers in the 100 kb

region upstream of the *lacZ+* and *mhpT-* protospacers. We then fit an exponential decay model to

each dataset using Microsoft Excel ("best fit line" function). Using the parameters associated

with these models, we determined the expected usage frequency for pre-spacers at every position

in the 100 kb windows, and we then took the ratio of the actual usage frequency to the expected

usage frequency. Pre-spacers with ratios >4 or <0.25 were selected for analysis of tetranucleotide

and trinucleotide content.

Identification of pre-spacers with higher-than-expected or lower-than-expected usage

frequency, independent of internal AAG sequences

We took the frequency of pre-spacer usage for all AAG-flanked 33 nt pre-spacers in the 100 kb

region upstream of the *lacZ+* and *mhpT-* protospacers, excluding pre-spacers with an internal

AAG. We then fit an exponential decay model to each dataset using Microsoft Excel ("best fit

line" function). Using the parameters associated with these models, we determined the expected

usage frequency for pre-spacers at every position in the 100 kb windows, and we then took the

ratio of the actual usage frequency to the expected usage frequency. Pre-spacers with ratios >4 or

<0.25 were used as input for Weblogo (Crooks et al., 2004) to create the sequence logos shown

in Figure 14A-B.

ACKNOWLEDGEMENTS, ACCESSION NUMBER, AND FUNDING

We thank the Wadsworth Center Applied Genomic Technologies Core Facility for DNA

sequencing. We thank the Wadsworth Center Immunology Core Facility for flow cytometry

analysis. We thank the Wadsworth Center Tissue Culture and Media Core Facility and

Glassware Facility for technical support. We thank Randy Morse, Keith Derbyshire and Todd

Gray for helpful discussions.

Raw Illumina sequencing data are available from EBI ArrayExpress using accession number

E-MTAB-8700.

This work was supported by National Institutes of Health grants R01GM122836 (JTW) and

R21AI126416 (JTW).

REFERENCES

Amlinger, L., Hoekzema, M., Wagner, E.G.H., Koskiniemi, S., and Lundgren, M. (2017).

Fluorescent CRISPR Adaptation Reporter for rapid quantification of spacer acquisition. Sci. Rep.

7, 10392.

Blattner, F.R., Plunkett, G., Bloch, C.A., Perna, N.T., Burland, V., Riley, M., Collado-Vides, J.,

Glasner, J.D., Rode, C.K., Mayhew, G.F., et al. (1997). The complete genome sequence of

Escherichia coli K-12. Science *277*, 1453–1462.

Brouns, S.J.J., Jore, M.M., Lundgren, M., Westra, E.R., Slijkhuis, R.J.H., Snijders, A.P.L.,

Dickman, M.J., Makarova, K.S., Koonin, E.V., and van der Oost, J. (2008). Small CRISPR

RNAs guide antiviral defense in prokaryotes. Science *321*, 960–964.

Caliando, B.J., and Voigt, C.A. (2015). Targeted DNA degradation using a CRISPR device

stably carried in the host genome. Nat. Commun. *6*, 6989.

Cooper, L.A., Stringer, A.M., and Wade, J.T. (2018). Determining the Specificity of Cascade

Binding, Interference, and Primed Adaptation In Vivo in the Escherichia coli Type I-E CRISPR-

Cas System. MBio *9*.

Crooks, G.E., Hon, G., Chandonia, J.-M., and Brenner, S.E. (2004). WebLogo: a sequence logo

generator. Genome Res. *14*, 1188–1190.

Datsenko, K.A., Pougach, K., Tikhonov, A., Wanner, B.L., Severinov, K., and Semenova, E.

(2012). Molecular memory of prior infections activates the CRISPR/Cas adaptive bacterial

immunity system. Nat. Commun. *3*, 945.

Dillard, K.E., Brown, M.W., Johnson, N.V., Xiao, Y., Dolan, A., Hernandez, E., Dahlhauser,

S.D., Kim, Y., Myler, L.R., Anslyn, E.V., et al. (2018). Assembly and Translocation of a

CRISPR-Cas Primed Acquisition Complex. Cell *175*, 934-946.e15.

Fagerlund, R.D., Wilkinson, M.E., Klykov, O., Barendregt, A., Pearce, F.G., Kieper, S.N.,

Maxwell, H.W.R., Capolupo, A., Heck, A.J.R., Krause, K.L., et al. (2017). Spacer capture and

integration by a type I-F Cas1–Cas2-3 CRISPR adaptation complex. Proc. Natl. Acad. Sci. *114*,

E5122–E5128.

Fineran, P.C., Gerritzen, M.J.H., Suárez-Diez, M., Künne, T., Boekhorst, J., Hijum, S.A.F.T.

van, Staals, R.H.J., and Brouns, S.J.J. (2014). Degenerate target sites mediate rapid primed

CRISPR adaptation. Proc. Natl. Acad. Sci. *111*, E1629–E1638.

Goren, M.G., Yosef, I., Auster, O., and Qimron, U. (2012). Experimental definition of a

clustered regularly interspaced short palindromic duplicon in Escherichia coli. J. Mol. Biol. *423*,

14–16.

Guzman, L.M., Belin, D., Carson, M.J., and Beckwith, J. (1995). Tight regulation, modulation,

and high-level expression by vectors containing the arabinose PBAD promoter. J. Bacteriol. *177*,

4121–4130.

Jackson, S.A., McKenzie, R.E., Fagerlund, R.D., Kieper, S.N., Fineran, P.C., and Brouns, S.J.J.

(2017). CRISPR-Cas: Adapting to change. Science *356*.

Jackson, S.A., Birkholz, N., Malone, L.M., and Fineran, P.C. (2019). Imprecise Spacer

Acquisition Generates CRISPR-Cas Immune Diversity through Primed Adaptation. Cell Host

Microbe *25*, 250-260.e4.

Jore, M.M., Lundgren, M., van Duijn, E., Bultema, J.B., Westra, E.R., Waghmare, S.P.,

Wiedenheft, B., Pul, Ü., Wurm, R., Wagner, R., et al. (2011). Structural basis for CRISPR RNA-

guided DNA recognition by Cascade. Nat. Struct. Mol. Biol. *18*, 529–536.

Jung, C., Hawkins, J.A., Jones, S.K., Xiao, Y., Rybarski, J.R., Dillard, K.E., Hussmann, J.,

Saifuddin, F.A., Savran, C.A., Ellington, A.D., et al. (2017). Massively Parallel Biophysical

Analysis of CRISPR-Cas Complexes on Next Generation Sequencing Chips. Cell *170*, 35-

47.e13.

Koonin, E.V., Makarova, K.S., and Zhang, F. (2017). Diversity, classification and evolution of

CRISPR-Cas systems. Curr. Opin. Microbiol. *37*, 67–78.

Künne, T., Kieper, S.N., Bannenberg, J.W., Vogel, A.I.M., Miellet, W.R., Klein, M., Depken,

M., Suarez-Diez, M., and Brouns, S.J.J. (2016). Cas3-Derived Target DNA Degradation

Fragments Fuel Primed CRISPR Adaptation. Mol. Cell *63*, 852–864.

Leenay, R.T., and Beisel, C.L. (2017). Deciphering, Communicating, and Engineering the

CRISPR PAM. J. Mol. Biol. *429*, 177–191.

Levy, A., Goren, M.G., Yosef, I., Auster, O., Manor, M., Amitai, G., Edgar, R., Qimron, U., and

Sorek, R. (2015). CRISPR adaptation biases explain preference for acquisition of foreign DNA.

Nature *520*, 505–510.

Li, M., Wang, R., Zhao, D., and Xiang, H. (2014). Adaptation of the Haloarcula hispanica

CRISPR-Cas system to a purified virus strictly requires a priming process. Nucleic Acids Res.

42, 2483–2492.

Li, M., Gong, L., Zhao, D., Zhou, J., and Xiang, H. (2017). The spacer size of I-B CRISPR is

modulated by the terminal sequence of the protospacer. Nucleic Acids Res. *45*, 4642–4654.

Luo, M.L., Mullis, A.S., Leenay, R.T., and Beisel, C.L. (2014). Repurposing endogenous type I

CRISPR-Cas systems for programmable gene repression. Nucleic Acids Res. gku971.

Musharova, O., Klimuk, E., Datsenko, K.A., Metlitskaya, A., Logacheva, M., Semenova, E.,

Severinov, K., and Savitskaya, E. (2017). Spacer-length DNA intermediates are associated with

Cas1 in cells undergoing primed CRISPR adaptation. Nucleic Acids Res. *45*, 3297–3307.

Musharova, O., Vyhovskyi, D., Medvedeva, S., Guzina, J., Zhitnyuk, Y., Djordjevic, M.,

Severinov, K., and Savitskaya, E. (2018). Avoidance of Trinucleotide Corresponding to

Consensus Protospacer Adjacent Motif Controls the Efficiency of Prespacer Selection during

Primed Adaptation. MBio *9*.

Musharova, O., Sitnik, V., Vlot, M., Savitskaya, E., Datsenko, K.A., Krivoy, A., Fedorov, I.,

Semenova, E., Brouns, S.J.J., and Severinov, K. (2019). Systematic analysis of Type I-E

Escherichia coli CRISPR-Cas PAM sequences ability to promote interference and primed

adaptation. Mol. Microbiol. *111*, 1558–1570.

Rao, C., Chin, D., and Ensminger, A.W. (2017). Priming in a permissive type I-C CRISPR-Cas

system reveals distinct dynamics of spacer acquisition and loss. RNA N. Y. N *23*, 1525–1538.

Redding, S., Sternberg, S.H., Marshall, M., Gibb, B., Bhat, P., Guegler, C.K., Wiedenheft, B.,

Doudna, J.A., and Greene, E.C. (2015). Surveillance and Processing of Foreign DNA by the

Escherichia coli CRISPR-Cas System. Cell *163*, 854–865.

Richter, C., Dy, R.L., McKenzie, R.E., Watson, B.N.J., Taylor, C., Chang, J.T., McNeil, M.B.,

Staals, R.H.J., and Fineran, P.C. (2014). Priming in the Type I-F CRISPR-Cas system triggers

strand-independent spacer acquisition, bi-directionally from the primed protospacer. Nucleic

Acids Res. *42*, 8516–8526.

Savitskaya, E., Semenova, E., Dedkov, V., Metlitskaya, A., and Severinov, K. (2013). High-

throughput analysis of type I-E CRISPR/Cas spacer acquisition in E. coli. RNA Biol. *10*, 716–

725.

Schneider, T.D., Stormo, G.D., Gold, L., and Ehrenfeucht, A. (1986). Information content of

binding sites on nucleotide sequences. J. Mol. Biol. *188*, 415–431.

Semenova, E., Savitskaya, E., Musharova, O., Strotskaya, A., Vorontsova, D., Datsenko, K.A.,

Logacheva, M.D., and Severinov, K. (2016). Highly efficient primed spacer acquisition from

targets destroyed by the Escherichia coli type I-E CRISPR-Cas interfering complex. Proc. Natl.

Acad. Sci. 201602639.

Shiriaeva, A.A., Savitskaya, E., Datsenko, K.A., Vvedenskaya, I.O., Fedorova, I., Morozova, N.,

Metlitskaya, A., Sabantsev, A., Nickels, B.E., Severinov, K., et al. (2019). Detection of spacer

precursors formed in vivo during primed CRISPR adaptation. Nat. Commun. *10*, 4603.

Shmakov, S., Savitskaya, E., Semenova, E., Logacheva, M.D., Datsenko, K.A., and Severinov,

K. (2014). Pervasive generation of oppositely oriented spacers during CRISPR adaptation.

Nucleic Acids Res. *42*, 5907–5916.

Staals, R.H.J., Jackson, S.A., Biswas, A., Brouns, S.J.J., Brown, C.M., and Fineran, P.C. (2016).

Interference-driven spacer acquisition is dominant over naive and primed adaptation in a native

CRISPR–Cas system. Nat. Commun. *7*, 12853.

Sternberg, S.H., Richter, H., Charpentier, E., and Qimron, U. (2016). Adaptation in CRISPR-Cas

Systems. Mol. Cell *61*, 797–808.

Stringer, A.M., Singh, N., Yermakova, A., Petrone, B.L., Amarasinghe, J.J., Reyes-Diaz, L.,

Mantis, N.J., and Wade, J.T. (2012). FRUIT, a scar-free system for targeted chromosomal

mutagenesis, epitope tagging, and promoter replacement in Escherichia coli and Salmonella

enterica. PloS One *7*, e44841–e44841.

Strotskaya, A., Savitskaya, E., Metlitskaya, A., Morozova, N., Datsenko, K.A., Semenova, E.,

and Severinov, K. (2017). The action of Escherichia coli CRISPR-Cas system on lytic

bacteriophages with different lifestyles and development strategies. Nucleic Acids Res *45*, 1946–

1957.

Swarts, D.C., Mosterd, C., van Passel, M.W.J., and Brouns, S.J.J. (2012). CRISPR Interference

Directs Strand Specific Spacer Acquisition. PLoS ONE *7*, e35888.

Westra, E.R., van Erp, P.B.G., Künne, T., Wong, S.P., Staals, R.H.J., Seegers, C.L.C., Bollen, S.,

Jore, M.M., Semenova, E., Severinov, K., et al. (2012). CRISPR immunity relies on the

consecutive binding and degradation of negatively supercoiled invader DNA by Cascade and

Cas3. Mol. Cell *46*, 595–605.

Wright, A.V., Nuñez, J.K., and Doudna, J.A. (2016). Biology and Applications of CRISPR

Systems: Harnessing Nature's Toolbox for Genome Engineering. Cell *164*, 29–44.

Xue, C., Seetharam, A.S., Musharova, O., Severinov, K., Brouns, S.J., Severin, A.J., and

Sashital, D.G. (2015). CRISPR interference and priming varies with individual spacer sequences.

Nucleic Acids Res *43*, 10831–10847.

Yosef, I., Shitrit, D., Goren, M.G., Burstein, D., Pupko, T., and Qimron, U. (2013). DNA motifs

determining the efficiency of adaptation into the Escherichia coli CRISPR array. Proc. Natl.

Acad. Sci. U. S. A. *110*, 14396–14401.

Table 1. Strains used in this study.

Name	Description	Source
MG1655	Wild-type *E. coli*	(Blattner et al., 1997)
CB386	MG1655 [Δ*cas3* P*cse1*]::[cat PJ23119]	(Luo et al., 2014)
AMD536	MG1655 [Δ*cas3* P*cse1*]::[PJ23119]	(Cooper et al., 2018)
AMD688	MLS1003 [Δ*cas3* P*cse1*]::[PJ23119]	(Cooper et al., 2018)
1XDNAi	MG1655 Δ*araC-araBAD*, ΔP*lacI*:*lacI*, Δ*cas-*CRISPR::pACT-01 (Kan-)	(Caliando and Voigt, 2015)
AMD671	MG1655 Δ*araC-araBAD*, ΔP*lacI*:*lacI*, Δcas-CRISPR::pACT-01, *cas1*$^{+}$ *cas2*$^{+}$, Δ*lacZ*::*thyA*	This Study
AMD052	MG1655 Δ*thyA*	(Stringer et al., 2012)

920   **Table 2. Plasmids used in this study.**

Name	Description	Source
pBAD24 amp	Empty pBAD24 amp	(Guzman et al., 1995)
pBAD33 cam	Empty pBAD33 cam	(Guzman et al., 1995)
pAMD179	Parent vector for cloning crRNAs	(Cooper et al., 2018)
pAMD189	stp; pAMD179 expressing a self-targeting crRNA	(Cooper et al., 2018)
pAMD191	pBAD33-*cas3*	(Cooper et al., 2018)
pCB380	pcrRNA.con-*lacZ*	(Luo et al., 2014)
pAMD211	pAMD179 expressing a crRNA targeting *mhpT*	This Study
pAMD212	pAMD179 expressing a crRNA targeting *codA*	This Study
pSDS009	Parent vector for cloning pre-spacers	This Study
pSK013	pSDS009 stp2-mut1 pre-spacer	This Study
pSK014	stp2; pAMD179 with an added pre-spacer	This Study
pSK015	pSDS009 stp2-mut2 pre-spacer	This Study
pSK035	pSDS009 stp2-mut3 pre-spacer	This Study
pSK016	pSDS009 stp2-mut5 pre-spacer	This Study
psK017	pSDS009 stp2-mut4 pre-spacer	This Study

923   **Table 3. Oligonucleotides and synthesized dsDNA used in this study.**

Name	Sequence
JW472	CCGACGCGCAGTTTA
JW473	CACGTTGTGTTTTCATGC
JW6518	CAGCGGGGATAAACC
JW7816	TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGAAGGTTGGTGGGTTGTTTTTATGGG
JW7817	GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGTTATCAATTACAACCGACAGGGAGCC
JW7818	TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGAAAGTTGGTAGATTGTGACTGGC
JW7819	GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGCAACAGCAGCACCCATGAC
JW7820	AATGATACGGCGACCACCGAGATCTACACGCGTAAGATCGTCGGCAGCGTC
JW8010	GCGCGGGGAACTCGAGCAGTCGCGCTTTGTCGAAACCGTTGCTGCCGGTTTATCCCCGC
JW8011	GCGCGGGGAACTCGAGGTGCCAGGGCATACAAAACGCTTTGCCCACGGTTTATCCCCGC
JW8053	AATGATACGGCGACCACCGAGATCTACACTCCAGGTATCGTCGGCAGCGTCAGATGTG
JW8054	CAAGCAGAAGACGGCATACGAGATGGATTCACGTCTCGTGGGCTCGGAGATGTG
JW8057	CAAGCAGAAGACGGCATACGAGATCGCATTAGGTCTCGTGGGCTCGGAGATGTG
JW8062	CAAGCAGAAGACGGCATACGAGATATGGACTCGTCTCGTGGGCTCGGAGATGTG
JW8476	CAAGCAGAAGACGGCATACGAGATTAAGGCGAGTCTCGTGGGCTCGGAGATGTG
JW8477	CAAGCAGAAGACGGCATACGAGATCGTACTAGGTCTCGTGGGCTCGGAGATGTG
JW8478	CAAGCAGAAGACGGCATACGAGATAGGCAGAAGTCTCGTGGGCTCGGAGATGTG
JW8479	CAAGCAGAAGACGGCATACGAGATTCCTGAGCGTCTCGTGGGCTCGGAGATGTG
JW8480	CAAGCAGAAGACGGCATACGAGATGGACTCCTGTCTCGTGGGCTCGGAGATGTG
JW8481	CAAGCAGAAGACGGCATACGAGATTAGGCATGGTCTCGTGGGCTCGGAGATGTG
JW8485	CAAGCAGAAGACGGCATACGAGATCGAGGCTGGTCTCGTGGGCTCGGAGATGTG
JW9009	ATTGGGCCAGCTAAATCG
JW9010	GGCTCATTATACCAGTCAGGACGTTGGGAAGAGGCCGCTCAAACAGGTAAAAAAGACACC

JW9016	GCTAAATCGATGGGATGTGGCTTGCTATCTTTGGCTCCACTGTGATAGACAGCTGCATGCAT
JW9017	AGAACTGGCTCATTATACCAGTCAGGACGTTGGGAAGAGGCCGCGTGTAGGCTGGAGCTG
JW9066	TGTGGAATTGTGAGCGGATAACAATTTCACACAGGAAACAGCTTAGACAGCTGCATGCAT
JW9067	TTCCTTACGCGAAATACGGGCAGACATGGCCTGCCCGGTTATTAGTGTAGGCTGGAGCTG
JW9076 (dsDNA)	TGTTTGACAGCTTATCATCGATTTGACAGCTAGCTCAGTCCTAGGTATAATGCTAGCATAAACCGCGGTACTTAGCTCCTCAGATTAGGATTGCGGAGAATAACAACCGCCGTTCTCATCGAGTAATCTCCGGATATCGACCCATAACGGGCAATGATAAAAGGAGTAACCTGTGAAAAGATGCAATCTATCGTACTCGCACTTTCCCTGGTTCTGGTCGCTCCCATGGCAGCACAGGCTGCGGAAATTACGTTAGTCCCGTCAGTAAAATTACAGATAGGGAGGAGCGATCCTGCAGTGTTCCCCGCGCCAGCGGGGATAAACCGAGGGAACTGCCAGGCATCAAATAAAACGAAAGGCTCAGTCGAAAGACTGGGCCTTTCGTTTTATCTGTTGTTTGTCGGTGAACGCTCTCCTGAGTAGGACAAGCTTGGCTGTTTTGGCGGA
JW10009	TGAGTAGGACAAGCTTACTAATTATAATAGAAGCCAGATACTAAAAGCTTGGCTGTTTT
JW10010	AAAACAGCCAAGCTTTTTAGTATCTGGCTTCTATTATAATTAGTAAGCTTGTCCTACTCA
JW10011	TGAGTAGGACAAGCTTACTAATTATAATAGAAACCAGATACTAAAAGCTTGGCTGTTTT
JW10012	AAAACAGCCAAGCTTTTTAGTATCTGGTTTCTATTATAATTAGTAAGCTTGTCCTACTCA
JW10013	TGAGTAGGACAAGCTTGCCGGCTGCAGCAGAAGCCAGATACTAAAAGCTTGGCTGTTTT
JW10014	AAAACAGCCAAGCTTTTTAGTATCTGGCTTCTATTATATATAGTAAGCTTGTCCTACTCA
JW10015	TGAGTAGGACAAGCTTGCCGGCTGCAGCAGAAACCAGATACTAAAAGCTTGGCTGTTTT
JW10016	AAAACAGCCAAGCTTTTTAGTATCTGGCTTCTATTATATATAGTAAGCTTGTCCTACTCA
JW10017	TGAGTAGGACAAGCTTACTAATTATAATAGAAACCAGATACCCCGAAGCTTGGCTGTTTT
JW10018	AAAACAGCCAAGCTTCGGGGTATCTGGTTTCTATTATAATTAGTAAGCTTGTCCTACTCA
JW10021	TGAGTAGGACAAGCTTACTAATTATAATAGAAACCAGATACCCAAAGCTTGGCTGTTTT
JW10022	AAAACAGCCAAGCTTTTGGGTATCTGGTTTCTATTATAATTAGTAAGCTTGTCCTACTCA

 FIGURE LEGENDS

Figure 1. Primed adaptation is >1000-fold more efficient than naïve adaptation. (A)

Schematic showing the *yfp* reporter construct of strain AMD688 used to quantify adaptation. The

genome of the parent strain has a single CRISPR repeat embedded within a *yfp* open reading

frame such that the repeat causes a frame-shift, preventing translation of YFP. Expansion of the

CRISPR array by a single repeat and spacer restores the frame, permitting translation of YFP,

leading to detectable fluorescence. **(B)** Percentage of YFP$^+$ cells for an *E. coli* strain containing

the *yfp* reporter (AMD688), either the stp (pAMD189; "crRNA +") or an equivalent empty

vector (pBAD24; "crRNA -"), and either a *cas3*-expressing plasmid (pAMD191; "cas3 +") or an

equivalent empty vector (pBAD33; "cas3 -"), following addition of arabinose to the cultures.

Figure 2. Primed adaptation from a plasmid results in primed adaptation from shared

chromosomal locations. (A) Frequency of pre-spacer usage across the stp (pAMD189) during

primed adaptation in AMD536 cells containing pAMD189 and pAMD191. Values represent the

number of sequenced instances of each pre-spacer, plotted as a function of position on the

plasmid. Positive values indicate pre-spacers on the forward strand; negative values indicate pre-

spacers on the reverse strand. **(B)** Frequency of pre-spacer usage on the stp compared for spacers

acquired in the CRISPR-I and CRISPR-II arrays. Data are only plotted for 33 nt pre-spacers that

were sequenced at least once in both datasets. **(C)** Frequency of pre-spacer usage across

pAMD191 during primed adaptation in AMD536 cells containing pAMD189 and pAMD191.

Only pre-spacers unique to pAMD191 are shown; grey shaded regions indicate shared sequence

with pAMD189. **(D)** Frequency of pre-spacer usage across a chromosomal region encompassing

the *araC* gene during primed adaptation in AMD536 cells containing pAMD189 and pAMD191.

Values are plotted as a function of genome position. Only pre-spacers unique to the chromosome

are shown; the grey shaded region indicates shared sequence with pAMD189.

Figure 3. Primed adaptation on the chromosome occurs over regions of >100 kb. Frequency

of pre-spacer usage across a chromosomal region encompassing the *lacZ+* and *mhpT-*

protospacers for AMD536 cells containing pAMD191 (encodes Cas3), and either pCB380

(encodes crRNA targeting *lacZ+*; data shown in black) or pAMD212 (encodes crRNA targeting

mhpT-; data shown in blue). Values represent the relative number of sequenced instances of each

pre-spacer, plotted as a function of genome position. Positive values indicate pre-spacers on the

forward strand; negative values indicate pre-spacers on the reverse strand. Green boxes indicate

positions of repetitive insertion element sequences. The pink box indicates the region analyzed in

Figure 6B.

Figure 4. Off-target Cascade binding sites are not associated with detectable primed

adaptation. Frequency of pre-spacer usage in the 10 kb upstream of the on-target *lacZ+*

protospacer (orange datapoint), or off-target protospacers identified by (Cooper et al., 2018), on

the non-target strand, only counting 33 nt pre-spacers. Relative Cascade association with each

target sequence, as determined by ChIP-seq (Cooper et al., 2018), is shown on the x-axis.

Figure 5. Replicate primed adaptation datasets for chromosomal protospacers are highly

reproducible. Frequency of pre-spacer usage across the chromosome for each of two replicates

for AMD536 cells containing pAMD191 (encodes Cas3), and **(A)** pCB380 (encodes crRNA

targeting *lacZ+*), **(B)** pAMD212 (encodes crRNA targeting *mhpT-*), or **(C)** pAMD211 (encodes

crRNA targeting *codA+*). Values represent the number of sequenced instances of each pre-

spacer, for every 33 nt pre-spacer that mapped to the chromosome.

Figure 6. The majority of primed adaptation occurs on the non-target strand, upstream of

the protospacer. (A) Distribution of pre-spacer usage for *lacZ+* (black) and *mhpT-* (blue)

between the non-target (positive values) and target (negative values) strands, and between the

upstream (dark colors) and downstream (light colors) directions. **(B)** Frequency of pre-spacer

usage in the 10 kb upstream of the *mhpT-* protospacer on the target strand for AMD536 cells

containing pAMD191 (encodes Cas3), and pCB380 (encodes crRNA targeting *lacZ+*; x-axis) or

pAMD212 (encodes crRNA targeting *mhpT-*; y-axis). Values represent the number of sequenced

instances of each pre-spacer, for every 33 nt pre-spacer associated with an AAG PAM.

Figure 7. Slipping accounts for the majority of pre-spacers on the non-target strand that

are not associated with an AAG PAM. (A) Normalized frequency of distances ("slipping

distance"; x-axis) between uniquely positioned protospacers with an AAG PAM and uniquely

positioned protospacers with a non-AAG PAM, for the 10 kb upstream of the *lacZ+* (black

datapoints) and *mhpT-* (blue datapoints) protospacers, on the non-target strand. Values were

normalized to the total number of pairwise comparisons. **(B)** Distribution of pre-spacer usage for

all pre-spacers with a non-AAG PAM, for the 10 kb upstream of the protospacer on the non-

target strand. Values represent averages from the *lacZ+* and *mhpT-* experiments, with error

values representing one standard deviation from the mean. **(C)** DNA sequence logos for PAMs

from pre-spacers with a non-AAG PAM, for the 10 kb upstream of each of the *lacZ+* and *mhpT-*

protospacers on the non-target strand, excluding pre-spacers associated with -2, -1, +1 or +2

slipping events. Logos were generated from all pre-spacer usage (i.e. not just unique pre-

spacerse).

Figure 8. Flipping often occurs following a slipping event. (A) Schematic showing different

possible flipping events. The yellow highlighted sequences indicate the AAG PAM associated

with the non-flipped pre-spacer. **(B)** Normalized frequency of distances ("slipping distance"; x-

axis) between uniquely positioned protospacers with an AAG PAM on the non-target strand, and

uniquely positioned protospacers on the target strand, for the 10 kb upstream of the *lacZ+* (black

datapoints) and *mhpT-* (blue datapoints) protospacers on the non-target strand (i.e. 10 kb

downstream of the protospacers on the target strand). Values were normalized to the total

number of pairwise comparisons.

1006

Figure 9. Pre-spacers with non-canonical lengths are often associated with slipping. (A)

Schematic showing pre-spacers of non-canonical lengths associated with an AAG PAM in either

a slipped or non-slipped configuration. **(B)** Normalized frequency of distances ("slipping

distance"; x-axis) between uniquely positioned, 33 nt protospacers with an AAG PAM and

uniquely positioned 32 nt protospacers, or **(C)** 33 nt protospacers, for the 10 kb upstream of the

lacZ+ (black datapoints) and *mhpT-* (blue datapoints) protospacers, on the non-target strand.

Values were normalized to the total number of pairwise comparisons.

1014

Figure 10. Pre-spacers with 200 nt of the protospacer are used less frequently. (A)

Frequency of pre-spacer usage across a chromosomal region encompassing the *lacZ+* and *codA+*

protospacers for AMD536 cells containing pAMD191 (encodes Cas3), and either pCB380

(encodes crRNA targeting *lacZ*+; data shown in black) or pAMD211 (encodes crRNA targeting

codA+; data shown in purple). Values represent the relative number of sequenced instances of

each pre-spacer, plotted as a function of genome position. Data are only plotted for pre-spacers

on the reverse strand. The blue box indicates the 10 kb region analyzed in panel B. **(B)**

Frequency of pre-spacer usage in the 10 kb upstream of the *lacZ*+ protospacer on the non-target

strand for AMD536 cells containing pAMD191 (encodes Cas3), and pAMD211 (encodes crRNA

targeting *codA*+; x-axis) or pCB380 (encodes crRNA targeting *lacZ*+; y-axis). Values represent

the number of sequenced instances of each pre-spacer, for every 33 nt pre-spacer associated with

an AAG PAM. Data for pre-spacers within 200 nt of the *lacZ*+ protospacer are circled, and the

distance from the *lacZ*+ protospacer is indicated.

1028

Figure 11. Pre-spacers that contain an AAG are used less frequently. Frequency of pre-

spacer usage across the 100 kb upstream of the **(A)** *lacZ*+ and **(B)** *mhpT-* protospacers for

AMD536 cells containing pAMD191 (encodes Cas3), and either pCB380 (encodes crRNA

targeting *lacZ*+) or pAMD212 (encodes crRNA targeting *mhpT-*). One was added to all

frequency values to allow visualization of zero-scoring datapoints. Data are only shown for 33 nt

protospacers on the non-targets strand, associated with an AAG PAM. Orange datapoints

represent pre-spacers with an internal AAG. **(C)** Frequency of tetranucleotide sequence usage

within pre-spacers that were de-enriched (x-axis) or de-enriched (y-axis) relative to an

exponential decay model. Values represent the percentage of all tetranucleotides, averaged for

data from the *lacZ*+ and *mhpT-* protospacer datasets. Error bars represent one standard deviation

from the mean. Tetranucleotide sequences containing AAG are plotted in orange. **(D)** Same as

(C), but for trinucleotide sequences.

1041

Figure 12. Experimental evidence that pre-spacers containing an AAG are used less

frequently in primed adaptation, and to a lesser degree in naïve adaptation. **(A)** Schematic

showing the unmodified and modified pre-spacer sequences for the stp2, stp2-mut1, stp2-mut2

and stp2-mut3 plasmids. The PAM is indicated by a dashed box. Grey highlighting indicates

changes relative to the unmodified pre-spacer. **(B)** Frequency of pre-spacer usage for all pre-

spacers on the unmodified and mut1 stp2 plasmids, for pre-spacers on the non-target strand with

an AAG PAM. The single modified pre-spacer that differs between the stp2 and stp2-mut1 is

shown in orange. The frequency of pre-spacer usage from the stp2 and stp2-mut1 plasmids are

shown on the *x*-axis and *y*-axis, respectively. **(C)** Tukey boxplot showing the distribution of pre-

spacers usage or all chromosomal sequences with an AAG PAM that were detected at least once

in an assay of naïve adaptation (Yosef et al., 2013). Pre-spacers are separated depending on the

presence or absence of an internal AAG sequence.

1054

Figure 13. The impact of an AAG within a pre-spacer on primed adaptation efficiency is

dependent on position of the AAG relative to the PAM. **(A)** Ratio of actual pre-spacer usage

frequency to the frequency predicted by an exponential decay model for all pre-spacers with an

internal AAG. Data are only shown for pre-spacers with an AAG PAM, on the non-target strand,

<100 kb upstream of the *lacZ*+ (black datapoints) or *mhp*T- (blue datapoints) protospacer. Ratios

are shown as a function of the position of the internal AAG sequence within the pre-spacer. **(B)**

Frequency of pre-spacer usage for all pre-spacers on the stp2-mut1 and stp2-mut2 plasmids, for

pre-spacers on the non-target strand with an AAG PAM. The single modified pre-spacer that

differs between stp2-mut1 and stp2-mut2 is shown in orange. The frequency of pre-spacer usage

from the stp2-mut1 and stp2-mut2 plasmids are shown on the x-axis and y-axis, respectively. **(C)**

As for (B) but for the stp2 and stp2-mut3 plasmids.

1066

Figure 14. Sequences at the PAM-distal end of the pre-spacer impact the efficiency of

primed adaptation. (**A**) DNA sequence logo for the 133 "enriched" pre-spacers without an

internal AAG that are used >4-fold more frequently than expected based on the exponential

decay model. Note that the y-axis maximum is set to 0.5 bits. **(B)** DNA sequence logo for the

195 "de-enriched" pre-spacers without an internal AAG that are used >4-fold less frequently

than expected based on the exponential decay model. Note that the y-axis maximum is set to 0.5

bits. **(C)** Schematic showing the unmodified and modified pre-spacer sequences for the stp2,

stp2-mut4, and stp2-mut5 plasmids. The PAM is indicated by a dashed box. Grey highlighting

indicates changes relative to the unmodified pre-spacer. **(D)** Frequency of pre-spacer usage for

all pre-spacers on the stp2 and stp2-mut3 plasmids, for pre-spacers on the non-target strand with

an AAG PAM. The single modified pre-spacer that differs between stp2 and stp2-mut4 is shown

in orange. The frequency of pre-spacer usage from the stp2 and stp2-mut4 plasmids are shown

on the *x*-axis and *y*-axis, respectively. **(E)** As for (D) but for the stp2 and stp2-mut5 plasmids.

1080

A

B

Figure 2

Protospacer
(non-target strand = plus strand)

mhpT-

10 kb

Plus Strand

Minus Strand

Pre-Spacer Usage Frequency

Genome Position

Plus Strand

Minus Strand

lacZ+

Protospacer
(non-target strand = minus strand)

Figure 4

Figure 6

A

B

% of non-AAG
pre-spacers

-1 slip NAAGNNN... 22.8 ± 8.2%

Canonical pre-spacer AAGNNN... N/A

+1 slip AGNNN... 36.7 ± 2.7%

+2 slip GNNN... 1.5 ± 0.2%

C

A

Flip with
-1 slip

```
5'-PAM NNNNNNNNNNNNNNNNNNNNNNNNNNNCTT-3'
3'-CTTNNNNNNNNNNNNNNNNNNNNNNNNNNNNGAA-5'
```

Flip with
no slip

```
5'-PAM NNNNNNNNNNNNNNNNNNNNNNNNNNNNCTT-3'
3'-CTTNNNNNNNNNNNNNNNNNNNNNNNNNNNNGAA-5'
```

Flip with
+1 slip

```
5'-PAM NNNNNNNNNNNNNNNNNNNNNNNNNNNNNCTT-3'
3'-CTTNNNNNNNNNNNNNNNNNNNNNNNNNNNNNGAA-5'
```

B

Figure 9

A

32 nt, +1 slip	**AAG**N NNNNNNNNNNNNNNNNNNNNNNNNNNNNNNN
32 nt, no slip	**AAG**NNNNNNNNNNNNNNNNNNNNNNNNNNNNNN
Canonical length, no slip	**AAG**NNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNN
34 nt, no slip	**AAG**NNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNN
34 nt, -1 slip	N**AAG**NNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNN

B

32 nt pre-spacers

C

34 nt pre-spacers

A

10 kb

lacZ+

codA+

Frequency of pre-spacer usage (minus strand only)

B

Frequency of pre-spacer usage, *lacZ*+

Frequency of pre-spacer usage, *codA*+

188 nt

153 nt

57 nt

Distance from *lacZ*+ protospacer to pre-spacer

A