1   **TRPM4 conductances in thalamic reticular nucleus neurons generate persistent firing**

2   **during slow oscillations**

Abbreviated title: TRPM4 mediates PF in TRN

John J. O'Malley[1,2], Frederik Seibt[1], Jeannie Chin[3], and Michael Beierlein[1,2]

[1]Department of Neurobiology & Anatomy, McGovern Medical School at UTHealth

Houston, TX 77030

[2]MD Anderson Cancer Center UTHealth Graduate School of Biomedical Sciences,

Houston, TX 77030

Department of Neuroscience, Baylor College of Medicine

Houston, TX 77030

Address for Correspondence

Michael Beierlein
Department of Neurobiology & Anatomy
McGovern Medical School
6431 Fannin Street, Suite 7.046
Houston, TX 77030
Email: michael.beierlein@uth.tmc.edu

Number of pages: 43
Number of figures: 7
Abstract: 247 words
Introduction: 541 words
Discussion: 1438 words

Conflict of interests: The authors declare no competing financial interests.

Acknowledgments: Supported by funds from the Harry S. and Isabel C. Cameron Foundation

(J.O.), the Neurodegeneration Consortium and the Belfer Family Foundation (J.C. and M.B.),

NIH NS085171, NS086965 (J.C.), and NIH NS077989 (M.B.). We thank Michael Zhu and Oleh

Pochynyuk for donating mouse models, John Magnotti for statistical advice, and Rajan

Dasgupta and Fabricio Do Monte for helpful discussions.

Abstract

During sleep, neurons in the thalamic reticular nucleus (TRN) participate in distinct types of

oscillatory activity. While the reciprocal synaptic circuits between TRN and sensory relay nuclei

are known to underlie the generation of sleep spindles, the mechanisms regulating slow (<1 Hz)

forms of thalamic oscillations are not well understood. Under *in vitro* conditions, TRN neurons

can generate slow oscillations in a cell-intrinsic manner, with postsynaptic Group 1 metabotropic

glutamate receptor (mGluR) activation leading to the generation of plateau potentials mediated

by both T-type Ca^{2+} currents and Ca^{2+} -activated nonselective cation currents (I_{CAN}). However,

the identity of I_{CAN} and the possible contribution of thalamic circuits to slow rhythmic activity

remain unclear. Using thalamic slices derived from adult mice of either sex, we recorded slow

forms of rhythmic activity in TRN neurons, which were mediated by fast glutamatergic

thalamoreticular inputs but did not require postsynaptic mGluR activation. For a significant

fraction of TRN neurons, synaptic inputs or brief depolarizing current steps led to long-lasting

plateau potentials and persistent firing (PF), and in turn, resulted in sustained synaptic inhibition

in postsynaptic relay neurons of the ventrobasal thalamus (VB). Pharmacological approaches

indicated that plateau potentials were triggered by Ca^{2+} influx through T-type Ca^{2+} channels and

mediated by Ca^{2+} and voltage-dependent transient receptor potential melastatin 4 (TRPM4)

channels. Taken together, our results suggest that thalamic circuits can generate slow

oscillatory activity, mediated by an interplay of TRN-VB synaptic circuits that generate

rhythmicity and TRN cell-intrinsic mechanisms that control PF and oscillation frequency.

Significance Statement

Slow forms of thalamocortical rhythmic activity are thought to be essential for memory consolidation during sleep and the efficient removal of potentially toxic metabolites. *In vivo*, thalamic slow oscillations are regulated by strong bidirectional synaptic pathways linking neocortex and thalamus. Therefore, *in vitro* studies in the isolated thalamus can offer important insights about the ability of individual neurons and local circuits to generate different forms of rhythmic activity. We found that circuits formed by GABAergic neurons in the thalamic reticular nucleus (TRN) and glutamatergic relay neurons in the ventrobasal thalamus generated slow oscillatory activity, which was accompanied by persistent firing in TRN neurons. Our results identify both cell-intrinsic and synaptic mechanisms that mediate slow forms of rhythmic activity in thalamic circuits.

Introduction

Slow (< 1Hz) rhythmic activity in the thalamocortical system is a defining feature of natural sleep and is thought to be essential for the grouping of higher frequency activity such as sleep spindles (Beenhakker and Huguenard, 2009; Lüthi, 2014), network plasticity and memory consolidation (Steriade and Timofeev, 2003; Diekelmann and Born, 2010), and homeostatic processes including the clearance of metabolites (Varga et al., 2016). On a cellular level, slow oscillations are characterized by rapid transitions between periods of sustained depolarizations and persistent firing (Up states) and quiescence (Down states), in both neocortex (Steriade et al., 1993b; Sanchez-Vives and McCormick, 2000) and thalamus (Steriade et al., 1993a; Crunelli and Hughes, 2010). It is well established that neocortex plays an important role in the generation of slow oscillations (Steriade et al., 1993b; Sanchez-Vives and McCormick, 2000; Neske, 2016), with cortical activity leading to the recruitment of thalamic targets via extensive corticothalamic projections (Steriade et al., 1993a; Stroh et al., 2013). However, accumulating evidence shows that rather than being passively entrained by cortical afferents, thalamus can

generate distinct types of rhythmic activity which in turn shape cortical activity patterns (Rigas &

Castro-Alamancos, 2007; Lemieux et al., 2014; Halassa et al., 2011; David et al., 2013;

Fernandez et al., 2018). Thus, rhythms in the thalamocortical system might result from the

interplay of multiple distinct oscillators (Crunelli and Hughes, 2010). While there is strong

consensus on the mechanisms that underlie slow oscillations in neocortex (Crunelli and

Hughes, 2010; Neske, 2016), it is important to gain a better understanding of how thalamic

neurons and circuits generate slow forms of rhythmic activity.

In vitro studies in the isolated thalamus have shown that neurons in the TRN (Blethyn et

al., 2006) and in thalamic relay nuclei (Hughes et al., 2002) can act as cellular pacemakers of

slow rhythms, under conditions of sustained activation of postsynaptic Group I mGluRs by either

exogenous agonists or high frequency corticothalamic activity. This results in an increase in

excitability, due to the closure of a K^+ leak conductance, enabling low threshold Ca^{2+} currents

(I_T) and Ca^{2+}-activated non-selective cation currents (I_{CAN}) to generate long-lasting and rhythmic

plateau potentials, which in TRN neurons lead to persistent firing (Blethyn et al., 2006).

However, the precise role of I_T and the molecular mechanisms mediating I_{CAN} are not known

(Zylberberg & Strowbridge, 2017; Crunelli et al., 2018). Moreover, these findings do not directly

address a potential role of intrathalamic networks. TRN and relay neurons form powerful

reciprocal connections (Gentet and Ulrich, 2003; Pinault, 2004; Pita-Almenar et al., 2014) which

underlie the generation of sleep spindles (Beenhakker and Huguenard, 2009), but whether

these circuits can also generate slower forms of oscillatory activity is not known.

Here we addressed these questions by performing recordings in horizontal slices of

somatosensory thalamus of adult mice. Surprisingly, we observed highly robust slow oscillatory

activity, which was driven by fast synaptic transmission but did not require mGluR activation.

We found that a significant number of TRN neurons displayed synaptically-evoked persistent

firing (PF), which could also be evoked by brief depolarizing current steps in the absence of

synaptic signaling. PF was triggered by Ca^{2+} influx through T-type Ca^{2+} channels and generated

by long-lasting plateau potentials mediated by TRPM4 conductances. Our findings highlight how

intrinsic properties of TRN neurons and intrathalamic synaptic circuits interact to generate slow

thalamic oscillatory activity.

Materials and Methods:

Animals. We employed C57BL6/J mice (JAX Laboratories, Stock No: 000664) unless specified

otherwise. For some experiments, we used TRPC1,4,5,6 general quadruple knockout mice

(Sachdeva et al., 2018) kindly provided by Dr. Michael Zhu or TRPC3 general knockout mice

(Hartmann et al., 2008) kindly provided by Dr. Oleh Pochynyuk. To optogenetically activate

cholinergic inputs to TRN, we used bacterial artificial chromosome (BAC)-transgenic mice

expressing channelrhodopsin (ChR2) under the control of the choline acetyltransferase (ChAT)

promoter (ChAT–ChR2–EYFP; JAX Laboratories, Stock No: 014546, Zhao et al., 2011). All

animals used in this study were treated following procedures in accordance with National

Institutes of Health guidelines and approved by the University of Texas Health Science Center

at Houston (UTHealth) animal welfare committee.

Slice Preparation. Brain slices derived from adult animals (P90-120) were prepared as

described previously (Ting et al., 2014). Briefly, male or female mice were anesthetized using

isoflurane and perfused using an ice cold N-Methyl-D-Glutamine (NMDG) based solution

saturated with 95% O_2–5% CO_2 consisting of the following (in mM): 92 NMDG, 2.5 KCl, 1.25

NaH_2PO_4, 10 $MgSO_4$, 0.5 $CaCl_2$, 30 $NaHCO_3$, 20 glucose, 20 HEPES, 2 thiouera, 5 Na-

ascorbate, and 3 Na-pyruvate. Brains were removed and horizontal slices (300 µm) were cut in

ice-cold NMDG-based solution using a VT1200 S vibratome (Leica). Slices were then held in

NMDG-based solution maintained at 35°C for ~12 minutes before being transferred to a

modified artificial cerebrospinal fluid (ACSF) held at room temperature, consisting of the

following (in mM): 92 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 2 $MgSO_4$, 2 $CaCl_2$, 30 $NaHCO_3$, 25 glucose,

20 HEPES, 2 thiouera, 5 Na-ascorbate, and 3 Na-pyruvate. Slices from younger animals (P13-

38) were prepared as described previously (Agmon & Connors, 1991). Briefly, mice were

anesthetized using isoflurane and then decapitated. Slices were cut in ice-cold cutting solution

saturated with 95% O_2/5% CO_2 consisting of the following (in mM): 213.4 sucrose, 2.5 KCl, 1.25

NaH_2PO_4, 10 $MgSO_4$, 0.5 $CaCl_2$, 26 $NaHCO_3$, and 11 glucose. Slices were then transferred to

ACSF consisting of the following (in mM): 126 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 2 $MgCl_2$, 2 $CaCl_2$,

26 $NaHCO_3$ and 10 glucose, incubated at 35°C for 20 minutes and then held at room

temperature until used for experiments.

Electrophysiological Recordings. Recordings were performed in a chamber perfused with

ACSF held at 32-34°C using a Warner Instruments TC-324B in-line heater. Cells were

visualized via infrared differential interference contrast (IR-DIC) under an Olympus BX51WI

microscope equipped with a Dage-MTI IR-1000 camera. Whole-cell recordings were obtained

using glass pipettes with a tip resistance of 3-5 MΩ. For current clamp recordings in TRN, we

used a potassium-based internal solution consisting of (in mM): 133 K-Gluconate, 1 KCl, 2

$MgCl_2$, 0.16 $CaCl_2$, 10 HEPES, 0.5 EGTA, 2 Mg-ATP, and 0.4 Na-GTP (adjusted to 290 mOsm

and pH 7.3). For voltage clamp recordings of inhibitory postsynaptic currents in neurons of the

ventrobasal nucleus of the thalamus (VB), recording pipettes were filled with a cesium-based

internal solution consisting of (in mM): 120 $CsMeSO_3$, 1 $MgCl_2$, 1 $CaCl_2$, 10 CsCl, 10 HEPES, 3

QX-314, 11 EGTA, 2 Mg-ATP, and 0.3 Na-GTP (adjusted to 295 mOsm and pH 7.3). For loose-

patch recordings of TRN neuronal firing, glass pipettes (3-5 MΩ) were filled with ACSF, and

recordings were obtained in voltage-clamp mode, at seal resistances of 50–200 MΩ. To

minimize any influence on the cell membrane potential, the holding voltage was adjusted

continually to maintain a holding current near 0 pA (Perkins, 2006). In experiments probing TRN

oscillatory activity, glutamine (0.3 mM) was added to the ACSF to prevent a rundown of network

activity (Bryant et al., 2009; Pita-Almenar et al., 2014). For experiments examining the

mechanisms of PF, the bath contained NBQX. In some experiments we replaced extracellular

NaCl with equimolar NMDG and used an internal solution consisting of (in mM): 133 K-

Gluconate, 1 KCl, 4 NaCl, 2 $MgCl_2$, 0.16 $CaCl_2$, 10 HEPES, 0.5 EGTA, 2 Mg-ATP, and 0.4 Na-

GTP (adjusted to 290 mOsm and pH 7.3). For optogenetic activation of cholinergic afferents we

used 1 ms pulses of blue light generated by an LED light source (UHP-T-450-EP, Prizmatix)

delivered through a 60x, 0.9 NA water-immersion objective (Olympus) and centered locally in

the TRN.

140   NBQX, apamin, nimodipine, TTX, 2-Aminoethoxydiphenylborane (2-APB), flufenamic

acid (FFA), 9-phenthranol, glibenclamide, 4-Chloro-2-[[2-(2-chlorophenoxy)acetyl]amino]benzoic

acid (CBA), JNJ 16259685, MTEP hydrochloride, and atropine were obtained from R&D

Systems. ω-conotoxin MVII and TTA-P2 were obtained from Alomone labs. All other chemicals

were obtained from Sigma-Aldrich.

Data acquisition. Recordings were made using a Multiclamp 700B amplifier (Molecular

Devices), filtered at 3–10 kHz, and digitized at 20 kHz with a 16-bit analog-to-digital converter

(Digidata 1440A; Molecular Devices). Data were acquired using Clampex 10.3 software

(Molecular Devices) and analyzed using custom macros written in IGOR Pro (Wavemetrics).

Experimental design and statistical analysis. TRN neuronal firing evoked by brief (25 ms)

current steps was classified into two groups, bursting and persistent firing (PF), based on the

duration of the evoked membrane depolarization measured at half-amplitude (see *Results*).

Post-hoc K means cluster analysis confirmed that classification into two clusters explained the

largest amount of variance. We used a depolarization duration of 200 ms to distinguish between

bursting and PF. Due to rundown of PF in whole-cell mode, data acquisition in the TRN was

limited to 3 min following establishment of whole-cell configuration. Pharmacological

experiments examining the mechanisms underlying PF were carried out by performing

recordings in the presence of a given antagonist under study, and comparing results with

recordings performed under control conditions. All data in a given group were collected from

slices derived from a minimum of 2 animals and reported as mean ± SEM. All statistical

analyses were performed with Prism 7 software. To evaluate significant differences in the

proportion of PF between different experimental groups we performed a chi square test, and

report results in the following format: degrees of freedom in parentheses, followed by the chi-

square value and p value. To test for significant differences in the duration of evoked membrane

depolarization between groups of neurons, we performed an unpaired *t*-test and report results

as: degrees of freedom in parentheses, followed by *t* value and *p* value. Statistical significance

was set at p < 0.05 and adjusted for multiple statistical comparisons with a Bonferroni correction

to correct for pairwise error and to reduce Type I error.

RESULTS

Slow oscillatory activity and persistent firing (PF) in thalamic networks

Following activation of postsynaptic Group I mGluRs, individual neurons in the TRN can

generate slow oscillatory responses, reminiscent of thalamic activity during sleep (Blethyn et al.,

2006). It remains unclear how thalamic synaptic networks contribute to slow oscillatory activity.

To examine this issue, we performed experiments in horizontal thalamic slices derived from

adult (> 3 month) mice. We and others have previously employed extracellular or optogenetic

synaptic stimulation to trigger transient (<2 s) forms of rhythmic thalamic activity *in vitro*

(Huguenard and Prince, 1994; Pita-Almenar et al., 2014). Surprisingly, by performing loose-

patch recordings from TRN neurons, we observed spontaneous and highly rhythmic patterns of

action potential activity that could last several minutes (n = 47 TRN neurons, n = 17 slices). For

12/47 (26%) of recordings, action potential activity was organized as rhythmic bursting

(frequency: 1.9 ± 0.1 Hz, n = 12, Fig. 1A). Interestingly, a second group of recordings (17/47,

36%) revealed rhythmic patterns of sustained action potential activity (> 200 ms duration) we

termed persistent firing (PF), characterized by an initial high-frequency burst, followed by a long-

lasting train of action potentials (average duration: 964.9±256.4 ms, average firing frequency:

175±23.4 Hz, n = 17, Fig. 1B). For neurons displaying rhythmic PF, oscillation frequency was

0.5 ± 0.1 Hz (n = 17), significantly lower compared to the oscillation frequency of cells showing

burst firing ($t(28) = 6.5$, $p = 10^{-16}$, Fig. 1D). The remaining recordings (n=18) showed more

complex patterns of activity, and included neurons that displayed both rhythmic PF and burst

firing (n=4, Fig. 1C). These data indicate that burst firing and PF are prominent firing modes in

the TRN and are associated with oscillatory activity at distinct frequencies.

Relay neurons in the ventrobasal nucleus (VB) receive their sole inhibitory input from

neurons in the TRN. Therefore, the patterns of synaptic inhibitory responses recorded in VB

neurons should closely reflect rhythmic spike firing observed in TRN. To confirm this, we

performed voltage-clamp recordings from VB neurons at a holding potential of 0 mV to isolate

IPSCs. In 76% (n = 22/29) of all VB neurons that showed oscillatory synaptic activity, we

recorded synaptic responses organized as long-lasting high-frequency IPSC barrages (duration:

620.1±54.3 ms, n = 32, Fig. 1F) which occurred rhythmically over several minutes (0.5±0.1 Hz,

n = 32). A smaller fraction of recordings showed rhythmic IPSC bursts (n = 4, Fig.1E). Thus,

relay neurons are the target of persistent inhibition during slow oscillatory activity, likely

mediated by TRN neurons with PF.

Postsynaptic mGluR activation by exogenous agonists or high-frequency stimulation of

glutamatergic afferents can lead to slow oscillatory activity in TRN neurons, even in the absence

of fast synaptic signaling (Blethyn et al., 2006). In contrast, by performing whole-cell recordings

in TRN, we found that rhythmic patterns of PF did not occur spontaneously but instead were

evoked by brief bursts of large-amplitude thalamoreticular EPSPs (Fig. 1G). Furthermore,

rhythmic IPSCs recorded in VB cells were fully blocked by bath application of NBQX (n = 5, Fig.

1H), suggesting that similar to spindle-like activity (Huguenard and McCormick, 2007), slow

oscillatory activity relied on a network of interconnected TRN and relay neurons, at least under

our experimental conditions.

Anatomic and functional studies have shown considerable axonal divergence for TRN to

VB projections, resulting in highly overlapping TRN afferents to neighboring VB neurons

(Pinault, 2004; Pita-Almenar et al., 2014). In agreement, the timing and duration of oscillatory

IPSC barrages were highly synchronous for local VB neuronal pairs (< 50 μm somatic distance)

recorded simultaneously (Fig. 2A). Synchrony was maintained for individual IPSCs throughout

the entire barrage suggesting that the large majority of IPSCs were spike-mediated, without a

detectable contribution of asynchronous GABA release (Hefft & Jonas, 2005).

We found that the majority of VB recordings showed oscillatory IPSC events with little

variability of the initial IPSC amplitude and event duration (Fig. 1E,F), consistent with a single

presynaptic TRN neuron. However, for a number of recordings we observed either two (n=10

VB cells, Fig. 1H and 2B,C) or three (n=4 VB cells, *data not shown*) different types of rhythmic

IPSC events, with each type characterized by a distinct initial IPSC amplitude and event

duration, suggesting convergent inputs from two or three presynaptic TRN neurons,

respectively. For recordings with two types of events, event duration suggested convergent

input by TRN cells that either both displayed PF (n=6, Fig. 2C) or PF and bursting (n=4, Fig.

2B). Importantly, IPSC events evoked by different presynaptic TRN neurons displayed clear

rhythmicity but never occurred synchronously. These data confirm our previous findings derived

in thalamic slices of less mature animals (Pita-Almenar et al., 2014) and suggest that local

thalamic circuits can mediate ongoing oscillatory activity in the absence of widespread

synchrony.

Brief depolarizations trigger PF in TRN neurons

So far our data indicate that a significant fraction of TRN neurons shows PF during network

activity. To better characterize the properties and underlying mechanisms of PF, we performed

whole-cell recordings and blocked network activity by adding NBQX (10 μM) to the bath

solution. Neurons were held at membrane potential of -75 mV unless stated otherwise and

actions potentials were evoked using brief depolarizing current steps (25 ms, 400 pA). Such

stimuli led to bursts of action potentials (average number of spikes: 8.1±0.2, n = 74) in 67% of

neurons recorded (Fig. 3A). For the remaining neurons (37/111, 33%) we observed sustained

action potential activity (average number of spikes: 72.2±9.8, firing frequency: 99.1±7.5 Hz, n =

37) closely resembling PF observed during network activity. PF was mediated by long-lasting

plateau potentials (amplitude from rest: 29.3±0.8 mV, plateau duration: 807.1±41.7 ms, n = 37,

Fig. 3A,B) that terminated in a stepwise manner. Using the duration of the current-step evoked

membrane depolarization, we performed K means cluster analysis, confirming that TRN firing

patterns can be most readily classified into two distinct groups, bursting and PF, with the latter

group displaying membrane depolarizations of > 200 ms. Neurons with current-step evoked PF

were not restricted to a specific TRN subregion and were localized along the entire thickness of

the TRN shell (not shown). We found that PF remained stable for several minutes but then

decayed over time, regardless of composition of the internal solution (Fig. 3C). This might partly

explain why studies employing whole-cell methods have not reported PF in adult TRN neurons

(e.g. Clemente-Perez et al., 2017), while at least one study using sharp electrode recordings

described PF in a subset of neurons of the cat perigeniculate nucleus (Kim and Mccormick,

1998). For the remainder of this study we limited data acquisition to the first 3 minutes in whole-

cell mode.

PF occurs late during development and is controlled by SK conductances

Previous work characterizing TRN firing patterns in slices derived from younger (<P30) animals

has not reported PF (Lee et al., 2007), suggesting that PF is developmentally regulated. To

examine this possibility, we carried out additional recordings in slices prepared from mice aged

P13-18 and P28-38. Compared to adult mice, only 1/24 (4%) of cells at P13-18 and 1/23 (4%) of

cells at P28-38 displayed PF (Fig. 3D-F), indicating that PF is prominent only in fully mature

animals.

The generation of plateau potentials that underlie PF is likely to be tightly regulated by

K^+ conductances, including small conductance Ca^{2+}-activated K^+ channels (SK). To examine

this possibility, we performed recordings in the presence of the SK channel blocker apamin (100

nM). Under these conditions a much larger fraction of TRN cells in adult animals showed PF

(92%, n = 46/50 χ^2(1) = 47.5, p = 10^{-16}, Fig. 3D-F). Thus, excitatory conductances responsible

for PF are widely expressed in the adult TRN but appear to be strongly controlled by SK

conductances. Given the role of SK channels in regulating PF in the adult, the developmental

increase in PF we observed might be solely mediated by a progressive decrease of SK

expression. However, we found that even in the presence of apamin the incidence of PF

remained very low early in development and dramatically increased with animal age (P13-18:

12%, n = 25, P28-38: 41%, n = 24, P90-120: 92%, Fig. 3D-F), suggesting a developmental

upregulation of excitatory mechanisms that underlie PF. Finally, we examined if neurons in the

VB displayed plateau potentials in control or in the presence of apamin. In contrast to TRN, PF

was completely absent from VB under both conditions (Fig. 3E,G). Thus, the mechanisms

underlying PF do not appear to be uniformly expressed in all thalamic nuclei.

Metabotropic glutamate receptor activation is not necessary for PF

Previous work has highlighted a critical role of postsynaptic group I mGluR activation in enabling

intrinsic oscillations and Up states in TRN neurons (Blethyn et al., 2006). While PF under our

experimental conditions did not require exogenous or synaptic mGluR I activation, it is possible

that mGluRs were tonically activated by ambient glutamate (Crabtree et al., 2013) or in a ligand

independent manner (Sun et al., 2016). To examine this possibility, we probed TRN firing in the

presence of the non-competitive mGluR1 antagonist JNJ 16259685 (JNJ) and the non-

competitive mGluR5 antagonist MTEP. Block of group I mGluRs did not significantly change the

incidence of PF (control: 33%, n = 111, JNJ+MTEP: 48%, n = 27, χ^2(1) = 1.7, p = 0.19) or the

plateau potential duration for cells with PF (control: 807.1±41.7 ms, JNJ+MTEP: 599.8±64.3 ms,

t(46) = 1.85, p = 0.071, Fig. 4A,B). These data suggest that under our experimental conditions

Group I mGluR activation is not essential for PF.

Activation of T-type Ca^{2+} channels is required for PF

Next, we examined if PF is voltage-dependent, by evoking action potential activity from

two distinct holding potentials. TRN cells held initially at -60 mV never displayed PF or plateau

potentials (0/15 cells, Fig. 4C,D). However, when the holding potential was adjusted to -75 mV,

8/15 neurons generated PF (χ^2(1) = 1.7, p = 0.0013), suggesting that the underlying

mechanisms of PF are voltage dependent.

To further isolate the mechanisms responsible for PF, we employed a pharmacological

approach. Due to the washout of PF in whole-cell mode, it was not possible to quantify changes

in PF prior to and following bath application of antagonists for a given neuron. Therefore, we

performed all recordings in the presence of a given antagonist and made comparisons to

neurons recorded under control conditions (see *Materials and Methods*). To limit the number of

recordings required for meaningful statistical comparisons, we performed the following

experiments in the presence of apamin, which, as described above, results in PF in 92% of

neurons (Fig. 3E,F).

Previous studies have indicated that in thalamic neurons T-type Ca^{2+} channels are

critical for the generation of mGluR-dependent Up states (Hughes et al., 2002; Blethyn et al.,

2006). The voltage-dependence of PF we observed appears to be consistent with these results.

To confirm the involvement of T-type Ca^{2+} channels in generating PF, we performed recordings

in the presence of the specific antagonist TTA-P2 (1 µM). Under these conditions, PF was

completely eliminated (apamin: 92%, apamin+TTA-P2: 0%, n = 17, $\chi^2(1)$ = 46.5, p = 10^{-20},

Fig. 4E,F). Thus, T-type Ca^{2+} channels are required for the generation of PF.

The plateau potential underlying PF is mediated by a sodium current

It is possible that long-lasting plateau potentials in TRN neurons are entirely mediated by T-type

Ca^{2+} conductances (Zylberberg and Strowbridge, 2017). While these channels show rapid

inactivation following strong depolarization, their inactivation is not complete for smaller

depolarizations near the resting membrane potential, resulting in a steady-state inward

("window") current (Williams et al., 1997). However, PF observed in the present study was

accompanied by large membrane depolarizations that do not appear to be compatible with a

significant role of T-type Ca^{2+} channels. Therefore, we systematically examined the possible

involvement of other voltage- or Ca^{2+}-gated non-inactivating conductances. We first tested the

role of high-threshold Ca^{2+} channels (L, P/Q, and N). In the presence of the selective L-type

antagonist nimodipine (nim, 20 µM, Fig. 5A) the incidence of PF (apamin: 92%, apamin+Nim:

80%, n = 10, $\chi^2(1)$ = 0.3, p = 0.6, Fig. 5B) and plateau potential duration for cells with PF

(apamin: 1108.5±126.6 ms, apamin+Nim: 846.8±235 ms, $t(48)$ = 0.9, p = 0.38, Fig. 5C)

remained comparable to responses recorded in apamin. Similarly, incubating slices in the P/Q

and N antagonist ω-conotoxin MVIIC (ω-Ctx , 1 µM) did not lead to significant changes in the

incidence of PF (apamin: 92%, apamin+ω-Ctx: 70%, n = 10, $\chi^2(1)$ = 1.4, p = 0.23, Fig. 5B), and

plateau potential duration (apamin: 1108.5±126.6 ms, apamin+ω-Ctx: 726.7±196.2 ms, $t(47)$ =

1.2, p = 0.24, Fig. 5C), suggesting that high-threshold Ca^{2+} channels do not play a critical role in

PF.

It is possible that plateau potentials are mediated by activation of a persistent Na^+

current (I_{NaP}) as was suggested previously (Kim and Mccormick, 1998; Fuentealba et al., 2005).

We found that in the presence of apamin and TTX (500 nM), fast action potential activity was

completely blocked, revealing a plateau potential in 9/16 cells (amplitude: 30±1.2 mV, n = 16,

Fig. 5A), a smaller fraction compared to conditions in apamin (apamin: 92%, apamin+TTX: 56%,

$\chi^2(1)$ =10, p = 0.002, Fig. 5B). Plateau potential duration remained comparable to control

(apamin: 1108.5±126.6 ms, apamin+TTX: 912.8±325.7 ms, $t(49)$ = 0.7, p = 0.5, Fig. 5C). These

data indicate that voltage-gated Na$^+$ currents facilitate plateau potential generation, but are not

strictly required.

Next, we examined if Na$^+$ currents other than the ones carried by voltage-gated Na$^+$

channels are essential for plateau potentials, by substituting the majority of extracellular Na$^+$

with NMDG. We found that in the presence of NMDG (together with apamin and TTX) the

incidence of plateau potential generation was significantly reduced (apamin + TTX: 56%,

apamin + TTX + NMDG: 11%, n = 28, $\chi^2(1)$ = 10.6, p = 0.001, Fig. 5A-C), suggesting that PF is

mediated at least in part by a TTX-insensitive Na$^+$ current.

TRPM4 channels underlie PF in the TRN

Our data thus far are consistent with a role of the Ca^{2+}-activated nonselective cationic current

(I_{CAN}) in mediating plateau potentials. Transient receptor potential (TRP) channels have been

previously proposed as molecular mechanisms mediating I_{CAN} (Launay et al., 2002), with

transient receptor potential canonical (TRPC) and melastatin (TRPM) channels as the most

likely candidates. To examine the potential involvement of TRPC channels, we performed

recordings in slices derived from global TRPC3 null mutant mice (TRPC3 KO, Hartmann et al.,

2008) and from global quadruple TRPC1,4,5,6 null mutant mice (QKO, Tian & Zhu, 2018), in the

presence of apamin. For both mouse lines, the incidence of PF (apamin: 92%, TRPC3 KO:

70%, n = 10, $\chi^2(1)$ = 3.4, p = 0.65, QKO: 70%, n = 10, $\chi^2(1)$ = 3.4, p = 0.65, Fig. 6B) and

plateau potential duration (apamin: 1108.5±126.6 ms, TRPC3 KO: 841.1±136 ms, $t(47)$ = 0.9, p

= 0.39, QKO: 752.6±133.4 ms, $t(47)$ = 1.1, p = 0.27, Fig. 6C) were comparable to our

observations in WT animals, suggesting that at least 5 of the 7 TRPCs known to be expressed

in the brain are not involved in mediating PF.

Next, we investigated a possible role of TRPM channels in mediating PF. TRPM2 has

been shown to facilitate burst firing (Lee et al., 2013), while both TRPM4 and TRPM5 have been

implicated in the generation of plateau potentials in cortex (Lei et al., 2014) and of slow inward

currents in cerebellar Purkinje cells (Kim et al., 2013). We found that in the presence of the TRP

channel inhibitor FFA (100 µM), PF was completely eliminated (apamin: 92%, apamin+FFA:

0%, n = 17, $\chi^2(1)$ = 46.6, p = 10^{-20}, Fig. 6A-C). Next, we examined the effects of the broad-

spectrum inhibitor 2-APB which among TRP channels blocks TRPC7 and TRPM2, but does not

inhibit TRPM4 or TRPM5. 2-APB had no effect on PF (apamin: 92%, apamin+ 2-APB: 90%, n =

10, $\chi^2(1)$ = 0.4, p = 0.54, Fig. 6B), with plateau potential duration comparable to control

conditions (apamin: 1108.5±126.6 ms, apamin+2-APB: 758±132.4 ms, $t(49)$ = 1.3, p = 0.22, Fig.

6C), ruling out a role of TRPM2 or TRPC7 and suggesting that either TRPM4 or TRPM5 might

mediate PF. Both channels are voltage-dependent, monovalent-selective, and require increases

in intracellular Ca^{2+} for their activation (Launay et al., 2002; Nilius et al., 2003). We found that in

the presence of the TRPM4/5 antagonists 9-Phenthranol (9-Phen, 100 µM) and glibenclamide

(Glib, 100 µM) PF was completely eliminated (apamin: 92%, apamin+9-Phen: 15%, n = 15,

$\chi^2(1)$ = 34.2, p = 10^{-14}, Glib: 0%, n = 21, $\chi^2(1)$ = 51.4, p = 10^{-26}, Fig. 6B). Next, we performed

experiments using the recently developed selective TRPM4 antagonist CBA (Ozhatil et al.,

2018). In the presence of CBA (50 µM), PF was completely eliminated (apamin: 92%, apamin +

CBA: 0%, n = 15, $\chi^2(1)$ = 44, p = 10^{-17}, Fig. 6A,B). Taken together, our results indicate that

TRPM4 conductances mediate the plateau potential underlying PF.

Since increases in intracellular Ca^{2+} are required for TRPM4 activation, exogenous

buffers such as BAPTA should eliminate PF. However, due to the rundown of PF, experiments

relying on the infusion of Ca^{2+} chelators were not feasible. We therefore probed a possible Ca^{2+}

requirement of PF by replacing extracellular Ca^{2+} with equimolar Ba^{2+}. Ba^{2+} passes through T-

type Ca^{2+} channels and generates large inward currents (Huguenard & Prince, 1992), but unlike

Ca^{2+} does not lead to TRPM4 activation (Nilius et al., 2004; Yamaguchi et al., 2014). Under

these conditions, brief depolarizing steps still led to prominent burst firing, indicating that T-type

Ca^{2+} channel dependent dendritic depolarizations were not affected (Fig. 6A). However, none of

the cells displayed PF (apamin: 92%, apamin+Ba^{2+}: 0%, n = 15, $\chi^2(1)$ = 44, p = 10^{-17}, Fig. 6B,C).

These data further confirm that T-type Ca^{2+} channels alone are not sufficient to generate

plateau potentials. Instead, our data suggest that T-type Ca^{2+} channel activation and the

resulting Ca^{2+} increases lead to the activation of TRPM4 conductances and the generation of

PF.

Synaptic recruitment of muscarinic acetylcholine receptors suppresses PF

Finally, we examined the modulation of PF by synaptic activity. TRN neurons are the target of

cholinergic afferents from the basal forebrain and brainstem (Hallanger and Wainer, 1988). Our

previous studies have shown that acetylcholine (ACh) release leads to a biphasic postsynaptic

response, with fast excitation mediated by nicotinic ACh receptors (nAChRs) and slow and long-

lasting inhibition mediated by M2 muscarinic receptors (mAChRs) coupled G protein-gated

inward rectifying K^+ (GIRK) channels (Sun et al., 2013). To examine if cholinergic inputs can

trigger PF in the TRN, we performed experiments using BAC-ChAT-ChR2 mice that selectively

express ChR2 in cholinergic neurons (Zhao et al., 2011) and optogenetically evoked ACh

release with light pulses (1 ms) centered over the recorded neuron. To avoid decay of PF, TRN

neurons were recorded in loose-patch mode. Under these conditions, 2/5 of neurons displayed

PF (Fig. 7A,C). We reasoned that the activation of mAChRs and the resulting GIRK activation

might limit a full manifestation of PF. Indeed, following atropine application to block mAChRs,

the duration of action potential firing was significantly prolonged in all neurons recorded (control:

213.5±94 ms, atropine: 416.3±84.1 ms, $t(4)$ = 7.1, p = 0.002, Fig. 7A-C), and the incidence of

PF increased from 40% to 80%. These data show that the activation of K^+ channels by

synaptically released neuromodulators such as ACh can dynamically modulate the expression

of PF.

Discussion

Here we have identified TRPM4 as the mechanism underlying PF in adult neurons of the TRN.

Our data indicate that the large depolarizations and intracellular Ca^{2+} increases required for

TRPM4 activation are mediated by T-type Ca^{2+} channels. PF was associated with robust slow

oscillatory activity in thalamic circuits which required fast synaptic transmission but did not rely

on mGluR receptor signaling. Previous studies have shown that both TRN and thalamic relay

neurons can act as cellular pacemakers for slow thalamic oscillations. Our present findings

highlight intrathalamic circuits as an alternative mechanism capable of generating slow rhythmic

activity.

Mechanisms of PF in the TRN

Our results extend recent work that has implicated TRPM4 in controlling neuronal excitability in

cortex (Lei et al., 2014; Riquelme et al., 2018), hippocampus (Menigoz et al., 2016), substantia

nigra (Mrejeru et al., 2011), and brainstem (Picardo et al., 2019). A unique property of TRPM4

activation within physiological ranges is the requirement for both depolarization and large

increases in intracellular Ca^{2+} concentration (Launay et al., 2002; Nilius et al., 2003; Nilius et al.,

2004). Since dendritic Ca^{2+} signals can experience decay under whole-cell conditions, this might

partly explain why we consistently observed rundown of PF in the present study. Alternatively,

recordings in whole-cell mode might have led to the dialysis of critical signaling molecules.

TRPM4 conductances rapidly desensitize in excised patches (Zhang et al., 2005) and in whole-

cell mode, but modulators such as PIP2 can restore TRPM4 activity. Additional investigations

into the mechanisms of TRPM4 regulation under physiological conditions will be important.

What are the Ca^{2+} sources of TRPM4 activation? We found that pharmacological block

of T-type Ca^{2+} channel using the specific antagonist TTA-P2 completely eliminated PF. In

addition, replacing Ca^{2+} with Ba^{2+} abolished PF. These findings suggest that the Ca^{2+} increases

required for TRPM4 activation are mediated by influx through T-type Ca^{2+} channels although we

cannot rule out a contribution of R-type Ca^{2+} channels (Zaman et al., 2011) or release from

internal stores (Neyer et al., 2016). Ca$_V$3.3 channels mediating I$_T$ are primarily expressed in the

distal dendrites of TRN neurons and their activation during burst firing results in large dendritic

but more modest somatic Ca^{2+} signals (Crandall et al., 2010; Astori et al., 2011). This would

suggest a dendritic location of TRPM4. In this scenario, membrane depolarizations necessary

for TRPM4 activation would likely be mediated by synaptically-generated T-type dependent Ca^{2+}

spikes, which in contrast to fast Na$^+$ action potentials, propagate effectively into TRN dendrites

(Crandall et al., 2010; Connelly et al., 2015).

We found that in the presence of apamin, PF could be observed in most recordings,

suggesting that TRPM4 conductances are expressed in the large majority of TRN neurons.

Thus, SK channels control both the initiation and duration of PF, likely due to the close

functional association with T-type Ca^{2+} channels (Cueni et al., 2008). Similarly, we found that

the synaptic activation of M2 muscarinic receptors curtailed the duration of PF. It is therefore

likely that under physiological conditions, TRPM4 activation and the generation of PF in a given

neuron does not occur in all-or-none manner, but instead is constantly regulated by multiple

synaptic and intrinsic processes. For example, an increased likelihood and duration of PF might

be promoted by mechanisms that increase T-type mediated Ca^{2+} increases or directly facilitate

TRPM4 activation, and curtailed by modulatory synaptic systems that reduce TRN excitability

via opening of specific K conductances.

We want to emphasize that our findings do not provide evidence for a well-defined cell

type that displays PF. Previous studies have used molecular markers (Clemente-Perez et al.,

2017) or firing patterns (Lee et al., 2007) to delineate TRN neuronal subtypes that differ in the

expression of T-type Ca^{2+} channels, which might be an important factor determining the

incidence of PF. However, it is likely that a wide range of factors including the spatial

arrangement and density of postsynaptic receptors, intrinsic conductances, and intracellular

Ca^{2+} signaling mechanisms interact to control PF generation in a given cell.

Comparison to previous studies

Plateau potentials and PF have been observed throughout the brain, though in most cases

there is little consensus on the underlying mechanisms (Zylberberg and Strowbridge, 2017). In

the thalamus, Kim and McCormick (1998) reported plateau potentials in a fraction of neurons of

the perigeniculate sector of the ferret TRN, with very similar properties as described here.

However, they concluded that plateau potentials were mediated by I_{NaP}, since they were fully

blocked by TTX. The same conclusion was reached in an *in vivo* study of TRN in anesthetized

cats, which demonstrated block of persistent firing using intracellular dialysis with QX-314

(Fuentealba et al., 2005). Here we find that in apamin a significant number of TRN neurons

shows plateau potentials even in the presence of TTX. It is possible that under physiological

conditions transient or persistent forms of voltage-gated I_{Na} indirectly aid PF by increasing

activation of TRPM4 and T-type Ca^{2+} channels.

For both TRN and thalamic relay neurons, group I mGluR activation can trigger rhythmic

forms of PF, mediated by the closure of a K^+ leak conductance, allowing the generation of a

long-lasting inward current mediated by the non-inactivating portion of T-type Ca^{2+} channel

conductance (Hughes et al., 2002; Blethyn et al., 2006). Our present findings differ from these

results in four important aspects: First, PF did not require mGluR activation and the resulting

closure of K^+ leak conductances. Second, in the absence of mGluR activation, PF did not occur

in a cell-intrinsic manner but required fast synaptic inputs. Under our experimental conditions,

both cholinergic inputs and bursts of glutamatergic thalamic inputs triggered PF, likely by

generating global Ca^{2+} spikes (Connelly et al., 2015). Third, while T-type Ca^{2+} channels were

important for the initiation of PF, our results are inconsistent with a critical role of I_T in generating

long-lasting currents that lead to plateau potentials. We found that plateau potentials reached

levels of ~-45 mV, where T-type Ca^{2+} conductances at steady-state experience near full

inactivation. Furthermore, the lack of PF in the presence of Ba^{2+}, which should allow for the

generation of a I_T-mediated window current (Huguenard and Prince, 1992) suggests that T-type

Ca^{2+} currents alone are unlikely to mediate PF. Fourth, using the potent and selective

antagonist CBA (Ozhatil et al., 2018), we showed that I_{CAN} was essential for the generation of

plateau potentials and was mediated by TRPM4. It should be stated that the majority of TRP

antagonists employed in previous studies have a number of non-specific effects, so the

validation of CBA as a selective TRPM4 antagonist under our experimental conditions will

require independent confirmation.

504 Mechanisms of slow thalamic oscillations

Accumulating evidence indicates that both cortex and thalamus can generate oscillations in

isolation, suggesting that rhythmic activity in the intact thalamocortical system results from the

complex interplay of multiple distinct oscillators (Crunelli and Hughes, 2010). While there has

been extensive research on the cortical mechanisms mediating slow rhythms, the nature of

thalamic pacemakers are less well understood. Previous work has shown that both TRN and

relay neurons are capable of generating cell-intrinsic rhythms in the 1 Hz range, but only under

conditions of postsynaptic mGluR activation (Hughes et al., 2002; Blethyn et al., 2006). Under

physiological conditions, this would require sustained corticothalamic activity, indicating that

both thalamic cell types act as conditional oscillators. Our findings derived from adult thalamic

slices highlight an alternative mechanism and suggest that networks of TRN and VB cells can

generate robust and long-lasting slow oscillatory activity, mediated by powerful bidirectional

synaptic connectivity (Gentet and Ulrich, 2003; Pinault, 2004; Pita-Almenar et al., 2014). We

observed an inverse relationship between TRN firing duration and oscillatory frequency. It is

therefore tempting to suggest that changes in TRN activity as seen during PF acts to reduce the

frequency of thalamic oscillations, via recruitment of GABA$_B$Rs (Kim et al., 1997) or

extrasynaptic high-affinity GABA$_A$Rs (Herd et al., 2013), thereby changing the latency to

rebound burst generation (Schofield et al., 2009).

Our network data suggest that robust thalamic rhythmicity can occur in the absence of

precise synchrony, highlighting that both processes can occur independently. While thalamic

oscillatory activity is commonly referred to as synchronous (Fogerson and Huguenard, 2016),

there have been few studies that have directly examined the degree of precise thalamic

synchronization during distinct behavioral states. Our current data showing a lack of precise

synchrony in local thalamic circuits confirm previous findings (Pita-Almenar et al., 2014) and

indicate that the variability in TRN firing might act as an additional factor in preventing

synchrony in isolated thalamic networks. As has been suggested (Contreras and Steriade,

1996), this would argue that synchronization of thalamic networks is primarily regulated by

afferent inputs, particularly from cortex.

References:

Agmon A, Connors BW (1991) Thalamocortical responses of mouse somatosensory (barrel)

cortex *in vitro*. Neuroscience 41:365–379 Available at:

http://www.ncbi.nlm.nih.gov/pubmed/1870696.

Astori S, Wimmer RD, Prosser HM, Corti C, Corsi M, Liaudet N, Volterra A, Franken P, Adelman

JP, Lüthi A (2011) The Ca v3.3 calcium channel is the major sleep spindle pacemaker in

thalamus. Proc Natl Acad Sci U S A 108:13823–13828.

Beenhakker MP, Huguenard JR (2009) Neurons that Fire Together Also Conspire Together: Is

Normal Sleep Circuitry Hijacked to Generate Epilepsy? Neuron 62:612–632 Available at:

http://dx.doi.org/10.1016/j.neuron.2009.05.015.

Blethyn KL, Hughes SW, Tóth TI, Cope DW, Crunelli V (2006) Neuronal basis of the slow (<1

Hz) oscillation in neurons of the nucleus reticularis thalami *in vitro*. J Neurosci 26:2474–

2486.

Bryant AS, Li B, Beenhakker MP, Huguenard JR (2009) Maintenance of thalamic epileptiform

activity depends on the astrocytic glutamate-glutamine cycle. J Neurophysiol 102:2880–

2888.

Clemente-Perez A, Makinson SR, Higashikubo B, Brovarney S, Cho FS, Urry A, Holden SS,

Wimer M, Dávid C, Fenno LE, Acsády L, Deisseroth K, Paz JT (2017) Distinct Thalamic

Reticular Cell Types Differentially Modulate Normal and Pathological Cortical Rhythms.

Cell Rep.

Connelly WM, Crunelli V, Errington AC (2015) The global spike: Conserved dendritic properties

enable unique Ca^{2+} spike generation in low-threshold spiking neurons. J Neurosci

35:15505–15522.

Contreras D, Steriade M (1996) Synchronization of low-frequency rhythms in corticothalamic

networks. Neuroscience 76:11–24.

Crabtree JW, Lodge D, Bashir ZI, Isaac JTR (2013) GABAA, NMDA and mGlu2 receptors

tonically regulate inhibition and excitation in the thalamic reticular nucleus. Eur J Neurosci

37:850–859.

Crandall SR, Govindaiah G, Cox CL (2010) Low-threshold Ca 2+ current amplifies distal

dendritic signaling in thalamic reticular neurons. J Neurosci 30:15419–15429.

Crunelli V, Hughes SW (2010) The slow (1 Hz) rhythm of non-REM sleep: A dialogue between

three cardinal oscillators. Nat Neurosci 13:9–17.

Crunelli V, Larincz ML, Connelly WM, David F, Hughes SW, Lambert RC, Leresche N, Errington

AC (2018) Dual function of thalamic low-vigilance state oscillations: Rhythm-regulation and

plasticity. Nat Rev Neurosci 19:107–118.

Cueni L, Canepari M, Luján R, Emmenegger Y, Watanabe M, Bond CT, Franken P, Adelman

JP, Lüthi A (2008) T-type Ca^{2+} channels, SK2 channels and SERCAs gate sleep-related

oscillations in thalamic dendrites. Nat Neurosci 11:683–692.

David F, Schmiedt JT, Taylor HL, Orban G, Di Giovanni G, Uebele VN, Renger JJ, Lambert RC,

Leresche N, Crunelli V (2013) Essential thalamic contribution to slow waves of natural

sleep. J Neurosci 33:19599–19610.

Diekelmann S, Born J (2010) The memory function of sleep. Nat Rev Neurosci 11:114–126

Available at: http://dx.doi.org/10.1038/nrn2762.

Fernandez LMJ, Vantomme G, Osorio-Forero A, Cardis R, Béard E, Lüthi A (2018) Thalamic

reticular control of local sleep in mouse sensory cortex. Elife 7:1–25.

Fogerson PM, Huguenard JR (2016) Tapping the Brakes: Cellular and Synaptic Mechanisms

that Regulate Thalamic Oscillations. Neuron 92:687–704 Available at:

http://dx.doi.org/10.1016/j.neuron.2016.10.024.

Fuentealba P, Timofeev I, Bazhenov M, Sejnowski TJ, Steriade M (2005) Membrane bistability

in thalamic reticular neurons during spindle oscillations. J Neurophysiol 93:294–304.

Gentet LJ, Ulrich D (2003) Strong, reliable and precise synaptic connections between thalamic

relay cells and neurones of the nucleus reticularis in juvenile rats. J Physiol.

Halassa MM, Siegle JH, Ritt JT, Ting JT, Moore CI (2011) Selective optical drive of thalamic

reticular nucleus generates thalamic bursts & cortical spindles. Nat Neurosci 14:1118–

1120.

Hallanger AE, Wainer BH (1988) Ultrastructure of ChAT-immunoreactive synaptic terminals in

the thalamic reticular nucleus of the rat. J Comp Neurol.

Hartmann J, Dragicevic E, Adelsberger H, Henning HA, Sumser M, Abramowitz J, Blum R,

Dietrich A, Freichel M, Flockerzi V, Birnbaumer L, Konnerth A (2008) TRPC3 Channels Are

Required for Synaptic Transmission and Motor Coordination. Neuron 59:392–398.

Hefft S, Jonas P (2005) Asynchronous GABA release generates long-lasting inhibition at a

hippocampal interneuron-principal neuron synapse. Nat Neurosci 8:1319–1328.

Herd MB, Brown AR, Lambert JJ, Belelli D (2013) Extrasynaptic GABAA receptors couple

presynaptic activity to postsynaptic inhibition in the somatosensory thalamus. J Neurosci

33:14850–14868.

Hughes SW, Cope DW, Blethyn KL, Crunelli V (2002) Cellular Mechanisms of the Slow (<1Hz)

Oscillations in Thalamocortical Neurons *in Vitro*. Neuron 33:947–958 Available at:

http://www.sciencedirect.com/science/article/pii/S0896627302006232.

Huguenard JR, McCormick DA (2007) Thalamic synchrony and dynamic regulation of global

forebrain oscillations. Trends Neurosci 30:350–356.

Huguenard JR, Prince DA (1992) A novel T-type current underlies prolonged Ca^{2+}-dependent

burst firing in GABAergic neurons of rat thalamic reticular nucleus. J Neurosci 12:3804–

3817.

Huguenard JR, Prince DA (1994) Intrathalamic rhythmicity studied *in vitro*: Nominal T-current

modulation causes robust antioscillatory effects. J Neurosci 14:5485–5502.

Kim U, Mccormick DA (1998) Functional and ionic properties of a slow afterhyperpolarization in

ferret perigeniculate neurons *in vitro*. J Neurophysiol 80:1222–1235.

Kim U, Sanchez-Vives M V., McCormick DA (1997) Functional dynamics of GABAergic

inhibition in the thalamus. Science (80-) 278:130–134.

Kim YS, Kang E, Makino Y, Park S, Shin JH, Song H, Launay P, Linden DJ (2013)

Characterizing the conductance underlying depolarization-induced slow current in

cerebellar Purkinje cells. J Neurophysiol 109:1174–1181.

Launay P, Fleig A, Perraud AL, Scharenberg AM, Penner R, Kinet JP (2002) TRPM4 is a Ca^{2+}-

activated nonselective cation channel mediating cell membrane depolarization. Cell

109:397–407.

Lee CR, Machold RP, Witkovsky P, Rice ME (2013) TRPM2 channels are required for NMDA-

induced burst firing and contribute to H2O2-dependent modulation in substantia nigra pars

reticulata GABAergic neurons. J Neurosci 33:1157–1168.

Lee SH, Govindaiah G, Cox CL (2007) Heterogeneity of firing properties among rat thalamic

reticular nucleus neurons. J Physiol.

Lei YT, Thuault SJ, Launay P, Margolskee RF, Kandel ER, Siegelbaum SA (2014) Differential

contribution of TRPM4 and TRPM5 nonselective cation channels to the slow

afterdepolarization in mouse prefrontal cortex neurons. Front Cell Neurosci 8:1–14.

Lemieux M, Chen JY, Lonjers P, Bazhenov M, Timofeev I (2014) The impact of cortical

deafferentation on the neocortical slow oscillation. J Neurosci 34:5689–5703.

Lüthi A (2014) Sleep spindles: Where they come from, what they do.

Menigoz A, Ahmed T, Sabanov V, Philippaert K, Pinto S, Kerselaers S, Segal A, Freichel M,

Voets T, Nilius B, Vennekens R, Balschun D (2016) TRPM4-dependent post-synaptic

depolarization is essential for the induction of NMDA receptor-dependent LTP in CA1

hippocampal neurons. Pflugers Arch Eur J Physiol 468:593–607.

Mrejeru A, Wei A, Ramirez JM (2011) Calcium-activated non-selective cation currents are

involved in generation of tonic and bursting activity in dopamine neurons of the substantia

nigra pars compacta. J Physiol 589:2497–2514.

Neske GT (2016) The slow oscillation in cortical and thalamic networks: Mechanisms and

functions. Front Neural Circuits 9:1–25.

Neyer C, Herr D, Kohmann D, Budde T, Pape HC, Coulon P (2016) MGluR-mediated calcium

signalling in the thalamic reticular nucleus. Cell Calcium 59:312–323 Available at:

http://dx.doi.org/10.1016/j.ceca.2016.03.009.

Nilius B, Prenen J, Droogmans G, Voets T, Vennekens R, Freichel M, Wissenbach U, Flockerzi

V (2003) Voltage dependence of the Ca^{2+}-activated cation channel TRPM4. J Biol Chem

278:30813–30820.

Nilius B, Prenen J, Voets T, Droogmans G (2004) Intracellular nucleotides and polyamines

inhibit the Ca^{2+}- activated cation channel TRPM4b. Pflugers Arch Eur J Physiol 448:70–75.

Ozhatil LC, Delalande C, Bianchi B, Nemeth G, Kappel S, Thomet U, Ross-Kaschitza D,

Simonin C, Rubin M, Gertsch J, Lochner M, Peinelt C, Reymons J-L, Abriel H (2018)

Identification of potent and selective small molecule inhibitors of the cation channel

TRPM4. Br J Pharmacol 175:2504–2519.

Perkins KL (2006) Cell-attached voltage-clamp and current-clamp recording and stimulation

techniques in brain slices. J Neurosci Methods.

Picardo MCD, Sugimura YK, Dorst KE, Kallurkar PS, Akins VT, Ma X, Teruyama R, Guinamard

R, Kam K, Saha MS, Del Negro CA (2019) Trpm4 ion channels in pre-bötzinger complex

interneurons are essential for breathing motor pattern but not rhythm. PLoS Biol 17:1–30.

Pinault D (2004) The thalamic reticular nucleus: Structure, function and concept.

Pita-Almenar JD, Yu D, Lu HC, Beierlein M (2014) Mechanisms Underlying Desynchronization

of Cholinergic-Evoked Thalamic Network Activity. J Neurosci 34:14463–14474.

Rigas P, Castro-Alamancos MA (2007) Thalamocortical up states: Differential effects of intrinsic

and extrinsic cortical inputs on persistent activity. J Neurosci 27:4261–4272.

Riquelme D, Silva I, Philp AM, Huidobro-Toro JP, Cerda O, Trimmer JS, Leiva-Salcedo E

(2018) Subcellular localization and activity of TRPM4 in medial prefrontal cortex layer 2/3.

Front Cell Neurosci 12:1–13.

Sachdeva R, Schlotterer A, Schumacher D, Matka C, Mathar I, Dietrich N, Medert R, Kriebs U,

Lin J, Nawroth P, Birnbaumer L, Fleming T, Hammes HP, Freichel M (2018) TRPC proteins

contribute to development of diabetic retinopathy and regulate glyoxalase 1 activity and

methylglyoxal accumulation. Mol Metab 9:156–167 Available at:

https://doi.org/10.1016/j.molmet.2018.01.003.

Sanchez-Vives M V., McCormick DA (2000) Cellular and network mechanisms of rhytmic

recurrent activity in neocortex. Nat Neurosci 3:1027–1034.

Schofield CM, Kleiman-Weiner M, Rudolph U, Huguenard JR (2009) A gain in GABA A receptor

synaptic strength in thalamus reduces oscillatory activity and absence seizures. Proc Natl

Acad Sci U S A 106:7630–7635.

Steriade M, Contreras D, Dossi RC, Nunez A (1993a) The slow (<1 Hz) oscillation in reticular

thalamic and thalamocortical neurons: Scenario of sleep rhythm generation in interacting

thalamic and neocortical networks. J Neurosci 13:3284–3299.

Steriade M, Nunez A, Amzica F (1993b) A novel slow (< 1 Hz) oscillation of neocortical neurons

in vivo: Depolarizing and hyperpolarizing components. J Neurosci 13:3252–3265.

Steriade M, Timofeev I (2003) Neuronal plasticity in thalamocortical networks during sleep and

waking oscillations. Neuron 37:563–576.

Stroh A, Adelsberger H, Groh A, Rühlmann C, Fischer S, Schierloh A, Deisseroth K, Konnerth A

(2013) Making Waves: Initiation and Propagation of Corticothalamic Ca^{2+} Waves *In Vivo*.

Neuron 77:1136–1150.

Sun Y-G, Pita-Almenar JD, Wu C-S, Renger JJ, Uebele VN, Lu H-C, Beierlein M (2013)

Biphasic cholinergic synaptic transmission controls action potential activity in thalamic

reticular nucleus neurons. J Neurosci 33:2048–2059 Available at:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3711637&tool=pmcentrez&rend

ertype=abstract.

Sun YG, Rupprecht V, Zhou L, Dasgupta R, Seibt F, Beierlein M (2016) MGluR1 and mGluR5

synergistically control cholinergic synaptic transmission in the thalamic reticular nucleus. J

Neurosci 36:7886–7896.

Tian J, Zhu M (2018) GABAB Receptors Augment TRPC3-Mediated Slow Excitatory

Postsynaptic Current to Regulate Cerebellar Purkinje Neuron Response to Type-1

Metabotropic Glutamate Receptor Activation. Cells 7:90.

Ting JT, Daigle T, Chen Q, Feng G (2014) Acute brain slice method for adult and aging animals:

application of targeted patch clamp analysis and optogenetics. In: Methods in Molecular

Biology, Patch-Clam. (Martina M, Taverna S, eds), pp 221–242. New York: Humana Press.

Varga AW et al. (2016) Reduced Slow-Wave Sleep Is Associated with High Cerebrospinal Fluid

Aβ42 Levels in Cognitively Normal Elderly. Sleep 39:2041–2048.

Williams SR, Tóth TI, Turner JP, Hughes SW, Crunelli V (1997) The "window" component of the

low threshold Ca^{2+} current produces input signal amplification and bistability in cat and rat

thalamocortical neurones. J Physiol.

Yamaguchi S, Tanimoto A, Otsuguro KI, Hibino H, Ito S (2014) Negatively charged amino acids

near and in transient receptor potential (TRP) domain of TRPM4 channel are one

determinant of its Ca^{2+} sensitivity. J Biol Chem 289:35265–35282.

Zaman T, Lee K, Park C, Paydar A, Choi JH, Cheong E, Lee CJ, Shin HS (2011) CaV2.3

Channels Are Critical for Oscillatory Burst Discharges in the Reticular Thalamus and

Absence Epilepsy. Neuron 70:95–108 Available at:

http://dx.doi.org/10.1016/j.neuron.2011.02.042.

Zhang Z, Okawa H, Wang Y, Liman ER (2005) Phosphatidylinositol 4,5-Bisphosphate rescues

TRPM4 channels from desensitization. J Biol Chem 280:39185–39192.

Zhao S, Ting JT, Atallah HE, Qiu L, Tan J, Gloss B, Augustine GJ, Deisseroth K, Luo M,

Graybiel AM, Feng G (2011) Cell type-specific channelrhodopsin-2 transgenic mice for

optogenetic dissection of neural circuitry function. Nat Methods 8:745–755.

Zylberberg J, Strowbridge BW (2017) Mechanisms of persistent activity in cortical circuits:

Possible neural substrates for working memory. Annu Rev Physiol 40:60. – 627.

   Figure legends:

Figure 1. Rhythmic activity in thalamic networks. A. Loose-patch recording of TRN neuron

showing rhythmic burst firing. **B.** *Top,* TRN neuron showing rhythmic persistent firing (PF).

Bottom, Close-up highlighting an individual barrage of PF. **C.** TRN neuron showing rhythmic

activity consisting of both burst firing and PF. **D.** Summary data (n = 29) plotting the average

duration of TRN firing as a function of oscillation frequency, for neurons with either burst firing or

PF. Data points in red denote cells shown in A and B. Dashed line marks criterion separating

between rhythmic bursting and PF. **E.** Voltage-clamp recording of VB neuron held at 0 mV

displaying rhythmic bursts of IPSCs. **F.** *Top,* Recording of VB neuron displaying rhythmic

barrages of IPSCs. *Bottom*, close-up shows an individual barrage of IPSCs. **G.** *Top*, Current

clamp recording of TRN neuron, showing PF during network activity. *Bottom*, Close up shows

that bursts of EPSCs (marked by red arrows) trigger PF. **H.** Representative recording of

rhythmic barrages of IPSCs in VB neuron, under control conditions and following bath

application of NBQX (10 µM). Similar data were obtained from 4 other VB neurons.

Figure 2. Local TRN neurons display rhythmicity in the absence of precise synchrony. VB

recordings were performed in voltage clamp, at a holding potential of 0 mV. **A.** Divergent TRN

output to VB neurons. *Left*, Dual recordings of neighboring VB cells showing a single barrage of

IPSCs. *Middle*, Close-up of IPSCs (normalized to the initial IPSC amplitude) highlighting precise

IPSC synchrony across both cells. *Right*, Schematic outlines proposed divergent TRN

projections giving rise to VB responses. **B.** TRN neurons with PF and burst firing form

convergent input onto VB neuron**.** *Left,* Voltage-clamp recording of VB neuron with IPSC bursts

(red tick marks) and prolonged IPSC barrages (cyan tick marks) with distinct initial amplitudes.

Middle, For the same recording, overlay of representative IPSC events and summary data

(n=40 for each event) quantifying the initial amplitude of IPSC events**.** *Right*, Schematic of

proposed circuit. **C.** Two TRN neurons with PF form convergent input onto VB neuron**.** *Left*,

Recording of VB neuron showing IPSC barrages with two distinct initial amplitudes (green and

purple tick marks, respectively). *Middle*, For the same recording, overlay of representative IPSC

events and summary data (n=40 for each event) quantifying the initial amplitude of IPSC events.

Right, Schematic of proposed circuit.

Figure 3. PF occurs late in development and regulated by SK conductances. A. Brief

depolarizing current steps (25 ms, 400 pA, RMP = -75 mV) lead to two distinct firing patterns,

Persistent Firing (PF, *top*) or Burst Firing (Bu, *bottom*). **B.** Summary data (n = 111) showing

bimodal distribution of current-step evoked membrane depolarizations. Grey line indicates

criterion separating PF and Bu. **C.** Raster plot of AP activity in representative TRN neuron

showing rundown of PF. **D.** Representative recordings of TRN neurons with PF at three

developmental stages, in control conditions (*black*) and in the presence of apamin (500 nM,

red). **D.** Representative recordings of TRN neurons with PF at three developmental stages, in

control (black) and in the presence of apamin (500 nM, *red*). **E.** Representative recordings of VB

cells, in control (black) and in the presence of apamin (red) show lack of PF in both conditions.

F. Summary data showing incidence of PF in TRN across development (P13-18: n = 24 in

control, n = 25 in apamin, P28-38: n = 23 in control, n = 22 in apamin, P90-120: n = 111 in

control, n = 50 in apamin) and in VB (n = 15 in control, n = 16 in apamin). **G.** For the same cells

as in F, summary data quantifying membrane potential duration for all TRN cells across

development, in control conditions and in apamin.

Figure 4. T-type Ca^{2+} channels but not mGluRs are required for PF. A. Representative

recordings of PF from two different TRN neurons, in control conditions and in the presence of

the mGluR1 antagonist JNJ 16259685 (1 µM) and the mGluR5 antagonist MTEP (10 µM). **B.**

Summary data quantifying duration of evoked membrane depolarization in control (n = 111) and

in the presence of mGluR 1/5 antagonists (n = 23). **C.** Representative recording of neuron with

PF, initially held at a holding potential of -60 mV and then at -75 mV. **D.** Summary data

quantifying evoked depolarization duration as a function of holding potential (n = 15 TRN

neurons). **E.** Representative recordings in the presence of apamin and in the presence of

apamin and the T-type Ca^{2+} channel antagonist TTA-P2 (1 µM). **F.** Summary data quantifying

evoked depolarization duration (n = 50 in apamin, n = 17 in apamin + TTA-P2).

Figure 5. A TTX-insensitive sodium current generates plateau potential. All recordings

were performed in the presence of apamin. **A.** Representative TRN recordings in the presence

of the L-type Ca^{2+} channel antagonist nimodipine (20 µM, *top*), in TTX (500 nM, *middle*), and in

the presence of TTX and NMDG replacing extracellular Na^+ (*bottom*). **B.** Summary data

showing incidence of PF in the presence of nimodipine (Nim, n = 10), the P/Q- and N-type Ca^{2+}

channel antagonist ω-conotoxin MVIIC (ω-Ctx, 1 µM, n = 10), TTX (n = 16), and TTX and

NMDG (n = 28). Dashed line indicates incidence of PF in apamin. **C.** Summary data from the

same cells as in B quantifying evoked depolarization duration in each condition. Grey solid line

represents criterion for PF. * denotes statistical significance of $p < 0.05$ compared to recordings

in apamin group, # denotes statistical significance of $p < 0.05$ compared to recordings in TTX.

Figure 6. TRPM4 conductances mediate PF. All recordings were performed in the presence

of apamin. **A.** Representative recordings in control and in the presence of the non-selective

transient potential receptor (TRP) channel antagonist FFA (100 µM, *top*), the TRPM4 antagonist

CBA (50 µM, *middle*), and in 2 mM $BaCl_2$ replacing extracellular $CaCl_2$ (*bottom*). **B.** Summary

data showing incidence of PF in TRPC3 KO (KO, n = 10 cells) in TRPC1,4,5,6 quadruple KO

(QKO, n = 10 cells), FFA (n = 17), the non-selective TRP channel antagonist 2-APB (50 µM, n =

10), the TRPM4/5 antagonists 9-phenthranol (9-Phen, 100 µM, n = 15) and glibenclamide (Glib,

100 µM, n = 21), CBA (n = 15), and Ba^{2+}, (n = 15). Grey dashed line represents incidence of PF

in apamin. **C.** For the same cells as in B, summary data quantifying the evoked depolarization

duration. * denotes statistical significance of $p < 0.05$ compared to recordings in apamin.

Figure 7. Synaptic recruitment of muscarinic AChRs suppresses PF. Recordings were

carried out in slices of ChAT-ChR2-EYFP BAC mice. **A.** Loose-patch recording of a TRN neuron

showing brief barrage of ACh-evoked activity (*black*), which is prolonged following application of

the mAChR antagonist atropine (10 μM, *blue*). **B.** Raster plot of ACh-evoked neuronal activity

prior to and following atropine application, for the cell shown in A. **C.** Summary data quantifying

the average duration of ACh-evoked action potential activity before and after atropine

application (n = 5). * $p < 0.05$

Figure 1. Rhythmic activity in thalamic networks. A. Loose-patch recording of TRN neuron showing rhythmic burst firing. **B.** *Top,* TRN neuron showing rhythmic persistent firing (PF). *Bottom*, Close-up highlighting an individual barrage of PF. **C.** TRN neuron showing rhythmic activity consisting of both burst firing and PF. **D.** Summary data (n = 29) plotting the average duration of TRN firing as a function of oscillation frequency, for neurons with either burst firing or PF. Data points in red denote cells shown in A and B. Dashed line marks criterion separating between rhythmic bursting and PF. **E.** Voltage-clamp recording of VB neuron held at 0 mV displaying rhythmic bursts of IPSCs. **F.** *Top,* Recording of VB neuron displaying rhythmic barrages of IPSCs. *Bottom*, close-up shows an individual barrage of IPSCs. **G.** *Top*, Current clamp recording of TRN neuron, showing PF during network activity. *Bottom*, Close up shows that bursts of EPSCs (marked by red arrows) trigger PF. **H.** Representative recording of rhythmic barrages of IPSCs in VB neuron, under control conditions and following bath application of NBQX (10 μM). Similar data were obtained from 4 other VB neurons.

Figure 2. Local TRN neurons display rhythmicity in the absence of precise synchrony. VB
recordings were performed in voltage clamp, at a holding potential of 0 mV. **A.** Divergent TRN
output to VB neurons. *Left*, Dual recordings of neighboring VB cells showing a single barrage of
IPSCs. *Middle*, Close-up of IPSCs (normalized to the initial IPSC amplitude) highlighting precise
IPSC synchrony across both cells. *Right*, Schematic outlines proposed divergent TRN
projections giving rise to VB responses. **B.** TRN neurons with PF and burst firing form
convergent input onto VB neuron. *Left,* Voltage-clamp recording of VB neuron with IPSC bursts
(red tick marks) and prolonged IPSC barrages (cyan tick marks) with distinct initial amplitudes.
Middle, For the same recording, overlay of representative IPSC events and summary data
(n=40 for each event) quantifying the initial amplitude of IPSC events. *Right*, Schematic of
proposed circuit. **C.** Two TRN neurons with PF form convergent input onto VB neuron. *Left*,
Recording of VB neuron showing IPSC barrages with two distinct initial amplitudes (green and
purple tick marks, respectively). *Middle*, For the same recording, overlay of representative IPSC
events and summary data (n=40 for each event) quantifying the initial amplitude of IPSC events.
Right, Schematic of proposed circuit.

Figure 3. PF occurs late in development and regulated by SK conductances. A. Brief
depolarizing current steps (25 ms, 400 pA, RMP = -75 mV) lead to two distinct firing patterns,
Persistent Firing (PF, *top*) or Burst Firing (Bu, *bottom*). **B.** Summary data (n = 111) showing bimodal

distribution of current-step evoked membrane depolarizations. Grey line indicates criterion
separating PF and Bu. **C.** Raster plot of AP activity in representative TRN neuron showing rundown
of PF. **D.** Representative recordings of TRN neurons with PF at three developmental stages, in
control conditions (*black*) and in the presence of apamin (500 nM, *red*). **D.** Representative
recordings of TRN neurons with PF at three developmental stages, in control (black) and in the
presence of apamin (500 nM, *red*). **E.** Representative recordings of VB cells, in control (black) and
in the presence of apamin (red) show lack of PF in both conditions. **F.** Summary data showing
incidence of PF in TRN across development (P13-18: n = 24 in control, n = 25 in apamin, P28-38:
n = 23 in control, n = 22 in apamin, P90-120: n = 111 in control, n = 50 in apamin) and in VB (n =
15 in control, n = 16 in apamin). **G.** For the same cells as in F, summary data quantifying membrane
potential duration for all TRN cells across development, in control conditions and in apamin.

Figure 4. T-type Ca^{2+} channels but not mGluRs are required for PF. A. Representative
recordings of PF from two different TRN neurons, in control conditions and in the presence of
the mGluR1 antagonist JNJ 16259685 (1 µM) and the mGluR5 antagonist MTEP (10 µM). **B.**
Summary data quantifying duration of evoked membrane depolarization in control (n = 111) and
in the presence of mGluR 1/5 antagonists (n = 23). **C.** Representative recording of neuron with
PF, initially held at a holding potential of -60 mV and then at -75 mV. **D.** Summary data
quantifying evoked depolarization duration as a function of holding potential (n = 15 TRN
neurons). **E.** Representative recordings in the presence of apamin and in the presence of
apamin and the T-type Ca^{2+} channel antagonist TTA-P2 (1 µM). **F.** Summary data quantifying
evoked depolarization duration (n = 50 in apamin, n = 17 in apamin + TTA-P2).

859
860
Figure 5. A TTX-insensitive sodium current generates plateau potential. All recordings
were performed in the presence of apamin. **A.** Representative TRN recordings in the presence
of the L-type Ca^{2+} channel antagonist nimodipine (20 µM, *top*), in TTX (500 nM, *middle*), and in
the presence of TTX and NMDG replacing extracellular Na^+ (*bottom*). **B.** Summary data
showing incidence of PF in the presence of nimodipine (Nim, n = 10), the P/Q- and N-type Ca^{2+}
channel antagonist ω-conotoxin MVIIC (ω-Ctx, 1 µM, n = 10), TTX (n = 16), and TTX and
NMDG (n = 28). Dashed line indicates incidence of PF in apamin. **C.** Summary data from the
same cells as in B quantifying evoked depolarization duration in each condition. Grey solid line
represents criterion for PF. * denotes statistical significance of $p < 0.05$ compared to recordings
in apamin group, # denotes statistical significance of $p < 0.05$ compared to recordings in TTX.

Figure 6. TRPM4 conductances mediate PF. All recordings were performed in the presence
of apamin. **A.** Representative recordings in control and in the presence of the non-selective
transient potential receptor (TRP) channel antagonist FFA (100 μM, *top*), the TRPM4 antagonist
CBA (50 μM, *middle*), and in 2 mM BaCl$_2$ replacing extracellular CaCl$_2$ (*bottom*). **B.** Summary
data showing incidence of PF in TRPC3 KO (KO, n = 10 cells) in TRPC1,4,5,6 quadruple KO
(QKO, n = 10 cells), FFA (n = 17), the non-selective TRP channel antagonist 2-APB (50 μM, n =
10), the TRPM4/5 antagonists 9-phenthranol (9-Phen, 100 μM, n = 15) and glibenclamide (Glib,
100 μM, n = 21), CBA (n = 15), and Ba^{2+}, (n = 15). Grey dashed line represents incidence of PF
in apamin. **C.** For the same cells as in B, summary data quantifying the evoked depolarization
duration. * denotes statistical significance of $p < 0.05$ compared to recordings in apamin.

Figure 7. Synaptic recruitment of muscarinic AChRs suppresses PF. Recordings were
carried out in slices of ChAT-ChR2-EYFP BAC mice. **A.** Loose-patch recording of a TRN neuron
showing brief barrage of ACh-evoked activity (*black*), which is prolonged following application of
the mAChR antagonist atropine (10 µM, *blue*). **B.** Raster plot of ACh-evoked neuronal activity
prior to and following atropine application, for the cell shown in A. **C.** Summary data quantifying
the average duration of ACh-evoked action potential activity before and after atropine
application (n = 5). * $p < 0.05$