1   **TRPM4 conductances in thalamic reticular nucleus neurons generate persistent firing**

2   **during slow oscillations**

Abbreviated title: TRPM4 mediates PF in TRN

John J. O'Malley[1,2], Frederik Seibt[1], Jeannie Chin[3], and Michael Beierlein[1,2]

[1]Department of Neurobiology & Anatomy, McGovern Medical School at UTHealth

Houston, TX 77030

[2]MD Anderson Cancer Center UTHealth Graduate School of Biomedical Sciences,

Houston, TX 77030

Department of Neuroscience, Baylor College of Medicine

Houston, TX 77030

Address for Correspondence

Michael Beierlein
Department of Neurobiology & Anatomy
McGovern Medical School
6431 Fannin Street, Suite 7.046
Houston, TX 77030
Email: michael.beierlein@uth.tmc.edu

Number of pages: 43
Number of figures: 7
Abstract: 246 words
Introduction: 580 words
Discussion: 1255 words

Conflict of interests: The authors declare no competing financial interests.

Acknowledgments: Supported by funds from the Harry S. and Isabel C. Cameron Foundation

(J.O.), the Neurodegeneration Consortium and the Belfer Family Foundation (J.C. and M.B.),

NIH NS085171, NS086965 (J.C.), and NIH NS077989 (M.B.). We thank Michael Zhu and Oleh

Pochynyuk for help with mouse models, John Magnotti for statistical advice and Fabricio Do

Monte for helpful discussions.

Abstract

During slow-wave sleep (SWS) and anesthesia, thalamocortical networks generate slow (< 1

Hz) oscillations, with activity in individual neurons showing rhythmic fluctuations between Up

and Down States. Neocortical Up states are driven by persistent synaptic activity in recurrent

networks. In contrast, rhythmic up states in thalamic neurons are thought to be primarily

mediated by cell-intrinsic mechanisms, with postsynaptic Group 1 metabotropic glutamate

receptor (mGluR) activation leading to the generation of long-lasting plateau potentials mediated

by both T-type Ca^{2+} currents and calcium-activated nonselective cation currents (I_{CAN}). However,

the molecular identity of I_{CAN} and the role of thalamic synaptic networks in generating slow

oscillations remain unclear. By performing recordings in thalamic slices derived from adult mice

of either sex, we observed robust slow oscillatory network activity that did not require mGluR

signaling but was completely eliminated in the absence of fast glutamatergic signaling. We

found that thalamic glutamatergic inputs triggered persistent firing (PF) in neurons of the

thalamic reticular nucleus (TRN). For about 30% of TRN neurons, PF could also be evoked by

brief depolarizing steps, while the remaining neurons displayed burst firing. PF was driven by

long-lasting plateau potentials that were triggered by Ca^{2+} influx through T-type Ca^{2+} channels

and mediated by Ca^{2+} and voltage-dependent transient receptor potential melastatin 4 (TRPM4)

channels. Taken together, our results suggest that slow oscillatory activity in TRN neurons

typical for SWS is mediated by cell-intrinsic mechanisms that generate PF, and synaptic

networks interconnecting TRN with thalamic relay nuclei that generate rhythmicity.

Significance Statement

Slow-wave sleep (SWS) is essential for memory consolidation and efficient removal of potentially toxic metabolites. During SWS, thalamic neurons display slow rhythmic activity, and fluctuate between periods of sustained activity and complete quiescence. The cellular and network contributions to slow thalamic oscillations remain poorly understood. We found that, in thalamic slice preparations of adult mice, thalamic reticular nucleus (TRN) neurons show spontaneous slow oscillations accompanied by persistent firing (PF). While thalamic synaptic networks were critical for oscillatory activity, T-type Ca^{2+} and TRPM4 channels in TRN neurons were responsible for initiation and maintenance of PF, respectively. Our findings help to elucidate the mechanisms involved in TRN oscillatory activity during SWS.

Introduction

A defining feature of natural sleep and anesthesia are slow (< 1Hz) and synchronized rhythms in the thalamocortical system. Rather than acting as a "default state" in the absence of behaviorally relevant sensory inputs, slow oscillations during sleep are thought to be critical for the grouping of higher frequency activity such as sleep spindles (Lüthi, 2014), network plasticity and memory consolidation (Steriade & Timofeev, 2003; Diekelmann & Born, 2010), and for homeostatic processes including the clearance of metabolites (Varga et al., 2016). On a cellular level, slow oscillations are characterized by rapid transitions between periods of sustained depolarizations (Up states) and quiescence (Down states), in both neocortex (Steriade et al., 1993b; Sanchez-Vives and McCormick, 2000) and thalamus (Steriade et al., 1993a; Crunelli and Hughes, 2010). In neocortex, slow oscillations result from activity in extensively interconnected networks of excitatory neurons, leading to long-lasting, rhythmic barrages of synaptic activity which are highly synchronous across large neuronal populations (Sanchez-Vives & McCormick, 2000). It was originally assumed that slow oscillations initiate in neocortex and then spread to thalamic targets (Steriade et al., 1993a). Neocortical slow oscillations persist

in the absence of thalamic inputs (Steriade et al., 1993c; Sanchez-Vives and McCormick, 2000)

and can entrain thalamic activity through extensive corticothalamic projections (Stroh et al.,

2013). However, there is accumulating evidence indicating that thalamic circuits can generate

different types of oscillations, which in turn shape cortical activity (Rigas & Castro-Alamancos,

2007; Lemieux et al., 2014; Halassa et al., 2011; David et al., 2013; Fernandez et al., 2018).

One well-established form of rhythmic activity initiating in the thalamus are sleep spindles (7-14

Hz), produced by the reciprocal synaptic interactions of two populations of neurons that show

bursting activity during sleep, thalamocortical relay cells and inhibitory thalamic reticular nucleus

(TRN) neurons (Beenhakker & Huguenard, 2009). The cellular and network mechanisms that

underlie slower forms of thalamic oscillatory activity seen during SWS are not clearly defined

(Crunelli and Hughes, 2010). In contrast to neocortex, slow rhythms in the isolated thalamus are

not assumed to require synaptic networks but instead are thought to be generated by cell-

intrinsic mechanisms (Hughes et al., 2002; Blethyn et al., 2006). For both TRN and thalamic

relay cells, activation of postsynaptic Group I mGluRs by either exogenous agonists or high

frequency corticothalamic activity results in an increase in excitability, due to the closure of a K^+

leak conductance. This enables low threshold Ca^{2+} currents (I_T) and Ca^{2+}-activated non-

selective cation currents (I_{CAN}) to generate long-lasting and rhythmic plateau potentials, which in

TRN neurons lead to persistent firing (Blethyn et al., 2006). However, the precise role of I_T and

the molecular mechanisms mediating I_{CAN} are not known (Zylberberg & Strowbridge, 2017;

Crunelli et al., 2018). Furthermore, the requirements for both corticothalamic and intra-thalamic

synaptic activity in generating slow thalamic oscillations remain unclear.

Here we addressed these questions by performing recordings in horizontal slices of

somatosensory thalamus of adult mice. Surprisingly, we observed highly robust slow oscillatory

thalamic network activity in the absence of mGluR activation. However, slow oscillations were

completely eliminated by antagonists for fast synaptic transmission. We found that a significant

number of TRN neurons displayed synaptically-evoked persistent firing (PF), which could also

be evoked by brief depolarizing current steps in the absence of synaptic signaling. PF was

triggered by Ca^{2+} influx through T-type Ca^{2+} channels and generated by long-lasting plateau

potentials mediated by TRPM4 conductances. Our findings highlight how intrinsic properties of

TRN neurons and intra-thalamic synaptic networks interact to generate slow thalamic oscillatory

activity.

Materials and Methods:

Animals. We employed C57BL6/J mice (JAX Laboratories, Stock No: 000664) unless specified

otherwise. For some experiments, we used TRPC1,4,5,6 general quadruple knockout mice

(Sachdeva et al., 2018) kindly provided by Dr. Michael Zhu or TRPC3 general knockout mice

(Hartmann et al., 2008) kindly provided by Dr. Oleh Pochynyuk. To optogenetically activate

cholinergic inputs to TRN, we used bacterial artificial chromosome (BAC)-transgenic mice

expressing channelrhodopsin (ChR2) under the control of the choline acetyltransferase (ChAT)

promoter (ChAT–ChR2–EYFP; JAX Laboratories, Stock No: 014546, Zhao et al., 2011). All

animals used in this study were treated following procedures in accordance with National

Institutes of Health guidelines and approved by the University of Texas Health Science Center

at Houston (UTHealth) animal welfare committee.

Slice Preparation. Brain slices derived from adult animals (P90-120) were prepared as

described previously (Ting et al., 2014). Briefly, male or female mice were anesthetized using

isoflurane and perfused using an ice cold N-Methyl-D-Glutamine (NMDG) based solution

saturated with 95% O_2–5% CO_2 consisting of the following (in mM): 92 NMDG, 2.5 KCl, 1.25

NaH_2PO_4, 10 $MgSO_4$, 0.5 $CaCl_2$, 30 $NaHCO_3$, 20 glucose, 20 HEPES, 2 thiouera, 5 Na-

ascorbate, and 3 Na-pyruvate. Brains were removed and horizontal slices (300 µm) were cut in

ice-cold NMDG-based solution using a VT1200 S vibratome (Leica). Slices were then held in

NMDG-based solution maintained at 35°C for ~12 minutes before being transferred to a

modified artificial cerebrospinal fluid (ACSF) held at room temperature, consisting of the

following (in mM): 92 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 2 $MgSO_4$, 2 $CaCl_2$, 30 $NaHCO_3$, 25 glucose,

20 HEPES, 2 thiouera, 5 Na-ascorbate, and 3 Na-pyruvate. Slices from younger animals (P13-

38) were prepared as described previously (Agmon & Connors, 1991). Briefly, mice were

anesthetized using isoflurane and then decapitated. Slices were cut in ice-cold cutting solution

saturated with 95% O_2/5% CO_2 consisting of the following (in mM): 213.4 sucrose, 2.5 KCl, 1.25

NaH_2PO_4, 10 $MgSO_4$, 0.5 $CaCl_2$, 26 $NaHCO_3$, and 11 glucose. Slices were then transferred to

ACSF consisting of the following (in mM): 126 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 2 $MgCl_2$, 2 $CaCl_2$,

26 $NaHCO_3$ and 10 glucose, incubated at 35°C for 20 minutes and then held at room

temperature until used for experiments.

Electrophysiological Recordings. Recordings were performed in a chamber perfused with

ACSF held at 32-34°C using a Warner Instruments TC-324B in-line heater. Cells were

visualized via infrared differential interference contrast (IR-DIC) under an Olympus BX51WI

microscope equipped with a Dage-MTI IR-1000 camera. Whole-cell recordings were obtained

using glass pipettes with a tip resistance of 3-5 MΩ. For current clamp recordings, we used a

potassium-based internal solution consisting of (in mM): 133 K-Gluconate, 1 KCl, 2 $MgCl_2$, 0.16

$CaCl_2$, 10 HEPES, 0.5 EGTA, 2 Mg-ATP, and 0.4 Na-GTP (adjusted to 290 mOsm and pH 7.3).

For voltage clamp recordings of inhibitory postsynaptic currents in neurons of the ventrobasal

nucleus of the thalamus (VB), recording pipettes were filled with a cesium-based internal

solution consisting of (in mM): 120 $CsMeSO_3$, 1 $MgCl_2$, 1 $CaCl_2$, 10 CsCl, 10 HEPES, 3 QX-314,

11 EGTA, 2 Mg-ATP, and 0.3 Na-GTP (adjusted to 295 mOsm and pH 7.3). For loose-patch

recordings, 3-5 MΩ pipettes were filled with ASCF. To probe oscillatory network activity,

Glutamine (0.3 mM) was added to the ACSF (Bryant et al., 2009; Pita-Almenar et al., 2014). In

some experiments we replaced extracellular NaCl with equimolar NMDG and used an internal

solution consisting of (in mM): 133 K-Gluconate, 1 KCl, 4 NaCl, 2 MgCl$_2$, 0.16 CaCl$_2$, 10

HEPES, 0.5 EGTA, 2 Mg-ATP, and 0.4 Na-GTP (adjusted to 290 mOsm and pH 7.3). For

optogenetic experiments cholinergic afferents were activated using 1 ms pulses of blue light

using an LED light source (UHP-T-450-EP, Prizmatix) delivered through a 60x, 0.9 NA water-

immersion objective (Olympus) and centered locally in the TRN.

NBQX, apamin, nimodopine, nifedopine, TTX, 2-APB, flufenamic acid, 9-phenthranol,

glibenclamide, CBA, and atropine were obtained from R&D Systems. ω-conotoxin MVII and

TTA-P2 were obtained from Alomone labs. All other chemicals were obtained from Sigma-

Aldrich.

Data acquisition. Recordings were made using a Multiclamp 700B amplifier (Molecular

Devices), filtered at 3–10 kHz, and digitized at 20 kHz with a 16-bit analog-to-digital converter

(Digidata 1440A; Molecular Devices). Data were acquired using Clampex 10.3 software

(Molecular Devices) and analyzed using custom macros written in IGOR Pro (Wavemetrics).

Experimental design and statistical analysis. TRN neuronal firing evoked by brief (25 ms)

current steps was classified into two groups, bursting and persistent firing (PF), based on the

duration of the evoked membrane depolarization measured at half-amplitude (see *Results*).

Post-hoc K means cluster analysis confirmed that classification into two clusters explained the

largest amount of variance. We used a depolarization duration of 200 ms to distinguish between

bursting and PF. Due to rundown of PF in whole-cell mode, data acquisition in the TRN was

limited to 3 min following establishment of whole-cell configuration. Pharmacological

experiments examining the mechanisms underlying PF were carried out by performing

recordings in the presence of a given antagonist under study, and comparing results with

recordings performed under control conditions. All data in a given group were collected from

slices derived from a minimum of 2 animals and reported as mean ± SEM. All statistical

analyses were performed with Prism 7 software. To evaluate significant differences in the

proportion of PF between different experimental groups we performed a chi square test, and

report results in the following format: degrees of freedom in parentheses, followed by the chi-

square value and p value. To test for significant differences in the duration of evoked membrane

depolarization between groups of neurons, we performed an unpaired *t*-test and report results

as: degrees of freedom in parentheses, followed by *t* value and *p* value. Statistical significance

was set at p < 0.05 and adjusted for multiple statistical comparisons with a Bonferroni correction

to correct for pairwise error and to reduce Type I error.

RESULTS

Slow oscillatory activity and persistent firing (PF) in thalamic networks

Following activation of postsynaptic Group I mGluRs, individual neurons in the TRN and the

adjacent relay nuclei can generate slow oscillatory responses, reminiscent of thalamic activity

during SWS (Blethyn et al., 2006). It remains unclear how thalamic synaptic networks contribute

to slow oscillatory activity. To examine this issue, we performed experiments in horizontal

thalamic slices derived from adult (> 3 month) mice. We and others have previously employed

extracellular or optogenetic synaptic stimulation to trigger transient (<2 s) forms of rhythmic

thalamic activity *in vitro* (Huguenard and Prince, 1994; Pita-Almenar et al., 2014). Surprisingly,

by performing loose-patch recordings from TRN neurons, we observed spontaneous and highly

rhythmic patterns of action potential activity that could last several minutes (n = 47 TRN

neurons, n = 17 slices). For 12/47 (26%) of recordings, action potential activity was organized

as rhythmic bursting (frequency: 1.9 ± 0.1 Hz, n = 12, Fig. 1A). Interestingly, a second group of

recordings (17/47, 36%) revealed rhythmic patterns of sustained action potential activity we

termed persistent firing (PF), characterized by an initial high-frequency burst, followed by a long-

lasting train of action potentials (duration: 964.9±256.4 ms, average firing frequency: 175±23.4

Hz, n = 17, Fig. 1B). For neurons displaying rhythmic PF, oscillation frequency was 0.5 ± 0.1 Hz

(n = 17), significantly lower compared to the oscillation frequency of cells showing burst firing

(t(28) = 6.5, p = 10^{-16}; Fig. 1E). The remaining recordings showed more complex patterns of

rhythmic activity that included single spikes, bursts and/or PF (n = 18, *data not shown*). These

data indicate the presence of two firing modes in the adult TRN, burst firing and PF, which are

associated with oscillatory activity at distinct frequencies.

Relay neurons in the ventrobasal nucleus (VB) receive their sole inhibitory input from

neurons in the TRN. Therefore, the patterns of synaptic inhibitory responses recorded in VB

neurons should closely reflect rhythmic spike firing observed in TRN. To confirm this, we

performed voltage-clamp recordings from VB neurons at a holding potential of 0 mV to isolate

IPSCs. In 76% (n = 22/29) of all VB neurons that showed oscillatory synaptic activity indicative

of intact thalamic networks, we recorded synaptic responses organized as long-lasting high-

frequency IPSC barrages (duration: 620.1±54.3 ms, n = 32, Fig. 1D) which occurred

rhythmically over several minutes (0.5±0.1 Hz, n = 32). A smaller fraction of recordings showed

rhythmic IPSC bursts (n = 4, Fig.1C). Thus, relay neurons are the target of persistent inhibition

during slow oscillatory activity, likely mediated by TRN neurons with PF.

Postsynaptic mGluR activation by exogenous agonists or high-frequency stimulation of

glutamatergic afferents can lead to slow oscillatory activity in TRN neurons, even in the absence

of fast synaptic signaling (Blethyn et al., 2006). In contrast, by performing whole-cell recordings

in TRN, we found that rhythmic patterns of PF did not occur spontaneously but instead were

evoked by brief bursts of large-amplitude thalamoreticular EPSPs (Fig. 1F). Furthermore,

rhythmic IPSCs recorded in VB cells were fully blocked by bath application of NBQX (n = 5, Fig.

1G), suggesting that similar to spindle-like activity (Huguenard and McCormick, 2007), slow

oscillatory activity relied on a network of interconnected TRN and relay neurons, at least under

our experimental conditions.

Anatomic and functional studies have shown considerable axonal divergence for TRN to

VB projections, resulting in highly overlapping TRN afferents to neighboring VB neurons

(Pinault, 2004; Pita-Almenar et al., 2014). In agreement, the timing and duration of oscillatory

IPSC barrages were highly synchronous for VB neuronal pairs recorded simultaneously (Fig.

2A). Synchrony was maintained for individual IPSCs throughout the entire barrage suggesting

that the large majority of IPSCs were spike-mediated, without a detectable contribution of

asynchronous GABA release (Hefft & Jonas, 2005).

We found that the majority of VB recordings showed oscillatory IPSC barrages with little

variability of the initial IPSC amplitude, consistent with activity of a single presynaptic TRN

neuron (Fig. 1C,D). However, some recordings contained two distinct types of barrages, with

each barrage showing a distinct initial IPSC amplitude and duration Figs. 1G, 2B,C). These

responses are most readily explained as being generated by convergent inputs from two

presynaptic TRN neurons with distinct firing patterns, including bursting and PF. Importantly,

IPSC bursts and barrages evoked by independent TRN neurons displayed clear rhythmicity but

never occurred synchronously. These data confirm our previous findings derived in thalamic

slices of less mature animals (Pita-Almenar et al., 2014) and suggest that local thalamic

networks can mediate robust oscillatory activity in the absence of widespread synchrony.

Brief depolarizations trigger PF in TRN neurons

So far our data indicate that a significant fraction of TRN neurons shows PF during network

activity. To better characterize the properties and underlying mechanisms of PF, we performed

whole-cell recordings and blocked network activity by adding NBQX to the bath solution. Brief

depolarizing current steps (25 ms, 400 pA) from a resting potential of -75 mV generated bursts

of action potentials (average number of spikes: 8.1±0.2, n = 74) in 67% of neurons recorded

(Fig. 3A). For the remaining neurons (37/111, 33%) we observed sustained action potential

activity (average number of spikes: 72.2±9.8; firing frequency: 99.1±7.5 Hz, n = 37) closely

resembling PF observed during network activity. PF was mediated by long-lasting plateau

potentials (amplitude from rest: 29.3±0.8 mV, plateau duration: 807.1±41.7 ms, n = 37, Fig.

3A,B) that terminated in a stepwise manner. Using the duration of the current-step evoked

membrane depolarization, we performed K means cluster analysis, confirming that TRN firing

patterns are most readily classified into two distinct groups, bursting and PF, with the latter

group displaying membrane depolarizations of > 200 ms. We found that PF remained stable for

several minutes but then decayed over time, regardless of composition of the internal solution

(Fig. 3C). Therefore, for the remainder of this study we limited data acquisition to the first 3

minutes in whole-cell mode.

PF occurs late during development and is controlled by SK conductances

Previous studies performed under similar experimental conditions have not reported PF in

preparations derived from younger animals (<P30). To determine if PF is developmentally

regulated, we carried out additional recordings in slices prepared from mice aged P13-18 and

P28-38. Compared to adult mice, only 1/24 (4%) of cells at P13-18 and 1/23 (4%) of cells at

P28-38 displayed PF (Fig. 3D-F), indicating that PF is prominent only in fully mature animals.

The generation of plateau potentials that underlie PF is likely to be tightly regulated by

K^+ conductances, including small conductance Ca^{2+}-activated K^+ channels (SK). To examine

this possibility, we performed recordings in the presence of the SK channel blocker apamin (100

nM). Compared to control conditions, a much larger fraction of TRN cells in adult animals

showed PF (92%, n = 46/50 $\chi^2(1)$ = 47.5, p = 10^{-16}; Fig. 3D-F). Thus, excitatory conductances

responsible for PF are widely expressed in the adult TRN but appear to be strongly controlled

by SK conductances. Given the role of SK channels in regulating PF in the adult, the

developmental increase in PF we observed might be solely mediated by a progressive

downregulation of SK conductances. However, we found that even in the presence of apamin

the incidence of PF remained very low early in development and dramatically increased with

animal age (P13-18: 12%, n = 25, P28-38: 41%, n = 24, P90-120: 92%; Fig. 3D-F), suggesting a

developmental upregulation of excitatory mechanisms that underlie PF. Finally, we examined if

neurons in the ventrobasal nucleus of the thalamus (VB) displayed plateau potentials in the

presence of apamin. In contrast to TRN, PF was completely absent from VB (0/11 cells; *data*

not shown). Thus, the mechanisms underlying PF do not appear to be uniformly expressed in all

thalamic nuclei.

Metabotropic glutamate receptor activation is not necessary for PF

Previous work has highlighted a critical role of postsynaptic group I mGluR activation in enabling

intrinsic oscillations and Up states in TRN neurons (Blethyn et al., 2006). While PF under our

experimental conditions did not require exogenous or synaptic mGluR I activation, it is possible

that mGluRs were tonically activated by ambient glutamate (Crabtree et al., 2013) or in a ligand

independent manner (Sun et al., 2016). To examine this possibility, we probed TRN firing in the

presence of the non-competitive mGluR1 antagonist JNJ 16259685 (JNJ) and the non-

competitive mGluR5 antagonist MTEP. Block of group I mGluRs did not significantly change the

incidence of PF (Control: 33%, n = 111; JNJ+MTEP: 48%, n = 27, $\chi^2(1)$ = 1.7, p = 0.19) or the

plateau potential duration for cells with PF (Control: 807.1±41.7 ms, JNJ+MTEP: 599.8±64.3

ms, $t(46)$ = 1.85, p = 0.071; Fig. 4A,B). These data suggest that under our experimental

conditions Group I mGluR activation is not essential for PF.

Activation of T-type calcium channels is required for PF

Next, we examined if PF is voltage-dependent, by evoking action potential activity from

two distinct holding potentials. TRN cells held initially at -60 mV never displayed PF or plateau

potentials (0/15 cells; Fig. 4C,D). However, when the holding potential was adjusted to -75 mV,

8/15 neurons generated PF ($\chi^2(1)$ = 1.7, p = 0.0013), suggesting that the underlying

mechanisms of PF are voltage dependent.

To further isolate the mechanisms responsible for PF, we employed a pharmacological

approach. Due to the washout of PF in whole-cell mode, it was not possible to quantify changes

in PF prior to and following bath application of antagonists for a given neuron. Therefore, we

performed all recordings in the presence of a given antagonist and made comparisons to

neurons recorded under control conditions (see *Materials and Methods*). To limit the number of

recordings required for meaningful statistical comparisons, we performed the following

experiments in the presence of apamin, which, as described above, results in PF in 92% of

neurons (Fig. 3E,F).

Previous studies have indicated that T-type Ca^{2+} channels are critical for the generation

of mGluR-dependent Up states (Hughes et al., 2002; Blethyn et al., 2006). The voltage-

dependence of PF we observed appears to be consistent with these results. To confirm the

involvement of T-type Ca^{2+} channels in generating PF, we performed recordings in the presence

of the specific antagonist TTA-P2 (1 µM). Under these conditions, depolarizing current steps still

led to action potential firing, but PF was completely eliminated (Apamin: 92%, Apamin+TTA-P2:

0%, n = 17, $\chi^2(1)$ = 46.5, *p* = 10^{-20}; Fig. 4E,F). Thus, T-type Ca^{2+} channels are required for the

generation of PF.

The plateau potential underlying PF is mediated by a sodium current

It is possible that long-lasting plateau potentials in TRN neurons are entirely mediated by T-type

Ca^{2+} conductances (Zylberberg and Strowbridge, 2017). While these channels show rapid

inactivation following strong depolarization, their inactivation is not complete for smaller

depolarizations near the resting membrane potential, resulting in a steady-state inward

("window") current (Williams et al., 1997). However, PF observed in the present study was

accompanied by large membrane depolarizations that do not appear to be compatible with a

significant role of T-type Ca^{2+} channels. Therefore, we systematically examined the possible

involvement of other voltage- or Ca^{2+}-gated non-inactivating conductances. We first tested the

role of high-threshold Ca^{2+} channels (L, P/Q, and N). In the presence of the selective L-type

antagonist Nimodipine (Nim, 20 µM; Fig. 5A) the incidence of PF (Apamin: 92%, Apamin+Nim:

80%, n = 10, $\chi^2(1)$ = 0.3, p = 0.6; Fig. 5B) and plateau potential duration for cells with PF

(Apamin: 1108.5±126.6 ms, Apamin+Nim: 846.8±235 ms, $t(48)$ = 0.9, p = 0.38; Fig. 5C)

remained comparable to responses recorded in apamin. Similarly, incubating slices in the P/Q

and N antagonist ω-conotoxin MVIIC (ω-Ctx , 1 µM) did not lead to significant changes in the

incidence of PF (Apamin: 92%, Apamin+ω-Ctx: 70%, n = 10, $\chi^2(1)$ = 1.4, p = 0.23; Fig. 5B), and

plateau potential duration (Apamin: 1108.5±126.6 ms, Apamin+ω-Ctx: 726.7±196.2 ms, $t(47)$ =

1.2, p = 0.24; Fig. 5C), suggesting that high-threshold Ca^{2+} channels do not play a critical role in

PF.

It is possible that plateau potentials are mediated by activation of a persistent Na^+

current (I_{NAP}) (Kim and McCormick, 1998). We found that in the presence of apamin and TTX

(500 nM), fast action potential activity was completely blocked, revealing a plateau potential in

9/16 cells (amplitude: 30±1.2 mV, n = 16, Fig. 5A), a smaller fraction compared to conditions in

apamin (Apamin: 92%, Apamin+TTX: 56%, $\chi^2(1)$ =10, p = 0.002; Fig. 5B). Plateau potential

duration remained comparable to control (Apamin: 1108.5±126.6 ms, Apamin+TTX:

912.8±325.7 ms, $t(49)$ = 0.7, p = 0.5; Fig. 5C). These data indicate that while action potential

activity mediated by transient Na^+ currents or I_{NAP} might facilitate plateau potential generation,

neither are strictly required.

Next, we examined if Na^+ current flow via conductances other than voltage-gated Na^+

channels are essential for plateau potentials, by substituting the majority of extracellular Na^+

with NMDG. We found that in the presence of NMDG (together with apamin and TTX) the

incidence of plateau potential generation was significantly reduced (apamin + TTX: 56%,

apamin + TTX + NMDG: 11%, n = 28, $\chi^2(1)$ = 10.6, p = 0.001; Fig. 5A-C), suggesting that PF is

mediated at least in part by a TTX-insensitive Na^+ current.

TRPM4 channels underlie PF in the TRN

Our data thus far are consistent with a role of the Ca^{2+}-activated nonselective cationic current (I_{CAN}) in mediating plateau potentials. Transient receptor potential (TRP) channels have been previously proposed as molecular mechanisms mediating I_{CAN} (Launay et al., 2002), with transient receptor potential canonical (TRPC) and melastatin (TRPM) channels as the most likely candidates. To examine the potential involvement of TRPC channels, we performed recordings in slices derived from global TRPC3 null mutant mice (TRPC3 KO, Hartmann et al., 2008) and from global quadruple TRPC1,4,5,6 null mutant mice (QKO, Tian & Zhu, 2018), in the presence of apamin. For both mouse lines, the incidence of PF (Apamin: 92%, TRPC3 KO: 70%, n = 10, $\chi^2(1)$ = 3.4, p = 0.65; QKO: 70%, n = 10, $\chi^2(1)$ = 3.4, p = 0.65; Fig. 6B) and plateau potential duration (Apamin: 1108.5±126.6 ms, TRPC3 KO: 841.1±136 ms, $t(47)$ = 0.9, p = 0.39; QKO: 752.6±133.4 ms, $t(47)$ = 1.1, p = 0.27; Fig. 6C) were comparable to our observations in WT animals, suggesting that at least 5 of the 7 TRPCs known to be expressed in the brain are not involved in mediating PF.

Next, we investigated a possible role of TRPM channels in mediating PF. TRPM2 has been shown to facilitate burst firing (Lee et al., 2013), while both TRPM4 and TRPM5 have been implicated in e.g. the generation of plateau potentials in cortex (Lei et al., 2014) and of slow inward currents in cerebellar Purkinje cells (Kim et al., 2013). We found that in the presence of the broad spectrum TRP channel inhibitor flufenamic acid (FFA, 100 µM), PF was completely eliminated (Apamin: 92%, Apamin+FFA: 0%, n = 17, $\chi^2(1)$ = 46.6, p = 10^{-20}; Fig. 6A-C). Next, we examined the effects of the broad-spectrum inhibitor 2-Aminoethoxydiphenyl borate (2-APB) which among TRP channels blocks TRPC7 and TRPM2, but does not inhibit TRPM4 or TRPM5. 2-APB had no effect on PF (Apamin: 92%, Apamin+ 2-APB: 90%, n = 10, $\chi^2(1)$ = 0.4, p = 0.54; Fig. 6B), with plateau potential duration comparable to control conditions (Apamin: 1108.5±126.6 ms, Apamin+2-APB: 758±132.4 ms, $t(49)$ = 1.3, p = 0.22; Fig. 6C), ruling out a role of TRPM2 or TRPC7 and suggesting that either TRPM4 or TRPM5 might mediate PF. Both

channels are voltage-dependent, monovalent-selective, and require increases in intracellular

Ca^{2+} for their activation (Launay et al., 2002; Nilius et al., 2003). We found that in the presence

of the TRPM4/5 antagonists 9-Phenthranol (9-Phen, 100 µM) and glibenclamide (Glib, 100 µM)

PF was completely eliminated (Apamin: 92%, Apamin+9-Phen: 15%, n = 15, $\chi^2(1) = 34.2$, $p =$

10^{-14}; Glib: 0%, n = 21, $\chi^2(1) = 51.4$, $p = 10^{-26}$; Fig. 6B). As both antagonists are known to have

non-specific effects, we performed additional experiments using the recently developed

selective TRPM4 antagonist 4-Chloro-2-[[2-(2-chlorophenoxy) acetyl]amino]benzoic acid (CBA,

Ozhatil et al., 2018). In the presence of CBA (50 µM), cells never displayed PF (Apamin: 92%,

CBA: 0%, n = 15, $\chi^2(1) = 44$, $p = 10^{-17}$; Fig. 6A,B). Taken together, our results indicate that

TRPM4 conductances mediate the plateau potential underlying PF.

Since increases in intracellular Ca^{2+} are required for TRPM4 activation, chelation of Ca^{2+}

using exogenous buffers such as BAPTA should eliminate PF. However, due to the rundown of

PF, such experiments were not feasible. We therefore probed a possible Ca^{2+} requirement of

PF by replacing extracellular Ca^{2+} with equimolar Ba^{2+}. Ba^{2+} passes through T-type Ca^{2+}

channels and generates large inward currents (Huguenard & Prince, 1992), but unlike Ca^{2+}

does not lead to TRPM4 activation (Nilius et al., 2004; Yamaguchi et al., 2014). Under these

conditions, brief depolarizing steps still led to prominent burst firing, indicating that T-type Ca^{2+}

channel dependent dendritic depolarizations were not affected (Fig. 6A). However, none of the

cells displayed PF (Apamin: 92%, Apamin+Ba^{2+}: 0%, n = 15, $\chi^2(1) = 44$, $p = 10^{-17}$; Fig. 6B,C).

These data further confirm that T-type Ca^{2+} channels alone are not sufficient to generate

plateau potentials. Instead, our data suggest that T-type Ca^{2+} channel activation and the

resulting Ca^{2+} increases lead to the activation of TRPM4 conductances and the generation of

PF.

Synaptic recruitment of muscarinic acetylcholine receptors suppresses PF

Finally, we examined the modulation of PF by synaptic activity. TRN neurons are the target of

cholinergic afferents from the basal forebrain and brainstem (Hallanger and Wainer, 1988). Our

previous studies have shown that acetylcholine (ACh) release leads to a biphasic postsynaptic

response, with fast excitation mediated by nicotinic ACh receptors (nAChRs) and slow and long-

lasting inhibition mediated by M2 muscarinic receptors (mAChRs) coupled G protein-gated

inward rectifying K^+ (GIRK) channels (Sun et al., 2013). To examine if cholinergic inputs can

trigger PF in the TRN, we performed experiments using BAC-ChAT-ChR2 mice that selectively

express ChR2 in cholinergic neurons (Zhao et al., 2011) and optogenetically evoked ACh

release with light pulses (1 ms) centered over the recorded neuron. To avoid wash-out of PF,

TRN neurons were recorded in loose-patch mode. Under these conditions, 2/5 of neurons

displayed PF (Fig. 7A,C). We reasoned that the activation of mAChRs and the resulting GIRK

activation might limit the expression of PF. Indeed, following atropine application to block

mAChRs, action potential firing was significantly prolonged in all neurons recorded (Control:

213.5±94 ms, Atropine: 416.3±84.1 ms, $t(4) = 7.1$, $p = 0.002$; Fig. 7A-C), and the incidence of

PF increased from 40% to 80%. These data show that mAChRs can modulate the expression of

PF.

Discussion

Here we have identified TRPM4 as the mechanism underlying PF in adult neurons of the TRN.

Furthermore, our data indicate that the large depolarizations and intracellular Ca^{2+} increases

required for TRPM4 activation are generated by Ca^{2+} influx through T-type Ca^{2+} channels.

Previous studies have focused on cellular pacemakers as being primarily responsible for slow

thalamic oscillations. Instead, our findings highlight that intrathalamic networks are equally

capable of generating slow rhythmic activity.

Mechanisms of PF in the TRN

Our results extend recent work that has implicated TRPM4 in controlling neuronal excitability in

cortex (Lei et al., 2014; Riquelme et al., 2018), hippocampus (Menigoz et al., 2016), substantia

nigra (Mrejeru et al., 2011), and brainstem (Picardo et al., 2019). A unique feature of TRPM4 is

its dependence on both depolarization and large increases in intracellular Ca^{2+} concentration

(Launay et al., 2002; Nilius et al., 2003; Nilius et al., 2004). Since dendritic Ca^{2+} signals can

experience decay under whole-cell conditions, this might partly explain why we consistently

observed rundown of PF in the present study. Alternatively, recordings in whole-cell mode might

have led to the dialysis of critical signaling molecules. TRPM4 conductances rapidly desensitize

in excised patches (Zhang et al., 2005) and in whole-cell mode, but modulators such as PIP2

can restore TRPM4 activity. Additional investigations into the mechanisms of TRPM4 regulation

under physiological conditions will be important.

What are the Ca^{2+} sources of TRPM4 activation? We found that pharmacological block

of T-type Ca^{2+} channel using the specific antagonist TTA-P2 completely eliminated PF. In

addition, replacing Ca^{2+} with Ba^{2+} abolished PF. These findings suggest that the Ca^{2+} increases

required for TRPM4 activation are mediated by influx through T-type Ca^{2+} channels although we

cannot rule out a contribution of R-type Ca^{2+} channels (Zaman et al., 2011) or release from

internal stores (Neyer et al., 2016). $Ca_V3.3$ channels mediating I_T are primarily expressed in the

distal dendrites of TRN neurons and their activation during burst firing results in large dendritic

but more modest somatic Ca^{2+} signals (Crandall et al., 2010; Astori et al., 2011). This would

suggest a dendritic location of TRPM4. In this scenario, membrane depolarizations necessary

for TRPM4 activation would likely be mediated by synaptically-generated T-type dependent Ca^{2+}

spikes, which in contrast to fast Na^+ action potentials, propagate effectively into TRN dendrites

(Crandall et al., 2010; Connelly et al., 2015).

We found that in the presence of apamin, PF could be observed in most recordings,

suggesting that TRPM4 conductances are expressed in the large majority of TRN neurons.

Thus, SK channels control both the initiation and duration of PF, likely due to the close

functional association between T-type and SK2 channels (Cueni et al., 2008). Similarly, we

found that GIRK conductances activated by M2 muscarinic receptors controlled the duration of

PF, suggesting that *in vivo* PF is not all-or-none but under constant regulatory control. While PF

has been observed in anesthetized preparations (Fuentealba et al., 2005), the activity of a

number of modulatory systems converging on TRN might explain why PF has yet to be

described during natural sleep.

Plateau potentials and PF have been observed in neurons throughout the brain

(Zylberberg & Strowbridge, 2017), including in several types of thalamic neurons, though there

is no definitive understanding of the underlying mechanisms. In neurons of the anterodorsal

thalamus, brief hyperpolarizing steps can lead to persistent firing over several minutes via

undefined mechanisms (Kulkarni et al., 2011). In local interneurons of the lateral geniculate

nucleus, postsynaptic mGluR5s can trigger long-lasting plateau potentials and PF lasting

minutes, possibly via recruitment of TRP conductances (Pressler & Regehr, 2013). For both

TRN and thalamic relay neurons in a number of species, group I mGluR activation can trigger

rhythmic forms of PF, mediated by the closure of a K^+ leak conductance, allowing the

generation of a long-lasting inward current mediated by the non-inactivating portion of T-type

Ca^{2+} channel conductance (Hughes et al., 2002; Blethyn et al., 2006). Our present findings differ

from these results in four important aspects: First, PF did not require mGluR activation and the

resulting closure of K^+ leak conductances. Second, in the absence of mGluR activation, PF did

not occur in a cell-intrinsic manner but required fast synaptic inputs. Under our experimental

conditions, both cholinergic inputs and bursts of glutamatergic thalamic inputs triggered PF,

likely by generating global Ca^{2+} spikes (Connelly et al., 2015). Third, while T-type Ca^{2+} channels

were important for the initiation of PF, our results are inconsistent with a critical role of I_T in

generating long-lasting currents that lead to plateau potentials. We found that plateau potentials

reached levels of ~-45 mV, where T-type Ca^{2+} conductances at steady-state experience near

full inactivation. Furthermore, the lack of PF in the presence of Ba^{2+}, which should allow for the

generation of a I_T-mediated window current (Huguenard and Prince, 1992) suggests that T-type

Ca^{2+} currents alone are unlikely to mediate PF. Fourth, using the potent and selective

antagonist CBA (Ozhatil et al., 2018), we showed that I_{CAN} was essential for the generation of

plateau potentials and was mediated by TRPM4. It should be stated that the majority of TRP

antagonists employed in previous studies have a number of non-specific effects, so the

validation of CBA as a selective TRPM4 antagonist under our experimental conditions will

require independent confirmation in additional studies.

Mechanisms of slow thalamic oscillations

Accumulating evidence indicates that both cortex and thalamus can generate oscillations in

isolation, suggesting that rhythmic activity in the intact thalamocortical system results from the

complex interplay of multiple distinct oscillators (Crunelli and Hughes, 2010). While there has

been extensive research on the cortical mechanisms mediating slow rhythms, the nature of

thalamic pacemakers are less well understood (Neske, 2016). Previous work has shown that

both TRN and relay neurons are capable of generating cell-intrinsic rhythms in the 1 Hz range,

but only under conditions of postsynaptic mGluR activation (Hughes et al., 2002; Blethyn et al.,

2006). Under physiological conditions, this would require sustained corticothalamic activity,

indicating that both thalamic cell types act as conditional oscillators. Our findings derived from

adult thalamic slices highlight an alternative mechanism and suggest that networks of TRN and

VB cells can generate robust and long-lasting slow oscillatory activity, in the absence of

sustained afferent activity. We observed an inverse relationship between TRN firing duration

and oscillatory frequency. It is therefore tempting to suggest that changes in TRN activity as

seen during PF acts to reduce the frequency of thalamic oscillations, via recruitment of

$GABA_BRs$ (Kim et al., 1997) or extrasynaptic high-affinity $GABA_ARs$ (Herd et al., 2013), thereby

changing the latency to rebound burst generation (Schofield et al., 2009). Thus, both intrinsic

and network mechanisms might interact to generate thalamic rhythmicity.

Our network data suggest that robust thalamic rhythmicity can occur in the absence of

precise synchrony, highlighting that both processes can occur independently. While thalamic

oscillatory activity is commonly referred to as synchronous (Fogerson and Huguenard, 2016),

there have been few studies that have systemically examined the degree of precise thalamic

synchronization during distinct behavioral states. We have previously shown that synaptically

evoked thalamic oscillatory activity rapidly desynchronizes, due to a number of cellular and

network mechanisms, including variable latencies of rebound bursting in relay neurons, TRN-

mediated lateral inhibition, and powerful but sparse thalamoreticular connections (Pita-Almenar

et al., 2014). Our current data suggest that the heterogeneity in TRN firing might act as an

additional factor in preventing synchrony in isolated thalamic networks. As has been suggested

previously (Contreras and Steriade, 1996), this would argue that synchrony in thalamic networks

is primarily regulated by afferent inputs, particularly from cortex.

References:

Agmon A, Connors BW (1991) Thalamocortical responses of mouse somatosensory (barrel)

cortex in vitro. Neuroscience 41:365–379 Available at:

http://www.ncbi.nlm.nih.gov/pubmed/1870696.

Astori S, Wimmer RD, Prosser HM, Corti C, Corsi M, Liaudet N, Volterra A, Franken P, Adelman

JP, Lüthi A (2011) The Ca v3.3 calcium channel is the major sleep spindle pacemaker in

thalamus. Proc Natl Acad Sci U S A 108:13823–13828.

Beenhakker MP, Huguenard JR (2009) Neurons that Fire Together Also Conspire Together: Is

Normal Sleep Circuitry Hijacked to Generate Epilepsy? Neuron 62:612–632 Available at:

http://dx.doi.org/10.1016/j.neuron.2009.05.015.

Blethyn KL, Hughes SW, Tóth TI, Cope DW, Crunelli V (2006) Neuronal basis of the slow (<1

Hz) oscillation in neurons of the nucleus reticularis thalami in vitro. J Neurosci 26:2474–

2486.

Bryant AS, Li B, Beenhakker MP, Huguenard JR (2009) Maintenance of thalamic epileptiform

activity depends on the astrocytic glutamate-glutamine cycle. J Neurophysiol 102:2880–

2888.

Connelly WM, Crunelli V, Errington AC (2015) The global spike: Conserved dendritic properties

enable unique ca2+ spike generation in low-threshold spiking neurons. J Neurosci

35:15505–15522.

Contreras D, Steriade M (1996) Synchronization of low-frequency rhythms in corticothalamic

networks. Neuroscience 76:11–24.

Crabtree JW, Lodge D, Bashir ZI, Isaac JTR (2013) GABAA, NMDA and mGlu2 receptors

tonically regulate inhibition and excitation in the thalamic reticular nucleus. Eur J Neurosci

37:850–859.

Crandall SR, Govindaiah G, Cox CL (2010) Low-threshold Ca 2+ current amplifies distal

dendritic signaling in thalamic reticular neurons. J Neurosci 30:15419–15429.

Crunelli V, Hughes SW (2010) The slow (1 Hz) rhythm of non-REM sleep: A dialogue between

three cardinal oscillators. Nat Neurosci 13:9–17.

Crunelli V, Larincz ML, Connelly WM, David F, Hughes SW, Lambert RC, Leresche N, Errington

AC (2018) Dual function of thalamic low-vigilance state oscillations: Rhythm-regulation and

plasticity. Nat Rev Neurosci 19:107–118.

Cueni L, Canepari M, Luján R, Emmenegger Y, Watanabe M, Bond CT, Franken P, Adelman

JP, Lüthi A (2008) T-type Ca2+ channels, SK2 channels and SERCAs gate sleep-related

oscillations in thalamic dendrites. Nat Neurosci 11:683–692.

David F, Schmiedt JT, Taylor HL, Orban G, Di Giovanni G, Uebele VN, Renger JJ, Lambert RC,

Leresche N, Crunelli V (2013) Essential thalamic contribution to slow waves of natural

sleep. J Neurosci 33:19599–19610.

Diekelmann S, Born J (2010) The memory function of sleep. Nat Rev Neurosci 11:114–126

Available at: http://dx.doi.org/10.1038/nrn2762.

Fernandez LMJ, Vantomme G, Osorio-Forero A, Cardis R, Béard E, Lüthi A (2018) Thalamic

reticular control of local sleep in mouse sensory cortex. Elife 7:1–25.

Fogerson PM, Huguenard JR (2016) Tapping the Brakes: Cellular and Synaptic Mechanisms

that Regulate Thalamic Oscillations. Neuron 92:687–704 Available at:

http://dx.doi.org/10.1016/j.neuron.2016.10.024.

Fuentealba P, Timofeev I, Bazhenov M, Sejnowski TJ, Steriade M (2005) Membrane bistability

in thalamic reticular neurons during spindle oscillations. J Neurophysiol 93:294–304.

Halassa MM, Siegle JH, Ritt JT, Ting JT, Moore CI (2011) Selective optical drive of thalamic

reticular nucleus generates thalamic bursts & cortical spindles. Nat Neurosci 14:1118–

1120.

Hallanger AE, Wainer BH (1988) Ultrastructure of ChAT-immunoreactive synaptic terminals in

the thalamic reticular nucleus of the rat. J Comp Neurol.

Hartmann J, Dragicevic E, Adelsberger H, Henning HA, Sumser M, Abramowitz J, Blum R,

Dietrich A, Freichel M, Flockerzi V, Birnbaumer L, Konnerth A (2008) TRPC3 Channels Are

Required for Synaptic Transmission and Motor Coordination. Neuron 59:392–398.

Hefft S, Jonas P (2005) Asynchronous GABA release generates long-lasting inhibition at a

hippocampal interneuron-principal neuron synapse. Nat Neurosci 8:1319–1328.

Herd MB, Brown AR, Lambert JJ, Belelli D (2013) Extrasynaptic GABAA receptors couple

presynaptic activity to postsynaptic inhibition in the somatosensory thalamus. J Neurosci

33:14850–14868.

Hughes SW, Cope DW, Blethyn KL, Crunelli V (2002) Cellular Mechanisms of the Slow (<1Hz)

Oscillations in Thalamocortical Neurons in Vitro. Neuron 33:947–958 Available at:

http://www.sciencedirect.com/science/article/pii/S0896627302006232.

Huguenard JR, McCormick DA (2007) Thalamic synchrony and dynamic regulation of global

forebrain oscillations. Trends Neurosci 30:350–356.

Huguenard JR, Prince DA (1992) A novel T-type current underlies prolonged Ca2+-dependent

burst firing in GABAergic neurons of rat thalamic reticular nucleus. J Neurosci 12:3804–

3817.

Huguenard JR, Prince DA (1994) Intrathalamic rhythmicity studied in vitro: Nominal T-current

modulation causes robust antioscillatory effects. J Neurosci 14:5485–5502.

Kim U, McCormick DA (1998) Functional and ionic properties of a slow afterhyperpolarization in

ferret perigeniculate neurons in vitro. J Neurophysiol 80:1222–1235.

Kim U, Sanchez-Vives M V., McCormick DA (1997) Functional dynamics of GABAergic

inhibition in the thalamus. Science (80-) 278:130–134.

Kim YS, Kang E, Makino Y, Park S, Shin JH, Song H, Launay P, Linden DJ (2013)

Characterizing the conductance underlying depolarization-induced slow current in

cerebellar Purkinje cells. J Neurophysiol 109:1174–1181.

Kulkarni M, Zhang K, Kirkwood A (2011) Single-cell persistent activity in anterodorsal thalamus.

Neurosci Lett 498:179–184 Available at: http://dx.doi.org/10.1016/j.neulet.2011.02.051.

Launay P, Fleig A, Perraud AL, Scharenberg AM, Penner R, Kinet JP (2002) TRPM4 is a Ca2+-

activated nonselective cation channel mediating cell membrane depolarization. Cell

109:397–407.

Lee CR, Machold RP, Witkovsky P, Rice ME (2013) TRPM2 channels are required for NMDA-

induced burst firing and contribute to H2O2-dependent modulation in substantia nigra pars

reticulata GABAergic neurons. J Neurosci 33:1157–1168.

Lei YT, Thuault SJ, Launay P, Margolskee RF, Kandel ER, Siegelbaum SA (2014) Differential

contribution of TRPM4 and TRPM5 nonselective cation channels to the slow

afterdepolarization in mouse prefrontal cortex neurons. Front Cell Neurosci 8:1–14.

Lemieux M, Chen JY, Lonjers P, Bazhenov M, Timofeev I (2014) The impact of cortical

deafferentation on the neocortical slow oscillation. J Neurosci 34:5689–5703.

Lüthi A (2014) Sleep spindles: Where they come from, what they do.

Menigoz A, Ahmed T, Sabanov V, Philippaert K, Pinto S, Kerselaers S, Segal A, Freichel M,

Voets T, Nilius B, Vennekens R, Balschun D (2016) TRPM4-dependent post-synaptic

depolarization is essential for the induction of NMDA receptor-dependent LTP in CA1

hippocampal neurons. Pflugers Arch Eur J Physiol 468:593–607.

Mrejeru A, Wei A, Ramirez JM (2011) Calcium-activated non-selective cation currents are

involved in generation of tonic and bursting activity in dopamine neurons of the substantia

nigra pars compacta. J Physiol 589:2497–2514.

Neske GT (2016) The slow oscillation in cortical and thalamic networks: Mechanisms and

functions. Front Neural Circuits 9:1–25.

Neyer C, Herr D, Kohmann D, Budde T, Pape HC, Coulon P (2016) MGluR-mediated calcium

signalling in the thalamic reticular nucleus. Cell Calcium 59:312–323 Available at:

http://dx.doi.org/10.1016/j.ceca.2016.03.009.

Nilius B, Prenen J, Droogmans G, Voets T, Vennekens R, Freichel M, Wissenbach U, Flockerzi

V (2003) Voltage dependence of the Ca2+-activated cation channel TRPM4. J Biol Chem

278:30813–30820.

Nilius B, Prenen J, Voets T, Droogmans G (2004) Intracellular nucleotides and polyamines

inhibit the Ca2+- activated cation channel TRPM4b. Pflugers Arch Eur J Physiol 448:70–

75.

Ozhatil LC, Delalande C, Bianchi B, Nemeth G, Kappel S, Thomet U, Ross-Kaschitza D,

Simonin C, Rubin M, Gertsch J, Lochner M, Peinelt C, Reymons J-L, Abriel H (2018)

Identification of potent and selective small molecule inhibitors of the cation channel

TRPM4. Br J Pharmacol 175:2504–2519.

Picardo MCD, Sugimura YK, Dorst KE, Kallurkar PS, Akins VT, Ma X, Teruyama R, Guinamard

R, Kam K, Saha MS, Del Negro CA (2019) Trpm4 ion channels in pre-bötzinger complex

interneurons are essential for breathing motor pattern but not rhythm. PLoS Biol 17:1–30.

Pinault D (2004) The thalamic reticular nucleus: Structure, function and concept.

Pita-Almenar JD, Yu D, Lu HC, Beierlein M (2014) Mechanisms Underlying Desynchronization

of Cholinergic-Evoked Thalamic Network Activity. J Neurosci 34:14463–14474.

Pressler RT, Regehr WG (2013) Metabotropic glutamate receptors drive global persistent

inhibition in the visual thalamus. J Neurosci 33:2494–2506.

Rigas P, Castro-Alamancos MA (2007) Thalamocortical up states: Differential effects of intrinsic

and extrinsic cortical inputs on persistent activity. J Neurosci 27:4261–4272.

Riquelme D, Silva I, Philp AM, Huidobro-Toro JP, Cerda O, Trimmer JS, Leiva-Salcedo E

(2018) Subcellular localization and activity of TRPM4 in medial prefrontal cortex layer 2/3.

Front Cell Neurosci 12:1–13.

Sachdeva R, Schlotterer A, Schumacher D, Matka C, Mathar I, Dietrich N, Medert R, Kriebs U,

Lin J, Nawroth P, Birnbaumer L, Fleming T, Hammes HP, Freichel M (2018) TRPC proteins

contribute to development of diabetic retinopathy and regulate glyoxalase 1 activity and

methylglyoxal accumulation. Mol Metab 9:156–167 Available at:

https://doi.org/10.1016/j.molmet.2018.01.003.

Sanchez-Vives M V., McCormick DA (2000) Cellular and network mechanisms of rhytmic

recurrent activity in neocortex. Nat Neurosci 3:1027–1034.

Schofield CM, Kleiman-Weiner M, Rudolph U, Huguenard JR (2009) A gain in GABA A receptor

synaptic strength in thalamus reduces oscillatory activity and absence seizures. Proc Natl

Acad Sci U S A 106:7630–7635.

Steriade M, Contreras D, Dossi RC, Nunez A (1993a) The slow (<1 Hz) oscillation in reticular

thalamic and thalamocortical neurons: Scenario of sleep rhythm generation in interacting

thalamic and neocortical networks. J Neurosci 13:3284–3299.

Steriade M, Nunez A, Amzica F (1993b) A novel slow (< 1 Hz) oscillation of neocortical neurons

in vivo: Depolarizing and hyperpolarizing components. J Neurosci 13:3252–3265.

Steriade M, Nunez A, Amzica F (1993c) Intracellular analysis of relations between the slow (<1

Hz) neocortical oscillation and other sleep rhythms of the electroencephalogram. J

Neurosci.

Steriade M, Timofeev I (2003) Neuronal plasticity in thalamocortical networks during sleep and

waking oscillations. Neuron 37:563–576.

Stroh A, Adelsberger H, Groh A, Rühlmann C, Fischer S, Schierloh A, Deisseroth K, Konnerth A

(2013) Making Waves: Initiation and Propagation of Corticothalamic Ca2+ Waves In Vivo.

Neuron 77:1136–1150.

Sun Y-G, Pita-Almenar JD, Wu C-S, Renger JJ, Uebele VN, Lu H-C, Beierlein M (2013)

Biphasic cholinergic synaptic transmission controls action potential activity in thalamic

reticular nucleus neurons. J Neurosci 33:2048–2059 Available at:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3711637&tool=pmcentrez&rend

ertype=abstract.

Sun YG, Rupprecht V, Zhou L, Dasgupta R, Seibt F, Beierlein M (2016) MGluR1 and mGluR5

synergistically control cholinergic synaptic transmission in the thalamic reticular nucleus. J

Neurosci 36:7886–7896.

Tian J, Zhu M (2018) GABAB Receptors Augment TRPC3-Mediated Slow Excitatory

Postsynaptic Current to Regulate Cerebellar Purkinje Neuron Response to Type-1

Metabotropic Glutamate Receptor Activation. Cells 7:90.

Ting JT, Daigle T, Chen Q, Feng G (2014) Acute brain slice method for adult and aging animals:

application of targeted patch clamp analysis and optogenetics. In: Methods in Molecular

Biology, Patch-Clam. (Martina M, Taverna S, eds), pp 221–242. New York: Humana Press.

Varga AW et al. (2016) Reduced Slow-Wave Sleep Is Associated with High Cerebrospinal Fluid

Aβ42 Levels in Cognitively Normal Elderly. Sleep 39:2041–2048.

Williams SR, Tóth TI, Turner JP, Hughes SW, Crunelli V (1997) The "window" component of the

low threshold Ca2+ current produces input signal amplification and bistability in cat and rat

thalamocortical neurones. J Physiol.

Yamaguchi S, Tanimoto A, Otsuguro KI, Hibino H, Ito S (2014) Negatively charged amino acids

near and in transient receptor potential (TRP) domain of TRPM4 channel are one

determinant of its Ca2+ sensitivity. J Biol Chem 289:35265–35282.

Zaman T, Lee K, Park C, Paydar A, Choi JH, Cheong E, Lee CJ, Shin HS (2011) CaV2.3

Channels Are Critical for Oscillatory Burst Discharges in the Reticular Thalamus and

Absence Epilepsy. Neuron 70:95–108 Available at:

http://dx.doi.org/10.1016/j.neuron.2011.02.042.

Zhang Z, Okawa H, Wang Y, Liman ER (2005) Phosphatidylinositol 4,5-Bisphosphate rescues

TRPM4 channels from desensitization. J Biol Chem 280:39185–39192.

Zhao S, Ting JT, Atallah HE, Qiu L, Tan J, Gloss B, Augustine GJ, Deisseroth K, Luo M,

Graybiel AM, Feng G (2011) Cell type-specific channelrhodopsin-2 transgenic mice for

optogenetic dissection of neural circuitry function. Nat Methods 8:745–755.

Zylberberg J, Strowbridge BW (2017) Mechanisms of persistent activity in cortical circuits:

Possible neural substrates for working memory. Annu Rev Physiol 40:60. – 627.

Legends:

Figure 1. Rhythmic activity in thalamic networks. A. Loose-patch recording of TRN neuron

showing rhythmic burst firing. **B.** Another TRN neuron showing rhythmic persistent firing (PF,

left), with a close-up (*right*) showing an individual bout of PF. **C.** Voltage-clamp recording of VB

neuron held at 0 mV displaying rhythmic bursts of IPSCs. **D.** Recording of another VB neuron

displaying rhythmic barrages of IPSCs (*left*). Close-up (*right*) shows a single barrage of IPSCs.

E. Summary data (n = 29) plotting the average duration of TRN firing as a function of oscillation

frequency. Red asterisks denote recordings shown in A and B. Dashed line marks criterion

separating between bursting and PF. **F.** Current clamp recording of TRN neuron, showing PF

during network activity. Close up (*right*) shows that bursts of EPSCs (highlighted by the red

arrows) trigger PF. **G.** Representative recording of rhythmic barrages of IPSCs in VB neuron,

under control conditions (*left*) and following bath application of NBQX (10 µM, *right*). Similar

data were obtained from 4 other VB neurons.

Figure 2. Axonal divergence and convergence characterize TRN outputs to VB. VB

recordings were performed in voltage clamp, with cells held at 0 mV. **A.** Divergent TRN output

to VB neurons. Dual recordings of neighboring VB cells showing a single barrage of IPSCs, with

a close-up (*middle*) highlighting precise synchrony of individual IPSCs activity in both cells.

Traces are normalized to the initial IPSC amplitude for both recordings. Schematic (*right*)

outlines proposed divergent TRN projections giving rise to VB responses. **B.** Convergent TRN

input to VB neuron**.** (*left*) Voltage-clamp recording of VB neuron with IPSC bursts (red tick

marks) and prolonged IPSC barrages (cyan tick marks) with distinct initial amplitudes. Close-up

(*middle*) shows overlay of individual events**.** Schematic (*right*) indicates proposed circuits giving

rise to VB response. **C.** Another example of TRN convergence onto VB neuron. (*left*) Recording

of VB neuron showing IPSC barrages with two distinct initial amplitudes (green and purple tick

marks, respectively). Close-up (*middle*) shows overlay of individual events**.** Schematic (*right*)

indicates proposed circuits giving rise to VB response.

Figure 3. PF is developmentally regulated and is controlled by SK conductances. A. Brief

depolarizing current steps (25 ms, 400 pA, RMP = -75 mV) lead to two distinct firing patterns,

Persistent Firing (PF, *top*) or Burst Firing (Bu, *bottom*). **B.** Summary data (n = 111) showing

bimodal distribution of current-step evoked membrane depolarizations. Grey line indicates

criterion separating PF and Bu. **C.** Raster plot of AP activity in representative TRN neuron

showing rundown of PF. **D.** Representative recordings of TRN neurons with PF at three

developmental stages, in control conditions (*black*) and in the presence of apamin (500 nM,

red). **E.** Summary data showing incidence of PF across development (P13-18: n = 24 in control,

n = 25 in apamin; P28-38: n = 23 in control, n = 22 in apamin; P90-120: 111 in control, n = 50 in

apamin). **F.** For the same cells as in D, summary data quantifying membrane potential duration

for all TRN cells across development, in control conditions and in apamin.

Figure 4. T-type Ca^{2+} channels but not mGluRs are required for PF. A. Representative

recordings of PF from two different TRN neurons, in control conditions and in the presence of

the mGluR1 antagonist JNJ 16259685 (1 µM) and the mGluR5 antagonist MTEP (10 µM). **B.**

Summary data quantifying duration of evoked membrane depolarization in control (n = 111) and

in the presence of mGluR 1/5 antagonists (n = 23). **C.** Representative recording of neuron with

PF, initially held at a holding potential of -60 mV and then at -75 mV. **D.** Summary data

quantifying evoked depolarization duration as a function of holding potential for TRN cells (n =

15). **E.** Representative recordings in the presence of apamin and in the presence of apamin and

the T-type Ca^{2+} channel antagonist TTA-P2. **F.** Summary data quantifying evoked depolarization

duration (n = 50 in apamin, n = 17 in apamin + TTA-P2).

Figure 5. A TTX-insensitive sodium current generates plateau potentials. All recordings

were performed in the presence of apamin. **A.** Representative TRN recordings in the presence

of the L-type Ca^{2+} channel antagonist nimodipine (20 µM, *top*), in TTX (500 nM, *middle*), and in

the presence of TTX and NMDG replacing extracellular Na^+ (*bottom*). **B.** Summary data

showing incidence of PF in the presence of nimodipine (n = 10), the P/Q- and N-type Ca^{2+}

channel antagonist ω-conotoxin MVIIC (1 µM, n = 10), TTX (n = 16), and TTX and NMDG (n =

28). Dashed line indicates incidence of PF in apamin. **C.** Summary data from the same cells as

in B quantifying evoked depolarization duration in each condition. Grey solid line represents

criterion for PF.

Figure 6. TRPM4 conductances mediate PF. All recordings were performed in the presence

of apamin. **A.** Representative recordings in control and in the presence of the non-selective

transient potential receptor (TRP) channel antagonist flufenamic acid (FFA, 100 µM, *top*), the

TRPM4 antagonist CBA (50 µM, *middle*), and in 2 mM $BaCl_2$ replacing extracellular $CaCl_2$

(*bottom*). **B.** Summary data showing incidence of PF in TRPC3 KO (KO, n = 10 cells) in

TRPC1,4,5,6 quadruple KO (QKO, n = 10 cells), FFA (n = 17), the non-selective TRP channel

antagonist 2-APB (50 µM, n = 10), the TRPM4/5 antagonists 9-phenthranol (100 µM, n = 15)

and glibenclamide (100 µM, n = 21), CBA (n = 15), and Ba^{2+}, (n = 15). Grey dashed line

represents incidence of PF in apamin. **C.** For the same cells as in B, summary data quantifying

the evoked depolarization duration.

Figure 7. Synaptic recruitment of muscarinic AChRs suppresses PF. Recordings were

carried out in slices of ChAT-ChR2-EYFP BAC mice. **A.** Loose-patch recording of a TRN neuron

showing brief barrage of ACh-evoked activity (*black*), which is prolonged following application of

the mAChR antagonist atropine (10 µM, *blue*). **B.** Raster plot of ACh-evoked neuronal activity

prior to and following atropine application, for the cell shown in A. **C.** Summary data quantifying

766   the average duration of ACh-evoked action potential activity before and after atropine

767   application (n = 5).

Figure 1. Rhythmic activity in thalamic networks. A. Loose-patch recording of TRN neuron
showing rhythmic burst firing. **B.** Another TRN neuron showing rhythmic persistent firing (PF,
left), with a close-up (*right*) showing an individual bout of PF. **C.** Voltage-clamp recording of VB
neuron held at 0 mV displaying rhythmic bursts of IPSCs. **D.** Recording of another VB neuron
displaying rhythmic barrages of IPSCs (*left*). Close-up (*right*) shows a single barrage of IPSCs.
E. Summary data (n = 29) plotting the average duration of TRN firing as a function of oscillation
frequency. Red asterisks denote recordings shown in A and B. Dashed line marks criterion
separating between bursting and PF. **F.** Current clamp recording of TRN neuron, showing PF
during network activity. Close up (*right*) shows that bursts of EPSCs (highlighted by the red
arrows) trigger PF. **G.** Representative recording of rhythmic barrages of IPSCs in VB neuron,

779 under control conditions (*left*) and following bath application of NBQX (10 µM, *right*). Similar
780 data were obtained from 4 other VB neurons.

Figure 2. Axonal divergence and convergence characterize TRN outputs to VB. VB
recordings were performed in voltage clamp, with cells held at 0 mV. **A.** Divergent TRN output
to VB neurons. Dual recordings of neighboring VB cells showing a single barrage of IPSCs, with
a close-up (*middle*) highlighting precise synchrony of individual IPSCs activity in both cells.
Traces are normalized to the initial IPSC amplitude for both recordings. Schematic (*right*)
outlines proposed divergent TRN projections giving rise to VB responses. **B.** Convergent TRN
input to VB neuron**.** (*left*) Voltage-clamp recording of VB neuron with IPSC bursts (red tick
marks) and prolonged IPSC barrages (cyan tick marks) with distinct initial amplitudes. Close-up
(*middle*) shows overlay of individual events**.** Schematic (*right*) indicates proposed circuits giving
rise to VB response. **C.** Another example of TRN convergence onto VB neuron. (*left*) Recording
of VB neuron showing IPSC barrages with two distinct initial amplitudes (green and purple tick
marks, respectively). Close-up (*middle*) shows overlay of individual events**.** Schematic (*right*)
indicates proposed circuits giving rise to VB response.

Figure 3. PF is developmentally regulated and is controlled by SK conductances. A. Brief depolarizing current steps (25 ms, 400 pA, RMP = -75 mV) lead to two distinct firing patterns, Persistent Firing (PF, *top*) or Burst Firing (Bu, *bottom*). **B.** Summary data (n = 111) showing bimodal distribution of current-step evoked membrane depolarizations. Grey line indicates criterion separating PF and Bu. **C.** Raster plot of AP activity in representative TRN neuron showing rundown of PF. **D.** Representative recordings of TRN neurons with PF at three developmental stages, in control conditions (*black*) and in the presence of apamin (500 nM, *red*). **E.** Summary data showing incidence of PF across development (P13-18: n = 24 in control, n = 25 in apamin; P28-38: n = 23 in control, n = 22 in apamin; P90-120: 111 in control, n = 50 in apamin). **F.** For the same cells as in D, summary data quantifying membrane potential duration for all TRN cells across development, in control conditions and in apamin.

Figure 4. T-type Ca^{2+} channels but not mGluRs are required for PF. A. Representative recordings of PF from two different TRN neurons, in control conditions and in the presence of the mGluR1 antagonist JNJ 16259685 (1 µM) and the mGluR5 antagonist MTEP (10 µM). **B.** Summary data quantifying duration of evoked membrane depolarization in control (n = 111) and in the presence of mGluR 1/5 antagonists (n = 23). **C.** Representative recording of neuron with PF, initially held at a holding potential of -60 mV and then at -75 mV. **D.** Summary data quantifying evoked depolarization duration as a function of holding potential for TRN cells (n = 15). **E.** Representative recordings in the presence of apamin and in the presence of apamin and the T-type Ca^{2+} channel antagonist TTA-P2. **F.** Summary data quantifying evoked depolarization duration (n = 50 in apamin, n = 17 in apamin + TTA-P2).

818
819
Figure 5. A TTX-insensitive sodium current generates plateau potentials. All recordings
were performed in the presence of apamin. **A.** Representative TRN recordings in the presence
of the L-type Ca^{2+} channel antagonist nimodipine (20 µM, *top*), in TTX (500 nM, *middle*), and in
the presence of TTX and NMDG replacing extracellular Na^+ (*bottom*). **B.** Summary data
showing incidence of PF in the presence of nimodipine (n = 10), the P/Q- and N-type Ca^{2+}
channel antagonist ω-conotoxin MVIIC (1 µM, n = 10), TTX (n = 16), and TTX and NMDG (n =
28). Dashed line indicates incidence of PF in apamin. **C.** Summary data from the same cells as
in B quantifying evoked depolarization duration in each condition. Grey solid line represents
criterion for PF.

Figure 6. TRPM4 conductances mediate PF. All recordings were performed in the presence of apamin. **A.** Representative recordings in control and in the presence of the non-selective transient potential receptor (TRP) channel antagonist flufenamic acid (FFA, 100 µM, *top*), the TRPM4 antagonist CBA (50 µM, *middle*), and in 2 mM BaCl$_2$ replacing extracellular CaCl$_2$ (*bottom*). **B.** Summary data showing incidence of PF in TRPC3 KO (KO, n = 10 cells) in TRPC1,4,5,6 quadruple KO (QKO, n = 10 cells), FFA (n = 17), the non-selective TRP channel antagonist 2-APB (50 µM, n = 10), the TRPM4/5 antagonists 9-phenthranol (100 µM, n = 15) and glibenclamide (100 µM, n = 21), CBA (n = 15), and Ba^{2+}, (n = 15). Grey dashed line represents incidence of PF in apamin. **C.** For the same cells as in B, summary data quantifying the evoked depolarization duration.

Figure 7. Synaptic recruitment of muscarinic AChRs suppresses PF. Recordings were
carried out in slices of ChAT-ChR2-EYFP BAC mice. **A.** Loose-patch recording of a TRN neuron
showing brief barrage of ACh-evoked activity (*black*), which is prolonged following application of
the mAChR antagonist atropine (10 µM, *blue*). **B.** Raster plot of ACh-evoked neuronal activity
prior to and following atropine application, for the cell shown in A. **C.** Summary data quantifying
the average duration of ACh-evoked action potential activity before and after atropine
application (n = 5).