1 **Characterizing the tumor microenvironment of metastatic ovarian**
2 **cancer by single cell transcriptomics.**
3

4 Susan Olalekan[1*], Bingqing Xie[1], Rebecca Back[1], Heather Eckart[1], Anindita Basu[1,2*]
5
6 1 Department of Medicine, University of Chicago
7 2 Center for Nanoscale Materials, Argonne National Laboratory
8
*Correspondence:
Anindita Basu, PhD
5841 S. Maryland Avenue. N417B
Chicago, IL 60637
Telephone: 773-834-1512
Fax: 773-702-2567
Email: (onibasu@uchicago.edu)

16
17 Running title: scRNA-seq of ovarian cancer tumor microenvironment.
18
19 Keywords: Ovarian cancer, scRNA-seq, transcriptomics, immune cells, human cancer
20
21 Conflict of interest: The authors declare no potential conflict of interest.
22

23 Funding: This project is internally funded.

Abstract

Epithelial ovarian cancer is a highly heterogenous, metastatic and lethal disease. The presence of CD8$^+$T cells within ovarian tumors is positively associated with overall patient survival. Determining if a patient has T cells that respond to immunotherapies, their characteristics and how they can be manipulated to target cancer cells is an area of intense investigation in cancer therapy. This study determines the cellular composition and the transcriptional state of immune cells in metastatic ovarian cancer samples from patients using single cell RNA sequencing (scRNA-seq). Hierarchical clustering stratified our patient cohort into 2 main groups: 1) a high T cell infiltration (high T_{inf}) group and 2) a low T cell infiltration (low T_{inf}) group. To assess the immune response in these patient samples, we performed an unsupervised clustering of the T cell population in each group. The T cell population clustered into 4 and 3 subpopulations in the high T_{inf} T cell and low T_{inf} respectively. A granulysin expressing T cell cluster was identified and unique to the High T_{inf} group. Interestingly, although both groups had resident memory CD8$^+$T (CD8$^+$Trm) cells, only the CD8$^+$Trm cells in the high T_{inf} group expressed TOX, a recently described transcription factor. TOX confers longevity to T cells within immunosuppressive environment such as cancer. Interestingly, along with TOX$^+$ T cells, we found a unique plasmablast cluster and an IRF8$^+$ macrophage cluster unique to the high T_{inf} group. Our comprehensive scRNA-seq study provides important insights in elucidating the immune response in ovarian cancer patients.

Introduction

Ovarian cancer is the most lethal malignancy of the female reproductive tract[1]. Conventional therapy involving cytoreductive surgery and chemotherapy are 90% effective when cancer is diagnosed at the early stage when it is still restricted to one or both ovaries. Unfortunately, most ovarian cancer cases are diagnosed at stage III or IV when the cancer has metastasized and the diagnosis in these patients result in a 30% 5-year survival rate[2]. To develop efficacious therapies for metastatic ovarian cancer, we need to define the cellular heterogeneity and the transcriptional state within the tumor microenvironment. Immunohistochemical staining and flow cytometry have been useful in categorizing the cell types based on specific cell surface markers but mask intra-cellular heterogeneity. Bulk RNA profiling has been used to categorize high grade serous ovarian carcinoma (HGSOC), the most common and lethal histotype of ovarian cancer, into molecular subtypes[3,4]. However, bulk RNA sequencing averages gene expression and fails to identify the respective contribution of cell subsets. Alternatively, single cell RNA sequencing (scRNA-seq) has emerged as a powerful tool to interrogate tumor composition, revealing previously uncharacterized cellular heterogeneity and gene regulatory networks at single cell resolution[5-10]. Recently, a scRNA-seq study, investigated the heterogeneity in the proposed cell of origin of HGSOC and revealed a high epithelial-mesenchymal transition (EMT) high subtype associated with poor prognosis[11]. These transcriptomic studies have focused on ovarian cancer cells; however, the tumor microenvironment is made up of other cell types that can affect patient stratification, targeted treatment and outcome.

The omentum is mainly an adipose tissue in the peritoneal cavity containing aggregates of immune cells in areas called milky spots[12]. These milky spots act similar to lymph nodes, collecting and responding to antigen within the peritoneal cavity. Interestingly, ovarian cancer cells preferentially colonize adipose tissue with milky spots in the peritoneal cavity[13]. Additionally, adipocytes provide adipokines and act as a source of energy for ovarian cancer cells[14]. These factors prime the omentum as a premetastatic niche for ovarian cancer. The initial presence of ovarian cancer cells in the omentum leads to a recruitment of macrophages into the milky spots without anti-tumor effects. Contrastingly, the presence of tumor infiltrating CD8+ T cells in both the ovary and omentum is associated with significantly longer overall survival[15]. However, checkpoint inhibitors, a cancer immunotherapeutic approach, aimed at restoring CD8+ T cell function have had a low response rate[15]. We therefore need a better understanding of the tumor microenvironment to improve patient response rate to cancer immunotherapy.

In this current study, we use Drop-seq technology to examine the cells within omental tumors from 6 patients with different histotypes of ovarian cancer. We identified 12 cell clusters and patient samples were stratified into two groups based on immune signatures: 1) a high T cell infiltration (high T_{inf}) group and 2) a low T cell infiltration (low T_{inf}) group. This unprecedented concurrent transcriptomic analysis of all cells within metastatic ovarian cancer unravels patient-specific responses to cancer and identifies patients that might be suitable for immune checkpoint inhibition.

Materials and methods

Tissue collection

Ovarian cancer tissue was collected from women undergoing debulking surgery at the University of Chicago. Human tissue acquisition after patient deidentification was approved by the university of Chicago institutional review board for human research. Ovarian cancer tissue was histologically carcinoma classified and staged by a pathologist according to tumor-node-metastasis (TMN) and/or international federation of gynecology and obstetrics (FIGO) classifications (Table 1). Three of the ovarian cancer samples originated from the left fallopian tube with associated serous tubal intraepithelial carcinoma (STIC).

Immunohistochemistry

Ovarian cancer tissues obtained from the anatomic pathology department were fixed overnight in 4% formaldehyde at 4^0C. After serial dehydration, the tissues were embedded in paraffin and cut into 5 μm thick sections. Histological evaluation was done with hematoxylin and eosin (H&E). Immunohistochemical staining was performed to confirm the presence of cytokeratin-7 (Thermofisher), pan-vimentin (DAKO), CD45 (Agilent), CD3 (Human Envision) and CD19 (Abcam) cells. Briefly, sections were deparaffinized and rehydrated using xylene and serial dilutions of EtOH in distilled water. Tissue sections were incubated in citrate buffer, pH 6 and heated in a microwave oven. Anti-cytokeratin-7 (1:1000), anti-vimentin (1:100), anti-CD45 (1:100), anti-CD3 (1:100), and anti-CD19 (1:200) antibodies were applied on tissue sections with one-hour incubation at room temperature in a humidity chamber. The antigen-antibody bindings were detected with labeled polymer-HRP Envision system (DAKO, K4007) and DAB+ chromogen (DAKO, K3468) system. Tissue sections were briefly immersed in hematoxylin for counterstaining and were covered with cover glasses.

Tissue digestion, red blood cell lysis and dead cell removal

Ovarian cancer tissue was transported in DMEM/F12 containing 10% FBS and 1% P/S (10% DMEMF/12) on ice from the anatomic pathology department to the laboratory. The tissue was minced manually with a scalpel and enzymatically digested using 1.5 mg/ml collagenase IV (Sigma-Aldrich), 1 mg/ml hyaluronidase (Sigma-Aldrich) and 500 μg/ml DNase I (GoldBio) in Hank's balanced salt solution (HBSS) in a 37 ^{0}C shaker (200 rpm) for 0.5 – 2 h. Following digestion, cells were resuspended in 10% DMEMF/12 and filtered serially through 70 μm and 40 μm strainers. Red blood cells were lysed by incubating the cell suspension in RBC lysis buffer (Sigma-Aldrich) for 2-5 minutes. Lysis was quenched by adding excess 10% DMEMF/12. The number of live cells was enriched using the dead cell removal kit (Miltenyi, 130-090-101).

Drop-seq experiments

Drop-seq experiment was performed as previously described[16]. Briefly, cells and barcoded beads
were loaded at a concentration of 100,000 cells/ml in PBS-BSA and 120,000 beads/ml in Drop-
seq lysis buffer in 3 ml syringes. Droplets were generated using a 125-micron microfluidic device
at 16 ml/hr (oil), 4 ml/hr (cells) and 4 ml/hr (beads) with ~15 minutes per collection. Following
collection, drops were broken and barcoded beads with mRNA hybridized onto them were
collected and washed. Barcoded cDNA attached to the beads or STAMPS were generated by
reverse transcription, treated with exonuclease 1 and the number of STAMPs was counted. 5000
STAMPSs were aliquoted per well in a 96-well plate and the cDNA attached to the STAMPS were
amplified through 14 PCR cycles. Supernatants from each well were pooled and cleaned with
Ampure beads. Purified cDNA was quantified using Qubit 3.0 (Invitrogen) and 450-650 pg of each
sample was used as input for Nextera reactions (12 cycles). Tagmented libraries were quantified
using Agilent BioAnalyzer High sensitivity chip before submission for sequencing on Illumina's
NextSeq 500, using 75 cycle v3 kits. Paired end sequencing was performed with 20 bp for Read 1
and 64 bp for Read2 using Read1 primer, GCCTGTCCGCGGAAGCAGTGGTATCAACGCAGAGTAC
and 5% Illumina PhiX Control v3.
187
188 **Data processing, alignment and clustering analysis**
189
From alignment of the reads from the Dropseq experiments, read count matrices for unique
molecular identifiers (UMIs) were generated for both exon and intron regions in the human
genome (gencode hg38 v.27) using the STAR version 2.5.3 aligner[17,18]. The ovarian cancer samples
from six patients' metastatic omentum were sequenced with Drop-seq. A total of 13 sequencing
runs were performed for six Drop-seq samples where each sample was sequenced at least twice
to maximize the sequencing depth (PT-3401 was sequenced three times). Each run produced
paired-end reads, with Read 1 representing the 12 bp cell barcode and six bp unique molecular
identifier (UMI), and Read 2 representing a 60 bp mRNA fragment. Paired-end reads from the
same samples were merged to generate six paired-end fastq files. A *snakemake* pipeline was
applied to each sequencing run and produce a count matrix representing the gene expression in
every cell[19]. Individual count matrices were produced for each of the six patients after accounting
for UMI duplicates.
202
The individual count matrix was produced for each patient with a total number of six count
matrices. The summarized counts for each gene were inferred based on both exon and intron
reads to produce the gene expression matrix per sample. Expression matrices of six ovarian
cancer samples from the metastatic sites in the omentum were generated. To select high quality
cells, we applied a filter based on the number of genes detected per cell. Prior to filtering, each
sample produced approximately 5000 cells. Based on the median number of genes captured, cells
with less than 800 genes were removed from the data sets. For lower quality samples, the
threshold was lowered to 600 or 700 as shown in Table 2. We followed a standardized pipeline
using single cell analysis tool suite, Seurat v3.0.2[20,21]. A global-scaling logarithmic normalization
method was applied to all samples[21] that normalizes the feature expression counts for each cell
by the total expression counts, multiplies by a scale factor of 10,000 (TP10K), and natural-log
transforms, as ln(TP10K+1). Each normalization matrix was then scaled by a linear transformation
to center the mean gene expression of cells and normalized the gene expression variance. The

most variable genes were extracted for principal component analysis (PCA). We applied PCA on
the normalized expression matrix with the most variable genes to extract the top 50 components
in the data, followed by a heuristic elbow plot on the variance explained of each PC. We selected
the number of top variant PCs based on the elbow plot which varies from 10 to 20 depending on
the sample. The top PCs were used in further exploration of the data, such as UMAP /tSNE
dimension reduction, construction of K-nearest neighbor graph, etc. For analysis that included
multiple samples, integration through anchoring[22] was applied. Note that a subset of genes,
usually highly variable ones, was selected to perform the integration, where the integrated gene
expression matrix has a lesser number of features (genes) than the original gene expression
matrix. The samples from multiple patients were integrated before classifying cell types.
Differential expression analysis was done through *FindMarkers* function in Seurat V3 using the
Wilcoxon Rank Sum test, and statistically significant markers were extracted for sub-populations
or contrast groups based on an adjusted p-value threshold of 0.05. We used dimension reduction
methods, UMAP[23] and tSNE[24] to generate 2D plots to visualize different cell populations in the
experiment.
231
Cancer subtype classification and correlation with The Cancer Genome Atlas

234 Four cancer subtypes- differentiated, immunoreactive, mesenchymal, and proliferative were
235 determined from bulk sequencing study on ovarian cancer[4]. Modular scores[25] were generated
236 between gene expression levels for each cell with upregulated marker signatures on the four
237 subtypes. The subtypes were then assigned to individual cells by the highest positive modular
238 score and if no positive modular scores were found, the subtype was considered undecided.

Cell type classification and correlation with CellAtlas

To assign cell types to individual cells, we used a bulk RNA sequencing data-set from 95 cell lines
collected by CellAtlas[26], covering 33 major cell types in normal human tissue, including common
immune, endothelial, epithelial, fibroblast, mesodermal cells and so on. The cell lines can be
further divided into 60 subtypes. We inferred the similarity between individual cells in the
samples and cell lines by calculating the pairwise Pearson correlation matrix $C = \{cor(i,j)\}$
among any cell i from the Drop-seq experiments with any cell line, j in the CellAtlas. A shared
nearest neighbor (SNN) modularity optimization-based clustering[27] on the CellAtlas cell lines
grouped the cell-lines into larger and more general cell-type categories. This was done to
eliminate any bias in CellAtlas dataset and remove rare or under-powered cell-types. The
resulting clusters of cell-lines were annotated by the most frequent major cell-type from the cell-
line groups. For each cell in our cancer sample, cell-type was then assigned from top 5 highly
correlated major cell-type clusters. We collapsed ambiguous cell types with neighboring cell
types, based on expression profiles.

Classification with cluster markers, canonical genes, and genetic functions

259 Sometimes the Gene Ontology (GO) and Pathway analysis gives general/noisy functional
260 categories which make certain cell types hard to identify. By leveraging prior knowledge from the
261 CellAtlas mapping, we were able to locate the relevant function categories for those cell types
262 and narrow down their marker genes efficiently[26]. Moreover, by the consensus of both GO and
263 CellAtlas analyses, we obtained higher confidence in classifying cell types in each patient sample.
264
We used the cell types obtained from CellAtlas correlation as baseline and curated cell-types
manually using canonical genes and functional association. Differentially expressed marker genes
were extracted from cell clusters. Cross-referencing was done between gene markers and known
canonical genes to the cell-types. Gene ontology and pathway enrichment was performed on
gene markers to provide additional evidence for cell-type assignment. We identified 9 major cell
types including epithelial cells, fibroblast cells, three immune cell types; B cells/B cell plasma, T
cells, and monocytes, mesenchymal stem cells (MSC), embryonic stem cells (ESC), and
endothelial cells.
273
274
275 # Results
276
277 **Characteristics of the tumor microenvironment**
278
Ovarian cancer samples (1-3 different sections) were collected from the omental metastatic site
of six patients (Table 1). Four patients were diagnosed with advanced high-grade serous cancer,
one with serous carcinoma and one with malignant mixed Mullerian tumor (MMMT). Ages of the
patients ranged from 46 to 71 years. Four of the patients received neo-adjuvant therapy prior to
surgery, including the patient diagnosed with MMMT. Normal omentum is comprised mainly of
adipocytes as well as aggregates of leukocytes called milky spots[12]. As metastasis progresses,
there is an inverse relationship between ovarian tumor growth (cancer and cancer associated cell
growth) and adipocytes within the omentum[14]. Therefore, the area of tumor occupied by
adipocyte decreases with disease score[28]. To quantify the transformation of the omentum by
ovarian cancer cells, we used Imagescope, a digital histopathology software, to annotate and
calculate the area occupied by adipocytes in each patient sample on hematoxylin and eosin (H&E)
stained sections (Figure 1A). The area of adipocytes was reported as a percentage of the total
surface area of the sample and used as a measure of disease score (Figure 1B), similar to Pearce
et al.[28] We rated the patient samples from lowest to highest disease score; PT-5150, PT-3401, PT-
6885, PT-3232, PT-4806, PT-2834 respectively. The percentage of cancer cells, stromal cells and
immune cells were quantified by immunohistochemistry (IHC) using antibodies against
cytokeratin-7 (CK-7), vimentin and CD45 antibodies (Figure 1C-H). There was a varying proportion
of each cell type across patients. Notably, we observed that malignant cells in some patients (PT-
5150, PT- 3232, PT-4806 and PT-2834) double stained for CK-7 and vimentin, suggesting active
epithelial-mesenchymal-transition (Figure 1C-F). In addition, the MMMT (PT-2834) sample had
two different types of malignant cells that stained separately for CK-7 and vimentin. We assessed
the relationship between the area of adipocytes and the cancer and stromal compartments and
only observed a positive correlation with CD45+ immune cells (Supplementary Figure 1). Notably,

we observed aggregation of immune cells closer to adipocytes and sparsely otherwise
throughout the rest of the tissue. Together, these data reveal patient variability in the cellular
transformation of tumors from the same pathological stage.
305
306
307 **Generation of single cell data, cell clustering and cell type assignment**
308
For Drop-seq experiments, tumors samples were enzymatically digested into a single cell
suspension and processed[16]. We integrated all six omental samples into a gene expression matrix
which contains the expression values of 9885 cells and 40947 genetic features such as protein
coding genes, pseudogenes, lncRNA, mapped from the GENCODE human reference genome
(version GRCh38). A minimum cut off of 600-800 genes per cell was set, as shown in Table 2.
Hierarchical clustering was performed using resolution of 0.2 with 12 clusters detected
(Supplementary Figure 2A). Each cluster had cells from all samples (Supplementary Fig. 2B). The
cell types were assigned and curated by cell line correlation, canonical genes and functional
categories for the significantly differentially expressed genes on the detected clusters. 9 major
clusters out of 12 were identified with cell types (Figure 2A). This included 1 cluster for malignant
epithelial cells, 1 cluster for fibroblasts, 1 cluster for mesenchymal stem cells, 1 cluster for
embryonic stem cells, 1 cluster for endothelial cells. Each patient sample had a varying proportion
of the 9 identified cell types/clusters (Figure 2B). Previous bulk mRNA and microRNA expression
studies have established 4 transcriptional subtypes of HGSOC[3,29]. To assign molecular subtypes,
we extracted gene sets defining each molecular subtype and applied them to our Drop-seq data.
At the single cell level, each molecular subtype is represented in every patient tumor sample
(Figure 2C). We further analyzed which cell type/cluster belonged to each molecular subtype.
The differentiated and proliferative subtypes were mainly made up of epithelial cells,
mesenchymal and immunoreactive subtypes were mainly made up of fibroblasts and immune
cells respectively suggesting that TCGA classification is reflective of the proportion of cells types
within each tumor (Figure 2D).
330
It has been suggested that the TCGA subtypes may be useful in specifically categorizing cancer
cells within a tumor. However, most of the cancer cells belonged to the differentiated,
proliferative and undecided subtype. Therefore, to determine patient-derived differences in
cancer cells that might be useful in personalizing therapy, we generated a heatmap from
differentially expressed genes from each patient. The expression profile of all patients appeared
similar in all patients except PT-2834 (Figure 2E). This patient was histologically diagnosed with
MMMT with chondrosarcomatous elements, also confirmed by our Drop-seq data
(Supplementary Figure 3). Next, the central cluster (epithelial cells) and connecting ESCs were
selected and assessed for known ovarian cancer cell markers with 7 clusters detected (Figure 2F).
PAX8 and MUC16, indicative of advanced disease, was expressed by multiple clusters[30,31]. Cluster
1 expressed vimentin along with high Ki-67 and PCLAF indicative of a proliferating population.
Cluster 5 also expressed PCLAF, a marker of proliferation, as well as FN1 and ZEB1, markers of
EMT. Cluster 6 highly expressed vimentin and Ki-67 along with comparatively lower expression
of EMT markers (ZEB1 and ZEB2), suggestive of cells undergoing mesenchymal to epithelial
transition (Figure 3E). Ovarian cancer stem cells marked by co-expression of CD33, CD44, CD117

and CD24 expression were not observed[32]. Cells in this cluster expressed IL-6 which correlates
with poor prognosis in ovarian cancer and TGF-β which induces an aggressive cancer
phenotype[33-36]. These results reveal a snapshot of the heterogeneous processes that cancer cells
harness to progress disease.
350
351 **Immune cellular profile of patient samples**
352
To investigate the immune population in our patient samples, we selected the immune cells from
the six patient samples and they clustered into 4 main populations; T cells, B cells, plasma B cells
and monocytes/macrophages (Figure 3A). Similar samples were grouped by a dendrogram with
a major difference in T cell infiltration (T_{inf}) (Figure 3B). Interestingly, the high T_{inf} group (PT-5150
and PT-3401) had the lowest disease score while the low T_{inf} group (PT-4806, PT-6885, PT-2834,
PT-3233) had higher disease score. During the initial phase of metastasis, macrophages are
recruited from the peritoneal cavity without any anti-tumor effect on the cancer cells[37,38]. To
assess the differences in the macrophage population, we performed unsupervised clustering on
both groups separately (Figure 3C&D). Both groups had a $CD14^+CD16^+CD163^+CD204^+$ cluster
reminiscent of tumor associated macrophages or TAMs (Figure 3E&F, Supplementary Figure 4).
In the low T_{inf} group, there was also an immature $FLT3^+$ population. These $FLT3^+$ progenitors can
differentiate into osteoclasts, dendritic cells, microglia and macrophages[39]. Other clusters in the
high T_{inf} group included a $CD274^+$ cluster, suggesting a regulatory population similar to MDSCs
and an $IRF8^+$ cluster induced in the presence of IFN-γ and promotes the formation of
autophagosome (Figure 3E&F, Supplementary Figure 4)[40]. The final cluster highly expressed
NR1H2 which inhibits inflammatory genes in macrophages[41,42]. This clustering analysis reveals
phenotypically distinct macrophages present in the tumor in addition to the established TAMs.
370
371 **Differences in T cell clustering and subtype analysis**
372
To reveal the functional subtypes of T cells and any difference in the low T_{inf} versus high T_{inf} group,
we clustered 136 and 820 T cells respectively (Figure 5A&B). A total of 3 and 4 transcriptionally
distinct clusters emerged from low T_{inf} and high T_{inf} T cells respectively. The top genes revealed
clusters similar to previously described T cell phenotypes in breast cancer, including $CD4^+IL7R^+$,
$CD4^+FOXP3^+$, resident memory $CD8^+T$ cells ($CD8^+Trm$ cells) and one population described in lung
cancer with high granulysin expression $CD4^+GNLY^+$ (Figure 5C-F, Supplementary Figure 5)[43,44].
Both groups had $CD8^+Trm$ cells which expressed IFN-γ, $CD4^+IL7R^+$ and $CD4^+FOXP3^+$ T cells. The
high T_{inf} group had an extra $CD4^+GNLY^+$ cluster. Contrastingly, GNLY was expressed in the low T_{inf}
group by $CD8^+Trm$ cluster (Figure 5C&D). TOX was highly expressed by the $CD8^+Trm$ and the
$CD4^+GNLY^+$ clusters in the high T_{inf} group as compared to the low T_{inf} group (Figure 5E&F). These
TOX^+ T cells persist during chronic infection and TOX is expressed on the $CD8^+T$ cells that are
reactivated in response to PD-L1 immunotherapy[45,46]. Layilin, a previously described marker of
exhaustion was mainly expressed by the $CD4^+FOXP3^+$ cluster (Figure 5E&F). These data suggest
that scRNA-seq is capable of determining patients that will benefit from novel therapy.
387
388 **Transcriptionally distinct plasmablasts cluster in high T_{inf} group**
389

The high T_{inf} group had an accompanying high B cell signature (Figure 3B). To investigate the B
cell subtypes in our samples, naïve B cells and plasma cells were clustered yielding 4 and 2
clusters in high T_{inf} and low T_{inf} group respectively (Figure 6A&B). In the high and low T_{inf} group,
the naïve cluster expressed MHC class II genes highly with a difference in CD24 expression (Figure
6C&D, Supplementary Figure 6). Other clusters in both groups expressed BLIMP1 indicating they
are plasma cell clusters. In the high T_{inf} group, the BLIMP1$^+$ plasma cells (PC) segregated into 3
clusters. The CD38$^+$PC and CD138$^+$PC are indicative of post germinal center B cells. The final
cluster highly expressed MHC class II genes as well as proliferative genes suggesting a plasmablast
population.
399

Discussion

In this study, we characterized samples from 6 ovarian cancer patients. One to three sections of
tumors were processed to account for intratumor heterogeneity based on availability. Initial IHC
staining of these sections revealed a positive correlation between the area of tumor made up of
adipocytes and the number of immune cells within the tumor. We processed these tumor
sections and obtained Drop-seq data for a total of 9885 cells after quality control. Mapping our
cells to curated cell types from the CellAtlas along with gene ontology and pathway-based
enrichment was robust in assigning cells types to almost all clusters in our analysis. Our data also
revealed a positive correlation between the presence of adipocytes (by IHC) and immune cells
and stratified our patient cohort into two groups based on the intrinsic properties of the immune
response.

The largest cluster of cells was cancer epithelial cells, which composed about 50% of cells
analyzed. These cells expressed various genes associated with metastatic disease including
MUC16 and PAX8, consistent with previous reports. The gene expression matrix was similar
between cancer cells from each patient. The variety of cancer associated genes expressed by
these cancer cells confirm the compensatory mechanisms that continue to progress disease.
Another approach to suppressing tumor growth is by identifying and targeting cancer stem cell
(CSC) in ovarian cancer[47]. Like other studies, we were unable to identify cells that expressed
known makers of stem cells[48,49]. However, we identified a cell population closely resembling ESC
and connected to the epithelial cell cluster that highly expressed proliferative marker, Ki-67.
Successful identification for isolation and interrogation of putative ESC might provide useful
insight about CSC in ovarian cancer.

Infiltration of immune cells into the omentum during metastatic cancer does not always elicit
anti-tumor responses[38]. In ovarian cancer, high tumor infiltrating T cells is associated with
significantly longer overall survival[15]. There is evidence that associates immunogenicity of neo-
antigens generated due to high tumor mutational load and clinical response to
immunotherapies[50-53]. To this end, we sought to characterize the differences between the two
groups in our patient cohort stratified by scRNA-seq data. Macrophages make up 70% of the
leukocytes present in the normal omentum so we characterized them first. Although TAMs were
found in both groups, an extra 2 clusters expressing NR1H2 and CD274 were found in the high

T_{inf} group. Both these markers are upregulated in an activation dependent manner suggesting
that the high T_{inf} group is actually mounting an anti-tumor immune response. Flt3$^+$ progenitors
were unique to the low T_{inf} group. Along with the presence of TAMs, these data suggest that
macrophages in the low T_{inf} group acquired immunosuppressive phenotype in response to the
presence of cancer cells.
438
Advances in personalized medicine has revealed TOX as a transcription factor expressed by T cells
that respond to immune checkpoint block[45,54]. In our patient cohort, TOX was mainly expressed
by two T cell clusters, CD8$^+$Trm and CD4$^+$GNLY in the high T_{inf} group. The low T_{inf} group also had
the CD8$^+$Trm subset which expressed granulysin but not a separate CD4$^+$GNLY subset.
Interestingly, there was minimal TOX expression in the low T_{inf} group. Layilin, a marker gene
expressed by exhausted CD8$^+$T cells and regulatory T cells, was mainly expressed by the
CD4$^+$FOXP3$^+$ cluster in our study. Only a few CD8$^+$T cells expressed Layilin. Coincidentally, there
is also a difference in the B cell population between the low and high T_{inf} groups. Along with the
expression of TOX in T cells, the high T cell response group also had a unique subset of
BLIMP1$^+$CD38$^+$CD138$^+$ B cells and plasmablasts. It is likely that antigen-cognate plasmablasts are
involved in the generation or maintenance of TOX$^+$ T cells. A mechanistic study of these cells is
necessary as they may hold the key into inducing TOX$^+$T cells in patients who do not respond to
immune checkpoint inhibitors.
452
In addition to the major cell types analyzed, there were other clusters including 3 undecideds
(N1, N2, N3), 1 endothelial, 1 mesenchymal stem cell and 1 fibroblast cluster. For the undecided
clusters, by mapping them to the CellAtlas cell types, they best correlate with astrocytes,
CMP/bone marrow progenitor and plasmacytoid dendritic cells respectively. We did not recover
adipocytes as they were probably lost during the tissue dissociation. Endothelial cells are the
major cell types involved in angiogenesis which is necessary for metastasis. The mesenchymal
stem cells are connected to the fibroblasts. Inflamed omentum contains stem cells displaying
similar surface markers to MSCs. These stem cells are capable of differentiating into fat, cartilage
or bone depending on the secreted factors present[55,56]. Finally, there was a significant fibroblast
cluster in our sample population. Metastatic transformation of the omentum changes the cellular
composition from mainly adipocytes to cancer cells, immune cells and fibroblasts[28]. Cancer
associated fibroblasts mainly function to remodel the extracellular matrix in the tumor
microenvironment[57]. Recently, a scRNA-seq study of CAF in pancreatic cancer revealed a
LRRC15$^+$ CAF population that correlated with poor response in patients treated with anti-PD-L1
therapy[58]. A few of the fibroblasts in our study also express LRRC15 (data not shown). Ultimately,
more studies are needed to further elucidate these cells and how they can be manipulated to
enhance immunotherapeutic approaches.
470

471   ## Conclusion

472
To our knowledge, this is the first scRNA-seq study that allows stratification of patient samples
all initially pathologically classified as stage III, according to immune response in metastatic
ovarian cancer. Concurrent transcriptomic analysis of cancer and stromal cells revealed patient

heterogeneity highlighting the need for personalized medicine. Interrogating tumor infiltrating
lymphocytes at the single cell level also revealed patient-intrinsic responses that are previously
undescribed in ovarian cancer. Mechanistic studies are required to determine the role of these
cells in tumor progression and response to novel cancer immunotherapies.
480

Acknowledgment

We are grateful to Drs. Mengjie Chen and Ernst Lengyel for helpful discussions and Mark Lingen
for tissue access. All tissues were histologically graded, de-identified and obtained from the
University of Chicago Human Tissue Resource Center. Computational resources and data storage
were provided by the University of Chicago Research Computing Center. This work was
performed, in part, at the Center for Nanoscale Materials, a U.S. Department of Energy Office of
Science User Facility, and supported by the U.S. Department of Energy, Office of Science, under
Contract No. DE-AC02-06CH11357.

490
491
492

References

1   Siegel, R. L., Miller, K. D. & Jemal, A. Cancer Statistics, 2017. *CA: a cancer journal for*
clinicians **67**, 7-30, doi:10.3322/caac.21387 (2017).
2   Testa, U., Castelli, G. & Pelosi, E. Genetic Abnormalities, Clonal Evolution, and Cancer
Stem Cells of Brain Tumors. *Med Sci (Basel)* **6**, doi:10.3390/medsci6040085 (2018).
3   Tothill, R. W. *et al.* Novel molecular subtypes of serous and endometrioid ovarian cancer
linked to clinical outcome. *Clinical cancer research : an official journal of the American*
Association for Cancer Research **14**, 5198-5208, doi:10.1158/1078-0432.CCR-08-0196
(2008).
4   Network, C. G. A. R. Integrated genomic analyses of ovarian carcinoma. *Nature* **474**, 609
(2011).
5   Zheng, G. X. *et al.* Massively parallel digital transcriptional profiling of single cells. *Nat*
Commun **8**, 14049, doi:10.1038/ncomms14049 (2017).
6   Villani, A. C. *et al.* Single-cell RNA-seq reveals new types of human blood dendritic cells,
monocytes, and progenitors. *Science* **356**, doi:10.1126/science.aah4573 (2017).
7   Dixit, A. *et al.* Perturb-Seq: Dissecting Molecular Circuits with Scalable Single-Cell RNA
Profiling of Pooled Genetic Screens. *Cell* **167**, 1853-1866 e1817,
doi:10.1016/j.cell.2016.11.038 (2016).
8   Jaitin, D. A. *et al.* Dissecting Immune Circuits by Linking CRISPR-Pooled Screens with
Single-Cell RNA-Seq. *Cell* **167**, 1883-1896 e1815, doi:10.1016/j.cell.2016.11.039 (2016).
9   Shalek, A. K. *et al.* Single-cell transcriptomics reveals bimodality in expression and splicing
in immune cells. *Nature* **498**, 236-240, doi:10.1038/nature12172 (2013).
10   Zheng, C. *et al.* Landscape of Infiltrating T Cells in Liver Cancer Revealed by Single-Cell
Sequencing. *Cell* **169**, 1342-1356 e1316, doi:10.1016/j.cell.2017.05.035 (2017).

11   Hu, Z. *et al.* The Repertoire of Serous Ovarian Cancer Non-genetic Heterogeneity Revealed
by Single-Cell Sequencing of Normal Fallopian Tube Epithelial Cells. *Cancer cell* **37**, 226-
242 e227, doi:10.1016/j.ccell.2020.01.003 (2020).

12   Platell, C., Cooper, D., Papadimitriou, J. M. & Hall, J. C. The omentum. *World J*
Gastroenterol **6**, 169-176, doi:10.3748/wjg.v6.i2.169 (2000).

13   Clark, R. *et al.* Milky spots promote ovarian cancer metastatic colonization of peritoneal
adipose in experimental models. *The American journal of pathology* **183**, 576-591,
doi:10.1016/j.ajpath.2013.04.023 (2013).

14   Nieman, K. M. *et al.* Adipocytes promote ovarian cancer metastasis and provide energy
for rapid tumor growth. *Nature medicine* **17**, 1498-1503, doi:10.1038/nm.2492 (2011).

15   Santoiemma, P. P. & Powell, D. J., Jr. Tumor infiltrating lymphocytes in ovarian cancer.
Cancer Biol Ther **16**, 807-820, doi:10.1080/15384047.2015.1040960 (2015).

16   Macosko, E. Z. *et al.* Highly Parallel Genome-wide Expression Profiling of Individual Cells
Using Nanoliter Droplets. *Cell* **161**, 1202-1214, doi:10.1016/j.cell.2015.05.002 (2015).

17   Frankish, A. *et al.* GENCODE reference annotation for the human and mouse genomes.
Nucleic Acids Res **47**, D766-D773, doi:10.1093/nar/gky955 (2019).

18   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21,
doi:10.1093/bioinformatics/bts635 (2013).

19   Selewa, A. *et al.* Systematic Comparison of High-throughput Single-Cell and Single-
Nucleus Transcriptomes during Cardiomyocyte Differentiation. *BioRxiv*,
doi:10.1101/585901 (2019).

20   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. *Nature*
biotechnology **36**, 411-420, doi:10.1038/nbt.4096 (2018).

21   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902 e1821,
doi:10.1016/j.cell.2019.05.031 (2019).

22   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902 (2019).

23   McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and
Projection for Dimension Reduction. *Journal of Open Source Software* **29**, 861,
doi:https://doi.org/10.21105/joss.00861 (2018).

24   Maaten, L. v. d. & Hinton, G. isualizing data using t-SNE. *Journal of machine learning*
research 9 **9**, 2579-2605 (2008).

25   I, T. *et al.* Dissecting the multicellular ecosystem of metastatic melanoma by single-cell
RNA-seq. *Science* **352**, 189-196 (2016).

26   Mabbott, N. A., Baillie, J. K., Brown, H., Freeman, T. C. & Hume, D. A. An expression atlas
of human primary cells: inference of gene function from coexpression networks. *BMC*
Genomics **14**, 632, doi:10.1186/1471-2164-14-632 (2013).

27   Waltman, L. & Eck, N. J. V. A smart local moving algorithm for large-scale modularity-
based community detection. *The European physical journal B* **86**, 471 (2013).

28   Pearce, O. M. T. *et al.* Deconstruction of a Metastatic Tumor Microenvironment Reveals
a Common Matrix Response in Human Cancers. *Cancer Discov* **8**, 304-319,
doi:10.1158/2159-8290.CD-17-0284 (2018).

29   Network., C. G. A. R. Intergrated genomic analyses of ovarian carcinoma. *Nature* **474**, 609-
615, doi:https://doi.org/10.1038/nature10166 (2011).

30　Robertson, D. M. *et al.* Combined inhibin and CA125 assays in the detection of ovarian
cancer. *Clin Chem* **45**, 651-658 (1999).
31　Theriault, C. *et al.* MUC16 (CA125) regulates epithelial ovarian cancer cell growth,
tumorigenesis and metastasis. *Gynecologic oncology* **121**, 434-443,
doi:10.1016/j.ygyno.2011.02.020 (2011).
32　Klemba, A. *et al.* Surface markers of cancer stem-like cells of ovarian cancer and their
clinical relevance. *Contemp Oncol (Pozn)* **22**, 48-55, doi:10.5114/wo.2018.73885 (2018).
33　Coward, J. *et al.* Interleukin-6 as a therapeutic target in human ovarian cancer. *Clinical
cancer research : an official journal of the American Association for Cancer Research* **17**,
6083-6096, doi:10.1158/1078-0432.CCR-11-0945 (2011).
34　Lutgendorf, S. K. *et al.* Interleukin-6, cortisol, and depressive symptoms in ovarian cancer
patients. *Journal of clinical oncology : official journal of the American Society of Clinical
Oncology* **26**, 4820-4827, doi:10.1200/JCO.2007.14.1978 (2008).
35　Scambia, G. *et al.* Prognostic significance of interleukin 6 serum levels in patients with
ovarian cancer. *Br J Cancer* **71**, 354-356, doi:10.1038/bjc.1995.71 (1995).
36　Li, W. *et al.* TGFbeta1 in fibroblasts-derived exosomes promotes epithelial-mesenchymal
transition of ovarian cancer cells. *Oncotarget* **8**, 96035-96047,
doi:10.18632/oncotarget.21635 (2017).
37　Shimotsuma, M., Simpson-Morgan, M. W., Takahashi, T. & Hagiwara, A. Activation of
omental milky spots and milky spot macrophages by intraperitoneal administration of a
streptococcal preparation, OK-432. *Cancer research* **52**, 5400-5402 (1992).
38　Oosterling, S. J. *et al.* Insufficient ability of omental milky spots to prevent peritoneal
tumor outgrowth supports omentectomy in minimal residual disease. *Cancer Immunol
Immunother* **55**, 1043-1051, doi:10.1007/s00262-005-0101-y (2006).
39　Servet-Delprat, C. *et al.* Flt3+ macrophage precursors commit sequentially to osteoclasts,
dendritic cells and microglia. *BMC Immunol* **3**, 15, doi:10.1186/1471-2172-3-15 (2002).
40　Gupta, M. *et al.* IRF8 directs stress-induced autophagy in macrophages and promotes
clearance of Listeria monocytogenes. *Nat Commun* **6**, 6379, doi:10.1038/ncomms7379
(2015).
41　Castrillo, A., Joseph, S. B., Marathe, C., Mangelsdorf, D. J. & Tontonoz, P. Liver X receptor-
dependent repression of matrix metalloproteinase-9 expression in macrophages. *J Biol
Chem* **278**, 10443-10449, doi:10.1074/jbc.M213071200 (2003).
42　N, A. G. & Castrillo, A. Liver X receptors as regulators of macrophage inflammatory and
metabolic pathways. *Biochim Biophys Acta* **1812**, 982-994,
doi:10.1016/j.bbadis.2010.12.015 (2011).
43　Savas, P. *et al.* Publisher Correction: Single-cell profiling of breast cancer T cells reveals a
tissue-resident memory subset associated with improved prognosis. *Nature medicine* **24**,
1941, doi:10.1038/s41591-018-0176-6 (2018).
44　Guo, X. *et al.* Publisher Correction: Global characterization of T cells in non-small-cell lung
cancer by single-cell sequencing. *Nature medicine* **24**, 1628, doi:10.1038/s41591-018-
0167-7 (2018).
45　Yao, C. *et al.* Single-cell RNA-seq reveals TOX as a key regulator of CD8(+) T cell persistence
in chronic infection. *Nat Immunol* **20**, 890-901, doi:10.1038/s41590-019-0403-4 (2019).

46   Khan, O. *et al.* TOX transcriptionally and epigenetically programs CD8(+) T cell exhaustion.
Nature **571**, 211-218, doi:10.1038/s41586-019-1325-x (2019).
47   Bast, R. C., Jr., Hennessy, B. & Mills, G. B. The biology of ovarian cancer: new opportunities
for translation. *Nat Rev Cancer* **9**, 415-428, doi:10.1038/nrc2644 (2009).
48   Shah, M. M. & Landen, C. N. Ovarian cancer stem cells: are they real and why are they
important? *Gynecologic oncology* **132**, 483-489, doi:10.1016/j.ygyno.2013.12.001 (2014).
49   Burgos-Ojeda, D., Rueda, B. R. & Buckanovich, R. J. Ovarian cancer stem cell markers:
prognostic   and   therapeutic   implications.   *Cancer   letters*   **322**,   1-7,
doi:10.1016/j.canlet.2012.02.002 (2012).
50   Van Allen, E. M. *et al.* Genomic correlates of response to CTLA-4 blockade in metastatic
melanoma. *Science* **350**, 207-211, doi:10.1126/science.aad0095 (2015).
51   Rizvi, N. A. *et al.* Cancer immunology. Mutational landscape determines sensitivity to PD-
1   blockade   in   non-small   cell   lung   cancer.   *Science*   **348**,   124-128,
doi:10.1126/science.aaa1348 (2015).
52   Snyder, A. *et al.* Genetic basis for clinical response to CTLA-4 blockade in melanoma. *N
Engl J Med* **371**, 2189-2199, doi:10.1056/NEJMoa1406498 (2014).
53   Aguade-Gorgorio, G. & Sole, R. Genetic instability as a driver for immune surveillance. *J
Immunother Cancer* **7**, 345, doi:10.1186/s40425-019-0795-6 (2019).
54   Siddiqui, I. *et al.* Intratumoral Tcf1(+)PD-1(+)CD8(+) T Cells with Stem-like Properties
Promote   Tumor   Control   in   Response   to   Vaccination   and   Checkpoint   Blockade
Immunotherapy. *Immunity* **50**, 195-211 e110, doi:10.1016/j.immuni.2018.12.021 (2019).
55   Shah, S. *et al.* Cellular basis of tissue regeneration by omentum. *PloS one* **7**, e38368,
doi:10.1371/journal.pone.0038368 (2012).
56   Friedenstein, A. J., Petrakova, K. V., Kurolesova, A. I. & Frolova, G. P. Heterotopic of bone
marrow.   Analysis   of   precursor   cells   for   osteogenic   and   hematopoietic   tissues.
Transplantation **6**, 230-247 (1968).
57   Kalluri, R. The biology and function of fibroblasts in cancer. *Nat Rev Cancer* **16**, 582-598,
doi:10.1038/nrc.2016.73 (2016).
58   Dominguez, C. X. *et al.* Single-Cell RNA Sequencing Reveals Stromal Evolution into
LRRC15(+)   Myofibroblasts   as   a   Determinant   of   Patient   Response   to   Cancer
Immunotherapy. *Cancer Discov* **10**, 232-253, doi:10.1158/2159-8290.CD-19-0644 (2020).
636

Table 1. De-identified meta-data for all omental tumors collected from ovarian cancer patients.

Omentum	Age	Race	Origin	Histologic type	Histological grade	Neoadjuvant therapy	Stage (PMN/FIGO)
PT-3232	46	Asian	Left fallopian (STIC)	High grade serous carcinoma	High grade	No	pT3c Nx / IIIc
PT-3401	56	White	Left ovary	High grade serous carcinoma	High grade	No	pT3c Nx Mx/ IIIc
PT-6885	66	Black	Left fallopian (STIC)	High grade serous carcinoma	High grade	Yes	ypT3c N1a / IIIc
PT-5150	62	White	undetermined	Serous	Not applicable	Yes	ypT3a Nx M1/ Ivb
PT-2834	66	Asian	Fallopian	Malignant mixed Mullerian tumor	High grade	Yes	ypT3c Nx/ IIIc
PT-4806	71	Black	Left fallopian (STIC)	High grade serous carcinoma	High grade	Yes	pT3c, N1, M1/ IIIc

Figure 1. Immunohistochemical staining and sample description. A) The hematoxylin and eosin (H&E) stained sections with B) the histogram of percentage area of adipocytes on the patient sample. C) Cytokeratin-7 (CK-7) staining on patient sample with D) histogram of the percentage of CK-7 positive cells. E) Vimentin staining on patient sample with F) the histogram of percentage of Vimentin positive cells. G) CD45 staining on patient sample with H) histogram of the percentage of CD45 positive cells. Images were taken at A) 40x; C), E), G) 400x.

Table 2. Filter parameters applied to samples collected from six patients and the resulting number of cells and genes per sample.

	Gene Cutoff	No of Genes	No of Cells
PT-5150	600	24776	1244
PT-3401	800	32024	3451
PT-6885	800	24063	1071
PT-3232	700	19297	1102
PT-4806	700	26499	1108
PT-2834	800	26221	1909

Figure 2. Cell and molecular subtypes assignment using Drop-seq data. A) UMAP plot for high-quality cells, based on number of genes detected per cell in all six omentum samples, colored by clustering results. B) Ratio of cellular composition by patients. C) Cancer subtype designation for each patient sample based on TCGA. D) Cancer subtype designation by cell type based on TCGA. E) Heatmap of top 150 genes for single cells (cancer and ESCs) in each patient sample. F) UMAP and feature plots of relevant marker genes of 4733 annotated cancer cell-types only from all six omentum.

Figure 3. scRNA-seq of annotated immune population from omental tumors. A) UMAP plot separated by colors showing the 4 main immune cell-types based on correlation with Cell Atlas cell-type. B) Heatmap of immune cell types (T cell, B cell, plasma B cell, and Monocyte) in each patient sample with dendrograms on cell types (columns) and patients (rows), dividing the samples into high and low T cell infiltration groups. C & D) UMAP plots of unsupervised clustering of annotated macrophages from C) high T_{inf} (383 cells), and D) low T_{inf} (312 cells) groups. Heat maps from single cell analysis showing differentially expressed markers between clusters in the two groups: E) high T_{inf}, and F) low T_{inf} group. G&H) Feature plots demonstrating expression of relevant genes in macrophages in high and low high T_{inf} groups, respectively.

Figure 4. Characterization of annotated T cell population. UMAP plot for T cells in A) high T_{inf} (820 cells), and B) low T_{inf} (136) group. Heat maps from single-cell analysis of key genes in different clusters in C) high T_{inf}, and D) low T_{inf} groups. Clusters in heat map are indicated by the same color as in the UMAP plots. Feature plots showing expression of key genes in E) high T_{inf}, and F) low T_{inf} groups.

Figure 5. Identification of B cell clusters using single cell analysis. UMAP plot for B cells in A) high T_{inf} (396 cells) and B) low T_{inf} (124 cells) groups. Heat maps from single-cell analysis of key genes in different clusters in C) high T_{inf}, and D) low T_{inf} groups. Clusters in heat map are indicated by the same color as in the UMAP plots. Feature plots showing expression of key genes in E) high T_{inf}, and F) low T_{inf} groups.