Adhesivity and motility of a zoonotic spirochete: Implications in host-dependent pathogenicity

Jun Xu[1], Nobuo Koizumi[2], Shuichi Nakamura[3]

[1]Department of Animal Microbiology, Graduate School of Agricultural Science, Tohoku University, Sendai, Miyagi, Japan
[2]Department of Bacteriology I, National Institute of Infectious Diseases, Shinjuku-ku, Tokyo, Japan
[3]Department of Applied Physics, Graduate School of Engineering, Tohoku University, Sendai, Miyagi, Japan

Corresponding author: Shuichi Nakamura.
Address: Department of Applied Physics, Graduate School of Engineering, Tohoku University, 6-6-05 Aoba, Aoba-ku, Sendai, Miyagi 980-8579, Japan.
Contact: naka@bp.apph.tohoku.ac.jp

Keywords: Host preference; bacterial infection; adhesivity; motility; spirochete; leptospirosis

Bacterial pathogens can infect particular hosts or have a broad range of host species. Mechanisms of host preferences involve complicated host-pathogen associations such as bacterial adherence, gene regulation, and the host's immune responses[1]. Here, we show the role of adhesivity and motility of the zoonotic spirochete *Leptospira* in host-dependent pathogenicity. *Leptospira* spp. are classified into more than 300 serovars and whereas infection of susceptible mammals can cause clinical symptoms, it can also result in an asymptomatic colonization in some cases, thereby generating maintenance hosts[2–5]. Although the severity of leptospirosis is generally dependent on a combination of *Leptospira* serovars and host species, the scenarios leading to such diverse consequences of infection remain unclear. To investigate the underlying mechanisms of the different outcomes of *Leptospira* infection, we infected kidney cells with various prepared pairs of *Leptospira* serovars and analyzed the *Leptospira* dynamics on the host cells. Quantitatively monitoring the behavior of individual *Leptospira* cells showed that adherence and movement on the host cell surface, called "crawling" correlated with the severity of infection in the host. In the

most severe cases, a large fraction of leptospires attached to the host cells and persistently traveled long distances using the crawling mechanism. The biophysical implications of the kinetics and kinematic features of these bacterial pathogens were found to be critical factors for disease severity in their hosts.

Some bacterial pathogens are specialized to invade a very limited array of hosts, whereas others can infect multiple host species. The host range differs for each pathogen and the clinical symptoms depend on each host-pathogen combination. For example, *Bacillus anthracis* primarily infects livestock and wild herbivores, transmitting to humans via direct or indirect contact with infected animals or animal products. However, its occurrence in carnivores is rare. Salmonellosis is caused by over 2000 serovars of *Salmonella enterica* and its transmission occurs from reptile reservoirs to a wide range of susceptible hosts including humans and other mammals. Virulence in the case of plant bacterial pathogens like *Actinobacteria* spp. and *Pantoea* spp. is host specific, and is believed to be related to the structure of the root microbiome and other factors that optimize their symbiosis (e.g., plant hormones, drought stress, temperature and salinity)[6,7]. In some cases, the mechanisms correlating host dependency to bacterial pathogenicity have been investigated. A correlation was found between the host range of the opportunistic food-poisoning bacterium *Listeria monocytogenes* and its host binding affinity through the listerial protein internalin A, which interacted with E-cadherin in intestinal epithelial cells thereby facilitating its invasion[8]. Here, we show that the adhesivity and motility of the zoonotic spirochete *Leptospira interrogans* are crucial factors responsible for their host-dependent pathogenicity.

Leptospirosis is a worldwide zoonosis, affecting various mammalian hosts such as livestock (cattle, pigs, horses, and others), companion animals (dogs and others), and humans, with consequences varying from asymptomatic to mild and fatal symptoms[2–4]. The causative agents *Leptospira* spp. are Gram-negative bacteria belonging to the phylum Spirochaetes. The genus *Leptospira* is comprised by over 300 serovars defined based on the structural diversity of lipopolysaccharide (LPS)[2,9]. Humans and other animals develop clinical disease; however, some animal species can become an asymptomatic reservoir for particular *Leptospira* serovars. Thus, the outcome of *Leptospira* infection depends on the

host-serovar association. Some host-serovar pairs can cause severe symptoms, such as hemorrhage, jaundice, and nephritis. In contrast, the same serovars can be maintained in the renal tubules of reservoir animals that will then intermittently shed leptospires in their urine over long periods[3,5]. The urinary shedding of leptospires to the environment leads to infection in humans and other animals through contact with contaminated soil or water. *Leptospira* spp. do not have an exotoxin-secretion machinery such as the type III secretion system[2]. The pathogenic *Leptospira*-specific protein Loa22 is known to be an essential virulence factor[10], but other proteins such as the fibronectin-binding protein LigB and the outer membrane protein LipL32 expressed solely in pathogenic strains, are not involved in their pathogenicity[11,12]. Despite these virulence factors being common among pathogenic *Leptospira* serovars, the host-dependent mechanisms of pathogenicity remain unclear.

Studies in animal models have shown that motility is an important factor involved in leptospiral virulence[13,14]. Motility is a critical component that facilitates infection for many species of bacteria[15]. *Helicobacter pylori* requires motility to migrate towards the epithelial tissue in the stomach[16] and motility and chemotaxis are key factors that guide host invasion in different *Salmonella* serovars [17,18]. In the case of Lyme disease, movement and adhesion by the spirochete *Borrelia burgdorferi* in blood vessels are thought to be important during the process of host cell invasion[19]. *Leptospira* spp. have two periplasmic flagella (PFs) beneath the outer cell membrane (Fig. 1A). The rotation of the PFs in the periplasmic space gyrates both ends of the cell body and rotates the coiled cell body allowing *Leptospira* not only to swim in fluids but also to crawl over solid surfaces[20,21]. Adherence and entry of pathogenic leptospires in the conjunctival epithelium[4] and in the paracellular routes of hepatocytes[22] were previously observed using scanning electron microscopy, raising the possibility that adhesion to the host tissue surfaces and subsequent surface movements are determinants of the host-dependent leptospiral pathogenicity. To verify this hypothesis, assuming the transition of leptospires between swimming and adhesion states and between adhesion and crawling states in the equilibrium (Fig. 1B), we quantitatively determined the adhesion and crawling motility of three *Leptospira* strains: the pathogenic *L. interrogans* serovars Icterohaemorrhagiae and Manilae and the nonpathogenic *L. biflexa* serovar Patoc on

cultured kidney cells from six different host species. We then discussed the correlation of the

leptospiral dynamics with host-dependent outcomes using statistical clustering based on the

measured parameters.

We first measured the fractions of swimming [S], adhered [A], and crawling [C]

Leptospira cells on the kidney cells (Figs. 1C-D and 2A), showing the dependence of

leptospiral dynamics on the host-serovar combination (Fig. 2B). We calculated the

equilibrium constant between swimming and adhesion states (K_{S-A}) and between adhesion

and crawling states (K_{A-C}) from the cell fractions using K_{S-A} = [A] / [S] and K_{A-C} = [C] / [A],

respectively (Fig. 2C). The data points scattered on the ternary plot of the cell fraction and

the pairwise plot of the equilibrium constants were statistically classified into three clusters.

Next, we measured the crawling speed of each individual cell on the kidney cells (Fig.

3A). Concerning the crawling motility, we observed that some leptospiral cells attached to

the kidney cells and moved smoothly for periods of time, migrating over long-distances (Figz.

3B, D, and F; Supplementary Movie 2), whereas others frequently reversed the crawling

direction (Figs. 3C, E, and G; Supplementary Movie 3). Cell movements with reversals are

considered to be diffusive and thus can be evaluated by plotting the mean square

displacement (MSD) against time, a general methodology for diffusion (Brownian motion)

analysis[23]. The MSD of simple diffusion without directivity is proportional to time, and

therefore double-logarithmic MSD plots from such non-directional diffusion represent slopes

of ~1, whereas those from directive movements show MSD slopes of ~2, representing the

relatively long distance travelled by the cells (Supplementary Fig. 1). Double-logarithmic

MSD plots obtained from each individual leptospiral cells showed a wide range of MSD

slopes and differed for each host-serovar pair (Figs. 3H-J and Supplementary Fig. 2). The

data of the crawling speeds and MSD slopes (i.e. crawling persistency) were also grouped

into statistical clusters.

To investigate the relevance of the measured parameters on the outcomes of leptospiral

infection, we translated Figs. 2–3 into Figs. 4A–D, where the mean values obtained for each

statistical cluster were indicated, and the serovar-host pairs were discriminated by their

clinical aspects, according to a literature search: red, pairs that exhibit any clinical signs;

yellow, isolation of leptospires from asymptomatic animals or natural reservoirs (i.e., rats); green, pairs containing the nonpathogenic *Leptospira* strain (i.e., non-infection); grey, no report (Table 1). Red, green and yellow pairs were mostly grouped together, although yellow pairs fluctuated between clusters with a red or green pair majority. Given that the determined clusters reflect the differences in outcomes of the host after infection by each *Leptospira* strain, our results suggest the involvement of adhesion and crawling in leptospiral pathogenicity. Fig. 4A shows the increased crawling fraction of the virulent cluster, whereas >75% cells remained in the swimming state in the non-infection cluster. This is consistent with the observation that the virulence pairs had a significantly larger K_{A-C} in comparison with asymptomatic and non-pathogenic pairs ($P < 0.05$); K_{S-A} did not seem to correlate to virulence (Fig. 4B). These thermodynamic parameters suggest that the biased transition from adhesion to crawling would be responsible for the virulence of *Leptospira*. Although the crawling speed could not be correlated to pathogenicity (Fig. 4C), a detailed analysis of the crawling patterns revealed that long-distance directive crawling was a crucial virulence factor (Fig. 4D). The crawling motility of *Leptospira* is caused by the attachment of the spirochete cell body to surfaces via adhesive cell surface components[20,24]. The successive alternation in the attachment and detachment of adhesins allows for this progressive movement by the spirochetes, however, an excessively strong adhesion can inhibit crawling[20]. LPS, the molecular basis for the identification of the different *Leptospira* serovars[2,9], is thought to be a crucial adhesin important for this crawling motion[20]. Thus, it is possible that compatibility of the serological characteristics of leptospires and the surface properties of the host tissue might affect the crawling behavior over the tissue surfaces and the subsequent clinical consequences. The results of our biophysical experiments outline a plausible framework for the adhesion and crawling-dependent pathogenicity of *Leptospira* (Fig. 4E). The biased transition from the adhesion to the crawling state and the long-distance, persistent crawling allows for the leptospires to explore the host's cell surfaces, increasing the probability of encountering routes for invasion through their intracellular tight junctions (Fig. 4E, left). In contrast, the weakly attached leptospires can detach easily and be swept by external forces, such as intermittent urination (Fig. 4E, right).

Adherence of microbes is generally important in the early stage of infection. For example,

the enteropathogenic *Escherichia coli* (EPEC) uses H2 or H6 flagella for adhesion[25], and

host-dependent pathogenicity of the EPEC strain RDEC-1 infection is associated to the

adherence affinity of the bacteria to the host intestinal epithelium[26]. Our results showed the

importance of adhesion to and persistent crawling on the host tissue for disease severity of

Leptospira infection, and concurrently highlighted the close relationship of such surface

dynamics of the spirochetal pathogen to their host-dependent pathogenicity. These findings

could lead to the development of antimicrobials for leptospirosis targeting the pathogens'

adhesion and crawling motility and provides a significant step forward to confront other

host-dependent bacterial infections. Meanwhile, microarray analyses have revealed that

regulation of gene expression in *Leptospira* is affected by its interaction with host cells[27,28].

Although host immune responses against *Leptospira* infection remain mostly unknown,

pathogenic leptospires interfere with the complement system through the degradation of

complement proteins[29] and inhibition of coagulation via binding to thrombin[30]. Thus, in

addition to bacterial dynamics, the molecular and immunological understanding of the

host-pathogen association will realize the establishment of more effective prevention

strategy.

Methods

***Leptospira* strains and growth conditions.** Pathogenic serovars of *L. interrogans*

serovar Icterohaemorrhagiae (strain WFA135), an isolate from *Rattus norvegicus* in Tokyo,

Japan, serovar Manilae (strain UP-MMC-NIID)[31,32] and a saprophytic *L. biflexa* serovar

Patoc (strain Patoc I) were used in this study. The serovar of WFA135 was determined by

multiple loci variable number of tandem repeats analysis (MLVA) and DNA sequencing of the

lic12008 gene[32,33]. Bacteria were cultured in enriched

Ellinghausen-McCullough-Johnson-Harris (EMJH) liquid medium (BD Difco, NJ, USA)

containing 25 µg/mL spectinomycin at 30 °C for 2 (*L. biflexa*) or 4 (*L. interrogans*) days until

the stationary phase. To track *Leptospira* cells when in co-culture with mammalian kidney

cells, a green fluorescent protein (GFP) was constitutively expressed in each strain (Fig. 1D;

Supplementary Movie 1).

Mammalian cells and media. Mammalian kidney epithelial cell lines used included
MDCK-NBL2 (dog), NRK-52E (rat), Vero (monkey), MDBK-NBL1 (cow), TCMK-1 (mouse),
and HK-2 (human). MDCK, Vero, TCMK, and MDBK cells were maintained in Eagle's
minimum essential media (MEM) (Sigma-Aldrich, Darmstadt, Germany) containing 10% fetal
bovine serum or 10% horse serum (Nacalai Tesque, Kyoto, Japan) at 37 °C and 5% CO_2.
NRK cells were maintained in Dulbecco's modified Eagle's media (DMEM) (Thermo Fisher
Scientific, MA, USA) with 4 mM L-glutamine, adjusted to contain 1.5 g/L sodium bicarbonate
and 4.5 g/L glucose, and with 5% bovine calf serum (Nacalai Tesque) at 37 °C and 5% CO_2.
HK-2 cells were maintained in keratinocyte serum-free (KSF) media with 0.05 mg/mL bovine
pituitary extract and 5 ng/mL human recombinant epidermal growth factor (Gibco - Thermo
Fisher Scientific, MA, USA). All culture conditions contained a 5% antibiotic / antimycotic
mixed solution (Nacalai Tesque). Cells were treated with a 0.1% trypsin - EDTA solution
(Nacalai Tesque) to dislodge the cells during each passage process.

GFP expression in *L. interrogans* and *L. biflexa*. For the construction of a replicable
plasmid in *L. interrogans*, the *rep-parB-parA* region of the *L. interrogans* serovar Canicola
strain Gui44[34] plasmid pGui1 was amplified from a *L. interrogans* serogroup Canicola isolate,
and the amplified product was cloned into the PCR-generated pCjSpLe94[35] by NEBuilder
HiFi DNA Assembly cloning (New England BioLabs), generating pNKLiG1. The *flgC*
promoter region and *gfp* were amplified from pCjSpLe94 and pAcGFP1 (Clontech),
respectively, and the amplified products were cloned into the *Sal*I-digested pNKLiG1 for *L.*
interrogans or the *Sal*I-digested pCjSpLe94 for *L. biflexa*. The plasmids were transformed
into strains WFA135, UP-MMC-NIID or Patoc I by conjugation with *E. coli* β2163 harboring
the plasmid[36]. Primer sequences used in this study are listed in Supplementary Table 1.
Expression of GFP did not affect motility in the *Leptospira* serovars (Supplementary Fig. 3).

Preparation of kidney cells and *Leptospira* cells in a chamber slide. Kidney cells were

harvested with 0.1% trypsin and 0.02% EDTA in a balanced salt solution (Nacalai Tesque) and plated onto a chamber slide (Iwaki, Tokyo, Japan) using their corresponding media without antibiotics. The slides were incubated for 48 h until a monolayer was formed and washed twice with media to remove non-adherent cells. The cells were incubated for a further 2 h at 37 °C and 5% CO_2. Approximately 500 ml of stationary phase *Leptospira* cells were harvested by centrifugation at 1,000 *g* for 10 min at room temperature, washed twice in PBS, then resuspended in the corresponding kidney cell culture media without antibiotics at 37 °C to a concentration of 10^7 cells / ml. These suspensions (1 ml) were then added into the corresponding chamber slides containing the kidney cell layer, and the chamber slides were incubated at 37 °C for 1 h.

Microscopy observation and adhesion-crawling assay. The movement behaviors, swimming, adhesion and crawling of the *Leptospira* cells on the kidney cells were observed using a dark-field microscope (BX53, Splan 40×, NA 0.75, Olympus, Tokyo, Japan) with an epi-fluorescent system (U-FBNA narrow filter, Olympus) and recorded by a CCD-camera (WAT- 910HX, Watec Co., Yamagata, Japan) at 30 frames per second. *Leptospira* cells were tracked using an ImageJ (NIH, MD, USA)-based tracking system and the motion parameters such as motile fraction, velocity and MSD were analyzed using Excel-based VBA (Microsoft, WA, USA). The two-dimensional MSD of individual leptospiral cells during a period Δt was calculated by the following equation: $MSD(\Delta t) = \langle (x_{i+\Delta t} - x_i)^2 + (y_{i+\Delta t} - y)^2 \rangle$, where ($x_i$, y_i) is the bacterial position at *I* (see also Supplementary Fig. 1).

Statistical analysis. All experiments were performed in triplicate. Statistical differences between data were evaluated using a Student's *t*-test. The data clustering was performed independently for each experiment using the k-means method in OriginPro. The clustering method grouped the data population into a specified number of clusters referring to the Euclidian distance of each data point from the centroid of the cluster, calculated at every clustering process, and reclassifying the data point to the nearest cluster.

References

1. Bäumler, A. & Fang, F. C. Host specificity of bacterial pathogens. *Cold Spring Harb. Perspect. Med.* **3**,
(2013).
2. Picardeau, M. Virulence of the zoonotic agent of leptospirosis: still terra incognita? *Nat. Rev. Microbiol.*
15, 297–307 (2017).
3. Adler, B. & de la Peña Moctezuma, A. *Leptospira* and leptospirosis. *Vet. Microbiol.* **140**, 287–296
(2010).
4. Bharti, A. R. *et al.* Leptospirosis: a zoonotic disease of global importance. *Lancet Infect. Dis.* **3**, 757–
771 (2003).
5. Levett, P. N. Leptospirosis. *Clin. Microbiol. Rev.* **14**, 296–326 (2001).
6. Naylor, D., DeGraaf, S., Purdom, E. & Coleman-Derr, D. Drought and host selection influence
bacterial community dynamics in the grass root microbiome. *ISME J.* **11**, 2691–2704 (2017).
7. Barash, I. & Manulis-Sasson, S. Virulence mechanisms and host specificity of gall-forming *Pantoea*
agglomerans. *Trends Microbiol.* **15**, 538–545 (2007).
8. Lecuit, M. *et al.* A single amino acid in E-cadherin responsible for host specificity towards the human
pathogen *Listeria monocytogenes*. *EMBO J.* **18**, 3956–3963 (1999).
9. Cerqueira, G. M. & Picardeau, M. A century of *Leptospira* strain typing. *Infect. Genet. Evol. J. Mol.*
Epidemiol. Evol. Genet. Infect. Dis. **9**, 760–768 (2009).
10. Ristow, P. *et al.* The OmpA-like protein Loa22 is essential for leptospiral virulence. *PLoS Pathog.* **3**,
e97 (2007).
11. Murray, G. L. *et al.* Major surface protein LipL32 is not required for either acute or chronic infection
with *Leptospira interrogans*. *Infect. Immun.* **77**, 952–958 (2009).
12. Croda, J. *et al.* Targeted mutagenesis in pathogenic *Leptospira* species: disruption of the LigB gene
does not affect virulence in animal models of leptospirosis. *Infect. Immun.* **76**, 5826–5833 (2008).
13. Lambert, A. *et al.* FlaA proteins in Leptospira interrogans are essential for motility and virulence but
are not required for formation of the flagellum sheath. *Infect. Immun.* **80**, 2019–2025 (2012).
14. Wunder, E. A. *et al.* A novel flagellar sheath protein, FcpA, determines filament coiling, translational
motility and virulence for the *Leptospira* spirochete. *Mol. Microbiol.* **101**, 457–470 (2016).
15. Josenhans, C. & Suerbaum, S. The role of motility as a virulence factor in bacteria. *Int. J. Med.*
Microbiol. **291**, 605–614 (2002).
16. Clyne, M., Ocroinin, T., Suerbaum, S., Josenhans, C. & Drumm, B. Adherence of isogenic
flagellum-negative mutants of *Helicobacter pylori* and *Helicobacter mustelae* to human and ferret
gastric epithelial cells. *Infect. Immun.* **68**, 4335–4339 (2000).
17. Olsen, J. E. *et al.* The role of flagella and chemotaxis genes in host pathogen interaction of the host
adapted *Salmonella enterica* serovar Dublin compared to the broad host range serovar
S.Typhimurium. *BMC Microbiol.* **13**, 67 (2013).
18. Siitonen, A. & Nurminen, M. Bacterial motility is a colonization factor in experimental urinary tract
infection. *Infect. Immun.* **60**, 3918–3920 (1992).
19. Ebady, R. *et al.* Biomechanics of *Borrelia burgdorferi* vascular interactions. *Cell Rep.* **16**, 2593–2604
(2016).
20. Tahara, H. *et al.* The mechanism of two-phase motility in the spirochete *Leptospira*: Swimming and
crawling. *Sci. Adv.* **4**, eaar7975 (2018).
21. Cox, P. J. & Twigg, G. I. Leptospiral motility. *Nature* **250**, 260–261 (1974).
22. Miyahara, S. *et al.* Destruction of the hepatocyte junction by intercellular invasion of *Leptospira*
causes jaundice in a hamster model of Weil's disease. *Int. J. Exp. Pathol.* **95**, 271–281 (2014).

23. Kusumi, A., Sako, Y. & Yamamoto, M. Confined lateral diffusion of membrane receptors as studied by
single particle tracking (nanovid microscopy). Effects of calcium-induced differentiation in cultured
epithelial cells. *Biophys. J.* **65**, 2021–2040 (1993).

24. Charon, N. W., Lawrence, C. W. & O'Brien, S. Movement of antibody-coated latex beads attached to
the spirochete *Leptospira interrogans*. *Proc. Natl. Acad. Sci. U. S. A.* **78**, 7166–7170 (1981).

25. Girón, J. A., Torres, A. G., Freer, E. & Kaper, J. B. The flagella of enteropathogenic *Escherichia coli*
mediate adherence to epithelial cells. *Mol. Microbiol.* **44**, 361–379 (2002).

26. Cheney, C. P., Schad, P. A., Formal, S. B. & Boedeker, E. C. Species specificity of *in vitro Escherichia*
coli adherence to host intestinal cell membranes and its correlation with *in vivo* colonization and
infectivity. *Infect. Immun.* **28**, 1019–1027 (1980).

27. Satou, K. *et al.* Complete genome sequences of low-passage virulent and high-passage avirulent
variants of pathogenic *Leptospira interrogans* serovar Manilae strain UP-MMC-NIID, originally
isolated from a patient with severe leptospirosis, determined using PacBio single-molecule real-time
technology. *Genome Announc.* **3**, (2015).

28. Toma, C. *et al.* Leptospiral outer membrane protein LMB216 is involved in enhancement of
phagocytic uptake by macrophages. *Cell. Microbiol.* **16**, 1366–1377 (2014).

29. Fraga, T. R. *et al.* Immune evasion by pathogenic *Leptospira* strains: the secretion of proteases that
directly cleave complement proteins. *J. Infect. Dis.* **209**, 876–886 (2014).

30. Fernandes, L. G., de Morais, Z. M., Vasconcellos, S. A. & Nascimento, A. L. T. O. *Leptospira*
interrogans reduces fibrin clot formation by modulating human thrombin activity via exosite I. *Pathog.*
Dis. **73**, (2015).

31. Koizumi, N. & Watanabe, H. Leptospiral immunoglobulin-like proteins elicit protective immunity.
Vaccine **22**, 1545–1552 (2004).

32. Santos, L. A. *et al.* Genomic comparison among global isolates of *L. interrogans* serovars
Copenhageni and Icterohaemorrhagiae identified natural genetic variation caused by an Indel. *Front.*
Cell. Infect. Microbiol. **8**, 193 (2018).

33. Koizumi, N. *et al.* Multiple-locus variable-number tandem repeat analysis of *Leptospira interrogans*
and *Leptospira borgpetersenii* isolated from small feral and wild mammals in East Asia. *Infect. Genet.*
Evol. J. Mol. Epidemiol. Evol. Genet. Infect. Dis. **36**, 434–440 (2015).

34. Zhu, W.-N. *et al.* Isolation and characterization of two novel plasmids from pathogenic *Leptospira*
interrogans serogroup Canicola serovar Canicola strain Gui44. *PLoS Negl. Trop. Dis.* **8**, e3103
(2014).

35. Picardeau, M. Conjugative transfer between *Escherichia coli* and *Leptospira* spp. as a new genetic
tool. *Appl. Environ. Microbiol.* **74**, 319–322 (2008).

36. Slamti, L. & Picardeau, M. Construction of a library of random mutants in the spirochete *Leptospira*
biflexa using a *mariner* transposon. *Methods Mol. Biol. Clifton NJ* **859**, 169–176 (2012).

37. Faine, S., Adler, B., Bolin, C. & Perolat, P. '*Leptospira*' *and leptospirosis*. (MediSci, 1999).

38. Villanueva, M. A. *et al.* Molecular epidemiology of pathogenic *Leptospira* spp. among large ruminants
in the Philippines. *J. Vet. Med. Sci.* **78**, 1649–1655 (2016).

39. Villanueva, S. Y. A. M. *et al.* Serologic and molecular studies of *Leptospira* and leptospirosis among
rats in the Philippines. *Am. J. Trop. Med. Hyg.* **82**, 889–898 (2010).

40. Gomes-Solecki, M., Santecchia, I. & Werts, C. Animal models of leptospirosis: Of mice and hamsters.
Front. Immunol. **8**, (2017).

41. Ratet, G. *et al.* Live imaging of bioluminescent *Leptospira interrogans* in mice reveals renal

colonization as a stealth escape from the blood defenses and antibiotics. *PLoS Negl. Trop. Dis.* **8**, e3359 (2014).

42. Fanton d'Andon, M. *et al. Leptospira Interrogans* induces fibrosis in the mouse kidney through Inos-dependent, TLR- and NLR-independent signaling pathways. *PLoS Negl. Trop. Dis.* **8**, e2664 (2014).

43. Yanagihara, Y., Villanueva, S. Y. A. M., Yoshida, S.-I., Okamoto, Y. & Masuzawa, T. Current status of leptospirosis in Japan and Philippines. *Comp. Immunol. Microbiol. Infect. Dis.* **30**, 399–413 (2007).

44. Masuzawa, T., Dancel, L. A., Miyake, M. & Yanagihara, Y. Serological analysis of human leptospirosis in the Philippines. *Microbiol. Immunol.* **45**, 93–95 (2001).

45. Organization, W. H. *Human leptospirosis : guidance for diagnosis, surveillance and control*. (World Health Organization, 2003).

46. Koizumi, N. *et al.* Human leptospirosis cases and the prevalence of rats harbouring *Leptospira interrogans* in urban areas of Tokyo, Japan. *J. Med. Microbiol.* **58**, 1227–1230 (2009).

47. Geisen, V. *et al.* Canine leptospirosis infections - clinical signs and outcome with different suspected *Leptospira* serogroups (42 cases). *J. Small Anim. Pract.* **48**, 324–328 (2007).

48. Rentko, V. T., Clark, N., Ross, L. A. & Schelling, S. H. Canine leptospirosis. A retrospective study of 17 cases. *J. Vet. Intern. Med.* **6**, 235–244 (1992).

49. Adin, C. A. & Cowgill, L. D. Treatment and outcome of dogs with leptospirosis: 36 cases (1990-1998). *J. Am. Vet. Med. Assoc.* **216**, 371–375 (2000).

50. Asai, T. *et al.* Prevalence of antibodies to five selected zoonosis agents in monkeys. *J. Vet. Med. Sci.* **53**, 553–559 (1991).

51. Perolat, P. *et al.* Occurrence of severe leptospirosis in a breeding colony of squirrel monkeys. *Am. J. Trop. Med. Hyg.* **46**, 538–545 (1992).

52. Szonyi, B., Agudelo-Flórez, P., Ramírez, M., Moreno, N. & Ko, A. I. An outbreak of severe leptospirosis in capuchin (Cebus) monkeys. *Vet. J. Lond. Engl. 1997* **188**, 237–239 (2011).

53. Pereira, M. M., Andrade, J., Marchevsky, R. S. & Ribeiro dos Santos, R. Morphological characterization of lung and kidney lesions inC3H/HeJ mice infected with *Leptospira interrogans* serovar icterohaemorrhagiae: Defect of CD4+ and CD8+ T-cells are prognosticators of the disease progression. *Exp. Toxicol. Pathol.* **50**, 191–198 (1998).

54. Hässig, M. & Lubsen, J. Relationship between abortions and seroprevalences to selected infectious agents in dairy cows. *Zentralblatt Vet. Reihe B J. Vet. Med. Ser. B* **45**, 435–441 (1998).

55. Guerra, M. A. Leptospirosis. *J. Am. Vet. Med. Assoc.* **234**, 472–478 (2009).

Acknowledgments   We thank Dr. H. Nishimura (Sendai Medical Center) and Dr. C. Toma (University of the Ryukyus) for the generous gift of animal cell lines; and Dr. E. Isogai (Tohoku University) and Dr. H. Yoneyama (Tohoku University) for the experiment reagents and the insightful discussion. This work was supported by the JSPS KAKENHI: 18K07100 for SN and 18J10834 for JX.

Author contributions J.X., N.K. and S.N. planned the project; J.X and N.K. carried out the experiments; S.N. set up the optical system and programs for data analysis; J.X. and S.N. analyzed the data; J.X., N.K. and S.N. wrote the paper.

Competing interests The authors declare that they have no competing interests.

Table 1. Host-dependent pathogenicity of *L. interrogans* serovars Manilae and Icterohaemorrhagiae

Serovar	Host	Major findings of infection (kidney)	References
Manilae	Rat	Persist in the kidneys causing either slight or no harm (maintenance hosts)	Faine et al. (1999), Villanueva et al. (2010), Villanueva et al. (2016)
	Dog	Unknown	NA
	Monkey	Unknown	NA
	Mouse	Pyelonephritis, chronic infection	Gomes-Solecki et al. (2017), Ratet et al. (2014), Fanton d'Andon et al. (2014)
	Cow	Asymptomatic carrier	Villanueva et al. (2016), Yanagihara et al. (2007)
	Human	Oliguria and/or proteinuria, hematuria, renal failure	Yanagihara et al. (2007), Masuzawa et al. (2001)
Icterohaem-orrhagia	Rat	Persist in the kidneys causing either slight or no harm (maintenance hosts)	WHO (2003), Faine et al. (1999), Koizumi et al. (2009)
	Dog	Dehydration, pyelonephritis, hematuria	Geisen et al.(2007), Rentko et al. (1992), Adin et al. (2000)
	Monkey	Hemorrhagic syndrome, renomegaly	Asai et al. (1991), Perolat et al. (1992), Szonyi et al. (2011)
	Mouse	Hemorrhage, pyelonephritis	Pereira et al. (1998), Gomes-Solecki et al. (2017)
	Cow	Hemoglobinuria, anemia, abortion, chronic carrier	Hassig et al. (1998), Guerra. (2009)
	Human	Oliguria and/or proteinuria, hematuria, renal failure	Levett (2001), Guerra. (2009), Faine (1999)

Faine et al. (1999)[37]; Villanueva et al. (2016)[38]; Villanueva et al. (2010)[39]; Gomes-Solecki et al. (2017)[40]; Ratet et al. (2014)[41]; Fanton d'Andon et al. (2014)[42]; Yanagihara et al. (2007)[43]; Masuzawa et al. (2001)[44]; World Health Organization (2003)[45]; Koizumi et al. (2009)[46]; Geisen et al. (2007)[47]; Rentko et al. (1992)[48]; Adin et al. (2000)[49]; Asai et al. (1991)[50]; Perolat et al. (1992)[51]; Szonyi et al. (2011)[52]; Pereira et al. (1998)[53]; Hässig et al. (1998)[54]; Guerra (2009)[55]; Levett (2001)[5].

381
Fig. 1. Working model and schematic representation of the experimental setup. (A)
Schematic diagram of the *Leptospira* cell structure. (B) A three-state kinetic model assuming
a transition between "swimming" (floating above cell layers without physical contact to the
cells) and "adhesion" (attachment to the cell layer without migration), and between
"adhesion" and "crawling" (attachment to the cell layer and movement over surfaces), with
K_{S-A} and K_{A-C} represent the equilibrium constants of each transition, respectively. (C) Fully
confluent kidney cells from either animals or humans were cultured in an observation
chamber, and *Leptospira* cells that constitutively expressed GFP were added to the cultures.
Three *Leptospira* strains (the pathogenic *L. interrogans* serovars Icterohaemorrhagiae and
Manilae, and the saprophytic *L. biflexa* serovar Patoc) and kidney epithelial cell lines from
six species (MDCK-NBL2, dog; NRK-52E, rat; Vero, monkey; MDBK-NBL1, cow; TCMK-1,
mouse; HK-2, human) were used in the experimental model. (D) Example of an
epi-fluorescence image of the *L. interrogans* serovar Icterohaemorrhagiae, as indicated by
the yellow triangles, on the rat kidney cell line, NRK-52E. The nucleus of the kidney cell is
shown by the white arrow (Supplementary Movie 1).
397

Fig. 2. Steady-state motility and adhesion of *Leptospira* cells on kidney cells. (A) Image sequences of swimming (top), adhered (middle), and crawling (bottom) leptospiral cells. Yellow and white arrows indicate *Leptospira* cells and kidney cells, respectively. No kidney cells were observed due to out of focus during measurement of *Leptospira* cells swimming in liquid media (**Fig. 1B**). (B) Ternary plot of the cell fractions in a state of swimming, adhesion, or crawling. The inset schematically explains how to read the ternary plot using an example plot with 30% for I, 45% for II, and 25% for III. Legend, is shown to the right of the ternary plot, and M, I and P indicate the *L. interrogans* serovar Manilae, *L. interrogans* serovar Icterohaemorrhagiae, and *L. biflexa* serovar Patoc, respectively. Each data point represents the mean of triplicate experiments and ~ 2,400 leptospiral cells were measured per host-serovar pair. Dashed circles indicate the results of the k-means clustering. (C) Pairwise plot of the equilibrium constants K_{S-A} and K_{A-C}, calculated from the data shown in **B**. Dashed circles are the results of k-means clustering performed independently of **B**. The statistical analysis used the Student's *t*-test (* $P < 0.05$).

Fig. 3. Analysis of the crawling movement of *Leptospira* on kidney cell layers. (A)
Average crawling speeds determined for each host-serovar pair. Examples of persistent
crawling (B) and diffusive crawling (C) of *Leptospira* cells obtained by single-cell tracking in
10 s. Colors indicate time courses in the order of red orange yellow green blue, and indigo.
(D-E) Examples of crawling trajectories of leptospires on the kidney cell surfaces. (F-G)
Time course of the displacement obtained from **D** and **E**; inset, expanded view of cell
displacements and schematics of motion patterns. **B**, **D**, and **F** show the long-distance
migration represented by directive crawling; **C**, **E**, and **G** show the limited migration due to
frequent reversal. (H-I) Evaluation of the directivity of individual leptospiral cell movements
by MSD vs time plots. MSD plots with slopes ~ 2 indicate persistently directive crawling (**H**),
whereas plots with slopes ~ 1 indicate a non-directional movement (**I**), i.e., motion with high
frequency of reversal patterns and short net migration distance (refer to the schematic
explanation of motion pattern in insets of **F** and **G**. In **B** and **C**, the values of MSD slopes of
each cell are shown in yellow numbers. (J) Average values of MSD slopes as determined
from the MSD analysis. Refer to legend on **Fig. 2B** for symbols shown in **A** and **J**. Each data
point was obtained from at least 90 leptospiral cells. Gray circles in **A** and **J** represent the
results of the k-means clustering. Statistical analysis used the Student's *t*-test, (* $P < 0.05$).

433
434

Fig. 4. Correlation of adhesion, motility and pathogenicity. According to the statistical classification in **Figs 2-3**, host-serovar pairs are listed below each figure. Based on a literature search summarized in **Table 1**, the examined pairs are indicated in red (with registered symptoms), yellow (asymptomatic or natural reservoir), green (non-pathogenic *Leptospira*), or grey (unknown infection outcomes). The correlations of (A) the cell fractions in each state, (B) the thermodynamic parameters, (C) the crawling speed, and (D) the MSD slopes (i.e., the directivity, or persistency of crawling) to the reported clinical consequences (**Table 1**) are shown. **A-D** show the mean values and standard deviations (error bars) calculated for each cluster using the values shown in **Fig. 2** and **Fig. 3**. (E) A plausible model for crawling-dependent pathogenicity of *Leptospira*. In the cases that lead to severe symptoms (left), leptospiral cells were biased to the crawling state, and most of the crawling cells showed directional translation persistently over host tissue surfaces, increasing the invasion probability. For the asymptomatic or non-infectious cases, many leptospires remained in the swimming state and might be removed through body fluids or urination. Some fractions of the adhered leptospires were able to crawl, but their migration distances were limited due to frequent reversal.