Domestication of the emblematic white cheese-making fungus *Penicillium camemberti* and its diversification into two varieties

Jeanne Ropars[1], Estelle Didiot[1], Ricardo C. Rodríguez de la Vega[1], Bastien Bennetot[1], Monika Coton[2], Elisabeth Poirier[2], Emmanuel Coton[2], Alodie Snirc[1], Stéphanie Le Prieur[1] and Tatiana Giraud[1]

[1]Ecologie Systematique et Evolution, Universite Paris-Saclay, CNRS, AgroParisTech, 91400 Orsay, France
[2]Univ Brest, Laboratoire Universitaire de Biodiversité et Écologie Microbienne, F-29280 Plouzané, France

Correspondence: Jeanne Ropars, jeanne.ropars@u-psud.fr

Keywords: domestication, fungi, cheese, *Penicillium,* mycotoxin, horizontal gene transfer

Summary

Domestication involves recent adaptation under strong human selection and rapid diversification, and therefore constitutes a good model for studies of these processes. We studied the domestication of the emblematic white mold *Penicillium camemberti*, used for the maturation of soft cheeses, such as Camembert and Brie, about which surprisingly little was known, despite its economic and cultural importance. Whole genome-based analyses of genetic relationships and diversity revealed that an ancient domestication event led to the emergence of the gray-green *P. biforme* mold used in cheese-making, from the blue-green wild *P. fuscoglaucum* fungus. Another much more recent domestication event led to the generation of the *P. camemberti* clonal lineage from *P. biforme*. Both these domestication events were associated with strong bottlenecks. *Penicillium biforme* displayed signs of phenotypic adaptation to cheese-making relative to *P. fuscoglaucum*, in terms of its whiter color, faster growth on cheese medium under cave conditions, lower levels of toxin production and greater ability to prevent the growth of other fungi. The *P. camemberti* lineage displayed even stronger signs of domestication for all these phenotypic features. We also identified two differentiated *P. camemberti* lineages linked to different kinds of cheeses with contrasted phenotypic features in terms of color, growth, toxin production and competitive ability. We have, thus, identified footprints of domestication in these fungi, with genetic differentiation between cheese and wild strains and specific phenotypic traits beneficial for cheese-making. This study has not only fundamental implications for our understanding of domestication but can also have important impacts on cheese-making.

Introduction

Understanding how organisms adapt to their environment is a key issue in evolutionary biology, requiring investigations of population subdivision, and levels of genetic and phenotypic diversity or adaptive divergence. Domestication is a good model for studies of adaptive divergence, as it involves recent adaptation events affecting known traits under strong human selection and rapid diversification. Several studies on domesticated animals (e.g. horse, dog, pig[1–3]) and plants (e.g. maize, apricot[4,5]) have improved our understanding of adaptive divergence by revealing how adaptation and diversification into varieties occurred. Maize, for example, has undergone major changes in phenotype compared to its wild relative (teosinte), including a decrease in tillering, and the development of larger, non-dehiscent grains[6]. Several different maize varieties have been selected for different usages, with sugar-rich varieties grown for human consumption as kernels and field corn varieties grown for animal feed. Similarly, a number of different *Brassica* varieties have been selected for their leaves (cabbage and kale), stems (kohlrabi), flower shoots (broccoli and cauliflower) or buds (Brussels sprouts). Dog breeds have also diversified greatly under human selection, with different breeds displaying very different behaviors and phenotypes. Such notable adaptation is often associated with a decrease in fitness in natural environments, with, for example, a decrease in or loss of sexual reproduction ability in bulldogs[7] and bananas[8], smaller brains and less acute sense organs in most domesticated animals[9].

Fungi are excellent models for studying evolution and adaptation in eukaryotes, given their many experimental assets[10], including their small genomes and tractability for laboratory experiments. They can therefore be used to address questions in evolutionary biology through complementary approaches, including genomic and experimental methods. Humans have domesticated several fungi for the fermentation of foods (e.g. for beer, bread, wine, dried sausage and cheese), the production of secondary metabolites used in pharmaceutics (e.g. penicillin), and for their nutritional and gustatory qualities (e.g. button and shiitake mushrooms)[11]. Despite their economic and industrial importance, and their utility as biological models for studying adaptive divergence, fungi used by humans have been little studied, with the exception of the budding yeast *Saccharomyces cerevisiae* used for beer, wine and bread production[12–22], the filamentous fungus *Aspergillus oryzae* used to ferment soy and rice products in Asia[23–25] and the blue-cheese mold *Penicillium roqueforti*[26–28]. Whole-genome analyses have revealed that *P. roqueforti* has been domesticated twice, in two independent events, resulting in one population specific to the Roquefort protected designation of origin (PDO), the other population being used worldwide for all types of blue cheeses[26,29]. The Roquefort population displays some genetic diversity and had beneficial traits for cheese

production before the industrial era, growing more slowly in cheese and displaying greater spore

production on bread, the traditional multiplication medium. By contrast, the non-Roquefort cheese

population is a clonal lineage with traits beneficial for industrial cheese production, such as high

levels of lipolytic activity, efficient cheese cavity colonization and high salt tolerance. Some of these

beneficial traits have been conferred by two large horizontally transferred genomic regions, *Wallaby*

and *CheesyTer*[27,28], in the clonal non-Roquefort cheese population. The non-Roquefort cheese *P.*

roqueforti population grows more slowly in harsh conditions (minimal medium) than the Roquefort

cheese population[28]. Whole-genome analyses of *S. cerevisiae* have also revealed the existence of a

cheese-specific population of this yeast differentiated from the populations used for alcohol or

bread production. The cheese *S. cerevisiae* population assimilates galactose more rapidly than *S.*

cerevisiae populations thriving in other food environments (e.g. beer, bread, wine) or in natural

environments (oak)[30].

The white mold *Penicillium camemberti* is used for the maturation of soft cheeses, such as Camembert,

Brie and Neufchatel (Figure 1A-C). It is thought to be a white mutant selected from the gray-green

species *P. commune* for its color at the start of the 20th century[31], and cultured clonally ever since.

However, very little is known about its taxonomic status, origin and diversity, despite its great

economic and cultural importance. In particular, its relationships to the closely related species *P.*

biforme, P. caseifulvum, P. commune and P. *fuscoglaucum,* and even possible overlaps with these species,

remain unclear. *Penicillium camemberti* has only ever been found in the cheese/meat environment.

Penicillium commune is used for the maturation of other types of cheese (e.g. hard and blue cheeses)

and in the production of dried sausages, is commonly found as spoiler of dairy products[32] and is

also widespread in non-food environments. Genetic analyses however suggested that *P. commune*

was an invalid species and proposed to reinstate two ancient species, *P. biforme* (named by Thom in

1910) and *P. fuscoglaucum* (named by Biourge in 1923), based on a few genetic sequences separating

P. commune strains isolated from cheese (including the *P. biforme* type) from *P. commune* strains

isolated from other environments (including the *P. fuscoglaucum* type)[33]. However, the most recent

taxonomic reference study of the *Penicillium* genus did not recognize *P. fuscoglaucum* as a valid

species[34]. *Penicillium caseifulvum* has also sometimes been advocated to constitute a separate species

in this clade[34], including cheese strains isolated from Danish blue and other German and French

cheeses, on the basis of colony morphology and their lack of production of cyclopiazonic acid, a

mycotoxin produced by *P. camemberti*. However, a study based on a small number of genetic

markers was unable to differentiate these putative *P. caseifulvum* strains from *P. camemberti*[33]. Parts of

the horizontally transferred *CheesyTer* and *Wallaby* regions have also been found in the reference *P.*

camemberti and *P. biforme* genomes[27,28].

The taxonomy of this clade thus remains unclear, and population structure has not been studied

with powerful genetic markers, despite the paramount importance of genetic relationships to our

understanding of domestication. Differences in phenotypic traits have not been extensively studied

either, but are also essential for determining whether or not domestication has occurred. Humans

may, indeed, have selected strains for specific traits beneficial for cheese production, such as faster

growth in cheese, attractive color and a greater ability to compete against food-spoiling

microorganisms, but without the production of detrimental extrolites, called mycotoxins.

Mycotoxins are natural secondary metabolites produced by fungi under certain conditions that can

inhibit competitor growth and reproduction and which can have a toxicological impact on humans

depending on the exposure conditions[35].

We addressed these questions of genetic and phenotypic differentiation, by collecting and

sequencing the genomes (ca. 35 Mb) of 35 strains isolated from the crusts of different cheeses from

around the world (e.g. Camembert, tommes, blue cheeses) and 26 strains from other environments

(e.g. leaf litter, sausages, ice and wood). We performed a whole genome-based analysis of

population structure, which revealed that the three species *P. biforme, P. camemberti* and *P. fuscoglaucum*

formed separate and specific genetic clusters, thus confirming the division of *P. commune* into *P.*

biforme and *P. fuscoglaucum*. The species name *P. commune* should therefore be avoided. We found

that *P. camemberti* and *P. biforme* were sister species, both specific to the cheese environment, whereas

the more distantly related species *P. fuscoglaucum* was mostly found in natural environments. These

relationships suggest an ancient domestication event separating *P. biforme* from *P. fuscoglaucum* and

a much more recent domestication event generating the *P. camemberti* clonal lineage from *P. biforme.*

Consistent with this scenario, we found evidence of phenotypic adaptation to cheese-making in *P.*

biforme relative to *P. fuscoglaucum*, with a whiter color, faster growth on cheese medium under cave

conditions and lower levels of toxin production. We also identified two differentiated *P. camemberti*

lineages with contrasting phenotypic features. One of these *P. camemberti* lineages contained the

fluffy white strains isolated from Camembert or Brie including the *P. camemberti* type strain, which

grew more slowly in cave conditions than the other tested *Penicillium* strains, and produced similar

amounts of cyclopiazonic acid to *P. biforme*. This *P. camemberti* lineage carried the horizontally

transferred *CheesyTer* and *Wallaby* regions except three strains isolated from Camembert cheese

before 1905. The gray-green *P. camemberti* lineage was isolated from cheeses other than Camembert,

carries *CheesyTer* but not the entire *Wallaby* region, grew faster than the white *P. camemberti* lineage,

produced no cyclopiazonic acid, due to a frameshift mutation, and included the *P. caseifulvum* type

strain. Both *P. camemberti* lineages grew more slowly in harsh conditions than the other two clades,

suggesting that adaptation to cheese-making has led to a degeneration of functions useful in wild

environments. Both *P. camemberti* lineages excluded fungal competitors more effectively than *P.*
biforme, the fluffy white Camembert lineage being more effective at exclusion than the gray-green
P. camemberti lineage. This study thus reveals footprints of domestication in *P. camemberti* and *P.*
biforme, with genetic differentiation between cheese and wild strains, and the identification of
specific phenotypic traits beneficial for cheese-making. We also reveal the existence of
diversification into several varieties, also displaying genetic and phenotypic differentiation (in terms
of growth rate, color, competitive abilities and production of mycotoxin), used for the production
of different cheese types.

Results and Discussion

Penicillium camemberti, P. biforme and *P. fuscoglaucum* each form separate specific genetic clusters

We collected and sequenced 61 strains with Illumina® technology, including 35 strains isolated
from the rinds of various types of cheese (e.g., tommes, Camembert, blue cheeses), 9 from
sausages, and 17 from non-food environments (e.g., wood or leaf litter), which were attributed by
their collectors to the species *P. camemberti, P. biforme, P. commune* or *P. fuscoglaucum* (Table S1). We
resequenced the reference genome of *P. camemberti* (LCP06093, also known as FM013, and initially
sequenced with 454 technology[27]) with PacBio® long-read technology and used the sequence
obtained for mapping. We identified 392,072 SNPs across all strains (Supplemental Data 1). We
investigated population structure with two clustering methods based on genetic differences and
free from assumptions about mating system and mode of reproduction. Principal component
analysis (PCA; Figure 2A) and neighbor net (splitstree) analysis (Figure 2B) identified three genetic
clusters, corresponding to *P. camemberti* (*n*=19)*, P. biforme* (*n*=28) and *P. fuscoglaucum* (*n*=14). A
population structure analysis also identified the same three genetic clusters at *K*=3 (Figure 2C), the
K value at which the structure was the strongest and clearest.
The strains present in public collections under the name "*P. commune*" did not form a single cluster
nor even a monophyletic clade (black arrows in Figures 2A-C); indeed, the *P. commune* strains
isolated from natural environments clustered with *P. fuscoglaucum* whereas the *P. commune* strains
isolated from cheeses grouped with *P. biforme*, being genetically closer to *P. camemberti* than to *P.*
fuscoglaucum (Figures 2A-C). The *P. caseifulvum* type strain (LCP05630) and the *P. camemberti* type

strain belonged to different clusters at K values ≥ 5 in the population structure analysis (red stars
on Figure 2C).

Penicillium camemberti was found only in food (cheese, sausage, food waste or animal feed; Figure 2).
Only three of the 28 strains in the *P. biforme* cluster were isolated from environments other than
food: the atmosphere, ice and leaf litter. These three strains were not genetically differentiated from
those isolated from food (bread, cheese or sausage) and may therefore be feral strains, i.e. escaped
from food. The *P. fuscoglaucum* cluster included three strains isolated from food environments (two
strains from cheese and one from sausage) and 13 strains from natural environments and was thus
likely to constitute a genuine wild population. The three food strains clustered with the wild strains.

Penicillium fuscoglaucum displayed the highest degree of nucleotide diversity level ($\pi = 0.177$), with
about 196,000 SNPs. The *P. biforme* cluster had a higher genetic diversity ($\pi = 0.095$; 142,771 SNPs;
wider point dispersion in the PCA on Figure 2A and in the neighbor net on Figure 2B) than *P.*
camemberti ($\pi = 0.005$). The very low genetic diversity detected within *P. camemberti,* with only 0.02%
sites identified as polymorphic (i.e., only 8,180 SNPs), suggests that *P. camemberti* is a clonal lineage.

The long branches and the lack of cross-linking observed in the neighbor-net analysis further
confirmed the clonality of *P. camemberti*, contrasting with the footprints of recombination detected
within both *P. biforme* and *P. fuscoglaucum* (Figure 2B). In *P. fuscoglaucum,* abundant cross-linking was
observed, right to the branch edges (Figure 2B), reinforcing the view that *P. fuscoglaucum*
corresponded to a sexual population thriving in natural environments, as strains used in the food
industry are replicated clonally. We studied the mating-type genes controlling fungal mating
compatibility to determine the likelihood of sexual reproduction occurrence within each species.
In heterothallic fungal species*,* sexual reproduction occurs only between two haploid individuals
carrying different alleles (MAT1-1 and MAT1-2) at the mating-type locus. A lack of sexual
reproduction leads to relaxed selection on the mating-type genes, as they are no longer used, and a
departure from a balanced ratio between mating types. Neither mating-type allele presented any
evidence of loss-of-function mutations or SNPs in any of the genomes. In *P. fuscoglaucum,* we found
no significant departure from the 1:1 ratio of the two mating-type alleles expected under conditions
of regular sexual reproduction ($\chi_2 = 1.14$; df = 1; P = 0.29). By contrast, *P. biforme* displayed a
significant departure from the 1:1 ratio ($\chi_2 = 14.29$; df = 1; P < $1e_{-3}$), with MAT1-2 strains
predominating. All 19 *P. camemberti* strains carried the MAT1-2 allele, providing further evidence
of the clonality of *P. camemberti*[36,37]. The observed clonality in *P. camemberti* may be explained by the
recent selection of a white mutant and its use in industrial production[38], whereas *P. biforme* is more
widely used in the production of artisanal cheeses and farmers may make use of diverse local strains.

Two genetic clusters within *Penicillium camemberti*

The population structure analysis at $K = 5$ (Figure 2C), the PCA and the splitstree without *P. fuscoglaucum* (Figure S1) supported the existence of two genetic clusters within *P. camemberti*, separating strains isolated from soft cheeses, such as Camembert, and strains isolated from other kinds, such as Rigotte de Condrieu and Saint Marcellin. The *P. caseifulvum* type strain (red star on Figure 2C) clustered with *P. camemberti* strains isolated from non-soft cheeses. The genetic cluster including the *P. caseifulvum* type strain will be referred to hereafter as *P. caseifulvum,* and the genetic cluster including the *P. camemberti* type strain will be referred to as *P. camemberti sensu stricto*. A recent study described rapid phenotypic change from green-gray to white in a "wild *P. commune"* strain[39], interpreting these changes as evidence that domestication can occur within four weeks. However, we actually found here this *P. commune* strain to belong to the cheese *P. biforme* clade and therefore to an already domesticated clade (Figure S2).

The population structure analysis (Figure 2C) also suggested further genetic subdivision within each of *P. fuscoglaucum* and *P. biforme*; these subdivisions did not correspond to the environment from which the isolate was obtained, type of cheese, or any other variable for which we had data; it did not correspond to either a strong subdivision in the PCA or the neighbor-net (Figures 2 and S1).

Presence/absence polymorphism of the two horizontally transferred regions likely to be advantageous in cheese

By contrast to the first 454-sequenced FM013 reference genome[27], in which the *Wallaby* horizontally transferred region was found to be fragmented, the obtained PacBio genome assembly of the same strain showed that the part of *Wallaby* present in the genome formed a single block (scaffold 18, positions 102,711 to 538,265). Two genes in *Wallaby* have been highlighted as putatively important for cheese-making. One, encoding the *Penicillium* antifungal protein (PAF)[27], was absent from FM013, whereas the other, encoding Hce2, a protein with known antimicrobial activities, was present. All *P. camemberti sensu stricto* strains carried the same *Wallaby* region as FM013, whereas strains from the *P. caseifulvum* lineage carried fragmented and duplicated segments of the *Wallaby* fragment present in FM013, in which the Hce2 gene was systematically present. We also found *Wallaby* fragmented with some duplicated regions in 18 *P. biforme* strains; it was completely absent from the 10 remaining strains (Figure 3 and Figure S3). *Wallaby* appeared to be even more fragmented in *P. fuscoglaucum* strains, always lacking Hce2 but in one strain, and it was completely absent in nine strains.

The other horizontally transferred region thought to be beneficial for growth on cheese[28], the 57
kb *CheesyTer* region, was not located at the very end of a scaffold in *P. camemberti* (scaffold 17,
positions 690,636 to 751,978 in FM013), like its position in *P. roqueforti*. We found no strict
association between *CheesyTer* and *Wallaby* in the clades studied here, whereas all strains of *P.*
roqueforti studied so far carried either both or neither of these regions[28]. In *P. camemberti*, 16 of 19
strains carried the whole *CheesyTer* region, with an identity of 100% (Figures 3 and S4). The other
three *P. camemberti* strains (LCP00584T, LCP05527 and UBOCC-A-108096), belonging to the *P.*
camemberti s.s. cluster, completely lacked *CheesyTer* and an 80 kb-downstream fragment. These three
strains were isolated from Camembert cheeses around 1905, suggesting that the horizontal transfer
of *CheesyTer* to *P. camemberti* occurred after 1905, in a single strain that was then clonally cultured.
In *P. biforme*, 25 of 28 strains carried *CheesyTer*, with an identity >99% between strains. The identity
with *P. camemberti s.l.* was also >99%, except in the 5 kb region at the start of *CheesyTer*, in which
the similarity between the two species dropped to 90%. The 26 kb between *CheesyTer* and the end
of scaffold 17 in *P. camemberti s.l.* also displayed lower similarity (around 90%) between *P. camemberti*
s.l. and *P. biforme*, whereas it was identical within species. The differences between *P. camemberti* and
P. biforme in the flanking regions and at the start of *CheesyTer* appeared to be mainly due to C:G to
T:A mutations in *P. camemberti*, corresponding to the repeat-induced point (RIP) mutations specific
to fungi occurring in repeated sequences during sexual reproduction[40] (Figure S5). As *P. camemberti*
s.l. was found to be a clonal lineage and no RIP footprints were detected elsewhere in the genome
of FM013, these findings suggest that *CheesyTer* was transferred to *P. camemberti s.l.* with these RIPed
sequences already present. *CheesyTer* was completely absent from six strains of *P. fuscoglaucum*
(Figure 3) and partially present in the other strains. One *P. fuscoglaucum* strain (LCP00218, isolated
from rubber) carried two copies of a 30 kb *CheesyTer* terminal region. The two *CheesyTer* genes
involved in lactose metabolism[28], one encoding a lactose permease and the other a β-galactosidase,
were present in all strains carrying at least one part of *CheesyTer* in *P. camemberti*, *P. biforme* and *P.*
fuscoglaucum (Figure 3). *Penicillium biforme*, *P. fuscoglaucum* and *P. roqueforti* had no RIP footprints in the
CheesyTer region for which RIP mutations were detected in *P. camemberti*, so it was not possible to
identify the species or strain from which *CheesyTer* was transferred to *P. camemberti*.

Better phenotypes for cheese production in *P. camemberti s.l.* and *P. biforme*
than in the closely related wild species *P. fuscoglaucum*

Strains selected by humans would be expected to display specific traits beneficial for cheese
production, such as faster growth in cheese, colonies of an attractive color, salt tolerance, an
absence of harmful toxin production and efficiency at excluding undesirable microorganisms. We

performed a set of laboratory experiments to determine whether cheese strains had evolved traits beneficial for cheese production not present in the closely related wild *P. fuscoglaucum* lineage occurring in other environments.

We first investigated whether *P. camemberti s.l.* and *P. biforme* had acquired traits enabling them to grow more rapidly than *P. fuscoglaucum* in cheese-making conditions. Rapidly growing fungi can be beneficial for cheese-making, as they are more capable of excluding spoiler bacteria, yeasts and molds[11] and promote faster cheese maturation. The process of cheese maturation begins with the transfer of salted milk curd to dark caves at a low temperature and high humidity. We therefore grew 61 strains on various media (unsalted cheese, salted cheese, minimal and malt media), in the dark and at different temperatures and humidity levels (25°C at ambient relative humidity, 10°C at 98% humidity or 8°C at 100% humidity), and we measured colony radial growth after 10 days. We found no significant effect on growth of the substrate from which the isolate was originally obtained (i.e. food versus non-food). We found a significant effect on growth of the culture media, species, temperature/humidity conditions, lineage within *P. camemberti s.l.* and the interaction between culture medium and species (Table S2, Figure 4A). The significance of the interaction between culture medium and temperature/humidity conditions (Table S2) reflected the very slow growth at low temperatures, making it difficult to detect significant differences between media. The two *P. camemberti s.l.* genetic clusters were genetically very closely related, but the *P. caseifulvum* lineage grew similarly to *P. biforme* under Camembert cave conditions (i.e. 10°C at 98% humidity, post-hoc Tukey honestly significant difference - HSD test $P = 1$), with greater radial growth on unsalted cheese and salted cheese, and slower growth on malt and minimal media than *P. fuscoglaucum*. By contrast, the *P. camemberti s.s.* lineage displayed weaker radial growth than any other lineage on all media, but was fluffier. Furthermore, *P. camemberti s.l.* and *P. biforme* grew less rapidly than *P. fuscoglaucum* on minimal and malt media, indicating a disadvantage for growth in harsh conditions, as expected for cheese strains, due to relaxed selection on functions useful only in wild environments. These findings demonstrated that *P. caseifulvum* and *P. biforme* had acquired growth traits that were beneficial under cheese maturation conditions. In addition, the two *P. camemberti s.l.* lineages displayed opposite growth patterns on unsalted and salted cheese, with strains used for making Camembert or Brie (*P. camemberti s.s.*) growing much less radially on cheese than any other strains, but growing much more vertically (i.e. being fluffier), with the mycelium growing up to the lid of the Petri dishes (Figure 1C).

High salt concentrations in cheeses prevent the growth of contaminants and cheese fungi may have

adapted to such conditions. We found an effect of salt on *P. biforme* growth, with strains growing

more rapidly on salted than unsalted cheese medium (post-hoc Tukey HSD test $P = 0.01$) and

more rapidly on salted cheese medium than *P. camemberti s.l.* (post-hoc Tukey HSD test $P = 0.007$).

Salt had no effect on the growth of *P. camemberti s.l.* (post-hoc test $P = 1$) or *P. fuscoglaucum* (post-

hoc test $P = 1$). There may have been stronger selection for salt tolerance in *P. biforme,* which is

used to inoculate blue and goat cheeses, both being more salty than soft cheeses, such as Brie or

Camembert[41], for which *P. camemberti s.s.* is used.

We investigated whether cheese lineages had become whiter, which can be more attractive to some

consumers than gray-green mold on cheese, by comparing the opacity of lineages on cheese

medium, as opacity increases with the brightness and fluffiness of a colony. We found significant

effects of species and lineage on color within *P. camemberti s.l.* (Figure 4C and Table S2)*,* with *P.*

camemberti s.l. being significantly more opaque (i.e. brighter and fluffier) and *P. camemberti s.s.* even

more so, than *P. caseifulvum.* This is consistent with the white and fluffy aspect of the crust of

Camembert and Brie, made with *P. camemberti s.s.*, whereas *P. caseifulvum* is found in cheeses with a

grayer and less fluffy crust, such as Saint Marcellin or Rigotte de Condrieu (Figures 1 and 4B-C).

The *P. caseifulvum* and *P. biforme* lineages did not differ significantly from each other (post-hoc Tukey

HSD test $P = 1$) and both were brighter than the wild blue-green *P. fuscoglaucum* (Figure 4B).

We also investigated whether cheese lineages produced smaller amounts of cyclopiazonic acid

(CPA), which is cytotoxic to humans[35,42], than strains isolated from other environments. *Penicillium*

camemberti has been reported to produce CPA on yeast extract sucrose (YES) medium and at very

low, non-toxic concentrations in cheese[43]. None of the *P. caseifulvum* strains tested here produced

CPA on YES (Figure 4D), consistent with findings for the type strain[44]. By contrast, *P. camemberti*

s.s., P. biforme and *P. fuscoglaucum* produced CPA, the highest levels being obtained with *P. biforme*

and *P. fuscoglaucum.* The CPA biosynthesis cluster in the genome appeared to be functional, with six

genes present (*cpaA*, *cpaD*, *cpaO*, *cpaH*, *cpaM* and *cpaT*) but not the *cpaR* gene encoding a regulatory

protein (Figure S6). The only exceptions were the six *P. caseifulvum* strains, in which a 2 bp deletion

in the *cpaA* gene led to a frameshift. The *cpaA* gene encodes a polyketide synthase/non-ribosomal

peptide synthase responsible for the first step of the CPA biosynthetic pathway so that a non-

functional protein probably prevents CPA production on all substrates. Humans have often

selected fungal strains unable to produce harmful toxins for use in food, as exemplified by the

Aspergillus oryzae strains used to ferment Asian food products, which do not produce aflatoxins,

whereas the wild ancestor, *A. flavus,* does[45]. In the blue cheese fungus *P. roqueforti*, strains belonging

to the non-Roquefort population were also found unable to produce mycophenolic acid due to a

174 bp deletion in the *mpaC* gene[46].

Cheese is a nutrient-rich environment in which many microorganisms can thrive, including

undesirable food spoilage organisms. We therefore investigated whether the cheese lineages were

better able to outcompete challengers. We allowed strains of *P. camemberti s.s.* (*n*=3), *P. caseifulvum*

(*n*=2) or *P. biforme* (*n*=10) to grow as lawns on the surface of the cheese medium, which were

inoculated 24 h later with a single spot, in the middle of the Petri dish, using a competitor:

Geotrichum candidum (*n*=5, two strains isolated from cheese and three from other environments)*, P.*

biforme (*n*=6, three strains isolated from cheese and three from other environments)*, P. fuscoglaucum*

(*n*=5, two strains isolated from cheese and three from other environments)*,* or *P roqueforti* (*n*=12,

three strains from each of the four known genetic clusters[26]). The species and the lineages of *P.*

camemberti s.l. used as the lawn had significant effects on the growth of the challenger (Table S2,

Figure 5). The two lineages of *P. camemberti s.l.* prevented the growth of all challengers more

effectively than *P. biforme*, and *P. camemberti s.s.* was even more effective. Furthermore, we observed

significant differences between the species used as challengers, with *Geotrichum candidum* having the

highest growth differential on the various lawns, growing less well on a *P. camemberti s.s.* lawn than

other lawns. *Geotrichum candidum* is a fungus that is also present on the surface of soft cheeses. The

exclusion effect may be mediated by the fluffy morphology, resulting in the fungus occupying more

space and using more resources, and/or by biochemical interactions.

Conclusion

Population structure analyses based on whole-genome sequences revealed that the three species,

P. biforme, P. camemberti s.l. and *P. fuscoglaucum,* each formed a specific genetic cluster, confirming the

separation of *P. commune* into *P. biforme* and *P. fuscoglaucum; P. commune* should therefore be

considered as an invalid species name and should not be used anymore as considering all *"P.*

commune" strains to be wild leads to wrong inferences about trait evolution[39]. We found that *P.*

camemberti s.l. and *P. biforme* were sister clades specific to the cheese environment, whereas the more

distantly related species *P. fuscoglaucum* was found mostly in natural environments. These

relationships and the lower diversity of *P. biforme* than *P. fuscoglaucum,* with even lower levels of

diversity in *P. camemberti s.l.*, suggest that an ancient domestication event led to the generation of *P.*

biforme from *P. fuscoglaucum* and that a much more recent domestication event led to *P. camemberti*

s.l. emerging from *P. biforme*. Domestication often leads to bottlenecks[47–50], which can be very severe

in clonally multiplied fungi[26,51]. Consistent with this scenario, we found evidence of phenotypic

adaptation to cheese-making in *P. biforme* relative to the wild *P. fuscoglaucum* species, with a whiter

color, faster growth on cheese medium under cave conditions and lower levels of toxin production.

These signs of domestication were even more marked in *P. camemberti s.l.* than in *P. biforme.*

We observed a similar evolution of traits as in populations of the domesticated blue-cheese fungus

P. roqueforti, which grows more rapidly on cheese medium, is more competitive against contaminant

microorganisms, and grows less well under harsh conditions than wild strains[26,28]. Such convergent

evolution under similar selective pressures suggests that evolution may repeat itself, as already

reported in natural populations of *Anolis* lizards[52,53], three-spine sticklebacks[54], Mexican

cavefishes[55], and cichlid fishes in African rifts[56] or crater lakes in central America[57,58]. Phenotypic

convergent evolution has been also reported in domesticated organisms, particularly in crop plants,

for the loss of seed shattering, minimization of seed dormancy and increase in seed size and

number, for example, with most of these changes resulting from different genomic alterations in

different species[59,60].

We identified two genetically differentiated *P. camemberti* varieties, *P. camemberti s.s.* and *P. caseifulvum,*

with contrasting phenotypic features, used in the production of different kinds of cheese. The *P.*

camemberti s.s. strains were white and were isolated from Camembert or Brie. Their radial growth

was slower but their mycelia were much fluffier than the other *Penicillium* strains tested (Figure 1C)

and they produced similar amounts of CPA compared to *P. biforme* strains. They excluded fungal

competitors effectively, as previously suggested[61], probably due to their fluffy morphology, taking

up the available space and monopolizing resources. *Penicillium caseifulvum* strains were gray-green,

unable to produce CPA, and were isolated from cheeses other than Camembert, such as St

Marcellin or Rigotte de Condrieu (Figure 1A). They displayed more rapid radial growth than *P.*

camemberti s.s. and grew similarly to *P. biforme* strains on cheese medium in cave conditions. The

existence of two genetically and phenotypically differentiated lineages with different uses in cheese-

making suggested that these lineages emerged as a result of different human selection pressures, as

reported for the domesticated blue-cheese fungus *P. roqueforti*[26], wine-making yeasts[12,15], maize, rice,

tomatoes, dogs, chickens and horses[1–6].

Penicillium camemberti s.s. is the emblematic species used to inoculate soft cheeses, such as

Camembert and Brie. According to the technical specifications for Brie de Meaux and Brie de

Melun PDOs and Camembert, the crust of the cheese must be white and fluffy, and inoculation

with *P. candidum* (a synonym of *P. camemberti s.l.*) is even specified for Brie de Meaux. These specifications are recent (20th century) and seem to have had a negative impact on the diversity of the fungi used for making these kinds of cheeses, as a single clonal lineage now predominates. The genetic diversity of *P. roqueforti* has also been greatly reduced by recent industrialization[26,29], although PDO specifications that local strains must be used have protected diversity to some extent in Roquefort cheeses. Camembert and Brie cheeses were gray-green before the 20th century[38], as illustrated by a 19th century painting by Marie Jules Justin entitled "Symphonie des fromages en Brie majeur - Nature morte au fromage". These historical records are consistent with our findings, which suggest a first domestication event leading to the emergence of the gray-green mold *P. biforme,* subsequently followed, around 1900, by the domestication of *P. camemberti* from *P. biforme,* with the selection of two different varieties displaying more marked signs of domestication than *P. biforme.*

These findings have industrial implications, as they reveal the existence of three closely related but different lineages that have evolved traits beneficial for cheese-making, with different phenotypic traits selected according to usage. This study should foster further research, as it would be interesting to assess other important traits, such as volatile compound production, and the efficiencies of lipolysis and proteolysis. Our findings raise questions about the use of limited numbers of clonal strains for cheese-making, which tends to lead to degeneration, limiting the possibilities for further improvement, which is currently a major concern in the agrofood sector[62], despite the great geneticist Vavilov known for having identified the centers of origin of cultivated plants long ago highlighting the importance of genetic diversity in domesticated organisms for variety improvement and diversification[63].

Materials and Methods

Strain collection and DNA extraction

We analyzed strains isolated from 34 cheeses from five countries around the world (e.g. Camembert, Saint Marcellin, tomme, Sein de Nounou). We also collected strains from dried sausages and moldy food (e.g. bread). Spores were sampled from the food sources and spread on Petri dishes containing malt-agar medium, which were then incubated for five days at 25°C. A total of 42 strains were obtained from public strain collections (the Laboratoire de Cryptogamie (LCP strains) at the National Museum of Natural History in Paris (France), the Laboratoire Universitaire de Biodiversité et Ecologie Microbienne (LUBEM, UBOCC strains) in Plouzané (France) and the

Westerdijk Fungal Biodiversity Institute (CBS strains) in Utrecht (The Netherlands)). We obtained

strains from natural environments (e.g. wood or natural cave walls) from nine different countries.

Detailed information about the strains used in this study can be found in the Table S1. For each

strain, single-spore cultures were generated by a dilution method, to ensure that only a single

haploid genotype was cultured for each strain. We checked the species identification of all strains

by Sanger sequencing of the β-tubulin gene (primers Bt2a -

GGTAACCAAATCGGTGCTGCTTTC and bt2b - AACCTCAGTGTAGTGACCCTTGGC[64])

and the PC4 microsatellite flanking regions[33] (primers PC4F – CAAGCTGGCCGATAACCTG

and PC4R – CCATCCGCTTGATTTCTCCT), to distinguish between *P. biforme* and *P. camemberti*

s. l. All the strains were added to the ESE (Ecology Systematics Evolution Laboratory) collection,

under an ESE accession number, and are available from the corresponding collections. For each

strain, we used the Nucleospin Soil Kit (Machernary-Nagel, Düren, Germany) to extract DNA

from fresh haploid mycelium grown for five days on malt agar.

Genome sequencing, assembly and mapping

Sequencing was performed with Illumina HiSeq 2500 paired-end technology (Illumina Inc.), with

a mean insert size of 400 bp, at the INRA GenoToul platform, to obtain 10-50X coverage (Table

S1). In addition, the genome of the *P. camemberti* LCP06093 (also known as FM013) reference

strain, initially sequenced with 454 technology[27], was resequenced at 34x using the PacBio's long-

read technology, assembled with Canu 1.8[65] and polished with pilon 1.23[66]. The PacBio assembly

and reads for all samples have been deposited at the NCBI Sequence Read Archive under

BioProject ID XXXX.

Reads were trimmed with Trimmomatic[67] (cut adapter, options PE, LEADING:3 TRAILING:3

SLIDINGWINDOW:4:25 MINLEN:36) and mapped onto the high-quality PacBio reference

genome of LCP06093 with Bowtie2[68] (options very-sensitive-local, --phred33 PL:ILLUMINA -X

1000). Mapped reads were filtered for PCR duplication with picard tools

(http://broadinstitute.github.io/picard) and realigned on the PacBio reference genome with

GATK[69]. Single-nucleotide polymorphisms (SNPs) were called with GATK HaplotypeCaller,

which provides one gVCF per strain (option -ERC GVCF). GVCFs were combined with GATK

CombineGVCFs, genotyped with GATK genotype and hard filtered in accordance with GATK

best practice workflow recommendations (https://software.broadinstitute.org/gatk/best-

practices/).

Statistics of population genetics

469   Nucleotide diversity (π) was calculated with the R package pegas[70] for the three species *P. biforme,*

470   *P. fuscoglaucum* and *P. camemberti s.l.*. We also calculated π considering *P. camemberti s.s.* and *P.*

471   *caseifulvum* as separate genetic clusters.

Genetic structure

474   We used the dataset for 392,072 SNPs to infer a finer population structure. We inferred individual

475   ancestry from genotype likelihoods based on realigned reads, by assuming a known number of

476   admixing populations, ranging from $K = 2$ to $K = 6$, using NgsAdmix from the ANGSD package[71].

477   Neighbor-joining trees were generated with the R package ape[72]. We used the R package phangorn[73]

478   for neighbor-net analyses, and the prcomp function of R for principal component analysis (PCA).

Laboratory experiments

Sampling and calibration

All experiments were performed on the whole collection of *P. camemberti s.l.* (*n* = 20)*, P. biforme* (*n*

= 28) and *P. fuscoglaucum* (*n* = 13) strains, including 34 strains isolated from cheeses, 8 from dried

sausages and 19 strains isolated from environments other than food. Experiments were initiated

with spore suspensions calibrated to 10^7 spores/mL with a hemocytometer, under the constraint

of the low rate of sporulation in *P. camemberti s.l.*.

Growth in different conditions and on different media

We investigated whether strains isolated from cheese displayed faster or slower radial growth than

strains isolated from other environments when cultured on cheese or other substrates, and under

conditions similar to those in maturation caves or other conditions. We prepared four different

culture media: a cheese medium without salt, a salted cheese medium (17g/L, corresponding to a

typical cheese), a malt medium and a minimal medium. The cheese media were produced from an

unsalted drained cow's milk cheese from Coubertin Farm in Saint Rémy-les-Chevreuse (France),

as previously described[61]; we added five drops of blue food coloring to these media, to make it

easier to distinguish white fungal colonies from the medium. The cheese media were rich in lipids

and proteins, whereas the malt medium (20 g/L) was rich in carbohydrates. The minimal medium[74]

contained only the trace elements necessary for fungal survival (i.e. iron sulfate, zinc sulfate, boric

acid, magnesium chloride, copper sulfate, ammonium heptamolybdate, cobalt chloride, EDTA,

sodium nitrate, potassium chloride, potassium phosphate and magnesium sulfate). All media were

sterilized in an autoclave (121°C for 20 min for the malt and minimal media, 110°C for 15 min for

the cheese media). Each 90 mm-diameter Petri dish was filled with 20 mL of the appropriate medium.

We allowed the fungi to grow in the dark, under three different conditions for each strain and each medium: 10°C with 98% humidity (Camembert cave conditions), 8°C with 85% humidity (cave conditions for other cheeses) and 25°C with ambient relative humidity (ambient conditions). We had 244 Petri dishes in total for each set of conditions, and we left the colonies to grow for 10 days. Images of the Petri dishes were obtained with a Scan 1200 from Interscience and analyzed with IRIS[75] for growth and color-opacity measurements.

Mycotoxin production

For measurements of mycotoxin production, we used 1 μL of calibrated spore suspension (10^7 spores/mL) for each of the 61 strains to inoculate YES (yeast extract sucrose) agar medium buffered at pH 4.5 with phosphate-citrate buffer and characterized by a high C/N ratio to favor mycotoxin production as already described[46]. Each culture was performed in triplicate for myctoxin analyses as well as for fungal dry weight measurements. The plates were incubated at 25°C in the dark for 10 days and were then stored at -20°C until mycotoxin analysis.

For mycotoxin extractions[46], we homogenized the thawed samples with an Ultraturrax T25 (IKA, Heidelberg, Germany) before recuperating 4g aliquots. Then, 25 mL of acetonitrile (ACN) supplemented with 0.1% formic acid (v/v) was added, samples were vortexed for 30 sec followed by 15 min sonication. Extracts were then centrifuged for 10 min at 5000g and the recuperated supernatants were directly filtered through 0.2 μm PTFE membrane filters (GE Healthcare Life Sciences, UK) into amber vials. All samples were stored at -20°C until analyses.

Mycotoxin detection and quantification were performed using an Agilent 6530 Accurate-Mass Quadropole Time-of-Flight (Q-TOF) LC/MS system equipped with a Binary pump 1260 and degasser, well plate autosampler set to 10°C and a thermostated column compartment. Filtered samples (2μl) were injected into a ZORBAX Extend C-18 column (2.1x50mm and 1.8 μm, 600 bar) maintained at 35°C with a flow rate of 0.3 ml/min using mobile phase A (milli-Q water + 0.1% formic acid (v/v) and 0.1% ammonium formate (v/v) and mobile phase B (ACN + 0.1% formic acid). Mobile phase B was maintained at 10% for 4 min followed by a gradient from 10 to 100% for 16 min. Then, mobile phase B was maintained at 100% for 2 min before a 5 min post-time. Cyclopiazonic acid (CPA) was ionized in electrospray ionization mode ESI+ in the mass spectrometer with the following parameters: capillary voltage 4 kV, source temperature 325°C, nebulizer pressure 50 psig, drying gas 12 l/min, ion range 100-1000 m/z. CPA had a retention time

of 15.3 min and was quantified using the $[M+H]_+$ 337.1545 m/z ion and $[M+Na]_+$ 359.1366

qualifier ion (ESI+). Other extrolites included in our internal database that were targeted (in ESI+

and ESI- modes) were andrastin A, citreoviridin, citrinin, ermefortins A & B, (iso)-fumigaclavin A,

griseofulvin, meleagrin, mycophenolic acid, ochratoxin A, patulin, penitrem A, PR toxin,

roquefortin C, sterigmatocystin.

We used a matrix matched calibration curve (R_2 >0.99) for reliable mycotoxin quantification with

final concentrations ranging from 10 to 10000 ng/ml. Method performance and mycotoxin

determination was carried out as previously described[46]. Specific mycotoxin production was

expressed as ng per g fungal dry weight.

Competition

We inoculated salted cheese with 150 μL of a calibrated spore solution (10_7 spores/mL), which we

allowed to grow as a lawn. We used two *P. caseifulvum* strains (LCP05630 and ESE00019), three *P.*

camemberti s.s. strains (FM013, LCP04810 and LCP05527), five *P. biforme* strains isolated from cheese

(ESE00018, ESE00086, ESE00089, ESE00126, ESE00139) and five *P. biforme* strains isolated from

other environments (LCP05531, LCP06701, ESE00154, LCP06620 and LCP05496). After 24 h of

growth, we deposited a single 20 μL drop of calibrated spore solution from a challenger species

(10_7 spores/mL) in the middle of the Petri dish. The challenger species used were *G. candidum* (*n*=4,

ESE00163 and ESE00165 isolated from cheese, CBS11628 and VTTC4559 from other

environments)*, P. biforme* (*n*=6, three isolated from cheese -ESE00126, ESE00089, ESE00018- and

three from other environments -ESE00154, LCP005531, LCP06701-)*, P. fuscoglaucum* (*n*=5, two

isolated from cheese -ESE00090, LCP06621- and three from other environments -LCP000797,

LCP04799, LCP03239)*, and P roqueforti* (*n*=12, three from each of the four genetic clusters

described in [26]: LCP06036, LCP06037, LCP06128, LCP06043, LCP06098, LCP06064, FM164,

LCP06137, LCP06131, LCP06173).

Statistical tests

All data were normalized with the bestNormalize R package[76]. Analysis of variance (ANOVA) was

performed, followed by post-ANOVA tests such as Tukey's HSD (honest significant difference)

test, performed with R.

We used the following normalizations for the various phenotypic traits tested. Ordered quantile

(ORQ) normalization, a one-to-one transformation for vectors with an arbitrary distribution that

generates normally distributed vectors, was applied to diameter measurements and competition

data. Yeo-Johnson transformation was used to obtain uniform distributions for color opacity and
mycotoxin data.
572

Acknowledgments

574 We thank everyone who sent us cheese crusts, Steve Labrie for the LMA strain, and Joëlle Dupont
575 for strains from the National Museum of Natural History in Paris.
576 This work was supported by the Fungadapt ANR-19-CE20-0002-02 ANR grant to TG, JR, EC,
577 MC and a Fondation Louis D. grant from the Institut de France to TG.

Author contributions

580 JR and TG designed and supervised the study, and obtained funding. JR, ED, AS and SLP
581 generated the data. JR, RRdlV and BB analyzed the genomes. JR, ED and AS performed the
582 experiments. MC, EC and EP performed the mycotoxin and contributed to CPA biosynthetic gene
583 cluster analyses. JR and ED analyzed the data from the laboratory experiments. JR and TG wrote
584 the manuscript, with contributions from all the authors.
585

References

1. Warmuth, V. *et al.* European Domestic Horses Originated in Two Holocene Refugia. *PLoS*
ONE **6**, e18194 (2011).
2. Axelsson, E. *et al.* The genomic signature of dog domestication reveals adaptation to a starch-
rich diet. *Nature* **495**, 360–364 (2013).
3. Frantz, L. A. F. *et al.* Evidence of long-term gene flow and selection during domestication
from analyses of Eurasian wild and domestic pig genomes. *Nat Genet* **47**, 1141–1148 (2015).
4. Hufford, M. B. *et al.* Comparative population genomics of maize domestication and
improvement. *Nat Genet* **44**, 808–811 (2012).
5. Decroocq, S. *et al.* New insights into the history of domesticated and wild apricots and its
contribution to Plum pox virus resistance. *Mol. Ecol.* **25**, 4712–4729 (2016).
6. Doebley, J. The Genetics of Maize Evolution. *Annual Review of Genetics* **38**, 37–59 (2004).
7. Wydooghe, E., Berghmans, E., Rijsselaere, T. & Soom, A. V. International breeder inquiry
into the reproduction of the English bulldog. *Vlaams Diergeneeskundig Tijdschrift* **82**, 38–43
(2013).
8. Perrier, X. *et al.* Multidisciplinary perspectives on banana (*Musa* spp.) domestication. *PNAS*
108, 11311–11318 (2011).
9. Diamond, J. Evolution, consequences and future of plant and animal domestication. *Nature*
418, 700–707 (2002).
10. Gladieux, P. *et al.* Fungal evolutionary genomics provides insight into the mechanisms of
adaptive divergence in eukaryotes. *Molecular Ecology* **23**, 753–773 (2014).
11. Dupont, J. *et al.* Fungi as a Source of Food. in *Microbiology Spectrum* vol. 5 (2017).
12. Almeida, P. *et al.* A Gondwanan imprint on global diversity and domestication of wine and
cider yeast *Saccharomyces uvarum*. *Nature Communications* **5**, (2014).
13. Fay, J. C. & Benavides, J. A. Evidence for domesticated and wild populations of *Saccharomyces*

cerevisiae. PLoS Genet **1**, e5 (2005).

14. Gallone, B. *et al.* Domestication and divergence of *Saccharomyces cerevisiae* beer yeasts. *Cell* **166**,
1397-1410.e16 (2016).

15. Legras, J.-L., Merdinoglu, D., Cornuet, J.-M. & Karst, F. Bread, beer and wine: *Saccharomyces*
cerevisiae diversity reflects human history. *Molecular Ecology* **16**, 2091–2102 (2007).

16. Libkind, D. *et al.* Microbe domestication and the identification of the wild genetic stock of
lager-brewing yeast. *PNAS* **108**, 14539–14544 (2011).

17. Marsit, S. *et al.* Evolutionary advantage conferred by an eukaryote-to-eukaryote gene transfer
event in wine yeasts. *Molecular Biology and Evolution* **32**, 1695–1707 (2015).

18. Novo, M. *et al.* Eukaryote-to-eukaryote gene transfer events revealed by the genome sequence
of the wine yeast *Saccharomyces cerevisiae* EC1118. *PNAS* **106**, 16333–16338 (2009).

19. Peter, J. *et al.* Genome evolution across 1,011 *Saccharomyces cerevisiae* isolates. *Nature* **556**, 339–
344 (2018).

20. Liti, G. *et al.* Population genomics of domestic and wild yeasts. *Nature* **458**, 337–341 (2009).

21. Spor, A. *et al.* Phenotypic and genotypic convergences are influenced by historical contingency
and environment in yeast. *Evolution* **68**, 772–790 (2014).

22. Dunn, B. *et al.* Recurrent rearrangement during adaptive evolution in an interspecific yeast
hybrid suggests a model for rapid introgression. *PLoS Genetics* **9**, e1003366 (2013).

23. Galagan, J. E. *et al.* Sequencing of *Aspergillus nidulans* and comparative analysis with *A. fumigatus*
and *A. oryzae. Nature* **438**, 1105–1115 (2005).

24. Gibbons, J. G. *et al.* The evolutionary imprint of domestication on genome variation and
function of the filamentous fungus *Aspergillus oryzae. Current Biology* **22**, 1403–1409 (2012).

25. Machida, M. *et al.* Genome sequencing and analysis of *Aspergillus oryzae. Nature* **438**, 1157–1161
(2005).

26. Dumas, E. *et al.* Independent domestication events in the blue-cheese fungus *Penicillium*
roqueforti. Mol Ecol mec.15359 (2020) doi:10.1111/mec.15359.

27. Cheeseman, K. *et al.* Multiple recent horizontal transfers of a large genomic region in cheese
making fungi. *Nat Commun* **5**, 2876 (2014).

28. Ropars, J. *et al.* Adaptive horizontal gene transfers between multiple cheese-associated Fungi.
Current Biology **25**, 2562–2569 (2015).

29. Gillot, G. *et al.* Insights into *Penicillium roqueforti* morphological and genetic diversity. *PLoS One*
10, e0129849 (2015).

30. Legras, J.-L. *et al.* Adaptation of *S. cerevisiae* to fermented food environments reveals remarkable
genome plasticity and the footprints of domestication. *Mol Biol Evol* **35**, 1712–1727 (2018).

31. Pitt, J. I. *The genus Penicillium and its teleomorphic states Eupenicillium and Talaromyces.* (Academic
Press, 1979).

32. Garnier, L. *et al.* Diversity of spoilage fungi associated with various French dairy products.
International Journal of Food Microbiology **241**, 191–197 (2017).

33. Giraud, F. *et al.* Microsatellite loci to recognize species for the cheese starter and contaminating
strains associated with cheese manufacturing. *International Journal of Food Microbiology* **137**, 204–
213 (2010).

34. Visagie, C. M. *et al.* Identification and nomenclature of the genus *Penicillium. Studies in Mycology*
78, 343–371 (2014).

35. Hymery, N. *et al.* Filamentous Fungi and Mycotoxins in Cheese: A Review. *Comprehensive*
Reviews in Food Science and Food Safety **13**, 437–456 (2014).

36. Dyer, P. S. & O'Gorman, C. M. Sexual development and cryptic sexuality in fungi: insights
from Aspergillus species. *FEMS Microbiol. Rev.* **36**, 165–192 (2012).

37. Barve, M. P., Arie, T., Salimath, S. S., Muehlbauer, F. J. & Peever, T. L. Cloning and
characterization of the mating type (MAT) locus from *Ascochyta rabiei* (teleomorph: *Didymella*
rabiei) and a MAT phylogeny of legume-associated *Ascochyta* spp. *Fungal Genetics and Biology* **39**,
151–167 (2003).

38. Boisard, P. *Le camembert, mythe français.* (2007).

39. Bodinaku, I. *et al.* Rapid Phenotypic and Metabolomic Domestication of Wild *Penicillium* Molds on Cheese. *mBio* **10**, e02445-19 (2019).

40. Selker, E. U., Cambareri, E. B., Jensen, B. C. & Haack, K. R. Rearrangement of duplicated DNA in specialized cells of Neurospora. *Cell* **51**, 741–752 (1987).

41. Hashem, K. M., He, F. J., Jenner, K. H. & MacGregor, G. A. Cross-sectional survey of salt content in cheese: a major contributor to salt intake in the UK. *BMJ Open* **4**, e005051–e005051 (2014).

42. Hymery, N., Masson, F., Barbier, G. & Coton, E. Cytotoxicity and immunotoxicity of cyclopiazonic acid on human cells. *Toxicol In Vitro* **28**, 940–947 (2014).

43. Le Bars, J. Cyclopiazonic acid production by *Penicillium camemberti* Thom and natural occurrence of this mycotoxin in cheese. *Appl Environ Microbiol* **38**, 1052–1055 (1979).

44. Lund, F., Filtenborg, O. & Frisvad, J. C. *Penicillium caseifulvum*, a new species found on *Penicillium roqueforti* fermented cheeses. *Journal of Food Mycology* **1**, 95–101 (1998).

45. Barbesgaard, P., Heldt-Hansen, H. P. & Diderichsen, B. On the safety of Aspergillus oryzae: a review. *Appl Microbiol Biotechnol* **36**, 569–572 (1992).

46. Gillot, G. *et al.* Functional diversity within the *Penicillium roqueforti* species. *Int. J. Food Microbiol.* **241**, 141–150 (2017).

47. Zhu, Q., Zheng, X., Luo, J., Gaut, B. S. & Ge, S. Multilocus analysis of nucleotide variation of Oryza sativa and its wild relatives: severe bottleneck during domestication of rice. *Mol. Biol. Evol.* **24**, 875–888 (2007).

48. Marsden, C. D. *et al.* Bottlenecks and selective sweeps during domestication have increased deleterious genetic variation in dogs. *PNAS* 201512501 (2015).

49. Eyre-Walker, A., Gaut, R. L., Hilton, H., Feldman, D. L. & Gaut, B. S. Investigation of the bottleneck leading to the domestication of maize. *PNAS* **95**, 4441–4446 (1998).

50. Gaut, B. S., Díez, C. M. & Morrell, P. L. Genomics and the Contrasting Dynamics of Annual and Perennial Domestication. *Trends in Genetics* **31**, 709–719 (2015).

51. Duan, S.-F. *et al.* The origin and adaptive evolution of domesticated populations of yeast from Far East Asia. *Nat Commun* **9**, (2018).

52. Mahler, D. L., Ingram, T., Revell, L. J. & Losos, J. B. Exceptional convergence on the macroevolutionary landscape in island lizard radiations. *Science* **341**, 292–295 (2013).

53. Thorpe, R. S., Barlow, A., Malhotra, A. & Surget-Groba, Y. Widespread parallel population adaptation to climate variation across a radiation: implications for adaptation to climate change. *Mol Ecol* **24**, 1019–1030 (2015).

54. Hohenlohe, P. A. *et al.* Population Genomics of Parallel Adaptation in Threespine Stickleback using Sequenced RAD Tags. *PLoS Genetics* **6**, e1000862 (2010).

55. Duboué, E. R., Keene, A. C. & Borowsky, R. L. Evolutionary convergence on sleep loss in cavefish populations. *Current Biology* **21**, 671–676 (2011).

56. Salzburger, W. The interaction of sexually and naturally selected traits in the adaptive radiations of cichlid fishes. *Molecular Ecology* **18**, 169–185 (2009).

57. Elmer, K. R. *et al.* Parallel evolution of Nicaraguan crater lake cichlid fishes via non-parallel routes. *Nature Communications* **5**, 5168 (2014).

58. Elmer, K. R., Kusche, H., Lehtonen, T. K. & Meyer, A. Local variation and parallel evolution: morphological and genetic diversity across a species complex of neotropical crater lake cichlid fishes. *Philos Trans R Soc Lond B Biol Sci* **365**, 1763–1782 (2010).

59. Martínez-Ainsworth, N. E. & Tenaillon, M. I. Superheroes and masterminds of plant domestication. *Comptes Rendus Biologies* **339**, 268–273 (2016).

60. Gross, B. L. & Olsen, K. M. Genetic perspectives on crop domestication. *Trends in Plant Science* **15**, 529–537 (2010).

61. Decker, M. & Nielsen, P. V. The inhibitory effect of *Penicillium camemberti* and *Geotrichum candidum* on the associated funga of white mould cheese. *International Journal of Food Microbiology*

104, 51–60 (2005).

62. Gouyon, P.-H., Leriche, H., Civard, A., Reeves, H. & Hulot, N. *Aux origines de l'environnement.*
(Fayard, 2010).

63. Vavilov, N. I., Vavylov, M. I., Vavílov, N. Í. & Dorofeev, V. F. *Origin and Geography of Cultivated*
Plants. (Cambridge University Press, 1992).

64. Glass, N. L. & Donaldson, G. C. Development of primer sets designed for use with the PCR
to amplify conserved genes from filamentous ascomycetes. *Appl Environ Microbiol* **61**, 1323–
1330 (1995).

65. Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive k-mer weighting
and repeat separation. *Genome Res.* **27**, 722–736 (2017).

66. Walker, B. J. *et al.* Pilon: An Integrated Tool for Comprehensive Microbial Variant Detection
and Genome Assembly Improvement. *PLoS One* **9**, e112963 (2014).

67. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**, 2114–2120 (2014).

68. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**,
357–359 (2012).

69. McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-
generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

70. Paradis, E. pegas: an R package for population genetics with an integrated–modular approach.
Bioinformatics **26**, 419–420 (2010).

71. Korneliussen, T. S., Albrechtsen, A. & Nielsen, R. ANGSD: Analysis of Next Generation
Sequencing Data. *BMC Bioinformatics* **15**, 356 (2014).

72. Paradis, E., Claude, J. & Strimmer, K. APE: Analyses of Phylogenetics and Evolution in R
language. *Bioinformatics* **20**, 289–290 (2004).

73. Schliep, K. P. phangorn: phylogenetic analysis in R. *Bioinformatics* **27**, 592–593 (2011).

74. Hill, T. W. & Kafer, E. Improved protocols for *Aspergillus* minimal medium: trace element and
minimal medium salt stock solutions. *Fungal Genetics Reports* **48**, 20–21 (2001).

75. Kritikos, G. *et al.* A tool named Iris for versatile high-throughput phenotyping in
microorganisms. *Nature Microbiology* **2**, 17014 (2017).

76. Peterson, R. A. & Cavanaugh, J. E. Ordered quantile normalization: a semiparametric
transformation built for the cross-validation era. *Journal of Applied Statistics* **0**, 1–16 (2019).

Figure Legends

Figure 1: Cheeses and fungi used for cheese-making.
A) A panel of soft cheeses with *Penicillium camemberti* or *P. biforme* on the crust. Left: Brie de Meaux protected designation of origin, top middle: Saint Marcellin, bottom middle: Camembert-like (buffalo milk), top right: Camembert protected designation of origin, bottom right: Selles fermier, Right picture: crottin de Chavignol (goat milk). B) Pictures of *P. biforme, P. camemberti* and *P. caseifulvum* on malt agar at 25°C after 10 days of growth. C) Petri dish with a white and fluffy *P. camemberti s.s.* strain (FM013) and a green rough *P. fuscoglaucum* strain (LCP06617).

Figure 2: Population structure of 61 strains from the *Penicillium camemberti* cheese-making fungal species complex, based on whole-genome data.
A) Principal component analysis (PCA), B) Neighbor-net analysis based on SNP data. Cross-linking indicates the likely occurrence of recombination. Branch lengths are shown and the scale bar represents 0.03 substitutions per site. C) Population subdivision inferred from $K = 2$ to $K = 6$. Each column represents a strain and colored bars represent their coefficients of membership for the various gene pools. For all panels, genetic clusters are represented by the same colors: dark purple for *P. fuscoglaucum,* light blue for *P. biforme,* dark green for *P. camemberti sensu stricto* and light green for *P. caseifulvum.* The strains identified as *P. commune* (invalid name) by collectors are shown in light purple. For PCA and neighbor-net analysis, symbols correspond to the environment of collection: circles for cheese, triangles for dried sausages and squares for other environments. On the structure analysis, letters below the plots indicate the origin of the strains: C for cheese, F for food strains other than cheese and E for other environments. Red asterisks indicated the type strains of *P. biforme, P. camemberti and P. caseifulvum.*

Figure 3: Presence/absence of the *Wallaby* and *CheesyTer* horizontally transferred regions in the 61 studied strains of the *Penicillium camemberti* species complex.
A) Neighbor-joining tree for the 61 strains. The strain name color indicates the environment of collection, black for food and blue for other environments. B) Presence/absence of the *Wallaby* region, as defined in the reference genome *P. camemberti s.s.* FM013. C) Presence/absence of the *CheeysyTer* region, as defined in the reference genome *P. camemberti s.s.* FM013. On panels B and C, the red bars represent the presence of the region in a given strain, white its absence, yellow the presence of a RIP region, light gray the presence of a highly fragmented region, dark gray the presence of a partially fragmented region and dark red a partially duplicated region. U before the strain ID indicates UBOCC strains.

Figure 4: Phenotypic traits distinguishing the four lineages *Penicillium biforme, P. caseifulvum, P. camemberti sensu stricto* and *P. fuscoglaucum.* A) Mean radial growth of the four lineages on unsalted cheese, salted cheese, malt and minimal media. B) Pictures of colonies on unsalted cheese, salted cheese, malt and minimal media; *P. fuscoglaucum*: LCP05471; *P. camemberti s.s.*: UBOCC-A-113052; *P. biforme*: ESE00211; *P. caseifulvum*: LCP06622. C) Difference in opacity between the four lineages on salted cheese at 25°C. D) Difference in production of cyclopiazonic acid between the four lineages on yeast extract sucrose. C-D) Horizontal lines of the boxplots represent the upper quartile, the median and the lower quartile. Each dot represents a strain and p-values are given for the tests of differences between the lineages (Wilcoxon test). The color indicates assignments to the four lineages as in other figures.

Figure 5: Competitive abilities of *Penicillium biforme, P. caseifulvum* and *P. camemberti sensu stricto* against *Geotrichum candidum, P. biforme, P. fuscoglaucum* and *P. roqueforti* competitors. Top: Pictures of *P. roqueforti* FM164, *Geotrichum candidum* ESE00163, *P. biforme* ESE00154 and *P. fuscoglaucum* ESE00221 on no lawn (first line), *P. biforme* ESE00086 lawn (second

line), *P. camemberti s.s.* LCP04810 (third line) and *P. caseifulvum* ESE00019 (fourth line) lawns, on salted cheese. Bottom: Boxplots representing the differences of growth abilities of the competitors on different species lawn. Horizontal lines of the boxplots represent the upper quartile, the median and the lower quartile. Each dot represents a strain and p-values are given for the tests of differences between the lineages (Wilcoxon test). The color indicates assignments to the three lineages as in other figures.

Figure S1: Population structure of 47 strains from the *Penicillium camemberti sensu lato* and *P. biforme* lineages, based on whole-genome data.
A) Principal component analysis; B) Neighbor-net analysis. Cross-linking indicates the likely occurrence of recombination Branch lengths are shown and the scale bar represents 0.01 substitutions per site. For both panels, the genetic clusters are represented by the same colors: light blue for *P. biforme,* dark green for *P. camemberti sensu stricto* and light green for *P. caseifulvum.* The strains identified as *P. commune* (invalid name) by collectors are shown in light purple. The symbols correspond to the environment of collection: circles for cheese, triangles for dried sausages and squares for other environments.

Figure S2: Population structure of 61 strains from the *Penicillium camemberti* cheese-making fungal species complex, based on whole-genome data with the addition of the genome of the *P. commune* strain published in [39].
A) Principal component analysis; B) Neighbor-net analysis. Cross-linking indicates the likely occurrence of recombination Branch lengths are shown and the scale bar represents 0.08 substitutions per site. For both panels, the genetic clusters are represented by the same colors: light blue for *P. biforme,* dark green for *P. camemberti sensu stricto* and light green for *P. caseifulvum.* The strain added in this analysis, 162 3FA WT8, identified as *P. commune* (invalid name) is shown in red. The symbols correspond to the environment of collection: circles for cheese, triangles for dried sausages and squares for other environments.

Figure S3: Genomic scans of Illumina read coverage along the scaffold 18 where *Wallaby* is located.
Wallaby is shown by red bars above plots. The vertical red line represents the position of the Hce2 gene, involved in the competition against micro-organisms. When several strains showed similar coverage patterns along the scaffold, only one representative plot is shown.

Figure S4: Genomic scans of Illumina read coverage along the scaffold 17 from position 600,000 until the end of the scaffold where *CheesyTer* is located.
CheesyTer is shown by gray rectangles on plots. Vertical red lines represent the position of the two genes with relevant function for cheese-making previously identified in *CheesyTer*, i.e. a lactose permease and a beta-galactosidase. When several strains showed similar coverage patterns along the scaffold, only one representative plot is shown.

Figure S5: Illustration of repeat-induced point mutation (RIP) footprints along the first 5kb of *CheesyTer* by comparing *Penicillium camemberti sensu lato, P. biforme, P. fuscoglaucum* and *P. roqueforti* genomes. The nucleotide differences between species corresponding to RIP, i.e., CpA to TpA or TpG to TpA are shown in red, while the other types of variation are shown in blue and green.
A) Alignments of genomes from *P. camemberti sensu lato, P. biforme, P. fuscoglaucum* and *P. roqueforti* genomes, showing in that mutations typical of RIP mostly occurred in *P. camemberti sensu lato* (red traits). B) Percentages of nucleotide differences in *P. camemberti sensu lato* compared to the other genomes showing that mutations typical of RIP are most abundant.

848 **Figure S6: Cyclopiazonic acid (CPA) biosynthesis cluster in the *Penicillium camemberti*
849 species complex.** A) Schematic representation of the CPA biosynthesis cluster in the *P.
850 camemberti* reference genome FM013. B) Amino-acid alignment of the *CpaA* gene showing a 2bp
851 deletion in strains belonging to the *P. caseifulvum* genetic cluster, leading to a truncated protein.
852
853 **Table S1:** Description of the origin, donor and genome coverage of the 61 strains of the
854 *Penicillium camemberti* species complex used in this study.
855
856 **Table S2:** Results of statistical analyses performed for testing differences in traits between
857 *Penicillium camemberti s.s., P. caseifulvum, P. biforme* and *P. fuscoglaucum.* Significant P-values are indicated
858 with an asterisk and highlighted in gray.
859

A

B

P. biforme   *P. caseifulvum*   *P. camemberti s.s.*

C