Long-term repeatability in social behaviours suggests

stable social phenotypes in wild chimpanzees

Patrick J. Tkaczynski[1]*, Alexander Mielke[2]*, Liran Samuni[1,3,4], Anna Preis[5], Roman Wittig[1,4]*, Catherine

Crockford[1,4]*

[1] Department of Human Behaviour, Ecology and Culture, Max Planck Institute for Evolutionary Anthropology,

Leipzig, Germany

[2] School of Anthropology & Museum Ethnography, University of Oxford, Oxford, UK

[3] Department of Human Evolutionary Biology, Harvard University, Cambridge, MA, USA

[4] Taï Chimpanzee Project, Centre Suisse de Recherches Scientifiques, Abidjan, Côte d'Ivoire

[5] Wild Chimpanzee Foundation, Conakry, Guinea

* These authors contributed equally to the manuscript

Corresponding author: Patrick J. Tkaczynski (pjtresearchltd@gmail.com).

Abstract

Animals living in social groups navigate challenges when competing and cooperating with other group
members. Changes in demographics, dominance hierarchies or ecological factors, such as food
availability or disease prevalence, are expected to influence decision-making processes regarding social
interactions. Therefore, it could be expected individuals show flexibility in social behaviour over time to
maximise the fitness benefits of social living. To date, research across species has shown that stable
inter-individual differences in social behaviour exist, but mostly over relatively short data collection time
periods. Using data spanning over 20 years, we demonstrate that multiple social behaviours are
repeatable over the long-term in wild chimpanzees, a long-lived species occupying a complex fission-
fusion society. We controlled for temporal, ecological and demographic changes, limiting pseudo-
repeatability. We conclude that chimpanzees living in natural ecological settings have relatively stable
long-term social phenotypes over years that may be independent of life history stage or strategies. Our
results add to the growing body of literature suggesting consistent individual differences in social
tendencies are more likely the rule rather than the exception in group-living animals.

Introduction

The fitness benefits of social bonds and social connectivity are well established across group-living
animals, including in humans (1–11). Despite the adaptive advantages of maintaining social
relationships, short-term strategies of social interaction avoidance could also improve fitness, such as
during periods of disease outbreak (12). Similarly, during periods of social upheaval, such as instability in
dominance hierarchies, individuals may change how they distribute their affiliative or aggressive social
investment to reinforce key relationships or dominance position respectively (13,14). Fluctuations in
resource availability (e.g. number of available mating partners or food availability) may also influence
time allocation to affiliative social interactions, rates of aggression, social partner choice or general
gregariousness (15,16). Lastly, an individual's physiological state may fluctuate over time, e.g. during
pregnancy and/or the rearing of offspring, in turn influencing motivation for social behaviours (17).

Substantial variation in socioecological settings and internal state suggests individuals should show
flexibility in their social interaction patterns (18). However, consistent individual differences in social
behaviour have been identified across various animal taxa, suggesting group-living individuals tend to
show stable tendencies in solving social problems and interacting with other group members (19–26).
This raises the question of why some individuals appear to be consistently more or less cooperative,
gregarious, or aggressive than others.

A high degree of repeatability in a trait, i.e. the proportion of variation attributable to between-
individual differences (27), may reflect underlying stable factors such as genetics and/or irreversible
developmental adjustment to early life conditions (28–32). Under such a framework, individual patterns
of social behaviour, and thus social phenotypes, emerge as a consequence of the interaction between
genetics and exposure to the physical and social environment during development (30,32). The "social
niche hypothesis" specifically proposes that consistent individual differences in behaviour arise due to
niche specialisation to ameliorate within-species and/or within-group competition for resources (33).
Here, an individual adopts a behavioural or social strategy to acquire resources dependent upon its
existing competition-related characteristics, such as body size, health, dominance etc.

Alternatively, consistent between-individual differences in behaviour and sociality may arise due to behavioural tendencies associated with specific life history stages or dominance positions, which appear to be individual phenotypes because the data collection protocol does not extend across the lifespan of the species (34–36). Indeed, within the human literature, longitudinal studies suggest human behavioural tendencies and personality may be more labile than previously thought, with shifts in these tendencies predicted by a combination of age-related change or adjustment to particular life events, e.g. marriage (37–39). Distinguishing individual differences that are independent of life history or artefacts of socio-demographic variables requires long-term data reflecting changing life history within a particular study species.

The majority of non-human animal (hereafter animal) studies examining the repeatability of social behaviour have used comparatively short-term datasets, representing a limited period within the lifespan of the study species (34). In our study, we examine how social behaviours vary between individuals in wild, adult chimpanzees, utilising a behavioural dataset spanning over 20 years and more than 7500 full-day focal follows. We examine whether social behaviours (grooming, aggression and association) are repeatable over time both on a daily scale and when aggregated on a yearly level, with each sex analysed separately. By controlling for demographic and intrinsic factors, such as group size and age, we aim to determine the amount of variation in these social behaviours that is most likely attributable to individually stable phenotypes, thus limiting pseudo-repeatability (29).

Chimpanzees are an interesting study species to discern whether and how consistent individual differences in sociality arise and are maintained. These primates live in multi-female, multi-male groups with a high degree of fission-fusion dynamics (16,40), allowing potential for considerable fluctuation in social organisation during the course of their long lifespans (41). Chimpanzee societies also feature various cooperative behaviours, such as alliance formation, food sharing, group hunting, and territorial patrols, which likely impact on fitness and for which strong social relationships are required (42–45). Therefore, these individuals face diverse social environments and important choices regarding their social behaviour (46), which may predict a degree of flexibility in social tendencies.

In terms of dominance structure, when compared to female hierarchies, male hierarchies are dynamic and defined by high male-male competition, and there tends to be considerable reproductive skew towards high-ranking males (47–49). Although female dominance ranks do change during their lifespan, they are comparatively stable (50–53). Therefore, we expected a high degree of within-individual variation in aggressive tendencies in males in relation to lifetime fluctuations in dominance rank, whereas within-individual variation in aggression would be lower in females due to relatively stable hierarchy structure. Individual chimpanzees are highly strategic and flexible in grooming partner choice (46), while time devoted to grooming in this species can vary depending on demographic factors such as group size (54). Therefore, we expected low repeatability in grooming behaviour across long time periods. Lastly, chimpanzees sociality is characterised by a high degree of fission-fusion, allowing individuals to adjust to variation in within-group competition arising from ecological constraints, such as the availability of receptive mating partners or food (40). Competition will vary both seasonally and in the longer-term due changing group sizes or sex ratios. Therefore, we anticipated low repeatability for association behaviour as chimpanzees adjust to these fluctuations in competition.

Methods

Study Groups and Data Collection

Daily focal follow (55) data have been systematically collected by the Taï Chimpanzee Project, Côte d'Ivoire, since 1992 (56,57). We focussed on data collected since 1996, when data collection was consistent for behaviours relevant to this study and the control factors included in the models (see Statistical Analyses). This data includes observations of adult (>12 years) males and females from three fully habituated communities of chimpanzees: North (1996 – 2016), South (2002 – 2016), and East (2012 – 2016).

We collated these data to identify repeatability in interaction rates on two levels: daily and yearly (with year ending on 31st August), allowing us to control for socioecologial variables at different temporal scales. For each of these levels, we restricted the dataset to individuals with regular focal follows to ensure that the data were sufficient to capture their typical social behaviour.

For the analysis on the daily level, we included focal follow days of adult individuals that lasted at least three hours, and included individuals for whom at least 10 focal follow days were available, resulting in a dataset of 70 individuals (45 females, 25 males) and 7615 individual focal follow days (individual mean = 109 days, max = 413 days).

For the analysis on the yearly level, we included adult individuals who were followed at least nine times in the time period from 1st September to 31st August and were present in their group for all this period, and who fulfilled these criteria at least in 3 years. This resulted in a dataset of 45 individuals (24 females, 21 males) and 272 years (individual mean = 6 years, max = 15 years). The number of years or days per subject of the study are detailed in Table S1 of the supplementary materials.

Social Phenotype

On both the daily and the yearly level, we focused on three social variables: grooming, aggression, and association. We calculated variables from the perspective of the focal individual and with the focal as the actor (in case of the interactions) to ensure that they represent the individual's social propensity as much as possible.

For grooming, we extracted the time (in minutes) focal individuals spent grooming adult partners (we focused on grooming given to others rather than overall time grooming, i.e. including grooming received, as this would reflect a tendency to attract grooming partners rather than an individual tendency to groom). For aggression, we extracted the number of aggressive interactions in which the focal individual was the aggressor. Both these variables were calculated the same for the daily and yearly level. We calculated association differently for the two levels: on the daily level, we extracted the cumulative number of individuals with whom the focal individual was seen in the same 'party' (defined as all individuals within visual contact of the observer – usually < 35 m) during that day. On the yearly level, as cumulatively individuals will likely associate with all other partners during this timeframe limiting variation, we instead calculated the total strength of connections of individuals by summing their association with all adult group members, using the simple ratio index (58) for that year based on common party membership of individuals. We standardized association strength by dividing it by group size.

As association indices have to be standardized by an appropriate null model (59), we conducted
permutation analyses to confirm that associations were different than would be expected by random.
For this analysis, we generated 1000 permutations of randomizations of party membership in which the
number of individuals per party was kept constant and subsequent parties that originally had the same
party membership also remained the same after randomization to account for autocorrelation (60). The
overall number of times individuals appeared was not held constant, as we were interested in whether
they would be more central/gregarious than expected. We subtracted the mean strength of individual's
connections with all other group members arising from the permutations from the observed strength,
effectively creating an index of gregariousness (above 0 = more connected than would be expected in
that group in that year; below 0 = less connected than expected).

Statistical Analyses

All data were prepared and analyses were conducted using R 3.6.1 statistical software (61), with
Generalized Linear Mixed Models (GLMM) fitted using the 'lme4' package (62). Models were fitted
separately for males and females, for the dependent variables of aggression, grooming, and association,
and for the daily and yearly levels of data aggregation, resulting in 12 models.

The **age** of all individuals was either known (for individuals who were not yet adult at the beginning of
habituation) or was estimated in the beginning of the habituation period by experienced observers. Age
ranged from 12 years (cut-off value) to 52 years, and the value assigned was either for the day (daily
level) or the beginning of the yearly period (yearly level). We included age as a squared term in all
models to account for potentially non-linear developmental patterns, e.g. individuals in their prime
being more aggressive than younger and older individuals.

Dominance rank was calculated using a modification of the Elo rating method (63,64) based on
unidirectional pant grunt vocalizations within each sex and standardized between 0 and 1 within each
group. For the daily level, rank was extracted for each day of data an individual was observed. Rank was
extracted on 31st August for each yearly level.

For females, as **reproductive state** and **infant age** influences gregariousness and sociality (65–69), on
the daily level, we included a factor with five levels: maximum tumescence of sexual swellings, mother
with a new-born infant (below 3 months of age), mother with an un-weaned offspring (below 4 years of
age), mother with weaned but immature offspring, or none of the above. In instances where females
had weaned or un-weaned offspring but were also maximally tumescent or had a new-born infant, they
were classified as maximally tumescent or with a new-born infant, as these reproductive states were
anticipated to have a greater effect on behaviour than the presence of older offspring (81-85). On the
yearly level, we included whether they had no offspring throughout the year, had a new-born at any
point, or un-weaned offspring. On the daily level for the analyses concerning males, we included the
number of females with maximally tumescent sexual swellings on that day as a variable.

As the **sex ratio** of adult individuals in a group can influence social behaviour, we calculated the adult
sex ratio (number of males/number of females) of each group at the start of the yearly period (for the
yearly level) or of those individuals present in the focal individual's party during the day (for the daily
level).

As interaction rates could be determined by the **number of available partners** (on the daily level) or
group size (on the yearly level), we included these variables into the models. The yearly group size
variable was standardized within groups, as there would otherwise be complete separation between the
group identity and group size (as North group was considerably smaller than the others).

For both daily and yearly levels of analysis, we included an offset term for focal observation time (in
hours; log-transformed) to account for differences in base rates of how often individuals could be
observed interacting. We accounted for seasonality effects in the daily level analyses by including the
radians of Julian date as a control variable (70). All models included a random effect of group by year: as
group dynamics change over time, and we do not expect the three communities to show the same
interaction rates, this term accounts for overall differences that could be influencing how social
individuals are.

Individual Model Structures

For **aggression** (both on the daily and yearly level, and males and females), we fitted GLMMs with
Poisson error structure and log link (71), using the number of bouts per day/year as outcome variable,
and having a log-transformed offset term for observation time to control for differences in observation
effort (72). For the daily level, we included age as a quadratic term, dominance rank, available partners
(daily level) or group size (yearly level), sex ratio, and group identity as fixed effect predictors. For the
daily level, we also included the number of females with maximum swellings and the cosine and sine
functions to capture seasonal effects. We included the random intercept of Year within Group, to
account for the fact that data collected in the same time period in the same group are not independent,
and the random intercept of Individual Identity. We also included the random slopes for rank, age,
available partners, and sex ratio in Individual Identity, and rank and available partners in Group Year
(73).

For **grooming**, on both the daily and yearly level, we fitted LMMs with Gaussian error structure, using
the log-transformed hourly grooming rate as outcome variable. The fixed effects and random effect
structure were the same as described for the aggression models.

For **association** on the daily level, we fitted GLMMs with Poisson error structure, using the number of
individuals that the focal associated with on the day as outcome variable, and the number of available
partners in the group as offset term. The fixed effects were the same as described for aggression, bar
the removal of the available partners from the fixed effects. For gregariousness on the yearly level, we
used the difference between observed and expected association strength as outcome variable, fitting
LMMs with Gaussian error structure, and the same fixed effects as described above.

For all models, we z-standardized continuous predictor variables to facilitate interpretation (74). We
compared the full models including all predictors and random intercepts and slopes with a null model
including all fixed and random effects except the random intercept and slope for the individual identity
(19,75), to test whether stable individual differences had an impact on interaction distributions. To
establish the strength of the observed effect of individual identity only (76), we applied the
'r.squaredGLMM' function of the 'MuMIN' package (77) on a reduced model containing only the fixed
effects and the random intercept and slopes for individual identity. We assumed that the variance in
social behaviour that can be attributed to intra-individual stability can be captured by subtracting the
variance explained by the fixed effects alone (marginal effect size) from that explained by the fixed

effects and the individual random effect structure (conditional effect size) (76,78). Thus, for both the
daily and the yearly level, both sexes, and all three social behaviours, we report whether the random
effect significantly improved predictability, the marginal and conditional effect size, and their difference.

To avoid problems due to multicollinearity, we established the Variance Inflation Factors (VIF) of each
model (79) using a simple linear regression of the fixed effects and applying the 'vif' function of the 'car'
package (80). The maximum VIF for any model was 3.13, indicating that collinearity was not an issue. For
all Poisson models, we tested for overdispersion, which was not an issue in any model.

Results

We report the effect sizes attributed to the individual random effects and the comparison of the full
model containing the random effect and the null model without it for the different models in Table 2
and Figure 1.

In all models, including the random intercept and slopes of individual identity significantly improved
model fit, indicating that for all social behaviours involved in this study, chimpanzees showed inter-
individual differences that were consistent over time and could be detected on the daily and yearly
level.

The variation explained by the random effect of individual identity varied between 0.14 and 0.61, and
inter-individual differences tended to be more pronounced on the yearly level (mean = 0.40) than on the
daily level (mean = 0.25). Thus, knowing the individual identity, for example, would allow one to predict
50% of variation in female aggression rates across years (Table 1). For grooming, the results for both
sexes were almost identical in terms of the impact of the random effect of identity (daily = 0.18/0.19,
yearly = 0.40/0.42). In general, inter-individual differences between females were more stable than
between males (overall female mean = 0.38 vs overall male mean = 0.28), especially on the yearly level
(female yearly mean = 0.50 vs male yearly mean = 0.30). Thus, after accounting for age, dominance rank,
reproductive state (for females), group-level sex ratio and group size, stable inter-individual differences
in yearly interaction rates where more pronounced in females than in males. For grooming and
aggression, the residuals of individuals that were represented in both the yearly and daily models were
highly correlated (see Table 1), while this was not the case for gregariousness, where different outcome
variables were used. Figures 2 and 3 illustrate inter-individual variation in grooming, aggression and
association at the daily and yearly levels in male and female chimpanzees respectively.

270 **Table 2: Repeatability coefficients of chimpanzee social behaviours.** *Effect sizes and results of full null model*
271 *comparisons for the models testing the impact of the individual random intercept and slopes on grooming rates,*
272 *aggression bouts, and association in male and female Taï chimpanzees on a daily and yearly level. The correlation*
273 *between the two levels was conducted on the random intercept residuals of individuals who were present in both*
274 *datasets.*

			Marginal R^2	Conditional R^2	Individual R^2	$X^2_{df=5}$	p	Correlation Daily – Yearly
Female	grooming	daily	0.08	0.26	0.19	50.87	<0.001	0.73
		Yearly	0.20	0.60	0.40	44.42	<0.001	
	aggression	Daily	0.11	0.25	0.14	13.55	0.019	0.70
		yearly	0.17	0.67	0.50	13.04	0.011	
	association	daily	0.25	0.67	0.42	450.93	<0.001	0.30
		yearly	0.11	0.72	0.61	36.29	<0.001	
Male	grooming	daily	0.10	0.28	0.18	50.71	<0.001	0.93
		yearly	0.33	0.74	0.42	28.32	<0.001	
	aggression	daily	0.30	0.54	0.24	150.69	<0.001	0.55
		yearly	0.67	0.93	0.26	202.19	<0.001	
	association	daily	0.47	0.83	0.35	71.08	<0.001	0.20
		yearly	0.24	0.45	0.21	9.61	0.022	

Figure 1: Repeatability of Chimpanzee Social Behaviours. *Overview of the effect sizes attributed to the individual random effect, delineated by behaviour, sex, and timeframe (yearly vs daily) over 20 years of data.*

Figure 2: Inter-individual variation in grooming, aggression and association in male chimpanzees. *Plots include random effect coefficients for individual identity from models examining variation in grooming, aggression and association in male at both the daily (a, b, c) and yearly (d, e, f) level of aggregation. The x-axis indicates the random effect coefficient; the y-axis indicates individual identities. Blue individuals expressed the variable of interest (grooming, aggression, and association) more than the average expression of the variable given all fixed and random effects within the model, red individuals expressed the variables less than the given average.*

Figure 2: Inter-individual variation in grooming, aggression and association in female chimpanzees. Plots include random effect coefficients for individual identity from models examining variation in grooming, aggression and association in male at both the daily (a, b, c) and yearly (d, e, f) level of aggregation. The x-axis indicates the random effect coefficient; the y-axis indicates individual identities. Blue individuals expressed the variable of interest (grooming, aggression, and association) more than the average expression of the variable given all fixed and random effects within the model, red individuals expressed the variables less than the given average.

Discussion

Our study reveals a high degree of repeatability in social behaviour in wild chimpanzees over several
years and in many of our subjects, over a sizeable proportion of the adult lifespan of the species. The
repeatability estimates for social behaviour in our study were comparable to an average repeatability
estimate of behavioural traits across numerous animal taxa formerly generated by a meta-analytical
study (R^2=0.37; (81)). Importantly, unlike many former studies, using a comprehensive dataset, we were
able to control for temporal, seasonal and demographic changes across the long-term study period that
may influence variation in social behaviours, and thus limit pseudo-repeatability (29). As such, our
method provides reasonable confidence that our repeatability estimates reflect stable social behaviour
phenotypes that may be independent of life history stage or strategy.

The three behaviours of aggression, grooming (affiliation) and association represent three important
components of sociality for chimpanzees, facilitating social goals such as dominance rank attainment
(82,83) and the formation of social bonds (42,84), both of which influence fitness (43,44,85–88). Our
study expands on former research in this field in terms of its scale, both temporally and in the range of
behaviours examined, in a wild population, demonstrating that social phenotypes are a phenomenon in
long-lived species occupying complex social environments. Indeed, given the results of our study and
work conducted in taxa ranging from insects (23) to fish (24,25) to birds (19) to primates (20,89),
consistent individual differences in social phenotypes seem the norm for group-living animals regardless
of the degree of social complexity.

Repeatability was generally lower in the daily versus the yearly level, with implications for data
aggregation and interpretation in repeatability analyses. The daily measures incorporate more data
points with potentially fewer social interactions within them compared to the yearly measures. This may
result in more random error and thus lower repeatability estimates for the daily level. Nevertheless, the
differences between the daily and yearly measures are still informative.

The contrast between daily and yearly levels of repeatability was strongest in female aggression, with
much higher repeatability estimates in the latter. Male aggression repeatability was low for both the
daily and yearly levels. This likely reflects sex-differences in dominance hierarchies. Males use
aggression to improve their position within the dominance hierarchy and to increase their reproductive
opportunities (90–93). Therefore, their aggression rates can fluctuate due to variation in hierarchy
stability and male-male competition (14,92). Compared to males, female chimpanzee hierarchies are
relatively stable (51,53) and female aggression rates are less likely to be determined by variability in
these factors. Instead, female aggression in chimpanzees is more likely determined by individual
tendencies and seasonal, but comparatively predictable, variation in food availability (51). However,
aggression is a costly behaviour in terms of energetics (94,95) and comes with inherent risks of injury,
which could even be lethal in certain chimpanzee populations (96,97). Therefore, although in the long-
term certain females are more aggressive than others, on a day-to-day basis, individuals are likely to be
selective in when to be aggressive, reducing inter-individual differences and repeatability measures.

Females had higher repeatability estimates than those calculated for males for association. However, for
both sexes, within-individual variation appeared much lower for association compared to the other
social measures. We did not observe sex differences in the repeatability estimates for grooming and
within-individual variation in this behaviour again seems comparable for each sex. Compared to other

populations, Taï chimpanzees are considered highly gregarious, with the contrast in this population-level
difference most apparent when examining female gregariousness (65). Our results suggest that these
chimpanzees have relatively invariable preferential party sizes and rates of association. While grooming
forms an important component of social bond formation (84), chimpanzees also make contingent
grooming choices based on a range of parameters, such as audience, partner rank or context (e.g.
reconciliation after an aggression) (68). Therefore, our results highlight this contrast in social decision
making between levels of association (constrained preferences) and grooming (flexible choices).

Importantly, we found consistent individual differences in multiple forms of social interaction. Much
research to date on social phenotypes has focused solely on association patterns (19,25), or on single
forms of dyadic interaction (20, 34), with a minority of studies examining consistency in multiple forms
of social behaviours (89,98,99) . Our study shows that consistent individual differences in social
behaviour extends to patterns of aggression and affiliation, both of which should influence fitness more
than association alone (43,44,85–88,100). Indeed, both aggression and grooming involve direct, typically
physical interactions with other group members, meaning that variation in these phenotypes will be
important for factors such as rank acquisition (51,64,68,101,102), disease transmission (103–105) and
group cohesion (45,106,107). Given the potential significance of these social tendencies, understanding
how certain individuals come to be more aggressive or affiliative, as well as gregarious, than others,
requires further empirical exploration.

The social niche hypothesis proposes that individuals adopt particular social strategies to ameliorate the
resource competition based on individual idiosyncrasies that might predict competitive ability, such as
body size (33). Chimpanzees have a protracted immature period, lasting around 10-15 years, during
which, young chimpanzees consistently associate with their mothers (108). That some adult females
(which will also typically be mothers) are more aggressive, affiliative and gregarious than others is
significant to offspring development. Given repeatability, mothers consistently expose their offspring to
particular environments, whether more or less social (109,110). In certain chimpanzee populations,
high-ranking females tend to occupy areas within a group's territory with more food resources, with
low-ranking females typically occupying more resource poor areas (50). Furthermore, high-ranking
chimpanzee mothers tend to have larger offspring (111), a pattern evident in our study population
(112). This suggests that young chimpanzees may have to adopt particular social strategies arising from
both their experience of competition via social exposure and their own competitive ability, which may
also be determined by maternal effects. Therefore, having revealed that stable social phenotypes can be
maintained over years in one of our closest living relatives, chimpanzees are promising model species to
disentangle the genetic and environmental contributions to these phenotypes by taking a
developmental approach to their emergence in younger subjects.

Our approach tested for stable individual differences in social tendencies that are independent of
socioecological influences on the rates of behaviour. As a growing number of longitudinal studies have
highlighted, age- or life event-related changes in behavioural tendencies are apparent in humans (37–
39). One approach to further characterise and understand social phenotypes in chimpanzees may be to
explore whether certain social phenotypes respond to comparable life events that occur within the
species and to what degree. For example, the stable social phenotypes generated in the present study
may predict behavioural and physiological responses to the loss of positions within a dominance
hierarchy. This would allow the testing of consistent individual differences in *plasticity* to changing

environments or internal state and provide insight on the mechanisms maintaining individual
differences in social phenotypes over time.

In summary, our results add to the well-established literature on the repeatability of behaviour and
social tendencies in group-living animals. Our study informs specifically within chimpanzees on
important sex-differences in social tendencies. Furthermore, given the long development period in this
species, our results present exciting opportunities to explore in detail the factors that contribute to the
emergence of social phenotypes, such as via social niche specialisation.

Acknowledgments

We thank the Ministère de l'Enseignement supérieur et de la Recherche scientifique and the Ministère
des Eaux et Forêts of Côte d'Ivoire, and the Office Ivoirien des Parcs et Réserves for permission to
conduct the study. We are grateful to the staff of the Taï Chimpanzee Project and the Centre Suisse de
Recherches Scientifiques for support. We are indebted to the numerous field staff that have collected
the data presented.

Funding

Core funding for the Taï Chimpanzee Project is provided by the Max Planck Society since 1997. PT, CC
and RMW were supported by the European Research Council (ERC; Grant Agreement No 679787). LS
was supported by the Minerva Foundation. AP was supported by the Leakey Foundation.

References

1. McFarland R, Majolo B. Coping with the cold: predictors of survival in wild Barbary macaques,
Macaca sylvanus. Biology Letters. 2013 Aug 23;9(4):20130428.

2. Lehmann J, Majolo B, McFarland R. The effects of social network position on the survival of wild
Barbary macaques, Macaca sylvanus. Behav Ecol. 2016 Jan 1;27(1):20–8.

3. Nuñez CMV, Adelman JS, Rubenstein DI. Sociality increases juvenile survival after a catastrophic
event in the feral horse (Equus caballus). Behav Ecol. 2015 Jan 1;26(1):138–47.

4. Cameron EZ, Setsaas TH, Linklater WL. Social bonds between unrelated females increase
reproductive success in feral horses. Proc Natl Acad Sci USA. 2009 Aug 18;106(33):13850–3.

5. McFarland R, Murphy D, Lusseau D, Henzi SP, Parker JL, Pollet TV, et al. The 'strength of weak ties'
among female baboons: fitness-related benefits of social bonds. Animal Behaviour. 2017 Apr
1;126:101–6.

6. Silk J, Alberts SC, Altmann J. Social Bonds of Female Baboons Enhance Infant Survival. Science.
2003 Nov 14;302(5648):1231–4.

7.   Silk J, Beehner J, Thore T, Crockford C, Engh A, Moscovice L, et al. The benefits of social capital:
close social bonds among female baboons enhance offspring survival. Proceedings of the Royal
Society B: Biological Sciences. 2009 Sep 7;276(1670):3099–104.

8.   Silk JB, Beehner JC, Bergman TJ, Crockford C, Engh AL, Moscovice LR, et al. Strong and consistent
social bonds enhance the longevity of female baboons. Curr Biol. 2010 Aug 10;20(15):1359–61.

9.   Archie E, Tung T, Clark M, Altmann J, Alberts SC. Social affiliation matters: both same-sex and
opposite-sex relationships predict survival in wild female baboons. Proceedings of the Royal
Society B: Biological Sciences. 2014 Oct 22;281(1793):20141261.

10.   House JS, Landis KR, Umberson D. Social relationships and health. Science. 1988 Jul
29;241(4865):540–5.

11.   Campbell LAD, Tkaczynski PJ, Lehmann J, Mouna M, Majolo B. Social thermoregulation as a
potential mechanism linking sociality and fitness: Barbary macaques with more social partners
form larger huddles. Scientific Reports. 2018 Apr 17;8(1):6074.

12.   Cauchemez S, Bhattarai A, Marchbanks TL, Fagan RP, Ostroff S, Ferguson NM, et al. Role of social
networks in shaping disease transmission during a community outbreak of 2009 H1N1 pandemic
influenza. PNAS. 2011 Feb 15;108(7):2825–30.

13.   Wittig RM, Crockford C, Lehmann J, Whitten PL, Seyfarth RM, Cheney DL. Focused grooming
networks and stress alleviation in wild female baboons. Hormones and Behavior. 2008 Jun
1;54(1):170–7.

14.   Muller MN, Wrangham RW. Dominance, aggression and testosterone in wild chimpanzees: a test
of the 'challenge hypothesis.' Animal Behaviour. 2004 Jan 1;67(1):113–23.

15.   Dunbar RIM, Korstjens AH, Lehmann J. Time as an ecological constraint. Biological Reviews.
2009;84(3):413–29.

16.   Lehmann J, Korstjens AH, Dunbar RIM. Fission–fusion social systems as a strategy for coping with
ecological constraints: a primate case. Evol Ecol. 2007 Sep 1;21(5):613–34.

17.   Seebacher F, Ward AJW, Wilson RS. Increased aggression during pregnancy comes at a higher
metabolic cost. Journal of Experimental Biology. 2013 Mar 1;216(5):771–6.

18.   Oliveira RF. Social plasticity in fish: integrating mechanisms and function. Journal of Fish Biology.
2012;81(7):2127–50.

19.   Aplin LM, Firth JA, Farine DR, Voelkl B, Crates RA, Culina A, et al. Consistent individual differences
in the social phenotypes of wild great tits, Parus major. Animal Behaviour. 2015 Oct 1;108:117–27.

20.   Blaszczyk MB. Consistency in social network position over changing environments in a seasonally
breeding primate. Behav Ecol Sociobiol. 2017 Dec 27;72(1):11.

21.   Blumstein DT, Petelle MB, Wey TW. Defensive and social aggression: repeatable but independent.
Behav Ecol. 2013 Mar 1;24(2):457–61.

22.   Finger JS, Guttridge TL, Wilson ADM, Gruber SH, Krause J. Are some sharks more social than
others? Short- and long-term consistencies in the social behavior of juvenile lemon sharks. Behav
Ecol Sociobiol. 2017 Dec 29;72(1):17.

23.   Fisher DN, Rodríguez-Muñoz R, Tregenza T. Wild cricket social networks show stability across
generations. BMC Evolutionary Biology. 2016 Jul 27;16(1):151.

24.   Jacoby DMP, Fear LN, Sims DW, Croft DP. Shark personalities? Repeatability of social network traits
in a widely distributed predatory fish. Behav Ecol Sociobiol. 2014 Dec 1;68(12):1995–2003.

25.   Krause S, Wilson ADM, Ramnarine IW, Herbert-Read JE, Clément RJG, Krause J. Guppies occupy
consistent positions in social networks: mechanisms and consequences. Behav Ecol. 2017 Apr
1;28(2):429–38.

26.   Wuerz Y, Krüger O. Personality over ontogeny in zebra finches: long-term repeatable traits but
unstable behavioural syndromes. Frontiers in Zoology. 2015 Aug 24;12(1):S9.

27.   Nakagawa S, Schielzeth H. Repeatability for Gaussian and non-Gaussian data: a practical guide for
biologists. Biol Rev Camb Philos Soc. 2010 Nov;85(4):935–56.

28.   Mitchell DJ, Dujon AM, Beckmann C, Biro PA. Temporal autocorrelation: a neglected factor in the
study of behavioral repeatability and plasticity. Behav Ecol [Internet]. 2019 [cited 2019 Nov 4];
Available from: https://academic.oup.com/beheco/advance-
article/doi/10.1093/beheco/arz180/5610999

29.   Wilson AJ. How should we interpret estimates of individual repeatability? Evolution Letters.
2018;2(1):4–8.

30.   Monaghan P. Early growth conditions, phenotypic development and environmental change.
Philosophical Transactions of the Royal Society B: Biological Sciences. 2008 May
12;363(1497):1635–45.

31.   Wilson V, Guenther A, Øverli Ø, Seltmann MW, Altschul D. Future Directions for Personality
Research: Contributing New Insights to the Understanding of Animal Behavior. Animals. 2019
May;9(5):240.

32.   Kaiser A, Merckx T, Van Dyck H. Behavioural repeatability is affected by early developmental
conditions in a butterfly. Animal Behaviour. 2019 Nov 1;157:219–26.

33.   Bergmüller R, Taborsky M. Animal personality due to social niche specialisation. Trends in Ecology
& Evolution. 2010 Sep 1;25(9):504–11.

34.   Brent LJN, Ruiz-Lambides A, Platt ML. Persistent social isolation reflects identity and social context
but not maternal effects or early environment. Scientific Reports. 2017 Dec 19;7(1):1–11.

35.   Sih A, Mathot KJ, Moirón M, Montiglio P-O, Wolf M, Dingemanse NJ. Animal personality and state–
behaviour feedbacks: a review and guide for empiricists. Trends in Ecology & Evolution. 2015 Jan
1;30(1):50–60.

36. Réale D, Garant D, Humphries MM, Bergeron P, Careau V, Montiglio P-O. Personality and the
emergence of the pace-of-life syndrome concept at the population level. Philosophical
Transactions of the Royal Society B: Biological Sciences. 2010 Dec 27;365(1560):4051–63.

37. Costa PT, McCrae RR, Löckenhoff CE. Personality Across the Life Span. Annual Review of
Psychology. 2019;70(1):423–48.

38. Roberts BW, Mroczek D. Personality Trait Change in Adulthood. Curr Dir Psychol Sci. 2008 Feb
1;17(1):31–5.

39. Denissen JJA, Luhmann M, Chung JM, Bleidorn W. Transactions between life events and
personality traits across the adult lifespan. Journal of Personality and Social Psychology.
2019;116(4):612–33.

40. Lehmann J, Boesch C. To fission or to fusion: effects of community size on wild chimpanzee (Pan
troglodytes verus) social organisation. Behav Ecol Sociobiol. 2004 Jul 1;56(3):207–16.

41. Shizuka D, Johnson AE. The long view on demographic effects on social networks: a response to
comments on Shizuka and Johnson. Behav Ecol [Internet]. [cited 2019 Nov 21]; Available from:
https://academic.oup.com/beheco/advance-article/doi/10.1093/beheco/arz184/5618606

42. Mitani JC. Male chimpanzees form enduring and equitable social bonds. Animal Behaviour. 2009
Mar 1;77(3):633–40.

43. Gilby IC, Brent LJN, Wroblewski EE, Rudicell RS, Hahn BH, Goodall J, et al. Fitness benefits of
coalitionary aggression in male chimpanzees. Behav Ecol Sociobiol. 2013 Mar 1;67(3):373–81.

44. Samuni, Preis A., Mielke A., Deschner T., Wittig R. M., Crockford C. Social bonds facilitate
cooperative resource sharing in wild chimpanzees. Proceedings of the Royal Society B: Biological
Sciences. 2018 Oct 10;285(1888):20181643.

45. Samuni L, Preis A, Mundry R, Deschner T, Crockford C, Wittig RM. Oxytocin reactivity during
intergroup conflict in wild chimpanzees. PNAS. 2017 Jan 10;114(2):268–73.

46. Mielke A, Preis A, Samuni L, Gogarten JF, Wittig RM, Crockford C. Flexible decision-making in
grooming partner choice in sooty mangabeys and chimpanzees. Royal Society Open Science.
2018;5(7):172143.

47. Surbeck M, Langergraber KE, Fruth B, Vigilant L, Hohmann G. Male reproductive skew is higher in
bonobos than chimpanzees. Current Biology. 2017 Jul 10;27(13):R640–1.

48. Newton-Fisher NE, Thompson ME, Reynolds V, Boesch C, Vigilant L. Paternity and social rank in
wild chimpanzees (Pan troglodytes) from the Budongo Forest, Uganda. American Journal of
Physical Anthropology. 2010;142(3):417–28.

49. Wroblewski EE, Murray CM, Keele BF, Schumacher-Stankey JC, Hahn BH, Pusey AE. Male
dominance rank and reproductive success in chimpanzees, Pan troglodytes schweinfurthii. Animal
Behaviour. 2009 Apr 1;77(4):873–85.

50.   Murray CM, Mane SV, Pusey AE. Dominance rank influences female space use in wild
chimpanzees, Pan troglodytes: towards an ideal despotic distribution. Animal Behaviour. 2007 Dec
1;74(6):1795–804.

51.   Wittig RM, Boesch C. Food Competition and Linear Dominance Hierarchy Among Female
Chimpanzees of the Taï National Park. International Journal of Primatology. 2003 Aug 1;24(4):847–
67.

52.   Pusey A, Williams J, Goodall J. The Influence of Dominance Rank on the Reproductive Success of
Female Chimpanzees. Science. 1997 Aug 8;277(5327):828–31.

53.   Mielke A, Crockford C, Wittig RM. Rank Changes in Female Chimpanzees in Tai National Park. In:
Wittig RM, Boesch C, editors. The Chimpanzees of the Tai Forest: 40 Years of Research. Cambridge
University Press; 2019.

54.   Lehmann J, Korstjens AH, Dunbar RIM. Group size, grooming and social cohesion in primates.
Animal Behaviour. 2007 Dec 1;74(6):1617–29.

55.   Altmann J. Observational study of behavior: Sampling methods. Behaviour. 1974;49(3–4):227–66.

56.   Boesch C, Boesch-Achermann H. The chimpanzees of the Taï Forest: Behavioural ecology and
evolution. Oxford: Oxford University Press; 2000.

57.   Wittig RM, Boesch C. Observation Protocol and Long-Term Data Collection in Taï. In: Boesch C,
Wittig RM, Crockford C, Vigilant L, Deschner T, Leendertz F, editors. The Chimpanzees of the Taï
Forest: 40 Years of Research [Internet]. Cambridge University Press; 2019 [cited 2019 Dec 11].
Available from: https://www.nhbs.com/the-chimpanzees-of-the-tai-forest-40-years-of-research-
book

58.   Whitehead H. Analyzing animal societies: quantitative methods for vertebrate social analysis.
Chicago, USA: University of Chicago Press; 2008.

59.   Farine DR, Whitehead H. Constructing, conducting and interpreting animal social network analysis.
Altizer S, editor. Journal of Animal Ecology. 2015 Sep;84(5):1144–63.

60.   Surbeck M, Girard-Buttoz C, Boesch C, Crockford C, Fruth B, Hohmann G, et al. Sex-specific
association patterns in bonobos and chimpanzees reflect species differences in cooperation. Royal
Society Open Science. 2017;4(5):1–20.

61.   R Development Core Team. R: A language and environment for statistical computing. R Foundation
for Statistical Computing Vienna Austria. 2016;0:{ISBN} 3-900051-07-0.

62.   Bates D, Mächler M, Bolker BM, Walker SC. Fitting linear mixed-effects models using lme4. Journal
of Statistical Software. 2015;67(1):1–48.

63.   Foerster S, Franz M, Murray CM, Gilby IC, Feldblum JT, Walker KK, et al. Chimpanzee females
queue but males compete for social status. Scientific Reports. 2016 Dec 14;6(1):1–11.

564 64. Mielke A, Samuni L, Preis A, Gogarten JF, Crockford C, Wittig RM. Bystanders intervene to impede
565   grooming in Western chimpanzees and sooty mangabeys. Royal Society Open Science.
566   2017;4(11):171296.

567 65. Riedel J, Franz M, Boesch C. How feeding competition determines female chimpanzee
568   gregariousness and ranging in the Taï National Park, Côte d'Ivoire. American Journal of
569   Primatology. 2011 Apr;73(4):305–13.

570 66. Wittiger L, Boesch C. Female gregariousness in Western Chimpanzees (Pan troglodytes verus) is
571   influenced by resource aggregation and the number of females in estrus. Behavioral Ecology and
572   Sociobiology. 2013;67(7):1097–111.

573 67. Foerster S, McLellan K, Schroepfer-Walker K, Murray CM, Krupenye C, Gilby IC, et al. Social bonds
574   in the dispersing sex: Partner preferences among adult female chimpanzees. Animal Behaviour.
575   2015;105:139–52.

576 68. Mielke A, Preis A, Samuni L, Gogarten JF, Wittig RM, Crockford C. Flexible decision-making in
577   grooming partner choice in sooty mangabeys and chimpanzees. Royal Society Open Science.
578   2018;5(172143).

579 69. Pepper JW, Mitani JC, Watts DP. General Gregariousness and Specific Social Preferences among
580   Wild Chimpanzees. International Journal of Primatology. 1999;20(5).

581 70. Wessling EG, Deschner T, Mundry R, Pruetz JD, Wittig RM, Kühl HS. Seasonal Variation in
582   Physiology Challenges the Notion of Chimpanzees (Pan troglodytes verus) as a Forest-Adapted
583   Species. Frontiers in Ecology and Evolution. 2018;6.

584 71. Dobson AJ. An introduction to generalized linear models. New York. London: Chapman & Hall/CRC;
585   2002. 225 p.

586 72. Baayen RH, Davidson DJ, Bates DM. Mixed-effects modeling with crossed random effects for
587   subjects and items. Journal of Memory and Language. 2008;59(4):390–412.

588 73. Schielzeth H, Forstmeier W. Conclusions beyond support: Overconfident estimates in mixed
589   models. Behavioral Ecology. 2009;20(2):416–20.

590 74. Schielzeth H. Simple means to improve the interpretability ofregression coefficients. Methods in
591   Ecology and Evolution. 2010 Feb 10;1(2):103–13.

592 75. Nakagawa S, Schielzeth H. Repeatability for Gaussian and non-Gaussian data: A practical guide for
593   biologists. Vol. 85, Biological Reviews. 2010. p. 935–56.

594 76. Nakagawa S, Schielzeth H. A general and simple method for obtaining R2 from generalized linear
595   mixed-effects models. Methods in Ecology and Evolution. 2013;4(2):133–42.

596 77. Bartoń K. MuMIn: Mutli-Model Inference. 2018.

597 78. Dingemanse NJ, Dochtermann NA. Quantifying individual variation in behaviour: mixed-effect
598   modelling approaches. Journal of Animal Ecology. 2013;82(1):39–54.

79.   Field AP, Miles J, Field Z. Discovering Statistics Using R. Vol. 58, Statistics. 2012. 53–57 p.

80.   Fox J, Weisberg S, Adler D, Bates DM, Baud-Bovy G, Ellison S, et al. Package 'car.' R topics
documented. 2014;167.

81.   Bell AM, Hankison SJ, Laskowski KL. The repeatability of behaviour: a meta-analysis. Animal
Behaviour. 2009 Apr 1;77(4):771–83.

82.   Nishida T. Alpha status and agonistic alliance in wild chimpanzees (Pan troglodytes schweinfurthii).
Primates. 1983 Jul 1;24(3):318–36.

83.   Duffy KG, Wrangham RW, Silk JB. Male chimpanzees exchange political support for mating
opportunities. Current Biology. 2007 Aug 7;17(15):R586–7.

84.   Crockford C, Wittig RM, Langergraber K, Ziegler TE, Zuberbühler K, Deschner T. Urinary oxytocin
and social bonding in related and unrelated wild chimpanzees. Proc R Soc B. 2013 Mar
22;280(1755):20122765.

85.   Cowlishaw G, Dunbar RIM. Dominance rank and mating success in male primates. Animal
Behaviour. 1991 Jun 1;41(6):1045–56.

86.   Silk JB. Social Components of Fitness in Primate Groups. Science. 2007 Sep 7;317(5843):1347–51.

87.   Langergraber K, Mitani J, Vigilant L. Kinship and social bonds in female chimpanzees (Pan
troglodytes). American Journal of Primatology. 2009;71(10):840–51.

88.   Lemoine S, Preis A, Samuni L, Boesch C, Crockford C, Wittig RM. Between-Group Competition
Impacts Reproductive Success in Wild Chimpanzees. Current Biology. 2020 Jan 20;30(2):312-
318.e3.

89.   Tkaczynski P. The behavioural ecology of personality in wild Barbary macaques [Internet] [Ph.D.].
University of Roehampton; 2016 [cited 2018 Jul 30]. Available from:
https://pure.roehampton.ac.uk/portal/en/studentthesis/the-behavioural-ecology-of-personality-
in-wild-barbary-macaques(023582d2-2214-448c-bf12-c1bef7d5549e).html

90.   Muller MN, Wrangham RW. Dominance, cortisol and stress in wild chimpanzees (<Emphasis
Type="Italic">Pan troglodytes schweinfurthii</Emphasis>). Behav Ecol Sociobiol. 2004 Feb
1;55(4):332–40.

91.   Anestis SF. Testosterone in juvenile and adolescent male chimpanzees (Pan troglodytes): Effects of
dominance rank, aggression, and behavioral style. American Journal of Physical Anthropology.
2006;130(4):536–45.

92.   Preis A, Samuni L, Deschner T, Crockford C, Wittig RM. Urinary Cortisol, Aggression, Dominance
and Competition in Wild, West African Male Chimpanzees. Front Ecol Evol [Internet]. 2019 [cited
2019 Nov 21];7. Available from:
https://www.frontiersin.org/articles/10.3389/fevo.2019.00107/full

93.   Wittig RM, Boesch C. "Decision-making" in conflicts of wild chimpanzees (Pan troglodytes): an
634   extension of the Relational Model. Behav Ecol Sociobiol. 2003 Sep 1;54(5):491–504.

635   94.   Georgiev AV, Klimczuk ACE, Traficonte DM, Maestripieri D. When Violence Pays: A Cost-Benefit
636   Analysis of Aggressive Behavior in Animals and Humans. Evol Psychol. 2013 Jul
637   1;11(3):147470491301100320.

638   95.   Wittig RM, Crockford C, Weltring A, Deschner T, Zuberbühler K. Single Aggressive Interactions
639   Increase Urinary Glucocorticoid Levels in Wild Male Chimpanzees. PLOS ONE. 2015 Feb
640   25;10(2):e0118695.

641   96.   Wrangham RW, Wilson ML, Muller MN. Comparative rates of violence in chimpanzees and
642   humans. Primates. 2006 Jan 1;47(1):14–26.

643   97.   Wilson ML, Boesch C, Fruth B, Furuichi T, Gilby IC, Hashimoto C, et al. Lethal aggression in *Pan* is
644   better explained by adaptive strategies than human impacts. Nature. 2014 Sep;513(7518):414–7.

645   98.   Madlon-Kay S, Brent L, Montague M, Heller K, Platt M. Using Machine Learning to Discover Latent
646   Social Phenotypes in Free-Ranging Macaques. Brain Sciences. 2017 Jul;7(7):91.

647   99.   Sanderson JL, Stott I, Young AJ, Vitikainen EIK, Hodge SJ, Cant MA. The origins of consistent
648   individual differences in cooperation in wild banded mongooses, Mungos mungo. Animal
649   Behaviour. 2015 Sep 1;107:193–200.

650   100.   Wittig RM, Crockford C, Weltring A, Langergraber KE, Deschner T, Zuberbühler K. Social support
651   reduces stress hormone levels in wild chimpanzees across stressful events and everyday
652   affiliations. Nature Communications. 2016 Nov 1;7:13361.

653   101.   Watts DP. Grooming Between Male Chimpanzees at Ngogo, Kibale National Park. II. Influence of
654   Male Rank and Possible Competition for Partners. International Journal of Primatology. 2000 Apr
655   1;21(2):211–38.

656   102.   Muller MN, Mitani JC. Conflict and Cooperation in Wild Chimpanzees. In: Advances in the Study of
657   Behavior [Internet]. Academic Press; 2005 [cited 2019 Nov 21]. p. 275–331. Available from:
658   http://www.sciencedirect.com/science/article/pii/S0065345405350078

659   103.   Salathé M, Kazandjieva M, Lee JW, Levis P, Feldman MW, Jones JH. A high-resolution human
660   contact network for infectious disease transmission. PNAS. 2010 Dec 21;107(51):22020–5.

661   104.   Duboscq J, Romano V, Sueur C, MacIntosh AJJ. Network centrality and seasonality interact to
662   predict lice load in a social primate. Scientific Reports. 2016 Feb 26;6(1):1–13.

663   105.   MacIntosh AJJ, Jacobs A, Garcia C, Shimizu K, Mouri K, Huffman MA, et al. Monkeys in the Middle:
664   Parasite Transmission through the Social Network of a Wild Primate. PLOS ONE. 2012 Dec
665   5;7(12):e51144.

666   106.   Grueter CC, Bissonnette A, Isler K, van Schaik CP. Grooming and group cohesion in primates:
667   implications for the evolution of language. Evolution and Human Behavior. 2013 Jan 1;34(1):61–8.

107. Radford AN, Majolo B, Aureli F. Within-group behavioural consequences of between-group
conflict: a prospective review. Proceedings of the Royal Society B: Biological Sciences. 2016 Nov
30;283(1843):20161567.

108. Boesch C, Boesch-Achermann H. The chimpanzees of the Taï Forest: Behavioural ecology and
evolution. Oxford University Press, USA; 2000.

109. Siegeler K, Sachser N, Kaiser S. The social environment during pregnancy and lactation shapes the
behavioral and hormonal profile of male offspring in wild cavies. Developmental Psychobiology.
2011;53(6):575–84.

110. Sachser N, Hennessy MB, Kaiser S. The adaptive shaping of social behavioural phenotypes during
adolescence. Biology Letters. 2018 Nov 30;14(11):20180536.

111. Pusey AE, Oehlert GW, Williams JM, Goodall J. Influence of Ecological and Social Factors on Body
Mass of Wild Chimpanzees. Int J Primatol. 2005 Feb 1;26(1):3–31.

112. Samuni L, Tkaczynski P, Deschner T, Löhrrich T, Wittig RM, Crockford C. Maternal effects on
offspring growth indicate post-weaning juvenile dependence in chimpanzees (Pan troglodytes
verus). Frontiers in Zoology. 2020 Jan 7;17(1):1.

683