Gut microbial genes are associated with neurocognition and brain development in healthy children

Kevin S. Bonham [1], Muriel M. K. Bruchhage [2], Sophie Rowland [1], Lexie Volpe [2], Kellyn Dyer [2], RESONANCE Consortium, Viren D'Sa [2,3], Curtis Huttenhower [4], Sean C. L. Deoni [2,3,5,6], Vanja Klepac-Ceraj* [1]

1. Department of Biological Sciences, Wellesley College, Wellesley, MA, 02481, USA

2. Advanced Baby Imaging Lab, Hasbro Children's Hospital, Rhode Island Hospital, Providence, RI, 02903, USA

3. Department of Pediatrics, Warren Alpert Medical School at Brown University, Providence, RI, 02912, USA

4. Department of Biostatistics, Harvard T. H. Chan School of Public Health, Boston, MA, USA

5. Department of Radiology, Warren Alpert Medical School at Brown University, Providence, RI, 02912, USA

6. Maternal, Newborn, and Child Health Discovery & Tools, Bill & Melinda Gates Foundation; Seattle WA, USA

ORCID:
Kevin: 0000-0003-3200-7533
Muriel: 0000-0001-9637-0951
Curtis: 0000-0002-1110-0096
Vanja: 0000-0001-5387-5706

* Address correspondence to vklepacc@wellesley.edu

Abstract

Both the brain and microbiome of humans develop rapidly in the first years of life, enabling extensive signaling between the gut and central nervous system (dubbed the "gut-brain axis"[1,2]). While emerging evidence implicates gut microorganisms and microbiota composition in cognitive outcomes and neurodevelopmental disorders (e.g., autism), the role of the gut microbial metabolism on typical neurodevelopment has not been explored in detail.

To examine the relationships between early childhood gut microbiome and neurocognitive development, we collected stool samples from 251 pregnant women and 313 children enrolled in the RESONANCE study of child development - part of the NIH Environmental Influences on Child Health Outcomes (ECHO) Initiative (Table 1). RESONANCE is a longitudinal observational study of healthy and neurotypical brain development that spans the fetal and infant to adolescent life stages, combining neuroimaging (magnetic resonance imaging, MRI), neurocognitive assessments, bio-specimen analyses, and rich demographic, socioeconomic, family and medical history information (Supplementary Figure 1).

Table 1 - Baseline characteristics of ECHO RESONANCE participants	
All samples (n)	802
Total subjects (n)	561
Pregnant women (n)	251
Kids (n)	310
Kids under 1yo (n)	81
Kids over 2yo (n)	200
Kids with high resolution scan (n)	141
Kids with cognitive function score (n)	263
Kids with both (n)	133
Non-white kids (%)	40.19
Mixed race kids (%)	23.49
Age in years (mean, SD)	4.27, 3.58
BMI (mean, SD)	16.66, 2.68
Maternal SES (mean, SD)	51.68, 10.86

As an initial characterization step, we used shotgun metagenomic sequencing to

generate taxonomic and functional profiles for each of our maternal and child fecal samples.

Participant age was the greatest driver of both taxonomic and functional diversity, as expected

(Figure 1a, b). Children under one year of age formed a distinct cluster from older children,

characterized by high aerobe load and low alpha (within-sample) diversity (Figure 1b,

Supplementary Figure 1), and samples from children over two years old were similar to

maternal samples (Figure 1a, c, Supplementary Figure 1a).

Figure 1: The gut microbiome of children is associated with neurocognition.

a, Principal coordinates analysis (PCoA) for all subjects based on species-level Bray-Curtis dissimilarity; most variation is driven by age and the presence of *Prevotella*[3,4]. **b**, PCoA based on Bray-Curtis dissimilarity in functional profiles (UniRef90 accessory genes) after removing gene families that were present in >90% of subjects in a given age group; variation is driven by similar effects as for taxonomic profiles. **c**, First principle coordinate axis of species PCoA for child samples vs. age; younger children cluster away from older children, and are lower in diversity. **d**, First principle coordinate axis of accessory gene family PCoA for child samples vs. age; functional diversity increases with age. **e**, PERMANOVA analysis for selected subject metadata vs. pairwiase Bray-Curtis dissimilarity for species-level taxonomic or functional profiles. Functional profiles include Kegg-Orthology (KO), Pfams or accessory UniRef90s; subject and subject type include all subjects, 2+ includes only children over 2 years old, others include all children for which the measure was available; stars indicate significance after Benjamini-Hochberg FDR correction. (* <0.1, ** < 0.01, *** < 0.001) **f**, Child age vs. normalized cognitive function score; age-appropriate assessments achieve similar score distributions for their age groups. **g**, FSEA analysis for gene sets with neuroactive potential (see Methods).

As in previous adult and infant cohorts, functional beta diversity was generally lower than

taxonomic diversity (Supplementary Figure 1), suggesting that healthy guts select for similar

gene functions even when different species contribute those functions. However, this

interpretation may be complicated by the fact that as many as 50% of sequencing reads in

some samples are not mapped to any of the reference genes used, and are thus unclassified

(Figure 1b). Interestingly, though children under 1 tended to have substantially fewer species

and, therefore, fewer total genes (Figure 1d, right), those genes tended to be better

characterized (Figure 1b, d)[5]. Consistent with previous studies of adult cohorts from

industrialized countries[4], another major driver of variation visible from principal coordinates

analysis was the presence of Prevotella copri (Supplementary Figure 3). Like samples from very

young children, samples with P. copri had reduced diversity compared with samples from older

children and pregnant mothers without P. copri (Figure 1a, Supplementary Figure 3). Overall,

these results are consistent with prior studies of adult and childhood gut microbiomes.

To assess the potential role of the microbiome in these neurocognitive processes, child

stool samples were collected alongside MRI and age-appropriate neurocognitive evaluations

(Figure 1f). Throughout childhood, a child's brain undergoes remarkable anatomical,

microstructural, organizational, and functional change. By age 5, a child's brain has reached

>85% of its adult size, has achieved near-adult levels of myelination, and the pattern of axonal

connections has been established[6]. Measures of overall cognitive ability (e.g., intelligence

quotient, IQ)[7,8], MRI measures of cortical volume and morphometry, as well as other potentially

relevant clinical metadata, were compared to taxonomic and functional profile dissimilarity by

PERMANOVA[9,10]. Consistent with previous studies, inter-individual variation accounted for the

majority of variation in microbial taxonomic and functional profiles ([79%, 77%], q < 0.001)

(Figure 1e, Supplementary Table 1). Subject type (child or mother) accounted for a large

amount of variation (4-6% q < 0.001), but this effect dropped to 2% when children under 2 years

of age were excluded, suggesting that age, rather than subject type, is actually responsible for

driving the taxonomic and functional variation. Among children's samples, age accounted for

over 20% of variation in both taxonomic and functional profiles, but this effect also largely

disappeared when children under 2 were excluded (Figure 1e, Supplementary Table 1),

suggesting that the age effect is primarily driven by the enormous changes in the microbiome

over the first year.

Microbiome taxonomic and functional variation was also associated with moderate but

significant differences in several neurocognitive measures including age-appropriate measures

of cognitive ability ([2.04%, 2.08%], q < 0.01, N=263). We also found significant associations

between regional brain volumes and microbial taxonomic and functional variation, including the

sizes of the cerebellum ([3.63%, 2.82%], q < 0.01), the subcortex ([3.64% 5.42%], q < [0.01,

0.001]), neocortex ([1.95% 1.1%], q < [0.01, NS]), and limbic regions ([4.31% 5.9%], q < [0.01,

0.001]) (N = 141 for all high resolution scans) even after controlling for the effect of age on brain

volume. These results suggest that there is a strong relationship between the gut microbiome

and neurocognitive development (Figure 1e). Though the direction of causality cannot be

determined, experimental models of brain development neurological disorders have

demonstrated that microbes in the intestine may have causal effects on the functioning of the

central nervous system[11–14].

To investigate potential mechanisms through which the gut microbiome might affect

neurostructural and neurocognitive development in infants and young children, we focused on a

group of microbial genes with neuroactive potential identified by Valles-Colomer et al.[15] which

code for enzymes that metabolize neuroactive compounds. We analyzed the association of

each of these gene sets with our neurocognitive measures using feature set enrichment

analysis (FSEA)[16,17] (Figure 1g, Supplementary Table 2). Briefly, we calculated the Pearson

correlation between the relative abundance of all identified UniRef90 gene families with each

neurocognitive measure, then calculated the Mann-Whitney U statistic for each potentially

neuroactive subset. Using this analysis, we observed that catabolic and anabolic pathways for

several molecules known to be important in the developing brain were significantly associated

with overall cognitive function scores and the size of brain subregions.

In particular, microbial genes for GABA synthesis were positively associated with

neocortical (q < 0.01, Figure 1g, Supplementary Table 2), subcortical (q < 0.01), and limbic (q <

0.001) volume, and negatively associated with cerebellar volume (q < 0.01) and overall

cognitive function (q < 1e-5). Interestingly, GABA degradation genes were also positively

associated with the size of the subcortex and negatively associated with cognitive function (q <

0.05). This may be due to higher GABA synthesis selecting for the ability to catabolize this

molecule, making it difficult to assess how actual GABA concentrations in the gut are associated

with brain development.

Unlike the metabolism of GABA, glutamate synthesis and degradation genes have an

inverse relationship (Figure 1g, Supplementary Table 2). The glutamate degradation gene set

was negatively associated with cognitive function (q < 1e-4) and cerebellar volume (q < 0.05)

and positively associated with the size of the neocortex (q < 0.05), while glutamate synthesis

was marginally negatively associated with overall cognitive function and the size of the

neocortex, while positively associated with the size of the cerebellum (q < 0.05). However, it

remains difficult to predict how gut concentrations of glutamate might be related to microbial

metabolism; while it might be intuitive to expect that higher glutamate synthesis and lower

glutamate degradation would lead to higher gut concentrations of glutamate, it might also be the

case that lower glutamate concentrations select for microbes that can synthesize it and against

those that break it down. Glutamate is also far more prevalent in the diet and can be rapidly

metabolized by gut epithelial cells, making the relationship between gut concentrations and

microbial metabolism even more complex[18].

Understanding the relationship of intestinal GABA and glutamate metabolism may be

particularly important to understanding the role of the gut microbiome in early childhood

cognitive development. Together, GABA and glutamate make up the main cerebellar output

neurons[19,20] with inhibitory GABAergic cells being widely distributed in the cerebellum projecting

out of the cerebellar cortex and excitatory glutamatergic neurons receiving afferent input. The

cerebellum, coined 'the little brain', is one of the earliest brain regions to develop[6,21], but

continues to grow into adulthood[22], making it especially vulnerable for disorder and disease[23].

Specifically, neurodevelopmental disorders, such as autism spectrum disorder, have been

associated with an imbalance of the inhibitory/excitatory system regulated by glutamate and

GABA, with recent evidence suggesting an impaired conversion of glutamate to GABA in the

disorder[24].

This is the first look at an ongoing study of child neurocognitive and microbiome

development. Using cross-sectional data, we have shown that differences in gut microbial taxa

and genes are associated with the structural development of the brain and with cognitive

development. In addition, we have shown that particular microbial gene sets with neuroactive

potential are associated with neurocognitive development, thus perhaps playing a direct role in

the gut luminal exposure of children to neuroactive metabolites. This study is ongoing, and we

are collecting additional clinical data such as participant genetic profiles, lead exposure, air

quality data and more to understand how the environment, microbiome, and biological

development interact to shape neurocognition. Future studies assessing gut metabolite pools

combined with MR spectroscopic methods to quantify concentrations of neurotransmitters such

as GABA and glutamate in the brain, as well humanized mouse models and longitudinal human

data, will provide further insight into the interactions of microbial metabolism and neurocognitive

development. As we continue to follow these subjects, we will be able to identify how early life

microbial exposures, including exposures of mothers *in utero*, might affect future neurocognitive

outcomes.

165   Materials and methods

166   Cohort description

Data used in this study were drawn from the ongoing longitudinal RESONANCE study of healthy

and neurotypical brain and cognitive development, based at Brown University in Providence, RI,

USA. From the RESONANCE cohort, 310 typically-developing children and 251 healthy pregnant

women were selected for analysis in this study. General participant demographics are provided

in **Table 1**, with children being representative of the RI population. As a broad background,

children in the BAMBAM cohort were born full-term (>37 weeks gestation) with height and weight

average for gestational age, and from uncomplicated singleton pregnancies. Children with known

major risk factors for developmental abnormalities at enrollment were excluded. In addition to

screening at the time of enrollment, on-going screening for worrisome behaviors using validated

tools were performed to identify at-risk children and remove them from subsequent analysis.

177   Additional data collection

178   Demographic and other non-biospecimen data such as race and ethnicity, parental education

179   and occupation, feeding behavior (breast- and formula-feeding), child weight and height, were

180   collected through questionnaires or direct examination as appropriate. All data were collected at

181   every assessment visit, scheduled on the same day of the MRI scan or at least within 2 weeks

182   of the scan date.

Approval for human patient research

All procedures for this study were approved by the local institutional review board at Rhode Island Hospital, and all experiments adhered to the regulation of the review board. Written informed consent was obtained from all parents or legal guardians of enrolled participants.

Stool sample collection and handling

Stool samples (n=802) were collected by parents in OMR-200 tubes (OMNIgene GUT, DNA Genotek, Ottawa, Ontario, Canada), stored on ice, and brought within 24 hrs to the lab in RI where they were immediately frozen at -80˚C. Stool samples were not collected if the infant had taken antibiotics within the last two weeks.

DNA extraction and sequencing of metagenomes

All processing of the samples was done at Wellesley College (Wellesley, MA). Nucleic acids were extracted from stool samples using the RNeasy PowerMicrobiome kit automated on the QIAcube (Qiagen, Germantown, MD), excluding the DNA degradation steps. Extracted DNA was sequenced at the Integrated Microbiome Resource (IMR, Dalhousie University, NS, Canada).

Shotgun metagenomic sequencing was performed on all samples. A pooled library (max 96 samples per run) was prepared using the Illumina Nextera Flex Kit for MiSeq and NextSeq from 1 ng of each sample. Samples were then pooled onto a plate and sequenced on the Illumina NextSeq 550 platform using 150+150 bp paired-end "high output" chemistry, generating ~400 million raw reads and ~120 Gb of sequence.

Computational methods, statistical analyses and data availability

Raw and processed data (excluding PHI) is available through SRA and Zenodo.org[25]. All code used for statistical and other analysis was version controlled and is available on github[26].

Software packages included vegan (R package) for PERMANOVAs, MultivariateStats.jl for MDS

analysis, HypothesisTests.jl for Mann-Whitney U tests (used in FSEA analysis),

MultipleTesting.jl for false discovery rate correction, and Makie.jl for plotting.

Metagenomic data were analyzed using the bioBakery[27] family of tools with default parameters.

Briefly, KneadData (v0.7.1) was used to trim and filter raw sequence reads and to separate human

reads from bacterial sequences. Samples that passed quality control were taxonomically profiled

to the species level using MetaPhlAn2 (v2.7.7) and functionally profiled by HUMAnN2 (v0.11.1).

212   MRI Acquisition and data processing

Structural T_1-weighted MRI scans were acquired on a 3T Siemens Trio scanner with a 12-

channel head RF array, preprocessed using a multistep registration procedure. Cortical

reconstruction and volumetric segmentation was performed with the Freesurfer image analysis

suite, which is documented and freely available for download online

(http://surfer.nmr.mgh.harvard.edu/). Brain regions were divided into neocortex, cerebellum,

limbic and subcortical regions (for more details on acquisition and processing, see extended

methods).

220   Neurocognitive assessments

Overall cognitive function was defined by the Early Learning Composite as assessed via the

Mullen Scales of Early Learning (MSEL)[7], a standardized and population normed tool for

assessing fine and gross motor, expressive and receptive language, and visual reception

functioning in children from birth through 68 months of age.

The third edition of the Bayley Scales of Infant and Toddler Development (Bayley's III) is a

standard series of measures primarily to assess the development of infants and toddlers,

ranging from 1 to 42 months of age[28].

The Wechsler Intelligence Quotient for Children (WISC)[29] is an individually administered

standard intelligence test for children aged 6 to 16 years. It derives a full scale intelligence

quotient (IQ) score, which we used to assess overall cognitive functioning.

The fourth edition of the Wechsler Preschool and Primary Scale of Intelligence (WPPSI-IV)[29] is

an individually administered standard intelligence test for children aged 2 years 6 months to 7

years 7 months, trying to meet the increasing need for the assessment of preschoolers. Just as

the WISC, it derives a full scale IQ score, which we used to assess overall cognitive functioning.

Acknowledgments

236   The authors would like to thank Christopher Loiselle and Jennifer Beauchemin for assistance

237   with biospecimen and metadata collection, Nisreen Abo-Sido for substantial sample preparation

238   and DNA extraction, and Julius Krumbiegel for extensive help with figure generation code.

239   This work was supported by the National Institutes of Health (UH3 ODD023313 and R01

240   MH087510).

References

1. Cryan, J. F. & O'Mahony, S. M. The microbiome-gut-brain axis: from bowel to behavior.

Neurogastroenterol. Motil. **23**, 187–192 (2011).

2. Grenham, S., Clarke, G., Cryan, J. F. & Dinan, T. G. Brain–Gut–Microbe Communication in

Health and Disease. *Front. Physiol.* **2**, (2011).

3. Arumugam, M. *et al.* Enterotypes of the human gut microbiome. *Nature* **473**, 174–180

(2011).

4. Tett, A. *et al.* The Prevotella copri complex comprises four distinct clades that are

underrepresented in Westernised populations. (2019) doi:10.1101/600593.

5. Vatanen, T. *et al.* The human gut microbiome in early-onset type 1 diabetes from the

TEDDY study. *Nature* **562**, 589–594 (2018).

6. Silbereis, J. C., Pochareddy, S., Zhu, Y., Li, M. & Sestan, N. The cellular and molecular

landscapes of the developing human central nervous system. *Neuron* **89**, 248–268 (2016).

7. Mullen, E. M. & others. *Mullen scales of early learning*. (AGS Circle Pines, MN, 1995).

8. Wechsler, D. Wechsler intelligence scale for children. (1949).

9. McArdle, B. H. & Anderson, M. J. Fitting Multivariate Models to Community Data: A

Comment on Distance-Based Redundancy Analysis. *Ecology* **82**, 290–297 (2001).

10. Anderson, M. J. Permutational Multivariate Analysis of Variance (PERMANOVA). in *Wiley*

StatsRef: Statistics Reference Online 1–15 (American Cancer Society, 2017).

doi:10.1002/9781118445112.stat07841.

11. Clarke, G. *et al.* The microbiome-gut-brain axis during early life regulates the hippocampal

serotonergic system in a sex-dependent manner. *Mol. Psychiatry* **18**, 666–673 (2013).

12. Hsiao, E. Y. *et al.* Microbiota modulate behavioral and physiological abnormalities

associated with neurodevelopmental disorders. *Cell* **155**, 1451–1463 (2013).

13. Blacher, E. *et al.* Potential roles of gut microbiome and metabolites in modulating ALS in

mice. *Nature* **572**, 474–480 (2019).

14. Gao, W. *et al.* Gut microbiome and brain functional connectivity in infants-a preliminary

study focusing on the amygdala. *Psychopharmacology (Berl.)* (2019) doi:10.1007/s00213-

018-5161-8.

15. Valles-Colomer, M. *et al.* The neuroactive potential of the human gut microbiota in quality of

life and depression. *Nat. Microbiol.* **4**, 623–632 (2019).

16. de Leeuw, C. A., Neale, B. M., Heskes, T. & Posthuma, D. The statistical properties of

gene-set analysis. *Nat. Rev. Genet.* **17**, 353–364 (2016).

17. Metwally, A. A., Aldirawi, H. & Yang, J. A review on probabilistic models used in microbiome

studies. (2018).

18. Reeds, P. J., Burrin, D. G., Stoll, B. & Jahoor, F. Intestinal Glutamate Metabolism. *J. Nutr.*

130, 978S-982S (2000).

19. Hoshino, M. Molecular mechanisms underlying glutamatergic vs. GABAergic neuronal

subtype specification in the cerebellum. *Seikagaku* **78**, 130 (2006).

20. Carletti, B. & Rossi, F. Neurogenesis in the cerebellum. *The Neuroscientist* **14**, 91–100

(2008).

21. Leto, K. *et al.* Consensus paper: cerebellar development. *The Cerebellum* **15**, 789–828

(2016).

22. Tiemeier, H. *et al.* Cerebellum development during childhood and adolescence: a

longitudinal morphometric MRI study. *Neuroimage* **49**, 63–70 (2010).

23. Wang, S. SH, Kloth, AD, & Badura, A.(2014). *Cerebellum Sensitive Periods Autism Neuron*

83, 518–532.

24. Fatemi, S. H. *et al.* Consensus paper: pathological role of the cerebellum in autism. *The*

Cerebellum **11**, 777–807 (2012).

25. Bonham, K. S. *et al.* Data for 'Gut microbial genes are associated with neurocognition and

brain development in healthy children'. (2020) doi:10.5281/zenodo.3633793.

26. Kevin Bonham. *Klepac-Ceraj-Lab/resonance_paper1: Zenodo Release*. (Zenodo, 2020).

doi:10.5281/zenodo.3648677.

27. McIver, L. J. *et al.* bioBakery: a meta'omic analysis environment. *Bioinforma. Oxf. Engl.* **34**,

1235–1237 (2018).

28. Bayley, N. *Bayley scales of infant and toddler development*. (PsychCorp, Pearson, 2006).

29. Wechsler, D. Wechsler preschool and primary scale of intelligence—fourth edition. *Psychol.*

Corp. San Antonio TX (2012).

Figure 1: The gut microbiome of children is associated with neurocognition.

a, Principal coordinates analysis (PCoA) for all subjects based on species-level Bray-Curtis dissimilarity; most variation is driven by age and the presence of Prevotella[3,4]. **b**, PCoA based on Bray-Curtis dissimilarity in functional profiles (UniRef90 accessory genes) after removing gene families that were present in >90% of subjects in a given age group; variation is driven by similar effects as for taxonomic profiles. **c**, First principle coordinate axis of species PCoA for child samples vs. age; younger children cluster away from older children, and are lower in diversity. **d**, First principle coordinate axis of accessory gene family PCoA for child samples vs. age; functional diversity increases with age. **e**, PERMANOVA analysis for selected subject metadata vs. pairwiase Bray-Curtis dissimilarity for species-level taxonomic or functional profiles. Functional profiles include Kegg-Orthology (KO), Pfams or accessory UniRef90s; subject and subject type include all subjects, 2+ includes only children over 2 years old, others include all children for which the measure was available; stars indicate significance after Benjamini-Hochberg FDR correction. (* <0.1, ** < 0.01, *** < 0.001) **f**, Child age vs. normalized cognitive function score; age-appropriate assessments achieve similar score distributions for their age groups. **g**, FSEA analysis for gene sets with neuroactive potential (see Methods).