# Poor availability of context-specific evidence hampers decision-making in conservation

Authors:
Alec P. Christie[1] (Corresponding author: apc58@cam.ac.uk),
Tatsuya Amano[1,3] (t.amano@uq.edu.au),
Philip A. Martin[1,2] (pam79@cam.ac.uk),
Silviu O. Petrovan[1] (sop21@cam.ac.uk),
Gorm E. Shackelford[2] (ges47@hermes.cam.ac.uk),
Benno I. Simmons[1,4,5] (benno.simmons@gmail.com),
Rebecca K. Smith[1] (r.k.smith@zoo.cam.ac.uk),
David R. Williams[6] (d.r.williams@leeds.ac.uk),
Claire F. R. Wordley[1] (clairefrwordley@gmail.com) and
William J. Sutherland[1,2] (w.sutherland@zoo.cam.ac.uk).

Affiliations:
[1]Conservation Science Group, Department of Zoology, University of Cambridge, The David Attenborough Building, Downing Street, Cambridge CB3 3QZ, UK.
[2]BioRISC, St. Catharine's College, Cambridge CB2 1RL, UK.
[3]School of Biological Sciences, University of Queensland, Brisbane, 4072 Queensland, Australia.
[4]Department of Animal and Plant Sciences, University of Sheffield, Sheffield, S10 2TN United Kingdom.
[5]Centre for Ecology and Conservation, College of Life and Environmental Sciences, University of Exeter, Penryn, UK
[6]Sustainability Research Institute, School of Earth and Environment, University of Leeds. LS2 9JT United Kingdom

Running title: Lack of relevant evidence limits conservation

Keywords: conservation evidence, conservation intervention, evidence-based conservation, external validity, generalizability, local context, prioritization, relevant evidence, study design, synthesis

Article type: Review

Abstract length: 200 words
Manuscript length: 4857 words
Number of references: 70
Number of figures and tables: 4

Corresponding author: Alec P. Christie, apc58@cam.ac.uk, 07955184785, Conservation Science Group, Department of Zoology, University of Cambridge, The David Attenborough Building, Downing Street, Cambridge CB3 3QZ, UK

Abstract

Evidence-based conservation relies on robust and relevant evidence. Practitioners often prefer locally relevant studies whose results are more likely to be transferable to the context of planned conservation interventions. To quantify the availability of relevant evidence for amphibian and bird conservation we reviewed Conservation Evidence, a database of quantitative tests of conservation interventions. Studies were geographically clustered and found at extremely low densities - fewer than one study was present within a 2,000 km radius of a given location. The availability of relevant evidence was extremely low when we restricted studies to those studying biomes or taxonomic orders containing high percentages of threatened species, compared to the most frequently studied biomes and taxonomic orders. Further constraining the evidence by study design showed that only 17-20% of amphibian and bird studies used robust designs. Our results highlight the paucity of evidence on the effectiveness of conservation interventions, and the disparity in evidence for local contexts that are frequently studied and those where conservation needs are greatest. Addressing the serious global shortfall in context-specific evidence requires a step change in the frequency of testing conservation interventions, greater use of robust study designs and standardized metrics, and methodological advances to analyze patchy evidence bases.

# Introduction

Tackling the biodiversity crisis with limited resources requires efficient and effective
conservation action (Dirzo et al., 2014; Sutherland, Pullin, Dolman, & Knight, 2004). To inform
which conservation actions ('interventions') are effective and which are not, we need a large and
robust evidence base, ideally including large numbers of studies (replication of evidence;
Fig.1A) with high internal validity (quality; Fig.1A) and external validity (relevance; Fig.1A).
However, the limited resources available for conservation research mean that the evidence
base for conservation is geographically and taxonomically biased (Christie, Amano, Martin,
Petrovan, et al., 2019; Fazey, Fischer, & Lindenmayer, 2005; Hickisch et al., 2019; Spooner,
Smith, & Sutherland, 2015). This is likely to limit the quality and relevance of evidence and
impair effective decision-making (Cook, Possingham, & Fuller, 2013). Quantifying the availability
of relevant, reliable studies is necessary to understand the strength of evidence upon which
decisions are made, and to prioritize research on the effectiveness of conservation
interventions.

Practitioners and policymakers typically prefer to base their decisions on studies that are
relevant (i.e., with high external validity; Fig.1) to their local context (Addison, Cook, & de Bie,
2016; Cook & Sgrò, 2017; Geijzendorffer et al., 2017). Using context-specific studies as
evidence helps to ensure that results are likely to be repeated if the intervention is implemented
again. The relevance of conservation studies to a given context will span multiple dimensions,
including: (i) bioclimatic (i.e., similarity between habitats or regions); (ii) taxonomic/functional
(i.e., similarity between taxa in terms of ecological function or taxonomic groups); and (iii) which
metric was used to quantify the effectiveness of an intervention (i.e., the response variables or
metrics of interest; Fig.1B). Other dimensions may also be important, such as the similarity
between a study's and a practitioner's socioeconomic and political contexts, but we focus on the
three dimensions above.

Figure 1 - Framework of the desirable aspects of an ideal evidence base (stronger colors = more desirable). Fig.1A shows the three major desirable factors that an evidence base should have; large replication of evidence that is highly reliable (high internal validity) and highly relevant (high external validity). Fig.1B refers to the three dimensions that we will focus on that influence the overall relevance of evidence: i) bioclimatic (e.g., the study system), ii) taxonomic/functional (the study taxa) and iii) effectiveness measure (how you define and measure conservation success).

The first of these dimensions - bioclimatic relevance - refers to the similarity between the study ecosystem and the practitioner's ecosystem (Fig.1B). The second dimension - taxonomic/functional relevance - concerns the similarity between the focal taxa of a study and the taxa of interest to the practitioner (Fig.1B). Together, these determine the ecological similarity between study and practitioner local contexts. This is vital because responses to interventions will vary between ecosystems and taxa. For example, the effectiveness of artificial

nest boxes varies between different countries and habitats (Finch et al., 2019), while the
effectiveness of translocation for New Zealand robins (*Petroica australis*) is unlikely to be
relevant to a practitioner translocating Kakapo (*Strigops habroptila*). Practitioners who are
interested in broader functional groups (e.g., seed dispersers or pollinators), taxa (e.g., birds,
amphibians), or even whole ecosystems, may focus more on the functional relevance rather
than taxonomic similarity of studied species.

The third dimension of relevance is the metric used to measure the effectiveness of an
intervention. Practitioners may be interested in different responses to interventions depending
on their focus (e.g., species or ecosystem-level responses) and effectiveness may vary
depending on the metric used (Capmourteres & Anand, 2016; Marshall, Wintle, Southwell, &
Kujala, 2019). For example, at the ecosystem-level, the effectiveness of bird boxes may be
measured using the species richness or diversity of birds using them (Caine & Marion, 1991),
while at the species-level, the number of individuals (Brawn & Balda, 1988), fledglings (Male,
Jones, & Robertson, 2006; Purcell, Verner, & Lewis W, 1997), or brood size (Browne, 2006)
may be measured. Similarly, the effectiveness of road mitigation interventions (e.g., tunnels or
bridges) may be measured by the numbers of individuals of different species using the
structures, but could also be measured in terms of levels of road mortality (Helldin & Petrovan,
2019). Therefore, the type of metric used by studies to measure effectiveness can have a major
influence on the relevance of evidence.

The reliability of an evidence base - the internal validity of its studies - ultimately determines the
overall quality of the evidence base and depends to a large extent on study design (Christie,
Amano, Martin, Shackelford, et al., 2019; De Palma et al., 2018; Spake & Doncaster, 2017). As
the conservation evidence base contains a wide variety of study designs (De Palma et al.,
2018), there is likely to be variation in the reliability of inferences that can be drawn (Christie,
Amano, Martin, Shackelford, et al., 2019). This variation may lead scientists to make misleading
recommendations to practitioners, ultimately reducing the effectiveness of conservation
practice, and making it difficult for decision-makers to weigh the strength of evidence provided
by different studies.

The replication of evidence - the number of studies in the evidence base - is also important as
greater numbers of studies demonstrating repeatable and reproducible effectiveness will give us
greater confidence in the overall strength of the evidence. Decision-makers should rightly be
wary of basing decisions on a low number of studies where reproducible effectiveness has not
been or cannot be demonstrated - particularly given the current reproducibility crisis (Begley &
Ioannidis, 2015; Nosek & Errington, 2017; Open Science Collaboration, 2015). However, the
overall number of studies is not the only indicator of the strength of the evidence, since studies
with low internal validity (e.g., poor study designs) and/or external validity (i.e., low relevance)
may not constitute reliable evidence. Currently, we have a poor quantitative understanding of
the availability of relevant and reliable studies in the conservation literature.

In this study, we assess whether studies testing conservation interventions are distributed
across different contexts (bioclimatically, taxonomically, and by the metric used to measure
effectiveness) in ways that reflect the needs of conservation. We also quantify other desirable

aspects of the evidence base for conservation in terms of the quantity and quality of available
studies; i.e., the number of studies that have tested different conservation actions, and how
many of these use robust study designs.

# Methods
171

### Conservation Evidence database
173

We assessed the availability of relevant evidence for conservation practice using Conservation
Evidence, a database of 5,525 publications as of January 2020 (Conservation Evidence, 2020a)
that have quantitatively assessed the effectiveness of conservation interventions. Interventions
are defined as management actions that a practitioner may undertake to benefit biodiversity
(see Sutherland et al. (2019) for detailed methods). When we refer to the number of studies per
intervention, we refer to the number of different tests of interventions - single publications may
report multiple tests of different interventions. We assessed the availability of evidence for
amphibians and birds based on synopses compiled in 2014 (n=419 studies; Smith &
Sutherland, 2014) and 2012 (n=1,232 studies; Williams et al., 2013), respectively. More recent
publications will obviously have increased the evidence base, but the broad patterns we quantify
are unlikely to have changed in the intervening years. We excluded meta-analyses or
systematic reviews from our analyses as these typically cannot be attributed to a particular local
context (e.g., biome or taxon). We also only included interventions for which studies were
present in the database. Since 32% (n=33) of interventions for amphibians and 25% (n=80) of
interventions for birds had no associated studies in the database (i.e., were untested or tests
were unpublished) or only included reviews or meta-analyses, the following analyses are likely
to be an optimistic assessment of the availability of evidence in conservation. We used R
statistical software version 3.5.1 (R Core Team, 2019) for all analyses.

### Local availability of studies by geographical distance
195

To calculate the average availability of studies within a certain distance of a given practitioner's
location, we generated 1,000 regularly spaced coordinates across certain parts of the world. For
amphibians, we spaced these coordinates over the combined extent of all amphibian species
ranges (IUCN, 2019) as this represents the possible range of locations in which a practitioner
might conduct an intervention to conserve amphibians. For birds, we spaced these coordinates
across the world's terrestrial land masses (using "OpenStreetMap" 2019; see Appendix S1 for
maps of coordinates) since although the combined distribution of all bird species is almost
global, most practitioners are likely to conduct interventions to conserve birds terrestrially.
Although non-terrestrial interventions are carried out by practitioners, the vast area covered by
the ocean would severely underestimate the availability of studies to a practitioner's likely
location. 19 non-terrestrial interventions for birds were found in the database (e.g., 'use
streamer lines to reduce seabird bycatch on longlines' or 'use high-visibility mesh on gillnets to
reduce seabird bycatch') containing 33 studies in total - these were still included in our analysis

as these studies tended to be conducted within close proximity to a terrestrial landmass (i.e.,
coastal).

We then calculated the Great Circle Distance from each study to each coordinate (see Appendix
S1 for details), binning distances into a series of categories (100 km, 1,000 km and then every
1,000 km up to and including 19,000 km). We also calculated the 'Global Mean', which is the
mean number of studies per intervention in the entire database - equivalent to approximately
20,000 km at the equator, the maximum distance separating any two coordinates. We then
calculated the mean number of studies within each distance bin across all coordinates, as well
as the number of studies that used different categories of study designs: i) any design, ii)
Before-After (BA), Control-Impact (CI), Before-After Control-Impact (BACI) or Randomized
Controlled Trial (RCT); iii) CI, BACI or RCT; iv) BACI or RCT designs (see Methods in Christie,
Amano, Martin, Petrovan, et al. 2019 for definitions of each design).

We then repeated this analysis using the same number of coordinates (n=1,000), but this time
by randomly selecting coordinates from amphibian and bird studies in the database (sampling
with replacement from amphibian studies as there were fewer than 1,000). Using both
approaches provided likely upper and lower bounds of evidence availability: regular coordinates
likely underestimated the availability of evidence to practitioners, giving equal weighting to
locations where conservation interventions are unlikely to occur (e.g., Antarctica) and those that
are more intensively managed (e.g., Europe). In contrast, using locations from existing
publications will likely overestimate study availability as this assumes that practitioners only
conduct interventions in locations where they have previously been tested.

We compared the results of the first analysis (regularly spaced coordinates) to the expected
patterns we would observe if studies were regularly distributed. We did this by generating equal
numbers of regularly spaced coordinates ('expected studies') as the number of amphibian and
bird studies (419 and 1,232 coordinates, respectively) using the same methods and shapefiles
as before. We then calculated the mean number of these 'expected studies' within each
distance bin.

**Context-specific availability of studies**

To quantify the amount of relevant and robust evidence on the effectiveness of different
conservation interventions, we required metadata that described each study's local context and
study design. By adapting previously described methods (Christie, Amano, Martin, Petrovan, et
al. 2019; Appendix S2), we extracted the biome, taxonomic order and reported metric type used
by each study (to quantify the number of relevant studies), as well as the broad category of
study design used (to quantify the number of robustly designed studies). When metric metadata
was extracted, we grouped similar metrics into the following nine metric types: count-based,
diversity, activity-based, physiological, survival, reproductive success, education-based,
regulation-based, and biomass (Appendix S2).

We quantified the number of studies per conservation intervention that met certain relevance
and study design criteria, to give an estimate of the availability of relevant and robust evidence.
To ensure that we did not artificially constrain the number of studies per intervention for different
subsets of studies (e.g., taxonomic order or biome), we grouped certain interventions that were
focused on single taxa or habitats but were fundamentally the same type of intervention (e.g.,
'create ponds for newts' and 'create ponds for toads' would be grouped into 'create ponds'; see
Acknowledgements and Data for files describing these groupings). This resulted in a total of 71
and 226 interventions for amphibians and birds, respectively.

Using these interventions, we then undertook two analyses to quantify the availability of
evidence under different scenarios: i) where we optimistically assume a given practitioner is
interested in the most frequently studied local context; and ii) where we assume that a given
practitioner is interested in local contexts in which a greater percentage of species are
threatened (i.e., those classified as Vulnerable, Endangered or Critically Endangered status on
the (IUCN, 2019) Red List).

The first analysis calculated the mean number of studies per intervention for both scenarios in
terms of three separate relevance criteria: biome, taxonomic order and metric. For the first
scenario we calculated the number of studies with the most frequently studied biome, order or
metric relative to each intervention. For the second scenario (to reflect conservation needs), we
calculated the number of studies with a randomly selected biome, taxonomic order or metric
from a weighted list (averaged over 1,000 repeated runs). This weighted list was generated so
that the probability of selection was determined by the percentage of species that are
threatened (i.e., those classified as Vulnerable, Endangered or Critically Endangered status on
the (IUCN, 2019) Red List) for each biome and taxonomic order, and the percentage usage of
each metric within each intervention in the database. We intersected shapefiles from the (IUCN,
2019) Red List with shapefiles of the world's terrestrial biomes (Dinerstein et al., 2017) to
determine the proportion of threatened species in each biome. We assumed that interventions
could be tested by studies in any biome and on any taxonomic order - this will likely mean that
our estimates for the second scenario are underestimates of study availability, for example, as
certain interventions are unlikely to be conducted in certain biomes. However, we grouped
interventions so they were not defined as taxon or habitat-specific and used coarse criteria
(biome and taxonomic order) to limit this underestimation.

For the second analysis, we used a stepwise process to calculate the number of studies that
met one or more of the relevance criteria - only carrying forward studies if they met all previous
criteria. For example, considering the first scenario (most frequently studied context), we
counted the number of studies featuring the most frequently studied biome, then studies
featuring the most frequently studied biome AND taxonomic order, and then studies featuring
the most frequently studied biome AND taxonomic order AND metric. We also repeated this for
all possible orderings of biome, taxonomic order and metric (Fig.3 and Figs.S1-S5), as well as
for the second scenario (weighting towards biomes and taxonomic orders with greater
percentages of threatened species). Taxonomic orders could only be selected if at least one
species in that order was present in the previously selected biome - we determined which

orders were present in each biome by intersecting shapefiles from the (IUCN, 2019) Red List
with shapefiles of terrestrial biomes (Dinerstein et al., 2017). The same was true for biomes
when taxonomic order was the first relevance criteria to be selected (i.e., only biomes where
that taxonomic order is present could be selected). In the final step, we also calculated the
number of studies that used different categories of study designs (any design; BA, CI, BACI or
RCT; CI, BACI or RCT; BACI or RCT).
303
304
305 **Results**
306
We considered a total of 71 and 226 interventions for amphibians and birds (mean = 7.9 and 6.9
studies per intervention; Fig.2), respectively, that contained at least one study. Studies were not
evenly distributed geographically; the mean number of amphibian and bird studies per
intervention (black large circles in Fig.2) deviated, particularly for amphibians, from what we
would have expected if the same number of studies were regularly distributed (orange triangles
in Fig.2). On average, there was less than one study per intervention available within 2,000km
from a given regular point. When restricting analyses to more robust designs, the availability of
studies decreased substantially, with a higher proportion of amphibian studies using BA
designs, compared to birds, but a smaller proportion using CI (see drop-offs from orange to
blue, and blue to green lines, respectively; Fig.2).
317
When considering distance of studies to randomly selected study coordinates, the mean
number of studies per intervention generally declined more gradually compared to a regular grid
of coordinates (Fig.2), implying that studies are clustered in space. At distances below 5,000km
these differences were particularly pronounced; for example, on average, 2.2 amphibian studies
and 1.5 bird studies were within 2,000km of a random study coordinate, compared to only 0.3
amphibian studies and 0.2 bird studies within 2,000km of regularly spaced coordinate. This
suggests that studies are slightly more clustered for amphibians than birds.
325

Figure 2 - The mean number of amphibian and bird studies per intervention using different study designs found within a certain distance of different sets of coordinates. The maximum distance that a study can be is shown on the x axis, starting with the Global Mean (mean number of studies per intervention considering all studies in the database) and decreasing to a distance of 100 km. Regular coordinates (large circle, thick line) show the mean number of studies within a certain distance from a set of regularly distributed coordinates. Expected coordinates (orange triangle) mimic how the availability of studies would be expected to change if studies were regularly distributed (this is only shown for studies using any study design). Random Study coordinates (small circle, thin line) show the mean number of studies within a certain distance from a set of randomly selected coordinates where previous studies have been conducted.

The mean number of studies per intervention was substantially greater for the most frequently
studied biome (Amphibians: 5.0; Birds: 3.5), relative to each intervention, compared to biomes
with higher percentages of species that are threatened (Amphibians: 0.4; Birds: 0.4; Fig.3).
Similarly, the mean number of studies per intervention was substantially greater for the most
frequently studied order in each intervention (Amphibians: 7.2; Birds: 4.4), compared to a
taxonomic orders with higher percentages of species that are threatened (Amphibians: 0.4;
Birds: 0.01; Fig.3). There was a smaller difference in the mean number of studies per
intervention between studies that used the most frequently used metric (Amphibians: 5.2; Birds:
4.8), relative to each intervention, and studies that used a randomly selected metric from within
each intervention (Amphibians: 4.5; Birds: 3.9; Fig.3). The mean numbers of biomes, taxonomic
orders and metrics per intervention were 2.7, 2.6, and 3.1 for amphibians, respectively, and 2.4,
6.1, and 2.6 for birds, respectively.
352
353
354

355
Figure 3 - Mean number of studies per intervention when studies were counted based on
whether they considered the most frequently studied biome, metric or order, and whether they
considered a randomly selected biome, metric or taxonomic order from a weighted list. These
weightings were based on the proportion of threatened species found in each biome or
taxonomic order. 'All' indicates the mean number of studies per intervention when considering
all studies.
362
The mean number of studies per intervention was also greater when we constrained by the
most frequently studied biome, taxonomic order and metric in a stepwise process Fig.4A),
compared to biomes and taxonomic orders with higher percentages of threatened species
(Fig.4B). When we constrained by the most frequently studied biome, taxonomic order and
metric, the greatest proportional decrease in the number of studies occurred once we further
constrained by study design, by only counting studies using robust BACI or RCT designs (on

average, ~20% of amphibian studies and ~17% of bird studies that had met all previous criteria;
Fig.4A). When we constrained by biomes and taxonomic orders with higher percentages of
threatened species, the greatest proportional decreases occurred when constraining by
taxonomic order, most notably for birds, and by biome (Fig.4B).
373
The sequence in which criteria were applied did not substantially affect the magnitude of the
decrease in the number of studies - e.g., when biome was selected before or after taxonomic
order and metric (Supporting Information Fig.S1-5). The overall decrease in studies from
applying all relevance criteria (biome, taxonomic order and metric) was similarly severe
regardless of the sequence in which the criteria were applied (Supporting Information Fig.S1-5).
For all sequences, constraining the evidence to studies that used robust BACI or RCT designs
reduced the mean number of studies to less than one study after constraining by the most
frequently studied biome, taxonomic order and metric (Fig.4A; Supporting Information Fig.S1-5).
Doing the same after instead constraining by the biomes and taxonomic orders with higher
percentages of threatened species reduced the mean number of studies to fewer than 0.01
studies with BACI or RCT designs (Fig.4B; Supporting Information Fig.S1-5).

397
Figure 4 - Mean numbers of amphibian and bird studies per intervention when only considering
studies that meet certain relevance criteria. In panel A, studies with the most frequently studied
biome, taxonomic order and metric relative to each intervention were counted - here we assume
practitioners are interested in the most frequently studied local context. At each step (left to
right) we add a further criterion, carrying forward relevant studies from the previous step - for
example, only studies conducted in the most frequently studied biome were carried forward into
the biome and order category. In panel B, studies with a selected biome, taxon and metric were
counted (y axis has a square root transformation). Here we assume practitioners are more likely
to be interested in: biomes that are inhabited by higher proportions of threatened species;
taxonomic orders that have higher relative proportions of threatened species; and metrics that
are most frequently used within each intervention. At the final step, studies are counted based
on the study design they use (see Methods for details of study designs).

## Discussion

Our work demonstrates that not only is there a general paucity of studies testing conservation interventions, but that the distribution of these studies does not reflect conservation needs. Specifically, there is a lack of studies testing conservation interventions in biomes and for taxonomic orders containing high percentages of threatened amphibian and bird species. Given substantial declines of bird fauna (Rosenberg et al., 2019) and severe threats to amphibians (Grant, Muths, Schmidt, & Petrovan, 2019), a better understanding of the effectiveness of interventions targeting threatened species is urgently required. Furthermore, a given decision-maker is likely to struggle to find robust studies addressing their local context. Addressing this deficit will be challenging, but there are several possible ways to improve the evidence base for conservation.

A fundamental problem that needs to be overcome in the long-term is the lack of studies testing conservation interventions. Williams, Balmford, & Wilcove (*in review*) found that only 15% of studies from a representative sample of the conservation literature tested interventions. Evaluation of interventions should become mainstream, both as a topic of academic research and as an activity for on-the-ground conservationists (Baylis et al., 2016). The publication of these tests, whether the results are positive, negative, or neutral, is critical to building a strong evidence base for conservation (Catalano, Lyons-White, Mills, & Knight, 2019). Current efforts to facilitate this include the Applied Ecology Resources repository (British Ecological Society, 2020), 'Evidence' articles in the journal Conservation Science and Practice (Society for Conservation Biology, 2020), and the journal Conservation Evidence (Conservation Evidence, 2020b).

Simply publishing more tests of conservation interventions, even at an increasing rate, is however unlikely to solve the paucity of locally relevant studies. For example, even though adding 1,000 studies testing interventions on birds would increase the mean number of studies to approximately 11 studies across the current 226 interventions, these studies would still be spread thin across a myriad of local contexts where the need for conservation is often not the greatest (see also Wilson et al., 2016). Although Reboredo Segovia, Romano, & Armsworth (2020) suggest that the number of general conservation studies in tropical locations correlates with the number of threatened species, the results of this study and (Christie, Amano, Martin, Petrovan, et al., 2019) suggest this is not the case for conservation studies testing interventions. Therefore, we need concrete solutions enabling conservationists to generate and collate more experimental evidence on the effectiveness of conservation interventions in underrepresented locations and on underrepresented taxa (Christie, Amano, Martin, Petrovan, et al., 2019; Donaldson et al., 2016; Murray, Green, Williams, Burfield, & de Brooke, 2015). For example, funders, principal investigators and heads of conservation organizations need to enhance and prioritize funding to test interventions in underrepresented areas. Evidence synthesis also needs to incorporate more evidence from non-English language and grey literature publications to help address underrepresented local contexts (Amano, González-Varo, & Sutherland, 2016; Amano & Sutherland, 2013) - for example, publications from over 317 non-English language journals are starting to be added to the Conservation Evidence database through the Transcending

454 Language Barriers to Environmental Sciences project (TRANSLATE, 2020). Making concerted
455 efforts to acquire grey literature from organizations and groups outside academia will also be
456 important.
457
The low proportion of studies using robust study designs, regardless of their relevance to a local
context, is also challenging. That more robustly designed studies are concentrated in North
America, Europe and Australia also compounds earlier taxonomic and biogeographical biases
(Christie, Amano, Martin, Petrovan, et al., 2019). If few robustly designed studies are available
for informing conservation, decision-makers may have to consider a wider range of studies that
may be less robust or relevant, potentially reducing the effectiveness of decision-making and
future practice (Slavin, 1995; Tugwell & Haynes, 2006; Whittaker, 2010). To increase the quality
of studies available for decision-making, we must recognize that the quality of studies testing
interventions may be limited in different ways. Studies evaluating mitigation efforts are often not
constrained by cost, but rather by short timescales and their focus on meeting legislative
requirements (for example, conserving legally protected species). Studies testing non-mitigation
interventions will likely be more constrained by cost, as well as short timescales (e.g., PhD
funding). Acknowledging how real-world constraints affect the choice of study design is
essential to devising approaches to improving the evidence base for conservation. While better
training of early career scientists, consultants and researchers in appropriate study designs for
causal inference may help, ultimately more regulatory and funder-led measures (e.g., requiring
grantees to demonstrate rigorous study design) will be required (De Palma et al., 2018; Grant et
al., 2019).
476
Given the general lack of evidence across conservation, there is also a need to use a
standardized set of metrics to evaluate conservation effectiveness (McQuatters-Gollop et al.,
2019). Using a diversity of metrics may be necessary to assess multiple important aspects of an
intervention's effectiveness, but a lack of consistency in the metrics used to report results often
makes the evidence base difficult to synthesize - especially if different metrics yield different
results (Mace & Baillie, 2007). Prioritisation of the most relevant metrics of effectiveness for
different interventions with input from decision-makers and practitioners is essential to facilitate
inter-study comparisons (McQuatters-Gollop et al., 2019). Initiatives aiming to do this are
underway in topics such as fishery habitats (Lederhouse & Link, 2016) and protected areas
(Nolte & Agrawal, 2013; Pomeroy, Parks, & Watson, 2004), and are supported by the Essential
Biodiversity Variables framework (Jetz et al., 2019). Funders could help strengthen these efforts
by requiring grantees to follow such initiatives and use consistent metrics when evaluating
interventions.
490
Increasing the size and quality of the evidence base for conservation decision-making will be a
slow process, but conservation practitioners need to make decisions now. Until the evidence
base improves, excluding studies from evidence syntheses because they do not meet certain
quality or relevance criteria could lead to little or no evidence being used to inform conservation
efforts (Davies & Gray, 2015; Gurevitch & Hedges, 1999; Lortie, Stewart, Rothstein, & Lau,
2015). Moreover, studies that do not meet these criteria may still provide useful evidence,

particularly in the absence of more relevant and robust studies (Burivalova et al., 2019; Cook,
Mascia, Schwartz, Possingham, & Fuller, 2013; Gough & White, 2018).

Therefore, we need novel approaches to rigorously synthesizing studies that vary considerably
in their relevance and robustness to maximize the use of the current imperfect evidence base.
We believe that weighting approaches in both quantitative meta-analyses and more qualitative
evidence synthesis would help maximize the number of studies available, while giving greater
influence to studies with desirable characteristics. This could involve giving greater influence to
more robustly designed studies (e.g., using accuracy weights from Christie, Amano, Martin,
Shackelford, et al. 2019 and evidence hierarchies from Mupepele, Walsh, Sutherland, &
Dormann 2016), and giving more weight to more relevant studies (e.g., weighting by the
relevance of studies to a decision-maker's local context, as proposed in healthcare by Kneale,
Thomas, O'Mara-Eves, & Wiggins 2019). To generate objective weights of study relevance that
reflect the likely generalizability of study results, we need studies which help us to understand
how generalizability varies between interventions for different ecological (e.g., artificial nest
boxes; Finch et al. 2019), socioeconomic, and political contexts. Understanding why some
interventions work in certain contexts and not others is fundamentally important for evidence-
based decision-makers (Grant et al., 2019).

Overall, we have shown that the evidence base for conservation does not reflect the needs of
conservation. When this is combined with the general paucity of robust studies testing
conservation interventions, we conclude that there is a serious lack of locally relevant and
robust studies to inform decision-making in conservation. We hope that the conservation
community can work together to improve the state of the conservation evidence base. Doing so
will require much greater collaboration between research and practice. Testing interventions
needs to become more routine, use a more standardized suite of metrics and robust study
designs, and, most importantly, focus on the locations and taxa where evidence is most needed
to inform conservation action. In the meantime, we need to explore ways to better analyze the
current patchy evidence base of conservation and ensure that we can support the shift towards
more evidence-based policy and practice.

## Acknowledgements and Data

529 We would like to thank Anne Mupepele for their useful comments on the manuscript and all past
530 and present members of the Conservation Evidence project. All data analyzed in this study and
531 code to repeat analyses are available from https://doi.org/10.5281/zenodo.3634780.

## Author funding sources

534 TA was supported by the Grantham Foundation for the Protection of the Environment, the
535 Kenneth Miller Trust and the Australian Research Council Future Fellowship (FT180100354);
536 WJS, PAM, CFRW, SOP and GES are supported by Arcadia and The David and Claudia

Harding Foundation; RKS was supported by the MAVA Foundation; BIS and APC were supported by the Natural Environment Research Council as part of the Cambridge Earth System Science NERC DTP [NE/L002507/1]. BIS is also supported by the Natural Environment Research Council [NE/S001395/1] and a Royal Commission for the Exhibition of 1851 Research Fellowship.

## References

1. Addison, P. F. E., Cook, C. N., & de Bie, K. (2016). Conservation practitioners' perspectives on decision triggers for evidence-based management. *Journal of Applied Ecology*, *53*(5), 1351–1357. https://doi.org/10.1111/1365-2664.12734
2. Amano, T., González-Varo, J. P., & Sutherland, W. J. (2016). Languages Are Still a Major Barrier to Global Science. *PLOS Biology*, *14*(12), e2000933. Retrieved from https://doi.org/10.1371/journal.pbio.2000933
3. Amano, T., & Sutherland, W. J. (2013). Four barriers to the global understanding of biodiversity conservation: wealth, language, geographical location and security. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1756), 20122649.
4. Baylis, K., Honey-ros, J., Corbera, E., Ezzine-de-blas, D., Ferraro, P. J., Jan, B., … Wunder, S. (2016). *Mainstreaming Impact Evaluation in Nature Conservation 3. 9*(February), 58–64. https://doi.org/10.1111/conl.12180
5. Begley, C. G., & Ioannidis, J. P. A. (2015). *Review Reproducibility in Science Improving the Standard for Basic and Preclinical Research*. 116–126. https://doi.org/10.1161/CIRCRESAHA.114.303819
6. Bivand, R., Keitt, T., & Rowlingson, B. (2019). *rgdal: Bindings for the "Geospatial" Data Abstraction Library R package version 1.4-8*. Retrieved from https://cran.r-project.org/package=rgdal
7. Bivand, R., & Rundel, C. (2019). *rgeos: Interface to Geometry Engine - Open Source ('GEOS'). R package version 0.5-2*. Retrieved from https://cran.r-project.org/package=rgeos%0A
8. Bivand, R. S., Pebesma, E., & Gomez-Rubio, V. (2013). *Applied spatial data analysis with R* (Second). Retrieved from http://www.asdar-book.org/
9. Brawn, J. D., & Balda, R. P. (1988). Population biology of cavity nesters in northern Arizona: do nest sites limit breeding densities? *The Condor*, *90*(1), 61–71.
10. British Ecological Society. (2020). Applied Ecological Resources repository and Ecological Solutions and Evidence journal. Retrieved January 10, 2020, from https://besjournals.onlinelibrary.wiley.com/journal/26888319
11. Browne, S. J. (2006). Effect of nestbox construction and colour on the occupancy and breeding success of nesting tits Parus spp. *Bird Study*, *53*(2), 187–192.
12. Burivalova, Z., Allnutt, T., Rademacher, D., Schlemm, A., Wilcove, D. S., & Butler, R. A. (2019). What works in tropical forest conservation, and what does not: Effectiveness of four strategies in terms of environmental, social, and economic outcomes. *Conservation Science and Practice*, *in press*(March), 1–15. https://doi.org/10.1111/csp2.28

13. Caine, L. A., & Marion, W. R. (1991). Artificial Addition of Snags and Nest Boxes to Slash Pine Plantations (Colocacion de maderos y cajas de anidamiento en plantaciones de Pinus elliottii). *Journal of Field Ornithology*, 97–106.

14. Capmourteres, V., & Anand, M. (2016). "Conservation value": a review of the concept and its quantification. *Ecosphere*, *7*(10), e01476. https://doi.org/10.1002/ecs2.1476

15. Catalano, A. S., Lyons-White, J., Mills, M. M., & Knight, A. T. (2019). Learning from published project failures in conservation. *Biological Conservation*, *238*, 108223. https://doi.org/https://doi.org/10.1016/j.biocon.2019.108223

16. Christie, A. P., Amano, T., Martin, P. A., Petrovan, S. O., Shackelford, G. E., Simmons, B. I., … Sutherland, W. J. (2019). The challenge of heterogeneous evidence in conservation. *BioRxiv*, 797639. https://doi.org/10.1101/797639

17. Christie, A. P., Amano, T., Martin, P. A., Shackelford, G. E., Simmons, B. I., & Sutherland, W. J. (2019). Simple study designs in ecology produce inaccurate estimates of biodiversity responses. *Journal of Applied Ecology*, *56*(12), 2742–2754. https://doi.org/10.1111/1365-2664.13499

18. Conservation Evidence. (2020a). Conservation Evidence. Retrieved February 4, 2020, from www.conservationevidence.com

19. Conservation Evidence. (2020b). Conservation Evidence journal. Retrieved from https://www.conservationevidence.com/collection/view

20. Cook, C. N., Mascia, M. B., Schwartz, M. W., Possingham, H. P., & Fuller, R. A. (2013). Achieving Conservation Science that Bridges the Knowledge–Action Boundary. *Conservation Biology*, *27*(4), 669–678. https://doi.org/10.1111/cobi.12050

21. Cook, C. N., Possingham, H. P., & Fuller, R. A. (2013). Contribution of Systematic Reviews to Management Decisions. *Conservation Biology*, *27*(5), 902–915. https://doi.org/10.1111/cobi.12114

22. Cook, C. N., & Sgrò, C. M. (2017). Aligning science and policy to achieve evolutionarily enlightened conservation. *Conservation Biology*, *31*(3), 501–512. https://doi.org/10.1111/cobi.12863

23. Davies, G. M., & Gray, A. (2015). Don't let spurious accusations of pseudoreplication limit our ability to learn from natural experiments (and other messy kinds of ecological monitoring). *Ecology and Evolution*, *5*(22), 5295–5304. https://doi.org/10.1002/ece3.1782

24. De Palma, A., Sanchez-Ortiz, K., Martin, P. A., Chadwick, A., Gilbert, G., Bates, A. E., … Purvis, A. (2018). Challenges With Inferring How Land-Use Affects Terrestrial Biodiversity: Study Design, Time, Space and Synthesis. In *Next Generation Biomonitoring: Part 1* (1st ed., pp. 163–199). Elsevier Ltd.

25. Dinerstein, E., Olson, D., Joshi, A., Vynne, C., Burgess, N. D., Wikramanayake, E., … Saleem, M. (2017). An Ecoregion-Based Approach to Protecting Half the Terrestrial Realm. *BioScience*, *67*(6), 534–545. https://doi.org/10.1093/biosci/bix014

26. Dirzo, R., Young, H. S., Galetti, M., Ceballos, G., Isaac, N. J. B., & Collen, B. (2014). Defaunation in the Anthropocene. *Science*, *345*(6195), 401–406. https://doi.org/10.1126/science.1251817

27. Donaldson, M. R., Burnett, N. J., Braun, D. C., Suski, C. D., Hinch, S. G., Cooke,

S. J., & Kerr, J. T. (2016). Taxonomic bias and international biodiversity
conservation research. *FACETS*, *1*(1), 105–113. https://doi.org/10.1139/facets-
2016-0011

28. Fazey, I., Fischer, J., & Lindenmayer, D. B. (2005). What do conservation
biologists publish? *Biological Conservation*, *124*(1), 63–73.
https://doi.org/https://doi.org/10.1016/j.biocon.2005.01.013

29. Finch, T., Branston, C., Clewlow, H., Dunning, J., Franco, A. M. A., Račinskis, E.,
… Butler, S. J. (2019). Context-dependent conservation of the cavity-nesting
European Roller. *Ibis*, *161*(3), 573–589. https://doi.org/10.1111/ibi.12650

30. Geijzendorffer, I. R., van Teeffelen, A. J. A., Allison, H., Braun, D., Horgan, K.,
Iturrate-Garcia, M., … Quatrini, S. (2017). How can global conventions for
biodiversity and ecosystem services guide local conservation actions? *Current
Opinion in Environmental Sustainability*, *29*, 145–150.

31. Gough, D., & White, H. (2018). *Evidence standards and evidence claims in web
based research portals*. Retrieved from https://uploads-
ssl.webflow.com/59f07e67422cdf0001904c14/5bfffe39daf9c956d0815519_CFHI
_EVIDENCE_STANDARDS_REPORT_V14_WEB.pdf

32. Grant, E. H. C., Muths, E., Schmidt, B. R., & Petrovan, S. O. (2019). Amphibian
conservation in the Anthropocene. *Biological Conservation*, *236*, 543–547.
https://doi.org/https://doi.org/10.1016/j.biocon.2019.03.003

33. Gurevitch, J., & Hedges, L. V. (1999). Statistical issues in ecological
meta-analyses. *Ecology*, *80*(4), 1142–1149.

34. Helldin, J. O., & Petrovan, S. O. (2019). Effectiveness of small road tunnels and
fences in reducing amphibian roadkill and barrier effects at retrofitted roads in
Sweden. *PeerJ*, *7*, e7518. https://doi.org/10.7717/peerj.7518

35. Hickisch, R., Hodgetts, T., Johnson, P. J., Sillero-Zubiri, C., Tockner, K., &
Macdonald, D. W. (2019). Effects of publication bias on conservation planning.
Conservation Biology, 33(5), 1151–1163. https://doi.org/10.1111/cobi.13326

36. Hijmans, R. J. (2017). *geosphere: Spherical Trigonometry. R package version
1.5-7*. Retrieved from https://cran.r-project.org/package=geosphere

37. IUCN. (2019). IUCN Red List. Retrieved November 12, 2019, from
https://www.iucnredlist.org/

38. Jetz, W., McGeoch, M. A., Guralnick, R., Ferrier, S., Beck, J., Costello, M. J., …
Merow, C. (2019). Essential biodiversity variables for mapping and monitoring
species populations. *Nature Ecology & Evolution*, *3*(4), 539–551.

39. Kneale, D., Thomas, J., O'Mara-Eves, A., & Wiggins, R. (2019). How can
additional secondary data analysis of observational data enhance the
generalisability of meta-analytic evidence for local public health decision making?
*Research Synthesis Methods*, *10*(1), 44–56. https://doi.org/10.1002/jrsm.1320

40. Lederhouse, T., & Link, J. S. (2016). A Proposal for Fishery Habitat Conservation
Decision-Support Indicators. *Coastal Management*, *44*(3), 209–222.
https://doi.org/10.1080/08920753.2016.1163176

41. Lortie, C. J., Stewart, G., Rothstein, H., & Lau, J. (2015). How to critically read
ecological meta-analyses. *Research Synthesis Methods*, *6*(2), 124–133.
https://doi.org/10.1002/jrsm.1109

42. Mace, G. M., & Baillie, J. E. M. (2007). The 2010 biodiversity indicators:

challenges for science and policy. *Conservation Biology*, *21*(6), 1406–1413.

43. Male, S. K., Jones, J., & Robertson, R. J. (2006). Effects of nest-box density on the behavior of Tree Swallows during nest building. *Journal of Field Ornithology*, *77*(1), 61–66.

44. Marshall, E., Wintle, B. A., Southwell, D., & Kujala, H. (2019). What are we measuring? A review of metrics used to describe biodiversity in offsets exchanges. *Biological Conservation*, 108250. https://doi.org/https://doi.org/10.1016/j.biocon.2019.108250

45. McQuatters-Gollop, A., Mitchell, I., Vina-Herbon, C., Bedford, J., Addison, P. F. E., Lynam, C. P., … Otto, S. A. (2019). From Science to Evidence – How Biodiversity Indicators Can Be Used for Effective Marine Conservation Policy and Management  . *Frontiers in Marine Science*  , Vol. 6, p. 109. Retrieved from https://www.frontiersin.org/article/10.3389/fmars.2019.00109

46. Mupepele, A.-C., Walsh, J. C., Sutherland, W. J., & Dormann, C. F. (2016). An evidence assessment tool for ecosystem services and conservation studies. *Ecological Applications*, *26*(5), 1295–1301. https://doi.org/10.1890/15-0595

47. Murray, H. J., Green, E. J., Williams, D. R., Burfield, I. J., & de Brooke, M. L. (2015). Is research effort associated with the conservation status of European bird species? *Endangered Species Research*, *27*(3), 193–206. https://doi.org/10.3354/esr00656

48. Nolte, C., & Agrawal, A. (2013). Linking Management Effectiveness Indicators to Observed Effects of Protected Areas on Fire Occurrence in the Amazon Rainforest. *Conservation Biology*, *27*(1), 155–165. https://doi.org/10.1111/j.1523-1739.2012.01930.x

49. Nosek, B. A., & Errington, T. M. (2017). Making sense of replications. *ELife*, *6*, e23383. https://doi.org/10.7554/eLife.23383

50. Open Science Collaboration. (2015). Estimating the reproducibility of psychological science. *Science*, *349*(6251), aac4716. https://doi.org/10.1126/science.aac4716

51. OpenStreetMap. (2019). Retrieved December 14, 2019, from http://openstreetmapdata.com/data/land-polygons

52. Pebesma, E. J., & Bivand, R. S. (2005). Classes and methods for spatial data in R. *R News*, *5*(2).

53. Pomeroy, R. S., Parks, J. E., & Watson, L. M. (2004). *How is your MPA doing?: a guidebook of natural and social indicators for evaluating marine protected area management effectiveness*. IUCN.

54. Purcell, K. L., Verner, J., & Lewis W, O. (1997). A comparison of the breeding ecology of birds nesting in boxes and tree cavities. *The Auk*, *114*(4), 646–656.

55. R Core Team. (2019). *R: A language and environment for statistical computing. R Foundation for Statistical Computing*. Retrieved from https://www.r-project.org/

56. Reboredo Segovia, A. L., Romano, D., & Armsworth, P. R. (2020). Who studies where? Boosting tropical conservation research where it is most needed. *Frontiers in Ecology and the Environment*, fee.2146. https://doi.org/10.1002/fee.2146

57. Rosenberg, K. V, Dokter, A. M., Blancher, P. J., Sauer, J. R., Smith, A. C., Smith, P. A., … Marra, P. P. (2019). Decline of the North American avifauna. *Science*,

720    eaaw1313. https://doi.org/10.1126/science.aaw1313

58. Slavin, R. E. (1995). Best evidence synthesis: An intelligent alternative to meta-analysis. *Journal of Clinical Epidemiology*, *48*(1), 9–18. https://doi.org/10.1016/0895-4356(94)00097-A

59. Smith, R. K., & Sutherland, W. J. (2014). *Amphibian conservation: global evidence for the effects of interventions* (Vol. 4). Pelagic Publishing Ltd.

60. Society for Conservation Biology. (2020). Conservation Science and Practice. Retrieved January 10, 2020, from https://conbio.onlinelibrary.wiley.com/journal/25784854

61. Spake, R., & Doncaster, C. P. (2017). Use of meta-analysis in forest biodiversity research: key challenges and considerations. *Forest Ecology and Management*, *400*, 429–437. https://doi.org/10.1016/j.foreco.2017.05.059

62. Spooner, F., Smith, R. K., & Sutherland, W. J. (2015). Trends, biases and effectiveness in reported conservation interventions. *Conservation Evidence*, *12*, 2–7.

63. Sutherland, W. J., Pullin, A. S., Dolman, P. M., & Knight, T. M. (2004). The need for evidence-based conservation. *Trends in Ecology & Evolution*, *19*(6), 305–308.

64. Sutherland, W. J., Taylor, N. G., MacFarlane, D., Amano, T., Christie, A. P., Dicks, L. V, … Wordley, C. F. R. (2019). Building a tool to overcome barriers in research-implementation spaces: The Conservation Evidence database. *Biological Conservation*, *238*, 108199. https://doi.org/10.1016/j.biocon.2019.108199

65. TRANSLATE. (2020). TRANSLATE - Transcending Language Barriers to Environmental Sciences. Retrieved January 11, 2020, from https://researchers.uq.edu.au/research-project/35572

66. Tugwell, P., & Haynes, R. B. (2006). Assessing claims of causation. *Clinical Epidemiology: How to Do Clinical Practice Research*, 356–387.

67. Whittaker, R. J. (2010). Meta-analyses and mega-mistakes: calling time on meta-analysis of the species richness–productivity relationship. *Ecology*, *91*(9), 2522–2533.

68. Williams, D. R., Balmford, A., & Wilcove, D. S. (n.d.). The past and future role of conservation science in protecting biodiversity. In *Conservation Letters*.

69. Williams, D. R., Pople, R. G., Showler, D. A., Dicks, L. V, Child, M. F., Zu Ermgassen, E. K. H. J., & Sutherland, W. J. (2013). *Bird Conservation: Global evidence for the effects of interventions* (Vol. 2). Pelagic Publishing.

70. Wilson, K. A., Auerbach, N. A., Sam, K., Magini, A. G., Moss, A. S. L., Langhans, S. D., … Meijaard, E. (2016). Conservation Research Is Not Happening Where It Is Most Needed. *PLOS Biology*, *14*(3), e1002413. Retrieved from https://doi.org/10.1371/journal.pbio.1002413