Formicine ants swallow their highly acidic poison for gut microbial selection and control

Simon Tragust[1†*], Claudia Herrmann[1], Jane Häfner[1], Ronja Braasch[1], Christina Tilgen[1], Maria Hoock[1], Margarita Artemis Milidakis[1], Roy Gross[2], Heike Feldhaar[1]

[1] Animal Ecology I, Bayreuth Center for Ecology and Environmental Research (BayCEER), University of Bayreuth, Universitätsstraße 30, 95447 Bayreuth, Germany

[2] Microbiology, Biocenter, University of Würzburg, Am Hubland, 97074 Würzburg

[†] Present address: General Zoology, Hoher Weg 8, Martin-Luther University, 06120 Halle (Saale), Germany

[*] Correspondence to: simon.tragust@zoologie.uni-halle.de

Abstract

Animals continuously encounter microorganisms that are essential for health or cause disease. They are thus challenged to control harmful microbes while allowing acquisition of beneficial microbes. This challenge is likely especially important for social insects with respect to microbes in food, as they often store food and exchange food among colony members. Here we show that formicine ants actively swallow their antimicrobial, highly acidic poison gland secretion. The ensuing acidic environment in the stomach, the crop, limits establishment of pathogenic and opportunistic microbes ingested with food and improves survival of ants when faced with pathogen contaminated food. At the same time, crop acidity selectively allows acquisition and colonization by Acetobacteraceae, known bacterial gut associates of formicine ants. This suggests that swallowing of the poison gland secretion acts as a microbial filter in formicine ants and indicates a potentially widespread but so far underappreciated dual role of antimicrobials in host-microbe interactions.

Introduction

Animals commonly harbor gut associated microbial communities (Engel and Moran, 2013, Moran et al., 2019). Patterns of recurring gut microbial communities have been described for many animal groups (Brune and Dietrich, 2015, Kwong et al., 2017, Ochman et al., 2010). The processes generating these patterns are however often not well understood. They might result from host filtering (Mazel et al., 2018), a shared evolutionary history between gut associated microbes and their hosts (Moeller et al., 2016) involving microbial adaptations to the host environment (McFall-Ngai et al., 2013), simply be a byproduct of similar host dietary preferences (Anderson et al., 2012, Hammer et al., 2017), or result from interactions between microbes in the gut associated microbial community (Brinker et al., 2019a, García-Bayona and Comstock, 2018).

Food is an important environmental source of microbial gut associates (Blum et al., 2013,

Broderick and Lemaitre, 2012, David et al., 2014, Hammer et al., 2017, Perez-Cobas et al.,

2015) but also poses a challenge, the need to discriminate between harmful and beneficial

microbes, as food may contain microbes that produce toxic chemicals or that are pathogenic

(Burkepile et al., 2006, Demain and Fang, 2000, Janzen, 1977, Trienens et al., 2010). In social

animals, control of harmful microbes in food while allowing the acquisition and transmission

of beneficial microbes from and with food, is likely especially important. Eusocial

Hymenoptera not only transport and store food in their stomach, the crop, but also distribute

food to members of their colony via trophallaxis, i.e. the regurgitation of crop content from

donor individuals to receiver individuals through mouth-to-mouth feeding (Gernat et al.,

2018, Greenwald et al., 2018, LeBoeuf et al., 2016). While trophallaxis can facilitate the

transmission of beneficial microbes, from an epidemiological perspective it can also entail

significant costs, as it might open the door to unwanted microbial opportunists and pathogens

that can take advantage of these transmission routes (Onchuru et al., 2018, Salem et al., 2015).

Here we investigate how formicine ants, specifically the Florida carpenter ant *Camponotus*

floridanus, solve the challenge to control harmful microbes in their food while allowing

acquisition and transmission of beneficial microbes from and with their food. Apart from

specialized intracellular endosymbionts associated with the midgut in the ant tribe

Camponotini (Degnan et al., 2004, Feldhaar et al., 2007, Russell et al., 2017, Williams and

Wernegreen, 2015), formicine ant species have only low abundances of microbial associates

in their gut lumen but carry members of the bacterial family Acetobacteraceae as a recurring

part of their gut microbiota (Brown and Wernegreen, 2016, Chua et al., 2018, He et al., 2011,

Ivens et al., 2018, Russell et al., 2017). Some formicine gut associated Acetobacteraceae show

signs of genomic and metabolic adaptations to their host environment indicating coevolution

(Brown and Wernegreen, 2019, Chua et al., 2020). But the recurrent presence of

Acetobacteraceae in the gut of formicine ants potentially also reflects direct transmission of

bacteria among individuals, selective uptake on the part of the ants, specific adaptation for

colonizing ant guts on the part of the bacteria, or some combination of all three (Engel and

Moran, 2013).

Generally, the immune system together with physiochemical properties of the gut

environment maintains homeostasis between gut associated microbes and the host (Chu and

Mazmanian, 2013, McFall-Ngai et al., 2013, Rakoff-Nahoum et al., 2004, Slack et al., 2009,

Watnick and Jugder, 2020, Xiao et al., 2019, see also Foster et al., 2017). Highly acidic

stomach lumens are ubiquitous in higher vertebrates, including amphibians, reptiles, birds and

mammals (Beasley et al., 2015, Koelz, 1992), while in insects acidic regions have rarely been

described from midgut regions (Chapman, 2013, Holtof et al., 2019). However in both, higher

vertebrates and the fruit fly *Drosophila melanogaster*, acidic gut compartments together with

the immune system serve microbial control and prevent infection by pathogens (Giannella et

al., 1972, Howden and Hunt, 1987, Martinsen et al., 2005, Overend et al., 2016, Rakoff-

Nahoum et al., 2004, Slack et al., 2009, Tennant et al., 2008, Watnick and Jugder, 2020).

Formicine ant species possess a highly acidic poison gland secretion containing formic acid as

its main component (Lopez et al., 1993, Osman and Brander, 1961, Schmidt, 1986). Although

the poison is presumably foremost used as a defensive weapon (Osman and Kloft, 1961), it is

also distributed to the environment of these ants as an external immune defence trait (*sensu*

Otti et al., 2014) to protect their offspring and the nest and to limit disease spread within the

society (see references in Tragust, 2016, Brütsch et al., 2017, Pull et al., 2018). To this end,

ants take up their poison from the acidopore, the opening of the poison gland at the gaster tip,

into their mouth (Tragust et al., 2013) during a specialized behaviour existing only in a subset

of ant families among all Hymenopterans (Basibuyuk and Quicke, 1999, Farish, 1972),

termed acidopore grooming.

Here we first investigate whether the poison is also swallowed during acidopore grooming in

C. floridanus and seven other formicine ant species from three genera in a comparative survey

through measurement of pH levels in the crop and midgut lumen, experimental manipulation

of acidopore access, and behavioural observations. In loss of acidopore and thus poison access

during survival experiments and in *in vitro* and *in vivo* bacterial viability and growth

experiments, we then investigate whether swallowing of the poison can serve gut microbial

control and prevent bacterial pathogen infection analogous to acidic stomachs of higher

vertebrates and acidic midgut regions in the fruit fly. Complementing these experiments, we

also investigate whether acidopore access has the potential to limit pathogen transmission

during trophallactic food exchange. Finally, we explore whether swallowing of the poison

acts as a microbial filter that is permissible to gut colonization by bacteria from the family

Acetobacteraceae.

Results

Swallowing of the poison gland secretion leads to acidic crops

Formicine ants naturally perform a specialized grooming behaviour, acidopore grooming,

during which they take up their poison from the acidopore, the opening of the poison gland at

the gaster tip, into their mouth (Basibuyuk and Quicke, 1999, Farish, 1972, Tragust et al.,

2013). To reveal whether the poison is also swallowed during acidopore grooming, we

monitored acidity levels in the crop lumen of the Florida carpenter ant *Camponotus floridanus*

after feeding them 10% honey water (pH = 5). We found that after feeding the crop became

increasingly acidic over time, reaching highly acidic values 48h after feeding (median pH = 2;

95% CI: 1.5-3.4), whilst renewed access to food after 48h raised the pH to levels recorded

after the first feeding (Fig. 1a; LMM, LR-test, $\chi^2 = 315.18$, df = 3, $P < 0.001$; Westfall

corrected post-hoc comparisons: 0+4h vs. 48h+4h: $P = 0.317$, all other comparisons: $P <$

0.001). This acidification after feeding was limited to the crop and did not extend to the

midgut (Fig. 1 – figure supplement 1; pH-measurements at four points along the midgut 24h

after access to 10% honey-water; mean ± se; midgut position 1 = 5.08 ± 0.18, midgut position

2 = 5.28 ± 0.17, midgut position 3 = 5.43 ± 0.16, midgut position 4 = 5.31 ± 0.19). Prevention

of acidopore grooming in *C. floridanus* ants through immobilisation (see Materials and

Methods) for 24h after feeding resulted in a significantly diminished acidity in the crop

compared to non-prevented ants (Fig. 1b; LMM, LR-test, $\chi^2 = 44.68$, df = 1, $P < 0.001$). A

similar, significantly diminished acidity in crop lumens was ubiquitously obtained in a

comparative survey across seven formicine ant species (genera: *Camponotus*, *Lasius* and

Formica) when ants were prevented from acidopore grooming through immobilisation

compared to non-prevented ants (Fig. 1c; two-sided Wilcoxon rank sum tests, comparisons

for all ant species: $P \leq 0.036$). This indicates that the causative agent of crop acidity in

formicine ants is not an internal source but an external one and that acidity of crop lumens is

achieved through swallowing of the acidic poison during acidopore grooming.

Although venomous animals often bear a cost of venom production and express behavioural

adaptations to limit venom expenditure (Casewell et al., 2013), *C. floridanus* increases the

frequency of acidopore grooming within the first 30 min. after ingesting food but also after

ingesting water compared to unfed ants (Fig. 1 - figure supplement 2; GLMM, LR-test, $\chi^2 =$

33.526, df = 2, $P < 0.001$; Westfall corrected post-hoc pairwise comparisons, water vs. 10%

honey-water: $P = 0.634$, unfed vs water and unfed vs 10% honey-water: $P < 0.001$). This

suggests that the behaviour of acidopore grooming is temporarily upregulated after fluid

ingestion, irrespective of the fluid's nutritional value. Moreover, pH levels were highly acidic

in the crop of *C. floridanus* ants taken directly out of a satiated colony (Fig. 1 – figure

supplement 3; major workers: median pH = 2, 95% CI: 2-3; minor workers: median = 3, CI:

2.5-3.6) and in worker cohorts that were satiated for three days and then starved for 24h

before measurements (majors: pH = 2, 95% CI: 2-3; minors: median = 2, CI: 2-3). This

indicates that the increased frequency of acidopore grooming after fluid ingestion and

swallowing of the acidic poison serves to maintain an optimal, acidic baseline pH in the crop

lumen of formicine ants following perturbation thereof through ingested fluids.

Poison acidified crops can serve microbial control

Acidic stomachs of vertebrates and acidic midgut regions in the fruit fly *Drosophila*

melanogaster can prevent infection by bacterial pathogens (Giannella et al., 1972, Howden

and Hunt, 1987, Martinsen et al., 2005, Overend et al., 2016, Tennant et al., 2008). For

swallowing of the poison to play a similar role in formicine ants and for the ensuing

acidification to take effect after perturbation of the crop pH through ingested fluids, ingested

fluids need to remain in the crop for a minimum time before being passed to the midgut. We

therefore estimated food passage over time through the gut of *C. floridanus* ants with food

containing fluorescent particles. We found that only a small amount of ingested food is passed

from the crop to the midgut 2-4h after feeding, while thereafter food is steadily passed from

the crop to the midgut until 18h after feeding (Fig. 2 – supplementary figure 1).

We then tested whether crop acidification after feeding could inhibit *Serratia marcescens*, an

insect pathogenic bacterium (Grimont and Grimont, 2006), ingested together with food and

prevent its passage from the crop to the midgut in *C. floridanus* ants. We measured this at two

time points before (0.5h and 4h) and after (24h and 48h) main food passage from the crop to

the midgut, with the time directly after food ingestion (0h) serving as a reference. When fed to

C. floridanus, *S. marcescens* presence decreased sharply over time in the crop (Fig. 2a;

GLMM, LR-test, $\chi^2 = 220.78$, df = 4, $P < 0.001$). The proportion of CFUs that we were able to

retrieve from the crop relative to the mean at 0h in the crop diminished from 43% at 0.5h

post-feeding (median, CI: 0-543%) to 0% at 4h (CI: 0-4%), 24h (CI: 0-1.8%), and 48h (CI: 0-

18%) post-feeding. In addition, relative to the mean at 0h in the crop, *S. marcescens* could

only be detected at extremely low numbers in the midgut (median 0%) at 0h (CI: 0-4%), 0.5h

(CI: 0-1%) and 24h (CI: 0-1%) post-feeding and not at all at 4h and 48h post-feeding (Fig. 2b;

GLMM, LR-test, $\chi^2 = 1.044$, df = 2, $P = 0.593$). This suggests that a lowering of the crop

lumen pH through swallowing of the poison quickly and effectively reduces the viability of $S.$

$marcescens$ in the crop thus preventing further transport to the midgut.

An in $vitro$ experiment on the ability of $S.$ $marcescens$ to withstand acidic conditions created

with formic acid, the main component of the formicine poison gland secretion (Lopez et al.,

1993, Osman and Brander, 1961, Schmidt, 1986), supports this interpretation. Incubation of $S.$

$marcescens$ for 2h in 10% honey water acidified with formic acid to pH 4 resulted in a

significantly lower number of CFUs relative to pH 5 and in zero growth for incubations at

pH-levels that were lower than 4 (Fig. 2 – figure supplement 2; GLM, LR-test, $\chi^2 = 79.442$, df

$= 1$, $P < 0.001$). This in $vitro$ experiment together with our in $vivo$ experiment on the ability of

$S.$ $marcescens$ to withstand acidic conditions in the crop of $C.$ $floridanus$, indicates that poison

acidified crops can serve microbial control in formicine ants.

Acidopore access improves survival in the face of pathogen contaminated food

To test whether the ability to swallow the acidic poison can limit infection by oral pathogens,

we prevented acidopore grooming through immobilisation in $C.$ $floridanus$ ants for 24h after

feeding them once with 5µl of either honey water contaminated with $S.$ $marcescens$, or non-

contaminated honey water and monitored their survival thereafter without providing

additional food. We found that acidopore access after pathogen ingestion significantly

increased the survival probability of ants (Fig. 3; COXME, LR-test, $\chi^2 = 20.95$, df = 3, $P =$

0.0001). The survival of ants prevented from acidopore grooming and fed once with the

pathogen contaminated food was significantly lower than that of non-prevented ants fed the

same food source (Westfall corrected post-hoc comparisons: FA - | $Serratia$ presence + vs. all

other ant groups: $P \leq 0.027$). Contrary, non-prevented ants fed once with the pathogen

contaminated food source did not differ in survival to prevented and non-prevented ants fed

the non-contaminated food source (Westfall corrected post-hoc comparisons: FA + | *Serratia*

presence + vs. FA + | *Serratia* presence – and FA + | *Serratia* presence + vs. FA + | *Serratia*

presence –: $P \geq 0.061$ both). This indicates that acidopore access and thus swallowing of the

poison provides a fitness benefit in terms of survival to formicine ants upon ingestion of

pathogen contaminated food.

Acidopore access can limit pathogen transmission

The ability to swallow the acidic poison may not only improve survival of formicine ants

feeding directly on pathogen contaminated food but also of ants that share the contaminated

food via trophallaxis. To test this, we created two types of donor-receiver ant pairs. Donor

ants in both pairs were directly fed *S. marcescens* contaminated food every other day, while

receiver ants obtained food only through trophallaxis from their respective donor ants.

Receiver ants in both pairs were precluded from swallowing of the poison through blockage

of their acidopore opening (see Materials and Methods), while donor ants were blocked in one

pair but only sham blocked in the other pair. We found that the duration of trophallaxis

between the two donor-receiver ant pairs during the first 30min. of the first feeding bout did

not significantly differ (Fig. 4 – figure supplement 1; LMM, LR-test, $\chi^2 = 1.23$, df = 1, $P =$

0.268), indicating that trophallactic behaviour was not influenced through acidopore blockage

in donor ants at the beginning of the experiment. Over the next 12 days we found that

acidopore blockage *per se* had a significant negative effect on the survival of donor as well as

receiver ants (Fig. 4; COXME, LR-test, $\chi^2 = 66.68$, df = 3, $P < 0.001$). Importantly however,

although receiver ants that obtained food every other day from donors with the ability to

swallow the poison died at a higher rate than their respective donor counterparts (hazard ratio:

1.81; Westfall corrected post-hoc comparison: $P < 0.001$) they were only half as likely to die

compared to receiver ants that obtained pathogen contaminated food from blocked donors

unable to swallow the poison (hazard ratio: 0.56; Westfall corrected post-hoc comparison: $P <$

0.001). This indicates that swallowing of the poison after feeding on pathogen contaminated

food does not only improve survival of formicine ants directly feeding on pathogen

contaminated food but also of ants that share the contaminated food via trophallaxis. Hence,

swallowing of the poison and the ensuing crop acidity have the potential to limit oral disease

transmission during food distribution within a formicine ant society.

Poison acidified crops can serve as a chemical filter for microbial selection

In addition to pathogen control, poison acidified formicine ant crops might act as a chemical

filter for gut associated microbial communities, similar to gut morphological structures that

can act as mechanical filters in ants and other insects (Itoh et al., 2019, Lanan et al., 2016,

Ohbayashi et al., 2015). To investigate the idea of a chemical filter, we tested the ability of the

insect gut associated bacterium *Asaia* sp. (family Acetobacteraceae) (Crotti et al., 2009, Favia

et al., 2007) to withstand acidic environments *in vitro* and *in vivo*. In contrast to *S. marcescens*

(Fig. 2 – supplementary figure 2), *Asaia* sp. was not affected by an incubation for 2h in 10%

honey water acidified with formic acid to a pH of 4 and was still able to grow when incubated

at a pH of 3 in *in vitro* tests (Fig. 5 – figure supplement 1; GLM, overall LR-test $\chi^2 = 21.179$,

df = 2, $P < 0.001$; Westfall corrected post hoc comparisons: pH = 5 vs. pH = 4: $P = 0.234$, all

other comparisons: $P < 0.001$). Moreover, in *in vivo* tests, *Asaia* sp. only gradually diminished

over time in the crop (Fig. 5a; GLMM; LR-test, $\chi^2 = 124.01$, df = 4, $P < 0.001$) with the

proportion of CFUs that we were able to retrieve from the crop relative to the mean at 0h in

the crop diminishing to only 34% (median, CI: 3-85%) and 2% (CI: 0-7%) at 4h and 24h post-

feeding, respectively. At the same time, relative to the mean at 0h in the crop, *Asaia* sp.

steadily increased in the midgut (Fig. 5b; GLMM; LR-test, $\chi^2 = 59.94$, df = 3, $P < 0.001$) from

its initial absence at 0h post-feeding to 2% (median, CI: 0-5%) at 48h post-feeding. This

indicates that a lowering of the crop lumen pH through swallowing of the acidic poison is

permissible to the passage of *Asaia* sp.from the crop to the midgut. Together our *in vitro* and

in vivo experiments with *S. marcescens* and *Asaia* sp. suggest that in formicine ants, poison

acidified crops might act as a chemical filter that works selectively against the establishment

of opportunistic and potentially harmful bacteria but allows entry and establishment of

members of the bacterial family Acetobacteraceae. This view is also supported by results

obtained *in vivo* for *E. coli,* a bacterium that is not a gut associate of insects (Blount, 2015)

and which, similarly to *S. marcescens,* was not able to pass from the crop to the midgut in *C.*

floridanus ants (Fig. 5 – figure supplement 2; crop: GLMM, LR-test, $\chi^2 = 156.74$, df $= 4$, *P* <

0.001; midgut: GLMM, LR-test, $\chi^2 = 14.898$, df $= 3$, *P* $= 0.002$).

Discussion

The results of our study indicate that formicine ants not only distribute their highly acidic,

antimicrobial poison to their environment as an external immune defence trait (see references

in Tragust, 2016, Brütsch et al., 2017, Pull et al., 2018), but also swallow the poison resulting

in the creation of an acidic environment in their stomach, the crop (Fig. 1). This acidic

environment can serve microbial control, limiting establishment of orally acquired,

pathogenic microbes with ingested food (Fig. 2). In line with this, survival of formicine ants

that fed pathogen contaminated food was improved if the ants were able to swallow their

poison (Fig. 3). This fitness benefit also extended to ants that received pathogen contaminated

food via trophallaxis from donor ants with the ability to swallow their poison (Fig. 4).

Additionally, poison acidified crop lumens can act as chemical filters that selectively allow

gut entry and establishment of members of the bacterial family Acetobacteraceae, a recurring

part of the gut microbiota of formicine ants (Fig. 5) (Brown and Wernegreen, 2016, Chua et

al., 2018, He et al., 2011, Ivens et al., 2018, Russell et al., 2017).

Instances of extreme pH conditions, especially alkaline, have been described in

gastrointestinal tract compartments of several insects (Chapman, 2013, Holtof et al., 2019).

Our study is however the first to report highly acidic conditions in insect crops and in animal

stomachs in general apart from higher vertebrate species (Beasley et al., 2015, Koelz, 1992).

Acidic conditions in the crop and other gut compartments of insects might be achieved in

several ways. In principle, a digestive compartment with a certain pH can be generated

through physiological mechanisms involving a transport-loop of acid-base equivalents across

epithelia (Onken and Moffett, 2017). Insects could thus regulate the pH of their crop or of

other gut compartments through active uptake and excretion of acid–base equivalents across

the gut epithelium (Matthews, 2017). With few notable exceptions (Flower and Filshie, 1976,

Miguel-Aliaga et al., 2018), the exact physiological mechanisms responsible for the creation

of a gut lumen compartment with a certain pH are however often unknown (Harrison, 2001).

Alternatively, insect gut associated microbes might contribute to the gut pH. Many bacteria

produce short chain fatty acids and are thus able to acidify their environment (Ratzke et al.,

2018, Ratzke and Gore, 2018). Members of the bacteria family Acetobacteraceae, and various

Lactobacilli, gut microbial associates of Hymenoptera (McFrederick et al., 2013), release

acetic acid as a waste product of their fermentative metabolism (Oude Elferink et al., 2001,

Wolfe, 2005) and might thus influence pH conditions in the gut of their insect host. Finally,

glandular secretions might contribute to the pH of the insect gut. In grasshoppers and crickets

it has been suggested that salivary glands are responsible for slightly acidic (pH 5-6) crop

lumens (Cooper and Vulcano, 1997, Harrison et al., 1992) and a recent study in the honeybee

found that larval food has a pH of 4 due to the addition of fatty acids from a cephalic gland

connected to the oral cavity (Muresan and Buttstedt, 2019). The oral cavity of ants is

connected to several cephalic and thoracic glands (Emmert, 1968, Hölldobler and Wilson,

1990) that apart from lubricating compounds and digestive enzymes (Billen, 2009) can also

produce acidic substances in some ants (Attygalle and Morgan, 1984, Morgan, 2008, Vander

Meer, 2012). Physiological mechanisms, gut associated microbes or cephalic and thoracic

glands connected to the oral cavity are however unlikely to be responsible for crop lumen

acidity in our study, as immobilisation upon food ingestion severely impaired acidification of

the crop in all formicine ants tested. This rather indicates that formicine ants use an external

source for acidification but not an internal one.

Ants possess a huge diversity of exocrine glands all over their body (Billen, 2009, Billen and

Sobotnik, 2015). Several of these glands produce acidic secretions (Attygalle and Morgan,

1984, Morgan, 2008, Vander Meer, 2012) and could thus serve as external sources for crop

acidity. Most notable among these exocrine glands are the metapleural gland and the poison

gland whose openings are located on the thorax and the gaster tip, respectively. Both produce

acidic secretions in several ant species (Fernández-Marín et al., 2015, Lopez et al., 1993,

Osman and Brander, 1961, Yek and Mueller, 2011) and are actively groomed with

movements involving the mouth (Basibuyuk and Quicke, 1999, Farish, 1972, Fernández-

Marín et al., 2009, Fernández-Marín et al., 2006, Tragust et al., 2013). Although

immobilisation of ants in our experiments effectively prevented grooming of both glands,

acidic metapleural gland secretions cannot serve as a source for crop acidity in the focal ant

species of our study, *Camponotus floridanus*, and in the ant *Camponotus maculatus* in our

comparative survey as ants of the genus *Camponotus* show, with few exceptions, an

evolutionary loss of the metapleural gland (Yek and Mueller, 2011). In addition, in the

formicine ant *Lasius neglectus,* grooming of the poison gland opening, the acidopore, results

in the uptake of the poison into the mouth, while metapleural gland grooming was never

observed (Tragust et al., 2013). Therefore, acidity of formicine ant crops in our study is likely

the result of acidopore grooming and swallowing of the highly acidic poison, although our

experiments cannot completely rule out the use of other sources.

Interestingly, although immobilisation of ants upon food ingestion led to a significantly

impaired acidification of the crop in all formicine ants tested, we found a variable crop acidity

in immobilised but also non-immobilised ants in our comparative survey. While this could

indicate the use of acidic sources other than the poison gland, it could also indicate species

specific differences in acidopore grooming, differences in the composition of the poison gland

secretion or, as the ant crop is seldom completely empty (Greenwald et al., 2015, Greenwald

et al., 2018) simply be the result of differences in crop filling status before feeding in our

experimental setup. In *C. floridanus* we found that ants increased the frequency of acidopore

grooming following ingestion of water or food. Together with the observed baseline level of

acidity in the crop of around pH 2-3 under fully satiated and starved conditions, this indicates

that *C. floridanus* ants strive to maintain an acidic, likely optimal pH in their crop through

swallowing of their poison following perturbation of the crop pH after the ingestion of fluids.

Although digestion is initiated in the crop, the midgut is the primary site of digestion in

insects and the midgut epithelium plays a pivotal role in maintaining an optimal pH, as the pH

is one of the most important regulators of digestive enzyme activity (Holtof et al., 2019, Terra

and Ferreira, 1994). This might be the reason why the midgut pH of *C. floridanus* shows only

slightly acidic levels (pH 5) after highly acidic levels in the crop; a change in pH that could be

achieved via physiological mechanisms. While the variable acidity of crop lumens in our

comparative survey will have to await further investigations, in *C. floridanus* ants, both, the

increased frequency of acidopore grooming upon fluid ingestion and the change of pH to only

slightly acidic levels in the midgut, the primary site of digestion, indicate that poison acidified

crop lumens in formicine ants serve other functions than digestion, although an additional

digestive function will have to be investigated.

Acidic fermentation of food is, for humans, a common practice which dates back centuries

(Hutkins, 2019). Due to their antimicrobial activity (Cherrington et al., 1991, Hirshfield et al.,

2003, Lund et al., 2014), the addition of organic acids to food represents the best practice to

preserve food together with refrigeration (Theron and Rykers Lues, 2010). In *C. floridanus*

ants ingested food largely remained in the crop for at least 2-4h before being passed to the

midgut, which agrees with food passage times through the gastrointestinal tract of other ants

(Cannon, 1998, Gößwald and Kloft, 1960, Gösswald and Kloft, 1960a, Howard and

Tschinkel, 1981, Markin, 1970). This would enable swallowing of the poison and the ensuing

crop acidity to take effect after perturbation of the crop pH through feeding. Acidic crop

lumens in formicine ants might therefore play a role in the sanitation of ingested fluids similar

to the ubiquitous, antimicrobials enabled sanitation of food in animals that provision food to

their offspring or that store, cultivate, develop or live in food (Cardoza et al., 2006, Currie et

al., 1999, Herzner et al., 2013, Herzner and Strohm, 2007, Joop et al., 2014, Milan et al.,

2012, Mueller et al., 2005, Scott et al., 2008, Shukla et al., 2018a, Vander Wall, 1990, Vogel

et al., 2017). A food sanitation function is supported by the rapid reduction of the insect

bacterial pathogen *Serratia marcescens* in the crop of *C. floridanus* ants *in vivo* and the

sensitivity of this bacterial pathogen to acidic conditions created with formic acid *in vitro*. The

sensitivity of *S. marcescens* to acidic conditions created with formic acid *in vitro* likely even

underestimates the antimicrobial effect of the formicine ant poison *in vivo*. Although formic

acid is the main component of the formicine ant poison (Schmidt, 1986), the poison gland also

contains acetic acid, hexadecanol, hexadecyl formate, hexadecyl acetat (Lopez et al., 1993)

and a small fraction of unidentified peptides (Hermann and Blum, 1968, Osman and Brander,

1961). Additionally, the poison of formicine ants is usually expelled from the acidopore

together with contents of the Dufour gland (Regnier and Wilson, 1968, Schoeters and Billen,

1996, but see Billen, 1982), which serve as wetting agents for the poison (Löfqvist, 1976, see

also Kohl et al., 2001). In a previous study we found that formic acid alone could explain 70%

of the antimicrobial activity of the poison against an entomopathogenic fungus, but that the

combination of formic acid with other components of the poison gland and the Dufour gland

could explain 94% (Tragust et al., 2013). In addition to the likely higher antimicrobial activity

of the natural poison compared to the activity of formic acid alone, *in vivo* immune system

effectors released into the gut lumen might contribute to the inability of *S. marcescens* to

establish in the gastrointestinal tract of *C. floridanus*. Highly acidic stomachs in vertebrates

and acidic midgut regions in the fruit fly *Drosophila melanogaster* serve together with

immune system effectors microbial control and prevent infection by oral pathogens (Giannella

et al., 1972, Howden and Hunt, 1987, Martinsen et al., 2005, Overend et al., 2016, Rakoff-

Nahoum et al., 2004, Slack et al., 2009, Tennant et al., 2008, Watnick and Jugder, 2020).

Future studies will need to investigate the contribution of immune system effectors released

into the gut lumen to the rapid reduction of *S. marcescens* in the crop of *C. floridanus* and its

inability to establish in the midgut.

Our study also found that access to the poison improved survival of *C. floridanus* ants after

ingestion of food contaminated with the bacterial pathogen *S. marcescens*. Mortality of ants

was generally high in this experimental setup irrespective of whether access to the poison was

manipulated or not. This is unlikely the consequence of stress induced mortality through the

method of immobilisation employed to prevent poison access in this experiment, as

immobilised and non-immobilised ants did not differ in their survival upon ingestion of non-

contaminated food. Rather, starvation following the one time feeding of contaminated and

non-contaminated food likely increased mortality in all ants. In a similar experiment involving

the formicine ant species *Formica exsecta*, oral exposure to *S. marcescens* contaminated food

followed by starvation as well as starvation alone led to a high mortality of ants with no

additive effects of pathogen exposure combined with starvation (Stucki et al., 2019). In

addition to starvation, social isolation of *C. floridanus* ants might have increased mortality, as

ants were kept individually in petri dishes. Social isolation has been shown to increase

mortality in ants (Koto et al., 2015) and social isolation has been shown to reduce an

individual's capacity to fight infections in other group living animals (Kohlmeier et al., 2016).

Despite the high mortality observed under these experimental conditions, our experiments on

the ability of *S. marcescens* to withstand acidic conditions *in vitro* and *in vivo* and our survival

experiment upon ingestion of *S. marcescens* contaminated food, suggest that swallowing of

the poison and the ensuing crop acidity can indeed serve microbial control in formicine ants

and improve survival in the face of pathogen contaminated food. In ants and other animals

that lack an acidic poison, acids produced by other glands or acidic derivatives produced by

defensive symbionts (Florez et al., 2015) or other environmental or gut associated bacteria,

might provide functionally similar roles to the acidic poison in formicine anrts, as indicated in

bees (Palmer-Young et al., 2018) and termites (Inagaki and Matsuura, 2018).

In addition to improve survival of ants that directly ingested pathogen contaminated food, the

ability of donor ants to access their poison also improved survival of receiver ants without

access to their poison when receiver ants shared pathogen contaminated food during

trophallactic food exchange. Although our experiments on the ability of *S. marcescens* to

withstand acidic conditions *in vivo* and *in vitro* indicate that this is likely due to fewer viable

bacteria that are passed on from donor to receiver ants during trophallactic food exchange, it

remains to be established whether this is indeed the case or whether this might be due to

obtaining trophallactic fluids with antimicrobial activity. Antimicrobial activity of formicine

ant trophallactic fluids has been described in previous studies (Hamilton et al., 2011, LeBoeuf

et al., 2016). These studies linked the antimicrobial activity of trophallactic fluids to the

presence of proteins related to cathepsin D, a lysosomal aspartic protease that can exhibit

antibacterial effector activity and the proteolytic production of antimicrobial peptides (Ning et

al., 2018). Our results however suggest a major role of swallowing the acidic poison to the

antimicrobial activity of trophallactic fluids in formicine ants. Future studies will need to

disentangle the relative contributions of crop acidity, proteins related to cathepsin D and, as

previously pointed out, other immune effectors that are released into the insect gut to the

antimicrobial activity of formicine ant trophallactic fluids.

The sensitivity of the bacterial pathogen *S. marcescens* to acidic conditions and the fitness

benefit bestowed to ants with direct and indirect access to the poison after feeding on or

receiving of pathogen contaminated food, might also indicate that swallowing of the poison

and the ensuing crop acidity can act as an important barrier to oral disease spread within

formicine ant societies. In the formicine ant *Formica polyctena,* food passage from the crop to

the midgut is dependent upon whether food is directly eaten or is transferred via trophallaxis

with only 20% of the honey water consumed directly seen in the midgut 4.5 h after feeding,

while 77% of the honey water received during trophallaxis reaching the midgut 2.5 hours after

feeding (Gösswald and Kloft, 1960a). Poison acidified crop lumens might therefore alleviate

the cost of sharing pathogen contaminated food (Onchuru et al., 2018, Salem et al., 2015) and

effectively counteract the generally increased risk of pathogen exposure and transmission

associated with group-living (Alexander, 1974, Boomsma et al., 2005, Kappeler et al., 2015).

On the other hand, trophallactic food exchange is a dynamic process that is dependent upon

the functional role of the worker (Greenwald et al., 2015), food type (Buffin et al., 2011), and

likely many other contexts. For example, it has repeatedly been reported that after a time of

starvation food is distributed extremely quickly and efficiently via trophallaxis within an ant

colony (Buffin et al., 2009, Gösswald and Kloft, 1960b, Markin, 1970, Sendova-Franks et al.,

2010, Traniello, 1977, Wilson and Eisner, 1957). While this increases the threat of pathogen

dissemination together with food, it has been suggested that dilution and mixing of food

together with the existence of disposable ants specialised in food storage can mitigate the

threat of pathogenic and noxious substances distributed together with food (Buffin et al.,

2011, Sendova-Franks et al., 2010). Recently, it has been reported that ant social networks can

plastically respond to the presence of a pathogen and that ants alter their contact network to

contain the spread of a disease (Stroeymeyt et al., 2018). In our study, the first bout of

trophallactic food exchange between donor and receiver ants was not affected by the

manipulation of poison access. However, especially at later time points, potential changes in

the amount of food transmitted cannot be excluded, as trophallaxis and feeding behaviour in

general might depend on the infection status of ants engaging in trophallactic food exchange

(Hite et al., 2020). In future studies, it will therefore be interesting to examine whether

swallowing of the poison and the ensuing crop acidity truly acts as a barrier to oral disease

spread within formicine ant societies, especially given the technical advances to track multiple

individuals of a group simultaneously over time that have been made in recent years (Gernat

et al., 2018, Greenwald et al., 2015, Imirzian et al., 2019, Stroeymeyt et al., 2018).

Our results on the ability of bacteria to withstand acidic environments *in vitro* and *in vivo* also

provide evidence that swallowing of the acidic poison might not only serve microbial control

but might also act as a chemical filter for microbes, working selectively against the

establishment of pathogenic or opportunistic bacteria but allowing entry and establishment of

species from the bacteria family Acetobacteraceae in *C. floridanus* ants. In ants, gut

morphological structures, i.e. the infrabuccal pocket, an invagination of the hypopharynx in

the oral cavity (Eisner and Happ, 1962), and the proventriculus, a valve that mechanically

restricts passage of fluids from the crop to the midgut (Eisner and Wilson, 1952), can

consecutively filter particles and microbes out of ingested food (Cannon, 1998, Glancey et al.,

1981, Lanan et al., 2016, Little et al., 2006, Quinlan and Cherret, 1978). In other insects, gut

morphological structures have been described to act as mechanical filters for gut associated

microbes (Itoh et al., 2019, Ohbayashi et al., 2015). Given that acidic gut compartments in

humans and fruit flies not only serve prevention of oral diseases but generally structure the

gut microbiota of their hosts (Imhann et al., 2016, Overend et al., 2016), it is likely that poison

acidified crop lumens can explain at least part of the recurrent presence of Acetobacteraceae

in the gut of formicine ants and the otherwise reduced microbial diversity and abundance of

gut associated microbes (Brown and Wernegreen, 2016, Brown and Wernegreen, 2019, Chua

et al., 2018, Chua et al., 2020, Ivens et al., 2018, Russell et al., 2017).

Acetobacteraceae often thrive in sugar-rich environments (Mamlouk and Gullo, 2013). As

many ants predominantly feed on sugars, e.g. extrafloral nectaries and aphid honeydew

(Blüthgen et al., 2004, Blüthgen et al., 2000, Way, 1963), the presence of Acetobacteraceae in

the gut of formicine ants could simply be a by-product of their dietary preferences (Anderson

et al., 2012). On the other hand, some members of the Acetobacteraceae are pathogenic in

humans and the fruit fly *Drosophila melanogaster* (Greenberg et al., 2006, Roh et al., 2008,

Ryu et al., 2008) and most Acetobacteraceae produce metabolites that can potentially interfere

with insect physiology and innate immunity (Chouaia et al., 2014). This may indicate that

Acetobacteraceae found in formicine ants can act as pathogens. The ability of *Asaia* sp., a

member of the Acetobacteraceae, to withstand acidic conditions *in vitro* and *in vivo* and to

establish in the midgut of *C. floridanus* however indicate that host filtering (Mazel et al.,

2018) might account for the pattern of recurring gut microbial communities in formicine ants.

In addition, where studied in detail, Acetobacteraceae found associated with the gut of

formicine ants show signs of genomic and metabolic adaptations to their host environment

(Brown and Wernegreen, 2019, Chua et al., 2020), indicating coevolution and potentially also

mutual benefit, though this has not formally been established (see also Mushegian and Ebert,

2016). The creation of an acidic pH crop environment in formicine ants that is easier to

endure if colonizing microbes are mutualists agrees with the theoretical concept of screening,

as opposed to signalling, as a means of partner choice in cross-kingdom mutualisms (Archetti

et al., 2011a, Archetti et al., 2011b, Biedermann and Kaltenpoth, 2014, Scheuring and Yu,

2012). Signalling is a well-known mechanism of partner selection in the context of mate

choice (Grafen, 1990, Zahavi, 1975) or plant-microbe mutualisms (Clear and Horn, 2019,

Rebolleda-Gomez and Wood, 2019) and signals play an important role in the cross talk

between the immune system and host associated microbes (Allaire et al., 2018, Fischbach and

Segre, 2016, Moura-Alves et al., 2019, Villena et al., 2018). If the signal has a cost, only good

quality individuals will find it profitable to advertise their quality, and therefore the signal will

be honest. Contrary to signalling, where costly information is displayed to partners, in

screening a costly environment is imposed on partners that excludes all but high-quality ones.

Partner choice in a number of cross-kingdom mutualisms is readily explained by screening

(see examples in Archetti et al., 2011a, Archetti et al., 2011b, Biedermann and Kaltenpoth,

2014, Scheuring and Yu, 2012) but experimental evidence is so far limited in insect-microbe

associations (Innocent et al., 2018, Itoh et al., 2019, Ranger et al., 2018). Although our

experiments can only indicate screening as a means of partner choice in formicine ants, the

results of our study would provide support for the prediction that screening is more likely to

evolve from a host's defence trait against parasites (Archetti et al., 2011a, Archetti et al.,

2011b), i.e. the highly acidic poison that creates a selective environment for microbes.

Conclusion

Overall, our study provides evidence that swallowing of the formic acid containing poison

gland secretion and the ensuing acidity in the crop lumen acts as a chemical filter for control

and selection of microbes ingested with food, protecting formicine ants from food borne

bacterial pathogens and structuring their gut associated microbial communities. Only recently,

studies have started to acknowledge that antimicrobials in the environment of an organism

have the potential dual role of microbial selection and control (Brinker et al., 2019b, Duarte et

al., 2018, Innocent et al., 2018, Shukla et al., 2018a, Shukla et al., 2018b, Strohm et al., 2019,

Tragust et al., 2020). Antimicrobials used as external immune defence traits (Otti et al., 2014)

may generally not only serve pathogen protection and microbial control but may act as

microbial filters to manage host associated microbes, be it in food or the environment, and

thus contribute to a host's ecological and evolutionary success. In the case of social species,

externally acting antimicrobials might contribute to the ecological and evolutionary success of

this group of insects by alleviating the increased risk of pathogen exposure and transmission

associated with group living but allowing the acquisition and transmission of microbial

mutualists.

Methods

Ant species and maintenance. Colonies of the carpenter ant *Camponotus floridanus* were

collected in 2001 and 2003 in Florida, USA, housed in Fluon® (Whitford GmbH, Diez,

Germany) coated plastic containers with plaster ground and maintained at a constant

temperature of 25°C with 70% humidity and a 12h/12h light/dark cycle. They were given

water *ad libitum* and were fed two times per week with honey water (1:1 tap water and

commercial quality honey), cockroaches (*Blaptica dubia*) and an artificial diet (Bhatkar and

Whitcomb, 1971). For comparison, workers of one other *Camponotus* species (*Camponotus*

maculatus), collected close to Kibale Forest, Uganda, in 2003 and housed under identical

conditions as *Camponotus floridanus* were used. Additionally, six other formicine ant species,

one *Lasius* and five *Formica* species (*Lasius fuliginosus, Formica cinerea, Formica*

cunicularia, Formica fuscocinerea, Formica pratensis and *Formica rufibarbis*) were collected

in Bayreuth, Germany in 2012 and 2018 and kept for approximately two weeks prior

experimental use at 20°C, 70% humidity and a 14h/10h light/dark cycle. Except otherwise

noted only the small worker caste ("minors") of *Camponotus* species was used.

Acidity of the crop lumen and pH measurements. To determine whether formicine ants

swallow their poison after feeding, we tracked changes in pH-levels of the crop lumen in *C.*

floridanus ants over time. Before use in experimental settings, cohorts of ~100 ants were

taken out of their natal colony (n = 6 colonies) into small plastic containers lined with Fluon®

and starved for 24-48h. Thereafter, ants were put singly into small petri dishes (Ø 55 mm)

with damp filter paper covered bottom, given access to a droplet of 10% honey water (w/v)

for 2h before removing the food source and measuring the pH of the crop lumen after another

2h (group 0+4h: n = 60 workers), after 24h (group 0+24h: n = 59 workers) or 48h (group

0+48h: n = 52 workers). To assess the effect of renewed feeding, a separate group of *C.*

floridanus ants were given access to 10% honey water 48h after the first feeding for 2h prior

to measuring the pH of their crop lumen after another 2h (group 48h+4h: n = 60 workers). To

measure the pH, ants were first cold anesthetized on ice, then their gaster was cut off with a

fine dissection scissor directly behind the petiole and leaking crop content (1-3µl) collected

with a capillary (5µl Disposable Micro Pipettes, Blaubrand intraMARK, Brand, Wertheim).

The collected crop content was then emptied on a pH sensitive paper to assess the pH

(Hartenstein, Unitest pH 1-11). This method of collecting crop content will invariably result

in some mixing of crop lumen content with haemolymph. As the pH of the insect

haemolymph ranges from only slightly acidic ($pH \geq 6.5$) to near-neutral or slightly alkaline

($pH \leq 8.2$) (Harrison, 2001, Matthews, 2017), this might have biased the results of our pH

measurements to slightly higher pH values. As a reference point for food pH, we also

measured the pH of 10% honey water on pH sensitive paper, which gave invariably $pH = 5$.

In addition, we measured the pH in the crop lumen and at four points in the lumen along the

midgut (1st measurement directly behind proventriculus to 4th measurement one mm apical

from insertion point of the Malpighian tubules) of *C. floridanus* workers that were fed 24 h

prior to measurements with 10% honey-water. For these measurements worker guts were

dissected as a whole and pH was measured in the crop (n = 2 workers from two colonies) and

along the midgut (all midgut points n = 10, except point four with n = 9 workers from four

different colonies) with a needle-shaped microelectrode (UNISENSE pH-meter;

microelectrode with needle tip of 20µm diameter).

In formicine ants, oral uptake of the poison into the mouth is performed via acidopore

grooming (Tragust et al., 2013). During this behavior ants bend their gaster forward between

the legs and the head down to meet the acidopore, the opening of the poison gland, at the

gaster tip (Basibuyuk and Quicke, 1999, Farish, 1972). In an additional experiment we

therefore compared the crop lumen pH of *C. floridanus* workers from four different colonies

that were either prevented to reach their acidopore (FA- ants) or could reach their acidopore

freely (FA+ ants). To do this, we again allowed single ants access to 10% honey water for 2h

after a starvation period, before cold anesthetizing them briefly on ice and immobilizing FA-

ants (n = 22 workers) in a pipetting tip, while FA+ ants (n = 23 workers) remained un-

manipulated. After 24h we measured the pH of the crop lumen as before.

To investigate whether swallowing of the acidic poison is widespread among formicine ants,

the latter experiment was repeated for six additional formicine ant species (FA- ants: n = 10

workers except for *Formica pratensis* with n = 21; FA+ ants: n = 10 workers except for

Formica pratensis with n=20; all ants: n = 1 colony) in the same fashion as described before

with the exception that apart from *Formica pratensis* the crop lumen was collected through

the mouth by gently pressing the ants' gaster. Crop lumen of *Formica pratensis* ants was

collected in the same fashion as crop lumen of *C. floridanus* ants.

To investigate whether the type of fluid and its nutritional value have an influence on the

frequency of acidopore grooming in *C. floridanus*, the following experiment was performed.

Cohorts of ~100 ants were taken out of their natal colony (n = 6 colonies) into small plastic

containers and starved for 24-48h. Thereafter, ants were again put singly into small petri

dishes (Ø 55 mm) and given access to either a 3µl droplet of 10% honey water (w/v, n = 126

ants, treatment: honey-water fed), a 3µl droplet of tap water (n = 128, water-fed) or to no fluid

(n = 125, unfed). After acclimatization (unfed ants) or after swallowing of the fluid (honey-

water and water-fed ants, both 1-2min.), all ants were filmed for the next 30min. (Logitech

webcam c910). These videos were then analyzed for the frequency of acidopore grooming.

Finally, we measured the pH in the crop lumen of *C. floridanus* ants (n = 3 colonies) under

satiated and starved conditions to estimate a baseline level of acidity in the crop. For this, ants

taken out of satiated, twice per week fed colonies on the day of feeding were compared to ants

that were maintained in cohorts of ~100 individuals for three days with access to 10% honey-

water and then starved for 24h before measuring the pH in their crop (n = 10 major and 10

minor workers per colony and condition). The pH in the crop lumen was measured as

described before by briefly cold anesthetizing ants an ice, collecting the crop content through

the mouth by gently pressing the ants' gaster and then emptying it on a pH sensitive paper

(Hartenstein, Unitest pH 1-11).

Bacterial strains and culture. As model entomopathogenic bacterium *Serratia marcescens*

DSM12481 (DSMZ Braunschweig, Germany) was used. This bacterium is pathogenic in a

range of insects (Grimont and Grimont, 2006) and has been detected in formicine ants, i.e.

Anoplolepis gracilipes (Cooling et al., 2018) and *Camponotus floridanus* (Ratzka et al.,

2011). While often non-lethal within the digestive tract, *S. marcescens* can cross the insect gut

wall (Mirabito and Rosengaus, 2016, Nehme et al., 2007) and is highly virulent upon entry

into the hemocoel (Flyg et al., 1980), not least due to the production of bacterial toxins

(Hertle, 2005). As a model bacterial gut-associate of ants *Asaia* sp. strain SF2.1 (Favia et al.,

2007), was used. *Asaia* sp. belongs to the family Acetobacteraceae, members of which often

thrive in sugar-rich environments (Mamlouk and Gullo, 2013), such as honey-dew that ants

like *C. floridanus* predominantly feed on. *Asaia* sp. is capable of cross-colonizing insects of

phylogenetically distant genera and orders (Crotti et al., 2009, Favia et al., 2007) and can be a

component of the gut associated microbial community of formicine and other ants (Chua et

al., 2018, Kautz et al., 2013a, Kautz et al., 2013b). In addition to *S. marcescens* and *Asaia sp.*,

Escherichia coli DSM6897 (DSMZ Braunschweig, Germany) was used as a model

opportunistic bacterium that is not a gut-associate of insects. *E. coli* bacteria are a principal

constituent of mammalian gut associated microbial communities but are commonly also found

in the environment (Blount, 2015).

Bacterial stocks of *S. marcescens, Asaia* sp., and *E. coli* were kept in 25% glycerol at -80°C

until use. For use, bacteria were plated on agar plates (LB-medium: 10g Tryptone, 5g Yeast

extract, 20g Agar in 1L MilliQ-water, and GLY-medium: 25g Gycerol, 10g Yeast extract, 20g

Agar in 1L MilliQ-water with pH adjusted to 5.0, for *S. marcescens/E. coli* and *Asaia* sp.

respectively), single colony forming units (CFUs) were picked after 24h (*S. marcescens/E.*

coli) or 48h (*Asaia* sp.) of growth at 30°C and transferred to 5ml liquid medium (LB-medium

and GLY-medium minus agar for *S. marcescens/E. coli* and *Asaia* sp. respectively) for an

overnight culture (24h) at 30°C. The overnight culture was then pelleted by centrifugation at

3000g, the medium discarded and resolved in 10% (w/v) honey water to the respective

working concentration for the experiments. The concentration of a typical overnight culture

was determined for *S. marcescens* and *Asaia* sp. by plating part of the overnight culture on

agar plates and counting CFUs after 24h or 48h of growth at 30°C, for *S. marcescens* and

Asaia sp. respectively. This yielded a concentration of $1.865 * 10^9 \pm 5.63 * 10^7$ (mean ± sd)

bacteria per ml for *S. marcescens* and $5.13 * 10^8 \pm 8.48 * 10^6$ (mean ± sd) bacteria for *Asaia*

sp.

Survival experiments. In a first survival experiment we tested whether the ability to perform

acidopore grooming within the first 24h after ingestion of pathogen contaminated food

provides a survival benefit for individual *C. floridanus* ants. Ants from eight colonies were

starved for 24-48h before start of the experiment, as described before, and then workers put

singly in small petri dishes were either given access to 5µl of *S. marcescens* contaminated

10% honey water ($9.33 * 10^9$ bacteria/ml; *Serratia+* ants: n = 127) or uncontaminated 10%

honey water (*Serratia-* ants: n = 135) for 2 min. Thereafter, all ants were cold anaesthetized

and approximately half of the *Serratia+* and the *Serratia-* ants (n = 65 and n = 69,

respectively) immobilized in a pipetting tip, thus preventing acidopore grooming (FA- ants: n

= 134) while the other half remained fully mobile (FA+ ants: n = 128). After 24h, FA- ants

were freed from the pipetting tip to minimize stress. Mortality of the ants was monitored over

5 days (120h) every 12h providing no additional food, except the one time feeding of 5µl

contaminated or uncontaminated honey water at the start of the experiment. We chose to

provide no additional food after the one time feeding at the beginning of the experiment, as an

altered feeding behavior, i.e. illness induced anorexia with known positive or negative effects

on survival (Hite et al., 2020), might otherwise have influenced our results.

In an additional survival experiment, we investigated whether the ability to acidify the crop

lumen has the potential to limit oral disease transmission during trophallactic food transfer.

To this end, *C. floridanus* ants from seven colonies were again starved, divided randomly in

two groups (donor and receiver ants, each n = 322) and their gaster marked with one of two

colours (Edding 751). Additionally, to prevent uptake of the poison, the acidopore opening of

all receiver ants (receiver FA-) and half of the donor ants (donor FA-) was sealed with

superglue, while the other half of the donor ants were sham treated (donor FA+) with a

droplet of superglue on their gaster (Tragust et al., 2013). We then paired these ants into two

different donor-receiver ant pairs. Pairs with both donor and receiver ants having their

acidopore sealed (donor FA- | receiver FA-) and pairs with only receiver ants having their

acidopore sealed (donor FA+ | receiver FA-). Six hours after pairing, donor ants from both

pairs were isolated and given access to 5µl of *S. marcescens* contaminated 10% honey water

$(1.865 * 10^9$ bacteria/ml) for 12h. Thereafter donor ants were again paired with the respective

receiver ants for 12 h and all pairs filmed for the first 30min. (Logitech webcam c910). These

videos were then analyzed for the duration of trophallaxis events donor-receiver ant pairs

engaged in during the first bout of trophallactic food exchange. After this first feeding round,

donor ants were fed in the same fashion, i.e. isolation for 12h with access to *S. marcescens*

contaminated 10% honey water, every 48h, while they were maintained with the respective

receiver ants for the rest of the time. This experimental design ensured that receiver ants were

fed only through the respective donor ants with pathogen contaminated food. Survival of both,

donor and receiver ants, was monitored daily for a total of 12 days.

Bacterial viability and growth assays. We tested the ability of *S. marcescens* and *Asaia* sp.

to withstand acidic environments *in vitro*, as well as their ability and the ability of *E. coli* to

pass from the crop to the midgut *in vivo* when ingested together with food. In ants, gut

morphological structures, i.e. the infrabuccal pocket, an invagination of the hypopharynx in

the oral cavity (Eisner and Happ, 1962), and the proventriculus, a valve that mechanically

restricts passage of fluids from the crop to the midgut (Eisner and Wilson, 1952),

consecutively filter solid particles down to 2µm (Lanan et al., 2016) which would allow *S.*

marcescens (Ø: 0.5-0.8µm, length: 0.9-2µm Grimont and Grimont, 2006), *Asaia* sp. (Ø: 0.4-

1µm, length: 0.8-2.5µm Komagata et al., 2014), and *E. coli* (length: 1µm, width: 0.35µm,

Blount, 2015) to pass. For the *in vitro* tests we incubated a diluted bacterial overnight culture

(10^5 and 10^4 CFU/ml for *S. marcescens* and *Asaia* sp., respectively) in 10% honey water (pH

= 5) and in 10% honey water acidified with commercial formic acid to a pH of 4, 3 or 2 for 2h

at room temperature (*S. marcescens*: n = 15 for all pH-levels, except pH = 4 with n = 13;

Asaia sp.: n = 10). Then we plated 100µl of the bacterial solutions on agar-medium (LB-

medium and GLY-medium for *S. marcescens* and *Asaia* sp., respectively) and incubated them

at 30°C for 24h (*S. marcescens*) or 48h (*Asaia* sp.) before counting the number of formed

CFUs. For the *in vivo* tests *C. floridanus* ants from five (*Asaia* sp.), four (*E. coli*) or from six

colonies (*S. marcescens*) were starved as before and then individually given access to 5µl of

bacteria contaminated 10% honey water (*Asaia* sp. and *E. coli*: $1 * 10^7$ CFU/ml, *S.*

marcescens: $1 * 10^6$ CFU/ml) for 2 min. To assess the number of CFUs in the digestive tract,

i.e. the crop and the midgut, ants were dissected either directly after feeding (0h; *S.*

marcescens: n = 60 workers; *Asaia* sp. and *E. coli*: n = 15 each), or at 0.5h (*S. marcescens*: n

= 60; *Asaia* sp. and *E. coli*: n = 15 each), 4h (*S. marcescens*: n = 60; *Asaia* sp. and *E. coli*: n =

15 each), 24h (*S. marcescens*: n = 53; *Asaia* sp. and *E. coli*: n = 15 each) or 48h (*S.*

marcescens: n = 19; *Asaia* sp. and *E. coli*: n = 15 each) after feeding. For dissection, ants were

cold anesthetized, the gaster opened and the whole gut detached. The crop and the midgut

were then separated from the digestive tract, placed in a reaction tube, mechanically crushed

with a sterile pestle and dissolved in 100µl (*Asaia* sp. and *E. coli*) or 150µl (*S. marcescens*)

phosphate buffered saline (PBS-buffer: 8.74g NaCl, 1.78g $Na_2HPO_4,2H_2O$ in 1L MilliQ-

water adjusted to a pH of 6.5). The resulting solutions were then thoroughly mixed, 100µl

streaked on agar-medium (LB-medium and GLY-medium for *S. marcescens*/*E.coli* and *Asaia*

sp., respectively) and incubated at 30°C for 24h (*S. marcescens* and *E. coli*) or 48h (*Asaia*

sp.), before counting the number of formed CFUs. No other bacteria (e.g. resident microbes)

were apparent in terms of a different CFU morphology on the agar plates which agrees with

the very low number of cultivable resident bacteria present in the midgut of *C. floridanus*

(Stoll and Gross, unpublished results). This methodology cannot completely exclude that

resident *S. marcescens* or species of Acetobacteraceae might have biased our count data by

adding a background level of CFUs at all timepoints or by adding random outlier CFUs at

specific timepoints. Both, background level CFU numbers and random outlier CFUs should

however not influence observed patterns over time. The timepoints of 0h, 0.5h, 4h, 24h and

48h in *in vivo* bacterial growth assays were chosen according to literature describing passage

of food from the crop to the midgut within 3-6h after food consumption in ants (Cannon,

1998, Gösswald and Kloft, 1960b, Gösswald and Kloft, 1960a, Howard and Tschinkel, 1981,

Markin, 1970). They should thus be representative of two time points before food passage

from the crop to the midgut (0.5h and 4h) and two time points after food passage from the

crop to the midgut (24h and 48h) together with the reference timepoint (0h).

Food passage experiment. To estimate food passage from the crop to the midgut and hindgut

of *C. floridanus* after feeding we performed the following experiment. We again took a cohort

of ~100 workers out of one natal colony of *C. floridanus*, starved them for 24h and then

offered them 200 µl of a 1:1 honey-water mix with 50mg of polymethylmethacrylate

(PMMA, aka acrylic glass) particles (size ≤ 40 µm). Thereafter, we dissected the digestive

tract of three major and three minor workers at each of the timepoints 2h, 4h, 6h, 8h, 12h,

14h, 16h, 18h, 24h, and 48h after feeding and placed each under a microscope (Leica DM

2000 LED) to detect and count the number of particles via fluorescence in the crop, the

midgut and the hindgut.

Statistical analyses. All statistical analyses were performed with the R statistical

programming language (version 3.6.1, R Core Team, 2019). All (zero-inflated) General(ized)

linear and mixed models and Cox mixed-effects models were compared to null (intercept

only) or reduced models (for those with multiple predictors) using Likelihood Ratio (LR) tests

to assess the significance of predictors. Pairwise comparisons between factor levels of a

significant predictor were performed using pairwise post-hoc tests adjusting the family-wise

error rate according to the method of Westfall (package "multcomp", Bretz et al., 2011). We

checked necessary model assumptions of (zero-inflated) General(ised) linear and mixed

models using model diagnostic tests and plots implemented in the package "DHARMa"

(Hartig, 2019). Acidity of the crop lumen (log transformed pH to normalize data) and midgut

lumen in *C. floridanus* was analyzed using linear mixed models (LMM, package "lme4",

Bates et al., 2015) including time since feeding (four levels: 0+4h, 0+24h, 0+48h, 48h+4h;

Fig. 1a), ant manipulation (two levels: FA+ and FA-, i.e. ants with and without acidopore

access; Fig. 1b) or digestive tract part (four levels: crop, midgut position 1, midgut position 2,

midgut position 3, midgut position 4; Fig. 1 – figure supplement 1) as predictors and natal

colony as a random effect. Due to non-normality and heteroscedasticity, the acidity of the

crop lumen in the seven formicine ant species other than *C. floridanus* (Fig. 1c) was analysed

using per species Wilcoxon Rank Sum tests with ant manipulation (FA+ and FA-) as

predictor. The frequency of acidopore grooming in *C. floridanus* upon feeding different types

of fluids was analyzed using Generalized linear mixed models (GLMM, package "lme4",

Bates et al., 2015) with negative binomial errors and type of fluid (three levels: unfed, water-

fed or 10% honey water fed) as predictor and natal colony as random effect (Fig. 1 – figure

supplement 2).

Survival data were analysed with Cox mixed effects models (COXME, package "coxme",

Therneau, 2019). For the survival of individual ants (Fig. 3), ant treatment (four levels:

Serratia- | FA-, *Serratia-* | FA+, *Serratia+* | FA-, *Serratia+* | FA+) was added as a predictor

and the three "blocks" in which the experiment was run and the colony ants originated from,

were included as two random intercept effects. For the survival of donor-receiver ant pairs

(Fig. 4), ant treatment (four levels: donor FA+, donor FA-, receiver FA+, receiver FA-) was

included as a predictor and the three "blocks" in which the experiment was run, the colony

ants originated from, and petri dish in which donor and receiver ants were paired, were

included as three random intercept effects. Survival of receiver ants was right censored if the

corresponding donor ant died at the next feeding bout (right censoring of both donor and

receiver ants in one pair upon death of one of the ants yielded statistically the same result:

COXME, overall LR χ^2 = 60.202, df = 3, P < 0.001; post-hoc comparisons: receiver FA- vs

donor FA-: *P* = 0.388, all other comparisons: P < 0.001). The duration of trophallaxis events

(square-root transformed to normalize data) between donor-receiver ant pairs was analysed

using a linear mixed model with ant pair type (two levels: donor FA+ | receiver FA- and

donor FA- | receiver FA-) as predictor and the three "blocks", in which the experiment was

run and the colony ants originated from as random effect (Fig. 4 – figure supplement 1).

Bacterial growth *in vitro* was analysed separately for *S. marcescens* and *Asaia* sp. using

Generalized linear models (GLM) with negative binomial errors and pH as predictor,

excluding pH levels with zero bacterial growth due to complete data separation (Fig. 2 -

supplementary figure 2 and Fig. 5 – figure supplement 1). Relative values shown in Fig. 2 -

supplementary figure 2 and Fig. 5 – figure supplement 1 were calculated by dividing each

single number of formed CFUs at the different pH-values through the mean of formed CFUs

at pH 5. Bacterial viability *in vivo* within the digestive tract of *C. floridanus* over time was

analysed separately for the crop and midgut for *S. marcescens* and *Asaia* sp. (Fig. 2 and Fig.

5, respectively) and for *E. coli* (Fig. 5 – figure supplement 2). Zero-inflated generalized linear

mixed models with negative binomial errors (package "glmmTMB", Brooks et al., 2017) were

used to model CFU number, with time after feeding as fixed predictor and ant colony as

random effect, except for the *E. coli* model in the crop where colony was included as fixed

factor as the model did not converge with colony as a random factor. Timepoints with zero

bacterial growth were again excluded in the models. Relative CFU values shown in Fig. 2,

Fig. 5, and Fig. 5 – supplementary figure 2 were calculated by dividing single CFU-values

through the mean of CFU-values at timepoint 0h in the crop. Proportions and percentages of

relative CFU change in the text are based on these relative CFU values.

Acknowledgements

We would like to thank Robert Paxton for English grammar and style check of a pre-submission version of the manuscript, Franziska Vogel, Marvin Gilliar, and Martin Wolak for part of the data collection, Elena Crotti and Daniele Daffonchio for providing the *Asaia* strain and Martin Kaltenpoth for access to the pH microelectrode.

Author contributions

S.T. and H.F. conceived the experiments. S.T. and M.A.M. performed the survival assays and the behavioral observations. C. H., C. T., R. B., H. F., and J. H. measured crop lumen acidity. H. F. measured pH in the midgut. M. H. and C. H. performed *in vivo* bacterial viability measurements. C. T. performed the *in vitro* bacterial growth measurements. S.T. analyzed the data and prepared the manuscript. S.T., C.H., J.H., R.B., C.T., M.H., M.A.M., R.G. and H.F. edited the manuscript.

Competing interests

The authors declare no competing interests.

Data and code availability

The authors declare that all data supporting the findings of this study and all code required to reproduce the analyses and figures of this study are available within the article and its supplementary information and will be made publicly available at the DRYAD digital repository upon acceptance.

References

ALEXANDER, R. D. 1974. The evolution of social behavior. *Annual Review of Ecology and Systematics,*
5, 325-383. doi: 10.1146/annurev.es.05.110174.001545
ALLAIRE, J. M., CROWLEY, S. M., LAW, H. T., CHANG, S. Y., KO, H. J. & VALLANCE, B. A. 2018. The
intestinal epithelium: central coordinator of mucosal immunity. *Trends in Immunology,* 39,
677-696. doi: 10.1016/j.it.2018.04.002
ANDERSON, K. E., RUSSELL, J. A., MOREAU, C. S., KAUTZ, S., SULLAM, K. E., HU, Y., BASINGER, U.,
MOTT, B. M., BUCK, N. & WHEELER, D. E. 2012. Highly similar microbial communities are
shared among related and trophically similar ant species. *Molecular Ecology,* 21, 2282-2296.
doi: 10.1111/j.1365-294X.2011.05464.x
ARCHETTI, M., SCHEURING, I., HOFFMAN, M., FREDERICKSON, M. E., PIERCE, N. E. & YU, D. W. 2011a.
Economic game theory for mutualism and cooperation. *Ecology Letters,* 14, 1300-12. doi:
10.1111/j.1461-0248.2011.01697.x
ARCHETTI, M., UBEDA, F., FUDENBERG, D., GREEN, J., PIERCE, N. E. & YU, D. W. 2011b. Let the right
one in: a microeconomic approach to partner choice in mutualisms. *American Naturalist,*
177, 75-85. doi: 10.1086/657622
ATTYGALLE, A. B. & MORGAN, E. D. 1984. Chemicals from the Glands of Ants. *Chemical Society*
Reviews, 13, 245-278. doi: 10.1039/cs9841300245
BASIBUYUK, H. H. & QUICKE, D. L. J. 1999. Grooming behaviours in the Hymenoptera (Insecta):
Potential phylogenetic significance. *Zoological Journal of the Linnean Society,* 125, 349-382.
doi: 10.1006/zjls.1997.0167
BATES, D., MAECHLER, M., BOLKER, B. & WALKER, S. 2015. Fitting Linear mixed-effects models using
lme4. *Journal of Statistical Software,* 67, 1-48. doi: 10.18637/jss.v067.i01
BEASLEY, D. E., KOLTZ, A. M., LAMBERT, J. E., FIERER, N. & DUNN, R. R. 2015. The evolution of
stomach acidity and its relevance to the human microbiome. *PloS One,* 10, e0134116. doi:

10.1371/journal.pone.0134116
BHATKAR, A. & WHITCOMB, W. H. 1971. Artificial diet for rearing various species of ants. *Florida*
Entomologist, 53, 229-232. doi: 10.2307/3493193
BIEDERMANN, P. H. & KALTENPOTH, M. 2014. New synthesis: the chemistry of partner choice in
insect-microbe mutualisms. *Journal of Chemical Ecology,* 40, 99. doi: 10.1007/s10886-014-
0382-8
BILLEN, J. 2009. *Diversity and morphology of exocrine glands in ants.,* Ouro Preto, Brazil, Universidade
Federal de Ouro Preto.
BILLEN, J. & SOBOTNIK, J. 2015. Insect exocrine glands. *Arthropod Structure & Development,* 44, 399-
400. doi: 10.1016/j.asd.2015.08.010
BILLEN, J. P. J. 1982. The Dufour gland closing apparatus in *Formica sanguinea* Latreille
(Hymenoptera, Formicidae). *Zoomorphology,* 99, 235-244. doi: 10.1007/BF00312297
BLOUNT, Z. D. 2015. The unexhausted potential of *E. coli. eLife,* 4, e05826. doi: 10.7554/eLife.05826
BLUM, J. E., FISCHER, C. N., MILES, J. & HANDELSMAN, J. 2013. Frequent replenishment sustains the
beneficial microbiome of *Drosophila melanogaster. mBio,* 4, e00860-13. doi:
10.1128/mBio.00860-13
BLÜTHGEN, N., STORK, N. E. & FIEDLER, K. 2004. Bottom-up control and co-occurrence in complex
communities: honeydew and nectar determine a rainforest ant mosaic. *Oikos,* 106, 344-358.
doi: 10.1111/j.0030-1299.2004.12687.x
BLÜTHGEN, N., VERHAAGH, M., GOITIA, W., JAFFE, K., MORAWETZ, W. & BARTHLOTT, W. 2000. How
plants shape the ant community in the Amazonian rainforest canopy: the key role of
extrafloral nectaries and homopteran honeydew. *Oecologia,* 125, 229-240. doi:
10.1007/s004420000449

BOOMSMA, J. J., SCHMID-HEMPEL, P. & HUGHES, W. O. H. 2005. Life histories and parasite pressure
across the major groups of social insects. *In:* FELLOWES, M. H. G. & ROLFF, J. (eds.) *Insect*
Evolutionary Ecology. Oxon, UK: CABI.
BRETZ, F., HOTHORN, T. & WESTFALL, P. 2011. *Multiple Comparisons using R.,* Boca Raton, FL, CRC
Press.
BRINKER, P., FONTAINE, M. C., BEUKEBOOM, L. W. & FALCAO SALLES, J. 2019a. Host, symbionts, and
the microbiome: the missing tripartite interaction. *Trends in Microbiology, 27,* 480-488. doi:
10.1016/j.tim.2019.02.002
BRINKER, P., WEIG, A., RAMBOLD, G., FELDHAAR, H. & TRAGUST, S. 2019b. Microbial community
composition of nest-carton and adjoining soil of the ant *Lasius fuliginosus* and the role of
host secretions in structuring microbial communities. *Fungal Ecology, 38,* 44-53. doi:
10.1016/j.funeco.2018.08.007
BRODERICK, N. A. & LEMAITRE, B. 2012. Gut-associated microbes of *Drosophila melanogaster. Gut*
Microbes, 3, 307-21. doi: 10.4161/gmic.19896
BROOKS, M. E., KRISTENSEN, K., VAN BENTHEM, K. J., MAGNUSSON, A., BERG, C. W., NIELSEN, A.,
SKAUG, H. J., MAECHLER, M. & BOLKER, B. M. 2017. glmmTMB balances speed and flexibility
among packages for zero-inflated generalized linear mixed modeling. *The R Journal, 9,* 378-
400. doi:
BROWN, B. P. & WERNEGREEN, J. J. 2016. Deep divergence and rapid evolutionary rates in gut-
associated Acetobacteraceae of ants. *BMC Microbiology, 16,* e140. doi: 10.1186/s12866-016-
0721-8
BROWN, B. P. & WERNEGREEN, J. J. 2019. Genomic erosion and extensive horizontal gene transfer in
gut-associated Acetobacteraceae. *BMC Genomics, 20,* e472. doi: 10.1186/s12864-019-5844-5
BRUNE, A. & DIETRICH, C. 2015. The gut microbiota of termites: Digesting the diversity in the light of
ecology and evolution. *Annual Review of Microbiology, 69,* 145-166. doi: 10.1146/annurev-
micro-092412-155715
BRÜTSCH, T., JAFFUEL, G., VALLAT, A., TURLINGS, T. C. & CHAPUISAT, M. 2017. Wood ants produce a
potent antimicrobial agent by applying formic acid on tree-collected resin. *Ecology and*
Evolution, 7, 2249-2254. doi: 10.1002/ece3.2834
BUFFIN, A., DENIS, D., VAN SIMAEYS, G., GOLDMAN, S. & DENEUBOURG, J. L. 2009. Feeding and
stocking up: Radio-labelled food reveals exchange patterns in ants. *PloS One, 4,* e5919. doi:
10.1371/journal.pone.0005919
BUFFIN, A., MAILLEUX, A.-C., DETRAIN, C. & DENEUBOURG, J. L. 2011. Trophallaxis in *Lasius niger*: a
variable frequency and constant duration for three food types. *Insectes Sociaux, 58,* 177-183.
doi: 10.1007/s00040-010-0133-y
BURKEPILE, D. E., PARKER, J. D., WOODSON, C. B., MILLS, H. J., KUBANEK, J., SOBECKY, P. A. & HAY,
M. E. 2006. Chemically mediated competition between microbes and animals: Microbes as
consumers in food webs. *Ecology, 87,* 2821-2831. doi: 10.1890/0012-
9658(2006)87[2821:Cmcbma]2.0.Co;2
CANNON, C. A. 1998. *Nutritional ecology of the carpenter ant Camponotus pennsylvanicus (De Geer):*
macronutrient preference and particle consumption., Virginia, USA, Doctor Thesis of
Philosophy in Entomology.
CARDOZA, Y. J., KLEPZIG, K. D. & RAFFA, K. F. 2006. Bacteria in oral secretions of an endophytic insect
inhibit antagonistic fungi. *Ecological Entomology, 31,* 636-645. doi: 10.1111/j.1365-
2311.2006.00829.x
CASEWELL, N. R., WUSTER, W., VONK, F. J., HARRISON, R. A. & FRY, B. G. 2013. Complex cocktails:
The evolutionary novelty of venoms. *Trends in Ecology & Evolution, 28,* 219-229. doi:
10.1016/j.tree.2012.10.020
CHAPMAN, R. F. 2013. *The insects: Structure and function.,* Cambridge, United Kingdom, Cambridge
University Press.

CHERRINGTON, C. A., HINTON, M., MEAD, G. C. & CHOPRA, I. 1991. Organic acids: chemistry,
antibacterial activity and practical applications. *Advances in Microbial Physiology,* 32**,** 87-108.
doi: 10.1016/s0065-2911(08)60006-5
CHOUAIA, B., GAIARSA, S., CROTTI, E., COMANDATORE, F., DEGLI ESPOSTI, M., RICCI, I., ALMA, A.,
FAVIA, G., BANDI, C. & DAFFONCHIO, D. 2014. Acetic acid bacteria genomes reveal functional
traits for adaptation to life in insect guts. *Genome Biology and Evolution,* 6**,** 912-920. doi:
10.1093/gbe/evu062
CHU, H. & MAZMANIAN, S. K. 2013. Innate immune recognition of the microbiota promotes host-
microbial symbiosis. *Nature Immunology,* 14**,** 668-75. doi: 10.1038/ni.2635
CHUA, K.-O., SEE-TOO, W.-S., TAN, J.-Y., SONG, S.-L., YONG, H.-S., YIN, W.-F. & CHAN, K.-G. 2020.
Oecophyllibacter saccharovorans gen. nov. sp. nov., a bacterial symbiont of the weaver ant
Oecophylla smaragdina with a plasmid-borne sole *rrn* operon. *bioRxiv***,** 2020.02.15.950782.
doi: 10.1101/2020.02.15.950782
CHUA, K. O., SONG, S. L., YONG, H. S., SEE-TOO, W. S., YIN, W. F. & CHAN, K. G. 2018. Microbial
community composition reveals spatial variation and distinctive core microbiome of the
weaver ant *Oecophylla smaragdina* in Malaysia. *Scientific Reports,* 8**,** e10777. doi:
10.1038/s41598-018-29159-2
CLEAR, M. R. & HORN, E. F. Y. 2019. The evolution of symbiotic plant-microbe signalling. *Annual Plant*
Reviews, 2**,** 1-52. doi: 10.1002/9781119312994.apr0684
COOLING, M. D., HOFFMANN, B. D., GRUBER, M. A. M. & LESTER, P. J. 2018. Indirect evidence of
pathogen-associated altered oocyte production in queens of the invasive yellow crazy ant,
Anoplolepis gracilipes, in Arnhem Land, Australia. *Bulletin of Entomological Research,* 108**,**
451-460. doi: 10.1017/S0007485317000967
COOPER, P. D. & VULCANO, R. 1997. Regulation of pH in the digestive system of the cricket,
Teleogryllus commodus Walker. *Journal of Insect Physiology,* 43**,** 495-499. doi:
10.1016/S0022-1910(97)85495-9
CROTTI, E., DAMIANI, C., PAJORO, M., GONELLA, E., RIZZI, A., RICCI, I., NEGRI, I., SCUPPA, P., ROSSI, P.,
BALLARINI, P., RADDADI, N., MARZORATI, M., SACCHI, L., CLEMENTI, E., GENCHI, M.,
MANDRIOLI, M., BANDI, C., FAVIA, G., ALMA, A. & DAFFONCHIO, D. 2009. *Asaia*, a versatile
acetic acid bacterial symbiont, capable of cross-colonizing insects of phylogenetically distant
genera and orders. *Environmental Microbiology,* 11**,** 3252-3264. doi: 10.1111/j.1462-
2920.2009.02048.x
CURRIE, C. R., SCOTT, J. A., SUMMERBELL, R. C. & MALLOCH, D. 1999. Fungus-growing ants use
antibiotic-producing bacteria to control garden parasites. *Nature,* 398**,** 701-704. doi:
10.1038/19519
DAVID, L. A., MAURICE, C. F., CARMODY, R. N., GOOTENBERG, D. B., BUTTON, J. E., WOLFE, B. E.,
LING, A. V., DEVLIN, A. S., VARMA, Y., FISCHBACH, M. A., BIDDINGER, S. B., DUTTON, R. J. &
TURNBAUGH, P. J. 2014. Diet rapidly and reproducibly alters the human gut microbiome.
Nature, 505**,** 559-63. doi: 10.1038/nature12820
DEGNAN, P. H., LAZARUS, A. B., BROCK, C. D. & WERNEGREEN, J. J. 2004. Host-symbiont stability and
fast evolutionary rates in an ant-bacterium association: cospeciation of *Camponotus* species
and their endosymbionts, *candidatus* Blochmannia. *Systematic Biology,* 53**,** 95-110. doi:
10.1080/10635150490264842
DEMAIN, A. L. & FANG, A. 2000. The Natural Functions of Secondary Metabolites. *In:* FIECHTER, A.
(ed.) *History of Modern Biotechnology I.* Berlin, Heidelberg: Springer Berlin Heidelberg.
DUARTE, A., WELCH, M., SWANNACK, C., WAGNER, J. & KILNER, R. M. 2018. Strategies for managing
rival bacterial communities: lessons from burying beetles. *Journal of Animal Ecology,* 87**,** 414-
427. doi: 10.1111/1365-2656.12725
EISNER, T. & HAPP, G. M. 1962. The infrabuccal pocket of a formicine ant: a social filtration device.
Psyche, 69**,** 107-116. doi: 10.1155/1962/25068
EISNER, T. & WILSON, E. O. 1952. The morphology of the porventriculus of a formicine ant. *Psyche,*
59**,** 47-60. doi: 10.1155/1952/14806

EMMERT, W. 1968. Die postembryonale Entwicklung sekretorischer Kopfdrüsen von *Formica*
pratensis Retz. und *Apis mellifica* L. (Ins., Hym.). *Zeitschrift für Morphologie der Tiere,* 63, 1-
62. doi: 10.1007/BF00343426
ENGEL, P. & MORAN, N. A. 2013. The gut microbiota of insects - diversity in structure and function.
FEMS Microbiology Reviews, 37, 699-735. doi: 10.1111/1574-6976.12025
FARISH, D. J. 1972. The evolutionary implications of qualitative variation in the grooming behaviour
of the Hymenoptera (Insecta). *Animal Behaviour,* 20, 662-76. doi: 10.1016/S0003-
3472(72)80139-8
FAVIA, G., RICCI, I., DAMIANI, C., RADDADI, N., CROTTI, E., MARZORATI, M., RIZZI, A., URSO, R.,
BRUSETTI, L., BORIN, S., MORA, D., SCUPPA, P., PASQUALINI, L., CLEMENTI, E., GENCHI, M.,
CORONA, S., NEGRI, I., GRANDI, G., ALMA, A., KRAMER, L., ESPOSITO, F., BANDI, C., SACCHI, L.
& DAFFONCHIO, D. 2007. Bacteria of the genus *Asaia* stably associate with *Anopheles*
stephensi, an Asian malarial mosquito vector. *Proceedings of the National Academy of*
Sciences of the United States of America, 104, 9047-9051. doi: 10.1073/pnas.0610451104
FELDHAAR, H., STRAKA, J., KRISCHKE, M., BERTHOLD, K., STOLL, S., MUELLER, M. J. & GROSS, R. 2007.
Nutritional upgrading for omnivorous carpenter ants by the endosymbiont *Blochmannia.*
BMC Biology, 5, 48. doi: 10.1186/1741-7007-5-48
FERNÁNDEZ-MARÍN, H., NASH, D. R., HIGGINBOTHAM, S., ESTRADA, C., VAN ZWEDEN, J. S.,
D'ETTORRE, P., WCISLO, W. T. & BOOMSMA, J. J. 2015. Functional role of phenylacetic acid
from metapleural gland secretions in controlling fungal pathogens in evolutionarily derived
leaf-cutting ants. *Proceedings of the Royal Society of London. Series B, Biological Sciences,*
282, 20150212. doi: 10.1098/rspb.2015.0212
FERNÁNDEZ-MARÍN, H., ZIMMERMAN, J. K., NASH, D. R., BOOMSMA, J. J. & WCISLO, W. T. 2009.
Reduced biological control and enhanced chemical pest management in the evolution of
fungus farming in ants. *Proceedings of the Royal Society of London. Series B, Biological*
sciences, 276, 2263-2269. doi: 10.1098/rspb.2009.0184
FERNÁNDEZ-MARÍN, H., ZIMMERMAN, J. K., REHNER, S. A. & WCISLO, W. T. 2006. Active use of the
metapleural glands by ants in controlling fungal infection. *Proceedings of the Royal Society of*
London. Series B, Biological sciences, 273, 1689-95. doi: 10.1098/rspb.2006.3492
FISCHBACH, M. A. & SEGRE, J. A. 2016. Signaling in host-associated microbial communities. *Cell,* 164,
1288-1300. doi: 10.1016/j.cell.2016.02.037
FLOREZ, L. V., BIEDERMANN, P. H. W., ENGL, T. & KALTENPOTH, M. 2015. Defensive symbioses of
animals with prokaryotic and eukaryotic microorganisms. *Natural Product Reports,* 32, 904-
936. doi: 10.1039/c5np00010f
FLOWER, N. E. & FILSHIE, B. K. 1976. Goblet cell membrane differentiations in the midgut of a
lepidopteran larva. *Journal of Cell Science,* 20, 357-75. doi:
FLYG, C., KENNE, K. & BOMAN, H. G. 1980. Insect pathogenic properties of *Serratia marcescens*:
Phage-resistant mutants with a decreased resistance to *Cecropia* immunity and a decreased
virulence to *Drosophila. Journal of General Microbiology,* 120, 173-181. doi:
10.1099/00221287-120-1-173
FOSTER, K. R., SCHLUTER, J., COYTE, K. Z. & RAKOFF-NAHOUM, S. 2017. The evolution of the host
microbiome as an ecosystem on a leash. *Nature,* 548, 43-51. doi: 10.1038/nature23292
GARCÍA-BAYONA, L. & COMSTOCK, L. E. 2018. Bacterial antagonism in host-associated microbial
communities. *Science,* 361. doi: 10.1126/science.aat2456
GERNAT, T., RAO, V. D., MIDDENDORF, M., DANKOWICZ, H., GOLDENFELD, N. & ROBINSON, G. E.
2018. Automated monitoring of behavior reveals bursty interaction patterns and rapid
spreading dynamics in honeybee social networks. *Proceedings of the National Academy of*
Sciences of the United States of America, 115, 1433-1438. doi: 10.1073/pnas.1713568115
GIANNELLA, R. A., ZAMCHECK, N. & BROITMAN, S. A. 1972. Gastric-acid barrier to ingested
microorganisms in man - studies i*n-vivo* and *in-vitro. Gut,* 13, 251-256. doi:
10.1136/gut.13.4.251

GLANCEY, B. M., VANDER MEER, R. K., GLOVER, A., LOFGREN, C. S. & VINSON, S. B. 1981. Filtration of microparticles from liquids ingested by the red imported fire ant *Solenopsis invicta* Buren. *Insectes Sociaux, 28,* 295-401. doi: 10.1007/BF02224196

GÖßWALD, K. & KLOFT, W. 1960. Untersuchungen mit radioaktiven Isotopen an Waldameisen. *Entomophaga, 5,* 33-41. doi: 10.1007/bf02376705

GÖSSWALD, K. & KLOFT, W. 1960a. Neuere Untersuchungen über die sozialen Wechselbeziehungen im Ameisenvolk, durchgeführt mit Radio-Isotopen. *Zoologische Beiträge, 5,* 519-556. doi:

GÖSSWALD, K. & KLOFT, W. 1960b. Untersuchungen mit radioaktiven Isotopen an Waldameisen. *Entomophaga, 5,* 33-41. doi: 10.1007/bf02376705

GRAFEN, A. 1990. Biological signals as handicaps. *Journal of Theoretical Biology,* 144, 517-46. doi: 10.1016/s0022-5193(05)80088-8

GREENBERG, D. E., PORCELLA, S. F., STOCK, F., WONG, A., CONVILLE, P. S., MURRAY, P. R., HOLLAND, S. M. & ZELAZNY, A. M. 2006. *Granulibacter bethesdensis* gen. nov., sp. nov., a distinctive pathogenic acetic acid bacterium in the family Acetobacteraceae. *International Journal of Systematic and Evolutionary Microbiology,* 56, 2609-2616. doi: 10.1099/ijs.0.64412-0

GREENWALD, E., SEGRE, E. & FEINERMAN, O. 2015. Ant trophallactic networks: simultaneous measurement of interaction patterns and food dissemination. *Scientific Reports,* 5, 12496. doi: 10.1038/srep12496

GREENWALD, E. E., BALTIANSKY, L. & FEINERMAN, O. 2018. Individual crop loads provide local control for collective food intake in ant colonies. *eLife,* 7, e31730. doi: 10.7554/eLife.31730

GRIMONT, F. & GRIMONT, P. A. D. 2006. The genus *Serratia. In:* DWORKIN, M., FALKOW, S., ROSENBERG, E., SCHLEIFER, K.-H. & STACKEBRANDT, E. (eds.) *The Prokaryotes: Volume 6: Proteobacteria: Gamma Subclass.* New York, NY: Springer New York.

HAMILTON, C., LEJEUNE, B. T. & ROSENGAUS, R. B. 2011. Trophallaxis and prophylaxis: social immunity in the carpenter ant *Camponotus pennsylvanicus. Biology Letters,* 7, 89-92. doi: 10.1098/rsbl.2010.0466

HAMMER, T. J., JANZEN, D. H., HALLWACHS, W., JAFFE, S. P. & FIERER, N. 2017. Caterpillars lack a resident gut microbiome. *Proceedings of the National Academy of Sciences of the United States of America,* 114, 9641-9646. doi: 10.1073/pnas.1707186114

HARRISON, J. F. 2001. Insect acid-base physiology. *Annual Review of Entomology,* 46, 221-250. doi: 10.1146/annurev.ento.46.1.221

HARRISON, J. F., WONG, C. J. & PHILLIPS, J. E. 1992. Recovery from acute haemolymph acidosis in unfed locusts: I. Acid transfer to the alimentary lumen is the dominant mechanism. *Journal of Experimental Biology,* 165, 85-96. doi:

HARTIG, F. 2019. DHARMa: Residual diagnostics for hierarchical (multi-level/mixed) regression models. R package version 0.2.4. *https://CRAN.R-project.org/package=DHARMa.* doi:

HE, H., CHEN, Y. Y., ZHANG, Y. L. & WEI, C. 2011. Bacteria associated with gut lumen of *Camponotus japonicus* Mayr. *Environmental Entomology,* 40, 1405-1409. doi: 10.1603/En11157

HERMANN, H. R. & BLUM, M. S. 1968. The hymenopterous poison apparatus. VI. *Camponotus pennsylvanicus* (Hymenoptera: Formicidae). *Psyche,* 75, Article ID 70198 doi: 10.1155/1968/70198

HERTLE, R. 2005. The family of *Serratia* type pore forming toxins. *Current Protein and Peptide Science,* 6, 313-325. doi: 10.2174/1389203054546370

HERZNER, G., SCHLECHT, A., DOLLHOFER, V., PARZEFALL, C., HARRAR, K., KREUZER, A., PILSL, L. & RUTHER, J. 2013. Larvae of the parasitoid wasp *Ampulex compressa* sanitize their host, the American cockroach, with a blend of antimicrobials. *Proceedings of the National Academy of Sciences of the United States of America,* 110, 1369-1374. doi: 10.1073/pnas.1213384110

HERZNER, G. & STROHM, E. 2007. Fighting fungi with physics: food wrapping by a solitary wasp prevents water condensation. *Current Biology,* 17, R46-7. doi: 10.1016/j.cub.2006.11.060

HIRSHFIELD, I. N., TERZULLI, S. & O'BYRNE, C. 2003. Weak organic acids: a panoply of effects on bacteria. *Science Progress,* 86, 245-69. doi: 10.3184/003685003783238626

HITE, J. L., PFENNING, A. C. & CRESSLER, C. E. 2020. Starving the enemy? Feeding behavior shapes
host-parasite interactions. *Trends in Ecology & Evolution*, 35, 68-80. doi:
10.1016/j.tree.2019.08.004
HÖLLDOBLER, B. & WILSON, E. O. 1990. *The ants*, Cambridge, MA, USA, Belknap Press.
HOLTOF, M., LENAERTS, C., CULLEN, D. & VANDEN BROECK, J. 2019. Extracellular nutrient digestion
and absorption in the insect gut. *Cell and Tissue Research*, 377, 397-414. doi:
10.1007/s00441-019-03031-9
HOWARD, D. F. & TSCHINKEL, W. R. 1981. The flow of food in colonies of the fire ant, *Solenopsis*
invicta: a multifactorial study. *Physiological Entomology*, 6, 297-306. doi: 10.1111/j.1365-
3032.1981.tb00274.x
HOWDEN, C. W. & HUNT, R. H. 1987. Relationship between gastric-secretion and infection. *Gut*, 28,
96-107. doi: DOI 10.1136/gut.28.1.96
HUTKINS, R. W. 2019. Microbiology and technology of fermented foods. IFT Press.
IMHANN, F., BONDER, M. J., VICH VILA, A., FU, J., MUJAGIC, Z., VORK, L., TIGCHELAAR, E. F.,
JANKIPERSADSING, S. A., CENIT, M. C., HARMSEN, H. J., DIJKSTRA, G., FRANKE, L., XAVIER, R.
J., JONKERS, D., WIJMENGA, C., WEERSMA, R. K. & ZHERNAKOVA, A. 2016. Proton pump
inhibitors affect the gut microbiome. *Gut*, 65, 740-8. doi: 10.1136/gutjnl-2015-310376
IMIRZIAN, N., ZHANG, Y., KURZE, C., LORETO, R. G., CHEN, D. Z. & HUGHES, D. P. 2019. Automated
tracking and analysis of ant trajectories shows variation in forager exploration. *Scientific*
Reports, 9, 13246. doi: 10.1038/s41598-019-49655-3
INAGAKI, T. & MATSUURA, K. 2018. Extended mutualism between termites and gut microbes:
Nutritional symbionts contribute to nest hygiene. *The Science of Nature*, 105, 52. doi:
10.1007/s00114-018-1580-y
INNOCENT, T., HOLMES, N., BASSAM, M. A., SCHIØTT, M., SCHEURING, I., WILKINSON, B.,
HUTCHINGS, M. I., BOOMSMA, J. J. & YU, D. W. 2018. Experimental demonstration that
screening can enable the environmental recruitment of a defensive microbiome. *bioRxiv*. doi:
10.1101/375634
ITOH, H., JANG, S., TAKESHITA, K., OHBAYASHI, T., OHNISHI, N., MENG, X. Y., MITANI, Y. & KIKUCHI, Y.
2019. Host-symbiont specificity determined by microbe-microbe competition in an insect
gut. *Proceedings of the National Academy of Sciences of the United States of America*, 116,
22673-22682. doi: 10.1073/pnas.1912397116
IVENS, A. B. F., GADAU, A., KIERS, E. T. & KRONAUER, D. J. C. 2018. Can social partnerships influence
the microbiome? Insights from ant farmers and their trophobiont mutualists. *Molecular*
Ecology, 27, 1898-1914. doi: 10.1111/mec.14506
JANZEN, D. H. 1977. Why fruits rot, seeds mold, and meat spoils. *American Naturalist*, 111, 691-713.
doi: 10.1086/283200
JOOP, G., ROTH, O., SCHMID-HEMPEL, P. & KURTZ, J. 2014. Experimental evolution of external
immune defences in the red flour beetle. *Journal of Evolutionary Biology*, 27, 1562-1571. doi:
10.1111/jeb.12406
KAPPELER, P. M., CREMER, S. & NUNN, C. L. 2015. Sociality and health: impacts of sociality on disease
susceptibility and transmission in animal and human societies. *Philosophical Transactions of*
the Royal Society of London. Series B, Biological Sciences, 370, 20140116. doi:
10.1098/Rstb.2014.0116
KAUTZ, S., RUBIN, B. E. R. & MOREAU, C. S. 2013a. Bacterial infections across the ants: Frequency and
prevalence of *Wolbachia*, *Spiroplasma*, and *Asaia*. *Psyche*, e936341. doi:
10.1155/2013/936341
KAUTZ, S., RUBIN, B. E. R., RUSSELL, J. A. & MOREAU, C. S. 2013b. Surveying the microbiome of ants:
Comparing 454 pyrosequencing with traditional methods to uncover bacterial diversity.
Applied and Environmental Microbiology, 79, 525-534. doi: 10.1128/Aem.03107-12
KOELZ, H. R. 1992. Gastric-acid in vertebrates. *Scandinavian Journal of Gastroenterology*, 27, 2-6. doi:
10.3109/00365529209095998

1106 KOHL, E., HÖLLDOBLER, B. & BESTMANN, H. J. 2001. Trail and recruitment pheromones in
1107   *Camponotus socius* (Hymenoptera: Formicidae). *Chemecology,* 67-73. doi:
1108   10.1007/PL00001834
1109 KOHLMEIER, P., HOLLANDER, K. & MEUNIER, J. 2016. Survival after pathogen exposure in group-living
1110   insects: don't forget the stress of social isolation! *Journal of Evolutionary Biology,* 29, 1867-
1111   1872. doi: 10.1111/jeb.12916
1112 KOMAGATA, K., IINO, T. & YAMADA, Y. 2014. The family Acetobacteraceae. *In:* ROSENBERG, E.,
1113   DELONG, E. F., LORY, S., STACKEBRANDT, E. & THOMPSON, F. (eds.) *The Prokaryotes:*
1114   *Alphaproteobacteria and Betaproteobacteria.* Berlin, Heidelberg: Springer Berlin Heidelberg.
1115 KOTO, A., MERSCH, D., HOLLIS, B. & KELLER, L. 2015. Social isolation causes mortality by disrupting
1116   energy homeostasis in ants. *Behavioral Ecology and Sociobiology,* 69, 583-591. doi:
1117   10.1007/s00265-014-1869-6
1118 KWONG, W. K., MEDINA, L. A., KOCH, H., SING, K. W., SOH, E. J. Y., ASCHER, J. S., JAFFE, R. & MORAN,
1119   N. A. 2017. Dynamic microbiome evolution in social bees. *Science Advances,* 3, e1600513.
1120   doi: 10.1126/sciadv.1600513
1121 LANAN, M. C., RODRIGUES, P. A. P., AGELLON, A., JANSMA, P. & WHEELER, D. E. 2016. A bacterial
1122   filter protects and structures the gut microbiome of an insect. *The ISME Journal,* 10, 1866-
1123   1876. doi: 10.1038/ismej.2015.264
1124 LEBOEUF, A. C., WARIDEL, P., BRENT, C. S., GONCALVES, A. N., MENIN, L., ORTIZ, D., RIBA-GROGNUZ,
1125   O., KOTO, A., SOARES, Z. G., PRIVMAN, E., MISKA, E. A., BENTON, R. & KELLER, L. 2016. Oral
1126   transfer of chemical cues, growth proteins and hormones in social insects. *eLife,* 5, e20375.
1127   doi: 10.7554/eLife.20375
1128 LITTLE, A. E., MURAKAMI, T., MUELLER, U. G. & CURRIE, C. R. 2006. Defending against parasites:
1129   fungus-growing ants combine specialized behaviours and microbial symbionts to protect
1130   their fungus gardens. *Biology Letters,* 2, 12-16. doi: 10.1098/rsbl.2005.0371
1131 LÖFQVIST, J. 1976. Formic acid and saturated hydrocarbons as alarm pheromones for the ant
1132   *Formica rufa. Journal of Insect Physiology,* 22, 1331-1346. doi: 10.1016/0022-1910(76)90155-
1133   4
1134 LOPEZ, L. C., MORGAN, E. D. & BRAND, J. M. 1993. Hexadecanol and hexadecyl formate in the venom
1135   gland of formicine ants. *Philosophical Transactions of the Royal Society B-Biological Sciences,*
1136   341, 177-180. doi: 10.1098/rstb.1993.0101
1137 LUND, P., TRAMONTI, A. & DE BIASE, D. 2014. Coping with low pH: molecular strategies in
1138   neutralophilic bacteria. *FEMS Microbiology Reviews,* 38, 1091-125. doi: 10.1111/1574-
1139   6976.12076
1140 MAMLOUK, D. & GULLO, M. 2013. Acetic acid bacteria: Physiology and carbon sources oxidation.
1141   *Indian Journal of Microbiology,* 53, 377-384. doi: 10.1007/s12088-013-0414-z
1142 MARKIN, G. P. 1970. Food distribution within laboratory colonies of the argentine ant,*Tridomyrmex*
1143   *humilis* (Mayr). *Insectes Sociaux,* 17, 127-157. doi: 10.1007/BF02223074
1144 MARTINSEN, T. C., BERGH, K. & WALDUM, H. L. 2005. Gastric juice: A barrier against infectious
1145   diseases. *Basic & Clinical Pharmacology & Toxicology,* 96, 94-102. doi: 10.1111/j.1742-
1146   7843.2005.pto960202.x
1147 MATTHEWS, P. G. D. 2017. Acid–base regulation in insect haemolymph. *In:* WEIHRAUCH, D. &
1148   O'DONNELL, M. (eds.) *Acid-base balance and nitrogen excretion in invertebrates.* Springer,
1149   Cham.
1150 MAZEL, F., DAVIS, K. M., LOUDON, A., KWONG, W. K., GROUSSIN, M. & PARFREY, L. W. 2018. Is host
1151   filtering the main driver of phylosymbiosis across the tree of life? *mSystems,* 3, e00097-18.
1152   doi: 10.1128/mSystems.00097-18
1153 MCFALL-NGAI, M., HADFIELD, M. G., BOSCH, T. C., CAREY, H. V., DOMAZET-LOSO, T., DOUGLAS, A. E.,
1154   DUBILIER, N., EBERL, G., FUKAMI, T., GILBERT, S. F., HENTSCHEL, U., KING, N., KJELLEBERG, S.,
1155   KNOLL, A. H., KREMER, N., MAZMANIAN, S. K., METCALF, J. L., NEALSON, K., PIERCE, N. E.,
1156   RAWLS, J. F., REID, A., RUBY, E. G., RUMPHO, M., SANDERS, J. G., TAUTZ, D. & WERNEGREEN,
1157   J. J. 2013. Animals in a bacterial world, a new imperative for the life sciences. *Proceedings of*

the National Academy of Sciences of the United States of America, 110**,** 3229-36. doi:
10.1073/pnas.1218525110
MCFREDERICK, Q. S., CANNONE, J. J., GUTELL, R. R., KELLNER, K., PLOWES, R. M. & MUELLER, U. G.
2013. Specificity between lactobacilli and hymenopteran hosts is the exception rather than
the rule. *Applied and Environmental Microbiology,* 79**,** 1803-1812. doi: 10.1128/AEM.03681-
12
MIGUEL-ALIAGA, I., JASPER, H. & LEMAITRE, B. 2018. Anatomy and physiology of the digestive tract
of *Drosophila melanogaster. Genetics,* 210**,** 357-396. doi: 10.1534/genetics.118.300224
MILAN, N. F., KACSOH, B. Z. & SCHLENKE, T. A. 2012. Alcohol consumption as self-medication against
blood-borne parasites in the fruit fly. *Current Biology,* 22**,** 488-493. doi:
10.1016/j.cub.2012.01.045
MIRABITO, D. & ROSENGAUS, R. B. 2016. A double-edged sword? The cost of proctodeal trophallaxis
in termites. *Insectes Sociaux,* 63**,** 135-141. doi: 10.1007/s00040-015-0448-9
MOELLER, A. H., CARO-QUINTERO, A., MJUNGU, D., GEORGIEV, A. V., LONSDORF, E. V., MULLER, M.
N., PUSEY, A. E., PEETERS, M., HAHN, B. H. & OCHMAN, H. 2016. Cospeciation of gut
microbiota with hominids. *Science,* 353**,** 380-382. doi: 10.1126/science.aaf3951
MORAN, N. A., OCHMAN, H. & HAMMER, T. J. 2019. Evolutionary and ecological consequences of gut
microbial communities. *Annual Review of Ecology, Evolution, and Systematics,* 50**,** 451-475.
doi: 10.1146/annurev-ecolsys-110617-062453
MORGAN, D. E. 2008. Chemical sorcery for sociality: exocrine secretions of ants (Hymenoptera:
Formicidae). *Myrmecological News,* 11**,** 79-90. doi:
MOURA-ALVES, P., PUYSKENS, A., STINN, A., KLEMM, M., GUHLICH-BORNHOF, U., DORHOI, A.,
FURKERT, J., KREUCHWIG, A., PROTZE, J., LOZZA, L., PEI, G., SAIKALI, P., PERDOMO, C.,
MOLLENKOPF, H. J., HURWITZ, R., KIRSCHHOEFER, F., BRENNER-WEISS, G., WEINER, J., 3RD,
OSCHKINAT, H., KOLBE, M., KRAUSE, G. & KAUFMANN, S. H. E. 2019. Host monitoring of
quorum sensing during *Pseudomonas aeruginosa* infection. *Science,* 366**,** eaaw1629. doi:
10.1126/science.aaw1629
MUELLER, U. G., GERARDO, N. M., AANEN, D. K., SIX, D. L. & SCHULTZ, T. R. 2005. The evolution of
agriculture in insects. *Annual Review of Ecology Evolution and Systematics,* 36**,** 563-595. doi:
10.1146/annurev.ecolsys.36.102003.152626
MURESAN, C. I. & BUTTSTEDT, A. 2019. pH-dependent stability of honey bee (*Apis mellifera*) major
royal jelly proteins. *Scientific Reports,* 9**,** 9014. doi: 10.1038/s41598-019-45460-0
MUSHEGIAN, A. A. & EBERT, D. 2016. Rethinking "mutualism'' in diverse host-symbiont communities.
Bioessays, 38**,** 100-108. doi: 10.1002/bies.201500074
NEHME, N. T., LIEGEOIS, S., KELE, B., GIAMMARINARO, P., PRADEL, E., HOFFMANN, J. A., EWBANK, J.
J. & FERRANDON, D. 2007. A model of bacterial intestinal infections in *Drosophila*
melanogaster. PLoS Pathogens, 3**,** e173. doi: 10.1371/journal.ppat.0030173
NING, M., YUAN, M., LIU, M., GAO, Q., WEI, P., GU, W., WANG, W. & MENG, Q. 2018.
Characterization of cathepsin D from *Eriocheir sinensis* involved in *Spiroplasma eriocheiris*
infection. *Developmental & Comparative Immunology,* 86**,** 1-8. doi:
10.1016/j.dci.2018.04.018
OCHMAN, H., WOROBEY, M., KUO, C. H., NDJANGO, J. B. N., PEETERS, M., HAHN, B. H. &
HUGENHOLTZ, P. 2010. Evolutionary relationships of wild hominids recapitulated by gut
microbial communities. *PLoS Biology,* 8**,** e1000546. doi: 10.1371/journal.pbio.1000546
OHBAYASHI, T., TAKESHITA, K., KITAGAWA, W., NIKOH, N., KOGA, R., MENG, X. Y., TAGO, K., HORI, T.,
HAYATSU, M., ASANO, K., KAMAGATA, Y., LEE, B. L., FUKATSU, T. & KIKUCHI, Y. 2015. Insect's
intestinal organ for symbiont sorting. *Proceedings of the National Academy of Sciences of the*
United States of America, 112**,** E5179-E5188. doi: 10.1073/pnas.1511454112
ONCHURU, T. O., MARTINEZ, A. J., INGHAM, C. S. & KALTENPOTH, M. 2018. Transmission of
mutualistic bacteria in social and gregarious insects. *Current Opinion in Insect Science,* 28**,** 50-
58. doi: 10.1016/j.cois.2018.05.002

ONKEN, H. & MOFFETT, D. F. 2017. Acid–base loops in insect larvae with extremely alkaline midgut
regions. *In:* WEIHRAUCH, D. & O'DONNELL, M. (eds.) *Acid-base balance and nitrogen*
excretion in invertebrates. Springer, Cham.
OSMAN, M. F. & BRANDER, J. 1961. Weitere Beiträge zur Kenntnis der chemischen
Zusammensetzung des Giftes von Ameisen aus der Gattung *Formica. Zeitschrift für*
Naturforschung Part B-Chemie Biochemie Biophysik Biologie und verwandten Gebiete, B 16**,**
749-753. doi: 10.1515/znb-1961-1108
OSMAN, M. F. H. & KLOFT, W. 1961. Untersuchungen zur insektiziden Wirkung der verschiedenen
Bestandteile des Giftes der kleinen roten Waldameise *Formica polyctena* Foerst. *Insectes*
Sociaux, 8**,** 383-395. doi: 0.1007/BF02226557
OTTI, O., TRAGUST, S. & FELDHAAR, H. 2014. Unifying external and internal immune defences. *Trends*
in Ecology & Evolution, 29**,** 625-634. doi: 10.1016/j.tree.2014.09.002
OUDE ELFERINK, S. J., KROONEMAN, J., GOTTSCHAL, J. C., SPOELSTRA, S. F., FABER, F. & DRIEHUIS, F.
2001. Anaerobic conversion of lactic acid to acetic acid and 1, 2-propanediol by *Lactobacillus*
buchneri. Applied and Environmental Microbiology, 67**,** 125-32. doi: 10.1128/AEM.67.1.125-
132.2001
OVEREND, G., LUO, Y., HENDERSON, L., DOUGLAS, A. E., DAVIES, S. A. & DOW, J. A. 2016. Molecular
mechanism and functional significance of acid generation in the *Drosophila* midgut. *Scientific*
Reports, 6**,** 27242. doi: 10.1038/srep27242
PALMER-YOUNG, E. C., RAFFEL, T. R. & MCFREDERICK, Q. S. 2018. pH-mediated inhibition of a
bumble bee parasite by an intestinal symbiont. *Parasitology,* 146**,** 380-388. doi:
10.1017/S0031182018001555
PEREZ-COBAS, A. E., MAIQUES, E., ANGELOVA, A., CARRASCO, P., MOYA, A. & LATORRE, A. 2015. Diet
shapes the gut microbiota of the omnivorous cockroach *Blattella germanica. FEMS*
Microbiology Ecology, 91**,** fiv022. doi: 10.1093/femsec/fiv022
PULL, C. D., UGELVIG, L. V., WIESENHOFER, F., GRASSE, A. V., TRAGUST, S., SCHMITT, T., BROWN, M.
J. & CREMER, S. 2018. Destructive disinfection of infected brood prevents systemic disease
spread in ant colonies. *eLife,* 7**,** e32073. doi: 10.7554/eLife.32073
QUINLAN, R. J. & CHERRET, J. M. 1978. Studies on the role of the infrabuccal pocket of the leaf-
cutting ant *Acromyrmex octospinosus* (Reich) (Hym., Formicidae). *Insectes Sociaux,* 25**,** 237-
245. doi: 10.1007/BF02224744
RAKOFF-NAHOUM, S., PAGLINO, J., ESLAMI-VARZANEH, F., EDBERG, S. & MEDZHITOV, R. 2004.
Recognition of commensal microflora by toll-like receptors is required for intestinal
homeostasis. *Cell,* 118**,** 229-41. doi: 10.1016/j.cell.2004.07.002
RANGER, C. M., BIEDERMANN, P. H. W., PHUNTUMART, V., BELIGALA, G. U., GHOSH, S., PALMQUIST,
D. E., MUELLER, R., BARNETT, J., SCHULTZ, P. B., REDING, M. E. & BENZ, J. P. 2018. Symbiont
selection via alcohol benefits fungus farming by ambrosia beetles. *Proceedings of the*
National Academy of Sciences of the United States of America, 115**,** 4447-4452. doi:
10.1073/pnas.1716852115
RATZKA, C., LIANG, C., DANDEKAR, T., GROSS, R. & FELDHAAR, H. 2011. Immune response of the ant
Camponotus floridanus against pathogens and its obligate mutualistic endosymbiont. *Insect*
Biochemistry and Molecular Biology, 41**,** 529-36. doi: 10.1016/j.ibmb.2011.03.002
RATZKE, C., DENK, J. & GORE, J. 2018. Ecological suicide in microbes. *Nature Ecology & Evolution,* 2**,**
867-872. doi: 10.1038/s41559-018-0535-1
RATZKE, C. & GORE, J. 2018. Modifying and reacting to the environmental pH can drive bacterial
interactions. *PloS Biology,* 16**,** e2004248. doi: 10.1371/journal.pbio.2004248
R CORE TEAM 2019. R: A language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing.
REBOLLEDA-GOMEZ, M. & WOOD, C. W. 2019. Unclear intentions: eavesdropping in microbial and
plant systems. *Frontiers in Ecology and Evolution,* 7**,** 395. doi: 10.3389/fevo.2019.00385
REGNIER, F. E. & WILSON, E. O. 1968. The alarm-defence system of the ant *Acanthomyops claviger.*
Journal of Insect Physiology, 14**,** 955-970. doi: 10.1016/0022-1910(68)90006-1

ROH, S. W., NAM, Y. D., CHANG, H. W., KIM, K. H., KIM, M. S., RYU, J. H., KIM, S. H., LEE, W. J. & BAE,
J. W. 2008. Phylogenetic characterization of two novel commensal bacteria involved with
innate immune homeostasis in *Drosophila melanogaster*. *Applied and Environmental*
Microbiology, 74**,** 6171-6177. doi: 10.1128/AEM.00301-08
RUSSELL, J. A., SANDERS, J. G. & MOREAU, C. S. 2017. Hotspots for symbiosis: function, evolution, and
specificity of ant-microbe associations from trunk to tips of the ant phylogeny (Hymenoptera:
Formicidae). *Myrmecological News,* 24**,** 43-69. doi: 10.25849/myrmecol.news_024:043
RYU, J. H., KIM, S. H., LEE, H. Y., BAI, J. Y., NAM, Y. D., BAE, J. W., LEE, D. G., SHIN, S. C., HA, E. M. &
LEE, W. J. 2008. Innate immune homeostasis by the homeobox gene *caudal* and commensal-
gut mutualism in *Drosophila*. *Science,* 319**,** 777-782. doi: 10.1126/science.1149357
SALEM, H., FLOREZ, L., GERARDO, N. & KALTENPOTH, M. 2015. An out-of-body experience: the
extracellular dimension for the transmission of mutualistic bacteria in insects. *Proceedings of*
the Royal Society B-Biological Sciences, 282**,** 20142957. doi: 10.1098/rspb.2014.2957
SCHEURING, I. & YU, D. W. 2012. How to assemble a beneficial microbiome in three easy steps.
Ecology Letters, 15**,** 1300-1307. doi: 10.1111/j.1461-0248.2012.01853.x
SCHMIDT, J. O. 1986. Chemistry, pharmacology, and chemical ecology of ant venoms. *In:* PIEK, T. (ed.)
Venoms of the Hymenoptera: biochemical, pharmacological and behavioral aspects. Orlando,
Florida: Academic Press.
SCHOETERS, E. & BILLEN, J. 1996. The control apparatus of the venom gland in formicine ants
(Hymenoptera: Formicidae). *Netherlands Journal of Zoology,* 46**,** 281-289. doi:
10.1163/156854295X00230
SCOTT, J. J., OH, D. C., YUCEER, M. C., KLEPZIG, K. D., CLARDY, J. & CURRIE, C. R. 2008. Bacterial
protection of beetle-fungus mutualism. *Science,* 322**,** 63-63. doi: 10.1126/science.1160423
SENDOVA-FRANKS, A. B., HAYWARD, R. K., WULF, B., KLIMEK, T., JAMES, R., PLANQUE, R., BRITTON,
N. F. & FRANKS, N. R. 2010. Emergency networking: Famine relief in ant colonies. *Animal*
Behaviour, 79**,** 473-485. doi: 10.1016/j.anbehav.2009.11.035
SHUKLA, S. P., PLATA, C., REICHELT, M., STEIGER, S., HECKEL, D. G., KALTENPOTH, M., VILCINSKAS, A.
& VOGEL, H. 2018a. Microbiome-assisted carrion preservation aids larval development in a
burying beetle. *Proceedings of the National Academy of Sciences of the United States of*
America, 115**,** 11274-11279. doi: 10.1073/pnas.1812808115
SHUKLA, S. P., VOGEL, H., HECKEL, D. G., VILCINSKAS, A. & KALTENPOTH, M. 2018b. Burying beetles
regulate the microbiome of carcasses and use it to transmit a core microbiota to their
offspring. *Molecular Ecology,* 27**,** 1980-1991. doi: 10.1111/mec.14269
SLACK, E., HAPFELMEIER, S., STECHER, B., VELYKOREDKO, Y., STOEL, M., LAWSON, M. A., GEUKING,
M. B., BEUTLER, B., TEDDER, T. F., HARDT, W. D., BERCIK, P., VERDU, E. F., MCCOY, K. D. &
MACPHERSON, A. J. 2009. Innate and adaptive immunity cooperate flexibly to maintain host-
microbiota mutualism. *Science,* 325**,** 617-20. doi: 10.1126/science.1172747
STROEYMEYT, N., GRASSE, A. V., CRESPI, A., MERSCH, D. P., CREMER, S. & KELLER, L. 2018. Social
network plasticity decreases disease transmission in a eusocial insect. *Science,* 362**,** 941-945.
doi: 10.1126/science.aat4793
STROHM, E., HERZNER, G., RUTHER, J., KALTENPOTH, M. & ENGL, T. 2019. Nitric oxide radicals are
emitted by wasp eggs to kill mold fungi. *eLife,* 8**,** e43718. doi: 10.7554/eLife.43718
STUCKI, D., FREITAK, D., BOS, N. & SUNDSTROM, L. 2019. Stress responses upon starvation and
exposure to bacteria in the ant *Formica exsecta*. *PeerJ,* 7**,** e6428. doi: 10.7717/peerj.6428
TENNANT, S. M., HARTLAND, E. L., PHUMOONNA, T., LYRAS, D., ROOD, J. I., ROBINS-BROWNE, R. M.
& VAN DRIEL, I. R. 2008. Influence of gastric acid on susceptibility to infection with ingested
bacterial pathogens. *Infection and Immunity,* 76**,** 639-645. doi: 10.1128/Iai.01138-07
TERRA, W. R. & FERREIRA 1994. Insect digestive enzymes: properties, compartmentalization and
function. *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry,* 109**,**
1-62. doi: 10.1016/0305-0491(94)90141-4
THERNEAU, T. 2019. coxme: Mixed Effects Cox Models. R package version 2.2-14. *http://CRAN.R-*
project.org/package=coxme. doi:

THERON, M. M. & RYKERS LUES, J. F. 2010. *Organic acids and food preservation,* Boca Raton, FL, USA,
CRC Press.
TRAGUST, S. 2016. External immune defence in ant societies (Hymenoptera: Formicidae): The role of
antimicrobial venom and metapleural gland secretion. *Myrmecological News,* 23**,** 119-128.
doi: 10.25849/myrmecol.news_023:119
TRAGUST, S., BRINKER, P., ROSSEL, N. & OTTI, O. 2020. Balancing life history investment decisions in
founding ant queens. *Frontiers in Ecology and Evolution,* 8**,** 76. doi: 10.3389/fevo.2020.00076
TRAGUST, S., MITTEREGGER, B., BARONE, V., KONRAD, M., UGELVIG, L. V. & CREMER, S. 2013. Ants
disinfect fungus-exposed brood by oral uptake and spread of their poison. *Current Biology,*
23**,** 76-82. doi: 10.1016/j.cub.2012.11.034
TRANIELLO, J. F. A. 1977. Recruitment behavior, orientation, and the organization of foraging in the
carpenter ant *Camponotus pennsylvanicus* de Geer (Hymenoptera: Formicidae). *Behavioral*
Ecology and Sociobiology, 2**,** 61-79. doi: 10.1007/BF00299289
TRIENENS, M., KELLER, N. P. & ROHLFS, M. 2010. Fruit, flies and filamentous fungi - experimental
analysis of animal-microbe competition using *Drosophila melanogaster* and *Aspergillus*
mould as a model system. *Oikos,* 119, 1765-1775. doi: 10.1111/j.1600-0706.2010.18088.x
VANDER MEER, R. 2012. Ant interactions with soil organisms and associated semiochemicals. *Journal*
of Chemical Ecology, 39**,** 728-745. doi: 10.1007/s10886-012-0140-8
VANDER WALL, S. B. 1990. *Food hoarding in animals,* Chicago, Illinois, USA, University of Chicago
Press.
VILLENA, J., KITAZAWA, H., VAN WEES, S. C. M., PIETERSE, C. M. J. & TAKAHASHI, H. 2018. Receptors
and signaling pathways for recognition of bacteria in livestock and crops: prospects for
beneficial microbes in healthy growth strategies. *Frontiers in Immunology,* 9**,** 2223. doi:
10.3389/fimmu.2018.02223
VOGEL, H., SHUKLA, S. P., ENGL, T., WEISS, B., FISCHER, R., STEIGER, S., HECKEL, D. G., KALTENPOTH,
M. & VILCINSKAS, A. 2017. The digestive and defensive basis of carcass utilization by the
burying beetle and its microbiota. *Nature Communications,* 8**,** 15186. doi:
10.1038/ncomms15186
WATNICK, P. I. & JUGDER, B. E. 2020. Microbial control of intestinal homeostasis via enteroendocrine
cell innate immune signaling. *Trends in Microbiology,* 28**,** 141-149. doi:
10.1016/j.tim.2019.09.005
WAY, M. J. 1963. Mutualism between ants and honeydew-producing homoptera. *Annual Review of*
Entomology, 8**,** 307-344. doi: DOI 10.1146/annurev.en.08.010163.001515
WILLIAMS, L. E. & WERNEGREEN, J. J. 2015. Genome evolution in an ancient bacteria-ant symbiosis:
parallel gene loss among *Blochmannia* spanning the origin of the ant tribe Camponotini.
PeerJ, 3**,** e881. doi: 10.7717/peerj.881
WILSON, E. O. & EISNER, T. 1957. Quantitative studies of liquid food transmission in ants. *Insectes*
Sociaux, 4**,** 157-166. doi: 10.1007/BF02224149
WOLFE, A. J. 2005. The acetate switch. *Microbiology and Molecular Biology Reviews,* 69**,** 12-50. doi:
10.1128/MMBR.69.1.12-50.2005
XIAO, R., WANG, X., XIE, E., JI, T., LI, X., MUHAMMAD, A., YIN, X., HOU, Y. & SHI, Z. 2019. An IMD-like
pathway mediates the intestinal immunity to modulate the homeostasis of gut microbiota in
Rhynchophorus ferrugineus Olivier (Coleoptera: Dryophthoridae). *Developmental &*
Comparative Immunology, 97**,** 20-27. doi: 10.1016/j.dci.2019.03.013
YEK, S. H. & MUELLER, U. G. 2011. The metapleural gland of ants. *Biological Reviews of the*
Cambridge Philosophical Society, 86**,** 774-91. doi: 10.1111/j.1469-185X.2010.00170.x
ZAHAVI, A. 1975. Mate selection-a selection for a handicap. *Journal of Theoretical Biology,* 53**,** 205-
14. doi: 10.1016/0022-5193(75)90111-3

1361

Fig. 1. Acidification of formicine ant crop lumens through swallowing of acidic poison

gland secretions. a, pH of crop lumens at 4h, 24h and 48h after feeding *C. floridanus* ants

10% honey water (pH = 5) at 0h and at 4h after re-feeding ants at 48h (LMM, LR-test, $\chi^2 =$

315.18, df = 3, *P* < 0.001, same letters indicate *P* = 0.317 and different letters indicate *P* <

0.001 in Westfall corrected post hoc comparisons). **b**, pH of crop lumens in *C. floridanus* ants

that were either prevented to ingest the formic acid containing poison gland secretion (FA-) or

not (FA+) for 24h after feeding (LMM, LR-test, χ^2 = 44.68, df = 1, *P* < 0.001). **c**, pH-value of

crop lumens 24h after feeding in seven formicine ant species that were either prevented to

ingest the formic acid containing poison gland secretion (FA-) or not (FA+). Wilcoxon rank

sum tests (two-sided). Lines and shaded boxes show median and interquartile range; points

show all data. Colours in shaded rectangles near y-axis represent universal indicator pH

colours. Colour filling of shaded boxes correspond to median pH colour of x-axis groups and

colour filling of points correspond to universal indicator pH colours. Border of shaded boxes

represents animal treatment (light grey: prevention of poison ingestion, FA-; dark grey:

poison ingestion not prevented, FA+).

1378

1379

1380

Fig. 1 – figure supplement 1. Acidity along the gastrointestinal tract of *C. floridanus*. pH-

measurements 24h after access to 10% honey-water in the crop (N = 2) and directly after the

proventriculus at four points along the midgut (N = 10 except position 4 with N = 9) (LMM,

LR-test, χ^2=22.152, df=4, *P* <0.001, same letters indicate *P* ≥ 0.443 and different letters

indicate *P* < 0.001 in Westfall corrected post hoc comparisons). Lines and shaded boxes show

median and interquartile range; points show all data. Colours in shaded rectangles near y-axis

represent universal indicator pH colours. Colour filling of shaded boxes correspond to median

pH colour of x-axis groups and colour filling of points correspond to universal indicator pH

colours.

1390

1391

Fig. 1 – figure supplement 2. Acidopore grooming frequency of *C. floridanus* after ingestion of different food types. Frequency of acidopore grooming within 30min. after fluid ingestion (water or 10% honey water) compared to ants that did not receive any fluid (unfed) (GLMM, LR-test, χ^2=33.526, df=2, *P* <0.001, same letters indicate *P* = 0.634 and different letters indicate *P* < 0.001 in Westfall corrected post hoc comparisons).

Fig. 1 – figure supplement 3. Baseline acidity of *C. floridanus* crop lumens under satiated and starved conditions. pH of crop lumens in *C. floridanus* workers (light grey: major workers, dark grey: minor workers) that were either taken directly out of a satiated colony or that were kept in cohorts of ~100 individuals under satiated conditions for three days and then starved for 24h before measuring the pH. Lines and shaded boxes show median and interquartile range; points show all data. Colours in shaded rectangles near y-axis represent universal indicator pH colours. Colour filling of shaded boxes correspond to median pH colour of x-axis groups and colour filling of points correspond to universal indicator pH

1406   colours. Border of shaded boxes represents animal caste (light grey: major workers; dark

1407   grey: minor workers).

1408

Fig. 2. Viability of *S. marcescens* over time in the digestive tract of *C. floridanus*. Change

in the number of colony forming units (CFUs) in the crop (**a**) and midgut (**b**) part of the

digestive tract (yellow colour in insert) relative to the mean CFU-number at 0h in the crop

(CFU change corresponds to single data CFU-value divided by mean CFU-value at 0h in the

crop), 0h, 0.5h, 4h, 24h, and 48h after feeding *Camponotus floridanus* ants 10% honey water

contaminated with *Serratia marcescens*. **a**, Change of *S. marcescens* in the crop (GLMM, LR-

test, $\chi^2 = 220.78$, df = 4, P <0.001, same letters indicate $P \geq 0.623$ and different letters indicate

$P < 0.001$ in Westfall corrected post hoc comparisons). **b**, Change of *S. marcescens* in the

midgut (GLMM, LR-test, $\chi^2 = 1.044$, df = 2, $P = 0.593$). Note that timepoints with zero

bacterial growth in the midgut (4h and 48h) were excluded from the statistical model.

1419

1420

Fig. 2 – figure supplement 1. Food passage of fluorescent particles through the digestive

tract of *C. floridanus***.** Number of fluorescent particles on a logarithmic scale in the crop

(dark blue), midgut (light blue) and hindgut (turquoise) part of the digestive tract of minor (**a**)

and major (**b**) ants 2h, 4h, 6h, 8h, 12h, 14h, 16h, 18h, 24h, and 48h after feeding them a 1:1

honey-water mix with polymethylmethacrylate (PMM) particles (size ≤ 40 µm). Note that for

displaying purposes and better visibility of zero values a value of one has been added to all

datapoints. Points represent the number of counted particles per individual ant and lines

connect the median value of particles at the different time points after feeding.

1429

1430

Fig. 2 – figure supplement 2. *S.* ***marcescens*** **growth** *in vitro***.** Change in the number of CFUs

relative to the mean at pH 5 (CFU change corresponds to single data CFU-value divided by

the mean CFU-value at pH 5) after incubation of *Serratia marcescens* in 10% honey water

(pH = 5) or in 10% honey water acidified with commercial formic acid to a pH of 4, 3 or 2 for

2h (GLM, LR-test, $\chi^2 = 79.442$, df = 1, $P < 0.001$). Note that pH-values with zero bacterial

growth (pH 2 and 3) were excluded from the statistical model.

1437

Fig. 3. Survival after ingestion of pathogen contaminated food. Survival of individual *C.*

floridanus ants that were either prevented to ingest the formic acid containing poison gland

secretion (FA-; ant outlines with blue dot) or not (FA+) after feeding them once either honey

water contaminated with *Serratia marcescens* (*Serratia+*, yellow circle with pink dots and

black ant outlines) or non-contaminated honey water (*Serratia-*) without providing food

thereafter (COXME, LR-test, χ^2 = 20.95, df=3, P = 0.0001, same letters indicate $P \geq 0.061$ and

different letters indicate $P \leq 0.027$ in Westfall corrected post hoc comparisons).

1446

1447

Fig. 4. Survival after sharing pathogen contaminated food via trophallaxis. Survival of

donor ants (light grey ant outlines) that were directly fed with pathogen contaminated food

(yellow circle with pink dots in insert) every other day and were either prevented to ingest

their formic acid containing poison gland secretion (FA-; ant outlines with blue dot) or not

(FA+) and survival of receiver ants (black ant outlines) that received pathogen contaminated

food only through trophallaxis with donor ants and were always prevented to ingest their

formic acid containing poison gland secretion (FA-) (COXME, LR-test, χ^2 = 66.68, df = 3, $P <$

0.001, same letters indicate $P = 0.309$ and different letters indicate $P \leq 0.002$ in Westfall

corrected post hoc comparisons).

1457

1458

Fig. 4 – figure supplement 1. Duration of trophallaxis in donor-receiver ant pairs. Total

duration of trophallaxis events within 30min. of the first bout of food exchange between

donor-receiver ant-pairs (LMM, LR-test, $\chi^2 = 1.23$, df = 1, $P = 0.268$). Donor ants in both

pairs were directly fed with *Serratia marcescens* contaminated 10% honey water and were

either prevented to ingest their formic acid containing poison gland secretion (FA-) or not

(FA+), while receiver ants received pathogen contaminated food only through trophallaxis

with the respective donor ants and were always prevented to ingest their formic acid

containing poison gland secretion (FA-).

1467

1468

Fig. 5. Viability of *Asaia* sp. over time in the digestive tract of *C. floridanus*. Change in the

number of colony forming units (CFUs) in the crop (**a**) and midgut (**b**) part of the digestive

tract (yellow colour in insert) relative to the mean CFU-number at 0h in the crop (CFU

change corresponds to single data CFU-values divided by mean CFU-value at 0h in the crop),

0h, 0.5h, 4h, 24h, and 48h after feeding ants 10% honey water contaminated with *Asaia* sp. **a**,

Change of *Asaia* sp. in the crop (GLMM; LR-test, $\chi^2 = 124.01$, df = 4,$P < 0.001$, same letters

indicate $P = 0.488$ and different letters indicate $P \leq 0.013$ in Westfall corrected post hoc

comparisons). **b**, Change of *Asaia* sp. in the midgut (GLMM; LR-test, $\chi^2 = 59.94$, df = 3,$P <$

0.001, same letters indicate $P = 0.116$ and different letters indicate $P \leq 0.005$ in Westfall

corrected post hoc comparisons). Note that timepoints with zero bacterial growth in the

midgut (0h) were excluded from the statistical model.

1480

Fig. 5 – figure supplement 1. *Asaia* **sp. growth** *in vitro***.** Change in the number of CFUs

relative to the mean at pH 5 (CFU change corresponds to single data CFU-value divided by

mean CFU-value at pH 5) after incubation of *Asaia* sp. in 10% honey water (pH = 5) or in

10% honey water acidified with commercial formic acid to a pH of 4, 3 or 2 for 2h (GLM,

LR-test χ^2 = 21.179, df = 2, *P* < 0.001, same letters indicate *P* = 0.234, and different letters

indicate *P* < 0.001 in Westfall corrected post hoc comparisons). Note that pH-values with zero

bacterial growth (pH 2) were excluded from the statistical model.

1488

1489

Fig. 5 – figure supplement 2. Viability of *E. coli* over time in the digestive tract of *C.*

***floridanus* over time.** Change in the number of colony forming units (CFUs) in the crop (**a**)

and midgut (**b**) part of the digestive tract (yellow colour in insert) relative to mean CFU-

number at 0h in the crop (CFU change corresponds to single data CFU-value divided by the

mean CFU-value at 0h in the crop), 0h, 0.5h, 4h, 24h, and 48h after feeding ants 10% honey

water contaminated with *Escherichia coli*. **a**, Change of *E. coli* in the crop (GLMM, LR-test,

$\chi^2 = 156.74$, df = 4, $P < 0.001$, same letters indicate $P = 0.979$ and different letters indicate $P <$

0.025 in Westfall corrected post hoc comparisons). **b**, Change of *E. coli* in the midgut

(GLMM, LR-test, $\chi^2 = 14.898$, df = 3, $P = 0.002$, same letters indicate $P \geq 0.629$ and different

letters indicate $P \leq 0.038$ in Westfall corrected post hoc comparisons). Note that timepoints

with zero bacterial growth in the midgut (0h) were excluded from the statistical model.

1501