Parallel processing of sensory cue and spatial information in the Dentate Gyrus

Sebnem N. Tuncdemir[1,2], Andres D. Grosmark[3], Gergely F. Turi[1,2], Amei Shank[4], Jack Bowler[3], Gokhan Ordek[2], Attila Losonczy[3], Rene Hen[1,2,5*], Clay Lacefield[1,2,5*]

[1]Department of Psychiatry, Columbia University, New York, NY 10032, USA
[2]Division of Systems Neuroscience, New York State Psychiatric Institute, New York, NY 10032, USA
[3]Department of Neuroscience, Mortimer B. Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY 10027, USA
[4]Columbia College, New York, NY 10027, USA
[5]Lead Contacts
*Correspondence: col8@cumc.columbia.edu or rh95@cumc.columbia.edu

Abstract

During exploration, animals form an internal map of an environment by combining information about specific sensory cues or landmarks with the animal's motion through space, a process which critically depends on the mammalian hippocampus. The dentate gyrus (DG) is the first stage of the hippocampal trisynaptic circuit where self-motion and sensory cue information are integrated, yet it remains unknown how neurons within the DG encode both cue related ("what") and spatial ("where") information during cognitive map formation. Using two photon calcium imaging in head fixed mice running on a treadmill, along with on-line sensory cue manipulation at specific track locations, we have identified robust sensory cue responses in DG granule cells largely independent of spatial location. Granule cell cue responses are stable for long periods of time, selective for the modality of the stimulus and accompanied by strong inhibition of the firing of other active neurons. At the same time, there is a smaller fraction of neurons whose firing is spatially tuned but insensitive to the presentation of nearby cues, similar to traditional place cells. These results demonstrate the existence of "cue cells" in addition to the better characterized "place cells" in the DG, an important heterogeneity that has been previously overlooked. We hypothesize that the granule cell population may support multiple channels of spatial and non-spatial information that contribute distinctly to local and down-stream computations and impact the role of the dentate gyrus in spatial navigation and episodic memory.

Introduction

An animal's location in an environment is highly relevant for guiding its behavior, both to find areas of potential reward and avoid areas of possible danger. The mammalian hippocampal formation plays a cardinal role in navigation and spatial memory by integrating self-motion and sensory cue information into a cognitive map of an environment, exemplified by the presence of "place cells" selective for specific locations within space[1]. As the initial stage in the 'trisynaptic circuit'[2], the dentate gyrus (DG) is the first region in the hippocampus to integrate sensory and self-motion information into a discrete spatial representation, and thus represents the most basic state of spatial map formation[3,4]. Yet causal evidence for conjunctive encoding of sensory and spatial information by the principle neurons of the DG, granule cells, is still lacking. While recent studies

46 have shown that granule cells may be activated by visual[5,6], tactile[7] and olfactory[8] cues, their
47 relationship to canonical 'place cells' and their influence on spatial map formation in the DG have
48 not been investigated.
49
The DG receives its main long-range excitatory inputs from the lateral and medial entorhinal
cortices (LEC and MEC, respectively), and sends mossy fiber projections only to area CA3[9]. The
LEC is thought to primarily represent information about sensory cues, while the MEC is thought
to more prominently encode self-motion information[10–12]. Since individual granule cell dendrites
receive projections from both of these areas, these inputs have the potential to be the basis for
dendritic computations that combine sensory and self-motion information into a discrete spatial
representation[6,13–16]. In concert with a highly effective winner-take-all process mediated by lateral
inhibition[17–19], these conjunctive representations could facilitate behavioral discrimination of
nearby salient locations with high spatial resolution[6,20,21]. Such processes may underlie proposed
computational roles of the DG in hippocampal information processing such as pattern separation,
where similar inputs are represented distinctly within the population in order to aid the selectivity
of spatial behavior and memory[18,22].
62
We sought to examine how the DG participates in spatial map formation by recording calcium
activity in large populations of granule cells in the mouse dorsal DG during head-fixed locomotion
on a treadmill. By controlling the administration of sensory cues and their pairing with the animal's
position on the treadmill, we were able to dissect sensory and spatial contributions to granule cell
firing. We found that surprisingly most of the task-associated neurons were highly sensitive to
specific sensory cues presented along the treadmill belt, rather than discrete locations. Cue
responses in single neurons were stable for long periods of time, sensory modality-selective and
led to inhibition of other active neurons. At the same time, a smaller fraction of neurons exhibited
robust spatial tuning independent of local cue presentation, yet were more context selective.
These two channels of information, sensory and spatial, were largely distinct within the granule
cell population, and led us to postulate the existence of "cue cells" in addition to the better
characterized "place cells" of the region. This work suggests that the DG maintains a largely
separate code for cues in an environment and their location; yet displays specific points of
integration, for example through mutual inhibition and spatial modulation of cue response
amplitudes. These properties of the heterogeneous population of cue cells and place cells may
play a role in higher level functions of the dentate gyrus such as pattern separation and contextual
encoding in spatial navigation.
80
81 **Results**
82
To investigate the interaction between sensory and spatial representations in the dentate gyrus
(DG), we recorded the activity of large populations of granule cells using two photon calcium
imaging in head-restrained mice running on a linear treadmill track[23]. Mice were injected
unilaterally with rAAV(1/2).Syn.GCaMP6s to express the genetically encoded calcium indicator
GCaMP6s in the dorsal DG. This was followed by the implantation of a chronic imaging window
above the hippocampal fissure, which allowed us to image the calcium dynamics of hundreds of
neurons in the granule cell layer simultaneously during exploratory behavior on the treadmill

(Figure 1a). After habituation to head fixation, mice were trained to run in order to receive
randomly delivered water rewards on a 2m-long treadmill belt, during 15 minute sessions (Figure
1b, Methods). Movies of population calcium imaging data were motion corrected offline[24], and the
activity of putative single neurons was isolated[25,26]. Neurons with significant spatial tuning of
calcium activity along the track were identified using previously described methods[23,27], and in a
subset of sessions activity of individual neurons was tracked in multiple conditions over multiple
sessions[28].
97
98   **Granule Cell Cue Responses**
99
Following initial training on the treadmill, we introduced a 1s odor pulse delivered in the middle of
the track on each lap as a dynamic spatial cue, in addition to an invariant tactile cue at the lap
boundary. We found that the majority of spatially selective neurons exhibited receptive fields near
the lap boundary and middle locations, corresponding to the lap cue and middle cue positions,
respectively (Figure 1c, 57% of cells with peak activity within 10cm of cues, 43% >10cm from
cues, $p<0.0001$, Mann-Whitney test). In order to normalize locomotion dependent modulation of
hippocampal activity[29,30], the treadmill was motorized at a constant speed, adjusted for each
mouse (motorized velocity = 10.11 ± 0.64 cm/s, self-driven velocity = 12.76 ± 2.45 cm/s, $p>0.05$,
Mann-Whitney test). In a separate cohort of mice advancing the treadmill belt through self-driven
locomotion, we confirmed that motorizing the treadmill does not alter the representation of
sensory cue information in the DG (Figures S1a-f).
111
Neurons responding reliably at cue locations could however be place cells that are enriched at
the locations of salient stimuli, as has been suggested in area CA1[31–36], or alternately could be
directly driven by the stimulus[8]. In order to dissociate cue responses from track location, the
olfactory cue was omitted or shifted $1/4^{th}$ of the track length (0.5m) once every 3-5 trials
interleaved throughout the session (41 ± 2 total laps/session). Under these conditions, a majority
of neurons normally active in the middle of the track shifted their firing position to match the new
location of the odor in cue-shift laps (Figure 1d, middle, 79.07 ± 2.76 %) and exhibited reduced
activity in cue-omitted laps (Figure 1d, right, 78.2 ± 3.87 %), compared with normal middle cue
laps (n = 285 spatially tuned neurons from 8 mice). In contrast, neurons firing at locations
corresponding to the invariant lap cue were unchanged in omit and shift laps. On trials with a
shifted middle cue, average firing rates were higher within the new cue region (Figure 1e, red,
inset $p=0.0234$ Wilcoxon signed-rank test), and lower at the normal cue location on cue-omitted
trials (blue, inset $p<0.00001$, Wilcoxon signed-rank test) compared to the corresponding positions
during normal trials. Rapid cue-associated activity and enrichment of highly tuned neurons at
positions corresponding to sensory cues were also seen in mice during self-driven locomotion on
the treadmill (Figure S1b,d). Thus, a substantial population of spatially tuned DG neurons in the
virtual linear track environment are in fact active directly in response to presentation of cues at
those locations, rather than the locations themselves.
130
Comparable population activity was found in response to cues of other sensory modalities, such
as visual or tactile cues (Figures S2a-d), as well for liquid rewards (Figures S2e-h), indicating
strong representation of diverse sensory variables in the DG. In addition, increasing the

complexity of the environment with two additional cues at other track locations results in an
additive pattern of single cue responses (Figures S2i-k), suggesting both that cue activity at
different locations is relatively independent and that strong cue responses are not limited to
situations with a single cue alone. Cue responses were also similar for cells imaged in the dorsal
DG of a granule cell-specific transgenic mouse (Dock10 Cre[37], Figures S3a-d,), indicating that
sensory cue representations are indeed a property of the granule cell population rather than
arising from other local neuron types such as mossy cells or inhibitory interneurons[7].
141
Next, we divided the spatially tuned population of granule cells into three groups for subsequent
analyses based on the position of their spatial fields and their activity during cue manipulation
(omit or shift) trials. The three response types are illustrated for one session (Figure 1f): 1) cells
with spatial fields within the middle cue region that closely track the changes in cue presentation
("odor-cue cells", left); 2) cells with spatial fields around the lap boundary cue on the treadmill belt
("lap-cue cells", center); and 3) the remaining spatially tuned cells with receptive fields outside of
the cue locations throughout the track ("place cells", right). These three groups, odor-cue, lap-
cue, and place cells, constituted $22.5 \pm 2\%$, $47.1 \pm 2\%$ and $30.4 \pm 1\%$ of the spatially tuned cells
within the imaging field of view, respectively (Figure S3e). We found that cells with similar
response types did not appear to cluster together spatially within the imaging field and the groups
did not exhibit significant differences in overall mean firing rates (Figures S3e-g). Spatial coding
properties however differed between cue and place coding populations of granule cells. Both
populations of cue cells (lap cue and olfactory cue) showed higher average spatial information
content[38] (Fig. 1g, $\chi^2=48.47$, $p<0.0001$, Kruskal-Wallis test) and had more consistent spatial firing
between the first and the second half of each session than place cells (Figure 1h, $\chi^2=32.60$,
$p<0.0001$, Kruskal-Wallis test).
158
In order to probe the emergence of selective firing in these populations, we identified the lap in
which responses began to robustly occur within the preferred spatial location of each cell during
the first session of exposure to the odor cue (field onset lap[39], see Methods). We found that for
odor and lap cue cells the majority had spatial fields that appeared within the first five laps (38/65
(58%), 75/136 (55%), respectively) while the majority of place cells emerged later in the session
(52/84 (62%) within 10 laps, Fig. 1i, $\chi^2=9.29$, $p=0.0096$, Kruskal-Wallis test). In agreement with
the later emergence of place cells, we found that the in-field firing rates of place cells were
significantly smaller than those of odor and lap cue cells only within the first 5 laps (Figure 1i,
inset, 2-way ANOVA, main effect of cell type: $F_{2,1473}=6.73$, $p=0.0012$; cell type × lap number
interaction $F_{4,1473}=2.44$, $p=0.04$; main effect of lap number: $F_{2,1473}=0.28$, $p=0.7$). Additional
experiments also showed that on an uncued treadmill belt prior to olfactory cue sessions, the
majority of 'future' odor cue cells were either inactive or had low spatial tuning (Figure S4),
suggesting that cue cells rapidly emerge *de novo* and are not place cells that remap to new cue
locations. Taken together, these results demonstrate that sensory cue representations are more
reliable and appear with less exposure than place cell representations, and arise from an
independent population of granule cells.
175
We performed two experiments in separate cohorts of mice to exclude the possibility that the cue
cells are tracking the periodicity of the stable locomotion on the motorized treadmill (i.e. are

instead 'time cells'). First, we directly compared the odor cue cell responses in mice running on a
motorized treadmill at a constant speed with responses from the same cells on a self-driven
treadmill at variable speeds in consecutive sessions (Figure 2a). Despite notable differences in
temporal occupancy of the cue location in the two conditions (Figure S1c), the same cells fired
reliably to the olfactory cue. Correlations in the spatial firing rates of individual cells show that
while both groups of cue cells were highly correlated between self-driven and motorized treadmill
conditions, place cells were significantly less consistent in sessions with different locomotive
behaviors (Figure 2b, p<0.001, Kruskall-Wallis test). Spatial population vector correlations of all
significantly tuned neurons between different locomotive behaviors also confirm highly correlated
activity selectively around the cue locations (Figure 2c). Second, we investigated the latency of
Ca^{2+} responses in sessions where the treadmill was motorized at different velocities each lap
(Figure 2c). We reasoned that if the cue cells were tracking the period from one cue delivery to
the next, cue-triggered response latencies would be proportional to the time it takes for the animal
to traverse a lap while position-triggered response latencies would be constant. On the contrary,
we found shorter response latencies from a constant position when the treadmill is motorized at
higher velocities and vice versa, with constant cue-triggered Ca^{2+} response latencies regardless
of the velocity of the animal (Figures 2e-f, S1g-h). And as in our previous experiments, omitting
the cue on intermittent laps still had the effect of eliminating the responses of cue cells on that
lap, confirming their acute cue sensitivity. These experiments demonstrate that sensory cue
responses occur independent of volitional locomotion and that cue cells are not in fact time cells.
Furthermore the differing responses of cue and place cells in these conditions supports our
previous findings that these are separate, dissociable population of granule cells in the DG. For
the rest of our study we will continue to call the remaining group of non cue-responsive neurons
"place cells", as they are spatially tuned cells that do not respond directly to local cues, and would
therefore be called place cells based upon standards used in previous work on hippocampal
spatial encoding. However, our results show that spatial responses in the majority of the dentate
gyrus place cells imaged in our experiments are highly sensitive to the behavioral mode as well
as the velocity of the mice, as has been shown in area CA1[36,40], suggesting that these "place
cells" respond to more than simple spatial information.
207
208   **Stability and Specificity of Sensory Cue Responses**
209
To further characterize the stability and coding specificity among the granule cell subpopulations,
we investigated the responses of individual neurons over time and with respect to different
sensory cues (Figure 3a, b). We tracked cells[28] over multiple sessions and were able to find
substantial numbers of the same cells active in different sessions within a day or 1 week later in
the same fields of view (Figures S5a, b). Although not all cells were identified in every session, a
similar percentage of spatially selective neurons was registered in all sessions, which was
confirmed by visual inspection to ensure that cells appeared consistent in the anatomical images
(Figures S5c-g).
218
Between any two sessions, over days or with different cue modalities, the cells encoding the
invariant lap cue were the largest fraction of cells that remained active and maintained their
response type (e.g. cue type and/or place, green circles in Figure 3c). Odor cue cells fired reliably

to the same olfactory cue over long periods of time (blue circles in red and blue shaded area,
Figure 3c), but were largely unresponsive to cues of other modalities presented at the same
position (green shaded area in Figure 3c, χ^2=19.78, p<0.0001, Kruskal-Wallis test). Conversely,
a lower percentage of place cells maintained their response type across days compared to cue
cells recorded in the same sessions using the same cues (orange circles in blue shaded area,
Figure 3c, χ^2=14.36, p<0.001, Kruskal-Wallis test). Furthermore when we examined specifically
whether cue cells become place cells between these sessions, or vice versa, we found that
categorical cue and place representations remain extremely stable within the DG population
(Figure 3d, p<0.001, Rank Sum test).

231
We further examined the degree of stability and specificity of cue and place cell tuning by cross-
correlating spatial firing rates for individual registered cells over time and with respect to different
sensory cues (Figure 3e). For this analysis, we computed the correlation of spatial firing rates for
cells registered during different sessions within a day and one week later, or with a different
sensory cue modality. In line with the stimulus selectivity analysis described above, we observed
that middle location odor cue cells displayed significantly lower correlations in sessions with a
different sensory cue, compared to separate sessions using the same olfactory cue on the same
day or one week later (left, χ^2=27.88, p<0.0001, Kruskal Wallis test). Thus, cells that responded
to a cue of one modality were unlikely to respond to cues of other modalities, despite a similar
spatial location of the cues. Correlations in the activity of individual place cells over 1 wk. were
significantly lower than on the same day, and were lower than both cue cell populations, again
indicating lower stability for place than cue representations (left, χ^2=18.79, p<0.0001, Kruskal
Wallis test). Lap cue cells did not display significant changes in their firing rate correlations
between sessions on the same day, across days or with different sensory cues, and were
therefore especially stable.

247
Responses of individual odor cue cells were also highly correlated in sessions recorded in
different rooms on the same day (Figures 3f-h, S5h), indicating that distal ambient cues that
differed between the rooms have a limited influence on the stability of sensory cue representation
in DG granule cells. Despite the high stability of cue cells, place cells were significantly less
consistent between rooms than cue cells in the same sessions (Figure 3g, h, p<0.001, Kruskall-
Wallis test), and were similar to that seen in place cells over 1wk when measured in the same
room (Figure 3e), suggesting that place cells are more context selective than cue cells.
Furthermore, cross-registered place cells did not become cue cells in the other room, and vice
versa (Figure 3i, p<0.01, Wilcoxon rank sum test). Taken together, these results suggest that
sensory cues are represented by a stable subpopulation of neurons that is highly selective for
specific cues while purely spatial representations are less stable and undergo progressive
reorganization over time and in different global contexts.

260
261 **Spatial Modulation of DG Cue Responses**

262
The juxtaposition of inputs from the lateral and medial entorhinal cortex onto the dendrites of
individual granule cells has been hypothesized to underlie a conjunctive code for sensory cues
and their spatial location[3,4]. We therefore examined the influence of spatial location on cue

responses ("spatial modulation") by tracking odor cue cells through multiple sessions with different
cue-location pairing, either with an intermittently shifted cue as in previous experiments or in
separate sessions with the same cue at random locations each lap (Figure 4a, b). On average,
cue-triggered Ca^{2+} response amplitudes for individual cue cells (a proxy for neuron action
potential burst firing rate) were smaller when cues were presented at the infrequent "shift" location
or at random locations, when compared to their responses at the more frequent middle location
(Figure 4c, d, $p<0.0001$, Friedman test, n=101, 5 mice). Thus while cue cells tend to respond to
the same sensory cue regardless of location, the strongest responses occur when a cue is
presented at the same place repeatedly.
275
To further examine the spatial modulation of cue responses, we performed a set of experiments
where a visual cue was delivered at multiple locations on the track to measure selectivity of
granule cell responses to cues presented consistently at distinct locations ("dual location cue",
Figure 4e), similar to recent work by other groups using virtual reality environments[41,42]. As in our
previous experiments with a single cue, each cue was omitted on a subset of laps to isolate cue
cells as opposed to place cells present in the vicinity of cues (Figures S6a, b). To measure the
extent of preference for one cue location versus the other, we calculated the spatial modulation
index for each cue cell as the ratio of the firing rates in the non-preferred cue location versus the
preferred cue location. Cue cells in the first session exhibited a range of spatial modulation
indices, but over the population their event rates were significantly modulated by the cue location
($p<0.00001$, Wilcoxon signed-rank test on day 1 spatial modulation index, 53% of individual
neurons significantly spatially modulated by location shuffle, Figure 4f). Spatial modulation also
increased over several days of dual cue presentations ($p=0.0125$ Wilcoxon rank sum test, day 1
vs 2, 3, or 4), with 81% of cue cells exhibiting significant spatial modulation by day 4 (Figures 4g-
h, S6c-d). This suggests that dentate cue cells are acutely modulated by the spatial location of
two identical cues but increase their preference for a single location over days.
292
293 **Effects of Cue Manipulation on Spatial Encoding in the DG**
294
While the most robust activity in the DG was found within the cue cell population, the existence
of the smaller population of place cells in uncued locations suggests that spatial activity in these
cells is referenced to one of the two cues on the otherwise featureless treadmill track in order to
encode a unique location. We therefore examined the effect of manipulations of the variable cue
on the subsequent spatial encoding of place cells, in order to judge the degree to which this cue
acts as a landmark. For example, if place cells were acutely referenced to the nearest cue we
would expect the place fields of all cells following the variable middle cue to shift on cue shifted
laps. To examine the relationship between spatial firing patterns in normal cue laps to those in
cue shifted laps, we first calculated population vector (PV) correlations of firing rates across all
spatially tuned cells (cue and place cells) on each lap (Figure 5a). While the correlation was higher
in the vicinity of the cues between normal laps, we observed a dramatic decrease in PV correlation
between normal and cue shifted laps which was confined to the area immediately around the cue
itself. However, this change in PV correlation did not extend much beyond the cue location,
consistent with a limited effect of cue shifting on subsequent place cell activity.
309

To further evaluate the manner in which cue associated activity contributes to spatial encoding, a
spatial Bayesian decoder was constructed from the firing rate vectors of all spatially tuned cells
(4,091 cells from 66 sessions)[27]. In order to establish a non-biased estimate of the position during
treadmill running, the decoding was performed using a 5 fold cross validation approach in which
1/5th of decoded laps were held out from the training set. Post-reconstruction, we divided the data
according to the lap types (Figure 5b). The decoder accuracy was higher in normal (median, 10.6
cm) compared to both cue shift (median, 19.6 cm) and cue omitted laps (median, 18.4 cm,
p<0.0001, Wilcoxon rank sum test, Figure 5c, data not shown for omit laps), indicating that cue
manipulations affect the accuracy of spatial coding by the DG population. However while the cue
shift strongly perturbed spatial decoding near the shifted cue, estimating the animal's position to
be farther along than the real position, the decoder error on shift laps soon converged to that of
normal laps, well before the subsequent lap cue (Figure 5d, dotted line). Together these results
suggest that as a population, the spatial encoding of granule cells is affected by sensory cues
within a localized area near the cues, rather than persistently altering their estimation of the
animal's position like a landmark.
325
Population vector correlations and Bayesian decoding are average measures of spatial
representation in the granule cell population as a whole, which might obscure differences in how
the activity of individual cells is referenced to local or distant cues. We therefore examined
responses of individual granule cells by measuring cross-correlation offsets between spatial firing
in normal and cue-shifted laps for each cell. In doing so we found two distinct classes of single-
cell shift responses: some cells shifted their firing precisely along with the odor cue shift distance
(0.5m), while others were consistently referenced to the stable lap boundary cue and did not shift
their response location during the odor cue location shift (Figure 5e, f). The majority of cells that
shifted their firing in reference to the new cue location were active immediately in response to the
middle cue ("middle cue-influenced"), while a small number were persistently affected by the cue
shift. In contrast, a relatively constant number of place cells along the track maintained their
normal firing location with reference to the distant lap boundary cue despite the shifted local cue
("non-middle cue-influenced"). Notably, there was also a small enrichment of place cells
immediately prior to the cue but unaffected by the cue shift (Figure 5f), indicating a spatial
response predictive of the normal cue location. The presence of two distinct groups of cells that
fire in reference to the two cues on the treadmill track (the variable middle cue and stable lap cue)
indicates that the DG population maintains independent spatial reference frames based upon
distinct cues within an environment. This code however appears to differ depending upon the
spatial stability of the cues: the effects of spatially variables cues are limited to the cue cell
population while place cells are referenced to global landmarks[43].
346
347
348 **Cue-related Inhibition of DG Responses**
349
Spatial receptive fields of granule cells are hypothesized to form within a competitive network in
which lateral inhibition mediated by GABAergic interneurons enforces sparse encoding that may
aid pattern separation[17,44]. The robust and highly selective cue responses of granule cells in our
behavioral paradigm allowed us to examine patterns of local-circuit inhibition within the DG

network. First, we sought to analyze the effect of cue responses on spontaneous firing observed
within the spatially tuned granule cell population. By comparing spontaneous firing rates at the
middle cue location outside of the primary spatial receptive field in normal laps ("out-of-field") with
laps where the cue is omitted, we identified a significant reduction in background firing of the
spatially tuned granule cell population during the cue presentation (Figure 6a, p=0.0004, Wilcoxon
signed-rank test). This suppression was absent on cue-omitted laps and generally co-varied with
the mean amplitude of cue-related excitation (within-field) on a session-by-session basis (Figure
6b, R=0.22, p<0.0001, Wilcoxon signed-rank test). Furthermore, the timing of the peak of this
inhibition was delayed with respect to the excitatory cue response in these sessions (Figure S6e,
f). These data show that sensory cue input is able to suppress 'noisy' spontaneous firing in the
granule cell population, potentially through lateral inhibition.
365
In addition to suppressing spontaneous firing activity in the DG network, we sought to determine
the effect of cue responses on spatially tuned activity in the place cell population and in cue cells
responding to other differing cues. By examining cells with receptive fields normally at the
intermittent shift location, we found that place cell firing rates were strongly suppressed on shift
laps when a cue was presented at this location compared to normal laps when a cue was not
present here (Figure 6d, e). Furthermore when two distinct cues were normally presented at
different locations, intermittent shifting of the position of one cue to overlap with the other cue
typically had the effect of suppressing the responses in both cue populations (Figure 6f, g),
indicating mutual inhibition among granule cells encoding different cues. Thus in addition to the
inhibition of spontaneous firing in the granule cell population, cue responses lead to overall
inhibition of place cell responses and responses to competing cues as well. Taken together, our
results suggest that cue-related activity in granule cells and the resulting suppression of
responses in neighboring cells via lateral inhibition may organize the dentate gyrus into a
competitive network that utilizes the strongest and perhaps the most informative parameters, such
as stable landmark cues, in order to accurately establish an animal's location in space.
381
382   **Discussion**
383
By imaging calcium activity in large populations of dentate granule cells during head-fixed spatial
behaviors on a treadmill, we have shown that a major population of task-selective neurons is
highly sensitive to specific sensory cues (non-spatial, or 'what' information) rather than to discrete
locations (spatial, or 'where' information). The ability to dynamically manipulate a sensory cue
and its association with locations on the treadmill track allowed us to isolate the population of cue
cells, in addition to the complementary population of place cells recorded in the same sessions.
390
Properties of cells within these groups also differed: sensory cue responses were highly-tuned
and remarkably robust across contexts and over time- significantly more so than the responses
of the canonical place cell population (Figures 1-3). And while cue cells did not have spatially
tuned responses in prior sessions on the same treadmill track without the odor cue (Figure S4),
or in the first several cue presentations, they emerged rapidly during the first session of exposures
to the cue while place cell responses emerged more slowly (Figure 1i). These findings suggest
that dentate cue cells stably represent both sensory cue and other non-spatial features in an

environment (e.g. reward, Figure S2e-h), while place cells constitute a more dynamic population
that gradually adapts to current conditions, for example, by integrating between stable cues in
order to provide an accurate estimate of the animal's location when no cues are present[45].
401
402   **Heterogeneous properties of Granule Cells**
403
The robust and consistent nature of GC cue responses suggests that cue-selective populations
may have been similarly present in previous experiments recording DG activity *in vivo*, however
the lack of precise stimulus control made it impossible to distinguish cue-responsive versus place-
responsive components[5,7,36,46]. Indeed, spatial receptive fields of individual granule cells
measured in different environments with similar sensory cues display mixed levels of changes in
their firing rates, with a subset of granule cells completely remapping their firing fields while others
show more stable dynamics[5,44,46–48]. Based upon our results, these subgroups may correspond to
place cells and cue cells, respectively. A major impact of sensory cues on DG firing largely
independent of spatial context might also explain the relatively lower context selectivity observed
in the DG compared with other hippocampal subfields, at least when measured in different
contexts that contain the same or similar sensory cues[5,44,46–48]. To support this notion, we found
that DG cue cells were largely stable when recorded over long periods of time, with different cue
positions, locomotion speeds, or in different room contexts, while place cells were not (Figures 2-
4). Previous work has demonstrated that the two major long range inputs to the dentate gyrus,
the lateral and medial entorhinal cortex (LEC and MEC), are involved with processing functionally
distinct information[10,11]. This raises the possibility that our "cue cells" may be driven primarily by
sensory inputs from the LEC, while "place cells" are driven by self-motion information relayed
from the MEC, for example in the form of grid cell responses. The segregation of these properties
within the overall granule cell population suggests that these streams of information remain largely
separated at the level of the DG[49], similar to the functional heterogeneity observed in other
hippocampal subfields[50,51]. In these regions such heterogeneity among principal neurons has
been hypothesized to be critical for efficient encoding of spatial representations within a local
population[49–52]. Heterogeneity has also been observed in immediate early gene expression
profiles within DG neurons encoding individual memory engrams[53] which are differentially
targeted by entorhinal afferent pathways[9,53–55]. These findings are in line with our study
demonstrating that sensory and spatial information remain partially separate at the level of the
DG, and suggest that multiple channels of spatial and non-spatial information contribute distinctly
to local and down-stream computations. This separation is also consistent with current ideas that
cue-based and path-integration based navigation are complementary, rather than integrated, in
order to produce place-specific firing both in locations where landmark cues are present as well
as in between cues[56].
435
Motorizing the treadmill, while different from previous studies[5,7,23,44]; does not affect our central
claim of the strong representation of sensory cue information in the DG (Figures 2 and S1).
Additionally, running on motorized treadmills does not influence the differences in the stability and
tuning properties between the cue and place encoding populations (Figures 1 and S1). Yet, we
note that these differences obtained during head-fixed navigation on a relatively cue impoverished
track at a constant speed may be larger than what would be observed during self-guided

exploration of cue-rich, multisensory real-world environments. Nevertheless, our ability to track
the same neurons across different locomotion conditions provides additional evidence for the
existence of a separate and non-overlapping population of cue cells in the DG compared to the
'classical' place cells reported in previous studies[5,7,23,44,47,48]. Indeed, we have found repeatedly
that cue cells displayed striking stability under different imaging conditions (locomotive behaviors,
across days, different rooms), while the spatial fields of place cells remapped in the same set of
conditions, suggesting that place cells integrate position and contextual information to a higher
degree than cue cells. Thus, we reason that the classification of individual cue cells and place
cells in the DG would likely remain consistent in more complex environments such as real-world
or visuospatial virtual environments. Future studies investigating whether cue and place encoding
populations receive different inputs from LEC and MEC, as well as the molecular basis of the
heterogeneity among granule cell populations, will allow for a more detailed inquiry into the
functional diversity in DG granule cells. Such studies may also allow us to selectively target cue
and place cells in order to manipulate their activity and determine their roles during discrimination
behaviors.
457
458 **Functional significance of dentate gyrus cue cell properties**
459
But how do these results inform our ideas about how the DG participates in pattern separation
and spatial map formation? First, we found that although individual granule cells respond to the
same cues when presented in different locations, the strength of this activity was spatially
modulated (Figures 4, S6). Differential encoding of cues based upon their spatial location may
contribute to pattern separation by allowing discrimination between similar cues present in
different places within an environment, or in distinct contexts, over the population. We observed
that spatially modulated cue responses also increased over days when a single cue was
repeatedly presented at multiple locations (Figure 4h), indicating plasticity in location specificity
and potentially progressive enhancement of spatial pattern separation. Notably, responses were
also on average larger for odor cues presented at a single location, as opposed to shifted or
random locations (Figure 4b-d). This suggests that cue responses are more robust for stimuli
paired repeatedly with one particular location, which may indicate dendritic integration of sensory
and spatial information. Conjunctive encoding of sensory cues and their locations within an
environment could play a role in establishing landmarks for spatial navigation. In support of this
idea, we found that most place cells were referenced to the stable lap cue rather than the variable
middle cue, indicating that this cue is preferentially utilized as a landmark (Figure 5).
476
Our study also shows that cue responses lead to potent inhibition of three distinct types of granule
cell activity: spontaneous "noisy" spiking activity, the spatial firing of place cells, and the activity
of cells responsive to other overlapping sensory cues (Figure 6). This supports the hypothesis of
a competitive network in the DG enforced by strong lateral inhibition, which has been suggested
to contribute to pattern separation. In addition, suppression of weaker place cell responses by
cue-related activity may indicate that the network is organized to utilize the strongest and perhaps
the most informative parameters, such as landmark cues, in order to establish an animal's location
in space.
485

Taken together, rate modulation of granule cell cue responses resulting from spatial location or
association with other cues is itself a form of "rate remapping", a feature frequently attributed to
the dentate gyrus in pattern separation[3,17,19,46]. Thus spatial location (Figures 4c,d, 5e,f), as well
as the juxtaposition of multiple cues (Figure 4e,f) and their associated patterns of inhibition (Figure
6), may create a rate modulated landscape of granule cell activity specific to the current context.
The overall pattern of contextually modulated cue responses in the DG may form the basis for
recruitment of distinct populations of neurons, or "global remapping", observed in area CA3[46].
Slower emergence of the place cell population, as well as lower stability over sessions and in
different room contexts, may also point to a role for these neurons in context selective encoding
of position based upon integration between current landmarks (Figures 1-3).
496
While previous work has shown that CA regions also have cue responses[31–35], all of these areas
receive direct (CA3) or indirect input (CA1) from the DG as well as entorhinal cortex, so it remains
unclear whether these properties arise as a result of direct inputs from entorhinal regions, or
through the hippocampal trisynaptic circuit. Thus, the DG is the best place to look specifically at
how the presumed different channels of information from the lateral and medial entorhinal cortex
are initially integrated by the hippocampus during spatial map formation. Our work suggests that
the DG encodes representations of sensory cues in the environment that interact weakly with
spatial representations, potentially through lateral inhibition. The rate-modulated code of DG
sensory cue representations could be refined in subsequent hippocampal areas such as the CA3
recurrent attractor network which may resolve conflicting cue and path-integration based
information to encode an integrated spatial map of things and places, important both for spatial
navigation as well as episodic memory. Future studies examining how these specific features
manifest in subsequent stages of the hippocampal circuit, will shed light on the progressive
processing and integration of sensory cue and spatial information in   hippocampal computation.
511
The properties of cue and place cells in the dentate gyrus have the hallmark of features one might
expect from the first stage of spatial map formation in the hippocampus: mostly distinct but slightly
mixed encoding of sensory cues, or "what", and place, or "where". Such complementary channels
of information may be further refined by downstream hippocampal areas into an integrated spatial
map that encodes relationships between important features such as spatial cues and goals, which
can be used to organize adaptive behavior. Uncovering the functional heterogeneity of cue and
place cells in the dentate gyrus helps us to better understand the basic neural mechanisms
underlying our ability to navigate in complex environments by utilizing both landmarks and self-
motion information to guide our movements through space.
521

Acknowledgements
We would like to thank Jack Berry and other members of the Hen lab for critical insight throughout
this project. This work was funded by NIMH T32 MH015144, S10 OD018464, NIH K99 MH12226
(S.N.T.), NYSTEM-C029157 (G.O., G.F.T., C.O.L, R.H.), Revson Senior Fellowship in Biomedical
Science (A.D.G.), NIMH R21MH122965,NARSAD Young Investigator Award (G.F.T.), NIMH R01
AG043688 (R.H.), MH068542 (R.H.), NIMH R01 MH100631 (A.L.), NINDS R01NS094668 (A.L.),
and NINDS U19NS104590 (A.L.).

Author Contributions

Conceptualization, C.O.L., S.N.T., and R.H.; Methodology, C.O.L., S.N.T., A.S.; Software, C.O.L., S.N.T., A.D.G., J.B.; Formal Analysis, S.N.T., C.O.L.; Investigation, C.O.L., S.N.T.; Resources, C.O.L., A.D.G., G.F.T., J.B., A.L., G.O.; Writing, Review and Editing, S.N.T., C.O.L., R.H., A.D.G., A.L.; Supervision, R.H and A.L.

Declaration of Interests

The authors declare no competing interests.

References

1. O'Keefe, J. & Dostrovsky, J. The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. *Brain Res.* **34**, 171–175 (1971).
2. Amaral, D. G., Scharfman, H. E. & Lavenex, P. The dentate gyrus: fundamental neuroanatomical organization (dentate gyrus for dummies). *Prog. Brain Res.* **163**, 3–22 (2007).
3. Kesner, R. P. An analysis of the dentate gyrus function. *Behavioural Brain Research* **254**, 1–7 (2013).
4. Lee, J. W. & Jung, M. W. Separation or binding? Role of the dentate gyrus in hippocampal mnemonic processing. *Neuroscience & Biobehavioral Reviews* **75**, 183–194 (2017).
5. Hainmueller, T. & Bartos, M. Parallel emergence of stable and dynamic memory engrams in the hippocampus. *Nature* **558**, 292–296 (2018).
6. Lee, J. W., Kim, W. R., Sun, W. & Jung, M. W. Role of dentate gyrus in aligning internal spatial map to external landmark. *Learn. Mem.* **16**, 530–536 (2009).
7. Jung, D. *et al.* Dentate granule and mossy cells exhibit distinct spatiotemporal responses to local change in a one-dimensional landscape of visual-tactile cues. *Sci Rep* **9**, 9545 (2019).
8. Woods, N. I. *et al.* The Dentate Gyrus Classifies Cortical Representations of Learned Stimuli. *Neuron* **107**, 173-184.e6 (2020).
9. Witter, M. P. The perforant path: projections from the entorhinal cortex to the dentate gyrus. *Prog. Brain Res.* **163**, 43–61 (2007).
10. Hargreaves, E. L., Rao, G., Lee, I. & Knierim, J. J. Major dissociation between medial and lateral entorhinal input to dorsal hippocampus. *Science* **308**, 1792–1794 (2005).
11. Knierim, J. J., Neunuebel, J. P. & Deshmukh, S. S. Functional correlates of the lateral and medial entorhinal cortex: objects, path integration and local-global reference frames. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **369**, 20130369 (2014).
12. Wang, C. *et al.* Egocentric coding of external items in the lateral entorhinal cortex. *Science* **362**, 945–949 (2018).
13. Kim, S., Kim, Y., Lee, S.-H. & Ho, W.-K. Dendritic spikes in hippocampal granule cells are necessary for long-term potentiation at the perforant path synapse. *Elife* **7**, (2018).
14. Krueppel, R., Remy, S. & Beck, H. Dendritic integration in hippocampal dentate granule cells. *Neuron* **71**, 512–528 (2011).
15. McNaughton, B. L., Douglas, R. M. & Goddard, G. V. Synaptic enhancement in fascia dentata: Cooperativity among coactive afferents. *Brain Research* **157**, 277–293 (1978).
16. Zhang, X. & Jonas, P. Integration of spatial and non-spatial information by heterogeneous dentate gyrus granule cells. *J Life Sci (Westlake Village)* **2**, 19–24 (2020).
17. de Almeida, L., Idiart, M. & Lisman, J. E. The input-output transformation of the hippocampal granule cells: from grid cells to place fields. *J. Neurosci.* **29**, 7504–7512 (2009).
18. Marr, D. Simple memory: a theory for archicortex. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **262**, 23–81 (1971).

19. Treves, A. & Rolls, E. T. Computational analysis of the role of the hippocampus in memory.
Hippocampus **4**, 374–391 (1994).
20. Hunsaker, M. R., Rosenberg, J. S. & Kesner, R. P. The role of the dentate gyrus, CA3a,b,
and CA3c for detecting spatial and environmental novelty. *Hippocampus* **18**, 1064–1073
(2008).
21. Xavier, G. F. & Costa, V. C. I. Dentate gyrus and spatial behaviour. *Prog.*
Neuropsychopharmacol. Biol. Psychiatry **33**, 762–773 (2009).
22. Knierim, J. J. & Neunuebel, J. P. Tracking the flow of hippocampal computation: Pattern
separation, pattern completion, and attractor dynamics. *Neurobiology of Learning and*
Memory **129**, 38–49 (2016).
23. Danielson, N. B. *et al.* Distinct Contribution of Adult-Born Hippocampal Granule Cells to
Context Encoding. *Neuron* **90**, 101–112 (2016).
24. Pnevmatikakis, E. A. & Giovannucci, A. NoRMCorre: An online algorithm for piecewise rigid
motion correction of calcium imaging data. *J. Neurosci. Methods* **291**, 83–94 (2017).
25. Giovannucci, A. *et al.* CaImAn an open source tool for scalable calcium imaging data
analysis. *Elife* **8**, (2019).
26. Pachitariu, M. *et al.* Suite2p: beyond 10,000 neurons with standard two-photon microscopy.
bioRxiv 061507 (2016) doi:10.1101/061507.
27. Grosmark, A. D. & Buzsáki, G. Diversity in neural firing dynamics supports both rigid and
learned hippocampal sequences. *Science* **351**, 1440–1443 (2016).
28. Sheintuch, L. *et al.* Tracking the Same Neurons across Multiple Days in Ca2+ Imaging Data.
Cell Rep **21**, 1102–1115 (2017).
29. Fuhrmann, F. *et al.* Locomotion, Theta Oscillations, and the Speed-Correlated Firing of
Hippocampal Neurons Are Controlled by a Medial Septal Glutamatergic Circuit. *Neuron* **86**,
1253–1264 (2015).
30. Terrazas, A. *et al.* Self-Motion and the Hippocampal Spatial Metric. *J. Neurosci.* **25**, 8085–
8096 (2005).
31. Bourboulou, R. *et al.* Dynamic control of hippocampal spatial coding resolution by local
visual cues. *eLife* **8**, e44487 (2019).
32. Hetherington, P. A. & Shapiro, M. L. Hippocampal place fields are altered by the removal of
single visual cues in a distance-dependent manner. *Behav. Neurosci.* **111**, 20–34 (1997).
33. Hollup, S. A., Molden, S., Donnett, J. G., Moser, M. B. & Moser, E. I. Accumulation of
hippocampal place fields at the goal location in an annular watermaze task. *J. Neurosci.* **21**,
1635–1644 (2001).
34. O'Keefe, J. & Conway, D. H. Hippocampal place units in the freely moving rat: why they fire
where they fire. *Exp Brain Res* **31**, 573–590 (1978).
35. Wiener, S. I., Paul, C. A. & Eichenbaum, H. Spatial and behavioral correlates of
hippocampal neuronal activity. *J. Neurosci.* **9**, 2737–2763 (1989).
36. O'Keefe, J. & Krupic, J. Do hippocampal pyramidal cells respond to nonspatial stimuli?
Physiol Rev **101**, 1427–1456 (2021).
37. Kohara, K. *et al.* Cell type-specific genetic and optogenetic tools reveal novel hippocampal
CA2 circuits. *Nat Neurosci* **17**, 269–279 (2014).
38. Skaggs, W. E., McNaughton, B. L. & Gothard, K. M. An Information-Theoretic Approach to
Deciphering the Hippocampal Code. in *Advances in Neural Information Processing Systems*
5 (eds. Hanson, S. J., Cowan, J. D. & Giles, C. L.) 1030–1037 (Morgan-Kaufmann, 1993).
39. Sheffield, M. E. J., Adoff, M. D. & Dombeck, D. A. Increased Prevalence of Calcium
Transients across the Dendritic Arbor during Place Field Formation. *Neuron* **96**, 490-504.e5
(2017).
40. Fenton, A. A. & Muller, R. U. Place cell discharge is extremely variable during individual
passes of the rat through the firing field. *Proc Natl Acad Sci U S A* **95**, 3182–3187 (1998).

41. Saleem, A. B., Diamanti, E. M., Fournier, J., Harris, K. D. & Carandini, M. Coherent
encoding of subjective spatial position in visual cortex and hippocampus. *Nature* **562**, 124–
127 (2018).
42. Zhao, X., Wang, Y., Spruston, N. & Magee, J. C. Membrane potential dynamics underlying
context-dependent sensory responses in the hippocampus. *Nature Neuroscience* **23**, 881–
891 (2020).
43. Wang, C., Chen, X. & Knierim, J. J. Egocentric and allocentric representations of space in
the rodent brain. *Curr. Opin. Neurobiol.* **60**, 12–20 (2019).
44. Danielson, N. B. *et al.* In Vivo Imaging of Dentate Gyrus Mossy Cells in Behaving Mice.
Neuron **93**, 552-559.e4 (2017).
45. Draht, F. *et al.* Experience-Dependency of Reliance on Local Visual and Idiothetic Cues for
Spatial Representations Created in the Absence of Distal Information. *Front Behav Neurosci*
11, 92 (2017).
46. Leutgeb, J. K., Leutgeb, S., Moser, M.-B. & Moser, E. I. Pattern separation in the dentate
gyrus and CA3 of the hippocampus. *Science* **315**, 961–966 (2007).
47. GoodSmith, D. *et al.* Spatial Representations of Granule Cells and Mossy Cells of the
Dentate Gyrus. *Neuron* **93**, 677-690.e5 (2017).
48. Senzai, Y. & Buzsáki, G. Physiological Properties and Behavioral Correlates of
Hippocampal Granule Cells and Mossy Cells. *Neuron* **93**, 691-704.e5 (2017).
49. Fernández-Ruiz, A. *et al.* Gamma rhythm communication between entorhinal cortex and
dentate gyrus neuronal assemblies. *Science* **372**, eabf3119 (2021).
50. Cembrowski, M. S. & Spruston, N. Heterogeneity within classical cell types is the rule:
lessons from hippocampal pyramidal neurons. *Nat. Rev. Neurosci.* **20**, 193–204 (2019).
51. Soltesz, I. & Losonczy, A. CA1 pyramidal cell diversity enabling parallel information
processing in the hippocampus. *Nat. Neurosci.* **21**, 484–493 (2018).
52. Chelaru, M. I. & Dragoi, V. Efficient coding in heterogeneous neuronal populations. *Proc.*
Natl. Acad. Sci. U.S.A. **105**, 16344–16349 (2008).
53. Sun, X. *et al.* Functionally Distinct Neuronal Ensembles within the Memory Engram. *Cell*
181, 410-423.e17 (2020).
54. Luna, V. M. *et al.* Adult-born hippocampal neurons bidirectionally modulate entorhinal inputs
into the dentate gyrus. *Science* **364**, 578–583 (2019).
55. Erwin, S. R. *et al.* A Sparse, Spatially Biased Subtype of Mature Granule Cell Dominates
Recruitment in Hippocampal-Associated Behaviors. *Cell Rep* **31**, 107551 (2020).
56. Poucet, B. *et al.* Independence of landmark and self-motion-guided navigation: a different
role for grid cells. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **369**, 20130370 (2014).
57. Grosmark, A. D., Sparks, F. T., Davis, M. J. & Losonczy, A. *Offline Memory Reactivation*
Promotes the Consolidation Of Spatially Unbiased Long-Term Cognitive Maps.
2020.08.20.259879 https://www.biorxiv.org/content/10.1101/2020.08.20.259879v1 (2020)
doi:10.1101/2020.08.20.259879.

667
668
Experimental Procedures

Mice
All procedures were conducted in accordance with the U.S. NIH Guide for the Care and Use of
Laboratory Animals and the Institutional Animal Care and Use Committees of New York State
Psychiatric Institute and Columbia University. Adult male C57BL/6J mice were supplied by
Jackson Laboratory and Dock10[Cre] mice were a gift from Susumu Tonegawa (Massachusetts
Institute of Technology). Mice were housed in a vivarium grouped 2-4 mice/cage enriched with
running wheels, maintained on a 12-hour light cycle and used at 8-10 weeks of age. Experiments

were conducted during the light portion of the cycle. Food and water were available *ad libitum*
until the beginning of the experiment, when they were placed under controlled water supply and
maintained at >90% of their pre-deprivation weight over the course of imaging experiments. In
total, imaging data from 18 mice were used in this study.
682
Surgery
Dentate gyrus virus injection and imaging window implantation surgeries were performed as
described previously (Danielson 2016, 2017). For all surgical procedures, mice were anesthetized
with 1.5% isoflurane at an oxygen flow rate of 1 L/min, and head-fixed in a stereotactic frame
(Kopf Instruments, Tujunga, CA). Eyes were lubricated with an ophthalmic ointment, and body
temperature maintained at 37°C with a warm water recirculator (Stryker, Kalamazoo, MI). The fur
was shaved and incision site sterilized prior to beginning surgical procedures, and subcutaneous
saline and carpofen were provided peri-operatively and for 3 days post-operatively to prevent
dehydration and for analgesia. Mice were unilaterally injected with recombinant adeno-associated
virus (rAVV) carrying the GCaMP6s transgene (pAAV.Syn.GCaMP6s.WPRE.SV40) purchased
from Addgene (viral prep #100843-AAV1) with titer of 1-5x10^{12} in dorsal dentate gyrus using a
Nanoject syringe (Drummond Scientific, Broomall, PA). Injection coordinates were -1.5 mm AP,
-2 mm ML, and -1.85, -1.7, -1.55 mm DV relative to the cortical surface. 30 nL of diluted virus was
injected at each DV location in 10 nL increments. Mice were allowed to recover for 3 days and
then were unilaterally implanted with an imaging window and stainless steel head-post for head
fixation. Imaging windows were constructed by adhering 2 mm diameter, 2.3 mm long stainless
steel hypodermic tubing (Ziggy's Tubes and Wires Inc, Pleasant Hill, TN) to 2 mm diameter glass
coverslips (Potomac Photonics, Halethorpe, MD). A 2 mm diameter craniotomy was made
centered on the previous injection site with a taper pointed-drill (Henry Schein Inc, 9004367) and
dura was removed with micro curette (FST, 10080-05). The overlying cortex was gently aspirated
to reveal capsular fibers with continuous irrigation with ice cold aCSF solution and bleeding was
controlled with a collagen gel sponge (Avitene). Under minimal bleeding, a 30g blunt syringe was
used to gently aspirate capsular and CA1 alveus fibers with white appearance and CA1
pyramidale and moleculare with pink appearance until vasculature of the hippocampal fissure
became visible (under bright light with low bleeding). The cannula, attached to the stereotactic
handle, was then gently lowered into the craniotomy and affixed to the skull using dental cement
(Unifast Trad powder and LC light cured acrylic UV, Henry Schein).
710
Behavioral training and apparatus
After a minimum of 1 week recovery period, mice underwent a water restriction scheme (1ml per
day) and trained to run on treadmill while head-restrained. The training period typically lasted a
week (2 training sessions/day, 15 min each) until the mice were able to run for at least 1 lap/
minute and seek reward from 3 randomly placed reward zones by licking the water delivery port.
We then initiated the motorized belt adjusted to the natural velocity of each mouse and proceeded
training for 1-2 more days. We discarded mice that did not perform sufficiently to receive getting
all of their daily water supply during treadmill training, and were not motivated to move on the
treadmill. A separate cohort of mice ran at will on the treadmill (Figures S1 and 2). During the
training period mice ran on a cue-less 'burlap' belt and progressed to a different belt containing
cues of different modalities as described below. For room switch experiments (Fig. 3f-h), mice

were transferred to a separate room with similar imaging equipment on another floor of the
building in their home cage. Mice were imaged for two consecutive sessions/day, each 15 minutes
long, for the duration of 7-10 days. Single odor, light, and whisker texture responses described in
Figures 1 and 3 and S2-5 are from the first time mice were exposed to these sensory stimuli. The
subsequent exposures to these sensory cues, they were either in different locations on the belt,
preceded or followed by additional cues (Figures 4, 6), reducing the effects of learning the location
of these cues. Additionally, in all experiments, the treadmill belt material was changed between
sessions to reduce the chances of urine contamination which might act as an additional olfactory
cue.
731
The behavioral apparatus consisted of 2m long, 3" wide fabric belt stretched between 6" diam.
laser-cut plastic wheels, mounted on an aluminum frame (8020.net). Spatial triggering of task
events was performed by custom software via serial communication with a microcontroller
(Arduino DUE) and an associated printed circuit board (OpenMaze OM4 PCB,
www.openmaze.org) on the treadmill. The axle of the treadmill wheel was attached to a
quadrature rotary encoder (US Digital part #: MA3-A10-125-B) connected to a custom quadrature
decoder board and Arduino Nano (courtesy of Wen Li). Angular displacement was converted into
a virtual linear distance based on the circumference of the treadmill. The errors were corrected
via a registration anchor marked by radio-frequency identification (RFID) buttons (SparkFun
Electronics) at the lap boundary of the belt and was detected when it passed over an RFID reader
(ID-12LA, SparkFun) affixed underneath the mouse. A 12V DC gear motor was attached to the
axle of the treadmill connected to a separate Arduino/OpenMaze shield using pulse-width
modulation to adjust the rotation speed. A water reservoir connected to a water delivery port
consisting of a small gavage needle (Cadence Science) was placed within reach of the mouse's
tongue. A capacitance touch sensor (Sparkfun MPR121) was attached to the water port to
measure licking and the sensor was connected to the Arduino/OM4 PCB. Small 2-3ml drops of
water were delivered by the brief opening a solenoid valve (Parker Hannefin) connected to the
water port. Rewards were triggered at random locations each lap when mice entered a 10cm long
reward zone on the track and were available until mice exited the reward zone or 3 sec had
elapsed. Olfactory stimuli consisted of undiluted isoamyl acetate (IAA, Sigma W205532) which
was added to syringe filters (Whatman #6888-2527) and delivered by opening a solenoid valve
(SMC) connected to a flow controller delivering constant airflow of compressed medical grade air
for 1s (~3psi). Visual and tactile stimulation consisted of a red LED contralateral to the imaged
region, or a 1" square piece of sand paper brushed by the contralateral whiskers using a stepper
motor, at approximately the speed of the treadmill belt. Custom written Be-Mate algorithm
implemented in Java was used for recording mice's licking, its position on the belt, and cue
delivery. Mice were monitored using an IR camera (PS3eye) and illuminated using an IR LED
array.
760
To isolate cue-selective responses among the granule cell population, normal cue laps in which
the olfactory, visual, or tactile cue was presented in the middle of the treadmill track (90-110cm)
were interspersed with occasional laps (10-20% of laps) in which the same cue was omitted
("omit" laps), or shifted forward ¼ of the track ("shift" laps). For a subset of sessions, the olfactory

cue was presented at one of 5 locations along the track randomly each lap, in order to examine
the effect of spatial pairing of the cue.
767
In vivo two-photon imaging
Imaging was conducted using a microscope setup which consists of 8kHz resonant galvanometer
(*Bruker*) mounted to a mirror-based multi-photon microscopy system (*Prairie Technologies*) and
an ultra-fast pulsed laser beam (920-nm wavelength; *Chameleon Ultra II, Coherent*, 20–40-
mW average power at the back focal plane of the objective) controlled with an electro-optical
modulator (*Conoptics*, Model 302 RM). GCaMP fluorescence was excited through a 40x water
immersion objective (Nikon NIR Apo, 0.8 NA, 3.5 mm WD) and fluorescence signals detected
with photomultiplier tubes (*Hamamatsu 7422P-40*), acquired with PrairieView software (*Prairie*)
at 30fps frame rate (512X512 pixels, 1.3 μm/pixel). A custom dual stage preamp (1.4X10^5 dB,
Bruker) was used to amplify signals prior to digitization. Two goniometers (Edmund Optics) were
used to adjust the angle of each mouse's head in order to achieve the same imaging plane over
multiple sessions.
780
Data processing for Ca^{2+} imaging. Movies were motion corrected using NoRMCorre algorithm
using a non-rigid registration method that splits the field of view (FOV) into overlapping patched
that are registered separately then merged by smooth interpolation[24]. Videos were then spatially
and temporally down-sampled by 2 to reduce noise and the computational power required for cell
segmentation. Spatial and temporal components for individual cells were extracted using large-
scale sparse non-negative matrix factorization (CNMF/CaImAn)[25] or using the singular value
decomposition method by Suite2p algorithm (https://github.com/cortex-lab/Suite2P), both of
which resulted in similar regions of interest (ROIs). We used Suite2p graphical user interface to
manually select small, densely packed DG granule cells and discard large isolated cell bodies
corresponding to mossy cells or other hilar interneurons. To obtain total number of DG granule
cells within the imaging fields of view in a subset of total sessions, time averaged images were
segmented using the Cellpose algorithm (https://github.com/MouseLand/cellpose) followed by
manual inspection. Ca^{2+} transient events were defined by a custom detection algorithm which
identifies fluorescence peaks with a rise slope greater than 4 standard deviations above an
iteratively refined baseline.
796
797 *Behavioral and Calcium Data Alignment.* Behavioral data was aligned to Ca^{2+} data using the
798 record of a synchronization signal between the two computers used for data collection. Behavioral
799 data was down-sampled to match Ca^{2+} imaging data.
800
801 **Data Analysis**
802 Data were analyzed using custom-written routines implemented in MATLAB. Plots were
803 generated in MATLAB and Prism.
804
Identification of spatially-tuned neurons. We restricted our analysis to continuous running at least
2 sec in duration and with a minimum peak speed of 5 cm/sec. For each lap crossing, position
data and Ca^{2+} transient events for each cell were binned into 2 cm-wide windows (100 bins),
generating raw vectors for occupancy-by-position and calcium transient numbers-by-position

which were then circularly smoothed with a Gaussian kernel (*SD* = 5 cm). A firing rate-by-position
vector was computed by dividing the smoothed transient number vector by the smoothed
occupancy vector. Within each lap, we circularly shuffled the positions 1000 times and
recomputed firing rate-by-position vectors to generate a null distribution for each spatial bin. A
spatially selective cell was defined that met the following criteria: (a) the cell should fire above its
mean firing rate within its spatial field in at least 20% of laps or for a minimum of 3 laps; and (b)
observed firing should be above 99% of the shuffled distribution for at least 5 consecutive spatial
bins (10 cm) wrapping around the two edges of the belt. We have identified spatially tuned
neurons by excluding bins in which sensory cues were omitted or shifted and calculated firing rate
vectors in these laps separately. Among all of the spatially tuned neurons, "middle cue cells" were
defined as those with averaged spatial fields that overlapped with at least 50% of the 45th and 55th
bins and had peak amplitude at least two times larger than those in cue-omitted laps. "Lap-cue
cells" were defined as those with averaged spatial fields overlapping at least 50% of the region
wrapping around the 90th and 10th bins in the normal laps and have peak amplitude in cue-omitted
laps and cue shifted laps not exceeding than at least two times of that in normal laps. The
remaining cells constituted the "place cells".

Spatial information, stability, consistency, and emergence of spatial fields. To calculate a measure
for spatial information content for granule cells in Fig. 1g, we adapted a traditional method of
spatial information assessment[23,38,57] to Ca^{2+} imaging data. For each cell, we used the firing rate-
by-position vector and shuffled null distribution computed above and calculated the spatial
information content for each as described previously[38,57]. To account for the fact that low firing
rates artificially produce high spatial information scores, we subtracted the mean of the shuffled
information per spike from observed information per spike, divided by the standard deviation of
the shuffled values to determine the spatial variance for each cell. Therefore, the amount of spatial
information is inferred from differences in normalized Ca^{2+} activity in each neuron and reported
as bits per seconds. The consistency of place field firing in Fig. 1h was determined as the cross-
correlation between the averaged firing rate-by-position vector of the first and the second halves
of the total number of cue normal laps within a session. We determined place field onset lap in
cue normal laps (Fig. 1i) as described previously[39]. Briefly, starting on lap 1 we searched for a
significant Ca^{2+} transient event present within the boundaries of the previously determined mean
spatial field calculated from all the laps in the session. If one were found we would then search
for Ca^{2+} transient event on each of the next 4 laps. If 3 of the 5 laps had Ca^{2+} transients within the
mean place field boundaries, lap 1 would be considered the place field onset lap. If either lap 1
had no Ca^{2+} transient or less than 3 of the 5 laps had Ca^{2+} transient, we would move to lap 2 and
repeat the search (Fig. 1i). The field onset laps, and the mean firing rates during first 5, 6-10 and
11-15 laps was then assessed for each group of cells.

Multi-Session Cell Tracking. Cells were tracked across sessions using CellReg[28]. Briefly, rigid
alignment with both translations and rotations was performed on spatial footprint projections of
each session and manually inspected for quality. To improve performance with our data, we
modified the CellReg source code to consider complete spatial footprints instead of centroids
during alignment. The centroid distance between neighbors was then calculated and used to
create a probabilistic model that estimated the expected error rate at different thresholds. The

optimal centroid distance threshold was chosen by the algorithm and used to match cells. A
clustering algorithm then refined these decisions previously made using pairwise comparisons.
855
Following cell registration, tracked cells were matched with their corresponding functional cell
types (i.e. mid-, lap-cue, place cells, as described above). All analyses presented in Figure 3 are
carried out in pairwise, to maximize the number of cells in each comparison and to minimize the
total number of comparisons. For multiday comparisons we used Day1, session 1 as the normal
session, and for multisession comparisons we used Visual stimulus session as the normal
session. To calculate the fraction of cells that maintain their identity, cell pairs that were counted
as being the same cell type in both sessions was divided by all of that cell type in the normal
session. In order to derive a null distribution for preservation of pairwise identity, we randomly
permutated the cell IDs of all the tracked cells in pairwise sessions 1000 times and calculated the
fraction of cells that were the same, among all of that cell type in the normal session. We
calculated p-values by comparing actual data to this null distribution, 97.5th% of the null
distribution is presented dotted lines in Figure 3.
868
Rate correlation and population vector (PV) analysis. Comparison of the activity between different
sessions was calculated using Pearson's correlation of the spatially binned, averaged firing rate-
by-position vector in cue normal laps in Figures 2,3, S4. The variability in neural activity between
self-driven and motorized treadmill sessions (Figure 2) and between lap types (in Figure 6) was
calculated by using Pearson's correlation on each 2 cm bins of the firing rate-by-position vector
along the treadmill during odor cue trials (mean for all spatially tuned cells).
875
Spatial modulation
To compare single cell cue responses at different locations, cue-triggered average z-scored
calcium transients were measured for cue cells, defined with reference to omit lap activity as done
previously, for middle cue sessions. For dual cue location experiments, cue cells were identified
as units with receptive field peaks in the cue location whose cue-zone calcium event rate was
greater than 95% of shuffled responses versus the cue-omitted laps. For statistics of spatial
modulation over days, cue cells that significantly preferred one cue location were counted when
responses at one location exceeded 95% of the shuffled cue position event rates. Spatial
modulation in this paradigm is listed as a ratio of mean spatial firing rates for the non-preferred
location over the preferred location rates for each cell (i.e. lower values correspond to higher
spatial selectivity of cue responses).
887
Bayesian Reconstruction Analysis. To calculate the probability of the animal's position given a
short time window of neural activity, we used a previously published method based on Bayesian
reconstruction algorithm[27]. Briefly, Ca^{2+} transient events for each cell were binned into 1 second
windows to construct firing rate vectors. For each of these binned firing rate vectors, Bayesian
classification of virtual position (posterior probability for each bin) was performed by a previously
described method[27] utilizing a template comprising of a cell's smoothed firing rate-by-position
vectors. In order to cross-validate our decoding procedure, we divided firing rate-by-position
template into lap crossings, used 1/5th of laps as "testing" dataset while the remaining 4/5th of laps
constituted the "training" dataset. For example, lap 1 was tested based on the firing rate-by-

position vectors calculated using laps 2,3,4,5, lap as template, and lap 6 was tested based on the
firing rate-by-position vectors calculated from laps 7,8,9,10, and so on. The resulting posterior
probability distribution for each bin is the likelihood for an animal is located in that bin, which adds
up to 1, and the bin with the maximum posterior probability is the estimated position of the animal.
To determine the decoding error we calculated the absolute difference between the animal's
actual position and the maximum posterior probability in that bin. Post-reconstruction, we divided
the time bins (excluding those with no activity) according to the lap types.
904
Single cell cue shift analysis
Spatial firing rates for each spatially tuned cell on normal middle cue laps were cross-correlated
with firing rates on shift laps in order to estimate the cross-correlation peak offset (i.e. shift
magnitude) for each cell after cue manipulation. Binned histograms of numbers of cells with
spatial receptive fields at particular locations along the track were plotted with respect to their shift
magnitudes. Numbers of cells were averaged over populations showing no shift (-5 to 5 cm shift)
or that shifted their firing along with the cue (50cm±5cm).
912
Inhibition analysis
Out-of-field firing was calculated for cells found to be significantly spatially tuned on normal middle
cue laps by extracting calcium event rates in the ~200cm track length excluding the peak place
field (+/- 10cm). Average out-of-field firing rates were then calculated across all cells for cue laps
and intermittent laps where the cue was omitted. For comparison of cue-associated excitation
and inhibition levels, average firing rates were computed by session for the 20cm region
surrounding the middle cue, with respect to the normal pre-cue baseline firing rate. Cue-related
inhibition of place cell firing was calculated by selecting spatially tuned cells whose firing field on
normal middle cue laps fell within the region of the cue on shift laps (50-80cm). Firing rates for
these cells were then averaged for normal laps where the cue was not presented in this region
and compared with laps where the cue was shifted to this region (50cm). Mutual inhibition
between cues was calculated by first selecting cells responsive to each of 2 cues of different
modalities (olfactory or visual) presented at 40cm and 120cm. Responses were then averaged
for each cue cell for laps in which the cues were presented alone at these locations versus
intermittent laps where the cues were presented together at one of the two former locations.
928
929 **Data and Software Availability**
930 Data and custom programs are available upon reasonable request.

Figures

Figure1: Robust representation of sensory cues in the dentate gyrus

a) Two-photon imaging of DG population calcium activity *in vivo*. Left, diagram of the imaging window implant in the dorsal dentate gyrus. Right, Time averaged two-photon image of GCaMP6s-expressing granule cells (Scale bar = 50μm)

b) Diagram of head-fixed treadmill apparatus for sensory cue delivery in the spatial cue task.

c) Fraction of spatially tuned cells at each treadmill position (bins=10cm, n = 4091 cells from 8 mice across 6-9 sessions/mouse), Locations of the odor and lap cue are shown blue and green shaded areas, respectively.

d) DG neuron activity during the spatial cue task. Top: Location of the lap cue (green boxes) and an odor cue (blue box), on normal, and odor cue shift or omit laps. Bottom: Lap-averaged spatial firing rates of 285 spatially tuned neurons (n=8 mice) during the first session with the odor cue on normal middle location (left), cue-shifted (middle) and cue-omitted (right) laps. Each row represents activity of a single cell in the different lap types, sorted by activity on normal middle cue laps.

e) Average spatial firing rates by position for neurons shown in panel "d." on normal (black), cue-shifted (red) and cue-omitted laps (blue, mean ± SEM). (Inset) averaged area under the firing rate curves (AUC, Hz*cm) within the middle cue region during normal (black bar), cue shifted (red bar), or cue omitted laps (blue bar). $P_{normal-omit}$ <0.0001, $P_{normal-shift}$ =0.02, Wilcoxon signed rank test; error bars are mean ± SEM.

f) Example fluorescence traces from representative odor-cue, lap-cue and place cells within a single session. Black, red, and blue traces represent normal, cue shifted and omitted laps, respectively. Scale bars are cm and standard deviation from each cell's baseline fluorescence.

g) Spatial information within cue and place cell populations. Z scored spatial information, χ^2=48.47, p<0.0001, $P_{OdorCue-LapCue}$=0.0387, $P_{OdorCue-Place}$=0.0028, $P_{LapCue-Place}$<0.0001, Kruskal Wallis and Dunn's multiple comparisons test.

h) Tuning consistency of cue and place cells. Firing rate correlation between first and last halves of the session, χ^2=32.60, p<0.0001, $P_{OdorCue-LapCue}$=0.2224, $P_{OdorCue-Place}$=0.0117, $P_{LapCue-Place}$<0.0001, Kruskal Wallis and Dunn's multiple comparisons test. Red dotted lines in violin plots show median, black dotted lines show quartiles.

i) Emergence of cue and place responses. Cumulative distribution of spatial field onset lap for cue and place cells, χ^2=9.29, p=0.0096, $P_{OdorCue-LapCue}$=0.8925, $P_{OdorCue-Place}$=0.0194, $P_{LapCue-Place}$=0.0091, Kruskal Wallis and Dunn's multiple comparisons test. (Inset) Spatial field firing rates of cue and place cells during -1-5th, 6-10th and 11-15th laps, main effect of cell type: $F_{2,1473}$=6.73, p=0.0012; cell type × lap number interaction $F_{4,1473}$=2.44, p=0.04; main effect of lap number: $F_{2,1473}$=0.28, p=0.7; Lap 1-5th, $P_{OdorCue-LapCue}$=0.99, $P_{OdorCue-Place}$=0.0011, $P_{LapCue-Place}$<0.0001, 2-way ANOVA and Tukey's multiple comparisons test. $N_{OdorCue}$=114, N_{LapCue}=220, $N_{PlaceCell}$=160, from 8 mice 2 sessions each. *P < 0.05; **P < 0.01; ***P < 0.001.

Figure 2: Stable encoding of sensory cues regardless of animal's locomotive behavior or velocity

a) Activity of DG granule cells in response to the same odor-cued track during self-driven (left) and motorized (right) treadmill locomotion. Top: diagram of treadmill track with lap cue and middle odor cue. Bottom: Spatial firing rates for neurons tracked through consecutive sessions of self-driven and motorized treadmills (n=67 cells, 5 mice), cells are sorted by their activity in the self-driven treadmill session. See also Figure S1a-f.

b) Comparison of average spatial rate correlations in individual neurons tracked across self-driven and motored treadmills (χ^2=10, p<0.01, $P_{Odor-Lap\ cue}$>0.99, $P_{Odor\ Cue-Place}$ =0.0059, $P_{Lap\ Cue-Place}$ =0.0421, $N_{Odor-Cue}$=30, $N_{Lap-Cue}$=22, N_{Place}=15). Kruskal-Wallis test, Dunn's multiple comparisons test. Boxes, 25th to 75th percentiles; bars, median; whiskers, 99% range. *$P < 0.05$; **$P < 0.01$.

c) Spatial population vector correlations of spatially tuned neurons between self-driven and motorized treadmills n=31±9 cells in 5 mice (red), shaded error bars represent mean ± SEM for each treadmill position. Locations of the middle and lap cue are shown by blue and green shaded areas, respectively. Mean population vector correlations by mice exceeded chance levels as estimated by shuffling cell identity in each mouse/session (gray).

d) DG neuron activity during the spatial cue task when the treadmill is motorized at different velocities in each lap, or the cue omitted, in a pseudorandom manner. Top: velocity1: 6cm/s, velocity2: 9cm/sec, velocity3: 12cm/sec, asterisks represent reference location for position-triggered averages in following panels; Bottom: Spatial firing rates of 89 spatially tuned neurons (n=3 mice) during different velocity and cue-omitted laps. Each row across all graphs represents a single cell, with the population sorted by each cell's activity on velocity1 laps. See also Figure S1g-h.

e) Example reference *position-triggered* average Ca²⁺ transients from six example cue cells on faster (red), normal (green), and slower (blue) velocity laps. Axes are seconds and standard deviation from each cell's baseline fluorescence for the session.

f) Example *cue-triggered* average Ca^{2+} transients from the same cue cells shown in panel "e." on faster (red), normal (green), and slower (blue) velocity laps.

Figure 3: Cue cell responses are modality-specific, and more stable across days and room contexts than place cells.

a) Spatial firing rates of individual DG granule cells matched between sessions within a day and after one week. Top: diagram of odor cue presentation during all sessions. Bottom: Spatial firing rates for spatially tuned neurons tracked during subsequent sessions on the same day or one week later, ordered according to the receptive field location of each cell during the first exposure (n=233 cells, 8 mice).

b) Activity of granule cells in response to different sensory cues. Top: diagram of sessions with cues of different sensory modalities in the middle position but a similar lap cue. Bottom: Spatial

rates for neurons tracked through consecutive sessions during exposures to different cues. Data are shown for neurons with significant tuning in the visual cue session and tracked in olfactory and whisker tactile cue sessions (n=196 cells, 6 mice). See also Figure S2a-d

c) Fraction of cross-session registered cells that encoded the same behavioral variable, lap cue (green), middle cue (blue), or place (orange), within same day (left), across days (1wk, middle), and during exposures to different sensory modalities (right) . Cells encoding the same variable in different sessions on the same day, χ^2=1.81, p=0.4039, $P_{LapCue-OdorCue}$=0.4221, $P_{LapCue-Place}$=0.549, $P_{OdorCue-Place}$=0.9713; cells encoding the same variable across days, χ^2=14.36, p<0.001, $P_{LapCue-OdorCue}$=0.1566, $P_{LapCue-Place}$<0.0001, $P_{OdorCue-Place}$=0.0313 (8 mice, 8 matched sessions, Day1-Day1', Day1-Day7); cells encoding the same variable across different sensory modalities, χ^2=19.78, p<0.00001, $P_{LapCue-OdorCue}$<0.00001, $P_{LapCue-Place}$=0.0766, $P_{OdorCue-Place}$=0.0464 (6 mice, 12 matched sessions, Odor-Vis. and Vis.-Tact.); Kruskal-Wallis test, Dunn's multiple comparisons test. Dashed lines represent 97.5th% of null distributions for each cell type. Error bars, mean ± SEM.

d) Fractions of spatially tuned cells that stably encoded only one variable (cue or place, pink) or switched response types (cue to place or vice versa, black) within same day, across days and in response to different sensory cues: $P_{Switch-Stable\ (Same\ Day)}$<0.001, $P_{Switch-Stable\ (Across\ Days)}$<0.001, $P_{Switch-Stable\ (Sensory\ Modalities)}$<0.001, Wilcoxon Rank Sum test. Dashed lines: 2.5th and 97.5th% of null distributions. Error bars, mean ± SEM.

e) Mean spatial firing rate correlations within the same day (red), different days (blue) and with different sensory cues (green) for mid-cue cells (left, χ^2=27.88, p<0.0001, $P_{Same\ Day-Across\ Days}$=0.2465, $P_{Same\ Day-Sensory\ Modalities}$<0.0001, $P_{Across\ Days-Sensory\ Modalities}$=0.0038, $N_{Same\ Day}$=37, $N_{Across\ Days}$=37, $N_{Sensory\ Modalities}$=55); place cells (middle, χ^2=18.79, p<0.0001, $P_{Same\ Day-Across\ Days}$<0.0001, $P_{Same\ Day-Sensory\ Modalities}$=0.7449, $P_{Across\ Days-Sensory\ Modalities}$=0.0055, $N_{Same\ Day}$=76, $N_{Across\ Days}$=76, $N_{Sensory\ Modalities}$=83), lap-cue cells (right, χ^2=5.096, p=0.0782, $P_{Same\ Day-Across\ Days}$=0.0723, $P_{Same\ Day-Sensory\ Modalities}$>0.9999, $P_{Across\ Days-Sensory\ Modalities}$=0.7932, $N_{Same\ Day}$=80, $N_{Across\ Days}$=80, $N_{Sensory\ Modalities}$=59). Kruskal-Wallis test, Dunn's multiple comparisons test. Boxes, 25th to 75th percentiles; bars, median; whiskers, 99% range. *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$.

f) Activity of DG granule cells in response to the same odor-cued track when recorded in different rooms. Top: diagram of treadmill track with an odor in the middle position and an invariant lap cue, performed in different rooms. Bottom: Spatial firing rates for tuned neurons tracked through consecutive sessions in different rooms (n=83 cells, 4 mice). See also Figure S5.

g) Fraction of individual cells that encoded the same variable, odor cue (blue), lap cue (green) or place (orange) in different recording rooms. χ^2=6.59, p=0.0372, $P_{OdorCue-Place}$=0.0490, $P_{LapCue-OdorCue}$=0.9557, $P_{LapCue-Place}$=0.0972. Dashed lines: 2.5th and 97.5th% of null distributions. Error bars, mean ± SEM.

h) Comparison of spatial firing rate correlations in all tuned cells across rooms (χ^2=15.85, p<0.001, $P_{Odor-Lap\ cue}$=0.27, $P_{Odor\ Cue-Place}$ <0.001, $P_{Lap\ Cue-Place}$ =0.03, $N_{Odor-Cue}$=25, $N_{Lap-Cue}$=33, N_{Place}=26).   Kruskal-Wallis test, Dunn's multiple comparisons test. Boxes, 25th to 75th percentiles; bars, median; whiskers, 99% range. *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$.

i) Fractions of spatially tuned cells that stably encode only one variable (cue or place, pink) or switched response types (cue to place or vice versa, black) across different rooms (p<0.01,

Wilcoxon Rank Sum test.) Dashed lines: 2.5th and 97.5th% of null distributions. Error bars, mean ± SEM.

Figure 4: Sensory cue responses are spatially modulated

a) Experimental setup to examine effects of cue-location pairing on cue responses. Odor cue response amplitudes are measured for the more frequent middle cue location (reference, black), an intermittently shifted cue location in the same sessions (shifted cue, red), or with random administration of the same cue at one of three positions on each lap in a subsequent session (random cue, cyan).

b) Example cue-triggered average Ca^{2+} transients from two example cue cells on normal (black), cue-shifted (red) and random-cue (cyan) laps/sessions.

c) Peak amplitudes of Ca^{2+} transients from individual odor cue cells in normal, shift and random cue presentation conditions; χ^2=49.36, p<0.001, $P_{Normal-Shift}$<0.0001, $P_{Normal-Random}$<0.0001, $P_{Shift-Random}$>0.9999, Friedman test and Dunn's multiple comparisons tests (n=101 cells, 5 mice). Red lines in violin plots show median, black lines show mean.

d) Difference in the magnitude of cue responses in normal laps from random-cue and cue-shift laps. Yellow line, linear regression (R^2= 0.546, p=3.5x10^{-9}). Pie chart shows the percentage of neurons with higher average responses in normal (black, 64%), cue-shift (red, 12%), random cue presentation conditions (cyan, 17%), and both shifted and random laps (purple, 7%).

e) Dual cue location task: (Top) An LED visual cue is given consistently at two positions on the track. Arrangement of cues for normal laps (left, 80% of laps), laps where first cue is omitted

(middle, 10%), and laps where the second cue is omitted (right, 10%). (Bottom) Spatial firing rates of 433 significantly tuned cells on first day of task (N=3 mice).

f) Average spatial firing rates for significant cue cells, for cells preferring cue position 1 (top) or cue position 2 (bottom) on first day of task (blue, n=25 pos1, n=25 pos2) and 4th day of task (pink, n=13 pos1, n=20 pos2 cells).

g) Spatial modulation index (rate non-preferred cue/preferred cue) for all cue cells on days 1-4. (p=0.0125, day 1 vs. days 2, 3, or 4, n= 50/40/50/33, Wilcoxon rank sum test). Note that lower values mean higher cue spatial selectivity.

h) Empirical cumulative probability distribution of spatial modulation index for all cue cells on days 1-4. See also Figure S6a-d.

Figure 5: Effects of cue manipulation on spatial encoding

a) Population vector correlations for all spatially tuned cells during normal middle cue laps and shifted cue laps. Pair-wise correlations of normal middle cue laps are shown in black, cue shifted laps are shown in red and between normal and shifted laps are shown in green. Large differences in pair-wise correlations are largely confined to locations of the middle/shift and lap cue that are shown blue and green shaded areas, respectively.

b) Bayesian decoding: example prediction of 4 normal and 1 cue shifted laps for a representative session, based upon activity from all spatially tuned granule cells. The magenta line shows predicted position at each time bin, while the green line shows the observed position of the animal.

c) Average prediction error for normal and cue shifted laps calculated as the absolute median distance between the decoded value in each time bin and the actual value of the position (p<0.0001, Wilcoxon signed-rank test, n=66 sessions). Boxes, 25th to 75th percentiles; bars, median; whiskers, 99% range.

d) Spatial decoding error for each treadmill position in normal (black) and cue shifted laps (red). Shaded error region represents SEM of the difference of the predicted position from the animal's actual position on the treadmill by session. Dotted lines indicates point of statistical equality (*p*>0.05, Wilcoxon Rank Sum test) between normal and cue shifted lap decoding.

e) Single-neuron receptive field shifts: Distribution of firing location shift on shifted cue laps for all spatially tuned cells, plotted based upon tuning location in normal middle cue laps (n=66 sessions). Red arrowhead indicates actual cue shift distance, and thus marks cells that are directly influenced by the cue. Black arrowhead indicates cells with zero shift, and thus not influenced by the odor cue.

f) Mean number of cells not influenced by the odor cue (shift magnitude<0.05m, "non middle cue influenced") and cells shifting precisely along with the cue (shift magnitude 0.5m+/-0.1m, "middle cue influenced") at each track position (corresponding to cells in regions of black and red arrowheads in "e.", respectively).

Figure 6: Cue presentation leads to suppression of diverse types of DG responses

a) Cue-related suppression of spontaneous, out-of-field firing of DG neurons: Spatial firing rates outside of the spatial receptive field center-of-mass (+/-10cm) for significantly tuned cells during normal and cue omitted laps (n=4,091 cells from 66 sessions).

b) Average out-of-field, spontaneous firing rates from all spatially tuned cells in the presence of the middle cue (black) compared to laps in which the cue is omitted (blue), within the same session. Blue shaded area shows the cue delivery position. (Inset) Mean ± SEM of firing rate within cue region, p<0.001, Signed-rank test, n=4,091 cells.

c) Comparison of the magnitude of cue-related excitation and inhibition by session: Average in-field spatial firing rates (i.e. cue-related excitation), compared with out-of-field, spontaneous firing rates (i.e. cue-related inhibition) at the middle cue location for each session (90-120cm), adjusted for pre-cue firing rate (66 sessions). Yellow, linear regression: p<0.0001, R^2=-0.22.

d) Cue-related suppression of place cell firing: Spatial firing rates of place cells with firing fields at 50-80cm during normal laps (left) and laps in which the cue is shifted to the same 50cm location (right).

e) (Left) Average spatial firing rate for the above 50-80cm receptive field place cells in normal laps where the cue is not presented at this location (Cue+, black), compared with laps in which the cue is shifted to this location (CueΔ, red; diff. in green). (Right), Average peak firing rate of the same cells in "d." and "e.", paired t-test, $p<0.001$, 645 cells.

f) Cue-related suppression of cells responding to other distinct cues: Top: Diagram of intermittent cue pairing experiment. Cues of two different modalities were presented at different locations, interspersed with paired presentation at one of these locations on intermittent laps. Bottom: Example cue-triggered averaged Ca^{2+} transients for a cue cell strongly active when the cue is presented alone (black) but with reduced responses when paired with a different cue, regardless of location (blue, pair location #1, purple, location #2).

g) (Left) Amplitudes of paired cue responses for individual cue cells at the two pairing locations with respect to the response to the cue alone. Gray line represents the diagonal, $p<0.0001$, $R^2=0.91$, n=56 cells, 3 mice. (Right) Average relative response amplitude at the two pairing locations (Location 1: 0.75 ± 0.061, $p1 = 1.45 \times 10^{-4}$; Location 2: 0.73 ± 0.086, $p2 = 0.003$, Wilcoxon Signed Rank Sum test, bar plots are mean ± SEM).

Supplementary Figures

Supplementary Figure 1. Cue associated activity and spatial properties of DG neurons are similar in self-driven and motorized treadmills, and at different treadmill motor velocity.
a) Spatial firing patterns of DG neurons during self-driven locomotion in the treadmill spatial cue task. (Left) normal middle cue lap responses, (middle) cue shift laps, (right) laps in which odor cue is omitted (n=234 cells from 10 sessions in 5 mice). Cells sorted by activity on normal middle cue laps.

b) Average spatial firing rates from all cells in "a." during normal middle cue (black), cue shift (red), and cue omitted laps (blue). (Inset) averaged area under the firing rate curves (Hz*cm) within the middle and shift location reference regions during normal (black bars), cue omitted (blue bar), and cue shifted laps (red bar). $P_{Normal-Omit}$ <0.001, $P_{Normal-Shift}$ =0.03, Wilcoxon signed rank test, error bars are mean ± SEM. Each row across all graphs represents a single cell from the same session.
c) Spatial occupancy of each mouse is calculated as the time spent in each 2cm bin along the 2 meter long treadmill during self-driven (red) or motorized (black) locomotion. Occupancy is shown for normal middle cue laps. During self-driven sessions mice tend to slow down upon odor cue delivery, which we aimed to normalize by using motorized treadmills.
d) Mean spatial information by position showing similar enrichment of neurons with high spatial information around the sensory cues in both self-driven (pink) and motorized (gray) treadmill running. Average Z scored spatial information for cells binned by tuning position during the spatial cue task performed on a motorized vs. self-driven treadmill. Error bars represent ± SEM.
e) Fraction of active cells that are spatially tuned (with at least 0.001 transients per s) in mice advancing the treadmill belt through self-driven locomotion (5 mice, 2 session/mouse) and mice running on the motorized treadmill (8 mice, 2 session/mouse, p=0.4508 Wilcoxon Rank Sum test)
f) Tuning consistency of cue and place cells during self-driven vs. motorized treadmill running. Firing rate correlation between first and last halves of the session, p=0.3197 Wilcoxon Rank Sum test.
g) Variable speed motorized treadmill: Averaged latencies of *reference position-triggered* Ca^{2+} transients of odor cue cells during running on treadmill motorized at fast, normal and slow velocities, χ^2=85.19, p<0.001, $P_{Fast-Normal}$<0.0001, $P_{Fast-Slow}$<0.0001, $P_{Normal-Slow}$=0.0295, Friedman test and Dunn's multiple comparisons tests.
h) Averaged latencies of *cue triggered* Ca^{2+} transients of same odor cue cells in "g." during running on treadmill motorized at fast, normal and slow velocities , χ^2=4.99, p=0.0823, Friedman test. In **(g)** and **(h)**, n=25 odor cue cells, 3 mice, red lines in violin plots show median, black lines show mean.

Supplementary Figure 2. Visual and tactile cues, reward and sequences of cues evoke robust responses in DG.

a) Different modality cues: Firing of DG neurons in response to an LED visual cue. Top: Location of lap cue (green boxes) and an LED cue (red box), during normal and cue-omitted laps. Bottom:

spatial firing rates of 188 spatially tuned neurons (n=6 mice) during the first session of exposure to the middle visual cue on normal (left), and cue-omitted (right) laps.

b) Firing of DG neurons in response to a whisker tactile cue. Top: Location of lap cue (green boxes) and a whisker tactile cue (a piece of rough sandpaper, brown box), during normal and cue omit laps. Bottom: spatial event rates of 199 spatially tuned neurons (n=8 mice) during the first session of exposure to the tactile cue on normal (left), and cue-omitted (right) laps. Each row represents an individual cell, and cells are sorted by their activity in middle cue laps.

c) Average spatial event rate for neurons shown in "a." (LED cue) on normal (black, mean ± SEM), and cue-omitted (blue) laps. Inset shows the averaged area under the firing rate curves (AUC, Hz*cm) within the visual cue region during normal laps (black bar) compared to the same region during cue omitted laps (blue, p<0.0001, Wilcoxon Signed Rank Sum test).

d) Average event rates of neurons shown in "b." (tactile cue) on normal (black), and cue-omitted (blue) laps. Inset shows the averaged area under the firing rate curves (AUC, Hz*cm) within the tactile cue region during normal laps (black) compared to the same region during cue omitted laps (brown, p<0.0001, Wilcoxon Signed Rank Sum test).

e) DG reward responses: Top: Behavioral task with a static operant reward at 75% (150cm) track length, preceded by an olfactory cue in middle of track. Bottom: Spatial event rates for all tuned cells in the static reward task. Laps with olfactory cue in middle of track and reward at 150cm (static rew., left), or laps with reward omitted (omit rew., right). (n=256 cells, 4 mice, 7 sessions).

f) Averaged spatial lick rates over laps, for behavioral sessions in "e.".

g) Average spatial event rates over all cells in "e." during static reward laps (black), and omit reward laps (blue). Olfactory cue zone shaded in yellow, reward zone shaded in blue.

h) Averaged area under the firing rate curves (AUC, Hz*cm) for cue zone and reward zone on static reward (black) and omit reward laps (blue). p = $3.8e^{-9}$, reward zone rate during rew+ vs. rew- (omit rew.) laps, Wilcoxon signed rank test.

i) Cue sequence task: (Top, left) Experimental setup with 3 cues, cue 1 = mint odor (olfactory), cue 2 = LED (visual), cue 3 = isoamyl acetate odor (olfactory). Individual cues are omitted on intermittent laps (right). (Bottom) Spatial event rates for the same tuned cells in normal 3-cue laps (left), and intermittent laps where individual cues are omitted (right 3 columns, as labeled, n=138 cells from 5 mice, 1 session/mouse, sorted by normal 3-cue lap activity).

j) Mean spatial event rates over all cells in "e." on normal laps (black), and laps where cue 1 is omitted (green), cue 2 omitted (red), or cue 3 omitted (blue).

k) Averaged area under the event rate curves (AUC, Hz*cm) within zones corresponding to cues 1-3 on normal laps (black), laps where cue 1 is omitted (green, p= 0.0031), cue 2 omitted (red, p<0.001), or cue 3 omitted (blue, p=0.0047). Wilcoxon signed rank test, error bars are mean ± SEM.

Supplementary Figure 3. Representations of sensory cues and place in a granule cell specific transgenic mouse, Dock10-cre

a) Confocal image projection of brain slice from Dock10-cre mouse injected with FLEX-eYFP AAV in the dorsal dentate gyrus, showing selective labeling of granule cells (blue=DAPI, green=eYFP).

b) Mean spatial information by position in Dock10-cre mice. Average Z scored spatial information for cells binned by tuning position during spatial cue task. WT dorsal dentate gyrus neurons express GCaMP6s (gray bars) and Dock10-cre labelled granule cells express FLEX-GcamP7s (purple bars). Error bars represent SEM, $P_{spkZ\text{-}DG - spkZ\text{-}GC}$ = 0.15, Wilcoxon rank-sum test calculated over cumulative mean spkZ.

c) Spatial firing rates of Dock10/GCaMP7s-expressing granule cells in the spatial cue task with an olfactory cue (IAA/banana). (Left) normal middle cue responses, (middle) cue shift laps, (right) laps in which odor cue is omitted (n=146 cells from 4 sessions in 2 mice). Cells sorted by activity on normal middle cue laps

d) Average spatial firing rates from all cells in "c." during normal middle cue (black), cue shift (red), and cue omitted laps (blue). (Inset) Averaged area under the event rate curves (AUC, Hz*cm) at the middle cue location during normal (black bar) and cue omitted laps (blue bar), and at the shifted cue location during normal (black bar) and cue shifted laps (red bar). $P_{Normal-Omit}$ <0.001, $P_{Normal-Shift}$ =0.03, Wilcoxon matched-pairs signed rank test, error bars are mean ± SEM.

e) A representative *in vivo* two-photon imaging field of view within the dentate gyrus of C57 mice expressing GCaMP6s (left) and Dock10-cre mouse expressing FLEX-GCaMP7s in granule cells (right), including spatially scattered odor (blue) and lap (green) cue cells and place cells (orange).

f) Fraction of inactive cells (gray), active cells (with at least 0.001 transients per s, pink) and spatially tuned cells (orange = place cells, green = lap cue cells, blue = odor cue cells). $Mean_{C57-GcamP6s}$ = 631.68±11.9 cells, from 8 mice 2 sessions each, $Mean_{Dock10Cre-GcamP7s}$ = 566.11±18.4 cells, from 2 mice 2 sessions each. Small, round somata corresponding to granule cells within the granule cell layer were counted using the *Cellpose* algorithm followed by manual inspection in a subset of total sessions imaged from one mouse.

g) Cumulative distribution curve of the firing rate of odor cue, lap cue and place cells in Dock10-cre mice (thin lines) vs. C57 mice (thick lines); Odor Cue $P_{C57-GcamP6s-Dock10Cre-GcamP7s}$=0.11 (blue), Lap Cue $P_{C57-GcamP6s-Dock10Cre-GcamP7s}$ = 0.05 (green), Place cell $P_{C57-GcamP6s-Dock10Cre-GcamP7s}$ = 0.05 (orange), Comparisons are Wilcoxon rank-sum test.

Supplementary Figure 4. Odor cue cells do not emerge from place cells.
a-c) Comparison of cell tuning properties between uncued vs. cued treadmill belts. **a)** Spatial firing rates of all tuned DG neurons within the same fields of view (FOV) recorded during paired sessions on an uncued belt (left) and subsequently an odor cued belt (right). Neurons are ordered

according to the position of their spatial receptive field in each belt (uncued: 155, cue: 249 spatially tuned neurons in 6 mice, 1 session/mouse), but are not matched between uncued and cue plots.
b) Fraction of active cells that are spatially tuned (with at least 0.001 transients per s) in matched FOVs recorded on uncued and cue belts (p=0.0313, Wilcoxon signed-rank test, n=6).
c) Spatial information of all spatially tuned neurons, excluding odor-cue responsive cells, in uncued (n=155) and cued sessions (n=193, p=0.0897, Wilcoxon rank-sum test).
d) Tuning consistency (firing rate correlation between first and last halves of the session) of all spatially tuned neurons in uncued and cued sessions, excluding odor-cue responsive neurons (p=0.5237, Wilcoxon rank-sum test). Boxes, 25th to 75th percentiles; bars, median; whiskers, 99% range.
e-h) Comparison of individual cue cell and place cell firing between uncued and cue belts. **e)** Activity of cross-registered odor cue cells in uncued and cue belts. Cells were classified as odor-cue cells with respect to the cued belt and then their responses were determined for the uncued belt, if active during this session. Note that these "future" cue cells were not spatially tuned in uncued sessions (i.e. not place cells).
f) Fraction of odor-cue (blue), lap-cue (green) and place cells (orange) identified on the cued belt that are cross-registered in uncued sessions (p=0.838, n=6 matched sessions, Friedman's test).
g) Fraction of registered cells that are spatially tuned in uncued belt sessions and encoded the odor cue (blue), lap cue (green), or place (orange) on the cued belt (χ^2=10.88, p=0.0043, $P_{OdorCue-LapCue}$=0.0066, $P_{OdorCue-Place}$=0.0286, $P_{LapCue-Place}$=0.9076). Gray area represent 2.5^{th} and 97.5^{th}% of null distributions for each cell type. Error bars, mean ± SEM.
h) The fraction of cross-registered odor cue cells that were spatially tuned (i.e. place cells or lap cue cells) on the uncued belt is significantly below the null distribution (left) while the fraction of lap-cue (middle) and place cells that are spatially tuned remain significantly above the null distribution. The null distributions are generated for each cell type by randomly permuting cell IDs of all cross-registered neurons and determining overlap among cell types. (Level of significance for 5,000 shufflings **p < 0.01; *p < 0.05, $N_{Odor-cue}$=35, $N_{Lap-cue}$=54, N_{Place}=58).

Supplementary Figure 5. Multisession tracking of individual cue and place cells over time, with different sensory cues, and in different rooms
a) Representative alignment of spatial footprints for cells segmented across sessions within a day or after 1wk. in a single imaging field.

b) Distribution of centroid distances between registered (blue, 1.45 ± 0.97, n=1604) and non-registered (pink, 7.07 ± 1.5, n=2214) neighboring cell pairs (p=6.9 10^{-43}, Wilcoxon Rank Sum test).

c) Fraction of all spatially tuned cells in day 1 session1 (D1), day 1 session 2 (D1') and day 7 (D7) that are registered to at least one other session, χ^2=0.25, p=0.9674, $P_{D1-D1'}$>0.9999, P_{D1-D7}>0.9999, $P_{D1'-D7}$>0.9999, n=8 matched sessions, Friedman and Dunn's multiple comparisons tests.

d) Comparison of average firing rates in all tuned cells across days regardless of tracking χ^2=4.993, p=0.0824, $P_{D1-D1'}$ >0.9999, P_{D1-D7} =0.0781, $P_{D1'-D7}$ =0.4789, N_{D1} =417, $N_{D1'}$ =365, N_{D7}=338, from 8 matched sessions, Kruskal Wallis and Dunn's multiple comparisons tests .

e) Representative fields of view with odor cue cells (blue), lap cue cells (green), and place cells (orange) tracked within a day and across days.

f) Representative Ca^{2+} transients for an odor cue cell that has stable cue-selective activity within one day and over 1wk (left). Transients for a place cell show relatively stable firing location within day but a reorganization of spatial selectivity across days (right). Black and red traces represent normal and cue shifted laps, respectively.

g) Spatial firing rates of neurons tracked between olfactory, visual and whisker tactile cue sessions ordered according to the position of peak activity in olfactory cue sessions (n=157 cells, 6 mice).

h) Photos of the two-photon behavioral setups in two different rooms, used in the room switch experiment (Fig.2f-i). Two-photon microscopes were identical and behavioral apparatuses on both rigs were similar including a similar configuration of lick port and location of odor delivery tube. However, the treadmills varied slightly in geometry and the cabinet in room 2 was larger than the setup in room1. We note that the difference in distal cues in two rig set ups were minor compared to typical arena change experiments in freely moving rodents[35,39,46].

Dual cue location task

Supplementary Figure 6. Spatial modulation of sensory cue responses and timing of cue-related inhibition in the DG.

a) Dual cue location task: Average spatial firing rates for cue cells preferring LED visual cue position 1 (top) and cue position 2 (bottom) on normal laps (blue), laps where the cue at position 1 is omitted (green), and laps where the cue at position 2 is omitted (cyan). (n=89 pos1 cue cells, 67 pos2 cue cells). Note that cue responses for both preferred and non-preferred cue locations are largely eliminated when the respective cue is omitted.

b) Average peak firing rates for all cue cells (both cue 1 and cue 2 preferring) for preferred cue location (left) and non-preferred location (right) on normal laps (blue), laps where the opposite cue is omitted (orange), and laps where the same cue (preferred or non-preferred) is omitted.

(n=156 cells, positions1&2) (mean ± SEM: 0.205 ± 0.017, 0.173 ± 0.020, 0.017 ± 0.008, 0.137 ± 0.016, 0.140 ± 0.024, 0.025 ± 0.008, $P_{Preferred\ Cue}$ = 7.16x10^{-9}, $P_{Non-preferred\ Cue}$ =3.48x10^{-9}, Wilcoxon sign rank test).

c) Average spatial firing rates for cue cells preferring cue position 1 (top) and cue position 2 (bottom) on days 1 (blue, n=50), 2 (green, n=40), 3 (cyan, n=50), and 4 (magenta, n=33) of dual cue location task. Rates for each cell are normalized to their receptive field center rate.

d) Cue location firing rates for preferred and non-preferred cue locations for all cue cells on days 1 (blue, n=50) and 4 (magenta, n=33) of the dual cue location task, showing increased cue spatial preference over the recording period.

e) Average in-field spatial firing rates (i.e. cue-related excitation) in the region around the middle cue location for all tuned neurons, compared with out-of-field, spontaneous firing rates for the same cells (i.e. cue-related inhibition), adjusted for pre-cue firing rate. Note that the position of the excitation peak precedes the trough of inhibition by 5-10cm, corresponding to 0.5-1s at typical treadmill speeds. Blue shaded area shows the cue delivery position.

f) (Left) Session averaged excitatory (within-field) peak position vs. inhibitory (out-of-field) peak position (66 sessions). Red = avg., diagonal (gray line). (Right) Quantification of the position of excitation peak (in cm) compared to the position of inhibition peak, p<0.01, Wilcoxon Signed Rank test.