1 **Parallel processing of sensory cue and spatial information in the Dentate Gyrus**
2

Sebnem N. Tuncdemir[1,2], Andres D. Grosmark[3], Gergely Turi[1,2], Amei Shank[4], Jack Bowler[3],
Gokhan Ordek[2], Attila Losonczy[3], Rene Hen[1,2,5], Clay Lacefield[1,2,5]
[1]Department of Psychiatry, Columbia University, New York, NY 10032, USA
[2]Division of Systems Neuroscience, New York State Psychiatric Institute, New York, NY 10032, USA
[3]Department of Neuroscience, Mortimer B. Zuckerman Mind Brain Behavior Institute, Columbia University,
New York, NY 10027, USA
[4]Columbia College, New York, NY 10027, USA
[5]Lead Contacts
Correspondence: col8@cumc.columbia.edu or rh95@cumc.columbia.edu

Abstract

During exploration, animals form an internal map of an environment by combining information
about specific sensory cues or landmarks with the animal's motion through space, a process
which critically depends on the mammalian hippocampus. The dentate gyrus (DG) is the first
stage of the hippocampal trisynaptic circuit where self-motion and sensory cue information are
integrated, yet it remains unknown how neurons within the DG encode both cue related ("what")
and spatial ("where") information during cognitive map formation. Using two photon calcium
imaging in head fixed mice running on a treadmill, along with on-line sensory cue manipulation at
specific track locations, we have identified robust sensory cue responses in DG granule cells
largely independent of spatial location. Granule cell cue responses are stable for long periods of
time, selective for the modality of the stimulus and accompanied by strong inhibition of the firing
of other active neurons. At the same time, there is a smaller fraction of neurons whose firing is
spatially tuned but insensitive to the presentation of nearby cues, similar to traditional place cells.
These results demonstrate the existence of "cue cells" in addition to the better characterized
"place cells" in the DG, an important heterogeneity that has been previously overlooked. We
hypothesize that the observed diversity of representations within the granule cell population may
support parallel processing of complementary sensory and spatial information and impact the role
of the dentate gyrus in spatial navigation and episodic memory.

Introduction

An animal's location in an environment is highly relevant for guiding its behavior, both to find areas
of potential reward and avoid areas of possible danger. The mammalian hippocampal formation
plays a cardinal role in navigation and spatial memory by integrating self-motion and sensory cue
information into a cognitive map of an environment, exemplified by the presence of "place cells"
selective for specific locations within space[1]. As the initial stage in the 'trisynaptic circuit'[2], the
dentate gyrus (DG) is the first region in the hippocampus to integrate sensory and self-motion
information into a discrete spatial representation, and thus represents the most basic state of
spatial map formation[3,4]. Yet causal evidence for conjunctive encoding of sensory and spatial
information by the principle neurons of the DG, granule cells, is still lacking. While recent studies
have shown that granule cells may be activated by visual[5], tactile[6] and olfactory[7] cues, their

relationship to canonical 'place cells' and their influence on spatial map formation in the DG have
not been investigated.
47
The DG receives its main long-range excitatory inputs from the lateral and medial entorhinal
cortices (LEC and MEC, respectively), and sends mossy fiber projections only to area CA3[8]. The
LEC is thought to primarily represent information about sensory cues, while the MEC is thought
to more prominently encode self-motion information[9–11]. Since individual granule cell dendrites
receive projections from both of these areas, these inputs have the potential to be the basis for
dendritic computations that combine sensory and self-motion information into a discrete spatial
representation[12–14]. In concert with a highly effective winner-take-all process mediated by lateral
inhibition[15–17], these conjunctive representations could facilitate behavioral discrimination of
nearby salient locations with high spatial resolution[18,19]. Such processes may underlie proposed
computational roles of the DG in hippocampal information processing such as pattern separation,
where similar inputs are represented distinctly within the population in order to aid the selectivity
of spatial behavior and memory[16,20].
60
We sought to examine how the DG participates in spatial map formation by recording calcium
activity in large populations of granule cells in the mouse dorsal DG during head-fixed locomotion
on a treadmill. By controlling the administration of sensory cues and their pairing with the animal's
position on the treadmill, we were able to dissect sensory and spatial contributions to granule cell
firing. We found that surprisingly most of the task-associated neurons were highly sensitive to
specific sensory cues presented along the treadmill belt, rather than discrete locations. Cue
responses in single neurons were stable for long periods of time, selective for the modality of the
stimulus and accompanied by strong inhibition of other active neurons. At the same time, a smaller
fraction of neurons exhibited robust spatial tuning independent of local cue presentation, yet were
more context selective. These two channels of information, sensory and spatial, were largely
distinct within the granule cell population, and led us to postulate the existence of "cue cells" in
addition to the better characterized "place cells" of the region. This work suggests that the DG
maintains a largely parallel code for cues in an environment and their location; yet, possesses
specific points of integration, for example through mutual inhibition and the spatial modulation of
cue responses. Such properties of the heterogeneous population of cue cells and place cells may
play a role in higher level functions of the dentate gyrus such as pattern separation and spatial
map formation.
78
79 **Results**
80
To investigate the interaction between sensory and spatial representations in the dentate gyrus
(DG), we used two photon calcium imaging of large populations of granule cells in head-restrained
mice running on a treadmill as a virtual linear track[21]. Mice were injected unilaterally with
rAAV.Syn.GCaMP6s to express the genetically encoded calcium indicator GCaMP6s in the dorsal
DG. This was followed by the implantation of a chronic imaging window above the hippocampal
fissure, which allowed us to image the calcium dynamics of neurons in the granule cell layer
during treadmill behavior (Fig. 1a). After habituation, mice were trained to run in order to receive
randomly delivered water rewards on a 2m-long treadmill belt, during 15 minute sessions (Fig.1b,

methods). Movies of population calcium imaging data were motion corrected offline[22], and the
activity of putative single neurons was isolated[23,24]. Neurons with significant spatial tuning of
calcium activity along the track were identified using previously described methods[21,25], and in a
subset of sessions activity of individual neurons was tracked in multiple conditions over multiple
sessions[26].
94
In order to isolate the effects of locomotion on hippocampal activity[27], the treadmill was motorized
at a constant speed, adjusted for each mouse (motorized velocity = 10.11 ± 0.64 cm/s, self-driven
velocity = 12.76 ± 2.45 cm/s). Despite high velocity dependence of DG activity in freely running
mice (Supplementary Fig. 1a), there were no significant differences in the fraction of spatially
selective neurons or their mean firing rates in mice running on the motorized treadmill compared
to mice advancing the treadmill belt through self-driven locomotion (Supplementary Fig. 1b, c),
however mean spatial tuning was higher in mice running on the motorized treadmill
(Supplementary Fig. 1d).
103
104 **Granule Cell Cue Responses**
105
To investigate the relationship between discrete sensory cues and spatial representations within
the DG, we introduced a 1s odor pulse delivered in the middle of the track on each lap as a
dynamic spatial cue, in addition to a tactile cue at the lap boundary. We found that the majority of
spatially selective neurons exhibited receptive fields near the lap boundary and middle locations,
corresponding to the lap cue and middle cue positions, respectively (Fig. 1c, 57% of cells with
peak activity within 10cm of cues, 43% >10cm from cues, $p<0.0001$, Mann-Whitney test). Across
the population, average firing rates (Fig. 1e) and spatial information (Supplementary Fig. 3a) of
spatially tuned neurons were also higher in positions corresponding to sensory cues. These
neurons could however be place cells that are enriched at the locations of salient stimuli, as has
been shown in area CA1[28–32] or alternately could be directly driven by the stimulus[7]. To test this,
the olfactory cue was omitted or shifted 1/4th of the track length once every 3-5 trials interleaved
throughout the session (41 ± 2 total laps/session) in order to dissociate cue responses from track
location. Under these conditions, a majority of neurons normally active in the middle of the track
shifted their firing position to match the new location of the odor in cue-shift trials (Fig. 1d, middle,
79.07 ± 2.76 %) and exhibited reduced activity in cue-omitted trials (Fig. 1d, right, 78.2 ± 3.87
%), compared with normal middle cue laps (n = 285 spatially tuned neurons from 8 mice). In
contrast, neurons firing at locations corresponding to the invariant lap cue were unchanged in
omit and shift laps. On trials with a shifted middle cue, average firing rates were higher within the
new cue region (red, inset p=0.0234 Signed Rank test), and lower at the normal cue location on
cue-omitted trials (blue, inset $p<0.00001$, Signed Rank Sum test) compared to the corresponding
positions during normal trials (Fig. 1e, inset). Thus, a substantial population of spatially tuned DG
neurons in the virtual linear track environment are in fact active directly in response to presentation
of cues at those locations, rather than the locations themselves.
129
130 Comparable population responses were found in response to cues of other sensory modalities,
131 such as visual or tactile cues (Supplementary Fig. 2a-d), as well for liquid rewards (Supplementary
132 Fig. 2e-h). In addition, increasing the complexity of the environment with two additional cues at

other track locations results in an additive pattern of single cue responses (Supplementary Fig. 2i-k), suggesting that strong cue responses are not limited to situations with a single cue alone. Cue responses were also similar for cells imaged in the dorsal DG of a granule cell-specific *Cre* transgenic mouse (Dock10 Cre[33], Supplementary Fig. 3b-d,), indicating that sensory cue representations are indeed a property of the granule cell population rather than arising from other local neuron types such as mossy cells or inhibitory interneurons[6].

Next, we divided the spatially tuned population of granule cells into three groups for subsequent analyses based on the position of their spatial fields and their activity during cue manipulation (omit or shift) trials. The three response types are illustrated for one session (Fig. 1f): 1) cells with spatial fields within the middle cue region that closely track the changes in cue presentation ("odor-cue cells", top); 2) cells with spatial fields around the lap boundary cue on the treadmill belt ("lap-cue cells", middle); and 3) the remaining spatially tuned cells with receptive fields outside of the cue locations throughout the track ("place cells", bottom). These three groups, odor-cue, lap-cue, and place cells, constituted 22.5 ± 2%, 47.1 ± 2% and 30.4 ± 1% of the spatially tuned cells within the imaging field of view, respectively (Supplementary Fig. 3e). We found that cells with similar response types did not cluster together spatially within the imaging field and the groups did not exhibit significant differences in overall mean firing rates (Supplementary Fig. 3e-g). Spatial coding properties however differed between cue and place coding populations of neurons. Both populations of cue cells (lap cue and olfactory cue) showed higher average spatial information (Fig. 1g, χ^2=48.47, p<0.0001, Kruskal-Wallis test) and had more consistent spatial firing between the first and the second half of each session than place cells (Fig. 1h, χ^2=32.60, p<0.0001, Kruskal-Wallis test).

In order to probe the emergence of selective firing in these populations, we identified the lap in which responses began to robustly occur within the preferred spatial location during the first session of exposure to the odor cue (field onset lap [34]). We found that for odor and lap cue cells the majority had spatial fields that appeared within the first five laps (38/65 (58%), 75/136 (55%), respectively) while the majority of place cells emerged later in the session (52/84 (62%) within 10 laps, Fig. 1i, χ^2=9.29, p=0.0096, Kruskal-Wallis test). In agreement with the later emergence of place cells, we found that the in-field firing rates of place cells were significantly smaller than those of odor and lap cue cells only within the first 5 laps (Fig. 1i, inset, 2-way ANOVA, main effect of cell type: $F_{2,1473}$=6.73, p=0.0012; cell type × lap number interaction $F_{4,1473}$=2.44, p=0.04; main effect of lap number: $F_{2,1473}$=0.28, p=0.7). Additional experiments also showed that on an uncued treadmill belt prior to olfactory cue sessions, the majority of 'future' odor cue cells were either inactive or had low spatial tuning (Supplementary Fig. 4), suggesting that cue cells rapidly emerge *de novo* and are not place cells that remap to new cue locations. Taken together, these results demonstrate that sensory cue representations are more reliable and appear with less exposure than place cell representations, and arise from an independent population of granule cells.

Stability and Specificity of Sensory Cue Responses

To further characterize the stability and coding specificity among the granule cell subpopulations, we investigated the responses of individual neurons over time and with respect to different

sensory cues (Fig. 2a, b). We utilized an analytical method [26] to track cells over multiple sessions
and were able to find substantial numbers of the same cells active in different sessions within a
day or 1 week later in the same fields of view (Supplementary Fig. 5a, b). Although not all cells
were identified in every session, a similar percentage of spatially selective neurons was registered
in all sessions, which was confirmed by visual inspection to ensure that cells appeared consistent
in the anatomical images (Supplementary Fig. 5c-g).

Between any two sessions, over days or with different cue modalities, the cells encoding the
invariant lap cue were the largest fraction of cells that remained active and maintained their
response type (e.g. cue type and/or place, green circles in Fig. 2c). Odor cue cells fired reliably
to the same olfactory cue over long periods of time (blue circles in red and blue shaded area, Fig.
2c), but were largely unresponsive to cues of other modalities presented at the same position
(green shaded area in Fig. 2c, χ^2=19.78, p<0.0001, Kruskal-Wallis test). Conversely, a lower
percentage of place cells maintained their response type across days compared to cue cells
recorded in the same sessions using the same cues (orange circles in blue shaded area, Fig. 2c,
χ^2=14.36, p<0.001, Kruskal-Wallis test). Furthermore when we examined specifically whether cue
cells become place cells between these sessions, or vice versa, we found that categorical cue
and place representations remain extremely stable within the DG population (Fig. 2d, p<0.001,
Rank Sum test).

We further examined the degree of stability and specificity of cue and place cell tuning by cross-
correlating spatial firing rates for individual registered cells over time and with respect to different
sensory cues (Fig. 2e). For this analysis, we computed the correlation of spatial firing rates for
cells registered during different sessions within a day and one week later, or with a different
sensory cue modality. In line with the stimulus selectivity analysis described above, we observed
that middle location odor cue cells displayed significantly lower correlations in sessions with a
different sensory cue, compared to separate sessions using the same olfactory cue on the same
day or one week later (left, χ^2=27.88, p<0.0001, Kruskal Wallis test). Thus, cells that responded
to a cue of one modality were unlikely to respond to cues of other modalities, despite a similar
spatial location of the cue between sessions. Correlations in the activity of individual place cells
over 1 wk. were significantly lower than on the same day, and were lower than both cue cell
populations, again indicating lower stability for place than cue representations (left, χ^2=18.79,
p<0.0001, Kruskal Wallis test). Lap cue cells did not display significant changes in their firing rate
correlations between sessions on the same day, across days or with different sensory cues, and
were therefore especially stable.

Responses of individual odor cue cells were also highly correlated in sessions recorded in
different rooms on the same day (Fig. 2f-h), indicating that distal ambient cues have a limited
influence on the stability of sensory cue representation in DG granule cells. Despite the high
stability of cue cells, place cells were significantly less consistent between rooms than cue cells
in the same sessions (Fig. 2g, h, p<0.001, Kruskal-Wallis test), and were similar to that seen in
place cells over 1wk when measured in the same room (Fig. 2e), suggesting that place cells are
more context selective than cue cells. Taken together, these results suggest that sensory cues
are represented by a stable subpopulation of neurons that is highly selective for specific cues

while purely spatial representations are less stable and undergo progressive reorganization over
time and in different global contexts.
223
Spatial Modulation of DG Cue Responses

The juxtaposition of inputs from the lateral and medial entorhinal cortex onto the dendrites of
individual granule cells has been hypothesized to underlie a conjunctive code for sensory cues
and their spatial location[3,4]. We therefore examined the influence of spatial location on cue
responses ("spatial modulation") by tracking odor cue cells through multiple sessions with different
cue-location pairing, either with an intermittently shifted cue as in previous experiments or in
separate sessions with the same cue at random locations each lap (Fig. 3a, b). On average, cue-
triggered Ca^{2+} response amplitudes for individual cue cells (a proxy for neuron action potential
burst firing rate) were smaller when cues were presented at the infrequent "shift" location or at
random locations, when compared to their responses at the more frequent middle location (Fig.
3c, d, p<0.0001, Friedman test, n=101, 5 mice). Thus while cue cells tend to respond to the same
sensory cue regardless of location, the strongest responses occur when a cue is presented at the
same place repeatedly.

To further examine the spatial modulation of cue responses, we performed a set of experiments
where a visual cue was delivered at multiple locations on the track to measure selectivity of
granule cell responses to cues presented consistently at distinct locations ("dual location cue",
Fig. 3e), similar to recent work by other groups using virtual reality environments[35,36]. As in our
previous experiments with a single cue, each cue was omitted on a subset of laps to isolate cue
cells as opposed to place cells present in the vicinity of cues (Supplementary Fig. 6a, b). To
measure the extent of preference for one cue location versus the other, we calculated the spatial
modulation index for each cue cell as the ratio of the firing rates in the non-preferred cue location
versus the preferred cue location. Cue cells in the first session exhibited a range of spatial
modulation indices, but over the population their event rates were significantly modulated by the
cue location (p<0.00001, Wilcoxon sign rank test on day 1 spatial modulation index, 53% of
individual neurons significantly spatially modulated by location shuffle). Spatial modulation also
increased over several days of dual cue presentations (p=0.0125 Wilcoxon ranked sum test, day
1 vs 2, 3, or 4), with 81% of cue cells exhibiting significant spatial modulation by day 4 (Fig. 3g, h
and Supplementary Fig. 6c, d). This suggests that dentate cue cells are acutely modulated by the
spatial location of two identical cues but increase their preference for a single location over days.
255
Effects of Cues on Spatial Encoding in the DG

While the most robust activity in the DG was found within the cue cell population, the existence
of the smaller population of place cells in uncued locations suggests that spatial activity in these
cells is referenced to one of the two cues on the otherwise featureless treadmill track in order to
encode a unique location. We therefore examined the effect of manipulations of the variable cue
on the subsequent spatial encoding of place cells, in order to judge the degree to which this cue
acts as a landmark. For example, if place cells were acutely referenced to the nearest cue we
would expect the place fields of cells following the variable middle cue to shift on cue shifted laps.

To examine the relationship between spatial firing patterns in normal cue laps to those in cue
shifted laps, we first calculated population vector (PV) correlations of firing rates across all
spatially tuned cells on each lap (Fig. 4a and Supplementary Fig. 7a). While the correlation was
higher in the vicinity of the cues between normal laps, we observed a dramatic decrease in PV
correlation between normal and cue shifted laps which was confined to the area immediately
around the cue itself, but did not extend much beyond the cue location.
271
To further evaluate the manner in which cue associated activity contributes to spatial encoding, a
spatial Bayesian decoder was constructed from the firing rate vectors of all spatially tuned cells
(4,091 cells from 66 sessions)[25]. In order to establish a non-biased estimate of the position during
treadmill running, the decoding was performed using a 5 fold cross validation approach in which
the 1/5th of decoded laps were held out from the training set. Post-reconstruction, we divided the
data according to the lap types (Fig. 4b and Supplementary Fig. 7b, c). The decoder accuracy
was higher in normal (median, 10.6 cm) compared to both cue shift (median, 19.6 cm) and cue
omitted laps (median, 18.4 cm, $p<0.0001$, Wilcoxon rank sum test, Fig. 4c and Supplementary
Fig. 7d, e), indicating that cue manipulations affect the accuracy of spatial coding by the DG
population. However, while spatial decoding was initially strongly perturbed by the cue shift, the
decoder error on shift laps soon converged to that of normal laps, well before the subsequent lap
cue (Fig. 4d, dotted line). Together these results suggest that as a population, the spatial encoding
of granule cells is affected by local cues for a limited time, rather than persistently altering their
estimate of the animal's position.
286
Population vector correlations and Bayesian decoding are average measures of spatial responses
for the granule cell population as a whole, which might obscure differences in how the activity of
individual cells is referenced to local or global cues to encode an animal's location. We therefore
examined responses of individual granule cells by measuring correlations between their spatial
firing in normal and cue-shifted laps. In doing so we found two general classes of single-cell
responses: some cells were acutely affected by the intermittently shifting odor cue, while others
were referenced to the stable lap boundary cue (Fig. 4e, f and Supplementary Fig. 7f, g). The
majority of cells that shifted their firing in reference to the new cue location (0.5m) were normally
active immediately in response to the middle cue ("cue-influenced"), with a smaller number
persistently affected by the shift prior to the next lap cue. In contrast, a relatively constant number
of tuned cells along the track maintained their normal firing location with reference to the distant
lap boundary cue despite the shifted local cue ("non cue-influenced"). Notably, there was a small
enrichment of such place cells immediately prior to the cue but unaffected by the cue shift,
indicating a spatial response predictive of salient cues. These data suggest that the DG population
maintains independent spatial reference frames based upon distinct local and global cues within
an environment. This code however differs depending upon the spatial stability of the cues: the
major effects of a variable cue presentation are limited to cells that fire directly in response to the
cue, while a majority of place cells over the track are preferentially influenced by the stable cue
as a landmark.
306
307
308

Cue-related Inhibition of DG Responses

Spatial receptive fields of granule cells are hypothesized to form within a competitive network in which lateral inhibition mediated by GABAergic interneurons enforces sparse encoding that may aid pattern separation[15]. The robust and highly selective cue responses of granule cells in our behavioral paradigm allowed us to examine patterns of local-circuit inhibition within the DG network. First, we sought to analyze the effect of cue responses on nonselective background firing observed within the spatially tuned granule cell population. By comparing out-of-field firing rates at the middle cue location in normal laps with laps where the cue is omitted, we identified a significant reduction in background firing of the granule cell population during the cue presentation (Fig. 5a, p=0.0004, Signed Rank Sum test). This suppression was absent on cue-omitted laps and generally co-varied with the mean amplitude of cue-related excitation on a session-by-session basis (Fig. 5b, R=0.22, p<0.0001, Signed Rank Sum test). Furthermore, the timing of the peak of this inhibition was delayed with respect to the excitatory cue response in these sessions (Supplementary Fig. 8a, b). These data show that sensory cue input is able to suppress 'noisy', nonspecific activity in the vicinity of the cue, potentially through lateral inhibition.

In addition to suppressing nonspecific activity in the DG network, we sought to determine the effect of cue responses on activity in place cells and in cells responding to other cues. By examining place cells with receptive fields normally at the shift location, we found that firing rates were strongly suppressed on shift laps when a cue was presented at this location compared to normal laps when a cue was not present here (Fig. 5d, e). Furthermore when two distinct cues were normally presented at different locations, shifting of the position of one cue to the other cue location typically had the effect of suppressing the responses in both cue populations (Fig. 5f, g), indicating mutual inhibition among granule cells encoding different cues. Thus in addition to the inhibition of nonspecific firing, cue responses lead to overall inhibition of place cell responses, and responses to competing cues as well. Taken together, our results suggest that cue-related activity in granule cells results in strong suppression of responses in neighboring cells via lateral inhibition.

Discussion

By imaging calcium activity in large populations of dentate granule cells during head-fixed spatial behaviors on a virtual linear track, we have shown that a major population of task-selective neurons is highly sensitive to specific sensory cues rather than to discrete locations. The ability to dynamically manipulate a sensory cue and its association with locations on the treadmill track allowed us to isolate the population of cue cells, in addition to the complementary population of spatially tuned place cells recorded in the same sessions.

Properties of cells within these groups also differed: sensory cue responses were highly-tuned and remarkably robust across contexts and over time- significantly more so than the responses of the canonical place cell population. And while cue cell responses were not present initially, they emerged rapidly during the first exposures to a cue while place cell responses emerged more slowly. These findings suggest that dentate cue cells stably represent basic features of sensory

353 cues in an environment, while place cells constitute a more dynamic population that gradually
354 adapts to current conditions, for example, by integrating between stable cues in order to provide
355 an accurate estimate of the animal's location when no cues are present[37].
356
357 **Heterogeneous properties of Granule Cells**
358
The robust and consistent nature of GC cue responses suggests that cue-selective populations
may have been similarly present in previous experiments recording DG activity *in vivo*, however
the lack of precise stimulus control made it impossible to distinguish cue-responsive versus place-
responsive components[5,6,38]. Indeed, spatial receptive fields of individual granule cells measured
in different environments with similar sensory cues display mixed levels of changes in their firing
rates, with a subset of granule cells completely remapping their firing fields while others show
more stable dynamics[5,38–41]. Based upon our results, these subgroups may correspond to place
cells and cue cells, respectively. A major impact of sensory cues on DG firing largely independent
of spatial context might also explain the relatively lower context selectivity observed in the DG
compared with other hippocampal subfields, at least when measured in different contexts that
contain the same or similar sensory cues[5,38–41]. To support this notion, we found that DG cue cells
were largely stable when recorded over long periods of time, with different cue positions, or in
different room contexts, while place cells were not.
372
Previous work has demonstrated that the two major long range inputs to the dentate gyrus, the
lateral and medial entorhinal cortex (LEC and MEC), are involved with processing functionally
distinct information[9,10]. This raises the possibility that our "cue cells" may be driven primarily by
sensory inputs from the LEC, while "place cells" are driven by self-motion information relayed
from the MEC, for example in the form of grid cell responses. The segregation of these properties
within the overall granule cell population suggests that these streams of information remain largely
separated at the level of the DG, similar to the functional heterogeneity observed in other
hippocampal subfields[42,44]. In these regions such heterogeneity among principal neurons has
been hypothesized to be critical for efficient encoding of spatial representations within a local
population[42–44]. Heterogeneity has also been observed in immediate early gene expression
profiles within DG neurons encoding individual memory engrams[45] which are differentially
targeted by entorhinal afferent pathways[8,45–47]. These findings are in line with our study
demonstrating that sensory and spatial information remain partially separate at the level of the
DG, and suggest that multiple channels of spatial and non-spatial information contribute distinctly
to local and down-stream computations. This separation is also consistent with current ideas that
cue-based and path-integration based navigation are complementary, rather than integrated, in
order to produce place-specific firing both in locations where landmark cues are present as well
as in between cues[48].
391
Furthermore, the separation of cue and place responses within the granule cell population
suggests that specific inputs from either LEC or MEC dominate individual granule cell firing. This
might result from the unique cellular properties of granule cells, local circuits such as inhibitory
networks[49], as well as the laminated pattern of LEC and MEC inputs onto granule cell dendrites[8].
Inputs from distinct cortical afferents can be strongly filtered due to hyperpolarized resting

membrane potentials and substantial attenuation of regenerative dendritic activity in granule
cells[12,13]. Furthermore, short coincidence-detection windows for EPSPs[50] may result in a high
activation threshold of each cortical input. Such gating of inputs by postsynaptic neuronal
excitability may provide a cellular substrate for our observation that these channels of information
remain largely separated at the level of the DG. Differences in input sources may also explain the
distinct time course of emergence of cue and place cell responses, which could reflect divergent
mechanisms for synaptic plasticity between LEC and MEC inputs, differential inhibition of these
pathways, or other factors such as associative vs. non-associative learning. Future studies
investigating whether cue and place encoding populations receive different inputs from LEC and
MEC, as well as the molecular basis of the heterogeneity among granule cell populations, will
allow for a more detailed inquiry into the functional diversity in DG granule cells. Such studies
may also allow us to selectively target cue and place cells in order to manipulate their activity and
determine their roles during discrimination behaviors.
410
411 **Functional significance of dentate gyrus cue cell properties**
412
But how do these results inform our ideas about how the DG participates in pattern separation
and spatial map formation? First, we found that although individual granule cells respond to the
same cues when presented in different locations, the strength of this activity was spatially
modulated. Differential encoding of cues based upon their spatial location may contribute to
pattern separation by differentiating between similar cues present in different places within an
environment, or in distinct contexts. We observed that spatially modulated cue responses also
increased over days when a single cue was repeatedly presented at multiple locations, indicating
plasticity in location specificity and potentially progressive enhancement of pattern separation.
Notably, responses were also on average larger for odor cues presented at a single location, as
opposed to shifted or random locations. This suggests that cue responses are more robust for
stimuli paired repeatedly with one particular location, which may indicate dendritic integration of
sensory and spatial information. Conjunctive encoding of sensory cues and their locations within
an environment could play a role in establishing landmarks for spatial navigation. In support of
this idea, we found that most place cells were referenced to the stable lap cue rather than the
variable middle cue, indicating that this cue is preferentially utilized as a landmark.
428
Our study also shows that cue responses lead to potent inhibition of three distinct types of granule
cell activity: nonspecific "noisy" spiking activity, the spatial firing of place cells normally active in
a location, and the activity of cells responsive to other distinct sensory cues. This supports the
hypothesis of a competitive network in the DG enforced by strong inhibition, which has been
suggested to contribute to pattern separation. In addition, suppression of weaker place cell
responses by cue-related activity may indicate that the network is organized to utilize the
strongest and perhaps the most informative parameters, such as landmark cues, in order to
establish an animal's location in space.
437
438 Furthermore, rate modulation of granule cell cue responses resulting from spatial location or
439 association with other cues is itself a form of "rate remapping", a feature frequently attributed to
440 the dentate gyrus in pattern separation. Thus spatial location, as well as the juxtaposition of

multiple cues and their associated patterns of inhibition, may create a rate modulated landscape of granule cell activity specific to the current context. The overall pattern of contextually modulated cue responses in the DG may form the basis for recruitment of distinct populations of neurons, or "global remapping", observed in area CA3. Slower emergence and lower stability over sessions in the place cell population may also point to a role for these neurons in context selectivity. Indeed place cells differed more greatly than cue cells recorded under similar conditions in different rooms.

The properties of cue and place cells in the dentate gyrus have the hallmark of features one might expect from the first stage of spatial map formation in the hippocampus: mostly distinct but slightly mixed encoding of sensory cues, or "what", and place, or "where". While place cells in the DG are referenced to stable spatial cues in order to determine a unique location of firing given movement-dependent grid cell inputs from MEC, cue cells are mostly driven by sensory input from the LEC, but modulated by the spatial context of the cues. Such complementary channels of information may be further refined by downstream hippocampal areas into an integrated spatial map that encodes relationships between important features such as spatial cues and goals, which can be used to organize adaptive behavior. Uncovering the functional heterogeneity of cue and place cells in the dentate gyrus helps us to better understand the basic neural mechanisms underlying our sense of orientation in complex environments by utilizing both landmarks and self-motion information to guide our movements through space.

Acknowledgements

We would like to thank Jack Berry for histological photos and other members of the Hen lab for critical insight throughout this project. This work was funded by NIMH T32 MH015144 (S.N.T.), NYSTEM-C029157 (G.O., G.T., C.O.L, R.H.), NIH S10 OD018464, Revson Senior Fellowship in Biomedical Science (A.D.G.), NIMH R01 AG043688 (R.H.), MH068542 (R.H.), NIMH R01 MH100631 (A.L.), NINDS R01NS094668 (A.L.), and NINDS U19NS104590 (A.L.).

Author Contributions

Conceptualization, C.O.L., S.N.T., and R.H.; Methodology, C.O.L., S.N.T., A.S.; Software, C.O.L., S.N.T., A.D.G., J.B.; Formal Analysis, S.N.T., C.O.L.; Investigation, S.N.T, C.O.L.; Resources, C.O.L., A.D.G., G.T., J.B., A.L., G.O.; Writing, Review and Editing, S.N.T., C.O.L., R.H., A.D.G., A.L.; Supervision, R.H and A.L.

Declaration of Interests

The authors declare no competing interests.

References

1. O'Keefe, J. & Dostrovsky, J. The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. *Brain Res.* **34**, 171–175 (1971).
2. Amaral, D. G., Scharfman, H. E. & Lavenex, P. The dentate gyrus: fundamental neuroanatomical organization (dentate gyrus for dummies). *Prog. Brain Res.* **163**, 3–22 (2007).
3. Kesner, R. P. An analysis of the dentate gyrus function. *Behavioural Brain Research* **254**, 1–7 (2013).

4. Lee, J. W. & Jung, M. W. Separation or binding? Role of the dentate gyrus in hippocampal
mnemonic processing. *Neuroscience & Biobehavioral Reviews* **75**, 183–194 (2017).
5. Hainmueller, T. & Bartos, M. Parallel emergence of stable and dynamic memory engrams in
the hippocampus. *Nature* **558**, 292–296 (2018).
6. Jung, D. *et al.* Dentate granule and mossy cells exhibit distinct spatiotemporal responses to
local change in a one-dimensional landscape of visual-tactile cues. *Sci Rep* **9**, 9545 (2019).
7. Woods, N. I. *et al.* The Dentate Gyrus Classifies Cortical Representations of Learned
Stimuli. *Neuron* **107**, 173-184.e6 (2020).
8. Witter, M. P. The perforant path: projections from the entorhinal cortex to the dentate gyrus.
Prog. Brain Res. **163**, 43–61 (2007).
9. Hargreaves, E. L., Rao, G., Lee, I. & Knierim, J. J. Major dissociation between medial and
lateral entorhinal input to dorsal hippocampus. *Science* **308**, 1792–1794 (2005).
10. Knierim, J. J., Neunuebel, J. P. & Deshmukh, S. S. Functional correlates of the lateral and
medial entorhinal cortex: objects, path integration and local-global reference frames. *Philos.*
Trans. R. Soc. Lond., B, Biol. Sci. **369**, 20130369 (2014).
11. Wang, C. *et al.* Egocentric coding of external items in the lateral entorhinal cortex. *Science*
362, 945–949 (2018).
12. Kim, S., Kim, Y., Lee, S.-H. & Ho, W.-K. Dendritic spikes in hippocampal granule cells are
necessary for long-term potentiation at the perforant path synapse. *Elife* **7**, (2018).
13. Krueppel, R., Remy, S. & Beck, H. Dendritic integration in hippocampal dentate granule
cells. *Neuron* **71**, 512–528 (2011).
14. McNaughton, B. L., Douglas, R. M. & Goddard, G. V. Synaptic enhancement in fascia
dentata: Cooperativity among coactive afferents. *Brain Research* **157**, 277–293 (1978).
15. de Almeida, L., Idiart, M. & Lisman, J. E. The input-output transformation of the hippocampal
granule cells: from grid cells to place fields. *J. Neurosci.* **29**, 7504–7512 (2009).
16. Marr, D. Simple memory: a theory for archicortex. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.*
262, 23–81 (1971).
17. Treves, A. & Rolls, E. T. Computational analysis of the role of the hippocampus in memory.
Hippocampus **4**, 374–391 (1994).
18. Hunsaker, M. R., Rosenberg, J. S. & Kesner, R. P. The role of the dentate gyrus, CA3a,b,
and CA3c for detecting spatial and environmental novelty. *Hippocampus* **18**, 1064–1073
(2008).
19. Xavier, G. F. & Costa, V. C. I. Dentate gyrus and spatial behaviour. *Prog.*
Neuropsychopharmacol. Biol. Psychiatry **33**, 762–773 (2009).
20. Knierim, J. J. & Neunuebel, J. P. Tracking the flow of hippocampal computation: Pattern
separation, pattern completion, and attractor dynamics. *Neurobiology of Learning and*
Memory **129**, 38–49 (2016).
21. Danielson, N. B. *et al.* Distinct Contribution of Adult-Born Hippocampal Granule Cells to
Context Encoding. *Neuron* **90**, 101–112 (2016).
22. Pnevmatikakis, E. A. & Giovannucci, A. NoRMCorre: An online algorithm for piecewise rigid
motion correction of calcium imaging data. *J. Neurosci. Methods* **291**, 83–94 (2017).
23. Giovannucci, A. *et al.* CaImAn an open source tool for scalable calcium imaging data
analysis. *Elife* **8**, (2019).
24. Pachitariu, M. *et al.* Suite2p: beyond 10,000 neurons with standard two-photon microscopy.
bioRxiv 061507 (2016) doi:10.1101/061507.
25. Grosmark, A. D. & Buzsáki, G. Diversity in neural firing dynamics supports both rigid and
learned hippocampal sequences. *Science* **351**, 1440–1443 (2016).
26. Sheintuch, L. *et al.* Tracking the Same Neurons across Multiple Days in Ca2+ Imaging Data.
Cell Rep **21**, 1102–1115 (2017).

27. Fuhrmann, F. *et al.* Locomotion, Theta Oscillations, and the Speed-Correlated Firing of
Hippocampal Neurons Are Controlled by a Medial Septal Glutamatergic Circuit. *Neuron* **86**,
1253–1264 (2015).
28. Bourboulou, R. *et al.* Dynamic control of hippocampal spatial coding resolution by local
visual cues. *eLife* **8**, e44487 (2019).
29. Hetherington, P. A. & Shapiro, M. L. Hippocampal place fields are altered by the removal of
single visual cues in a distance-dependent manner. *Behav. Neurosci.* **111**, 20–34 (1997).
30. Hollup, S. A., Molden, S., Donnett, J. G., Moser, M. B. & Moser, E. I. Accumulation of
hippocampal place fields at the goal location in an annular watermaze task. *J. Neurosci.* **21**,
1635–1644 (2001).
31. O'Keefe, J. & Conway, D. H. Hippocampal place units in the freely moving rat: why they fire
where they fire. *Exp Brain Res* **31**, 573–590 (1978).
32. Wiener, S. I., Paul, C. A. & Eichenbaum, H. Spatial and behavioral correlates of
hippocampal neuronal activity. *J. Neurosci.* **9**, 2737–2763 (1989).
33. Kohara, K. *et al.* Cell type-specific genetic and optogenetic tools reveal novel hippocampal
CA2 circuits. *Nat Neurosci* **17**, 269–279 (2014).
34. Sheffield, M. E. J., Adoff, M. D. & Dombeck, D. A. Increased Prevalence of Calcium
Transients across the Dendritic Arbor during Place Field Formation. *Neuron* **96**, 490-504.e5
(2017).
35. Saleem, A. B., Diamanti, E. M., Fournier, J., Harris, K. D. & Carandini, M. Coherent
encoding of subjective spatial position in visual cortex and hippocampus. *Nature* **562**, 124–
127 (2018).
36. Zhao, X., Wang, Y., Spruston, N. & Magee, J. C. Membrane potential dynamics underlying
context-dependent sensory responses in the hippocampus. *Nature Neuroscience* **23**, 881–
891 (2020).
37. Draht, F. *et al.* Experience-Dependency of Reliance on Local Visual and Idiothetic Cues for
Spatial Representations Created in the Absence of Distal Information. *Front Behav Neurosci*
11, 92 (2017).
38. Leutgeb, J. K., Leutgeb, S., Moser, M.-B. & Moser, E. I. Pattern separation in the dentate
gyrus and CA3 of the hippocampus. *Science* **315**, 961–966 (2007).
39. Danielson, N. B. *et al.* In Vivo Imaging of Dentate Gyrus Mossy Cells in Behaving Mice.
Neuron **93**, 552-559.e4 (2017).
40. GoodSmith, D. *et al.* Spatial Representations of Granule Cells and Mossy Cells of the
Dentate Gyrus. *Neuron* **93**, 677-690.e5 (2017).
41. Senzai, Y. & Buzsáki, G. Physiological Properties and Behavioral Correlates of
Hippocampal Granule Cells and Mossy Cells. *Neuron* **93**, 691-704.e5 (2017).
42. Cembrowski, M. S. & Spruston, N. Heterogeneity within classical cell types is the rule:
lessons from hippocampal pyramidal neurons. *Nat. Rev. Neurosci.* **20**, 193–204 (2019).
43. Chelaru, M. I. & Dragoi, V. Efficient coding in heterogeneous neuronal populations. *Proc.*
Natl. Acad. Sci. U.S.A. **105**, 16344–16349 (2008).
44. Soltesz, I. & Losonczy, A. CA1 pyramidal cell diversity enabling parallel information
processing in the hippocampus. *Nat. Neurosci.* **21**, 484–493 (2018).
45. Sun, X. *et al.* Functionally Distinct Neuronal Ensembles within the Memory Engram. *Cell*
181, 410-423.e17 (2020).
46. Luna, V. M. *et al.* Adult-born hippocampal neurons bidirectionally modulate entorhinal inputs
into the dentate gyrus. *Science* **364**, 578–583 (2019).
47. Erwin, S. R. *et al.* A Sparse, Spatially Biased Subtype of Mature Granule Cell Dominates
Recruitment in Hippocampal-Associated Behaviors. *Cell Rep* **31**, 107551 (2020).
48. Poucet, B. *et al.* Independence of landmark and self-motion-guided navigation: a different
role for grid cells. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **369**, 20130370 (2014).

49. Pernía-Andrade, A. J. & Jonas, P. Theta-gamma-modulated synaptic currents in
hippocampal granule cells in vivo define a mechanism for network oscillations. *Neuron* **81**,
140–152 (2014).
50. Schmidt-Hieber, C., Jonas, P. & Bischofberger, J. Subthreshold dendritic signal processing
and coincidence detection in dentate gyrus granule cells. *J. Neurosci.* **27**, 8430–8441
(2007).
51. Skaggs, W. E., McNaughton, B. L. & Gothard, K. M. An Information-Theoretic Approach to
Deciphering the Hippocampal Code. in *Advances in Neural Information Processing Systems*
5 (eds. Hanson, S. J., Cowan, J. D. & Giles, C. L.) 1030–1037 (Morgan-Kaufmann, 1993).

Experimental Procedures

Mice

All procedures were conducted in accordance with the U.S. NIH Guide for the Care and Use of
Laboratory Animals and the Institutional Animal Care and Use Committees of the New York State
Psychiatric Institute and Columbia University. Adult male C57BL/6J mice were supplied by
Jackson Laboratory and Dock10^{Cre} mice were a gift from Susumu Tonegawa (Massachusetts
Institute of Technology). Mice were housed in a vivarium grouped 2-4 mice/cage enriched with
running wheels, maintained on a 12-hour light cycle and used at 8-10 weeks of age. Experiments
were conducted during the light phase of the cycle. Food and water were available *ad libitum* until
the beginning of the experiment, when they were placed under controlled water supply and
maintained at >90% of their pre-deprivation weight over the course of imaging experiments. In
total, imaging data from 18 mice were used in this study.

Surgery

Dentate gyrus virus injection and imaging window implantation surgeries were performed as
described previously (Danielson 2016, 2017). For all surgical procedures, mice were anesthetized
with 1.5% isoflurane at an oxygen flow rate of 1 L/min, and head-fixed in a stereotactic frame
(Kopf Instruments, Tujunga, CA). Eyes were lubricated with an ophthalmic ointment, and body
temperature maintained at 37°C with a warm water recirculator (Stryker, Kalamazoo, MI). The fur
was shaved and incision site sterilized prior to beginning surgical procedures, and subcutaneous
saline and carpofen were provided peri-operatively and for 3 days post-operatively to prevent
dehydration and for analgesia. Mice were unilaterally injected with recombinant adeno-associated
virus (rAVV) carrying the GCaMP6s transgene (pAAV.Syn.GCaMP6s.WPRE.SV40), purchased
from Addgene (viral prep #100843-AAV1) with a titer of 1-5x10^{12}, in dorsal dentate gyrus using a
Nanoject syringe (Drummond Scientific, Broomall, PA). Injection coordinates were -1.5 mm AP,
-2 mm ML from the bregma skull suture, and -1.85, -1.7, -1.55 mm DV relative to the cortical
surface. 30 nL of diluted virus was injected at each DV location in 10 nL increments. Mice were
allowed to recover for 3 days and then were implanted with an imaging window over the injected
area and stainless steel head-post for head fixation. Imaging windows were constructed by
adhering 1.8 mm diameter, 2.3 mm long stainless steel hypodermic tubing (Ziggy's Tubes and
Wires Inc, Pleasant Hill, TN) to 1.8 mm diameter glass coverslips (Potomac Photonics,
Halethorpe, MD). A 1.8 mm diameter craniotomy was made centered on the previous injection
site with a taper pointed-drill (Henry Schein Inc, 9004367) and the dura was removed with a micro
curette (FST, 10080-05). The overlying cortex was gently aspirated to reveal capsular fibers

during continuous irrigation with ice-cold aCSF solution and bleeding was controlled with a
collagen gel sponge (Avitene). Under minimal bleeding, a 30g blunt syringe was used to gently
aspirate capsular and CA1 alveus fibers with white appearance and CA1 pyramidale and
moleculare with pink appearance until vasculature of the hippocampal fissure became visible
(under bright light with low bleeding). The imaging implant, attached to the stereotactic handle,
was then gently lowered into the craniotomy and affixed to the skull using dental cement (Unifast
Trad powder and LC light cured acrylic UV, Henry Schein).
637

Behavioral training and apparatus
After a minimum of 1 week recovery period, mice underwent a water restriction scheme
(1ml per day) and trained to run on treadmill while head-restrained. The training period typically
lasted 7-10 days (2 training sessions/day, 15 min each) until the mice were able to run for at least
1 lap/ minute and seek reward from 3 randomly placed reward zones by licking the water delivery
port. We then initiated the motorized belt adjusted to the natural velocity of each mouse and
proceeded training for 2-3 more days. We did not utilize mice that were not getting all of their daily
water supply during treadmill training, and were not motivated to move on the treadmill. A subset
of mice preformed treadmill running without the help of motorized belt (Supplementary Figure 1a).
The behavioral apparatus consisted of a 2m long, 3" wide fabric belt stretched between 6"
diam. laser-cut plastic wheels, mounted on an aluminum frame (8020.net). Spatial triggering of
task events was performed by custom software via serial communication with a microcontroller
(Arduino DUE) and an associated circuit board (OpenMaze OM4 PCB, www.openmaze.org) on
the treadmill. The axle of the treadmill wheel was attached to a quadrature rotary encoder (US
Digital part #: MA3-A10-125-B) connected to a custom quadrature decoder board and Arduino
Nano (courtesy of Wen Li). Angular displacement was converted into a linear distance based on
the circumference of the treadmill. The errors were corrected via a registration anchor marked by
radio-frequency identification (RFID) buttons (SparkFun Electronics) at the lap boundary of the
belt and was detected when it passed over an RFID reader (ID-12LA, SparkFun) affixed
underneath the mouse. A 12V DC gear motor was attached to the axle of the treadmill connected
to a separate Arduino/OpenMaze shield using pulse-width modulation to adjust the rotation
speed. A water reservoir connected to a water delivery port consisting of a small gavage needle
(Cadence Science) was placed within reach of the mouse's tongue. A capacitance touch sensor
(Sparkfun #MPR121) was attached to the water port to measure licking and the sensor was
connected to the Arduino/OM4 PCB. Small 5-10µl drops of water were delivered by the brief
opening a solenoid valve (Parker Hannefin) connected to the water port and controlled by the
OM4 H-bridge circuit. Rewards were triggered at random locations each lap when mice entered
a 10cm long reward zone on the track and were available until mice exited the reward zone or 3
sec had elapsed. Olfactory stimuli consisted of undiluted isoamyl acetate (IAA, Sigma W205532)
which was added to syringe filters (Whatman #6888-2527) and delivered by opening a solenoid
valve (SMC) connected to a flow controller delivering constant airflow of compressed medical
grade air for 1s (~3psi), controlled by another OM4 H-bridge output. Visual and tactile stimulation
consisted of a red LED contralateral to the imaged region, or a 1" square piece of sand paper
brushed by the contralateral whiskers using a stepper motor, at approximately the speed of the
treadmill belt. A custom-written B-Mate behavioral software application implemented in Java was

used for recording mice's licking, its position on the belt, and cue delivery. Mice were monitored
using an IR camera (PS3eye) and illuminated using an IR LED array.
675
To isolate cue-selective responses among the granule cell population, normal cue laps in which
the olfactory, visual, or tactile cue was presented in the middle of the treadmill track (90-110cm)
were interspersed with occasional laps (10-20% of laps) in which the same cue was omitted
("omit" laps), or shifted forward ¼ of the track (0.5m, "shift" laps). For a subset of sessions, the
olfactory cue was presented at one of 5 locations along the track randomly each lap, in order to
examine the effect of spatial pairing of the cue. In all experiments, the treadmill belt material was
changed between sessions to reduce the chances of urine contamination which might act as an
additional olfactory cue.
684
In vivo two-photon imaging
Imaging was conducted using a microscope setup which consists of 8kHz resonant galvanometer
scanner (*Bruker*) mounted to a mirror-based multi-photon microscopy system (*Prairie*
Technologies) and an ultra-fast pulsed laser beam (920-nm wavelength; *Chameleon Ultra II,*
Coherent, 20–40-mW average power at the back focal plane of the objective) controlled with
an electro-optical modulator (*Conoptics*, Model 302 RM). GCaMP fluorescence was
excited through a 40x water immersion objective (Nikon NIR Apo, 0.8 NA, 3.5 mm WD) and
fluorescence signals detected with photomultiplier tubes (*Hamamatsu 7422P-40*), acquired with
PrairieView software (*Prairie*) at 30fps frame rate (512X512 pixels, 1.3 mm/pixel). A custom dual
stage preamp (1.4×10^5 dB, Bruker) was used to amplify signals prior to digitization. Two
goniometers (Edmund Optics) were used to adjust the angle of each mouse's head in order to
achieve the same imaging plane over multiple sessions.
697
Data processing for Ca^{2+} imaging. Movies were motion corrected using the NoRMCorre algorithm
using a non-rigid registration method that splits the field of view (FOV) into overlapping patched
that are registered separately then merged by smooth interpolation[22]. Videos were then spatially
and temporally down-sampled by 2 to reduce noise and reduce the computational power required
for cell segmentation. Spatial and temporal components for individual cells were extracted using
large-scale sparse non-negative matrix factorization[23] or using the singular value decomposition
method by the Suite2p algorithm (https://github.com/cortex-lab/Suite2P), both of which resulted
in similar regions of interest (ROIs). We used Suite2p graphical user interface to manually select
small, densely packed DG granule cells and discard large isolated cell bodies corresponding to
mossy cells or other hilar interneurons. To obtain the total number of DG granule cells within the
imaging fields of view in a subset of total sessions, time averaged images were segmented using
the Cellpose algorithm (https://github.com/MouseLand/cellpose) followed by manual inspection.
Ca^{2+} transient events were defined by a custom detection algorithm which identifies fluorescence
peaks with a rise slope greater than 4 standard deviations above an iteratively refined baseline.
712
713 *Behavioral and Imaging Data Alignment.* Behavioral data were aligned to Ca^{2+} data using the
714 record of a synchronization signal between the two computers used for data collection. Behavioral
715 data were resampled to match Ca^{2+} imaging data frame times.
716

Data Analysis

Data were analyzed using custom-written routines implemented in MATLAB. Plots were generated in MATLAB and Prism and edited in Adobe Illustrator CC 2018.

Identification of spatially-tuned neurons. We restricted our analysis to data from epochs of continuous running at least 2 sec in duration and with a minimum peak speed of 5 cm/sec. For each lap, position data and Ca^{2+} transient events for each cell were binned into 2 cm-wide windows (100 bins), generating raw vectors for occupancy-by-position and calcium transient numbers-by-position which were then circularly smoothed with a Gaussian kernel (SD = 5 cm). A firing rate-by-position vector was computed by dividing the smoothed transient number vector by the smoothed occupancy vector. Within each lap, we circularly shuffled the positions 1000 times and recomputed firing rate-by-position vectors to generate a null distribution for each spatial bin. A spatially selective cell was defined that met the following criteria: (a) the cell should fire above its mean firing rate within its spatial field in at least 20% of laps or for a minimum of 3 laps; and (b) observed firing should be above 99% of the shuffled distribution for at least 5 consecutive spatial bins (10 cm) wrapping around the two edges of the belt. We identified spatially tuned neurons by excluding laps in which sensory cues were omitted or shifted and calculated firing rate vectors in these laps separately. Cells were classified as cue or place cells based upon their peak firing location and activity during laps in which the cue was omitted. Among all of the spatially tuned neurons, "middle cue cells" were defined as those with averaged spatial fields that overlapped with at least 50% of the 45th and 55th bins and had peak amplitude at least two times larger than those in cue-omitted laps. "Lap-cue cells" were defined as those with averaged spatial fields overlapping at least 50% of the region wrapping around the 90th and 10th bins in the normal laps. The remaining cells constituted the "place cells".

Spatial information, stability, consistency, and emergence of spatial fields. To calculate a measure for spatial information content for granule cells in Figure 1, we adapted a traditional method of spatial information assessment [21,51] to Ca^{2+} imaging data. For each cell, we used the firing rate-by-position vector and shuffled null distribution, as computed above, and calculated the spatial information content for each as described previously [51]. To account for the fact that low firing rates artificially produce high spatial information scores, we subtracted the mean of the shuffled information per spike from observed information per spike, divided by the standard deviation of the shuffled values to determine the spatial variance for each cell. Therefore, the amount of spatial information is inferred from differences in normalized Ca^{2+} activity in each neuron and reported as bits per s. To measure spatial field correlation across environments, after identifying spatially tuned neurons, we calculated a Pearson's correlation coefficient between a cell's averaged firing rate-by-position vector in different sessions. The consistency of place field firing was determined as the cross-correlation between the averaged firing rate-by-position vector of normal cue laps from the first and the second halves of the session. We determined place field onset lap in normal cue laps (Figure 1) as described previously [34]. Briefly, starting on lap 1 we searched for a significant Ca^{2+} transient event present within the boundaries of the previously determined mean spatial field calculated from all the laps in the session. If one were found we would then search for Ca^{2+} transient event on each of the next 4 laps. If 3 of the 5 laps had Ca^{2+} transients within the mean place field boundaries, lap 1 would be considered the place field onset lap. If either lap 1

had no Ca^{2+} transient or less than 3 of the 5 laps had Ca^{2+} transient, we would move to lap 2 and
repeat the search. To determine place field onset lap in cue shifted laps (<u>Figure 1</u>), we searched
for the first Ca^{2+} transient event on 2 of 4 consecutive cue shifted laps.

Multi-Session Cell Tracking. Cells were tracked across sessions using CellReg[26]. Briefly, rigid
alignment with both translations and rotations was performed on spatial footprint projections of
each session and manually inspected for quality. To improve performance with our data, we
modified the CellReg source code to consider complete spatial footprints instead of centroids
during alignment. The centroid distance between neighbors was then calculated and used to
create a probabilistic model that estimated the expected error rate at different thresholds. The
optimal centroid distance threshold was chosen by the algorithm and used to match cells. A
clustering algorithm then refined these decisions previously made using pairwise comparisons.
Following cell registration, tracked cells were matched with their corresponding functional
cell types (i.e. mid-, lap-cue, place cells, as described above). All analyses presented in Figure 2
are carried out in pairwise, to maximize the number of cells in each comparison and to minimize
the total number of comparisons. For multiday comparisons we used Day1, session 1 as the
normal session, and for multisession comparisons we used the visual stimulus session as the
reference. To calculate the fraction of cells that maintain their identity, cell pairs that were counted
as being the same cell type in both sessions was divided by all of that cell type in the normal
session. In order to derive a null distribution for preservation of pairwise identity, we randomly
permutated the cell IDs of all the tracked cells in pairwise sessions 1000 times and calculated the
fraction of cells that were the same, among all of that cell type in the normal session. We
calculated p-values by comparing actual data to this null distribution, 97.5th% of the null
distribution is presented dotted lines in Figure 2.

Spatial event rate and Population vector (PV) correlation analysis. Comparison of the activity
between different sessions was calculated using Pearson's correlation of the spatially binned,
averaged firing rate-by-position vector in cue normal laps of all recorded cells in 8 mice for
multiday analysis (1 session from each condition) and 6 mice for multisession analysis (1 session
from each condition) in Figure 2.The variability in neural activity between lap types was calculated
by using Pearson's correlation on each 2 cm bins of the firing rate-by-position vector along the
treadmill during odor cue trials (mean for all spatially tuned cells, 4,091 total spatially tuned cells
from 8 mice across 66 sessions).

Spatial modulation
To compare single cell cue responses at different locations, cue-triggered average z-scored
calcium transients were measured for cue cells, defined based upon omit laps as done previously.
For dual cue location experiments, cue cells were identified as cells that whose cue-zone calcium
event rate was greater than 95% of shuffled responses versus the cue-omitted laps. Cue cells
that significantly preferred one location were counted when responses at one cue location
exceeded 95% of the shuffled cue position event rates. Spatial modulation index in this paradigm
is listed as a ratio of mean spatial firing rates for the nonpreferred location over the preferred
location rates for each cell.

Bayesian Reconstruction Analysis. To calculate the probability of the animal's position given a
short time window of neural activity, we used a previously published method based on Bayesian
reconstruction algorithm [25]. Briefly, Ca^{2+} transient events for each cell were binned into 1 second
windows to construct firing rate vectors. For each of these binned firing rate vectors, Bayesian
classification of virtual position (posterior probability for each bin) was performed by a previously
described method [25] utilizing a template comprising of a cell's smoothed firing rate-by-position
vectors. In order to cross-validate our decoding procedure, we divided firing rate-by-position
template into lap crossings, used $1/5^{th}$ of laps as "testing" dataset while the remaining $4/5^{th}$ of laps
constituted the "training" dataset. For example, lap 1 was tested based on the firing rate-by-
position vectors calculated using laps 2,3,4,5, lap as template, and lap 6 was tested based on the
firing rate-by-position vectors calculated from laps 7,8,9,10, and so on. The resulting posterior
probability distribution for each bin is the likelihood for an animal is located in that bin, which adds
up to 1, and the bin with the maximum posterior probability is the estimated position of the animal.
To determine the decoding error we calculated the absolute difference between the animal's
actual position and the maximum posterior probability in that bin. Post-reconstruction, we divided
the time bins (excluding those with no activity) according to the lap types.
821
Cue shift analysis
Spatial firing rates for each spatially tuned cell on normal middle cue laps were cross-correlated
with firing rates on shift laps in order to estimate the shift magnitude for each cell after cue
manipulation. Binned histograms of numbers of cells with peak firing rates at particular locations
along the track were plotted with respect to their shift magnitudes on cue shift laps. The spatial
distribution of cells showing no shift (<5 cm shift, "non cue-influenced") or that shifted their firing
along with the cue (-50cm +/-5cm, "cue influenced") was then compared.
829
Inhibition analysis
Out-of-field firing was calculated for spatially tuned cells as the spatial event rate, excluding the
peak place field for that cell (+/- 10cm). Average out-of-field firing rates were then calculated
across all cells for normal middle cue laps and intermittent laps where the cue was omitted. For
comparison of cue-associated excitation and inhibition levels, average firing rates were computed
by session for the 20cm region surrounding the middle cue, with respect to the normal pre-cue
baseline firing rate. Cue-related inhibition of place cell firing was calculated by selecting spatially
tuned cells whose firing field on normal middle cue laps fell within the region of the cue on shift
laps (50-80cm). Firing rates for these cells were then averaged for normal laps where the cue
was not presented in this location and compared with laps where the cue was shifted to this region
(50cm). Mutual inhibition between cues was calculated by first selecting cue cells responsive to
each of 2 cues of different modalities (olfactory or visual, presented at 40cm and 120cm,
respectively) based upon their activity when each cue was omitted, as described above for single
cues. Responses were then averaged for each cue cell for laps in which the cues were presented
alone at these locations versus intermittent laps where the cues were presented together at one
of the two locations.
Data and Software Availability
Data and custom programs are available upon reasonable request.
848

Figures

Figure1: Robust representation of sensory cues in the dentate gyrus
a) Two-photon imaging of DG population calcium activity. Left, diagram of the imaging window implant in the dorsal dentate gyrus. Right, Time averaged *in vivo* two-photon image of GCaMP6s-expressing granule cells.
b) Diagram of head-fixed treadmill apparatus for spatial sensory cue delivery.

c) Fraction of spatially tuned cells at each linearized treadmill position (bins=10cm) per session, n = 4091 cells from 8 mice across 6-9 sessions/mouse, locations of the odor and lap cue are shown blue and green shaded areas, respectively.

d) Spatial patterns of DG neuron activity during cue task. Top: Location of lap cue (green boxes) and an odor cue (blue box), on normal, shift or omit laps. Bottom: Lap-averaged spatial firing rates of 285 spatially tuned neurons (n=8 mice) during their first session with the cue on normal middle location (left), cue-shifted (middle) and cue-omitted (right) laps. Each row across all graphs represents a single cell, and the x axis represents the treadmill position.

e) Average firing rates by position for neurons shown in panel D on normal (black), cue-shifted (red) and cue-omitted laps (blue, mean ± SEM). (Inset) averaged area under the firing rate curves (Hz*cm) within the middle cue region during normal (black bar), cue omitted (blue bar), and cue shifted laps (red bar). $P_{normal-omit}$ <0.0001, $P_{normal-shift}$ =0.02, Wilcoxon matched-pairs signed rank test, error bars are mean ± SEM.

f) Example fluorescence traces from odor-cue, lap-cue and place cells within a single session. Black, red, and blue traces represent normal, cue shifted and omitted laps, respectively. Scale bars are cm and standard deviation from each cell's baseline fluorescence.

g) Z scored spatial information of cue and place cell populations. Z scored spatial information, χ^2=48.47, p<0.0001, $P_{OdorCue-LapCue}$=0.0387, $P_{OdorCue-Place}$=0.0028, $P_{LapCue-Place}$<0.0001.

h) Tuning consistency of cue and place cells. Firing rate correlation between first and last halves of the session, χ^2=32.60, p<0.0001, $P_{OdorCue-LapCue}$=0.2224, $P_{OdorCue-Place}$=0.0117, $P_{LapCue-Place}$<0.0001.

i) Emergence of cue and place responses. Cumulative distribution of spatial field onset lap in normal laps for cue and place cells, χ^2=9.29, p=0.0096, $P_{OdorCue-LapCue}$=0.8925, $P_{OdorCue-Place}$=0.0194, $P_{LapCue-Place}$=0.0091. (Inset) Spatial field firing rates of cue and place cells during -1-5th, 6-10th and 11-15th laps, main effect of cell type: $F_{2,1473}$=6.73, p=0.0012; cell type × lap number interaction $F_{4,1473}$=2.44, p=0.04; main effect of lap number: $F_{2,1473}$=0.28, p=0.7; Lap 1-5th, $P_{OdorCue-LapCue}$=0.99, $P_{OdorCue-Place}$=0.0011, $P_{LapCue-Place}$<0.0001, 2-way ANOVA and Tukey's multiple comparisons test.

$N_{OdorCue}$=114, N_{LapCue}=220, $N_{PlaceCell}$=160, from 8 mice 2 sessions each. Red dotted lines in violin plots show median, black dotted lines show quartiles. Comparisons are Kruskal Wallis and Dunn's multiple comparisons tests unless otherwise noted. *P < 0.05; **P < 0.01; ***P < 0.001.

See also Supplementary Figures 1-4.

Figure 2: Sensory cue and place responses differ in stability and selectivity

a) Spatial firing rates of individual DG granule cells matched between sessions within a day and over one week. Top: diagram of odor cue presentation during all sessions. Bottom: Spatial firing rates for spatially tuned neurons tracked during subsequent sessions on the same day or one week later, ordered according to the position of peak activity during the first exposure (n=233 cells, 8 mice).

b) Activity of granule cells in response to different sensory cues. Top: diagram of behavior in sessions with cues of different sensory modalities in the middle position and an invariant lap cue.

Bottom: Spatial tuning for neurons tracked through consecutive sessions during exposures to different cues. Data are shown for neurons with significant tuning in visual cue session and tracked in olfactory and whisker tactile cue sessions (n=196 cells, 6 mice).

c) Fraction of cross-session registered cells that encoded the same variable, lap cue (green), middle cue (blue), or place (orange), within same day (left), across days (middle), and during exposures to different sensory modalities (right) . Cells encoding the same variable on the same day, χ^2=1.81, p=0.4039, $P_{LapCue-OdorCue}$=0.4221, $P_{LapCue-Place}$=0.549, $P_{OdorCue-Place}$=0.9713; cells encoding the same variable across days, χ^2=14.36, p<0.001, $P_{LapCue-OdorCue}$=0.1566, $P_{LapCue-Place}$<0.0001, $P_{OdorCue-Place}$=0.0313 (8 mice, 8 matched sessions, Day1-Day1', Day1-Day7); cells encoding the same variable with different sensory modalities, χ^2=19.78, p<0.00001, $P_{LapCue-OdorCue}$<0.00001, $P_{LapCue-Place}$=0.0766, $P_{OdorCue-Place}$=0.0464 (6 mice, 12 matched sessions, Odor-Vis. and Vis.-Tact.); Kruskal-Wallis test, Dunn's multiple comparisons test. Dashed lines represent 97.5th% of null distributions for each cell type. Error bars, mean ± SEM.

d) Fractions of spatially tuned cells that stably encode only one variable (cue or place, pink) or switched response types (cue to place or vice versa, black) within same day, across days and in response to different sensory cues: $P_{Switch-Stable\ (Same\ Day)}$<0.001, $P_{Switch-Stable\ (Across\ Days)}$<0.001, $P_{Switch-Stable\ (Sensory\ Modalities)}$<0.001, Wilcoxon Rank Sum test. Dashed lines: 2.5th and 97.5th% of null distributions. Error bars, mean ± SEM.

e) Mean rate correlations within the same day (red), different days (blue) and different sensory cues (green) for mid-cue cells (left, χ^2=27.88, p<0.0001, $P_{Same\ Day-Across\ Days}$=0.2465, $P_{Same\ Day-Sensory\ Modalities}$<0.0001, $P_{Across\ Days-Sensory\ Modalities}$=0.0038, $N_{Same\ Day}$=37, $N_{Across\ Days}$=37, $N_{Sensory\ Modalities}$=55); place cells (middle, χ^2=18.79, p<0.0001, $P_{Same\ Day-Across\ Days}$<0.0001, $P_{Same\ Day-Sensory\ Modalities}$=0.7449, $P_{Across\ Days-Sensory\ Modalities}$=0.0055, $N_{Same\ Day}$=76, $N_{Across\ Days}$=76, $N_{Sensory\ Modalities}$=83), lap-cue cells (right, χ^2=5.096, p=0.0782, $P_{Same\ Day-Across\ Days}$=0.0723, $P_{Same\ Day-Sensory\ Modalities}$>0.9999, $P_{Across\ Days-Sensory\ Modalities}$=0.7932, $N_{Same\ Day}$=80, $N_{Across\ Days}$=80, $N_{Sensory\ Modalities}$=59). Kruskal-Wallis test, Dunn's multiple comparisons test. Boxes, 25th to 75th percentiles; bars, median; whiskers, 99% range. *P < 0.05; **P < 0.01; ***P< 0.001.

f) Activity of DG granule cells in response to the same odor-cued track when recorded in different rooms. Top: diagram of exposures to an odor in the middle position and an invariant lap cue, performed in different rooms. Bottom: Spatial firing rates for tuned neurons tracked through consecutive sessions in different rooms (n=83 cells, 4 mice).

g) Fraction of cross-session registered cells that encoded the same variable, odor cue (blue), lap cue (green) or place (orange) in different recording rooms. χ^2=6.59, p=0.0372, $P_{OdorCue-Place}$=0.0490, $P_{LapCue-OdorCue}$=0.9557, $P_{LapCue-Place}$=0.0972. Dashed lines: 2.5th and 97.5th% of null distributions. Error bars, mean ± SEM.

h) Comparison of spatial firing rate correlations in all tuned cells across rooms (χ^2=15.85, p<0.001, $P_{Odor-Lap\ cue}$=0.27, $P_{Odor\ Cue-Place}$ <0.001, $P_{Lap\ Cue-Place}$ =0.03, $N_{Odor-Cue}$=25, $N_{Lap-Cue}$=33, N_{Place}=26). Kruskal-Wallis test, Dunn's multiple comparisons test. Boxes, 25th to 75th percentiles; bars, median; whiskers, 99% range. *P < 0.05; **P < 0.01; ***P< 0.001. See also Supplementary Fig. 5.

Figure 3: Sensory cue responses are spatially modulated

a) Experimental setup to examine effects of cue-location pairing on cue responses. Cue responses are measured with respect to a typical middle cue location or an intermittently shifted cue location in the same sessions, or with random administration of the same cue on each lap in a subsequent session.

b) Example cue-triggered average Ca^{2+} transients from two cue cells on normal (black), cue-shifted (blue) and random-cue (pink) laps. Scale bars are sec and standard deviation from each cell's baseline fluorescence for the session.

c) Z-scored peak amplitudes of Ca^{2+} transients from odor cue cells in normal, shift and random cue presentation conditions; χ^2=49.36, p<0.001, $P_{Normal-Shift}$<0.0001, $P_{Normal-Random}$<0.0001, $P_{Shift-Random}$>0.9999, Friedman test and Dunn's multiple comparisons tests (n=101 cells, 5 mice). Red lines in violin plots show median, black lines show mean.

d) Difference in the magnitude of cue responses in normal laps from random-cue and cue-shift laps. Yellow line, linear regression (R^2= 0.546, p=3.5x10^{-9}). Pie chart shows the percentage of neurons with higher average responses in normal (black), cue-shift (blue), random cue presentation conditions (red), and both shifted and random laps (purple).

e) (Top) Dual cue location task: LED visual cue is given consistently at two positions on the track. Arrangement of cues for normal laps (left, 80% of laps), laps where first cue is omitted (middle, 10%), and laps where the second cue is omitted (right, 10%). (Bottom) Spatial firing rates of 433 significantly tuned cells on first day of task (N=3 mice).

f) Average spatial firing rates for significant cue cells (vs. cue omitted laps), for cells preferring cue position 1 (top) or cue position 2 (bottom) on first day of task (blue, n=25 pos1, n=25 pos2) and 4th day of task (pink, n=13 pos1, n=20 pos2 cells).

g) Spatial modulation index (rate non-preferred cue/preferred cue) for all cue cells on days 1-4. (p=0.0125, day 1 vs. days 2, 3, or 4, n= 50/40/50/33, Wilcoxon rank sum test).

h) Empirical cumulative probability distribution for all cue cells on days 1-4.

See also Supplementary Figure 6.

Figure 4: Cue shifts have limited effects on spatial encoding

a) Population vector correlations for all spatially tuned cells for normal middle cue laps and shifted cue laps in cue shift task. Locations of the middle/shift and lap cue are shown blue and green shaded areas, respectively.

b) Bayesian decoding of 4 normal and 1 cue shifted laps for a representative session, based upon activity from all spatially tuned granule cells. The magenta line shows predicted position at each bin, while the green line shows the observed position of the animal.

c) Average decoding error for normal and cue shifted laps calculated as the absolute median distance between the decoded value in each time bin and the actual value of the position (p<0.0001, Wilcoxon signed-rank test, n=66 sessions). Boxes, 25th to 75th percentiles; bars, median; whiskers, 99% range.

d) Decoding error for each treadmill position in normal (black) and cue shifted laps (red). Shaded error region represents the difference of the median from the null distribution of median decoder error. Dotted line indicates point of statistical equality between normal and cue shifted laps.

e) Distribution of cell firing location shift on shifted cue laps for all spatially tuned cells, plotted based upon normal tuning location (n=66). Red arrowhead indicates cue shift distance, and thus marks cells that are directly influenced by the cue. Black arrowhead indicates cells with zero shift, and thus not influenced by the cue.

f) Mean number of cells not influenced by the odor cue (shift magnitude<0.05m) and cells shifting precisely along with the cue (shift magnitude 0.5m+/-0.1m) at each track position (corresponding to cells in regions of black and red arrowheads in "b", respectively). Cue influenced cells are largely confined to locations around the cue itself, while non cue-influenced place cells are more evenly distributed, with a slight enrichment immediately preceding the middle cue.

See also Supplementary Figure 7.

Figure 5: Cue presentation leads to suppression of diverse types of DG responses

a) Cue-related suppression of noisy, out-of-field firing. Spatial firing rates outside of the field center-of-mass (+/-10cm) for significantly tuned cells during normal and cue omitted laps (n=4,091 cells from 66 sessions).

b) Average out-of-field firing rates from spatially tuned cells in the presence of the middle cue (black) compared to laps in which the cue is omitted (blue). Blue shaded area shows the cue delivery position. Inset, mean ± SEM of firing rate, p<0.001, rank-sum test, n=4,091 cells.

c) Comparison of the magnitude of cue-related excitation and inhibition. Session averaged in-field firing rate vs. out-of-field firing rate (66 sessions). Yellow, linear regression: p<0.0001, R^2=-0.22.

d) Cue-related suppression of place cell firing. Spatial firing rates of place cells with firing fields at 40-80cm during normal laps (left) and laps in which the cue is shifted to 40cm (right).

e) (Left) Average firing rate by position for the above place cells in normal laps when the cue is not presented at this location (Cue⁺, black), compared with laps in which the cue is shifted to this location (Cue^Δ, red; diff. in green). (Right), Average peak firing rate of the same cells in "d" and "e", paired t-test, p<0.001, 645 cells.

f) Cue-related suppression of cells responding to other cues. (Top) diagram of intermittent cue pairing experiment. Cues of two different modalities were presented at different locations, interspersed with paired presentation of these two cues at one of these locations. (Bottom) example cue-triggered averaged Ca^{2+} transients for a cue cell strongly active when the cue is presented alone (black) but with reduced responses when paired with a different cue, regardless of location (blue, pair location #1, purple, location #2).

g) (Left) Amplitudes of paired cue responses at two pairing locations with respect to the response to the cue alone. Gray line represents the diagonal, $p<0.0001$, $R^2=0.91$, n=56 cells, 3 mice. (Right) Average relative response amplitude at the two pairing locations (Location 1: 0.75 ± 0.061, $p1 = 1.45\times10^{-4}$; Location 2: 0.73 ± 0.086, $p2= 0.003$, Wilcoxon Signed Rank Sum test, bar plots are mean ± SEM).

See also Supplementary Figure 8.

Supplementary Figures

Supplementary Figure 1. Motor driven treadmill does not influence spatial properties of DG neurons.
a) DG population activity is dependent on the locomotion of the animal, shaded error bar represents mean Ca²⁺ events rate and SEM binned by normalized velocity (190 spatially tuned neurons in 5 mice, 1 session/mouse). **b-d)** Comparison of cell firing for self-driven vs. motorized treadmill. **b)** Fraction of active cells that are spatially tuned (with at least 0.001 transients per s) in mice advancing the treadmill belt through self-driven locomotion (black, 190 spatially tuned neurons in 5 mice, 1 session/mouse) and mice running on the motorized treadmill (red, 285 spatially tuned neurons in 8 mice, 1 session/mouse,) p=0.435, Wilcoxon Rank Sum test.
c) Firing rate of spatially tuned cells in self-driven vs. motorized treadmill (p=0.1876, Wilcoxon Rank Sum test).
d) Z scored spatial information in mice running on motorized treadmill vs. self-driven locomotion (p<0.0001, Wilcoxon Rank Sum test).

Supplementary Figure 2. Single visual and tactile cues, reward and multiple salient cues evoke robust responses in DG.

a) Firing of DG neurons in response to an LED visual cue. Top: Location of lap cue (green boxes) and an LED cue (red box), during normal and cue-omitted trials. Bottom: spatial firing rates of 188

spatially tuned neurons (n=6 mice) during the first exposure to the middle visual cue on normal (left), and cue-omitted (right) laps.

b) Firing of DG neurons in response to a whisker tactile cue. Top: Location of lap cue (green boxes) and a whisker tactile cue (brown box), during normal and omit trials. Bottom: spatial firing rates of 199 spatially tuned neurons (n=8 mice) during the first exposure to the middle tactile cue on normal (left), and cue-omitted (right) laps. Each row represents a cell, and the x axis represents the treadmill position.

c) Average spatial firing rate for neurons shown in "e" (LED cue) on normal (black, mean ± SEM), and cue-omitted (blue) laps. Inset shows the averaged area under the firing rate curves (Hz*cm) within the visual cue region during normal laps (black bar) compared to the same region during cue omitted laps (p<0.0001, Wilcoxon Signed Rank Sum test).

d) Average firing rates of neurons shown in "f" (tactile cue) on normal (black), and cue-omitted (blue) laps. Inset shows the averaged area under the firing rate curves (Hz*cm) within the tactile cue region during normal laps (black bar representing blue area under the middle region of the treadmill) compared to the same region during cue omitted laps (p<0.0001, Wilcoxon Signed Rank Sum test).

e) Top: Behavioral task with a static operant reward at 75% track length, preceded by an olfactory cue in middle of track. Bottom: Spatial firing rates for all spatially tuned cells in the static reward task. Laps with olfactory cue in middle of track and reward at 75% track length (static rew., left), or laps with reward omitted (omit rew., right). (n=256 cells, 4 mice, 7 sessions).

f) Averaged spatial lick rates over laps, for behavioral sessions in "e".

g) Average spatial firing rates over all cells in "e" during static rew. laps (blue), and omit rew. laps (red). Olfactory cue zone shaded in yellow, reward zone shaded in orange.

h) Averaged area under the firing rate curves for cue zone and reward zone on static reward and omit reward laps. $p = 3.8e^{-9}$, reward zone rate during rew+ vs. rew- (omit rew.) laps, Wilcoxon signed rank test.

i) (Top, left) Experimental setup with 3 cues, cue 1 = mint odor (olfactory), cue 2 = LED (visual), cue 3 = isoamyl acetate odor (olfactory). Individual cues are omitted on intermittent laps (right). (Bottom) Spatial calcium event rates for the same tuned cells in normal 3-cue laps (left), and intermittent laps where individual cues are omitted (right 3 cols., as labeled, n=138 cells from 5 mice, 1 session/mouse, sorted by normal 3-cue lap activity).

j) Mean calcium event rates over all cells in "e" on normal laps (black), laps where cue 1 is omitted (green), cue 2 omitted (red), or cue 3 omitted (blue).

k) Averaged area under the firing rate curves within zones corresponding to cues 1-3 on normal laps (black), laps where cue 1 is omitted (green, p= 0.0031), cue 2 omitted (red, p<0.001), or cue 3 omitted (green, p=0.0047). Wilcoxon matched-pairs signed rank test, error bars are mean ± SEM.

Supplementary Figure 3. Representations of sensory cues and place in a granule cell specific transgenic mouse, Dock10-cre

a) Mean spatial information by position. Average Z scored spatial information for cells binned by tuning position during on-line cue manipulation task. WT dorsal dentate gyrus neurons express GCaMP6s (gray bars) and Dock10-cre labelled granule cells express FLEX-GcamP7s (purple bars). Error bars represent SEM, $P_{spkZ-DG-spkZ-GC}$ = 0.15, Wilcoxon rank-sum test calculated over cumulative mean spkZ.

b) Confocal image projection of brain slice from Dock10-cre mouse injected with FLEX-eYFP AAV in the dorsal dentate gyrus (blue=DAPI, green=eYFP).

c) Spatial firing rates of Dock10/GCaMP7s-expressing granule cells in treadmill task with an olfactory cue. (Left) normal middle cue responses, (middle) cue shift laps, (right) laps in which odor cue is omitted (n=146 cells from 4 sessions in 2 mice). Cells sorted by activity on normal middle cue laps

d) Average spatial firing rates from all cells in "c" during normal middle cue (black), cue shift (red), and cue omitted laps (blue). (Inset) averaged area under the firing rate curves (Hz*cm) within the

middle cue region during normal (black bar), cue omitted (blue bar), and cue shifted laps (red bar). $P_{Normal\text{-}Omit}$ <0.001, $P_{Normal\text{-}Shift}$ =0.03, Wilcoxon matched-pairs signed rank test, error bars are mean ± SEM.

e) A representative *in vivo* two-photon field of view within the dentate gyrus of C57 mice expressing GCaMP6s (left) and Dock10-cre mouse expressing FLEX-GCaMP7s in granule cells (right), including spatially scattered odor (blue) and lap (green) cue cells and place cells (orange).

f) Fraction of inactive cells (gray), active cells (with at least 0.001 transients per s, pink) and spatially tuned cells (orange, green, blue). $Mean_{C57\text{-}GcamP6s}$ = 631.68±11.9 cells, from 8 mice 2 sessions each, $Mean_{Dock10Cre\text{-}GcamP7s}$ = 566.11±18.4 cells, mice/session from 2 mice 2 sessions each. Small, round somata corresponding to granule cells within the granule cell layer were counted using the *Cellpose* algorithm followed by manual inspection in a subset of total sessions imaged from one mouse.

g) Cumulative distribution curve of the firing rate of odor cue, lap cue and place cells in Dock10-cre mice; Odor Cue $P_{C57\text{-}GcamP6s\text{-}Dock10Cre\text{-}GcamP7s}$=0.11, Lap Cue $P_{C57\text{-}GcamP6s\text{-}Dock10Cre\text{-}GcamP7s}$ = 0.05, Place cell $P_{C57\text{-}GcamP6s\text{-}Dock10Cre\text{-}GcamP7s}$ = 0.05, Comparisons are Wilcoxon rank-sum test.

Supplementary Figure 4. Odor responsive cells are not spatially tuned on an uncued belt.
a-c) Comparison of cell tuning properties between uncued vs. cued treadmill belts. **a)** Spatial firing rates of all tuned DG neurons within the same fields of view (FOV) recorded during paired sessions on an uncued belt (left) and odor cued belt (right). Neurons are ordered according to the

position of their peak rate in each belt (uncued: 155, cue: 249 spatially tuned neurons in 6 mice, 1 session/mouse), and are not matched between uncued and cue belts.

b) Fraction of active cells that are spatially tuned (with at least 0.001 transients per s) in matched FOVs recorded on uncued and cue belts (p=0.0313, Wilcoxon signed-rank test, n=6).

c) Z scored spatial information of all spatially tuned neurons in uncued (n=155) and cued sessions (n=193), excluding odor-cue responsive neurons (p=0.0897, Wilcoxon rank-sum test).

d) Tuning consistency (firing rate correlation between first and last halves of the session) of all spatially tuned neurons in uncued and cued sessions, excluding odor-cue responsive neurons (p=0.5237, Wilcoxon rank-sum test). Boxes, 25th to 75th percentiles; bars, median; whiskers, 99% range.

e-h) Comparison of registered cue and place cell responses between uncued and cue belts. **e)** Activity of cross-registered odor cue cells in uncued and cue belts. Cells were classified as odor-cue cells in cue belt and then their responses were determined in uncued belt. Note that these cue cells were active but not spatially tuned in uncued sessions.

f) Fraction of odor-cue (blue), lap-cue (green) and place cells (orange) that are cross- registered in uncued belt sessions, and are therefore active but not necessarily spatially tuned (p=0.838, n=6 matched sessions, Friedman's test).

g) Fraction of registered cells that are spatially tuned in uncued belt sessions and encoded the odor cue (blue), lap cue (green), or place cells (orange) on the cued belt (χ^2=10.88, p=0.0043, $P_{OdorCue-LapCue}$=0.0066, $P_{OdorCue-Place}$=0.0286, $P_{LapCue-Place}$=0.9076). Gray area represent 2.5th and 97.5th% of null distributions for each cell type. Error bars, mean ± SEM.

h) The fraction of cross-registered odor cue cells that were spatially tuned on the uncued belt is significantly below the null distribution (left) while the fraction of lap-cue (middle) and place cells that are spatially tuned remain significantly above the null distribution. The null distributions are generated for each cell type by randomly permutating cell IDs of all cross-registered neurons and determining overlap among cell types. (Level of significance for 5,000 shufflings **p < 0.01; *p < 0.05, $N_{Odor-cue}$=35, $N_{Lap-cue}$=54, N_{Place}=58).

Supplementary Figure 5. Multisession tracking of individual cue and place cells

a) Representative alignment of spatial footprints for cells segmented across sessions within a day or over 1wk. in a single imaging field.

b) Distribution of centroid distances between registered (blue, 1.45 ± 0.97, n=1604) and non-registered (pink, 7.07 ± 1.5, n=2214) neighboring cell pairs (p=6.9 10^{-43}, Wilcoxon Rank Sum test).

c) Fraction of all spatially tuned cells in day 1 session1 (D1), day 1 session 2 (D1') and day 7 (D7) that are registered to at least one other session, χ^2=0.25, p=0.9674, $P_{D1-D1'}$>0.9999, P_{D1-D7}>0.9999, $P_{D1'-D7}$>0.9999, n=8 matched sessions, Friedman and Dunn's multiple comparisons tests.

d) Comparison of average firing rates in all tuned cells across days regardless of tracking χ^2=4.993, p=0.0824, $P_{D1-D1'}$ >0.9999, P_{D1-D7} =0.0781, $P_{D1'-D7}$ =0.4789, N_{D1} =417, $N_{D1'}$ =365, N_{D7}=338, from 8 matched sessions, Kruskal Wallis and Dunn's multiple comparisons tests .

e) Representative fields of view with odor cue cells (blue), lap cue cells (green), and place cells (orange) tracked within a day and across days.

f) Representative Ca²⁺ transients for an odor cue cell that has stable cue-selective activity within one day and over 1wk (left). Transients for place cells show relatively stable firing location within

day but a reorganization of spatial selectivity across days (right). Black and red traces represent normal and cue shifted laps, respectively.

g) Firing rates of tracked neurons in visual and whisker tactile cue sessions ordered according to the position of peak activity in olfactory cue sessions (n=157 cells, 6 mice).

Supplementary Figure 6. Spatial modulation of sensory cue responses
a) Dual cue location task: Average spatial firing rates for cue cells preferring cue position 1 (top) and cue position 2 (bottom) on normal laps (blue), laps where the cue at position 1 is omitted (green), and laps where the cue at position 2 is omitted (cyan). (n=89 pos1 cells, 67 pos2 cue cells).
b) Average peak firing rates for all cue cells (both cue 1 and cue 2 preferring) for preferred cue location (left) and non-preferred location (right) on normal laps (blue), laps where the opposite cue is omitted (orange), and laps where the same cue (preferred or non-preferred) is omitted. (n=156 cells, positions1&2) (mean ± SEM: 0.205 ± 0.017, 0.173 ± 0.020, 0.017 ± 0.008, 0.137 ± 0.016, 0.140 ± 0.024, 0.025 ± 0.008, $P_{\text{Preferred Cue}}$ = 7.16x10^{-9}, $P_{\text{Non-preferred Cue}}$ =3.48x10^{-9}, Wilcoxon sign rank test).
c) Average spatial firing rates for cue cells preferring cue position 1 (top) and cue position 2 (bottom) on days 1 (blue, n=50), 2 (green, n=40), 3 (cyan, n=50), and 4 (magenta, n=33) of dual cue location task.
d) Cue location firing rates for preferred and non-preferred cue location for all cue cells on days 1 (blue, n=50) and 4 (magenta, n=33) of dual cue location task.

Supplementary Figure 7. Effect of sensory cues on spatial coding during cue-omitted laps
a) Population vector correlations for all spatially tuned cells at each treadmill position for normal middle cue laps (black) and omitted cue laps (blue). Locations of the middle and lap cue are shown blue and green shaded areas, respectively.
b) Bayesian decoding: Cumulative distribution functions of the error of the maximum likelihood estimate of position for 30 example recording sessions.
c) The decoder error for spatially tuned neurons plotted against their firing rate.
d) Decoding error for each treadmill position in normal (black) and cue omitted laps (blue). Shaded error region represents the difference of the median from the null distribution of median decoder error.
e) Average decoding error for normal and cue omitted laps calculated as the absolute median distance between the decoded value in each time bin and the actual value of the position. p<0.0001, Wilcoxon signed-rank test (n=66 sessions). Boxes, 25th to 75th percentiles; bars, median; whiskers, 99% range.
f) Effects of cue omission on individual cells: Color histogram of all spatially tuned cells, plotted based upon spatial field location and difference in firing rate between normal cue and cue-omitted laps.
g) Average difference in normal middle cue and cue-omitted lap firing rates based upon cell spatial field location. Note that the largest cue effects are confined to cells immediately responsive to the middle cue. With respect to pre-cue cells, Wilcoxon Rank Sum test: *** = p < 10e-10, * = p = 0.03, n.s.= not significant.

Supplementary Figure 8. Timing of cue-related inhibition in the DG

a) Average in-field spatial firing rates (i.e. cue-related excitation) in the region around the middle cue location for all tuned neurons, compared with out-of-field firing rates (i.e. cue-related inhibition), and adjusted for pre-cue firing rate. The position of the excitation peak precedes the nadir of inhibition. Blue shaded area shows the cue delivery position.

b) (Left) Session averaged excitatory (within-field) peak position vs. inhibitory (out-of-field) peak position (66 sessions). Red = avg., diagonal (gray line). (Right) Quantification of the position of excitation peak compared to the position of inhibition peak, p<0.01, Wilcoxon Signed Rank Sum test.